Title: Grasping with a twist: Dissociating action goals from motor actions in human frontoparietal circuits

Running Title: Decoding of isolated and task-directed grasping

Authors: Guy Rens[1,2,3], Teresa D. Figley[4], Jason P. Gallivan[5], Yuqi Liu[6,7], Jody C. Culham[1,4]

[1]Department of Psychology, University of Western Ontario, London, Ontario, Canada
[2]Laboratorium voor Neuro- en Psychofysiologie, Department of Neurosciences, KU Leuven, Belgium
[3]Leuven Brain Institute, KU Leuven, Belgium
[4]Graduate Program in Neuroscience, University of Western Ontario, London, Ontario, Canada
[5]Departments of Psychology & Biomedical and Molecular Sciences, Centre for Neuroscience Studies,
Queen's University, Kingston, Ontario, Canada
[6]Department of Neuroscience, Georgetown University Medical Center, Washington, DC, 20057, USA
[7]Institute of Neuroscience, CAS Center for Excellence in Brain Sciences and Intelligence Technology,
Chinese Academy of Sciences, Shanghai, China
*Corresponding Author:
Guy Rens
The Brain and Mind Institute
University of Western Ontario
Ontario N6A 3K7, Canada
grens@uwo.ca

Acknowledgements: This work was supported by grants from the Canadian Institutes of Health
Research Operating Grant (MOP84293 to J.C.C.), Western University Strategic Success Program
(J. C. C.), the Natural Sciences and Engineering Research Council of Canada (to J.C.C.) and a
CIHR-Doctoral Research Award 215131 (T.D.F).

Conflict of interest: The authors declare no conflicts of interest.

Abstract

In daily life, prehension is typically not the end goal of hand-object interactions but a precursor for

manipulation. Nevertheless, functional MRI (fMRI) studies investigating manual manipulation have

primarily relied on prehension as the end goal of an action. Here, we used slow event-related fMRI to

investigate differences in neural activation patterns between prehension in isolation and prehension for

object manipulation. Sixteen participants were instructed either to simply grasp the handle of a

rotatable dial (isolated prehension) or to grasp and turn it (prehension for object manipulation). We

used representational similarity analysis to investigate whether the experimental conditions could be

discriminated from each other based on differences in task-related brain activation patterns. We also

used temporal multivoxel pattern analysis to examine the evolution of regional activation patterns over

time. Importantly, we were able to differentiate isolated prehension and prehension for manipulation

from activation patterns in the early visual cortex, the caudal intraparietal sulcus, and the superior

parietal lobule. Our findings indicate that object manipulation extends beyond the putative cortical

grasping network (anterior intraparietal sulcus, premotor and motor cortices) to include the superior

parietal lobule and early visual cortex.

Significance statement

A simple act such as turning an oven dial requires not only that the central nervous system encode the

initial state (starting dial orientation) of the object but also the appropriate posture to grasp it in order

to achieve the desired end state (final dial orientation) and the motor commands to achieve that state.

Using advanced temporal neuroimaging analysis techniques, we reveal how such actions unfold over

time and how they differ between object manipulation (turning a dial) vs. grasping alone. We find that a

combination of brain areas implicated in visual processing and sensorimotor integration can distinguish

between the complex and simple tasks during planning, with neural patterns that approximate those

during the actual execution of the action.

1. Introduction

The hand is central in physical interactions with our environment. Typical hand-object interactions consist of sequential phases, starting with reaching and ending with object manipulation (Castiello, 2005). Electrophysiological studies in macaques have implicated a frontoparietal network in hand-object interactions (for a review see Gerbella et al., 2017). Previous functional MRI (fMRI) research (for a review see Errante et al., 2021) has identified a similar network in humans. This network comprises a dorsomedial pathway, consisting of the superior parietal occipital cortex (SPOC, corresponding to V6/V6A) and the dorsal premotor cortex (PMd), and a dorsolateral pathway, consisting of anterior intraparietal sulcus (aIPS) and ventral premotor cortex (PMv).

While past human neuroimaging studies revealed the neural substrates of grasping, most treated prehension as the end goal, not as a step towards meaningful hand-object interactions. In contrast, real-world hand-object interactions typically involve grasping only as a prelude to subsequent actions such as manipulating or moving objects. Importantly, previous studies have shown that the final action goal shapes prehension: Initial prehension strategies are affected by the goal in the 'end-state comfort' effect (Rosenbaum et al., 1990) and in actions like tool use (Comalli et al., 2016). Moreover, the end goal affects brain responses during action planning and prehension (Fogassi et al., 2005; Gallivan et al., 2016a).

The purpose of the current study was to investigate how isolated and sequential actions unfold differently on a moment-to-moment basis. The temporal unfolding of brain activation during actions has been largely overlooked (or only studied with EEG, without localizing the specific brain regions involved, e.g., Guo et al., 2019) . Most studies that have used multivoxel pattern analysis (MVPA) to investigate hand actions have averaged data within time bins for planning and execution (e.g., Gallivan et al., 2011). Notably, several studies have revealed the temporal unfolding of MVPA grasping representations for sequential timepoints (Ariani et al., 2018a; Gallivan et al., 2013). In addition, one study examined isolated (grasping an object) vs. sequential (grasping to move an object to one of two locations) actions, showing that activation patterns could be discriminated across the grasping network even during action planning (Gallivan et al., 2016).

Here we examined (univariate) activation levels and (multivariate) activation patterns (using representational similarity analysis, RSA (Kriegeskorte, 2008). Moreover, we also used a new methodological approach, temporal MVPA (tMVPA; Ramon et al., 2015; Vizioli et al., 2018) to examine the representation similarities across trials for the same and different points in time, separately for isolated vs. sequential actions.

We measured brain activation using fMRI while participants performed a motor task consisting

of either simple grasping of a dial (with two possible initial orientations) or grasping followed by rotation

of the dial (clockwise or counterclockwise), as one might turn an oven dial (See Figure 1). Based on

earlier findings that brain regions within the grasping network plan the full action sequence, and not just

the initial grasp (see Gallivan et al., 2016) , we expected that during the plan phase, as well as the

execute phase, tMVPA would reveal representations of the task (grasp vs. turn). Given that visual

orientation (Kamitani & Tong, 2005), surface/object orientation (Rice et al., 2007; Shikata et al., 2001;

Valyear et al., 2006) and grip orientation (Monaco et al., 2011) are represented in early visual cortex, the

caudal intraparietal sulcus (cIPS), and reach-selective cortex (SPOC), respectively, we predicted that the

representation of orientation would be biased to the start orientation early in planning, with a greater

emphasis on end orientation as the movement progressed. Our approach with tMVPA allowed us to

examine, across different regions, how representations unfolded over time for isolated vs. sequential

actions. Specifically, we expected that regions sensitive to the kinematics of executed actions (e.g., M1

and S1) would only show highly similar representations during action execution; whereas, regions

involved in more abstract features of action planning (e.g., aIPS for coding object shape) would show

similar representations across planning and execution.

2. Methods

2.1 Participants

Data from sixteen right-handed volunteers was utilized in the analysis (7 males, 9 females, mean

age: 24.4 years). Participants were recruited from Western University (London, Ontario, Canada) and

provided informed consent in accordance with procedures approved by the University's Health Sciences

Research Ethics Board. Data from an additional two subjects (one male and one female) was collected

but excluded due to excessive motion artifacts (see 2.6).

2.2 Setup and apparatus

The experimental setup is illustrated in Figure 1A-B. Participants lay supine in a 3-Tesla MRI

scanner with the head and head coil tilted approximately 30° to allow for direct viewing without mirrors

of a manipulandum positioned above the participant's hips. The manipulandum consisted of a black

rotatable dial (9-cm diameter; Figure 1B) with a yellow rectangular handle (5-cm length x 1-cm width x

2-cm depth). The dial was mounted on a black surface. The black surface was positioned such that the

dial was approximately perpendicular to the subject's line of gaze and comfortably within reach of their

right arm. Two yellow markers were put on the black surface (Figure 1B) indicating the start and end

positions for turning the dial. A grey platform (not shown in Figure 1A) was positioned above the

participant's lower torso serving as the home/resting position for the right arm between trials.

Participants' upper arms were braced above their torsos and just above their elbows (Figure 1A) to limit

movement of the shoulder, which can induce motion artifacts in fMRI signals. As such, participants

could only rely on elbow flexion/extension and forearm rotation to perform the experimental task.

Considering these constraints, the position and orientation of the dial were adjusted for each

participant to optimize participant comfort during task performance and ensure the dial remained fully

visible. The position of the yellow markers on the black surface was adjusted for each participant

individually so that dial rotation would not exceed 80% of the participant's maximum range of motion

when turning the dial clockwise or counterclockwise.

During the experiment, the dial was illuminated from the front by a bright yellow Light Emitting

Diode (LED) attached to flexible plastic stalks (Figure 1A; Loc-Line, Lockwood Products, Lake Oswego,

OR). A dim red LED (masked by a 0.1° aperture) was positioned approximately 15° of visual angle above

the rotatable dial and just behind it to provide a fixation point for participants (Figure 1A-B).

Experimental timing (see below) and lighting were controlled with in-house MATLAB scripts (The

MathWorks Inc., Natick, MA).

2.3 Experiment design and timing

Behavioral task. This experiment was a 2 (starting orientation: left or right) x 2 (action: grasp or

turn) delayed movement paradigm (Figure 1C). For each trial, the dial would appear in one of the two

yellow-marked starting positions. The grasp condition consisted of reaching towards the dial and

squeezing it between the middle phalanges of the index and middle fingers (trial end position shown in

Figure 1A; hand shape during grasp shown in Figure 1B). After grasp completion, participants returned

their arm back to the home position. In the turn condition, participants performed the same reach-to-

grasp action but would then subsequently rotate the dial clockwise or counterclockwise after they

grasped the dial in the left or right start position, respectively (Figure 1C). We decided on the index-

middle finger grip instead of a more natural variant of precision grasp as this would ensure the grip

would be highly similar in all conditions and limit changes to grip angle to optimize end-state comfort

(e.g., putting the right-hand thumb further up when planning to turn clockwise and further down when

planning to turn the dial counterclockwise). This avoids contaminating neural activation with low-level

sensorimotor confounds such as digit positioning. Participants were instructed to keep the timing of all

movements as similar as possible, such that the right hand reached from and returned to the home
position at the same time (see next paragraph 'Trial Design'). To isolate the visuomotor planning
response from the visual and motor execution responses, we used a slow event-related paradigm with
32-s trials, consisting of three phases: 'Presentation', 'Plan' and 'Execute' (see Figure 1D). We adapted
this paradigm from previous fMRI studies that successfully isolated delay-period activity from transient
neural responses following the onset of visual stimuli and movement execution (Beurze et al., 2007,
2009; Pertzov et al., 2011). Furthermore, using this paradigm in previous work from our group, we were
able to successfully isolate and decode planning-related neural activation prior to action execution
(Gallivan et al., 2011; Gallivan, McLean, et al., 2013).

Trial design. Before each trial, subjects were in complete darkness except for the fixation LED
upon which participants were instructed to maintain their gaze. The trial began with a 6-s (3 TRs of 2 s
each) Presentation phase in which the illumination LED lit up the rotatable dial. After the Presentation
phase, the 12-s (6-TR) Plan phase was initiated with a voice cue (0.5-s duration) saying either 'Grasp' or
'Turn' to instruct the upcoming action to the participant. Participants could see the object during the
presentation phase, thus perceiving the starting orientation of the yellow handle. However, they were
instructed to only begin the action after they received the 'Go' cue. After the Plan phase, a 2-s (1-TR)
Execute phase began with a 0.5-s beep (the 'Go' cue') cueing participants to initiate and execute the
instructed action. Performing the entire action, consisting of reaching and grasping (with or without dial
turning), took approximately 2 s. The dial remained illuminated for 2 s after the 'Go' cue allowing visual
feedback during action execution. After the 2-s Execution phase, the illumination LED was turned off,
cueing participants to let go of the dial and return their hand back to the home position. Participants
remained in this position for 12 s (6 TRs) in the dark (i.e., the intertrial interval) to allow the BOLD
response to return back to baseline before the next trial would be initiated.

Functional runs. Within each functional run, each of the four trial types (grasp: left or right –
turn: from left to right or right to left) was presented five times in a pseudo-randomized manner for a
total of 20 trials. Each participant performed eight functional runs, yielding a total of 40 trials/condition
(160 trials in total). Each functional run took between 10 and 11 minutes. For each participant, trial
orders were counterbalanced across all functional runs so that each trial type was preceded and
followed equally often by every trial type (including the same trial type) across the entire experiment.
During testing, the experimenter was positioned next to the set-up to ensure that the dial was in the
correct position before each trial, and manually adjust it if needed. In addition, the experimenter could
visually check whether the participant performed the trial correctly. Incorrectly trials were defined as

any trial where the participant did not perform the task as instructed: for instance, turning instead of
grasping, using incorrect grips, failing to turn in a smooth manner or having the dial slip from the fingers.

Participants performed a separate practice session before the actual experiment to familiarize
them with the motor task and ensure proper execution. The experimental session took approximately
three hours and consisted of preparation (i.e., informed consent, MRI safety, placing the participant in
the scanner and setting up the behavioral task), eight functional runs and one anatomical scan. The
anatomical scan was collected between the fourth and fifth functional runs to give participants a break
from the task.

2.4 MRI acquisition

Imaging was performed using a 3-Tesla Siemens TIM MAGNETOM Trio MRI scanner at the
Robarts Research Institute (London, ON, Canada). The T1-weighted anatomical image was collected
using an ADNI MPRAGE sequence (time to repetition (TR) = 2300 ms, time to echo (TE) = 2.98 ms, field of
view = 192 mm x 240 mm x 256 mm, matrix size = 192 x 240 x 256, flip angle = 9°, 1-mm isotropic
voxels). Functional MRI volumes sensitive to the blood oxygenation level-dependent (BOLD) signal were
collected using a T2*-weighted single-shot gradient-echo echo-planar imaging (EPI) acquisition sequence
(TR = 2000 ms, slice thickness = 3 mm, in-plane resolution = 3 mm x 3 mm, TE = 30 ms , field of view =
240 mm x 240 mm, matrix size = 80 x 80, flip angle = 90°, and acceleration factor (integrated parallel
acquisition technologies, iPAT) = 2 with generalized auto-calibrating partially parallel acquisitions
(GRAPPA) reconstruction. Each volume comprised 34 contiguous (i.e., with no gap) oblique slices
acquired at an approximate 30° caudal tilt with respect to the anterior-to-posterior commissure (ACPC)
plane, providing near whole brain coverage. We used a combination of parallel imaging coils to achieve
a good signal to noise ratio and to enable direct viewing of the rotatable dial without mirrors or
occlusion. Specifically, we placed the posterior half of the 12-channel receive-only head coil (6-channels)
beneath the head and tilted it at an angle of approximately 20°. To increase the head tilt to
approximately 30°, we put additional foam padding below the head. We then suspended a 4-channel
receive-only flex coil over the forehead (Figure 1A).

2.5 fMRI anatomical data processing

All fMRI preprocessing was performed in BrainVoyager version 22 (Brain Innovation, Maastricht
Netherlands). For the present study, we defined regions of interest (ROIs) in surface space instead of
volumetric space as it has been shown that cortical alignment improves group results by reducing

individual differences in sulcal locations (Fischl et al., 1999; Frost & Goebel, 2012). As such, we

performed surface reconstruction (mesh generation) and cortex-based alignment. Given that we relied

on the recommended approach and standard settings of BrainVoyager 22, these steps are explained

only in brief.

Folded mesh generation for each participant. In volumetric space we performed the following

steps. First, we corrected for intensity inhomogeneity. Then, we rotated the anatomical data in ACPC

space, due to the experimental head tilt, and normalized to Montreal Neurological Institute (MNI) space.

Next, we excluded the subcortical structures (labelled as white matter) and the cerebellum (removed

from anatomical) from mesh generation. We then defined the boundaries between white matter and

gray matter and between gray matter and cerebrospinal fluid. Finally, we created a folded mesh (surface

representation) of only the left hemisphere after removing topologically incorrect bridges (Kriegeskorte

& Goebel, 2001). We decided to only investigate the left hemisphere because the motor task involved

the right hand only, which previous work has shown predominantly activates the left (contralateral)

hemisphere (Cavina-Pratesi et al., 2010, 2018; Gallivan et al., 2011; Gallivan, McLean, et al., 2013).

Standardized folded mesh generation for each participant. Briefly, folded meshes created from

anatomical files often result in different numbers of vertices between participants. To facilitate cortex-

based alignment between participants, folded meshes were first transformed into high-resolution

standardized folded meshes. Each folded mesh was first morphed into a spherical representation by

smoothing (thus removing differences between sulci and gyri) and correcting for distortion. The

spherical representation of each participant mesh was then mapped to a high-resolution standard

sphere to create a high-resolution standardized spherical representation of the participant mesh. The

vertex position information of the original participant folded (not spherical) mesh was then used to

generate a standardized folded mesh for each participant.

Cortex-based alignment. Cortex-based alignment was performed following the approach of Frost

& Goebel (2012) and Goebel et al. (2006): We aligned all individual standard meshes to a dynamically

generated group average target mesh. Before aligning to the dynamic group average, we performed

pre-alignment (i.e., rigid sphere alignment). The actual steps of cortex-based alignment generate a

dynamic group average (a surface mesh based on all individual meshes) and sphere-to-sphere mapping

files for each participant that enable transporting the functional data from each individual to the

dynamic group average. Inverse sphere-to-sphere mapping files were also generated, which allows

transporting of data (such as regions of interest) from the dynamic group average back to individual

meshes.

2.6 fMRI functional data processing

General preprocessing. All functional runs were screened for motion and magnet artifacts by examining the movement time courses and motion plots created with the motion correction algorithms (three translation and three rotation parameters). Data from a run was discarded if translation exceeded 1 mm or rotation exceeded 1° between successive volumes. Based on this screening, all data from two participants and one run from an additional participant was discarded. Examination of the remaining data indicated negligible motion artifacts in time courses (as will be shown in later figures).

Functional runs were co-registered with the anatomical data using boundary-based registration (Greve & Fischl, 2009). We subsequently performed slice-scan time correction, motion correction and MNI normalization. Next, we performed linear-trend removal and temporal high-pass filtering (using a cut-off of three sine and cosine cycles on the fast Fourier transform of the time courses).

Volumetric to surface-based time courses. Volumetric time courses were first aligned with the respective participant's anatomical scan using boundary-based registration to ensure optimal alignment (Greve & Fischl, 2009). Volumetric time courses were then transformed to the respective participant's standardized folded mesh using depth integration along the vertex normal (sampling from -1 to 3 mm relative to the gray-white matter boundary). Once functional runs were transformed into surface space, they were spatially smoothed. Finally, functional runs in individual standard mesh space, could then be mapped onto the dynamic group average mesh using sphere-to-sphere mapping files that were generated during cortex-based alignment. This approach enabled us to model the averaged brain activation across all participants onto the group mesh and define ROIs based on hotspots of strongest activations in surface space.

2.7 Region of interest-based analysis

For the current project, we performed a ROI-based analysis of cortical regions within the hemisphere contralateral to the acting hand at the peak foci for task-related activity. To allow for better intersubject consistency and statistical power, we utilized cortex-based alignment for defining ROIs (Frost & Goebel, 2012; Goebel et al., 2006). We were primarily interested in cortical areas that constitute the cortical grasping and reaching networks that have been investigated in previous studies of our group (Cavina-Pratesi et al., 2018) and others (Ariani et al., 2018). As in previous projects (e.g., Gallivan et al., 2018; Fabbri et al., 2016), we considered only the contralateral hemisphere as it is well-established that hand-object actions are strongly lateralized and elicit only limited activation in the

ipsilateral hemisphere (Culham et al., 2006). Moreover, as our analysis of the left cerebral hemisphere

already included 17 regions and we performed multiple analyses, some at many time points, we wanted

to limit the complexity of the results and find a balance between statistical rigour and statistical power.

We chose an ROI approach rather than a whole-brain cortex-based searchlight to optimize

statistical power and computational processing time (Etzel et al., 2013; Frost & Goebel, 2012;

Kriegeskorte, 2008). Due to the large number of comparisons performed in a searchlight analysis, only

the most robust effects survive the correction for multiple comparisons (Kriegeskorte, 2008;

Kriegeskorte & Kievit, 2013). Moreover, tMVPA relies on single-trial correlations which are

computationally very intensive. Performing this analysis on the voxel-based searchlight level would likely

take multiple months.

<u>2.8 Defining regions of interest</u>

To localize our regions of interest on the group mesh, we applied a general linear model (GLM)

on our data in surface space. Predictors of interest were created from boxcar functions convolved with

the two-gamma haemodynamic response function (HRF). We aligned a boxcar function with the onset of

each phase of each trial within each run of each participant, with a height dependent upon the duration

of each phase. We used 3 TRs for the presentation phase, 6 TRs for the Plan phase and 1 TR for the

Execution phase. The six motion parameters (three translations and three rotations) were added as

predictors of no interest. Each incorrect trial was also assigned a unique predictor of no interest. All

regression coefficients (beta weights) were defined relative to the baseline activity during the intertrial

interval. In addition, time courses were converted to percent signal change before applying the random

effects GLM (RFX-GLM).

To specify ROIs for our analyses, we searched for brain areas, on the group level, involved in the

experimental task. We contrasted brain activation for all three phases with baseline. This contrast was

performed across all conditions to ensure that ROI selection based on local activation patterns was not

biased by differences between specific conditions (e.g., grasp versus turn). Specifically, the contrast was:

{Presentation [Grasp $_{left}$ + Grasp $_{right}$ + Turn $_{left\ to\ right}$ + Turn $_{right\ to\ left}$] + Plan [Grasp $_{left}$ + Grasp $_{right}$ + Turn $_{left\ to}$

$_{right}$ + Turn $_{right\ to\ left}$] + Execute [Grasp $_{left}$ + Grasp $_{right}$ + Turn $_{left\ to\ right}$ + Turn $_{right\ to\ left}$]} > baseline. This

contrast enabled us to identify regions that showed visual and/or motor activation associated with the

task.

In BrainVoyager, area selection on a surface generates a hexagon surrounding the selected

vertex. We decided on an area selection size of 100 (arbitrary units) as this value provided a good

balance between inclusion of a sufficient number of vertices/voxels around each hotspot for
multivariate analyses and avoiding overlap between ROIs (especially in the parietal lobe).

<u>2. 9 Regions of interest</u>

Selected regions of interest (Figure 2) were defined in the left hemisphere. Most regions of interest
were defined using the contrast above (all conditions > baseline), with two exceptions, as detailed
below.

First, the involvement of the cortical grasping network (Gerbella et al., 2017) was investigated by
including sensorimotor and visuomotor regions (Ariani et al., 2018; Cavina-Pratesi et al., 2018; Fabbri et
al., 2016; Gallivan et al., 2011; Gallivan, McLean, et al., 2013).

- Primary motor cortex (M1): Hotspot of strongest activation near the 'hand knob' landmark (Gallivan et al., 2013; Yousry et al., 1997).

- Dorsal premotor cortex (PMd): Hotspot of strongest activation near the junction of the precentral sulcus and the superior frontal sulcus (Picard & Strick, 2001; Pilacinski et al., 2018).

- Ventral premotor cortex (PMv): Hotspot of strongest activation inferior and posterior to the junction of the inferior frontal sulcus) (Tomassini et al., 2007).

- Primary somatosensory cortex (S1): Hotspot of strongest activation anterior to the anterior intraparietal sulcus, encompassing the postcentral gyrus and sulcus (Gallivan et al., 2011; Gallivan, McLean, et al., 2013)

- Anterior intraparietal sulcus (aIPS): Hotspot of strongest activation directly at the junction of the intraparietal sulcus and the postcentral sulcus (Culham et al., 2003).

- Caudal intraparietal sulcus (cIPS): Hotspot of strongest activation on the lateral side of the brain, anterior and superior to the junction between the intraparietal sulcus and the posterior occipital sulcus (Beurze et al., 2009; Grefkes & Fink, 2005). Comparisons with the Julich atlas (Richter et al., 2019) suggested overlap with hIP7 and/or hP01.

- Anterior superior parietal occipital sulcus (aSPOC): Hotspot of strongest activation on the medial side of the brain, anterior and superior to the parietal-occipital sulcus (Cavina-Pratesi et al., 2010), thought to correspond to area V6A(Pitzalis et al., 2015).

- Posterior superior parietal occipital sulcus (pSPOC): Hotspot of strongest activation on the medial side of the brain, posterior and inferior to the parietal-occipital sulcus (Cavina-Pratesi et al., 2010). thought to correspond to area V6 (Pitzalis et al., 2015).

Second, the following medial and frontal regions were selected due to their involvement in motor

planning and decision making (Ariani et al., 2015; Badre & Nee, 2018; Cavina-Pratesi et al., 2018).

• Dorsolateral prefrontal cortex (DLPFC): Hotspot of strongest activation near the middle

frontal gyrus (Mylius et al., 2013).

• Supplementary motor area (SMA): hotspot of strongest activation adjacent to the medial

end of the cingulate sulcus and posterior to the plane of the anterior commissure (Picard &

Strick, 2001).

• Pre-supplementary motor area (pre-SMA): Hotspot of strongest activation superior to the

cingulate sulcus, anterior to the plane of the anterior commissure and anterior and inferior

to the hotspot of strongest activation selected for SMA (Picard & Strick, 2001).

Third, we included the superior parietal lobule (SPL) due to the extensive activation evoked by our task

and in hand actions more generally (Ariani et al., 2018a; Cavina-Pratesi et al., 2018). We defined the SPL

as the area on the lateral/superior side of the brain that is bordered anteriorly by the postcentral sulcus,

inferiorly by the intraparietal sulcus and posteriorly by the parietal-occipital sulcus (Scheperjans, et al.,

2008a; Scheperjans et al., 2008b). Due to the large swathe of activation evoked by our contrast, we

defined four ROIs within the SPL based on their relative position to each other (to ensure minimal

overlap) and the anatomical landmarks bordering the SPL. We decided on four ROIs as well as their

names based on Scheperjans et al. (2008a).

• 7PC: Hotspot of strongest activation located on the posterior wall of the postcentral sulcus and

superior to the intraparietal sulcus. Given we defined post-aIPS as well, 7PC was also defined as

superior to post-AIPS.

• 5L: Hotspot of strongest activation located just posterior to the postcentral sulcus and superior

to area 7PC.

• 7P: Hotspot of strongest activation superior to the intraparietal sulcus and anterior to the

parietal-occipital sulcus.

• 7A: Hotspot of strongest activation in the postcentral gyrus, superior to 7PC, posterior to 5L and

anterior to 7P.

Finally, two visual regions were selected from a probabilistic functional atlas that utilized cortex-based

alignment (Rosenke et al., 2021), to which we aligned our participant surface meshes.

• Dorsal primary visual cortex (V1d): Given that primary visual cortex (V1) responds to visual

orientation (Kamitani & Tong, 2005b) we wanted to investigate its response here. Because our

target objects (and participants hands) fell within the lower visual field (below the fixation

point), they would stimulate the dorsal divisions of early visual areas (Wandell et al., 2007). As

such, we investigated only the dorsal division of V1, V1d, using the Rosenke atlas.

•   Extrastriate body area (EBA): We were also interested in examining the response of the

extrastriate body area, which has been implicated not only in the visual perception of bodies

(Greenfield et al., 1996) but also in computing goals during action planning (Astafiev et al., 2004;

Zimmermann et al., 2016). Because the EBA is not easy to distinguish from nearby regions of the

lateral occipitotemporal cortex, we utilized the EBA from the Rosenke atlas. Briefly, this process

is nearly identical for the other ROIs. The only difference was that we aligned all individual

standard meshes to the template of Rosenke (instead of the dynamic group average). This

provided sphere-to-sphere and inverse sphere-to-sphere mapping files. The latter could then be

used to transport EBA from the Rosenke template to the individual standard meshes and

perform all other steps identically as for the other ROIs. We defined EBA as the combination of

all body-selective patches, being OTS-bodies, MTG-bodies, LOS-bodies and ITG-bodies (for the

full explanation of these regions see Rosenke et al., 2021)

After ROIs were defined in group surface space, they were transformed to individual surface space,

using the inverse transformation files generated during cortex-based alignment and then to individual

volumetric MNI space, using depth expansion (inverse of depth integration; see 'fMRI functional data

processing: Volumetric to surface-based time courses') along the vertex normals (-1 to 3 mm). This

approach allowed us to define ROIs on the group surface but extract functional data from the individual

volumetric level.

<u>2.10 Analysis of functional data</u>

Functional data were extracted to perform deconvolution general linear models (deconvolution

GLMs; Hinrichs et al., 2000), representational similarity analysis (Kriegeskorte, 2008) and temporal

multivoxel pattern analysis (tMVPA; Vizioli et al., 2018). All analyses described below, excluding initial

processing of fMRI data and the univariate analyses (which where both done in BrainVoyager), were

done with in-house MATLAB scripts.

2.10.1 Univariate analysis.

We used a random-effects GLM with deconvolution to extract the time courses of activation in

each ROI during the experimental task. For each experimental condition, we used 15 matchstick

predictors (timepoint 0 to 14), the first predictor ("predictor 1") was aligned with time point zero (Figure

1D; "light on") and the last predictor ("predictor 15") was aligned with the drop in signal at the end of

the trial ("timepoint 14). Due to this clear drop in signal at timepoint 14, we did not include a predictor

(non-existing "predictor 16") for the last timepoint (existing timepoint 15; see Figure 1). For each

functional voxel in each ROI, baseline z-normalized estimates of the deconvolved BOLD response,

representing the mean-centered signal for each voxel and condition relative to the standard deviation of

signal fluctuations, were extracted for all conditions. Voxel activation was then averaged across voxels

within each ROI. This provided us with an estimate of the averaged time course of brain activation for

each condition for each ROI and for each participant.

Given previous work from our group demonstrating the involvement of our selected ROIs in

hand-object interactions (e.g., Fabbri et al., 2016; Gallivan et al., 2009; Monaco et al., 2011), we

included univariate time courses for comparison as they indicate changes in the signal and reveal the

degree to which differences in activation levels are present (or not). The univariate time courses also

demonstrate that our quality-assurance procedures, particularly motion screening, succeeded in

generating clean data.

Time courses were generated for each ROI (as shown in Figure 2) separately for each of the four

conditions. However, the differences in univariate signals between conditions were small or negligible.

Thus, to simply evaluate whether or not there was significant activation during each of the three phases

of the trial, we performed statistical comparisons of the average activation levels (collapsed across the

four conditions) compared to the intertrial baseline (time 0).

First, for each ROI in each participant we averaged the data across conditions. Second, to reduce

the number of statistical comparisons while targeting peak responses, we defined three distinct phases

by averaging two datapoints for each. These phases were response (i) to presentation, by averaging

timepoint 2 and 3, (ii) to planning, by averaging timepoint 6 and 7, (iii) and to execution by averaging

timepoint 11 and 12. Third, we investigated significant changes compared to baseline (i.e., > 0) using

one-sample t tests. Accordingly, per ROI we performed three one-sample t tests. Fourth, we corrected

for multiple comparisons. It is well-known that balancing type I and type II errors in fMRI research can

be challenging due to the amount of data and number of comparisons involved. Although data between

ROIs is independent, the number of comparisons in total across ROIs increase the likelihood of Type I

errors. Because of this predicament, we decided to correct the univariate analysis for multiple

comparisons using the Bonferroni method in two manners being (a) only including comparisons within

ROIs (i.e., 3 comparisons within each ROI thus $alpha$ = 0.05/3) and (b) including all comparisons (i.e., 3

comparisons within each of the 17 ROIs thus $alpha$ = 0.05/51). The results can be found in Figure 2: a

small asterisk indicates that the values are significant against the standard correction (3 comparisons), a

larger bold asterisk indicates that the values are significant against the strict correction (51

comparisons). The values in the results reflect mean ± SEM. *P* values will be referred to as *P* and *P*(strict)

for the 3-comparison and 51-comparison corrections, respectively. Note that we provide the *p* values

corrected for the comparisons. For instance, if we found a p-value of 0.0035, it would be reported as P =

0.0105 (corrected for 3 comparisons) and P(strict) = 0.1785 (corrected for 51 comparisons).

2.10.2 Temporal multivoxel pattern analysis.

For tMVPA we relied on the approach developed in Ramon et al. (2015) and Vizioli et al. (2018).

Briefly, tMVPA has been developed to investigate the temporal development of neural representations

by relying on multivariate analyses with a trial wise approach. The tMVPA methods are schematized in

Figure 3. For each voxel of each ROI of each participant, we performed a deconvolution GLM for every

trial separately, which was then transformed into BOLD percent signal change by dividing the raw BOLD

time course by its mean. Note that for tMVPA we performed deconvolution GLMs for each trial (of the

same condition) separately whereas for the univariate analysis and RSA (see below) we ran

deconvolution GLMs for each condition (i.e., across same-condition trials). After computing the single-

trial activation patterns, we computed single-trial representational dissimilarity matrices (stRDMs;

dissimilarity = 1 - *r*) using Pearson correlations for each condition of each ROI of each participant. As

shown in Figure 3A, a stRDM is generated by computing dissimilarity between the activation patterns of

voxels at each timepoint of a given trial ('trial m') with the activation pattern of the same voxels for each

of the timepoints for another trial ('trial n'). This process is iteratively repeated until all unique within-

condition trial pairings are run. stRDMs were then averaged across the main diagonal to yield a

diagonally symmetric matrix. That is, the data matrix was averaged position-wise with its transpose

matrix. We performed this averaging as we were only interested in between-condition differences and

not in differences between same-condition trials.

To further clarify calculation of the dissimilarity metric: in Figure 3B and Figure 3C, the green

highlighted square indicates the dissimilarity between all the voxels (of a given ROI) at timepoint 2 of

Turn Left to Right trial m with the activation of the same voxels at timepoint 2 of Turn Left to Right trial

n. The magenta highlighted squares indicate the averaged dissimilarity between timepoints 2 and 10 of

trial m and n. As explained before, we averaged the dissimilarity between timepoint 2 of trial m and

timepoint 10 of trial n (magenta highlighted square below diagonal) with the dissimilarity between

timepoint 10 of trial m and timepoint 2 of trial n (magenta highlighted square above diagonal). Finally,

the blue highlighted square indicates the dissimilarity between timepoints 10 of trial m and n. In sum,

values on the main diagonal show dissimilarity between within-condition trials at the same timepoint.

Values that are off the main diagonal show the dissimilarity between within-condition trials at different

(i.e., earlier or later) timepoints.

stRDMs were calculated for the four conditions separately, Fisher z-transformed and then

averaged (10 % trimmed mean; Vizioli et al., 2018) within conditions (e.g., average RDM for grasp left).

Finally, the averaged RDMs were then averaged across orientations (e.g., averaging of RDMs between

left and right start orientations to produce average Grasp and Turn RDMs; Figure 3D and 3E

respectively). This was done as we were primarily interested in investigating differences in neural

representations between singular (grasping) and sequential actions (grasping then turning) and we did

not expect that these differences would depend upon the start orientation. As such, this approach

resulted in two RDMs for each participant.

In line Vizioli et al. (2018), we performed statistical analysis on the Fisher z-transformed data;

however, we used the non-transformed data for visualization purposes to render the values visually

more interpretable. To test for statistically significant differences between the grasp and turn RDMs, we

subtracted for each participant the turn RDM from the grasp RDM resulting in a subtraction matrix (Turn

versus Grasp: Figure 3F) and investigated where the subtraction differed significantly from zero. As such,

a given value in the subtraction matrix that is positive indicates that grasping trials are more dissimilar

than turning trials at that given timepoint. Conversely, a value that is negative indicates that turning

trials are more dissimilar than grasping trials at that given timepoint. Note that we decided to

statistically test whether each cell differed from zero instead of the increasingly larger sliding window

analysis used in Ramon et al. (2015) and Vizioli et al. (2018). Our rationale was that the earlier studies

relied on a visual task (face recognition) whereas our study relied on a motor task. Arguably, during

motor preparation/execution, neural representations might evolve differently than purely visual

responses. As such, we argued that statistically testing each cell separately might reveal more

information on the temporal evolution of motor execution as, for instance, transitions between

activation patterns between planning and execution might be brief or abrupt. Note that because we had

averaged values across the main diagonal when calculating the dissimilarity matrices (Figure 3 and 5),

these matrices were symmetrical reflections across the diagonal. However, we performed statistical

analyses only on the values on and above the main diagonal of the Turn versus Grasp subtraction

matrices (Figure 3F).

To test for statistical significance, we performed (1 - alpha) bootstrap confidence-interval

analysis (critical alpha = 0.05) by sampling participants with replacement 500 times. As described above,

we investigated whether tMVPA differed significantly from zero in either direction. Importantly, we

excluded the first two timepoints (timepoints 0 and 1) and the last one (timepoint 14) from the

statistical analyses (resulting in 12 timepoints included in the analysis: timepoints 2 to 13) because no

differences were expected before the BOLD response to emerge at the start, because later timepoints

reflect the post-stimulus undershoot phase of the BOLD response, and to reduce the number of

comparisons requiring Bonferroni correction for multiple comparisons. See Figure 1 for the trial timing

and Figure 3 for the timepoints in the tMVPA analysis).

As for the univariate analysis, we accounted for multiple comparisons using Bonferroni

correction in two manners. Our first, standard, correction did not take multiple comparisons between

regions in account but only the number of comparisons within an ROI. This resulted in correcting for 78

tests: the binomial coefficient of 12 timepoints (which excludes the diagonal) plus the 12 included

timepoints on the diagonal. The second, strict, correction, included multiple comparisons between all

regions at all time points and resulted in 936 tests. Accordingly, the critical alpha was Bonferroni

corrected for the number of tests in the bootstrap confidence-interval analysis. Due to the

impracticability of reporting our tMVPA data in text or table format, we provide these results only in

figure format (see data availability statement). In Figure 6, cells that differ significantly from zero have a

dashed outline or solid outline for the standard and strict correction, respectively.

2.10.3 Hierarchical clustering.

The tMVPA matrices (stRDMs) for Grasp and Turn revealed that different ROIs showed different

temporal profiles. For example, some regions showed similar activation patterns throughout the entire

trial; whereas, other regions showed similar patterns only during motor execution (See Figure 5). To

make it easier to group these different ROI timing patterns for display and discussion, we used

hierarchical clustering. To do so, we averaged within-participants the grasp and turn RDMs to generate

one RDM per participant. Next, we calculated for each participant the Spearman correlations between

all ROIs using the averaged RDMs. The ROI correlation matrix for each participant was then transformed

into a dissimilarity measure (1-r). Finally, hierarchical clustering was then performed using the ROI

dissimilarity matrix of all participants.

2.10.4 *Representational similarity analysis.*

We used RSA to examine the degree to which the pattern of activation across voxels within each

ROI at each time point represented the start orientation, the end (goal) orientation, and the task, as

illustrated in Figure 4A-C. For RSA, we relied on the methods described in Kriegeskorte (2008). By

examining the degree to which different conditions evoke a similar pattern of brain activation within a

ROI, the nature of neural coding (or representational geometry; Kriegeskorte & Kievit, 2013) can be

assessed.

For each voxel of each ROI of each participant, we performed a deconvolution GLM. This was

done for each condition and each run (but not trial as for tMVPA) separately, providing the average

development of brain activation over time for each condition in each voxel in each ROI of each run of

each participant. Then for each voxel separately, activations of the four conditions were normalized by

subtracting the grand mean (i.e., average across conditions) from the value of each condition in that

voxel (Haxby et al., 2001). Note that this was done for each run separately. These steps resulted in the

normalized brain activation over time averaged across trials of the same conditions within one run. This

was done for each condition for each voxel of each ROI for each run of each participant.

Subsequently, we computed representational dissimilarity matrices (RDMs) within each ROI.

Pearson correlations were calculated at each timepoint between all voxels of a given ROI, i.e., "within

timepoint correlations", and then transformed into dissimilarity metric ($1-r$). This was done within-

condition (e.g., grasp left activations at timepoint 1 correlated with itself at timepoint 1) and between

conditions (e.g., grasp left activations at timepoint 1 correlated with turn left to right at timepoint 1). In

this manner, RDMs can quantify the dissimilarity in activation patterns between conditions, i.e. "how

well is activity at any timepoint in a given set of voxels in a given condition correlated with the activity at

the same timepoint of the same voxels during a different condition.". Please note that we did not

perform autocorrelations (e.g., correlate grasp left activations during run 1 with itself thus leading to

correlation values of 1). Instead to test within-subject reliability of the RDM calculations, we used cross-

validation by splitting data into all potential combinations of runs yielding two sets of four runs each

(two examples: split 1 = correlate all even runs with the uneven runs; split 2: correlate the first four runs

with the last four runs). Split data provides data-based estimates of dissimilarity even for the same

condition as no autocorrelations are performed whereas, in unsplit data, the dissimilarity is necessarily

zero. Data for all cells in the RDM is necessary for RSA on factorial designs to ensure that contrasts are

balanced across orthogonal factors. Finally, the RDMs were Fisher transformed to have a similarity

metric with a Gaussian distribution.

To test whether each region contained information about condition differences, we measured
correlations between the RDM in each region (Figure 4D-E) and three separate models that capture
orthogonal components of the experimental task (Figure 4A-C). This was done for each timepoint
separately (15 timepoints included in the GLM being timepoint 0-14; time-resolved decoding). Note that
in contrast to the tMVPA we included all timepoints due to our selection of models which included "two
visual models" which may decode already during the earliest timepoints. In total, we used three models
that capture specific task attributes: (A) Start orientation, irrespective of task (i.e., grasp or turning) — a
timepoint with an RDM that correlates with this model would indicate an encoding of the initial
orientation of the handle (i.e., left, or right), (B) end orientation, irrespective of the performed action
(i.e., grasp or turning) — a timepoint with an RDM that correlates with this model would indicate an
encoding of the final orientation of the handle (i.e., left, or right) after task execution, (C) motor task,
irrespective of the initial/final handle's orientation (i.e., left, or right) — a timepoint with an RDM that
correlates with this model would indicate an encoding of the task goal (i.e., grasping or turning). The
metric was calculated by computing the Spearman correlation (ρ) between the Fisher-transformed split
RDMs and each model for each ROI for each participant. For each ROI we also calculated the upper and
lower bound of the noise ceiling. Briefly, the noise ceiling is the expected RDM correlation achieved by
the (unknown) true model, given the noise of the data and provides an estimate of the maximum
correlations that could be expected for a given model in a given ROI (Kriegeskorte, 2008). RDMs were
first rank transformed, following the original z-transformation. The upper bound of the noise ceiling,
considered an overestimate of the maximum correlation, was calculated as iteratively correlating one
participant's RDM with the average RDM of all participants (thus including the given participant) and
then averaging across all participants. The lower bound of the noise ceiling, considered an
underestimate of the maximum correlation, was calculated as iteratively correlating between one
participant's RDM with the average RDM of all other participants (thus excluding the given participant),
and then averaging across participants.

We used one-way Student's t tests (alpha < 0.05) to assess whether model correlations were
significantly greater than zero for each timepoint in each ROI. Importantly, as for the tMVPA we decided
to exclude the first two timepoints (timepoints 0 and 1) and the last one (timepoint 15). This resulted in
13 timepoints included in the analysis. Note that we did not compare whether models outperformed
each other as we were primarily interested in whether which ROI embodied what type of information.
To be in line with previous studies combining motor tasks and RSA (e.g., Di Bono et al., 2015; Gallivan et
al., 2013; Monaco et al., 2020), we performed corrections for multiple comparisons using the false

discovery rate. As for the univariate analysis, we performed multiple comparisons for each ROI

separately (i.e., 12 timepoints x 3 models) and in a stricter manner across all regions (strict comparison:

12 timepoints x 3 models x 17 ROIs). The results can be found in Figure 6: models at any given timepoint

that are significantly larger than zero for only the within-ROI corrections are indicated with a "plus sign".

When the model at the timepoint also reaches significance for the strict threshold, this is indicated by an

asterisk instead. Due to the amount of data (i.e., 12 timepoints x 3 models x 17 ROIs), we provide a

specific selection of values in the results section. Akin to the univariate analysis, we provide data for

timepoints 2 and 3 for the presentation phase, 6 and 7 for the planning phase and 11 and 12 for the

execution phase in Table 2 for the subset of ROIs in Figure 6. Again, note that we performed statistical

tests and performed multiple comparisons for 12 timepoints and not just the six timepoints mentioned

here to keep the statistical analysis in line with our tMVPA approach. In addition, note that here we do

not average timepoints for each phase (e.g., no averaging of timepoint 2 and 3 into one "presentation

phase") for the same reasons. The values in Table 2 reflect the q values resulting from the strict false

discovery rate correction as well as the confidence intervals, given as [lower threshold mean upper

threshold], which are Bonferroni corrected for multiple comparisons in the strict manner (12 timepoints

x 3 models x 17 ROIs). Note that we had to correct confidence intervals using Bonferroni given the step-

by-step approach FDR, which is not common for confidence intervals. As such, it is possible that our

actual interpretation of the statistical analysis using FDR (shown in Figure 6) reveals significant effects

despite the confidence intervals using Bonferroni (shown in Table 2) containing zero.

To end, we will also qualitatively discuss the grasp and turn RDMs (which subtraction constitutes

the tMVPA subtraction matrix that is statistically tested) in order to facilitate conceptual understanding

of the data. Please note that this qualitative discussion will not be incorporated in the results and

discussion due to the lack of the statistical analysis (which has been done on the subtraction matrix

instead).

3. Results

Qualitative examination of the deconvolution time courses (Figure 2) and the tMVPA results

indicated that different regions had different temporal profiles of activity. As shown in Figure 5,

hierarchical clustering of the RDMs averaged across Grasp and Turn facilitated conceptual grouping of

the ROIs for further investigation.

Notably, ROIs differed in the time ranges over which they showed reliable activation patterns

across trials of the same type. Most strikingly, V1d and cIPS showed strong temporal similarity in

activity patterns throughout the Plan and Execute phases of the trial, as indicated by the large block of

high similarity (dark red) beginning early in planning and continuing through late execution (Figure 5).

That is, in these regions, voxel activation patterns were not only similar during the same phase (along

the diagonal cells) but were also similar across timepoints throughout planning and execution (off-

diagonal cells in the planning and execution phases). This indicates consistency in the neural

representation throughout the trials. Similar but weaker similarity patterns were also observed in other

regions, particularly aSPOC, pSPOC, 7P, EBA, 7A , 7PC, and 5L.

In contrast, other regions -- aIPS, M1, S1, pre-SMA, SMA, PMd -- showed trial-consistent

activation patterns predominantly for the peak execution period (dark red blocks in the execution

phase; i.e., in lower right corners of Grasp and Turn matrices). In some cases, there was also some

consistency of patterns between the peak execution phase and earlier timepoints during Plan and

Execute of which M1 is a clear example in Figure 5 (off-diagonal cells between planning and execution

phase, as indicated by the reddish "wings" above and left of the peak similarity). PMv and DLPFC

showed only weak consistency of patterns across trials, highest in the peak of execution.

Interesting patterns were also observed in the tMVPA difference in trial-by-trial consistency

between turn and grasp (final column of Figure 5). Notably many of the regions showed higher

consistency during Turn actions than Grasp actions.

Based on these observations from Figure 5, we focussed subsequent analysis and interpretation

on a subset of ROIs that is shown in Figure 6.

3.1 Visual regions of V1d and cIPS

Univariate analysis. The first column in Figure 6 shows the deconvolution time courses for V1d

and cIPS. V1d and cIPS show a steep increase in average voxel activation following object presentation

(V1d = 0.31 ± 0.06, P < 0.001,P(strict) = 0.008; cIPS = 0.46 ± 0.07, P < 0.001; P(strict) < 0.006 compared to

baseline. This initial increase is maintained throughout the planning (V1d = 1.31 ± 0.19, P <

0.001,P(strict) < 0.001; cIPS = 1.09 ± 0.19, P < 0.001; P(strict) < 0.001) and execution phases (V1d = 1.07

± 0.22, P < 0.001,P(strict) = 0.013; cIPS = 1.15 ± 0.18, P < 0.001; P(strict) < 0.001).

Qualitative (non-statistical) assessment of the condition RDMs. In Figure 6 the second and third

columns show the condition RDMs for grasp and turn, respectively. For both regions, activation pattern

similarity within conditions strongly increases following object presentation. Due to the sluggish nature

of the hemodynamic response and our findings for the univariate analysis, it is plausible that this early

increase of similarity in the activation patterns is primarily driven by a visual response. Notably, the cIPS

within-condition similarity in activation patterns increases towards the peak execution (approx.

timepoint 11-13).

Temporal multivoxel pattern analysis. The fourth column in Figure 6 depicts the subtraction Turn

minus grasp on which we performed statistical analyses as previously described. For V1d, tMVPA is

unable to discriminate the grasp and turning condition for matched timepoints (along the diagonal)

during motor planning. That is, subtracting the grasp dissimilarity from the turn dissimilarity (second and

third column in Figure 6) does not result in a difference that differs statistically from zero. Interestingly,

tMVPA is able to discriminate the activation patterns of grasping and turning during late execution (i.e.,

the "plus sign" of significant squares in the grasp versus turn RDM for V1d) which indicates that V1d may

encode change in visual information (e.g., increased motion during) or motor-components that are non-

visual. Interestingly, tMVPA was also able to discriminate activity patterns related to grasping versus

turning when dissimilarity calculations were done between timepoints of the late planning phase and

timepoints of the late execution phase (i.e., the three adjacent significant squares in the grasp versus

turn RDM for V1d). These findings suggest that the activation patterns in V1d become tuned towards

either grasping or turning during late motor planning and may not be limited to solely encoding visual

information. Similar but stronger effects were found for cIPS. tMVPA was able to discriminate cIPS

activation patterns during most of the execution phase as well as when dissimilarity calculations were

done between timepoints of the execution phase and early and late timepoints of the planning phase

(i.e., the vertical and horizontal "wings" of significant squares above and left of the execution phase).

This suggests that in cIPS, representations of the task emerge early in the planning period.

Representational similarity analysis. For the RSA, we tested how well multivariate activation

patterns fit three models for the representation of start orientation, end orientation, and task (Turn

versus Grasp) over time (Figure 6; last column). The confidence intervals and *q*-values for a selection of

timepoints (Figure 6; last column; timepoints within transparent rectangles with dashed grey outlines) is

provided in Table 2. Notably, although all three types of information appear to be represented in both

V1d and cIPS during action execution, starting orientation appeared to be represented during the early

planning phase. In addition, during the late planning phase, both starting orientation and grasp versus

turn also appeared to be represented in V1d (see Table 2 for values for Timepoints 6 and 7).Although

the decoding of start orientation during planning and end orientation during execution may reflect the

processing of simple visual orientation information (Kamitani & Tong, 2005), coding of both start and

end orientation overlapped during execution (Figure 6; last column; V1d and cIPS; green and red trace),

suggest a much more complex representation. Notably, V1d, represented the task during planning,

before any action had been initiated, perhaps due to anticipation of the visual consequences of the

upcoming action.

In sum, while our findings are consistent with a role for V1d and cIPS as predominantly visual

regions, the similarity in activation patterns (tMVPA) between the planning and the execution phase, in

particular for cIPS (Figure 6; fourth column) as well as V1d coding the motor task during planning (Figure

6; last column' blue trace) suggest that these regions may encode motor components already during the

planning phase.

<u>3.2 Regions of the SPL: 7A, 7PC, 5L and 7P</u>

Univariate analysis. Despite only 7A showing a significant increase following object presentation

(area 7P = 0.13 ± 0.05, P = 0.09, P(strict) = 1.00; area 7A = 0.16 ± 0.05, P = 0.037, P(strict) = 0.63 ; area

7PC = 0.11 ± 0.06, P = 0.36, P(strict) = 1.00 ; area 5L = 0.15 ± 0.05, 0.040. P(strict) = 0.679), all regions

showed a significant increase following action planning (area 7P = 0.40 ± 0.08, P < 0.001, P(strict) <

0.001; area 7A = 0.42 ± 0.08, P < 0.001, P(strict) = 0.01 ; area 7PC = 0.25 ± 0.07, P = 0.006, P(strict) =

0.103; area 5L = 0.28 ± 0.06, P = 0.001, P(strict) = 0.017), and execution (area 7P = 0.36 ± 0.07, P < 0.001,

P(strict) = 0.009; area 7A = 0.49 ± 0.09, P < 0.001, P(strict) = 0.003 ; area 7PC = 0.30 ± 0.08, P = 0.006,

P(strict) = 0.104; area 5L = 0.54 ± 0.09, P < 0.001, P(strict) = 0.002). As shown in the first column in Figure

6, given that the activity within this ROIs show a different trend than cIPS and V1d (i.e., primarily a

significant increase in activation following action planning, not object presentation), it seems plausible

that the subregions of the SPL have primarily a visuomotor involvement.

Qualitative (non-statistical) assessment of the condition RDMs. The condition RDMs of Figure 6

show that the ROIs within the SPL show activity patterns that are similar to each other. First, similarity

between trials within each condition increases following object presentation, in particular, during early

planning. Second, similarity in activation patterns is then maintained until a second increase during the

execution phase. Notably, this pattern (in the condition RDMs of the SPL regions) is similar to that of

V1d and cIPS, although it seems weaker (i.e., less red). Most importantly, the increase in within-

condition trial similarity following object presentation seems weaker for the SPL regions than for V1d

and cIPS suggesting that these regions are likely less modulated by this type of motor task.

Temporal multivoxel pattern analysis. In line with the assessments of the univariate analysis and

the condition RDMs, tMVPA indicate that the regions of the SPL may be involved in the motor task, in

particularly during motor execution (Figure 6; column 4). For area 7A, tMVPA was able to discriminate

activation patterns for grasping and turning during the later execution phase. Interestingly, we found

similar results for area 7PC as for cIPS (as explained in "Visual regions of V1d and cIPS"). tMVPA was able

to discriminate grasping and turning throughout both earlier and later stages of the execution phase as

well as when dissimilarity calculations were done between timepoints of the planning phase and

timepoints of the execution phase. These findings indicate that area 7PC might anticipate turning

actions already during the planning phase. The results for area 5L are somewhat similar to the findings

for areas 7A and 7PC. That is, tMVPA is primarily able to differentiate between grasping and turning

during the later execution phase but also, in a more limited manner than area 7PC, when dissimilarity

was calculated on timepoints of the early/late planning phase and timepoints of the execution phase.

For area 7P, tMVPA could significantly discriminate activation patterns associated with grasping and

turning only during the execution phase. Finally, it can be seen that tMVPA can discriminate between

grasping and turning by relying on the activation patterns of each of the four regions. However, this

ability to discriminate the task is the weakest in 7P, stronger in 7A and 5L and the strongest in 7PC.

Representational similarity analysis. The RSA results (Figure 6 column 5) indicate that the

regions of the SPL could rather have a visuomotor involvement than a purely visual one: starting

orientation cannot be decoded from the SPL (Figure 6; last column; 7P, 7A, 7PC and 5L; green trace)

during the planning phase. In support of our tMVPA results, grasping versus turning can be decoded

from all four regions in the SPL (Figure 6; last column; 7P, 7A, 7PC and 5L; blue trace). RSA is primarily

able to do this when relying on the execution phase. Significant decoding is also observed during the late

planning phase for the regions 7A and 7P (see Table 2 for values for timepoints 6 and 7). Interestingly, in

area 7PC only turning versus grasping can be decoded, suggesting that this region might be mainly

involved in rotating the wrist, irrespective of the direction, and may not integrate visual information.

Indeed, from the other regions (areas 7A, 5L and 7P) both start and end orientation can be decoded

during the action execution phase suggesting that these regions may encode specific wrist orientations,

or may still be involved in visuomotor integration.

In sum, both tMVPA and RSA seem to provide evidence for the involvement of the SPL in

decoding sequential actions. That is, sequential (grasping then turning) actions can be differentiated

from singular grasping actions based on the activation patterns from the regions 7A, 7PC, 5L and 7P. In

particular, for areas 7A and 7P our findings highlight that RSA is already able to do so during motor

planning.

3.3 EBA and aIPS

Univariate analysis. As shown in Figure 6, EBA shows an initial response to object presentation

(EBA = 0.19 ± 0.05, P = 0.011 , P(strict) = 0.184) which is maintained throughout action planning (EBA =

0.27 ± 0.07, P = 0.008 , P(strict) = 0.140) and execution (EBA = 0.29 ± 0.06, P = 0.002 , P(strict) = 0.037).

In contrast, for aIPS the univariate analysis reveal only a significant response to action execution (aIPS =

0.24 ± 0.07, P = 0.016, P(strict) = 0.273) that is absent for the presentation (aIPS = 0.03 ± 0.06, P = 1.00,

P(strict) = 1.00) and planning phase (aIPS = 0.25 ± 0.05, P = 0.07, P(strict) = 1.00).

Qualitative (non-statistical) assessment of the condition RDMs. In line with the univariate

analysis, the condition RDMs for aIPS show that activation patterns remain very dissimilar and then

show an increase following action execution. As such, the condition RDMs indicate that aIPS is primarily

involved in action execution. In the condition RDMs for EBA, a clear visual response can be seen as

within-condition similarity increases early during trials. Interestingly, in line with the univariate analysis,

within-condition similarity increases again following task execution supporting the notion of EBA's

involvement in motor execution and/or visual feedback of the hand, which in the case of the condition

RDMs would relate to the difference in perceived motion.

Temporal multivoxel pattern analysis. tMVPA can discriminate activation patterns in aIPS related

to grasping and turning during the early execution phase, again supporting the notion that aIPS is

involved in the execution of hand-object interactions. Interestingly, tMVPA was able to discriminate

activation patterns related to grasping and turning during the planning phase as well as when

dissimilarity was calculated between timepoints of the planning phase and timepoints of the execution

phase. EBA showed sporadically significant differences between the two tasks during the early visual

response and planning.

Representational similarity analysis. From aIPS, RSA could discriminate the activation patterns

associated with grasping and turning during the planning phase as well as during the execution phase

(Figure 6; last column; aIPS; blue trace; Table 2 for data of timepoints 6 and 7). For EBA, the RSA

revealed an early visual response as it was able to decode starting orientation from the activation

patterns following object presentation (Figure 6; last column; EBA; green trace). Interestingly, in with

the previous analyses, RSA was able to decode grasp versus turn in both the planning phase as well as

the execution phase (Figure 6; last column; EBA; blue trace).

In sum, our findings for aIPS are in line with previous studies that showed its involvement in

motor planning and execution (e.g., Singhal et al., 2013). Interestingly, our multivariate analyses support

the suggestion that EBA, perhaps along with hand-selective divisions of the lateral occipitotemporal

cortex, is not a purely visual region but is also involved in analyzing visual feedback of the body and hand

during motor execution (Astafiev et al., 2004; van den Heiligenberg et al., 2018; Zimmermann et al.,

2016)

<u>3.4 Other regions of interest</u>

We focussed the presentation of the results above in the most interesting findings about how

hand-object interactions unfold over time. Other regions showed effects that were absent, weaker or

less surprising. The condition RDMs and tMVPA results for the previously discussed ROIs and other ROIs

can be found in Figure 5. The RSA findings of all regions can be found on GitHub

(https://github.com/GuyRens/OvenDials).

Briefly, we found weak effects (no significant tMVPA) for the parietal regions of pSPOC and

aSPOC (tMVPA: no significant squares; RSA: no significant models) which have been considered to be

typically involved in hand-object interactions. For DLPFC the tMVPA revealed only one significant

difference for grasp vs turn during late execution. RSA revealed that both the end orientation and motor

task model were significant for at least one timepoint (when applying strict FDR, i.e., taking all ROIs in

account when correcting for multiple corrections) indicating the involvement of DLPFC in motor

execution. For SMA and pre-SMA, the RSA revealed that the start orientation model was significant for

SMA at timepoint 4 and the end orientation model at timepoint 10 for pre-SMA. Both regions showed

significant negative values (grasp > turn) for limited cells when correlations were computed on the

planning and execution phase. For S1, RSA showed significance for the start orientation and motor task

during late execution. This is supported by tMVPA results which showed higher similarity for turn than

for grasp during this phase. Finally, both PMv and M1 show no response in the tMVPA results. It should

be noted in Figure 5 that the subtraction matrix shows a significant cell for M1 being a correlation

between the presentation phase and the execution phase. Similarly, for the RSA results the end

orientation model is significant PMv timepoint 3. Both these results are likely to be analytical anomalies

as participants had no information about the upcoming motor task at these stages of the experimental

trials. To end, both PMv and M1 reached significance for the grasp versus turn model for one timepoint

during motor execution further supporting the well-accepted of these regions involvement in motor

control.

4. Discussion

Our results (summarized by the outlined circles in Figure 2) provide new insights into how neural

representations unfold over time for a simple prehension task vs. a more complex manipulation task.

While prior work has shown that different types of grasp sequences can be decoded (Gallivan et al.,
2016), the current study examines the temporal coding of this information across planning and
execution, as well as the nature of the information being represented in individual brain regions over
time. In addition, our study examines the activity of regions within the SPL, not previously explored in
the prior work. Our additions corroborate earlier findings (Gallivan et al., 2016, 2019) that action
sequences are represented in visual areas (such as V1 and EBA), but also implicate functional
subdivisions of the SPL (areas 7A, 7PC, 7P and 5L).

Using RSA, we found that while some regions, particularly V1, coded start orientation during
planning, multiple regions (V1, cIPS, SPL and EBA) coded a combination of start orientation, end
orientation and task during execution. This finding suggests that the activation patterns in these areas
reflect more than simple visual-perceptual information such, as the visual orientation of the dial
(Kamitani & Tong, 2005); instead, these patterns reflect the combination of the initial goal, the motor
specific act, and the final outcome. Moreover, the similarity of tMPVA activation patterns across
planning and execution, particularly in cIPS and area 7PC, suggests that an action and its outcome are
anticipated well before the movement begins.

<u>Temporal Unfolding of Complex vs. Simple Actions</u>

We investigated the temporal unfolding of complex vs. simple actions using multiple measures
of neural responses: univariate time courses (Figure 1), RSA for sequential volumes (Figures 4 and 6)
and, tMVPA (Figures 3, 5 and 6). These different measures provide a fuller picture of how actions unfold
over time than considering each approach in isolation. For example, univariate signals reveal that both
V1 and cIPS show robust visual signals in all three phases of the trial (presentation, plan and execute), as
do SPL regions and EBA to a lesser degree. tMVPA shows consistency in the patterns of activation in
these areas throughout the trial. However, RSA shows that different areas are representing different
kinds of information at different points in the trial. For example, V1 represents start orientation early in
the trial and task later, while 7PC represents task primarily starting from task onset. Moreover, tMVPA
shows that trial-by-trial similarity during execution and between execution and planning is higher for
Turn than Grasp trials.

These results build upon earlier work examining univariate and multivariate time courses during
motor tasks (Ariani et al., 2018b; Gallivan et al., 2016c). They also provide a valuable complement for
EEG studies that reveal temporal coding with even finer temporal resolution but poorer spatial
resolution. An EEG study by (Guo et al., 2019) showed that grasp orientation (defined by instruction

rather than object attributes) can be classified during both a visual preview and action execution, with
similar representations between the two phases; moreover, their most informative electrodes were
over left caudal parietal cortex, though source localization indicated diverse potential sources (that may
include cIPS, SPL, and EBA).

One of the more surprising results of our TMVPA and RSA findings was the involvement of SPL in
showing differences between Turn and Grasp actions, in particular areas 7A and 7PC. The role of the SPL
in hand actions has been relatively less studied than areas in the IPS (especially cIPS and aIPS) and SPOC
(V6/V6A), for which clear homologies between humans and other primates have been proposed
(Culham & Kanwisher, 2001; Grefkes & Fink, 2005). One partial exception is area 7PC, an SPL region
dorsal to aIPS, that has been proposed as the human homologue of the macaque medial intraparietal
sulcus (mIPS) and/or parietal reach region (PRR), albeit with limited consensus about locus (Gallivan &
Culham, 2015) and putative homologies (Culham et al., 2006). These SPL areas likely form part of the
dorsomedial (or dorsal-dorsal) visual stream (Rizzolatti & Matelli, 2003) and have been postulated as an
integration zone for visual, motor, and somatosensory information for goal-directed movements
(Passarelli et al., 2021). Human and macaque SPL have similar structural organization with the anterior
sector -- 7PC (macaque homologue: VIP; Caminiti et al., 2015) and 5L (macaque homologue: PE;
Gamberini et al., 2020) -- being predominantly somatosensory and the more caudal parts -- 7A
(macaque homologue: PEc; Gamberini et al., 2020) and 7P (macaque homologue: V6A; Gamberini et al.,
2020) -- being both somatosensory and visual (Gamberini et al., 2020). The anterior regions of the SPL
(5L and 7PC) fall on the more somatosensory side of the visual-to-somatosensory gradient from the
posterior to anterior SPL. As such, they may be involved in anticipating the sensory (perhaps largely
somatosensory) and motor requirements and consequences of the action.

<u>Orientation Processing</u>

Our experiment explicitly involved tasks that required processing of object orientation for
grasping and turning. While processing of object location for reaching and object shape and size for
grasping have been well characterized behaviorally and neurally (Cavina-Pratesi et al., 2010), the
influence of object orientation on reach-to-grasp actions has only recently been investigated in the
monkey and human. These studies have suggested that object orientation and/or wrist/grip orientation
rely on V6A/aSPOC (Battaglini et al., 2002; Monaco et al., 2011) and the adjacent cIPS (Rice et al., 2007;
Shikata et al., 2001; Valyear et al., 2006) perhaps along with aIPS (Breveglieri et al., 2022). One fMRI
study that had participants grasp a dial much like ours found that "bistable" object orientations (which

equally afforded two possible grip postures) evoked greater activation in cIPS than stable orientations
(which afforded only one grip posture) (Wood et al., 2017). Moreover, this study found that two
neuropsychological patients with unilateral damage to cIPS showed deficits in grasping bistable
orientations with the contralateral hand. We find that cIPS codes a combination of start orientation, end
orientation, and task during action execution. Moreover, cIPS activation is more consistent throughout a
trial in a complex turning task, where object orientation must be changed through a grip rotation, than a
simpler grasp-only task, corroborating the proposed role of cIPS in processing orientation for action. To
our surprise, we did not observe any orientation coding in aSPOC or pSPOC.

Akin to cIPS, early visual cortex has been increasingly implicated in motor planning and
execution (Gallivan et al., 2014, 2019; Gutteling et al., 2015; Monaco et al., 2020). Our findings suggest
that the visual cortex (V1) codes the start orientation of the object throughout the trial with increases in
the task representation toward the end of planning and during execution. This suggests that V1
activation patterns are not solely driven in a bottom-up fashion by current visual cues (Kamitani & Tong,
2005) but come to incorporate the complexity of the task and visual information as the action unfolds.

Although studies of EBA have focused on its role in body perception (Kontaris et al., 2009), EBA
has also been implicated in processing body posture during planning and execution of goal-directed
actions (Gallivan et al., 2016; Zimmermann et al., 2018). We found some evidence for this notion: EBA
has strong representations of task and orientations during execution but only limited ones during
planning. As such, EBA may be mainly driven by the perceptual consequences of the action.

<u>Methodological considerations</u>

Our "grasping" actions were atypical in that they required gripping the dial between the
knuckles of the index and middle fingers, in a way that was similar between grasping, turning to the left,
and turning to the right. This choice of posture was deliberate, to ensure that the representations of the
upcoming task were not confounded with differences in initial grip posture. Nevertheless, this strategy
differs considerably from the approach that participants would likely have undertaken without such
instruction (Rosenbaum et al., 1990). Nevertheless, we found multivariate differences even in the case
when the low-level grasp posture remained constant (Ariani et al., 2015; Ramon et al., 2015). Finally,
we focused on the dorsal pathway given the low motor complexity (Goodale & Milner, 1992). However,
using more complex objects could be used to explore ventral stream involvement: Dijkerman et al,
(2009) highlighted its involvement in high-level grip selection. In addition, it has been argued that the
ventral stream, in particular EBA (Zimmermann et al., 2016) provides an initial structure to the motor

plans for grasping whereas regions of the SPL might be more feedback-driven (Scheperjans, Hermann,

et al., 2008). Techniques with high temporal resolution such as TMS and EEG could explore the

relationship between these regions that have similar activation patterns in our study (Figure 6) but that

have been considered to be functionally different.

<u>Conclusion</u>

We investigated the neural underpinnings of how hand-object interactions unfold on a moment-

to-moment basis. Importantly, we found that the neural underpinnings of grasping and turning actions

could best be differentiated in occipital cortex (particularly cIPS and V1) and SPL. These results

corroborate the importance of cIPS in processing object and grip orientation and suggest that SPL may

play a larger role in action sequences than previously realized. Our results also show how tMVPA can be

used to understand how motor actions are represented across different phases of the trial.

Bibliography

Ariani, G., Oosterhof, N. N., & Lingnau, A. (2018). Time-resolved decoding of planned delayed and
immediate prehension movements. *Cortex, 99*, 330–345.
https://doi.org/10.1016/j.cortex.2017.12.007

Ariani, G., Wurm, M. F., & Lingnau, A. (2015). Decoding internally and externally driven movement
plans. *Journal of Neuroscience, 35*(42), 14160–14171. https://doi.org/10.1523/JNEUROSCI.0596-
15.2015

Astafiev, S. v., Stanley, C. M., Shulman, G. L., & Corbetta, M. (2004). Extrastriate body area in human
occipital cortex responds to the performance of motor actions. *Nature Neuroscience, 7*(5), 542–
548. https://doi.org/10.1038/nn1241

Badre, D., & Nee, D. E. (2018). Frontal Cortex and the Hierarchical Control of Behavior. *Trends in
Cognitive Sciences, 22*(2), 170–188. https://doi.org/10.1016/j.tics.2017.11.005

Battaglini, P., Muzur, A., Galletti, C., Skrap, M., Brovelli, A., & Fattori, P. (2002). Effects of lesions to area
V6A in monkeys. *Experimental Brain Research, 144*(3), 419–422. https://doi.org/10.1007/s00221-
002-1099-4

Beurze, S. M., De Lange, F. P., Toni, I., & Medendorp, W. P. (2007). Integration of target and effector
information in the human brain during reach planning. *Journal of Neurophysiology, 97*(1), 188–
199. https://doi.org/10.1152/jn.00456.2006

Beurze, S. M., De Lange, F. P., Toni, I., & Medendorp, W. P. (2009). Spatial and effector processing in the
human parietofrontal network for reaches and saccades. *Journal of Neurophysiology, 101*(6),
3053–3062. https://doi.org/10.1152/jn.91194.2008

Breveglieri, R., Borgomaneri, S., Filippini, M., Tessari, A., Galletti, C., Davare, M., & Fattori, P. (2022).
Complementary contribution of the medial and lateral human parietal cortex to grasping: a
repetitive TMS study. *Cerebral Cortex.* https://doi.org/10.1093/cercor/bhac404

Caminiti, R., Innocenti, G. M., & Battaglia-Mayer, A. (2015). Organization and evolution of parieto-
frontal processing streams in macaque monkeys and humans. *Neuroscience and Biobehavioral
Reviews, 56*, 73–96. https://doi.org/10.1016/j.neubiorev.2015.06.014

Castiello, U. (2005). The neuroscience of grasping. *Nature Reviews. Neuroscience*, 6(9), 726–736.
https://doi.org/10.1038/nrn1744
Cavina-Pratesi, C., Connolly, J. D., Monaco, S., Figley, T. D., Milner, A. D., Schenk, T., & Culham, J. C.
(2018). Human neuroimaging reveals the subcomponents of grasping, reaching and pointing
actions. *Cortex*, 98, 128–148. https://doi.org/10.1016/j.cortex.2017.05.018
Cavina-Pratesi, C., Monaco, S., Fattori, P., Galletti, C., McAdam, T. D., Quinlan, D. J., Goodale, M. A., &
Culham, J. C. (2010). Functional magnetic resonance imaging reveals the neural substrates of arm
transport and grip formation in reach-to-grasp actions in humans. *Journal of Neuroscience*, 30(31),
10306–10323. https://doi.org/10.1523/JNEUROSCI.2023-10.2010
Comalli, D. M., Keen, R., Abraham, E. S., Foo, V. J., Lee, M. H., & Adolph, K. E. (2016). The development
of tool use: Planning for end-state comfort. *Developmental Psychology*, 52(11), 878–1892.
https://doi.org/10.1037/dev0000207
Culham, J. C., Cavina-Pratesi, C., & Singhal, A. (2006). The role of parietal cortex in visuomotor control:
What have we learned from neuroimaging? *Neuropsychologia*, 44(13), 2668–2684.
https://doi.org/10.1016/j.neuropsychologia.2005.11.003
Culham, J. C., Danckert, S. L., DeSouza, J. F. X., Gati, J. S., Menon, R. S., & Goodale, M. A. (2003). Visually
guided grasping produces fMRI activation in dorsal but not ventral stream brain areas.
Experimental Brain Research, 153(2), 180–189. https://doi.org/10.1007/s00221-003-1591-5
Culham, & Kanwisher. (2001). Neuroimaging of cognitive functions in human parietal cortex. *Current
Opinion in Neurobiology*.
Di Bono, M. G., Begliomini, C., Castiello, U., & Zorzi, M. (2015). Probing the reaching-grasping network in
humans through multivoxel pattern decoding. *Brain and Behavior*, 5(11), 1–18.
https://doi.org/10.1002/brb3.412
Dijkerman, H. C., McIntosh, R. D., Schindler, I., Nijboer, T. C. W., & Milner, A. D. (2009). Choosing
between alternative wrist postures: Action planning needs perception. *Neuropsychologia*, 47(6),
1476–1482. https://doi.org/10.1016/j.neuropsychologia.2008.12.002
Errante, A., Ziccarelli, S., Mingolla, G., & Fogassi, L. (2021). Grasping and Manipulation: Neural Bases
and Anatomical Circuitry in Humans. *Neuroscience*, 458, 203–212.
https://doi.org/10.1016/j.neuroscience.2021.01.028
Etzel, J. A., Zacks, J. M., & Braver, T. S. (2013). Searchlight analysis: Promise, pitfalls, and potential. In
NeuroImage (Vol. 78, pp. 261–269). https://doi.org/10.1016/j.neuroimage.2013.03.041
Fabbri, S., Stubbs, K. M., Cusack, R., & Culham, J. C. (2016). Disentangling Representations of Object and
Grasp Properties in the Human Brain. *The Journal of Neuroscience*, 36(29), 7648–7662.
https://doi.org/10.1523/JNEUROSCI.0313-16.2016
Fischl, B., Sereno, M. I., Tootell, R. B. H., & Dale, A. M. (1999). High-Resolution Intersubject Averaging
and aCoordinate System for the Cortical Surface. *Human Brain Mapping*.
Fogassi, L., Ferrari, P. F., Gesierich, B., Rozzi, S., Chersi, F., & Rizzolotti, G. (2005). Parietal lobe: From
action organization to intention understanding. *Science*, 308(5722), 662–667.
https://doi.org/10.1126/science.1106138
Frost, M. A., & Goebel, R. (2012). Measuring structural–functional correspondence: Spatial variability of
specialised brain regions after macro-anatomical alignment. *NeuroImage*, 59(2), 1369–1381.
https://doi.org/10.1016/j.neuroimage.2011.08.035
Gallivan, J. P., Adam McLean, D., Valyear, K. F., & Culham, J. C. (2013). Decoding the neural mechanisms
of human tool use. *ELife*, 2013(2), 1–29. https://doi.org/10.7554/eLife.00425
Gallivan, J. P., Cant, J. S., Goodale, M. A., & Flanagan, J. R. (2014). Representation of object weight in
human ventral visual cortex. *Current Biology*, 24(16), 1866–1873.
https://doi.org/10.1016/j.cub.2014.06.046

Gallivan, J. P., Cavina-Pratesi, C., & Culham, J. C. (2009). Is that within reach? fMRI reveals that the human superior parieto-occipital cortex encodes objects reachable by the hand. *Journal of Neuroscience*, *29*(14), 4381–4391. https://doi.org/10.1523/JNEUROSCI.0377-09.2009

Gallivan, J. P., Chapman, C. S., Gale, D. J., Flanagan, J. R., & Culham, J. C. (2019). Selective Modulation of Early Visual Cortical Activity by Movement Intention. *Cerebral Cortex*, 1–17. https://doi.org/10.1093/cercor/bhy345

Gallivan, J. P., & Culham, J. C. (2015). Neural coding within human brain areas involved in actions. In *Current Opinion in Neurobiology* (Vol. 33, pp. 141–149). Elsevier Ltd. https://doi.org/10.1016/j.conb.2015.03.012

Gallivan, J. P., Johnsrude, I. S., & Randall Flanagan, J. (2016). Planning Ahead: Object-Directed Sequential Actions Decoded from Human Frontoparietal and Occipitotemporal Networks. *Cerebral Cortex*, *26*(2), 708–730. https://doi.org/10.1093/cercor/bhu302

Gallivan, J. P., McLean, D. A., Valyear, K. F., Pettypiece, C. E., & Culham, J. C. (2011). Decoding action intentions from preparatory brain activity in human parieto-frontal networks. *Journal of Neuroscience*, *31*(26), 9599–9610.

Gamberini, M., Passarelli, L., Fattori, P., & Galletti, C. (2020). Structural connectivity and functional properties of the macaque superior parietal lobule. *Brain Structure and Function*, *225*(4), 1349–1367. https://doi.org/10.1007/s00429-019-01976-9

Gerbella, M., Rozzi, S., & Rizzolatti, G. (2017). The extended object-grasping network. *Experimental Brain Research*, *235*(10), 2903–2916. https://doi.org/10.1007/s00221-017-5007-3

Goebel, R., Esposito, F., & Formisano, E. (2006). Analysis of Functional Image Analysis Contest (FIAC) data with BrainVoyager QX: From single-subject to cortically aligned group General Linear Model analysis and self-organizing group Independent Component Analysis. *Human Brain Mapping*, *27*(5), 392–401. https://doi.org/10.1002/hbm.20249

Goodale, M. A., & Milner, D. A. (1992). Separate visual pathways for perception and action. *Trends in Neurosciences*.

Greenfield, A., Jahnsen, H., Downing, P. E., Jiang, Y., Shuman, M., & Kanwisher, N. (1996). A Cortical Area Selective for Visual Processing of the Human Body. In *Trends Pharmacol. Sci* (Vol. 16). https://www.science.org

Grefkes, C., & Fink, G. R. (2005). *The functional organization of the intraparietal sulcus in humans and monkeys*. 3–17.

Greve, D. N., & Fischl, B. (2009). Accurate and Robust Brain Image Alignment using Boundary- based Registration. *Neuroimage*, *48*(1), 63–72. https://doi.org/10.1016/j.neuroimage.2009.06.060.Accurate

Guo, L. L., Nestor, A., Nemrodov, D., Frost, A., & Niemeier, M. (2019). Multivariate analysis of electrophysiological signals reveals the temporal properties of visuomotor computations for precision grips. *Journal of Neuroscience*, *39*(48), 9585–9597. https://doi.org/10.1523/JNEUROSCI.0914-19.2019

Gutteling, T. P., Petridou, N., Dumoulin, S. O., Harvey, B. M., Aarnoutse, E. J., Leon Kenemans, J., & Neggers, S. F. W. (2015). Action preparation shapes processing in early visual cortex. *Journal of Neuroscience*, *35*(16), 6472–6480. https://doi.org/10.1523/JNEUROSCI.1358-14.2015

Haxby, J. V., Gobbini, M. I., Furey, M. L., Ishai, A., Schouten, J. L., & Pietrini, P. (2001). Distributed and overlapping representations of faces and objects in ventral temporal corten. *Science*, *293*(September), 87–96. https://doi.org/10.4324/9780203496190

Hinrichs, H., Scholz, M., Tempelmann, C., Woldorff, M. G., Dale, A. M., & Heinze, H. J. (2000). Deconvolution of event-related fMRI responses in fast-rate experimental designs: Tracking amplitude variations. *Journal of Cognitive Neuroscience*, *12*. https://doi.org/10.1162/089892900564082

Kamitani, Y., & Tong, F. (2005). Decoding the visual and subjective contents of the human brain. *Nature*
Neuroscience, *8*(5), 679–685. https://doi.org/10.1038/nn1444
Kontaris, I., Wiggett, A. J., & Downing, P. E. (2009). Dissociation of extrastriate body and biological-
motion selective areas by manipulation of visual-motor congruency. *Neuropsychologia*, *47*(14),
3118–3124. https://doi.org/10.1016/j.neuropsychologia.2009.07.012
Kriegeskorte, N. (2008). Representational similarity analysis – connecting the branches of systems
neuroscience. *Frontiers in Systems Neuroscience*, *2*(November), 1–28.
https://doi.org/10.3389/neuro.06.004.2008
Kriegeskorte, N., & Goebel, R. (2001). An efficient algorithm for topologically correct segmentation of
the cortical sheet in anatomical MR volumes. *NeuroImage*, *14*(2), 329–346.
https://doi.org/10.1006/nimg.2001.0831
Kriegeskorte, N., & Kievit, R. A. (2013). Representational geometry: Integrating cognition, computation,
and the brain. *Trends in Cognitive Sciences*, *17*(8), 401–412.
https://doi.org/10.1016/j.tics.2013.06.007
Monaco, S., Cavina-Pratesi, C., Sedda, A., Fattori, P., Galletti, C., & Culhaml, J. C. (2011). Functional
magnetic resonance adaptation reveals the involvement of the dorsomedial stream in hand
orientation for grasping. *Journal of Neurophysiology*, *106*(5), 2248–2263.
https://doi.org/10.1152/jn.01069.2010
Monaco, S., Malfatti, G., Culham, J. C., Cattaneo, L., & Turella, L. (2020). Decoding motor imagery and
action planning in the early visual cortex: Overlapping but distinct neural mechanisms.
NeuroImage, *218*(May), 116981. https://doi.org/10.1016/j.neuroimage.2020.116981
Mylius, V., Ayache, S. S., Ahdab, R., Farhat, W. H., Zouari, H. G., Belke, M., Brugières, P., Wehrmann, E.,
Krakow, K., Timmesfeld, N., Schmidt, S., Oertel, W. H., Knake, S., & Lefaucheur, J. P. (2013).
Definition of DLPFC and M1 according to anatomical landmarks for navigated brain stimulation:
inter-rater reliability, accuracy, and influence of gender and age. *NeuroImage*, *78*, 224–232.
https://doi.org/10.1016/j.neuroimage.2013.03.061
Passarelli, L., Gamberini, M., & Fattori, P. (2021). The superior parietal lobule of primates: A sensory-
motor hub for interaction with the environment. In *Journal of Integrative Neuroscience* (Vol. 20,
Issue 1, pp. 157–171). IMR Press Limited. https://doi.org/10.31083/J.JIN.2021.01.334
Pertzov, Y., Avidan, G., & Zohary, E. (2011). Multiple reference frames for saccadic planning in the
human parietal cortex. *Journal of Neuroscience*, *31*(3), 1059–1068.
https://doi.org/10.1523/JNEUROSCI.3721-10.2011
Picard, N., & Strick, P. L. (2001). Imaging the premotor areas. *Current Opinion in Neurobiology*, *11*(6),
663–672. https://doi.org/10.1016/S0959-4388(01)00266-5
Pilacinski, A., Wallscheid, M., & Lindner, A. (2018). Human posterior parietal and dorsal premotor cortex
encode the visual properties of an upcoming action. *PLoS ONE*, *13*(10), 1–20.
https://doi.org/10.1371/journal.pone.0198051
Pitzalis, S., Fattori, P., & Galletti, C. (2015). The human cortical areas V6 and V6A. In *Visual Neuroscience*
(Vol. 32). Cambridge University Press. https://doi.org/10.1017/S0952523815000048
Ramon, M., Vizioli, L., Liu-Shuang, J., & Rossion, B. (2015). Neural microgenesis of personally familiar
face recognition. *Proceedings of the National Academy of Sciences of the United States of America*,
112(35), E4835–E4844. https://doi.org/10.1073/pnas.1414929112
Rice, N. J., Valyear, K. F., Goodale, M. A., Milner, A. D., & Culham, J. C. (2007). Orientation sensitivity to
graspable objects: An fMRI adaptation study. *NeuroImage*, *36*(SUPPL. 2).
https://doi.org/10.1016/j.neuroimage.2007.03.032
Richter, M., Amunts, K., Mohlberg, H., Bludau, S., Eickhoff, S. B., Zilles, K., & Caspers, S. (2019).
Cytoarchitectonic segregation of human posterior intraparietal and adjacent parieto-occipital

sulcus and its relation to visuomotor and cognitive functions. *Cerebral Cortex*, *29*(3), 1305–1327. https://doi.org/10.1093/cercor/bhy245

Rizzolatti, G., & Matelli, M. (2003). Two different streams form the dorsal visual system: Anatomy and functions. *Experimental Brain Research*, *153*(2), 146–157. https://doi.org/10.1007/s00221-003-1588-0

Rosenbaum, D. A., Marchak, F., Barnes, H. J., Vaughan, J., Slotta, J. D., & Jorgensen, M. J. (1990). Constraints for Action Selection: Overhand Versus Underhand Grips. *Attention and Performance XIII: Motor Representation and Control, August 2014*, 321–342. https://doi.org/10.4324/9780203772010-10

Rosenke, M., van Hoof, R., van den Hurk, J., Grill-Spector, K., & Goebel, R. (2021). A Probabilistic Functional Atlas of Human Occipito-Temporal Visual Cortex. *Cerebral Cortex*, *31*(1), 603–619. https://doi.org/10.1093/cercor/bhaa246

Scheperjans, F., Eickhoff, S. B., Hömke, L., Mohlberg, H., Hermann, K., Amunts, K., & Zilles, K. (2008). Probabilistic maps, morphometry, and variability of cytoarchitectonic areas in the human superior parietal cortex. *Cerebral Cortex*, *18*(9), 2141–2157. https://doi.org/10.1093/cercor/bhm241

Scheperjans, F., Hermann, K., Eickhoff, S. B., Amunts, K., Schleicher, A., & Zilles, K. (2008). Observer-independent cytoarchitectonic mapping of the human superior parietal cortex. *Cerebral Cortex*, *18*(4), 846–867. https://doi.org/10.1093/cercor/bhm116

Shikata, E., Hamzei, F., Glauche, V., Knab, R., Dettmers, C., Weiller, C., Bu¨chel, C., & Bu¨chel, B. (2001). *Surface Orientation Discrimination Activates Caudal and Anterior Intraparietal Sulcus in Humans: An Event-Related fMRI Study*. www.jn.physiology.org

Singhal, A., Monaco, S., Kaufman, L. D., & Culham, J. C. (2013). Human fMRI Reveals That Delayed Action Re-Recruits Visual Perception. *PLoS ONE*, *8*(9). https://doi.org/10.1371/journal.pone.0073629

Tomassini, V., Jbabdi, S., Klein, J. C., Behrens, T. E. J., Pozzilli, C., Matthews, P. M., Rushworth, M. F. S., & Johansen-Berg, H. (2007). Diffusion-weighted imaging tractography-based parcellation of the human lateral premotor cortex identifies dorsal and ventral subregions with anatomical and functional specializations. *Journal of Neuroscience*, *27*(38), 10259–10269. https://doi.org/10.1523/JNEUROSCI.2144-07.2007

Valyear, K. F., Culham, J. C., Sharif, N., Westwood, D., & Goodale, M. A. (2006). A double dissociation between sensitivity to changes in object identity and object orientation in the ventral and dorsal visual streams: A human fMRI study. *Neuropsychologia*, *44*(2), 218–228. https://doi.org/10.1016/j.neuropsychologia.2005.05.004

van den Heiligenberg, F. M. Z., Orlov, T., Macdonald, S. N., Duff, E. P., Henderson Slater, D., Beckmann, C., Johansen-Berg, H., Culham, J. C., & Makin, T. R. (2018). Artificial limb representation in amputees. *Brain*, *March*, 1–12. https://doi.org/10.1093/brain/awy054

Vizioli, L., Bratch, A., Lao, J., Ugurbil, K., Muckli, L., & Yacoub, E. (2018). Temporal multivariate pattern analysis (tMVPA): A single trial approach exploring the temporal dynamics of the BOLD signal. *Journal of Neuroscience Methods*, *308*(June 2018), 74–87. https://doi.org/10.1016/j.jneumeth.2018.06.029

Wandell, B. A., Dumoulin, S. O., & Brewer, A. A. (2007). Visual field maps in human cortex. In *Neuron* (Vol. 56, Issue 2, pp. 366–383). https://doi.org/10.1016/j.neuron.2007.10.012

Wood, D. K., Chouinard, P. A., Major, A. J., & Goodale, M. A. (2017). Sensitivity to biomechanical limitations during postural decision-making depends on the integrity of posterior superior parietal cortex. *Cortex*, *97*, 202–220. https://doi.org/10.1016/j.cortex.2016.07.005

Yousry, T. A., Schmid, U. D., Alkadhi, H., Schmidt, D., Peraud, A., Buettner, A., & Winkler, P. (1997). Localization of the motor hand area to a knob on the precentral gyrus A new landmark. *Brain*, 141–157.

Zimmermann, M., Mars, R. B., de Lange, F. P., Toni, I., & Verhagen, L. (2018). Is the extrastriate body
area part of the dorsal visuomotor stream? *Brain Structure and Function*, *223*(1), 31–46.
https://doi.org/10.1007/s00429-017-1469-0
Zimmermann, M., Verhagen, L., de Lange, F. P., & Toni, I. (2016). The extrastriate body area computes
desired goal states during action planning. *ENeuro*, *3*(2), 1918–1921.
https://doi.org/10.1523/ENEURO.0020-16.2016

1156
1157

Figure legends

Figure 1. Experimental set-up and design. **1A.** Picture of participant set-up in the fMRI scanner shown from side view. **1B.** Close up of participant using the instructed grip to grasp the experimental stimulus, i.e., rotatable dial with a handle. **1C.** Experimental conditions in a two (start orientation: left or right) by two (action: grasp or turn) design. **2D.** Timing of each event-related trial. Trials began with the stimulus being illuminated (TR 0-3) indicating the start of the preparation phase. At TR 3 participants received task instructions indicating the start of the planning phase (TR 3-9). At TR 9 participants received the "go" cue indicating onset of action execution (TR 9-10). After 2 s (1 TR) the light was turned off and the intertrial interval was initiated (TR 10-16)

Figure 2. Brain areas selected for multivariate analysis based on a univariate contrast. Cortical areas that exhibited larger responses during the experimental trials [(presentation + plan + execute) > baseline] are shown in yellow to red activation. Results calculated across all participants (RFX GLM) are displayed on the dynamic group average surface across participants. The selected ROIs were selected in BrainVoyager software, which uses a hexagonal selection area, and then transformed to individual volumetric MNI space. Each ROI is linked to the group average of corresponding % signal change of BOLD activity (Y-axis) over time (X-axis; each X-ticks represents one TR of 2 s, based on a deconvolution analysis) for each of the four conditions. Vertical dashed lines on the graphs indicate start of the planning phase (TR 3) and the execution phase (TR 9). For the time courses, statistical analysis was done for the shaded areas (e.g., average timepoint 2 and 3) averaged across the 4 conditions. If significantly > 0, when Bonferroni corrected within ROI-only, a small asterisk is depicted. If significantly > 0, when Bonferroni corrected for all comparisons, a large asterisk is depicted. On the surface brain, sulcal landmarks are denoted by white lines (stylized according to the corresponding legend). ROI acronyms are spelled out in the methods. tMVPA results are indicated using different types of circles (e.g., dashed vs. solid). RSA results are shown by color-coding the tMVPA circles using a Venn diagram. Note that the RSA and tMVPA results are discussed in the results section as well as in Figure 5 and Figure 6.

Figure 3. Overview of the steps to perform temporal multivoxel pattern analysis (see methods for full explanation). The figure represents the steps taken for each participant separately. Within each condition single trial representational distance matrices (RDMs) are calculated for each timepoint of each trial pairing (**3A** shows example for grasping trials and **3B** for turning trials). **3C:** Grey cubes represent voxels of the same ROI during different timepoints for two sample trials to exemplify how cells for each RDM are calculated. After calculating single trial RDMs, a condition average is calculated (middle column; **3D** and **3E** for grasping and turning respectively) which are then subtracted from each other (last column; **3F**; red and blue matrix). Not shown on picture (for results see Figure 4 and 5): the subtraction matrices are then used for bootstrapping to determine whether the group average differs significantly from zero and to investigate whether one condition is more similar than the other for each given timepoint

Figure 4. Models used for representational similarity analysis (RSA). The first row shows the models for start orientation (**4A**), end orientation (**4B**) and turn versus grasp (**4C**). The second row shows example data for one single timepoint for two regions of interest (**4D**: V1d); **4E**: Area-7A) of one participant.

Figure 5. Left: Hierarchical clustering of all the ROIs based on the averaged representational dissimilarity matrices (RDMs) of the **Grasp** and **Turn** condition. We refer the reader to the legend of Figure 3 for the full explanation of the heatmaps. Note that **T-G** represents 'Turn minus Grasp' and represents the final

tMVPA result showing statistically significant similarities (small red boxes = Turn > Grasp; small blue
boxes = Grasp > Turn). The outline of the significance box indicates significance level. When the outline
is dashed, the square was significant only when Bonferroni corrected for within-ROI comparisons. When
the outline is solid, the square was significant when Bonferroni corrected for all comparisons across all
ROIs. Note that we only tested whether boxes were significantly different from zero (and not from each
other).

Figure 6. Selection of ROIs that showed prominent differences for the tMVPA analysis. Some data are
reproduced from subsets of Figures 2 (Column 1) and 5 (Columns 2-4, at higher resolution) to facilitate
inspection and comparisons across analysis methods for key regions. **Column 1** shows the group average
deconvolution time courses for the four conditions. The X-axis represents time (in 2 s TRs) and the Y-axis
represents % signal change in the BOLD signal. Vertical dashed lines represent onset of the planning
phase (TR 3) (after the initial presentation phase) and the execution phase (TR 9) following the planning
phase. **Columns 2 and 3** show the group average condition representational dissimilarity matrices for
the grasp (column 2) and turn condition (column 3). Each cell/square of the matrix represents the
dissimilarity between trials of the same condition between two timepoints. Dashed horizontal and
vertical lines represent the same timepoints as those in Column 1, i.e., the first dashed
horizontal/vertical line represents the start of the planning phase (TR 3) and the second dashed
vertical/horizontal line represents the start of the execution phase (TR 9). **Column 4.** tMVPA results.
Dashed lines have the same representation as explained for columns 1-3. Red squares indicate
dissimilarity metrics between timepoints where turning trials are significantly more similar (less
dissimilar) than grasping trials. blue squares represent the opposite (grasping more similar than turning).
Column 5. Representational similarity analysis using three models, as explained in the color legend, for
each timepoint of the experimental trials. Solid lines represent the correlation between the data and the
model over time. The shaded grey area represents the noise ceiling with the lower and upper edges
representing the lower and upper bounds. Dashed vertical lines have the same representation as
explained for the previous columns. Colored "plus signs" indicate time points with statistically significant
correlations for the respective model when FDR corrected for all within-ROI comparisons against zero.
Colored asterisks indicate the same when FDR corrected for all comparisons (across all ROIs) against
zero.

1229

Number of Voxels Per Region of Interest

Region of Interest	Number of Functional Voxels
M1	98 ± 7
S1	123 ± 11
PMv	107 ± 8
PMd	217 ± 9
aSPOC	178 ± 14
pSPOC	138 ± 7
aIPS	268 ± 17
cIPS	172 ± 10
Pre-SMA	108 ± 7
SMA	132 ± 10
DLPFC	129 ± 9
V1d	152 ± 13
Area-5L	167 ± 10
Area-7A	194 ± 12
Area-7P	150 ± 9
Area-7PC	198 ± 13
EBA	1163 ± 23

1230   **Table 1.** The average number of functional (1-mm iso-)voxels per Region of Interest. Values shown are
1231   mean ± SEM.

ROI	Phase	Presentation				Planning				Execution			
		Timepoint 2		Timepoint 3		Timepoint 6		Timepoint 7		Timepoint 11		Timepoint 12	
		CI	Q	CI	Q	CI	Q	CI	Q	CI	Q	CI	Q
V1d	Start Ori	-0.728 -0.187 0.355	0.979	-0.411 0.231 0.873	0.192	**-0.246 0.36 0.966**	**0.048**	-0.337 0.155 0.648	0.244	**-0.033 0.302 0.637**	**0.008**	-0.164 0.187 0.538	**0.07**
	End Ori	-0.551 0.009 0.569	0.691	-0.494 -0.009 0.476	0.732	-0.395 0.115 0.626	0.365	-0.234 0.173 0.581	0.14	**-0.156 0.298 0.752**	0.034	-0.284 0.258 0.799	0.103
	GvT	-0.619 -0.018 0.583	0.753	-0.452 0.036 0.523	0.602	-0.201 0.067 0.334	0.331	**-0.164 0.24 0.644**	**0.048**	-0.136 0.315 0.767	0.026	-0.231 0.293 0.817	0.059
cIPS	Start Ori	-0.698 -0.049 0.6	0.81	-0.313 0.191 0.695	0.179	-0.407 0.133 0.673	0.333	-0.497 0.16 0.817	0.338	-0.183 0.209 0.6	0.069	**0.067 0.311 0.555**	**0.001**
	End Ori	-0.321 0.004 0.33	0.694	-0.163 0.111 0.385	0.155	-0.377 0.049 0.475	0.53	-0.253 0.044 0.342	0.478	**-0.072 0.209 0.49**	0.023	0.037 0.351 0.665	0.002
	GvT	-0.599 -0.036 0.528	0.795	-0.439 -0.018 0.403	0.773	-0.378 0.076 0.529	0.451	-0.397 0.195 0.788	0.229	**0.119 0.515 0.912**	**0.001**	0.243 0.609 0.974	0.001
7P	Start Ori	-0.573 -0.218 0.137	0.996	-0.7 -0.138 0.425	0.943	-0.253 0.124 0.502	0.229	-0.297 0.111 0.519	0.299	-0.358 0.062 0.482	0.479	-0.288 0.076 0.439	0.394
	End Ori	-0.59 0 0.59	0.699	-0.583 -0.062 0.459	0.858	-0.275 0.004 0.283	0.691	-0.342 0.098 0.537	0.371	-0.216 0.209 0.633	0.092	**-0.11 0.195 0.501**	**0.038**
	GvT	-0.611 -0.049 0.514	0.824	-0.521 -0.058 0.405	0.86	-0.691 -0.031 0.628	0.776	**-0.127 0.369 0.865**	**0.023**	**-0.031 0.484 1**	**0.008**	-0.156 0.404 0.964	0.025
7A	Start Ori	-0.722 -0.089 0.544	0.873	-0.328 0.213 0.754	0.164	-0.314 0.209 0.732	0.158	-0.278 0.133 0.545	0.235	**-0.037 0.311 0.659**	**0.009**	-0.2 0.169 0.537	0.116
	End Ori	-0.677 -0.111 0.455	0.916	-0.481 -0.04 0.401	0.829	-0.473 -0.044 0.384	0.842	-0.51 0 0.51	0.699	**-0.054 0.32 0.694**	**0.01**	-0.101 0.24 0.58	0.026
	GvT	-0.508 0.04 0.588	0.602	-0.278 0.2 0.678	0.145	-0.317 0.253 0.823	0.129	**-0.171 0.364 0.899**	**0.029**	-0.078 0.462 1.002	0.01	0.061 0.533 1.005	0.002
7PC	Start Ori	-0.57 -0.133 0.304	0.969	-0.53 -0.036 0.459	0.808	-0.383 -0.022 0.339	0.795	-0.473 0.004 0.482	0.699	-0.414 0.04 0.494	0.58	-0.403 0.062 0.527	0.504
	End Ori	-0.597 -0.142 0.313	0.97	-0.481 -0.031 0.419	0.806	-0.392 0.102 0.597	0.395	-0.528 -0.009 0.51	0.731	-0.175 0.138 0.45	0.131	-0.264 0.164 0.593	0.177
	GvT	-0.527 0.044 0.616	0.599	-0.656 -0.129 0.399	0.943	-0.26 0.258 0.775	0.09	-0.263 0.227 0.716	0.113	**-0.071 0.435 0.941**	**0.01**	-0.202 0.369 0.939	0.036
5L	Start Ori	-0.769 -0.227 0.316	0.988	-0.463 0.04 0.543	0.596	-0.395 0 0.395	0.699	-0.519 -0.004 0.51	0.716	-0.209 0.151 0.511	0.144	-0.3 0 0.3	0.699
	End Ori	-0.381 -0.053 0.274	0.887	-0.434 0.058 0.549	0.526	-0.236 0.062 0.36	0.393	-0.262 0.182 0.627	0.151	**-0.029 0.346 0.722**	**0.008**	-0.322 0.098 0.518	0.353
	GvT	-0.554 0.013 0.581	0.68	-0.513 0.098 0.709	0.461	-0.299 0.222 0.743	0.14	-0.717 -0.009 0.699	0.723	**0.026 0.426 0.826**	**0.003**	-0.328 0.329 0.986	0.088
aIPS	Start Ori	-0.543 -0.151 0.241	0.983	-0.567 -0.062 0.443	0.86	-0.192 0.244 0.68	0.059	-0.321 0.098 0.517	0.352	-0.245 0.195 0.636	0.129	-0.325 0.049 0.423	0.505
	End Ori	-0.39 -0.018 0.354	0.776	-0.593 -0.089 0.416	0.897	-0.579 0.04 0.659	0.613	-0.489 0.022 0.533	0.651	-0.25 0.169 0.588	0.156	-0.385 0.093 0.571	0.403
	GvT	-0.47 0.036 0.541	0.602	-0.665 -0.071 0.522	0.858	**-0.076 0.338 0.751**	**0.013**	-0.271 0.235 0.742	0.112	**-0.175 0.338 0.85**	**0.033**	-0.14 0.338 0.816	0.026
EBA	Start Ori	-0.172 0.191 0.554	0.072	-0.364 0.089 0.542	0.403	-0.476 0.004 0.485	0.699	-0.501 0.009 0.519	0.689	-0.239 0.107 0.452	0.25	-0.238 0.08 0.398	0.327
	End Ori	-0.502 0.027 0.555	0.641	-0.537 0.013 0.563	0.679	-0.661 -0.004 0.652	0.716	-0.479 -0.018 0.443	0.769	**-0.135 0.369 0.872**	0.024	-0.397 0.133 0.663	0.327
	GvT	-0.297 0.227 0.75	0.135	-0.439 0.138 0.715	0.343	**-0.17 0.267 0.703**	**0.044**	-0.439 0.107 0.652	0.403	**-0.058 0.413 0.884**	**0.009**	-0.047 0.48 1.007	0.008

RSA values for ROIs shown in Figure 6

Table 2. Values represent model correlation values for each ROI at specific timepoints. Values depict mean ± confidence interval and is written as [lower threshold mean upper threshold]. Note that the confidence intervals are Bonferroni corrected in the strict manner (critical alpha divided by 12 timepoints x 3 models x 17 ROIs). Q-statistic represents the p-value that is FDR corrected in the same strict manner. Significance was defined as a Q-value < 0.05 (see 2.10.4 Representational Similarity Analysis).

A

B

+LED

C

Start left   Start right

Grasp

Grasp
and
turn

D

Light on   "Grasp"   "Go"
            or
            "Turn"

Previous | Presentation | Plan | Execute | Inter-trial interval | Next
trial | | | | | trial

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16

Trial duration (in 2 s TRs)

A

Grasp left: trial m

Pres   Plan   Exe

Pres

Plan

Exe

Single trial RDMs (1-r)
Grasp left: trial n

0   0.5   1
Highly similar   Highly dissimilar

D

Single subject Grasp RDM (1-r)

0   0.5   1
Highly similar   Highly dissimilar

Average across
all trial pairs for
Grasp Left
and
Grasp Right

B

Turn left to right: trial m

Pres   Plan   Exe

Pres

Plan

Exe

Single trial RDMs (1-r)
Turn left to right: trial n

0 2 4 6 8 10 12 14

E

Single subject Turn RDM (1-r)

Average across
all trial pairs for
Turn Left to Right
and
Turn Right to Left

F

Turn versus Grasp

-0.1   0   0.1

Grasp trials are
more similar
than turn trials

Turn trials are
more similar
than grasp trials

Subtract condition RDMs

C

Timepoint 2   Timepoint 10

Trial m

Trial n

0 0.5 1

A

| | | Grasp | | Turn | |
| | | Left | Right | Left to right | Right to left |

Grasp Left
Grasp Right
Turn Left to right
Turn Right to left

Start Orientation

B

End Orientation

C

Turn Versus Grasp

0 — Highly similar

0.5

1 — Highly dissimilar

D

**Single subject
V1d
Time point 4**

E

**Single subject
Area-7A
Time point 12**

| | Decon time course | Grasp RDM | Turn RDM | Turn versus Grasp | RSA model over time |

Visual

V1d

cIPS

Superior parietal

7P

7A

7PC

5L

Other

aIPS

EBA

— Grasp left
···· Grasp right
— Turn left to right
···· Turn right to left

Highly similar — 0   0.5   1 — Highly dissimilar

Grasp more similar than turn — -0.1   0   0.1 — Turn more similar than grasp

— Start orientation
— End orientation
— Turn versus Grasp