LoxCode in vivo barcoding resolves epiblast clonal fate to fetal organs

Tom S. Weber[1,2]*^, Christine Biben[2,3]*, Denise C. Miles[1,2] , Stefan Glaser[4], Sara Tomei[1], Stephen Zhang[1], Patrick P. L. Tam[5,6], Samir Taoudi[2,3]#, Shalin H. Naik[1,2]#^

[1]Immunology Division, WEHI, Parkville, Melbourne. Australia. 3052
[2]Department of Medical Biology, University of Melbourne, Melbourne, Australia. 3052
[3]Epigenetics and Development Division, WEHI, Parkville, Melbourne. Australia. 3052
[4]Boehringer Ingelheim RCV GmbH & Co KG, Austria
[5]Embryology Research Unit, Children's Medical Research Institute, University of Sydney. Westmead, NSW. Australia. 2145
[6]School of Medical Sciences, Faculty of Medicine and Health, University of Sydney, Sydney NSW Australia 2006.
*Contributed equally
#Contributed equally
^Correspondence: naik.s@wehi.edu.au, weber.ts@wehi.edu.au

Abstract:

Contribution of the mammalian epiblast to fetal organs during embryogenesis has been investigated using reporters of marker genes, or through single cell or spatial RNA sequencing to infer differentiation trajectories. However, much remains to be learned about the clonal fate of mammalian epiblast cells *in vivo*. Here we develop a high diversity, high throughput, Cre recombinase-driven DNA LoxCode barcoding technology for *in vivo* clonal lineage tracing. Using this LoxCode mouse model, cells in E5.5 pre-gastrulation embryos were barcoded *in utero* and assessed in bulk via PCR or via single-cell RNA sequencing for their contribution to a comprehensive range of tissues and cell types in the E12.5 organogenesis-stage embryo. While a few typically large clones contributed to a diverse range of cell types of multiple germ layer derivatives, many clones displayed reproducible patterns of lineage restriction. Most prominent were clonal fate biases towards either blood, ectoderm lineages, mesenchymal tissues or limbs, likely reflecting branch points during development. In the context of a stochastic agent-based model of tissue development, clonal fate biases could be explained by early differentiation events occurring shortly after barcoding, and clonal similarities between tissues arose as a consequence of shared differentiation paths. At the single-cell level, clones exhibited heterogeneity in terms of tissue contributions, gene expression profiles, and in some instances left-right asymmetries and/or anterior-posterior segregation. Our study demonstrates the power and versatility of LoxCode barcoding in investigating native clonal fate and provides a deep clonal interrogation of the contribution of the mammalian epiblast to fetal organs.

Introduction:

Mammalian embryogenesis involves the generation and assembly of different cell types derived from the epiblast into a body plan that constitutes the blueprint of embryonic development. However, it is unclear whether all epiblast cells are pluripotent or whether certain cells are already restricted to certain lineages prior to gastrulation. Answering this question in the mammalian embryo has been hampered by the inability to track cells of the epiblast at single-cell resolution over a substantial timeline. Seminal studies have shed light on the early stages of mouse embryogenesis, including the production of a presumptive fate map for the epiblast embryo[1], the discovery of how spatial position of cells in the epiblast influences the acquisition of developmental fate[2], single-cell RNA sequencing and fluorescent in situ hybridization of embryonic cells and tissues at different stages during development[3-9], and the lineage tracing of single cell clones of the epiblast across the window of gastrulation[10,11]. Despite the advances made through these studies of prospective cell fate, including a more recent *in vivo* pedigree analysis[5], technical challenges remain for implementing clonal lineage analysis through both prospective and/or retrospective tracing of lineage trajectories *in vivo* across gastrulation to organogenesis[6]. As an imperative, parallel assessment of the development of multiple clones is required to answer such questions.

Technological advances in cellular barcoding have empowered the tracing of clonal lineages[12,13]. Hereby, a progenitor population of cells is tagged with unique and heritable barcodes prior to further proliferation and differentiation. Subsequent comparison of the barcode repertoire of multiple clones contributing to cellular progeny allows determination of clonal relationships, e.g., derivation from common versus separate ancestors, estimation of clone sizes, timing of lineage allocation and diversification, and spatial location of the lineage precursor. Unlike viral barcoding methods that transduce cells *ex vivo* and which are then engrafted into a recipient, recent *in vivo* technologies enable barcode generation in cells in their native environment. This feature offers the ability to investigate clonal biology in a *native* setting. These barcoding techniques encompass prospective (multispectral barcoding[14], *in vivo* viral delivery[15], transposons[16,17], CRISPR/Cas9[5,18-22], Cre/Lox recombination[23,24]) and retrospective (natural barcodes[25], cell history recorder[26,27]) approaches. Many also allow the simultaneous assessment of the transcriptome and clonal origin of single cells for leveraging the ability to link cell state to cell fate, or to determine shared molecular features among clones[28].

Previously, we established the foundational design principles for a Cre/Lox-based barcode generation system and mathematically determined that the arrangement of distinct DNA elements flanked by LoxP sites in alternating directions would lead to optimal barcode diversity[29]. Notably, our findings indicated that the number of theoretical barcodes would increase combinatorially with the addition of each barcode element to the cassette. For example, a configuration comprising 14 LoxP sites and 13 code elements could potentially produce a barcode diversity exceeding 30 billion upon partial recombination (*Methods*). Furthermore, our design incorporated a novel aspect: utilizing DNA elements of minimal length to create a compact construct. This feature facilitated efficient recombination at low Cre concentrations, generated over 2,000 size-stable barcodes resistant to further Cre excisions and ensured, in contrast to established Cre based barcoding systems[23], compatibility with high-throughput short-read sequencing technologies.

Here we describe how LoxCode clonal lineage tracing offers several critical attributes that permit resolution of clonal pattern of cell/tissue lineages. Upon LoxCode generation in the E5.5 epiblast and LoxCode extraction from tissue of E12.5 embryo either through PCR or single cell RNA-seq, and applying a suite of novel computational

methods, we have established i) the contribution of epiblast cellular descendants to a full range of embryonic tissues, ii) the inter-relationships of fetal tissues and organs, and iii) an appraisal of cell types derived from individual epiblast cells, which informed iv) cell type relationships at a clonal level. In parallel to the present study, we have applied LoxCode clonal lineage tracing to investigate lineage relationships during yolk sac haematopoiesis[30].

RESULTS:

LoxCode design and experimental setup

Based on optimal design principles for Cre/LoxP cellular barcoding[29], the LoxCode was constructed through serial integration of LoxP/Barcode paired inserts to generate a final cassette of 14 LoxP sites of alternating orientation flanking 13 short barcode elements of 8 bp and 14 bp (Fig. 1a, *Methods*). Upon Cre exposure, random recombination between the LoxP sites led to inversions and excisions, resulting in DNA sequence reshuffling with a theoretical DNA diversity of over 30 billion barcodes (Methods and Table 1). The LoxCode construct was introduced into the *Rosa26* safe harbor locus in C57BL/6 mice to enable *in vivo* lineage tracing though both PCR-based detection in genomic DNA as well as read-out of LoxCodes transcripts driven by the endogenous *Rosa26* promoter.

Figure 1. LoxCode design and experimental setup. a) The LoxCode cassette is composed of 14 LoxP sites in alternating orientation flanking 13 small (8-14 nucleotides) barcode elements indexed from 1 to 13. Cre exposure leads to random excisions and inversions, and different sized bands can be analysed via gel electrophoresis. Hypothetical LoxCodes of different sizes with inverted barcode elements depicted upside down. To annotate barcodes, elements are listed in order, with a plus or minus sign indicating if the orientation matches or is inverted compared to the original sequence. With this notation, the original cassette is written as '+1+2+3+4+5+6+7+8+9+10+11+12+13', and the first recombined example in panel a) is annotated as '+9+10-5-4-1-8-3-2-7-6+11+12+13' where inverted LoxCode elements are denoted by a minus (-) symbol. Computed theoretical barcode diversities for each

barcode size (containing 13, 9, 7, 5, 3, 1 elements) are provided (right column). Experimental design: **b)** Barcoding of all cells (epiblast
cells are highlighted in colour) in E5.5 pre-gastrulation embryos was activated by administration of 4-hydroxytamoxifen (4-OHT) to
the pregnant mice, followed by dissection of E12.5 embryos into **c)** 40 tissues and organs for barcode retrieval and single-cell RNA
sequencing analysis.

To study the clonal fate of E5.5 embryonic cells *in vivo*, LoxCode homozygous females were crossed to *Rosa26*-
Cre-ER^{T2} homozygous males to produce heterozygous LoxCode/*Rosa26*-Cre-ER^{T2} embryos (Fig. 1b). LoxCode
generation was induced at E5.5 via intravenous injection (IV) of low doses (50-100 µg) of 4-Hydroxytamoxifen
(4-OHT) to both minimise developmental toxicity (all embryos used were of stage, normal, and healthy) and to
allow partial LoxCode recombination – the latter attribute enhances barcode diversity and complexity. To assess
the clonal fate of E5.5 embryonic cells, E12.5 embryos were dissected into 40 tissues and organs (Fig. 1c) which
were either: i) lysed, prior to barcode PCR and sequencing (Fig. 2, 3); or ii) dissociated into single cell suspensions
for concurrent scRNA-seq and barcode detection (Fig. 5).

While barcoding of the E5.5 embryo using *Rosa26*-Cre-ER^{T2} would not be limited to the epiblast cells, prior
knowledge of lineage development during embryogenesis indicates that the extraembryonic ectoderm has no or
little contribution to the germ layer tissues while the visceral endoderm may contribute to the endoderm of the
gastrulating embryo[31,32]. Therefore, we can assume that most clonal descendants in the embryo during
organogenesis are derived from epiblast cells. Accordingly, we directed our analysis of cell clones in the late-
organogenesis stage of the embryo.

Epiblast clonal fate heterogeneity

From three embryos, we recovered a total of 703 barcodes present in both PCR replicates of each tissue (~200-
300 barcodes per embryo) of all expected barcode sizes with a minimum number of recombination events
(excisions and/or inversions) for each barcode ranging from one up to eight (Supplementary Fig. 1 a-b). After
removal of frequent barcodes that may arise from more likely recombination events ('repeat use' or 'barcode
collisions') using a new empirical method of filtering (see Methods) ~30% of barcodes remained, with an overlap
of 10 barcodes between any two embryos, and only 1% (5 out of 265) present in all three embryos (Fig. 2a).
Furthermore, clones contributing to at least two tissues represented one third of those barcodes, with very little
overlap between embryos. This indicated efficient enrichment of *bona fide* single-use barcodes (Fig. 2a).

In a pluripotent model where every epiblast clone contributes broadly to every cell/tissue type, albeit with varying
clone size, one would expect every barcode to be detected in all 40 different tissues or body parts. While we
detected a few, typically large, multi-lineage clones in each embryo (~5% of all clones), this was not the rule:
allocation of E5.5 epiblast cells ranged from only ~3.5% clones towards the gut up to a maximum of ~20% of all
clones towards the brain (Fig. 2b). A minority of clones were restricted to a single tissue type which ranged from
1-3% of all clones for most tissues. Strikingly, a significant fraction (12%) of LoxCodes were present exclusively
in the circulating blood cells, which at E12.5 is predominantly extra-embryonic yolk sac-derived erythroid cells[33],
indicating particularly early fate restriction (Fig. 2c). This is in line with previous studies[34,35].

Figure 2. Patterns of epiblast clonal fate to organs and tissues. a) LoxCodes were excluded based on empirical frequency (see Methods) resulting in few overlapping barcodes between biological replicates. **b)** Proportion of clones contributing to indicated tissue types. **c)** Proportion of clones uniquely detected in the indicated tissues. **d)** Comparison of the clone size and number of tissues each LoxCode was detected in, versus a simulation of random sampling of cells from all tissues of the corresponding clone sizes. Histograms on the right show the number of clones and average size for clones with more than 1000 cells. **e)** UMAP projection of clonal outcomes with 5 clusters of clonal output. **f)** Bubble plot/binary heatmap of clonal contribution to embryonic tissues. LoxCodes (columns) are ordered by 1D UMAP according to patterns of their clonal output to the indicated tissues (rows) from 3 embryos. Darkened rectangles indicate all detected clones (binary visualisation), and circles inside are proportional to clonal output to the indicated tissues (quantitative visualisation). Coloured backgrounds (six) correspond to the cluster annotations from e). **g)** Examples of clonal fate in a pseudo-anatomical map, where circles are scaled proportional to the logarithm of frequency of the indicated LoxCode and preceded by embryo of origin e1, e2, e3, respectively).

Overall, the majority of LoxCodes of varying clone sizes (spanning three orders of magnitude) were unequally detected in a restricted number of tissues (Fig. 2d), inconsistent with a scenario in which single-cell epiblast fate is homogeneous with contributions to all tissues. This observation was further corroborated by performing

simulations where cell numbers equivalent to the empirically derived clone sizes were drawn randomly from the distribution of cells across all 40 tissues (Fig. 2d, Kolmogorov–Smirnov, p-value<0.001).

Next, we explored whether, from the large number of possible clonal fates, a limited set of reproducible fate patterns were realised *in vivo*. For this, epiblast clones detected in at least two tissues were classified by Leiden clustering[36] into five main clusters of fate according to their contribution to 11 tissue types (Fig. 2b) and projected onto 2 dimensions by repurposing the Uniform Manifold Approximation and Projection (UMAP[37,38]) algorithm (Fig. 2e) and summarized in a bubble plot (Fig. 2f). We observed neural tissue-biased (cluster 1), multi-potent (cluster 2), limb-biased (cluster 3), and blood-biased (cluster 5) clones. Cluster 4 clones contributed to many tissues, but less to splanchnopleure and intermediate mesoderm derivatives (organs). Examples of Cluster 1-5 clonal contributions are shown (Fig. 2g).

Taken together, this analysis indicates that there were a small set of reproducible clonal fate patterns of E5.5 epiblast cells and that, within multi-lineage clones, strong biases were found among certain tissues, most notably between neural versus limb-biased multi-outcome clones (Fig. 2e-g).

Fetal organ lineage branch point hierarchies informed by clonal relatedness of tissues

The fate of clones can differ both in terms of the type of tissues they generate and the number of cells they contribute to in each tissue (Fig. 2). We reasoned that by comparing these clone-related features between pairs of tissues, it would be possible to determine the ancestral relationships of tissues and infer the developmental processes that underlie them. Within this concept, when two tissues have a similar clonal composition, they likely have a shared developmental history with a 'late' branching point, whereas dissimilar clonal compositions may reflect an 'early' branching point or late-stage clonal expansion.

As a proof-of-principle, we first assessed similarities between clonal frequencies (using the cosine similarity as a measure) in a selection of samples that are well studied in terms of segregation during development, namely blood, brain (fore- and hindbrain, left and right), limbs (forelimbs and hindlimbs, left and right), and gonad and kidneys. We found that samples of the same tissue type were more similar in their clonal composition compared to samples originating from a different tissue (e.g. brain tissues vs limbs), likely reflecting developmental branching between the tissue types (Fig. 3a). Unfortunately, clustering was relatively noisy both on individual embryos and for all three embryos pooled probably due to insufficient numbers of barcodes (Fig. 3a).

We noted however that the clustering of tissue types became more robust when all barcodes (including likely 'barcode collisions' otherwise excluded due to high frequency[39], see Methods) were incorporated for computing the cosine similarity scaled to range from 0 to 1 (Fig. 3b). Because frequent barcodes may correspond to several individual clones harbouring the same barcode by chance, i.e., what we refer to as *meta-clones*, we termed this approach **m**eta-**c**lonal **s**imilarity **a**nalysis (MCSA). To quantify the power of MCSA more systematically we compared branching hierarchies derived from filtered and unfiltered barcodes and from individual embryos and pooled against each other using the Robinson-Foulds similarity, a commonly used measure for the similarity between trees. In this comparison we also included a hierarchy, based on results from Fig 3b and prior knowledge from literature. This resulted in a hierarchy consisting of six sequential bifurcation points: initiated with early branching of circulating nucleated blood cells, followed by ectoderm and mesoderm specification, paraxial/lateral plate mesoderm and intermediate mesoderm, then anterior posterior, then left right, and finally proximal distal

branching (Fig. 3 c-d). This analysis highlighted that hierarchies derived from unfiltered barcodes were, despite the risk of barcode collisions, more consistent between individual embryos and the pool and were more similar to the model hierarchy in Fig. 3c.

The fact that stringent filtering, which is the norm in *in vivo* barcoding data analyses[20,23,24], lead to less robust results was counterintuitive initially due to technical artifacts potentially introduced by barcode collisions. However, we speculated that a trade-off exists between the number of clones that are available to compute the similarity and the chance that some barcodes may represent more than a single clone. Next, we applied MCSA to the full data set, and utilized 3 complementary approaches to visualize the results: through a hierarchically clustered heatmap (Fig. 3e), through UMAP to project tissues (rather than individual clones as in Fig 2e) in 2D and connect MCSA-based nearest neighbours (Fig. 3f), and a pseudo-anatomical representation of tissue interconnected using edge bundling of nearest neighbours and split by a reduced set of tissue classes (Methods) (Fig. 3g). For completeness we also provide the heatmap using *bona-fide* clones in Supplementary Fig. 2a with the same ordering as in Fig 3e.

All three visualizations revealed similar groupings, consisting of neural clusters, limb clusters, mesenchymal and organ cluster, and blood clearly isolated from the other tissues. However, there were also differences. While the heatmap separated splanchnopleure/extraembryonic mesoderm from neurectoderm/mesoderm derivatives, the UMAP suggested a more global neurectoderm/mesoderm separation, with splanchnopleural tissues closely associated with paraxial mesoderm derived tissues. Due to its robustness to noise and more flexible 2D layout rather than a linear ordering, the UMAP representation is more amenable for visualizing complex tissue relationships accurately.

The resulting data, amenable through *in vivo* LoxCode clonal lineage tracing, determined a comprehensive series of highly correlated and uncorrelated tissue groupings (Fig. 3e-f). These included:

(a) a brain tissues/neural tube/tail/face group that represent tissues of a neuroectoderm origin;

(b) limbs/neural tube/lateral mesoderm/medial mesoderm that represent tissues derived from mesoderm;

(c) gonad/kidney/gut containing derivatives of intermediate/lateral mesoderm.

These global groupings putatively demonstrate which tissue/organ groups have shared/similar clonal ancestry. Tissue groups that were less correlated such as liver/blood, brain/kidneys and gonad/lung were suggestive of distinct ancestry. Examples of the least correlated samples were:

(d) endoderm containing tissues (liver and lungs) that were distinct from mesodermal and neuroectodermal tissues, but are associated with gut and stomach;

(e) anterior neuroectodermal derivatives (forebrain and midbrain) were clustered separately from the neural tube. The separate clustering of these two groups may be due to the distinct origin of anterior neuroectoderm derived from the epiblast and the trunk neural tube derived from the neuromesodermal progenitor[11];

(f) neural tissues and limbs displayed distinct clonal composition;

(g) fore- and hindlimb displayed separate clonal origin and sidedness of clones in the limb tissues, possibly related to the separation of progenitors to contralateral lateral plate mesoderm.

Figure 3. Major developmental branch points established through E12.5 metaclonal similarity analysis. Clustered heatmap of cosine similarity matrix computed for a selection of tissues pooled from three embryos using **a)** bona-fide and **b)** all clones (meta-clonal similarity analysis, MCSA, see text and Methods). **c)** Differentiation hierarchy of tissue development comprising 6 sequential levels of bifurcation. **d)** Robinson-Foulds similarity between hierarchies derived from bona-fide and all barcodes. **e)** MCSA of all tissues visualised as a heatmap, **f)** UMAP with lines connecting to 4 nearest neighbours with an MCS >0.75, or **g)** a pseudo-anatomical map where tissues of a given type (legend in a), were connected to their 4 nearest neighbours with an MCS >0.75. Edge colour indicates the tissue type of the outgoing node per legend in e). Data represent the aggregation of 3 biological replicates that were also assessed individually (Supplementary Fig. 3). Abbreviations: Pxm, paraxial mesoderm; Lpm, lateral plate mesoderm; Intm, intermediate mesoderm.

To explore potential aetiologies of the clonal similarities between samples observed in our data, we implemented a generative stochastic agent-based model of tissue development (see *Methods* and Supplementary File 1). Starting from the zygote, we simulated stochastic cell divisions[40], barcoding at E5.5, as well as differentiation into a selection of tissues following the lineage rules informed by Fig. 3c (Fig. 4a). At day E5.5, cells (i.e., agents) were assigned a LoxCode with probabilities based on a stochastic recombination process[40]. Except for the first two divisions and blood progenitors (see *Methods*), division times were assumed to be 12 hours in average yielding cell numbers at E5.5 and E12.5 similar to previous estimates[3]. At day E12.5, with a simulated population size of typically over 6 million cells, clonal data from three simulated embryos (to match the actual data) was

used to generate an MCSA matrix. These simulations were then run thousands of times to fit the model output to the empirical data using the evolutionary covariance adaptation strategy (CMA-ES, Fig. 4b, Supplementary Fig. 4, [41]). A total of 33 parameters were allowed to vary including the timings t of fate bifurcations, p the probabilities to differentiate into a specific cell type at a junction, and the rate of fate determination.

Despite its simplicity and absence of spatial aspects, we found that the model could reproduce well not only clonal similarities summarized in the MCSA heatmap (Fig. 4c) but also fate biases at the clonal level (Supplementary Fig. 4a), which importantly were not used for model fitting. The blood-epiblast bifurcation was inferred to occur shortly after barcoding at E5.5 followed by the ectoderm-mesoderm split approximately 24 hours later (Fig. 3d). All remaining bifurcations were completed by the end of gastrulation (i.e., E9.5). Best-fit parameters from different random initialization showed a relatively narrow range indicating the parameters were well defined given the model and the data (Supplementary Fig. 4b-c). To visualize the full pedigree, which due to the large number of cells is impractical with ordinary hierarchical trees, we positioned cells on concentric circles according to generation, ordered within and between circles such that mother cells neighbour their daughter cells, and coloured according to the cell's state at division (Fig. 3e). Using a similar approach, we then visualized E5.5 clones (Fig. 4f), illustrating how fate diversity is generated at the clonal level. In summary, fate biases, in this model, were simulated to be consistent with early differentiation events that occur shortly after barcoding, while clonal similarities between tissues arise as a consequence of shared differentiation paths.

Overall, this analysis by MCSA suggested that a putative hierarchical model of developmental branch points and timings can be inferred from a single snapshot of barcode induction. In the future, incorporation of clonal data at additional time points could be used to refine the model and empower the discovery of so far uncharacterized branch points for elucidating the order and timing of each bifurcation.

Determination of cell type bias and clonal relationships at a single cell level

Bulk barcode analysis has the advantage of examining millions of cells, unlike single-cell analysis, which typically studies thousands due to cost and throughput limitations. This allows for a more comprehensive view of clones, including the detection of rare clones that might be missed with single-cell omics analysis. However, bulk barcode analysis doesn't reveal which cell types contain which barcodes within a tissue. As the LoxCode construct is actively transcribed under the endogenous Rosa26 promoter, combined with single-cell RNA-sequencing, it enables simultaneous identification of both cell type and clonal origin. In this study, we dissected E12.5 embryos into organs/tissues derived from an E5.5 barcoded embryo as outlined in Fig 1b. From this, single cell suspensions were generated, multiplexed using MultiSeq[42] (Methods, Supplementary Fig. 5a), and processed on a droplet-based scRNAseq platform to capture ~30,000 cells. Following quality control (Methods), 33 cell types (Fig. 5a) across all tissues (Fig. 5b, Supplementary Fig. 5b) were annotated (Methods, Supplementary File 1).

Figure 4. *In silico* **embryogenesis. a)** Agent-based model of tissue development incorporating differentiation rules in Fig. 3c. Model progenitor nomenclature introduced for illustrative purposes. **b)** Schematic of a realization of the model at the cellular level, including stochastic timings between divisions, differentiation and barcoding. Parameters are fitted iteratively using the CMA-ES evolutionary algorithm. **C)** Best fit MCSA matrix to empirical data shown in Fig. 3 panel b). **d)** Simulated kinetics of tissue differentiation from day of conception to E12.5 with frequencies on linear and population size on logarithmic scale. **e)** Nested-zoom into a simulated 22 generation pedigree, from the zygote to E12.5. Concentric circles correspond to generations (the centre is generation 0 and 1), each segment represents one cell coloured according to its cell state at division (legend, panel a). Two daughter cells are placed adjacent to the mother but on the next outer concentric circle (see *Methods*). f) Examples of simulated clonal pedigrees from single cells at generation 10, corresponding to the time of experimental LoxCode induction at ~E5.5. The binary string next to each example indicates the ancestry of the founder cell of that clone (see *Circular Pedigree plot* in *Methods*).

Full length barcodes were then selectively amplified from the single-cell full length cDNA libraries prior to fragmentation of the cDNA library and both were sequenced separately on a NextSeq (for cDNA) and MiSeq (for LoxCodes) and mapped *in silico* by virtue of their shared 10X cell barcode. Full length LoxCode capture efficiency was 27% (>8500 cells), but we noted recovery as low as 6% in neuronal tissues. After filtering out

frequent LoxCodes as in Fig 2, we were left with ~150 cells, spanning 20 bona-fide clones, where 5 of them contained >8 cells.

Cellular contribution of these 5 clones indicated contribution to a large range of cell types, but also biased distributions across tissues (p-value < 0.0001) and cell types (p-value < 0.0001) (Fig. 5c-d) At the tissue level, clone "+1+8-13" was enriched in lungs, gut and mesenchymal tissues, clone "+1-6+9" was prominent in fore- and hindlimbs, with low detection frequency across all organs. Clone "+1-6+9", "+1+4+5+2+3", and "+1+10-5-4-13" were detected in face, but only for the latter two were present in brain. At the transcriptional level, all 5 clones had representatives in the pulmonary/gastrointestinal mesenchyme and chondrocytes. Clone "+1+4+5+2+3" contributed to a diverse set of neuronal cells, while clone "+1-6+9" differentiated mostly to musculoskeletal mesenchyme, chondrocyte and fibroblast cells consistent with its limb-biased nature.

Overall, despite the low number of clones that could be assessed, heterogeneity in transcriptional fates was clearly evident. To explore if meta-clones could provide additional insights, we repeated the analysis for barcodes for which we were less confident that they originated from a single cell. First, we confirmed that tissue type contribution deviated from the population baseline (Fig. 5e). Bias towards forelimbs (light-brown color) or hindlimbs (dark-brown color) was apparent for meta-clone "+1+2+3+4+5" and "-7-2+13" respectively. Relative to baseline, meta-clone "+1+6+7+8+13" showed increased presence in mesodermal organs, coinciding with a low detection in endoderm organs such as lungs, liver and gut. Meta-clone "+1+4+13" was not detected in limbs but in blood (1 cell), although this was not statistically significant. Interestingly most meta-clones were detected in gut. Among meta-clones we identified 5 additional candidates with apparent transcriptional bias (Supplementary Fig. 6a-c). While we cannot directly comment on the fate of these clones (i.e., they could represent the combined fate of two or more epiblast cells), their heterogeneity adds support to the previous observations (Fig 3) that most E5.5 epiblast clones are biased in their fate.

Global relationships between cell types were analysed using a single cell version of MCSA (scMCSA). In this approach meta-clonal similarities between transcriptional cluster (instead of tissues as in Fig. 3f) were computed and projected onto 2D using UMAP (Fig. 5f). Cell types broadly clustered into 1) haematopoietic and cardiac, 2) neuronal, and 3) mesenchymal types. Strong relatedness between haematopoietic and cardiac tissues was consistent with the notion that mesodermal cells that contribute to both tissues, are the first to be allocated during gastrulation, and might share common epiblast ancestors.

Comparing single-cell clonal data from left and right sides, and anterior and posterior compartments of the embryo allowed identification of asymmetries along the left-right and anterior-posterior body axis. Some clones displayed asymmetrical distribution to fore- and hindlimbs, and gonad and kidney (Fig. 5g), while other clones showed no significant spatial bias (Supplementary Fig. 7). Importantly, instances of asymmetrical contribution to limbs, gonads, and kidney were also detected in the bulk tissue analysis (Fig. 2f).

In sum, through LoxCode barcoding of the mouse epiblast cells followed by barcode detection and analyses of the clonal contribution to tissues and cell types, we have uncovered the patterns of contribution of clones to the tissues of the late-organogenesis-stage embryo and the inter-relationship of fetal tissues and organs in terms of their clonal development.

Figure 5. Single cell readout of epiblast fate bias and clonal relationships. a) UMAP of scRNA-seq derived from ~30,000 cells and annotated cell types, with **b)** their distribution across tissues (left and right and mesenchymal tissues pooled). **c)** Examples of single cell bona-fide clonal output corresponding to the indicated LoxCodes. Distribution of single cells per detected LoxCode in c) according to **d)** tissue (top) or cell type (bottom). **e)** Tissue type distribution of meta-clones with at least 20 cells and ordered using 1D UMAP. * indicates barcodes where distribution significantly deviates from a base-line distribution (χ2 test, adj. p-value < 0.05 using Benjamini-Hochberg correction). **f)** scMCSA UMAP representation of cell types from b), with lines connecting each cell type's 3 nearest neighbours. Dot size is proportional to the cell number in each cluster, and colouring is according to broad cell type categories as indicated in the legend. **g)** Left-right, anterior/posterior and tissue asymmetry of clonal contribution (χ2 test, adj. p-value < 0.05 using Benjamini-Hochberg correction).

Discussion

Using the LoxCode mouse model presented herein, we find pre-gastrulation epiblast cells are biased in their clonal contributions across tissues and cell types in the germ layer derivatives: notably towards either blood, ectoderm lineages, mesenchymal tissues, or limbs, likely reflecting developmental branch points during development. How do these developmental branch points occur? Our simulated clonal pedigrees infer that fate restriction commences around E5.5 and completes by E9.5. However, presently, our results do not allow discrimination between several non-mutually exclusive aetiologies: epiblast molecular fate imprinting[43], multi-lineage epiblasts that, by virtue of their physical location, receive specific signals to later restrict their fate[44], lineage specific timing of allocation to specific cell/tissue type during development[45], and late-stage clonal expansion of the specified progenitor or the descendant tissue/cell types[46].

The notion of fate bias is well-established within the field of embryology, particularly during the initial phases of development stages[47,48]. In contrast to a binary approach that classifies cells according to their capacity to differentiate into a specific assortment of mature types (e.g., bi-potent, tri-potent), a framework of fate bias presents a more refined evaluation of relative contributions. This concept is key to the latest generation of revised models of hematopoiesis[49] that, instead of rigid binary bifurcations, favour a 'continuous' model and acknowledge the existence of multi- and oligopotent progenitors exhibiting varying degrees of lineage-bias early in the stem and progenitor cell hierarchy. Throughout development, each stage is characterized not only by the current cellular state but also by its origins and future differentiation potential. The LoxCode mouse model and the accompanying analytical tools introduced here now offer a higher-throughput approach to quantify these fate biases and their progression not only for embryogenesis, but also in other systems like hematopoiesis, immunology and cancer.

The key features of the LoxCode mouse that empower this approach include barcode generation that is expeditious (within 6-12 hrs of Cre-ER^{T2} activity most barcodes recombine at least once), using low-dose tamoxifen induction, and efficient (a stable identity and high diversity of >30 billion LoxCodes, >2000 size-stable barcodes resistant to further Cre recombination), with barcodes detectable through PCR or single cell RNA-seq. This model allows a significantly higher throughput compared to existing Cre LoxP based barcoding[23,24] by using short read sequencing for barcode read-out.

An important next challenge will be to connect fate bias across different stages of development. Pioneering technologies including evolvable hgRNA barcodes[19] already address this question, but are currently hampered by i) the non-randomness of NHEJ outcomes[50] which limits the diversity of founder cells, and ii) the lack of rapidly inducible Cas9 mouse models. Crossing the LoxCode mice with the MARC1 system or CARLIN[20] or other CRISPR based barcoding technologies could represent a powerful avenue to precisely map fate bias kinetics over time, by combining the best of Cre LoxP (i.e., efficiency, temporal control, high baseline diversity) with Cas9 based approaches (i.e., accumulation of mutations, exponential increase in diversity as a clone expands). In addition, constructs using alternative site-specific recombinases or integrase (e.g., Flp, Bxb1), or emerging technologies like transcriptional cell history recorders[26], could provide additional levels of resolution and the recording of molecular correlates of branching.

 While we are far away from replicating the notable work in C. elegans[51] for mouse development[52], the current study provides a comprehensive view into clonal fate heterogeneity of E5.5 epiblast in murine embryos and demonstrates the LoxCode technology as a powerful tool to characterize clonal fate at exceptional depth and higher throughput.

Methods:

LoxCode cassette restriction cloning:

Following multiple unsuccessful attempts to synthesize the LoxCode cassette through commercial service providers, presumably due to the palindromic characteristics of LoxP repeats, we pursued an alternative approach involving the assembly of the LoxCode cassette via sequential restriction cloning. To this end, we designed and procured two partially reverse complementary 5' phosphorylated 90 base pair oligonucleotide libraries (Integrated DNA Technologies, duplex oligos), which encompassed two LoxP sites in opposing orientations (underscored), an intervening 8-base sequence consisting of mixed bases (Ns), and flanking regions comprising 4 mixed bases. Additionally, each oligonucleotide contained a 5-base extension at the 5' terminus, facilitating the formation of sticky 5'-AATT-3' overhangs upon annealing, thereby rendering the duplex oligo compatible with ligation into EcoRI-digested plasmids. The nucleotide base adjacent to the sticky end was selectively designated as cytosine (C) and adenine (A) (highlighted in bold), ensuring that the EcoRI (GAATTC) recognition site would be reconstituted solely at one extremity following ligation, thereby permitting sequential integrations, i.e.

```
5'-AATTCNNNNATAACTTCGTATAATGTATGCTATACGAAGTTATNNNNNNNNATAACTTCGTATAGCATACATTATACGAAGTTATNNNNA-3'
     3'-GNNNNTATTGAAGCATATTACATACGATATGCTTCAATANNNNNNNNTATTGAAGCATATCGTATGTAATATGCTTCAATANNNNTTTAA-5'
```

Subsequently, the pUC18 plasmid was subjected to digestion with the EcoRI-HF restriction enzyme (New England Biolabs), followed by a standard ligation reaction. In order to minimize the likelihood of background recombination in bacterial cells, Sure 2 (Agilent) competent cells were transformed with the ligation product and cultured at reduced temperatures (27°C). Individual colonies were then screened for integration events and orientation via restriction digest analysis. Plasmids isolated from successful colonies, exhibiting the reconstituted EcoRI site at the 3' terminus of the cassette, underwent a subsequent round of EcoRI digestion for further cloning. This procedure was iteratively performed seven times, ultimately yielding a nucleotide sequence encompassing a total of 14 LoxP sites and 13 random barcode elements, with either 8 (odd position, NNNNNNNN) or 14 base pairs (even position, NNNNGAATTTNNNN).

```
ATAACTTCGTATAATGTATGCTATACGAAGTTATACTCCGCAATAACTTCGTATAGCATACATTATACGAAGTTATTCCAGAATTTGTATATAACTTCGTATAATGTATGCT
ATACGAAGTTATACATCCACATAACTTCGTATAGCATACATTATACGAAGTTATAAAGGAATTTCTCCATAACTTCGTATAATGTATGCTATACGAAGTTATATTTCCTCAT
AACTTCGTATAGCATACATTATACGAAGTTATGCCCGAATTTTTTCATAACTTCGTATAATGTATGCTATACGAAGTTATGCTACTGGATAACTTCGTATAGCATACATTAT
ACGAAGTTATATGAGAATTTATGGATAACTTCGTATAATGTATGCTATACGAAGTTATAACTAGAAATAACTTCGTATAGCATACATTATACGAAGTTATTGCAGAATTTCC
TCATAACTTCGTATAATGTATGCTATACGAAGTTATCGACACTTATAACTTCGTATAGCATACATTATACGAAGTTATAACGGAATTTTCAAATAACTTCGTATAATGTATG
CTATACGAAGTTATCGTGTTTGATAACTTCGTATAGCATACATTATACGAAGTTAT
```

LoxCode mouse generation:

The LoxCode mouse line was generated following a previously established protocol[53]. In brief, double strand breaks were introduced at the *Rosa26* locus in C57BL/6J oocytes via microinjection of Cas9-gRNA ribonucleoproteins (gRNA: 5'-CTCCAGTCTTTCTAGAAGAT -3'), co-injected with a circular HDR donor vector (pBlueScriptIISK backbone) containing a splice acceptor, the LoxCode sequence, and a polyA signal flanked by 1000-1500bp Rosa26 specific homology arms. Oocytes were transplanted into pseudopregnant females, and 2 out of 26 viable pups displayed the expected integration via PCR and gel electrophoresis. Sanger sequencing was performed to confirm the integrity of the LoxCode sequence. The LoxCode line was then bred to homozygosity on a C57BL/6 background.

LoxCode theoretical and practical diversity:

The LoxCode construct consists of 13 barcode elements, with 7 at odd positions (e.g., 1, 3, 5, 7, etc.) and 6 at even positions (e.g., 2, 4, …, 12). Recombination results in inversions and excisions that can alter the locations and orientations of individual barcode elements. However, by design, odd and even barcode elements always maintain their respective positions. We have previously demonstrated, through mathematical reasoning, that all barcodes can be generated via a series of inversions and excisions[29]. As a result, the theoretical diversity of a cassette containing n odd barcode elements and m = n - 1 even barcode elements can be calculated by enumerating all permutations and considering both forward and reverse orientations:

$$\frac{7!\,6!}{(7-n)!\,(6-m)!}2^{n+m}$$

Summing this formula for barcodes of size 1, 3, 5, 7, 9, and 13 yields a total of 30,204,722,030, or over 30 billion theoretical barcodes. It is important to note that barcodes with 11 elements are not expected to be generated due to the DNA size constraint of Cre recombination, although we detected them at low frequency.

# recombinations	1	3	5	7	9	13
0	0	0	0	0	0	1
1	2	4	6	8	10	37
2	7	52	106	162	253	912
3	14	360	1212	2378	4418	19812
4	14	1506	11333	31928	70538	317578
5	14	2006	56242	252490	728021	3988251
6	14	2016	154973	1403382	5643564	36058157
7	14	2016	199695	4789612	29519488	238462747
8	14	2016	201587	9951050	106581710	1149595506
9	14	2016	201600	12504942	256471281	4056050952
10	14	2016	201600	12889288	409271561	10401368516
11	14	2016	201600	12902328	461386076	19356268864
12	14	2016	201600	12902400	464458544	26701414952
13	14	2016	201600	12902400	464486400	29495220993
14	14	2016	201600	12902400	464486400	29724820834
15	14	2016	201600	12902400	464486400	29727129600

Table 1. Theoretical barcode diversity for barcodes with 1 to 13 elements (columns) per number of recombination events (rows).

The barcode diversity generated in an experiment depends on the number of recombination events (see Table 1), which are influenced by Cre activity. Insufficient Cre results in limited barcode diversity, while excessive Cre activity leads to an accumulation of less diverse barcodes with fewer elements. Theoretically, all barcodes can be reached with 15 recombination events (see Table 1); however, barcodes with fewer elements typically achieve full diversity sooner. For example, generating all possible 201,600 five-element barcodes requires nine recombinations, while five recombinations are sufficient to produce all possible 3-element barcodes.

The necessary barcode diversity for accurate lineage tracing is determined by the number of clones tracked simultaneously. Generally, we recommend maintaining a 100-fold excess of barcodes relative to founding cells, which for the E5.5 is approximately 20,000 barcodes or at least three recombination events per cell. We typically achieve a mean of around four recombinations with a maximum

450 of eight or nine. Across all our experiments to date (including those outside this work), we have identified over 125,000 unique barcodes,
451 with a maximum of 11 recombination events (e.g., -3+12-9-10+7+4-11+8+5).
452
453 In addition to theoretical barcode diversity, it is crucial to consider uneven probabilities. Some barcodes are much more likely to be
454 generated than others, which reduces effective diversity as well as the proportion of cells that can be confidently tracked within a given
455 population.
456
457
4-OH-Tamoxifen preparation and administration:
459
460 4-Hydroxytamoxifen Z-isomer (4-OHT, Sigma) was dissolved in 100% EtOH, diluted 1:1 with Kolliphor and stored in aliquots at –20
461 °C. On the day of the injection the dissolved 4-OHT was further diluted in PBS (50 µg/100 µl) and intravenously injected according to
462 a procedure developed in Chevalier *et al.*[54]. Pregnant females received 50-100 µg of 4-OHT via tail vein injection 5 days after a plug
463 was detected. Injected mice were separated from untreated females and monitored regularly. All animal experiments were approved by
464 The Walter and Eliza Hall Institute animal ethics committee.
465
466 **Embryo dissections:**
467
Rosa26CreERT2/Rosa26CreERT2 and *LoxCode/LoxCode* or *LoxCode/+* mice were set up and plug checked daily. Noon of the day of
plug was referenced as 0.5 embryonic day (E0.5). Embryos were dissected in their yolk sac at E12.5 in PBS with 7% Foetal Calf Serum.
The yolk sac was used as a genotyping sample when necessary (*LoxCode/+* mother). Embryos were staged using morphological criteria
and assessed for normal development. The head was set aside and spilling blood was collected (blood sample). Heart, right and left lung
lobes, liver, stomach (with spleen and pancreatic buds attached), intestine, left and right kidneys, left and right gonads, the region
containing the dorsal aorta at the level of the aorta-gonad-mesonephros (AGM) area were dissected next. The rest of the body was cut
transversely below the forelimb buds. From the anterior segment (forelimb level or level 1), left and right forelimb buds (termed L hand,
R hand) were removed, followed by the left and right arms (L arm, R arm). The neural tube was removed (neural T I segment).
Mesenchymal tissues were dissected as medial (left and right medial mesenchyme: L mid mes I and R mid mes I, encompassing somites)
and lateral (left and right lateral mesenchyme: L lat I and R lat I, encompassing everything lateral/medial beyond the somites). The rest
of the body was then cut just above the hindlimb buds level (between limb buds level or level 2), and corresponding neurectodermal or
mesenchymal segments (neural T II, L mid mes II, R mid mes II, L lat II, R lat II) were dissected as above. The tail was removed and
the third axial level (hindbud level or level 3) was dissected as per - level 1 (left and right hindlimbs (termed L foot, R foot), left and
right hips (termed L leg, R leg), neural T III, L mid mes III, R mid mes III, L lat III, R lat III). The facial mesenchyme (face) was
removed from the head, leaving the neurectoderm behind. Forebrain (brain I), midbrain (brain II) and hindbrain (brain III) segments
were dissected according to morphological landmarks and halved along the midline (L brain I, R brain I, L brain II, R brain II, L brain
III, R brain III). This resulted in a total of 43 embryonic samples. At the E12.5 stage, the primordia of the spleen and pancreas develop
on the outer surface of the stomach, making it challenging to dissect them without cross-contamination. Consequently, these tissues
were either co-dissected alongside the stomach or pooled together for subsequent analysis.
487
488 **Tissue lysis, PCR LoxCode amplification and sequencing library preparation:**
489
490 Tissue samples were lysed in DirectPCR lysis reagent (cell) (Viagen Biotech), and the LoxCode sequence amplified using a 3-round
491 PCR. The first round PCR (20 cycles) uses LoxCode specific primers that efficiently amplify the LoxCode sequence from genomic
492 DNA. As an internal quality control, for each sample, two PCR reactions were performed.
493
494 FWD primer: TCTAGAGGATCCCCGGGTACCGA
495 REV Primers: TGATCCGCGCCTGGATGAAT
496
497 In the second round (5 cycles), overhang primers add a partial Illumina Read 1 and Read 2 sequencing primer site, as well as a stagger
498 between the sequencing primers and the LoxCode sequence in order to increase diversity.

FWD Primers

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG TCTAGAGGATCCCCGGGTACCGA
TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG N TCTAGAGGATCCCCGGGTACCGA
TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG NN TCTAGAGGATCCCCGGGTACCGA
TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG NNN TCTAGAGGATCCCCGGGTACCGA

REV Primers

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG GCCTGGATGAATTCGTGTATAACTTCG
GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG N GCCTGGATGAATTCGTGTATAACTTCG
GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG NN GCCTGGATGAATTCGTGTATAACTTCG
GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG NNN GCCTGGATGAATTCGTGTATAACTTCG

The third round (5 cycles) completes the sequencing primers and adds the P5 and P7 sequencing adaptors as well as introduces a unique sample index.

FWD primer:

AATGATACGGCGACCACCGAGATCTACAC[i5]TCGTCGGCAGCGTC

REV Primers

CAAGCAGAAGACGGCATACGAGAT[i7]GTCTCGTGGGCTCGG

To remove unused primers, primer dimers, and to counteract preferential amplification of shorter LoxCodes, amplicons were purified using SPRI beads at a 0.9:1 ratio between each round. All PCR reactions are carried out in a 20 µl volume, using Kapa Hifi Hot-Start Mastermix (Merck), using the following program 95 °C 2', N times (98 °C 20'', 65 °C 20'', 72 °C 30''), 72 °C 5', 4 °C hold, where N varied depending on the PCR round. Sequencing libraries were pooled at equal volumes and sequenced on the Illumina MiSeq 600 cycle kit, using for Read 1 and Read 2 354 and 264 cycles respectively.

Dissociation of tissues:

For 10X analysis, dissected tissues were dissociated for 30 mins at 37 °C in 200ul of Liberase™ (Roche, 100ug/ml in PBS). 2 mL of cold PBS (without ca/Mg) + 7% FCS + 2.5mM EDTA (FE buffer) was added to the tubes and carefully removed to rinse the enzyme off (the blood sample was centrifuged at this step to avoid cell loss). 1 mL of fresh FE buffer was added. After 15-20 mins on ice, each tissue was mechanically dissociated and filtered on a 40 µm sieve. Cells were pelleted (5 mins, 1500 rpm, 4 °C), counted with an automated cell counter (Invitrogen). 100k cells from each tissue was processed for MultiSeq labelling following the protocol described in McGinnis *et al.*[42]. For each barcode oligo (5'-CCTTGGCACCCGAGAATTCCANNNNNNNNNA$_{30}$-3'), the oligo and anchor LMO (5'-TGGAATTCTCGGGTGCCAAGGgtaacgatccagctgtcact-Lipid-3') were diluted to 2 µM in 1:1 molar ratio in PBS. The co-anchor LMO (5'-Lipid-AGTGACAGCTGGATCGTTAC-3') was diluted to 2 µM in PBS. Cells were resuspended in 200 µL PBS, 23 µl of anchor-barcode mixture added, and incubated on ice for 5 minutes. Then 23 µl of co-anchor solution was added and kept on ice. After an additional 5 minutes, labelling was quenched by addition of 1 mL of 1% BSA in PBS (ice cold). In all subsequent steps, cells were kept in 1% BSA on ice. After MultiSeq labelling, cells were resuspended in FACS buffer (PBS (without Ca/Mg) + 7% FCS) and labelled with 7AAD. Live cells were sorted on an Aria II cell sorter. 10K cells of each tissue were used to create 4 pools for 4 10X lanes: Pool 1 (blood, heart, right and left lung, liver, stomach, intestine, right and left kidney, right and left gonad, face), Pool 2 (dorsal aorta, L hand, R hand, L arm, R arm, L foot, R foot, L leg, R leg, L lat I, L lat II, face), Pool 3 (L lat III, R lat I, R lat II, R lat III, L mid mes I, L mid mes II, L mid mes III, R mid mes I, R mid mes II, R mid mes III, L brain I, face), Pool 4 (R brain I, L brain II, R brain II, L brain III, R brain III, neural T I, neural T II, neural T III, tail, face). The "Face" sample was seeded into each pool to allow for batch effect correction. Each pool was centrifuged (5 mins, 1500 rpm, 4C), resuspended in 32ul of PBS + 1% Bovine Serum Albumin and counted

before loading on Chromium controller (10X Genomics) for scRNAseq. Sequencing libraries were prepared according to manufacturer's recommendations and sequenced on the Illumina platform (NextSeq 75 cycle kit).

Target amplification (scRNAseq):

The LoxCode sequence was amplified from full length cDNA using a combination of PCR, biotin capture, restriction digest and adaptor ligation. In a first round, transcripts contained the LoxCode sequence were enriched from full-length cDNA using a partial Illumina Read 1sequencing primer (present on every fragment in the full length cDNA library), and a reverse primer specific to exon 1 of the *Rosa26* locus. In a second round, overhanging forward primers completes the Read 1 sequencing primer binding site on the 5' end, and biotinylated LoxCode specific primers (located inside the LoxCode construct) further enriches LoxCode cDNA. In a next round, biotinylated amplicon is purified using MyONE C1 streptavidin beads (Thermofisher), removing unwanted DNA. Then LoxCode specific DNA was cleaved off the beads using an Eco53KI restriction site 5' of the LoxCode sequence. Finally, adaptors (IDT) were ligated using Blunt/TA ligation mix (NEB), and amplified using generic P5 and P7 primers. Libraries were sequenced on an Illumina MiSeq 600 cycle kit using custom sequencing primers.

LoxCode Analysis:

All analyses were performed in 'R 4.1.3' and RStudio, except for LoxCode alignment and agent-based simulations of tissue development (Fig. 4). Most graphs were created using the R package 'ggplot 2 3.3.6' [55].

LoxCode Alignment:

Raw Illumina MiSeq paired-end sequencing reads were aligned to LoxCode element sequences using custom C++ code and the library edlib[56]. Depending on the sequencing configuration, for full length LoxCodes with 13 barcode elements, 7 elements were determined from Read 1 and the remaining 6 elements from Read 2. For LoxCodes of size less than 13, a consensus barcode was determined, by comparing overlapping elements from Read 1 and Read 2. Average barcode reads for embryo 1, 2, 3, were 136K, 138K, and 130K per sample respectively. Barcode reads for all samples are provided in Supplementary Fig. 1b.

Bulk barcode filtering, normalization and scaling:

Several filtering steps were performed on the LoxCode alignment output in order to ensure only high-quality samples and barcodes were included in the analysis. Samples with less than 500 barcode reads were discarded (Supplementary Fig. 2a). Barcodes with repeated elements (likely due to PCR artifacts) and barcodes detected in only one of two PCR replicates were discarded, PCR duplicates normalized to counts per million (CPM), and barcode counts merged by taking the mean across the two samples. Next a CPM cut-off of 0.1 was applied to remove barcodes with very low frequencies. To filter out low-complexity LoxCodes for bona-fide clonal fate analysis, an empirical exclusion list of close to 30K common barcodes was generated from a total of 37 independent DNA samples from LoxCode-Rosa26CreErt2 cells (>50K cells per sample) induced at a similar recombination rate (average number of recombination: 5.14 vs 4.58 in actual data) in the embryos but in adult mice. Barcodes detected in more than 10 out of the 37 independent samples were excluded from the analysis. This cut-off provided a good compromise between stringency and the number of barcodes retained in the analysis. In some instances, only barcodes detected in at least two tissues were considered for the analysis (Fig. 2e-g).

Biomass estimation and random sampling:

To evaluate the biomass for each clone (Fig. 2d), relative DNA concentrations were quantified, except for blood, from the cellular lysates using a fluorometric assay (Qubit, Invitrogen). Based on the assumption that an E12.5 embryo contains approximately 6 million cells[3], the cell count for each clone was inferred by multiplying the barcode frequency present in the tissue sample by the relative DNA concentration and the total cell number.

In order to evaluate the presence of a clone across multiple tissues against a null hypothesis, which posits that cells are distributed randomly and uniformly across tissues according to tissue size, random samples were generated by employing the estimated cell count per clone and the relative cell count for each tissue. A comparison of the distribution of the number of tissues detected for both empirical and randomly sampled data was performed using a two-sided Kolmogorov-Smirnov test to assess the statistical significance of observed differences.

Metaclone similarity analysis (MCSA):

Barcode counts per sample were log(1+x) transformed, and the average cosine similarity ($E((A \cdot B) / (\|A\| \|B\|))$) between samples and across embryos (with missing samples removed) calculated and normalized to range from zero to one. Meta-clonal similarity (MCS) values were then ordered using hierarchical clustering (method="ward.D2") and plotted as a heatmap using the R package 'pheatmap 1.0.12'.

UMAP (tissues)

MCS values were computed as for the heatmap and projected onto 2 dimensions through Uniform Manifold Approximation and Projection using the R package 'umap 0.2.7.0' (parameters: n_neighbours_=20, spread_=2.0, min_dist_=0.1, metric="cosine").

Anatomical tissue connectivity map

An anatomical 2D tissue map was created using the locations of the analysed embryonic tissues and regions. A 6-nearest-neighbour network based on MCS values was then projected onto this map connecting nodes between most similar tissues and regions. Edges with an MCS <0.75 were removed. For the purpose of visualization, edges between nodes were drawn using the edge bundling algorithm using the R package 'ggraph 2.0.5'. To ease bundling intermediate nodes (not shown in Fig. 3g) were introduced for brain, neural tube, mesenchymal, organs, forelimbs, hindlimbs and blood.

UMAP (barcodes)

Barcode CPM counts across tissues were log(1+x) transformed and projected onto 2 dimensions through Uniform Manifold Approximation and Projection using the R package 'umap 0.2.7.0' (parameters: n_neighbours=10, spread=1.0, min_dist=0.2, metric="cosine"). Leiden clustering was performed from umap's computed NN-network using the R package 'leiden 0.4.3' (resolution_parameter = 1.0).

Comparison of bona-fide vs MCSA analysis

The Robinson-Foulds distances between trees generated from hierarchical clustering were computed using the Treedist function from the R package 'phangorn 2.10.0'.

Agent based modelling and parameter fitting

Tissue development was modelled using an agent-based approach written in C++ programming language, by combining previously published implementations of stochastic cellular growth[40] and LoxCode barcoding[29] with fate commitment. Heterogeneity in cell division times[57], random recombination during LoxCode barcoding and fate commitment over time were assumed independent random processes, with non-synchronous divisions modelled in continuous time, barcoding occurring at a single time-point (i.e., E5.5), and for ease of implementation differentiation happening at discrete 1 hour time intervals. Times between divisions were assumed log-normal distributed (mean=12 hours, variance=2.56 hours squared), except for the first two divisions which have been shown to last approximately 20 hours[58]. For blood progenitors an approximately two-times slower division time (a factor 1.3 in the parameters of the log-normal distribution) was assumed to prevent blood becoming the dominant tissue at E12.5. Recombination between pairs of LoxP sites were chosen uniformly from the original or already recombined LoxCode cassette with LoxP sites at a distance >82 nts[29] in the

same orientations leading to inversions and LoxP sites in opposite orientations leading to excisions (number of recombination events = 4). Fate commitment was assumed to follow the hierarchy shown in Fig. 3c, with two parameters per branch point determining the minimal maturation time from the previous branchpoint (or time of fertilization for the zygote) and the probability p to differentiate in one or the other branch. Beyond the minimal maturation time, the chance to differentiate was assumed constant per time. In total the model has 38 parameters (32 for the differentiation process, four for division time distributions (1 for the duration of the first two divisions, 2 for the log-normal distribution, 1 for the scaling factor for blood progenitors), one for the differentiation rate, and one for the number of LoxCode recombinations). The simulations were initiated after the first two divisions (4 cells) and run until E12.5, reaching typically between 6-7 million individual agents.

To fit 33 free model parameters (parameters associated with division times and LoxCode recombination were kept fixed) to the experimental data, we utilized the covariance-matrix adaptation evolutionary strategy (CMA-ES) as outlined in reference[41]. This strategy was implemented using the C++ library libcmaes (https://github.com/CMA-ES/libcmaes). CMA-ES is a robust algorithm known for its ability to handle noise and scale effectively to high-dimensional problems. By minimizing the sum-of-squares between the entries in the empirical and simulated MCSA matrix ((17^2-17)/2-1=135 degrees of freedom) and the relative number of barcodes per tissue (16 degrees of freedom), we identified the parameters that optimally reproduced the empirical observations. For performance, in some optimizations, the MCSA matrix was evaluated 250 hours post conception rather than 300 hours. From 20 random initialization, we selected the 5 best-performing runs, all of which reached apparent convergence after 3,000 evaluations (Supplementary Fig. 4b). The best parameters were subsequently plotted (Supplementary Fig. 4c).

Circular pedigree plots

Embryo-scale pedigree charts were produced by employing two simple rules to assign a binary code to each cell in the simulated pedigree. Firstly, the zygote received the binary code '0', while daughter cells inherited the same binary code as their mother, augmented by an additional digit ('0' or '1'). The fate at the time of division was also recorded together with the binary code. Subsequently, cells were classified by generation, arranged lexicographically, and displayed as stacked histograms with the generation on the x-axis (parameters: position='fill' and width scaled to generation) using circular coordinates within the ggplot2 R package and color according to fate at time of division. To generate circular pedigrees for barcoded cells at approximately E5.5, cells that underwent equal or fewer than 10 divisions (the average number of divisions at E5.5) since the zygote stage were filtered from the dataset prior to plot generation.

Single Cell RNA sequencing analysis and cluster annotation

Sequencing data was aligned to mouse genome (mm10) with the cellranger software (version 5.0) and analysed using the R package 'Seurat 4.0.4'. Samples were demultiplexed using the R package 'demultiplex 1.0.2' and custom code. Transcriptional clusters were annotated using established marker genes from the literature (Supplementary Data 2). E12.5 embryo cells could be found in two main clusters: a *Tubb2b*+ neuronal cluster and a mesenchymal cluster with high expression of *Col3a1, Col2a1, Col9a1, Col1a1, Prrx1, Foxd1*. Additional clusters corresponded to neural crest (*Sox10*), blood subsets (primitive erythrocytes (*Hbb-bh1, Hbb-y, Hba-a1, Hba-a2, Alas2, Gypa*), definitive erythrocytes (*Hbb-bh1-, Hbby-, Hba-a1, Hba-a2, Alas2, Gypa*), macrophages (*Cx3cr1, Csf1r, C1qa, C1qb, C1qc*), progenitors (*Mpo, Fcgr3*) and megakaryocytes (*Pf4, Cd9, Itga2b, Gp1bb*), striated muscle (skeletal (*Myf5, Myod, Myog, Acat1, Tnnt1*) and cardiac (*Actc1, Tnnc1, Nppb*)), endothelium (*Cdh5, Cldn5, Kdr, Pecam1, Ramp2, Cd93*), epithelial cells (*Krt5, Krt8, Krt18, Epcam*) and pericytes (*Pdgfrb, Kcnj8, Rgs5*). Tissue of origin was consistent with transcriptional identity.

The neuronal cluster could be further subsetted into neuronal/glial progenitors (*Sox2, Nes, hes5, Btg2, Ub2c, Cenpf*), neuroblasts (*Sox2, Tubb3, Neurog2*) and differentiating neurons (*Tubb3, Stmn2, Stmn3, Sox11*). Differentiating neurons, could be broadly grouped as *Nrx3n-* (differentiating neurons 1) and *Nrx3n+* (differentiating neurons 2) populations, with GABAergic neurons (*Dlx5*) forming a distinctive loop within the *Nrx3n+* neurons. Neuronal neural crest derivatives (dorsal root ganglia (*Isl1, Neurod1*) and enteric neurons (*Isl1, Phoxa1, Phoxa2, Ret, Th*)) were also identifiable.

The mesenchymal cluster contained differentiating visceral and musculoskeletal cells populating the upper and lower region of that cluster, respectively (Fig. 4). A largely undifferentiated/proliferative mesenchymal population (absent of differentiation markers, *Cenpf,*

Ube2c, Top2a) was found at the interface between visceral and musculoskeletal cell types. Within the lower section of the musculoskeletal region, undifferentiated musculoskeletal mesenchyme (*Msx1, Hoxa13, Hoxd13, Lhx2*), differentiating chondrocytes (*Col2a1, Col9a1, Col9a2, Col9a3, Sox9, Acan*), tenocytes/ligamentocytes (*Scx, Dcn, Postn, Fbn2, Col5a2, Col1a1, Col3a1*), fibroblasts (*Col1a1, Col1a2, Col3a1, Cd63, Cxcl14, Shox2, Ddr2, Tnc, Vim, Pdgfra*) and meningeal fibroblasts (*Col1a1, Col1a2, Col3a1, Cd63, Vim, Zic1, Foxd1*) could be identified. The upper visceral region of the mesenchymal cluster contained ovarian mesenchyme (*Col1a1, Col1a2, Col3a1, Tcf21, Wnt4, Wt1, Kitl, Amhr2, Wnt6*), pregranulosa cells (*Upk3b, Aldh1a2, Wt1, Krt18, Krt7, Krt19, Upk1b, Alcam*), smooth muscle (*Acta2, Tagln, Myh11, Cnn1, Myocd*), pulmonary/gastrointestinal mesenchyme (*Foxf1, Wnt2, Barx1*), renal mesenchyme (*Foxd1, Pbx1, Meis2, Hoxd11, Hoxc10, Wt1*), nephron progenitors (*Meox1, Hoxd11*), valve/endocardial cushion (*Col1a1, Col3a1, Col1a2, Acta2, Tagln2, Eln, Tbx20*) and perivascular cells (*Col1a1, Col3a1, Col1a2, Cxcl12*).

Single-cell MCSA

Similar to MCSA based on tissue level PCR barcoding data, frequencies of barcodes detected in single cells between cell types (instead of tissues) were log(1+x) transformed and used to compute scaled cosine similarity values. Only clones with at least five cells were included and to increase specificity the six most common barcodes were excluded ("+9","-9","+1","+13","+1+2+3","+11+12+13"). Cell types were then clustered using leiden clustering (R package 'leiden 0.4.3', resolution_parameter = 0.7) and projected onto 2 dimensions with the R package 'umap 0.2.7.0' (parameters: n_neighbors=6, spread=1, min_dist=0.01, metric="cosine").

Statistical analysis

Chi-square and Kolmogorov-Smirnov tests were computed in R. If appropriate, p-values were adjusted for multiple comparisons using the Benjamini-Hochberg correction[59]. P-values below 0.05 were assumed statistically significant.

References:

Lawson, K. A., Meneses, J. J. & Pedersen, R. A. Clonal analysis of epiblast fate during germ layer formation in the mouse embryo. *Development* **113**, 891-911 (1991). https://doi.org:10.1242/dev.113.3.891

Parameswaran, M. & Tam, P. P. Regionalisation of cell fate and morphogenetic movement of the mesoderm during mouse gastrulation. *Dev Genet* **17**, 16-28 (1995). https://doi.org:10.1002/dvg.1020170104

Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis. *Nature* **566**, 496-502 (2019). https://doi.org:10.1038/s41586-019-0969-x

Peng, G. *et al.* Molecular architecture of lineage allocation and tissue organization in early mouse embryo. *Nature* **572**, 528-532 (2019). https://doi.org:10.1038/s41586-019-1469-8

Chan, M. M. *et al.* Molecular recording of mammalian embryogenesis. *Nature* **570**, 77-82 (2019). https://doi.org:10.1038/s41586-019-1184-5

Tam, P. P. L. & Ho, J. W. K. Cellular diversity and lineage trajectory: insights from mouse single cell transcriptomes. *Development* **147** (2020). https://doi.org:10.1242/dev.179788

Grosswendt, S. *et al.* Epigenetic regulator function through mouse gastrulation. *Nature* **584**, 102-108 (2020). https://doi.org:10.1038/s41586-020-2552-x

Mittnenzweig, M. *et al.* A single-embryo, single-cell time-resolved model for mouse gastrulation. *Cell* **184**, 2825-2842 e2822 (2021). https://doi.org:10.1016/j.cell.2021.04.004

Argelaguet, R. *et al.* Multi-omics profiling of mouse gastrulation at single-cell resolution. *Nature* **576**, 487-491 (2019). https://doi.org:10.1038/s41586-019-1825-8

Saykali, B. *et al.* Distinct mesoderm migration phenotypes in extra-embryonic and embryonic regions of the early mouse embryo. *Elife* **8** (2019). https://doi.org:10.7554/eLife.42434

Tzouanacou, E., Wegener, A., Wymeersch, F. J., Wilson, V. & Nicolas, J. F. Redefining the progression of lineage segregations during mammalian embryogenesis by clonal analysis. *Dev Cell* **17**, 365-376 (2009). https://doi.org:10.1016/j.devcel.2009.08.002

VanHorn, S. & Morris, S. A. Next-Generation Lineage Tracing and Fate Mapping to Interrogate Development. *Dev Cell* **56**, 7-21 (2021). https://doi.org:10.1016/j.devcel.2020.10.021

Chen, C., Liao, Y. & Peng, G. Connecting past and present: single-cell lineage tracing. *Protein Cell* **13**, 790-807 (2022). https://doi.org:10.1007/s13238-022-00913-7

Livet, J. *et al.* Transgenic strategies for combinatorial expression of fluorescent proteins in the nervous system. *Nature* **450**, 56-62 (2007). https://doi.org:10.1038/nature06293

Ratz, M. *et al.* Clonal relations in the mouse brain revealed by single-cell and spatial transcriptomics. *Nat Neurosci* **25**, 285-294 (2022). https://doi.org:10.1038/s41593-022-01011-x

Sun, J. *et al.* Clonal dynamics of native haematopoiesis. *Nature* **514**, 322-327 (2014). https://doi.org:10.1038/nature13824

Rodriguez-Fraticelli, A. E. *et al.* Clonal analysis of lineage fate in native haematopoiesis. *Nature* **553**, 212-216 (2018). https://doi.org:10.1038/nature25168

Yang, D. *et al.* Lineage tracing reveals the phylodynamics, plasticity, and paths of tumor evolution. *Cell* **185**, 1905-1923 e1925 (2022). https://doi.org:10.1016/j.cell.2022.04.015

Kalhor, R. *et al.* Developmental barcoding of whole mouse via homing CRISPR. *Science* **361** (2018). https://doi.org:10.1126/science.aat9804

Bowling, S. *et al.* An Engineered CRISPR-Cas9 Mouse Line for Simultaneous Readout of Lineage Histories and Gene Expression Profiles in Single Cells. *Cell* **181**, 1693-1694 (2020). https://doi.org:10.1016/j.cell.2020.06.018

Quinn, J. J. *et al.* Single-cell lineages reveal the rates, routes, and drivers of metastasis in cancer xenografts. *Science* **371** (2021). https://doi.org:10.1126/science.abc1944

Frieda, K. L. *et al.* Synthetic recording and in situ readout of lineage information in single cells. *Nature* **541**, 107-111 (2017). https://doi.org:10.1038/nature20777

Pei, W. *et al.* Polylox barcoding reveals haematopoietic stem cell fates realized in vivo. *Nature* **548**, 456-460 (2017). https://doi.org:10.1038/nature23653

Pei, W. *et al.* Resolving Fates and Single-Cell Transcriptomes of Hematopoietic Stem Cell Clones by PolyloxExpress Barcoding. *Cell Stem Cell* **27**, 383-395 e388 (2020). https://doi.org:10.1016/j.stem.2020.07.018

Park, S. *et al.* Clonal dynamics in early human embryogenesis inferred from somatic mutation. *Nature* **597**, 393-397 (2021). https://doi.org:10.1038/s41586-021-03786-8

Choi, J. *et al.* A time-resolved, multi-symbol molecular recorder via sequential genome editing. *Nature* **608**, 98-107 (2022). https://doi.org:10.1038/s41586-022-04922-8

Loveless, T. B. *et al.* Molecular recording of sequential cellular events into DNA. *bioRxiv* (2021).

Weinreb, C., Rodriguez-Fraticelli, A., Camargo, F. D. & Klein, A. M. Lineage tracing on transcriptional landscapes links state to fate during differentiation. *Science* **367** (2020). https://doi.org:10.1126/science.aaw3381

Weber, T. S. *et al.* Site-specific recombinatorics: in situ cellular barcoding with the Cre Lox system. *BMC Syst Biol* **10**, 43 (2016). https://doi.org:10.1186/s12918-016-0290-3

Biben, C. *et al.* In vivo clonal tracking reveals evidence of haemangioblast and haematomesoblast contribution to yolk sac haematopoiesis. *Nat Commun* **14**, 41 (2023). https://doi.org:10.1038/s41467-022-35744-x

Gardner, R. L. & Rossant, J. Investigation of the fate of 4-5 day post-coitum mouse inner cell mass cells by blastocyst injection. *J Embryol Exp Morphol* **52**, 141-152 (1979).

Kwon, G. S., Viotti, M. & Hadjantonakis, A. K. The endoderm of the mouse embryo arises by dynamic widespread intercalation of embryonic and extraembryonic lineages. *Dev Cell* **15**, 509-520 (2008). https://doi.org:10.1016/j.devcel.2008.07.017

McGrath, K. E. *et al.* A transient definitive erythroid lineage with unique regulation of the beta-globin locus in the mammalian embryo. *Blood* **117**, 4600-4608 (2011). https://doi.org:10.1182/blood-2010-12-325357

Kinder, S. J. *et al.* The orderly allocation of mesodermal cells to the extraembryonic structures and the anteroposterior axis during gastrulation of the mouse embryo. *Development* **126**, 4691-4701 (1999). https://doi.org:10.1242/dev.126.21.4691

Padron-Barthe, L. *et al.* Clonal analysis identifies hemogenic endothelium as the source of the blood-endothelial common lineage in the mouse embryo. *Blood* **124**, 2523-2532 (2014). https://doi.org:10.1182/blood-2013-12-545939

Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-connected communities. *Sci Rep* **9**, 5233 (2019). https://doi.org:10.1038/s41598-019-41695-z

Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP. *Nat Biotechnol* (2018). https://doi.org:10.1038/nbt.4314

McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and projection for dimension reduction. *arXiv preprint arXiv:1802.03426* (2018).

Naik, S. H., Schumacher, T. N. & Perie, L. Cellular barcoding: a technical appraisal. *Exp Hematol* **42**, 598-608 (2014). https://doi.org:10.1016/j.exphem.2014.05.003

Weber, T. S., Perie, L. & Duffy, K. R. Inferring average generation via division-linked labeling. *J Math Biol* **73**, 491-523 (2016). https://doi.org:10.1007/s00285-015-0963-3

Hansen, N. The CMA Evolution Strategy: A Tutorial. arXiv:1604.00772 (2016). <https://ui.adsabs.harvard.edu/abs/2016arXiv160400772H>.

McGinnis, C. S. *et al.* MULTI-seq: sample multiplexing for single-cell RNA sequencing using lipid-tagged indices. *Nat Methods* **16**, 619-626 (2019). https://doi.org:10.1038/s41592-019-0433-8

Chen, L., Wang, D., Wu, Z., Ma, L. & Daley, G. Q. Molecular basis of the first cell fate determination in mouse embryogenesis. *Cell Res* **20**, 982-993 (2010). https://doi.org:10.1038/cr.2010.106

Bardot, E. S. & Hadjantonakis, A. K. Mouse gastrulation: Coordination of tissue patterning, specification and diversification of cell fate. *Mech Dev* **163**, 103617 (2020). https://doi.org:10.1016/j.mod.2020.103617

Zhu, M. *et al.* Developmental clock and mechanism of de novo polarization of the mouse embryo. *Science* **370** (2020). https://doi.org:10.1126/science.abd2703

Dumas, L., Clavreul, S., Michon, F. & Loulier, K. Multicolor strategies for investigating clonal expansion and tissue plasticity. *Cell Mol Life Sci* **79**, 141 (2022). https://doi.org:10.1007/s00018-021-04077-1

Tabansky, I. *et al.* Developmental bias in cleavage-stage mouse blastomeres. *Curr Biol* **23**, 21-31 (2013). https://doi.org:10.1016/j.cub.2012.10.054

Goolam, M. *et al.* Heterogeneity in Oct4 and Sox2 Targets Biases Cell Fate in 4-Cell Mouse Embryos. *Cell* **165**, 61-74 (2016). https://doi.org:10.1016/j.cell.2016.01.047

Cheng, H., Zheng, Z. & Cheng, T. New paradigms on hematopoietic stem cell differentiation. *Protein Cell* **11**, 34-44 (2020). https://doi.org:10.1007/s13238-019-0633-0

Leeper, K. *et al.* Lineage barcoding in mice with homing CRISPR. *Nat Protoc* **16**, 2088-2108 (2021). https://doi.org:10.1038/s41596-020-00485-y

Sulston, J. E., Schierenberg, E., White, J. G. & Thomson, J. N. The embryonic cell lineage of the nematode Caenorhabditis elegans. *Dev Biol* **100**, 64-119 (1983). https://doi.org:10.1016/0012-1606(83)90201-4

Domcke, S. & Shendure, J. A reference cell tree will serve science better than a reference cell atlas. *Cell* **186**, 1103-1114 (2023). https://doi.org:10.1016/j.cell.2023.02.016

Kueh, A. J. *et al.* An update on using CRISPR/Cas9 in the one-cell stage mouse embryo for generating complex mutant alleles. *Cell Death Differ* **24**, 1821-1822 (2017). https://doi.org:10.1038/cdd.2017.122

Chevalier, C., Nicolas, J. F. & Petit, A. C. Preparation and delivery of 4-hydroxy-tamoxifen for clonal and polyclonal labeling of cells of the surface ectoderm, skin, and hair follicle. *Methods Mol Biol* **1195**, 239-245 (2014). https://doi.org:10.1007/7651_2013_63

Wickham, H. in *Use R!,*   1 online resource (XVI, 260 pages 232 illustrations, 140 illustrations in color (Springer International Publishing : Imprint: Springer,, Cham, 2016).

Sosic, M. & Sikic, M. Edlib: a C/C ++ library for fast, exact sequence alignment using edit distance. *Bioinformatics* **33**, 1394-1395 (2017). https://doi.org:10.1093/bioinformatics/btw753

Weber, T. S., Jaehnert, I., Schichor, C., Or-Guil, M. & Carneiro, J. Quantifying the length and variance of the eukaryotic cell cycle phases by a stochastic model and dual nucleoside pulse labelling. *PLoS Comput Biol* **10**, e1003616 (2014). https://doi.org:10.1371/journal.pcbi.1003616

Ciemerych, M. A. & Sicinski, P. Cell cycle in mouse development. *Oncogene* **24**, 2877-2898 (2005). https://doi.org:10.1038/sj.onc.1208608

Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**, 289-300 (1995).

Data availability

Unfiltered and filtered barcoding data, barcode exclusion list, and raw as well as processed scRNA data generated in this study have been deposited at Zenodo (10.5281/zenodo.7840234). A description of the files is provided as Supplementary Notes 1.

Code availability

Custom C++ code for LoxCode alignment and agent-based simulations have been deposited at Zenedo (10.5281/zenodo.7840234). A description of the files is provided as Supplementary Notes 1.

Acknowledgements

We thank the WEHI genomics and single cell core (SCORE), especially Casey Anttila for 10X library preparations and Daniel Brown for multiplexing, CRISPR mouse facility MAGEC, especially Andrew Kueh for mouse generation, and mouse facilities for animal husbandry. We thank Illumina, Inc for sequencing support. We thank Chris McGinnis and the Gartner lab (UCSF) for sharing MultiSeq reagents. We would also like to thank Ken Duffy for discussions and input during the early conceptual phase of the LoxCode technology. This work was funded by the CASS Foundation (SM/18/7776 for T.S.W.), National Health and Medical Research Council, Australia (2012196 (T.S.W), 1184736 (T.S.W. & S.H.N), 2009675 and 1145184 (S.H.N.), 1052195 (D.C.M.), 1129012 and 2011770 (S.T), the DHB Foundation (S.T) Human Frontiers Special Program (RGP0060/2012).

Contributions

T.S.W. with S.H.N. conceived the LoxCode approach. D.C.M., T.S.W., and S.H.N. built the LoxCode construct with S.G.'s advice. T.S.W., C.B, D.C.M, P.T, S.T., S.H.N. conceived the study design. C.B. performed all embryonic dissections, with T.S.W. for tissue and single cell processing. T.S.W optimised library preparation and sequencing. T.S.W. performed all analysis and generated figures, with S.Z. and S.T. for the theoretical LoxCode recombination look-up table and empirical exclusion list respectively. C.B. provided cluster annotations, and T.S.W., C.B., S.T., P.T. and S.H.N. interpreted the data.

T.S.W, D.C.M., S.T., and S.H.N. sourced funding. S.T. and S.H.N. supervised the study. All authors contributed to writing the manuscript.

Supplementary Figures

Supplementary Fig 1. a) Distribution of barcode sizes for all (PCR-filtered) and bona-fide (empirical-frequency-filtered) clones. **b)** Distribution of the minimum number of recombinations (complexity) for all (PCR-filtered) and bona-fide clones. **c)** LoxCode sequencing reads for all samples (including PCR replicates).

Supplementary Fig 2. a) Cosine similarity matrix for all tissues using bona-fide barcodes ordered according to the corresponding

890　MCSA heatmap in Fig. 2e.

Supplementary Fig 3. MCSA heatmap, UMAP and pseudo-anatomical map of tissue relationships. These plots are as per Figure 2 and

893 represent the corresponding 3 biological replicates that were pooled to generate Figure 2.

Supplementary Fig 4. a) Heatmap showing empirical vs simulated barcode data pooled over three embryos. **b)** Convergence of the CMA-ES optimization routine (5 out of 20 best performing runs are shown). **c)** Best-fit parameters from 5 best performing runs (boxplot indicating median, 25th and 75th quantile). Populations and parameters are provided in Supplementary Methods 1.

Supplementary Fig 5. The 40 tissues, which were dissected and processed into a single-cell suspension as depicted in Figure 1, underwent multiplexing using the MULTI-seq method. Subsequently, the samples were pooled and analyzed using a droplet-based scRNA-seq platform with a 3' capture scRNA-seq kit. Demultiplexing via MULTI-seq proved effective in: a) distinguishing the tissues of origin, and b) differentially distributing them across the transcriptional space (as illustrated by the UMAP in Figure 4). It should be noted that distinct colors represent left and right samples (or I, II, III for mesenchymal tissues) of the specified tissue, such as left and right kidneys.

907 **Supplementary Fig 6. Meta-clonal fate heterogeneity. a)** Examples of single cell meta-clonal output corresponding to the indicated

908 LoxCodes. **b)** Distribution of single cells per detected LoxCode according to tissue (top) or cell type (bottom).

910 **Supplementary Fig 7. Asymmetry in clonal contribution.** Contributions of meta-clones to a) left and right fore- and hindlimbs, b)

911 left and right gonad and kidneys. Meta-clones with significant bias are shown in Fig 4. g.

917