Pan-genome study underlining the extent of genomic variation of invasive *Streptococcus pneumoniae* in Malawi

Arash Iranzadeh[1], Arghavan Alisoltani[2,3,4], Anmol M Kiran[5,6], Robert F Breiman[7], Chrispin Chaguza[8,9,10],

Chikondi Peno[5,6,9], Jennifer E Cornick[5,11], Dean B Everett[12&], Nicola Mulder[1&]

[1] Computational Biology Division, Department of Integrative Biomedical Sciences, Institute of Infectious Disease and Molecular Medicine, Faculty of Health Sciences, University of Cape Town, Western Cape, South Africa.

[2] Department of Microbiology-Immunology, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA.

[3] Department of Medicine, Division of Infectious Diseases, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA.

[4] Center for Pathogen Genomics and Microbial Evolution, Havey Institute for Global Health, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA.

[5] Malawi-Liverpool-Wellcome Trust, Queen Elizabeth Central Hospital, College of Medicine, Blantyre, Malawi.

[6] Centre for Inflammation Research, Queens Research Institute, University of Edinburgh, Edinburgh, United Kingdom.

[7] Rollins School of Public Health, Emory University, Atlanta, Georgia, United States of America.

[8] Parasites and Microbes Programme, Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, United Kingdom.

[9] Department of Epidemiology of Microbial Diseases, Yale School of Public Health, Yale University, New Haven, Connecticut, USA.

[10] Yale Institute for Global Health, Yale University, New Haven, Connecticut, USA.

[11] Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, United Kingdom.

[12] Department of Pathology, College of Medicine and Health Sciences, Khalifa University, Abu Dhabi, UAE.

[&] Joint last authors

Corresponding author:

30   Nicola Mulder: nicola.mulder@uct.ac.za

Impact Statement

Our research applied pan-genomics principles to comprehensively assess diversity within the

pneumococcus genome, with the primary objective of identifying pneumococcal virulence genes for

advancing vaccine design and drug development. Within this study, we identified Serotypes 1 and 5 as

the predominant and highly invasive pneumococcal strains in Malawi, characterized by a short

nasopharyngeal colonization period, suggesting their potential for rapid infection of sterile sites within

the human body such as blood and the central nervous system. These serotypes exhibited significant

genetic divergence from other serotypes in Malawi, notably lacking key genes within the RD8a operon

while harboring transporters functioning independently of ATP. It's important to note that these findings

are based on computational analysis, and further validation through laboratory experiments is essential

to confirm their biological significance and potential clinical applications. The implications of our

research offer potential avenues for more effective pneumococcal disease prevention and treatment,

not only in Malawi but also in regions facing similar challenges.

Abstract

Streptococcus pneumoniae is a common cause of acute bacterial infections in Malawi. Understanding

the molecular mechanisms underlying its invasive behavior is crucial for designing new therapeutic

strategies. We conducted a pan-genome analysis to identify potential virulence genes in *S. pneumoniae*

by comparing the gene pool of isolates from carriers' nasopharyngeal secretions to isolates from the

blood and cerebrospinal fluid of patients. Our analysis involved 1,477 pneumococcal isolates from

Malawi, comprising 825 samples from carriers (nasopharyngeal swab) and 652 from patients (368 from

blood and 284 from cerebrospinal fluid). We identified 56 serotypes in the cohort. While most serotypes

exhibited a similar prevalence in both carriage and disease groups, serotypes 1 and 5, the most

abundant serotypes in the entire cohort, were significantly more commonly detected in specimens from

patients compared to the carriage group. This difference is presumably due to their shorter

nasopharyngeal colonization period. Furthermore, these serotypes displayed genetic distinctiveness

from other serotypes. A magnificent genetic difference was observed in the absence of genes from the

RD8a genomic island in serotypes 1 and 5 compared to significantly prevalent serotypes in the

nasopharynx. RD8a genes play pivotal roles in binding to epithelial cells and performing aerobic

respiration to synthesize ATP through oxidative phosphorylation. The absence of RD8a from serotypes 1

and 5 may be associated with a shorter duration in the nasopharynx, theoretically due to a reduced

capacity to bind to epithelial cells and access free oxygen molecules required for aerobic respiration

(essential to maintain the carriage state). Serotypes 1 and 5, significantly harbor operons that encode

phosphoenolpyruvate phosphotransferase systems, which might relate to transporting carbohydrates,

relying on phosphoenolpyruvate as the energy source instead of ATP. In conclusion, serotypes 1 and 5 as

the most prevalent invasive pneumococcal strains in Malawi, displayed considerable genetic divergence

from other strains, which may offer insights into their invasiveness and potential avenues for further

research.

68   Author summary

Despite introducing the pneumococcal conjugate vaccine in 2011, *Streptococcus pneumoniae* remains a

major cause of bacterial infection in Malawi. Whilst some pneumococcal strains harmlessly colonize the

nasopharynx, others find their way into normally sterile sites, such as lungs, blood, and nervous system,

resulting in serious illness. Our study identified specific pneumococcal serotypes as the most invasive in

Malawi, characterized by a short colonization period and significant genetic distinctiveness from other

strains. This genetic divergence notably included the absence of several genes associated with aerobic

respiration and the presence of genes facilitating ATP-independent carbohydrate transport. The

presence or absence of these genes may underlie their heightened invasiveness and shorter colonization

period. This hypothesis positions these genes as potential candidates for future therapeutic research.

We propose that the specific gene gain and/or loss in invasive versus other serotypes may be linked to

the development of invasive pneumococcal diseases.

Introduction

Streptococcus pneumoniae, also known as *pneumococcus*, is a Gram-positive, facultatively anaerobic

bacteria and is one of the leading causes of mortality worldwide. Despite reductions in the incidence of

pneumococcal disease in countries that introduced pneumococcal conjugate vaccines (PCV), the

pneumococcal mortality rate is still high. Pneumococci are estimated to be responsible for 317,300

deaths in children aged 1 to 59 months worldwide in 2015 [1]. In the post-PCV era, a high disease

burden has still been reported in low-income countries in Africa, such as Malawi [2].

Although pneumococcal nasopharyngeal colonization is asymptomatic, it is a prerequisite for

transmission and disease development [3][4]. Symptoms appear when isolates from the nasopharynx

spread to normally sterile sites such as the lung, blood, and central nervous system. Depending on the

infected organ, *S. pneumoniae* can cause two types of infection: (i) non-invasive (mucosal)

pneumococcal diseases such as otitis media and sinusitis, and (ii) invasive pneumococcal diseases (IPD)

such as bacteremia and meningitis. IPD incidence is highest among infants, the elderly, and

immunosuppressed people, most likely due to their less efficient immune systems [5].

Pneumococci possess several virulence factors, including the polysaccharide capsule, surface proteins,

and enzymes [6][7]. The polysaccharide capsule is the most important virulence factor as it aids the

pneumococcus in evading the immune response during colonization and invasion [8]. Its biosynthesis is

regulated by a cluster of genes in the capsular polysaccharide (*cps*) locus [8][9]. Pneumococcal serotypes

are defined by the type and order of monosaccharides that compose the capsule structure. To date, one

hundred pneumococcal serotypes have been identified [10]. Each strain has a set of capsular synthesis

genes in the *cps* locus that determine its serotype. Immunogenic properties of the capsular

polysaccharide were utilized to develop all pneumococcal vaccines in use, including PCV7, PCV10, and

PCV13 that cover 7, 10, and 13 serotypes, respectively. PCV13 includes the following serotypes: 1, 3, 4,

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F. Although the introduction of PCVs has significantly reduced

the burden of disease caused by vaccine types (VTs), serotype replacement has increased the non-

vaccine types (NVTs) carriage rate and IPD incidence [11][12].

In November 2011, PCV13 was introduced in Malawi, which markedly decreased the health system

burden and rates of severe childhood pneumonia [13]. A case-control study in Malawi showed vaccine

effectiveness against VT-IPD of 80·7% [14]. A nasopharyngeal carriage survey conducted in the Karonga

district showed that although the vaccine reduced the VT colonization rate, a moderate level of serotype

replacement was observed among carriers [15]. Moreover, the emergence of antibiotic-resistant

pneumococci due to the overuse of antibiotics is a global concern in the 21st century [16]. To develop

new, more effective vaccines against the vaccine-escape clones and design effective drugs against

antibiotic-resistant strains, it is critical to understand the functions of genes involved in pneumococcal

colonization and pathogenesis. During the past decade, the evolution of high-throughput sequencing

technologies has generated enormous amounts of genomic data that have enabled researchers to

perform large-scale genomic analysis. A well-known example is pan-genome studies. The pan-genome is

the entire gene set in a collection of closely related strains within a specie [17]. Determining the genetic

drivers is an active and promising area of research that can provide insights into pneumococcal disease

prevention and treatment to reduce mortality rates. The pan-genome is useful for analyzing

recombinogenic pathogens such as *pneumococcus* [18]. The recombination level is high in thirteen

pneumococcal genomic loci known as regions of diversity (RDs) numbered from RD1 to RD13, some of

which are involved in virulence [19][20].

In this study, we conducted whole-genome sequencing (WGS) on 1477 pneumococcal samples from

residents of Blantyre, Karonga, and Lilongwe in Malawi. Our study aims to: (i) identify serotype

distribution, (ii) characterize the pneumococcal population structure, and (iii) identify potential driver

genes for invasion and their biological functions.

Materials and methods

Study design and sample collection

The study utilized archived samples maintained by the Malawi-Liverpool Wellcome Trust Clinical

Research Programme (MLW). Samples were collected from individuals residing in three distinct regions

of Malawi: Blantyre in the south, Karonga in the north, and Lilongwe in the central part of the country.

This cohort included isolates obtained from both asymptomatic carriers and symptomatic patients.

Carriage samples were collected from the nasopharynx of healthy individuals as part of the Health and

Demographic Surveillance System in Karonga and Blantyre. The collection process involved the use of

nasopharyngeal swabs. Subsequently, pneumococcal isolates were identified utilizing a previously

established protocol [21]. Briefly, the identification method involved culturing isolates on blood agar

supplemented with gentamicin, with further confirmation relying on optochin disc-based assays,

scrutinizing colony morphology, alpha-hemolysis, and optochin susceptibility, adhering to established

norms and practices for pneumococcal isolates. To account for the true diversity of carriages, only a

single isolated colony was sequenced and serotyped, therefore no carriage samples included in this

study had multiple serotypes.

Invasive pneumococcal samples were also sourced from archived bacterial isolates at MLW, which had

been collected from the blood and cerebrospinal fluid (CSF) of symptomatic patients attending Queen

Elizabeth Central Hospital in Blantyre and Kamuzu Central Hospital in Lilongwe. Notably, the selection of

isolates for this group was blind to their serotypes, ensuring an accurate representation of their

prevalence in the disease group without any influence from serotype inclusion criteria. These isolates

were subsequently streaked onto blood agar plates supplemented with gentamicin, and optochin tests

were conducted, according to the procedures outlined in reference [22].

It's important to note that this study did not involve paired samples. During data collection, each

individual contributed only one sample, which was either a nasopharyngeal swab from healthy

individuals or a blood or CSF sample from symptomatic patients. In the context of this study, the term

'sterile sites' refers to blood and CSF. The term 'invasive samples' specifically refers to those samples

obtained from these sterile sites (blood and CSF).

Whole-genome sequencing and quality control

Archived samples were sequenced under the Global Pneumococcal Sequencing project and

Pneumococcal African Genomic Consortium at the Wellcome Trust Sanger Institute in the United

Kingdom. Bacterial DNA was extracted using the QIAamp DNA mini kit and QIAgen Biorobot by QIAGEN.

Whole-genome sequencing was conducted on Illumina Genome Analyzer II and HiSeq platforms,

producing 125 nucleotide paired-end reads. Read quality was assessed using Fastqc [23].

In-silico serotyping, sequence typing, and quantification of serotype invasiveness

SeroBA version 1.23.4 was employed to infer the serotype of the samples [24]. SeroBA applies a k-mer

method to determine serotypes directly from the paired-end reads in FASTQ format. Any serotype with

a relative frequency greater than 5% was categorized as an abundant serotype. To identify serotypes

with a significant presence in the nasopharynx and sterile sites, Fisher's exact test was applied. P-values

were adjusted using the Benjamini-Hochberg method, and serotypes with an adjusted p-value less than

0.01 were considered significant. The odds ratio (OR) was calculated as follows: OR = (ad)/(bc), where 'a'

represents the number of invasive serotype k, 'b' is the number of carriage serotype k, 'c' is the number

of invasive non-serotype k, and 'd' is the number of carriage non-serotype k. Zero values were replaced

by 0.5 in OR calculations. Abundant serotypes with a significant presence in the nasopharynx were

considered to have low invasiveness, while abundant serotypes with a significant presence in sterile

sites were considered to have high invasiveness. Fisher's exact test was also used to identify serotypes

whose frequencies changed significantly after the introduction of PCV13 in 2011 (adjusted p-value <

0.01).

Genome assembly and annotation

176 Genomes were assembled using Velvet Optimiser version 2.2.5 [25] with settings to generate contigs

177 longer than 500 base pairs, employing a hash range from 61 to 119. The quality of the assembled

178 genomes was assessed using Quast version 5.2 [26], and annotation was performed with Prokka version

179 13.1 [27].

Pan-genome construction

181 The pan-genome for the samples was generated using Roary version 3.12.0 [28]. Roary was run to

182 perform the core gene alignments with Mafft version 7.313 [29]. Genes in the pan-genome were

183 categorized into three groups based on their abundance among samples: genes present in 100% of

184 samples were designated as core genes, those in more than 95% but not core were termed soft-core

185 genes, and the remaining genes were considered accessory genes.

Analysis of the population structure

Small-scale variations, such as single nucleotide polymorphisms (SNPs) and short indels, within the core

genes were analyzed to understand population diversity. A phylogenetic tree, illustrating the genetic

separation between samples, was constructed using SNPs and indels in the core gene alignment as

phylogeny markers. The core genome alignment served as input for IQ-TREE version 2 [30] to generate a

phylogenetic tree, which was visualized using iTol version 3 [31].

Diversity in the accessory genome manifests as large-scale gene presence-absence variations. The R

package Nonnegative Matrix Factorization [32] was used to create a gene presence-absence heatmap

from the pan-genome matrix. To determine the factors influencing gene distribution, including isolation

sites (nasopharynx, blood, and CSF), serotypes, geographical locations (Blantyre, Karonga, and

Lilongwe), and vaccination eras, a principal component analysis (PCA) of gene distribution was

conducted using the R package MixOmics version 6.20.0 [33].

Gene presence-absence statistical analysis

Phenotypic traits of samples were assigned based on population structure and invasiveness. To identify

putative virulence factors, a gene presence-absence statistical test was conducted using Scoary version

1.6.1643 [34]. This tool scores the components of the pan-genome for associations with observed

phenotypic traits while accounting for population stratification. The test was conducted across samples

from different sources and serotypes to find putative virulence factors. Genes with a Bonferroni-

corrected p-value less than 0.05 were deemed significant.

Functional and gene ontology (GO) enrichment analysis

The list of significant genes was submitted to STRING webtool version 11.5 [35] for functional

enrichment analysis. STRING is a network that integrates information from various protein-protein

interaction databases, predicting both direct (physical) and indirect (functional) interactions between

proteins from five sources, including genomic context predictions, lab experiments, co-expression,

automated text mining, and previous knowledge in databases. Functional enrichment analysis in STRING

utilizes information from classification systems such as the Kyoto Encyclopedia of Genes and Genomes

(KEGG) [36] and the Protein families database (Pfam) [37]. The tool entitled "Multiple Sequences" was

selected, and the *S. pneumoniae TIGR4* was chosen as the reference organism. STRING reports the

associated enriched pathways with a false discovery rate of less than 0.05.

Results

In total, 825 isolates from the nasopharynx of healthy carriers, 368 isolates from the blood of

bacteremia patients, and 284 isolates from the CSF of meningitis patients were sequenced. The

demographics of the samples are shown in Table 1.

Table 1. Demographics of 1477 pneumococcal isolates collected from Malawi.

Characteristics	Categories	Nasopharynx	Blood	CSF
Age (in years)	< 5	538	165	141
	5-19	109	42	60
	20-40	60	67	50
	> 40	7	24	7
	Unknown	111	70	26
Sex	Female	401	131	111
	Male	313	122	122
	Unknown	111	115	51
City	Blantyre	169	357	259
	Karonga	656	0	0
	Lilongwe	0	0	23
	Unknown	0	11	2
Sampling period		2009-2014	1997-2015	2000-2015

Serotypes 1, 5, and 12F had the highest invasiveness, likely with a short period of

nasopharyngeal colonization.

Altogether, we identified isolates belonging to 56 different serotypes. Irrespective of their isolation

sources, serotypes 1 (8.7%), 5 (7.8%), 6B (6.6%), 23F (6.3%), and 19F (5.5%) were the most prevalent,

each accounting for over 5% of the entire cohort. Of the samples, 66% were collected prior to the

introduction of PCV13 in 2011, while 27% were obtained in the post-PCV13 era (S1 Fig). In the pre-

PCV13 era, prominent serotypes with frequencies exceeding 5% included 5 (10.21%), 6B (8.72%), 1

(8.4%), 23F (7.23%), 6A (5.74%), and 16F (5.53%). In the post-PCV13 era, serotypes 1 (11.5%) and 12F

(5.4%) predominated. It is worth noting that serotype 1 exhibited sustained dominance, with its

frequency increasing following the vaccine rollout, while serotype 12F emerged as an abundant strain

after 2011 (S2 Fig). Nevertheless, a more extensive dataset encompassing vaccination information could

offer further insights into this phenomenon.

Within the carriage isolates, serotypes exhibited abundant frequencies, included 19F (7.88%), 6B

(7.27%), 16F (6.79%), and 23F (5.21%), with each surpassing a 5% frequency threshold. The distribution

of serotypes among carriers in Blantyre and Karonga largely mirrored each other, with the exception of

serotype 13, which displayed higher prevalence in Blantyre, and serotype 6B, which exhibited greater

dominance in Karonga (S3 Fig).

Among the blood samples, serotype 5 (20.38%) was the most dominant, followed by 1 (16.58%), 23F

(8.42%), and 6B (5.98%). In the cerebrospinal fluid (CSF) samples, serotypes 1 (21.13%), 5 (8.1%), 12F

(7.04%), 23F (6.69%), 6A (5.63%), and 6B (5.28%) predominated. When considering the combined blood

and CSF groups, the most frequently observed serotypes were 1 (18.56%), 5 (15.03%), 23F (7.67%), 6B

(5.67%), and 6A (5%). It's noteworthy that the majority of invasive samples were collected in Blantyre

(96.5%). The invasive samples from Lilongwe were exclusively CSF samples, with serotypes 1 and 12F

being dominant (S4 Fig).

As depicted in Fig 1 and detailed in Supplementary Table 1 (S1 Table), among the serotypes that were

abundant in either the blood or CSF, serotypes 1 (p = 1.96E-34), 5 (p = 3.97E-19), and 12F (p = 5.29E-06)

exhibited a significant presence among patients, with a low frequency of occurrence in the nasopharynx.

Given that the colonization phase is a prerequisite for infection, it is plausible that serotypes 1, 5, and

12F may have a short period of nasopharyngeal colonization before infecting sterile sites. Considering

that serotypes 1 and 5 were also the most prevalent across the entire cohort, they could be regarded as

the most common serotypes with the highest invasiveness. In this study, we categorize serotypes 1, 5,

and 12F as 'significant invasive serotypes' or 'hyper-invasive serotypes'.

In contrast, abundant serotypes 16F and 19F in the nasopharynx were significantly prevalent among

carriers, suggesting that they may have a lower potential to cause invasive disease. Other frequently

detected serotypes, such as 6A, 6B, and 23F, were both abundant and evenly distributed among carriers

and patients. It is conceivable that they might require a longer period of nasopharyngeal colonization

compared to the hyper-invasive serotypes (1, 5, and 12F) before causing infections at sterile sites.

Serotypes 6A, 6B, and 23F have previously been reported as common and abundant serotypes among

both non-invasive carriers and those with invasive infections [38][39].

Fig 1. The distribution of the 56 pneumococcal serotypes assigned to 1477 samples from Malawi. (a) The relative frequency of each serotype in the nasopharynx of carriers, the blood of bacteremia patients, and the CSF of meningitis patients is shown in blue, red, and yellow, respectively (UT: Un-Typeable). (b) The log-transformed odds ratio of the significantly over- and under-abundant serotypes in

the sterile sites (blood and CSF). Fisher's exact test was applied to identify serotypes with a significant differential abundance among carriers and patients (nasopharynx and sterile sites) at the significance level of the Benjamini-Hochberg adjusted p-value < 0.01 (BH: Benjamini-Hochberg).

As mentioned earlier, the temporal distribution of the hyper-invasive serotypes (1, 5, and 12F)

concerning the vaccine rollout timeline was noteworthy. The relative frequency of serotype 1 exhibited

a significant increase (pre-PCV13=8.6% and post-PCV13=11.3%), while serotype 5 displayed a significant

decrease (pre-PCV13=11% and post-PCV13=1.5%) following the introduction of the vaccination program

in Malawi. This suggests that vaccination may be effective against serotype 5 but did not alleviate the

burden of invasive pneumococcal diseases (IPDs) caused by serotype 1. Additionally, serotype 12F,

which is not included in PCV13, showed a significant increase (pre-PCV13=1.1% and post-PCV13=5.7%)

in the post-PCV13 era, indicating its potential emergence as an invasive strain. Nevertheless, a more

extensive dataset, containing more recent samples, is essential to comprehensively characterize the

long-term effects of PCV13.

High diversity in the pneumococcal pan-genome

The genome assembly produced an average optimized assembly hash value of 96 and an average N50 of

113,986. The mean assembled genome size was estimated to be 2,116,779 nucleotides, with a standard

deviation of 106,481. This length aligns within the range of previously reported *S. pneumoniae* genome

sizes [40].

The pan-genome spanned 5,178,167 base pairs and encompassed 6,803 genes, comprising 729 core

genes (10.7%), 820 soft-core genes (12.1%), and 5,254 accessory genes (77.2%). The pan-genome

remained open, demonstrating a continuous increase in the number of genes as the sample size

expanded (S5 Fig). The gene presence-absence heatmap in Fig 2 illustrates the pan-genome, revealing

serotypes as the primary factor influencing gene distribution. Notably, distinct clustering was observed

for hyper-invasive serotypes 1, 5, and 12F, forming unique clades. Specific sets of genes present in

different serotypes were represented as distinctive blue blocks within the heatmap.

Fig 2. The pan-genome matrix of 1477 pneumococcal isolates from Malawi. The pan-genome is
visualized as a gene presence-absence heatmap representing the hierarchical unsupervised clustering of
samples based on the distribution of genes in the pan-genome. Each row is a sample, and each column
is a gene. A blue dot denotes the presence of each gene. On the right side of the heatmap, the large blue
block shows core genes present in all samples. The left side of the heatmap represents the accessory
genome along with the clustering bands. In addition to the significant serotypes 1, 5, 12F, 16F, and 19F,
other abundant serotypes, including 6A, 6B, and 23F, as well as serotypes with source-based p-value <
0.05, including 21, 11A, and 15B, are also highlighted on the heatmap.

The significant invasive serotypes (1, 5, and 12F) showed the highest distinction in the core-

and accessory-genome

The maximum-likelihood tree, depicting the distribution of small-scale variants (SNPs and indels) within

the core-genome, highlighted the hyper-invasive serotypes 1, 5, and 12F as monophyletic clusters (Fig

3). These serotypes (1, 5, and 12F) were distinct, forming individual clusters that were more prominently

separated compared to other abundant serotypes, such as 6B, 19F, and 23F, which appeared as multiple

clusters on the phylogenetic tree (see Fig 3)

Fig 3. The phylogenetic population structure of 1477 pneumococcal samples from Malawi. The
phylogenetic tree was built based on the multiple sequence alignment of the core genome using the
maximum likelihood method. Colors on the loops show the serotypes, specimen sources (isolation sites),
and PCV13 eras. In addition to the significant serotypes 1, 5, 12F, 16F, and 19F, other abundant
serotypes, including 6A, 6B, and 23F, as well as serotypes with source-based p-value < 0.05, including 21,
11A, and 15B, are also highlighted on the tree.

291   The PCA of the large-scale variants in the accessory-genome (gene presence/absence) displayed

292   serotypes 1 and 5 as distantly clustered from other strains (Fig 4). Additionally, a moderate level of

293   separation was evident for serotypes 12F, 19F, and 23F. This distinct clustering of serotypes 1 and 5 is

294   aligned with profiles demonstrated by the phylogenetic tree.

Fig 4. The PCA of the gene distribution in the pan-genome of pneumococcal isolates from 1477 Malawians. The PCA of variants (gene presence-absence) in the accessory-genome indicates the influence of (a) specimen sources (isolation sites), (b) serotypes, (c) PCV13 (vaccination) era, and (d) geographical locations on the gene presence-absence profile of pneumococcal isolates in Malawi. Serotypes 1 and 5 were clearly separated from other samples.

Figures 3 and 4 indicate that factors such as time, locations, and isolation sites (specimen sources) failed

to sufficiently explain the small- and large-scale genetic variants in the pan-genome. Instead, the

serotype of the samples emerged as the primary driver of the population structure. The hyper-invasive

serotypes (1, 5, and 12F) exhibited the most significant core and accessory distances from other strains,

signaling their genetic distinctiveness among disease-associated serotypes. This heightened genetic

dissimilarity might be linked to their invasive potential.

It's crucial to thoroughly consider population structure and assess any associations with disease across

the population. An important observation is the absence of the same level of genetic distinction in

serotype 12F, potentially due to its smaller sample count (n=35) compared to serotypes 1 (n=129) and 5

(n=116). Additionally, the near-complete separation of serotypes 1 and 5 from the others may primarily

reflect their infrequent presence in the carriage group, suggesting that these patterns might stem more

from sampling biases than genuine genetic variations. To address these concerns, ten samples from

each hyper-invasive serotype and the PCV13 vaccine types were randomly selected from the

nasopharynx, blood, and CSF. A PCA of the gene distribution was performed on the downsampled

dataset, reiterated a noticeable pattern of clustering evident for serotypes 1, 5, and 12F each positioned

far from other strains (S6 Fig).

Another aspect to consider revolves around the differentiation of serotypes. Theoretically, serotypes are

differentiated due to their distinct serotype-defining capsule genes. However, a question emerges

regarding the notable distinction in clustering observed among hyper-invasive serotypes (1, 5, and 12F)

compared to other serotypes. Our investigation focused on the hypothesis that serotypes 1, 5, and 12F

might have undergone gene acquisitions or losses, potentially contributing to their invasiveness. While

there are other prevalent serotypes in blood and CSF, such as 6B and 23F, their similar prevalence in the

nasopharynx suggests they might persist in the nasopharynx for extended durations compared to the

hyper-invasive serotypes (1, 5, and 12F).

The gene presence-absence statistical analysis

The following issues could skew the gene presence-absence analysis:

a) **The batch effect introduced by geographical locations:** 85% of carriage samples were from

Karonga, whereas 95% of disease samples were from Blantyre (Table 1). A comparison between

carriage and disease groups may only identify the difference between pneumococcal genomes

from two geographical locations rather than between the non-invasive and invasive groups.

b) **Study limitation:** The likely presence of invasive serotypes in the carriage group made it unclear

which nasopharyngeal samples progressed to disease after collection. Indeed, the carriage

population likely contained invasive serotypes that could bias the test between carriage and

disease samples to identify potential virulence genes.

c) **Population structure:** The significant abundance of the hyper-invasive serotypes (1, 5, and 12F)

in the patient group and their highest genetic distinction would skew the test between the

carriage and patient groups. The difference between carriage and disease groups would actually

be the difference between the carriage and hyper-invasive serotypes (and not all strains in the

blood and CSF).

To assess the geographical batch effect, carriage isolates from Karonga were compared with those from

Blantyre. The gene presence-absence statistical test did not identify any significant genes differing

between the Blantyre and Karonga groups, indicating a similarity in gene content within the carriage

samples from both locations. Additionally, as previously observed, the serotype distributions in the

carriage groups from Karonga and Blantyre displayed similarities (S3 Fig). Consequently, the impact of

geographical location on the pneumococcal genomes was not substantial. This outcome aligns with

expectations considering that Karonga and Blantyre are approximately 830 kilometers apart, and the

demographic similarities between the populations in these cities.

While serotypes like 6A, 6B, and 23F, potentially associated with invasive traits, were found in both

carriers and patients, comparing the entirety of the carriage and patient groups remains important. This

is because serotypes identified as potential invasive in the nasopharynx might undergo genomic

alterations before reaching sterile sites. During the colonization phase in the nasopharynx, these

serotypes likely engage in genetic exchange via recombination and horizontal gene transfer with other

pneumococci or bacterial species. Therefore, the genomic profile of an invasive serotype in the

nasopharynx might differ from that in the blood and CSF.

To account for these complexities, our analysis involved an association test between the entire carriage

and disease groups, excluding the hyper-invasive serotypes 1, 5, and 12F (a location-based analysis). This

exclusion aimed to prevent these hyper-invasive serotypes from introducing biases when comparing the

carriage and patient groups. The location-based analysis identified 27 significant genes, including 11

genes significantly present in the blood and CSF and 16 genes in the nasopharynx (Table 2, Table 3, and

S2 Table for further details)

The most significant genes identified in both blood and CSF belonged to the cps locus (RD3), suggesting

a potentially increased level of encapsulation during disease. Specifically, genes SP_0357, SP_0358, and

SP_0360 encode epimerases involved in the biosynthesis of complex lipopolysaccharides, which are

essential components of the pneumococcal capsule. SP_0351 encodes a membrane protein

glycosyltransferase responsible for catalyzing glycosyl group transfer during capsule synthesis, and

SP_0359 encodes UDP-2-acetamido-2,6-beta-L-arabino-hexul-4-ose reductase, a crucial protein involved

in capsular polysaccharide biosynthesis. Other significant genes in the blood and CSF, including SP_1953,

SP_0535, SP_1037, and SP_1056, are involved in toxic secretion and recombination. SP_1056 is part of

the pneumococcal pathogenicity island 1 (PPI1) located within RD6. This gene encodes a mobilization

protein necessary for the horizontal transfer of genes and plasmids via bacterial conjugation. SP_1056

plays a role in forming the relaxation complex or relaxosome by interacting with other enzymes [41].

Table 2. Significant genes (p-value < 0.05) present in pneumococci in the blood and CSF (hyper-invasive serotypes 1, 5, and 12F were excluded) compared to the nasopharyngeal pneumococci.

ID	Annotation	P-value	Odds ratio
SP_0360	Capsular polysaccharide biosynthesis protein (from RD3)	8.17E-05	4.897059
SP_0358	Capsular polysaccharide biosynthesis protein (from RD3)	8.17E-05	4.897059
SP_0351	Capsular polysaccharide biosynthesis protein (from RD3)	8.17E-05	4.897059
SP_0359	Capsular polysaccharide biosynthesis protein (from RD3)	8.17E-05	4.897059
SP_0357	Capsular polysaccharide biosynthesis protein	0.000175	4.758421
SP_1037	Type II restriction endonuclease BcgI	0.001074	2.878307
SP_1953	Bacteriocin/lantibiotic secretion ABC transporter permease protein	0.001864	4.631892
SP_0535	Putative immunity protein	0.004674	1.988523
SP_1056	Relaxase/Mobilisation nuclease domain (From RD6)	0.012604	2.932153
SP_1656	Hypothetical protein	0.014050	1.963212
SP_0347	Capsular polysaccharide biosynthesis protein (from RD3)	0.020560	1.808997

367   Significant genes identified in the nasopharynx (absent in samples from blood and CSF) originated from

368   RD10, recognized as the SecY2A2 island responsible for the secretion of pneumococcal serine-rich

369   repeat protein (PsrP) (S7 Fig) [42]. RD10 contains several glycosyltransferases and secretory components

370   that were significantly missing from the genome of samples obtained from blood and CSF.

Table 3. Significant genes (p-value < 0.05) present in the nasopharyngeal pneumococci compared to pneumococci in the blood and CSF (hyper-invasive serotypes 1,5, and 12F were excluded).

ID	Annotation	P-value	Odds ratio
SP_1770	Glycosyl transferase, glyB (from RD10)	0.000132	0.502513
SP_1771	Glycosyl transferase, family 2/family 8 (from RD10)	0.012770	0.560236
SP_1763	Preprotein translocase secY family protein (from RD10)	0.017254	0.563256
SP_1765	Glycosyl transferase, glyF (from RD10)	0.018150	0.564542
SP_0939	Hypothetical protein	0.020740	0.483266
SP_1766	Glycosyl transferase, glyE (from RD10)	0.020740	0.483266
SP_1767	Glycosyl transferase, glyD (from RD10)	0.023196	0.568879
SP_1762	Accessory secretory protein asp1 (from RD10)	0.023196	0.568879
SP_1761	Accessory secretory protein asp2 (from RD10)	0.023196	0.568879
SP_1760	Accessory secretory protein asp3 (from RD10)	0.023196	0.568879
SP_1755	Hypothetical protein	0.031054	0.558554
SP_1757	Glycosyl transferase, glyB (from RD10)	0.031054	0.558554
SP_1764	Glycosyl transferase, glyG (from RD10)	0.023196	0.568879
SP_1768	Conserved hypothetical protein (from RD10)	0.031054	0.558553
SP_1758	Poly(glycerol-phosphate) alpha-glucosyltransferase, tagE (from RD10)	0.031122	0.571941
SP_1759	Preprotein translocase, secA subunit (from RD10)	0.031321	0.574557

To explore the divergence of the hyper-invasive serotypes (1, 5, and 12F), they were compared to

serotypes 16F and 19F, which were significantly present in the nasopharynx (serotype-based analysis).

Serotypes 16F and 19F could represent non-invasive strains better than the whole nasopharyngeal

population that potentially contained some invasive serotypes. Indeed, it was a test between serotypes

with the highest and lowest invasiveness to characterize the genomes of serotypes 1, 5, and 12F. The

gene gain and loss profiles in the hyper-invasive serotypes may include components contributing to their

virulence and short colonization period.

The serotype-based analysis identified 184, 157, and 186 significant genes (present/absent) in serotypes

1, 5, and 12F, respectively (S3 Table, S4 Table, and S5 Table) that were much larger than the number of

significant genes identified by the location-based analysis. The functional enrichment analysis identified

the phosphotransferase system (PTS, KEGG ID: spn02060) as over-represented and oxidative

phosphorylation (KEGG ID: spn00190) as under-represented pathways in the hyper-invasive serotypes

1, 5, and 12F (p-value < 0.05).

In total, there were 18 significant genes jointly present in the hyper-invasive serotypes (Fig 5.a),

including elements of the PTSs that transport sucrose and lactose across the membrane (SP_0302,

SP_0303, SP_0304, SP_0305, SP_0306, SP_0308, SP_0309, and SP_0310), bacteriocins (SP_0544 and

SP_1051) and a permease protein (SP_1527). The over-represented pathway (spn02060) was associated

with significant genes that code for PTS transporters involved in carbohydrate metabolism. Seven genes

were unannotated. The PTS transporters genes were also present in a high proportion of abundant

serotypes in sterile sites such as 6B (67%) and 23F (86%).

A total of 57 significant genes were absent in serotypes 1, 5, and 12F (Fig 5.a). The most significant

absences were observed within RD8a, consisting of two operons, RD8a1 (SP_1315-1324) and RD8a2

(SP_1325-SP_1331) (S8 Fig). RD8a1 harbors eight *ntp* genes, that code *V-type proton/sodium ATP*

synthase complex that produces ATP via oxidative phosphorylation in the presence of a Na+ gradient

across the membrane[43]. RD8a2 includes *neuraminidase, N-acetylneuraminate lyase (nanA),* and *N-*

acetylmannosamine-6-phosphate epimerase (nanE). These genes cleave carbohydrates from the

glycoproteins on the surface of epithelial cells. Other genes in RD8a2 encode the Sodium/solute

symporter subunits that use Na$^+$ gradient to import the carbohydrates [44]. Symporter refers to a

channel that transports the solute (carbohydrates) and co-solute (Na+) in the same direction by utilizing

the energy stored in an inwardly directed sodium gradient. Fundamentally, the genes within RD8a

operons collaborate to generate ATP, cleave carbohydrates from the host epithelial cells, and import

them into the bacterial cell. RD8a was absent in all samples associated with serotypes 1, 5, and 12F,

while it was present in other prevalent serotypes like 6A, 6B, 16F, and 19F. The pathway sp00190, which

was underrepresented in hyper-invasive serotypes, was associated with genes within RD8a. The absence

of RD8a in hyper-invasive serotypes (1,5, and 12F) may be linked to their rapid invasion into the blood

and CSF, where the availability of free oxygen molecules necessary for oxidative phosphorylation is

limited [45].

Other significant genes absent from the hyper-invasive serotypes were from RD4 and RD7. RD4 consists

of a cluster of sortase enzymes responsible for the assembly of pilins into pili and anchoring these

structures and other surface proteins to the cell wall [46][47]. The pilus is a hair-like structure associated

with bacterial adhesion and colonization [48]. Owing to the hypothesized short colonization period of

hyper-invasive serotypes, they may not harness the benefits of RD4 genes involved in pilus assembly.

RD7 genes remain uncharacterized as of the present date.

It is worth mentioning that there were similarities between hyper-invasive serotypes (1,5, and 12F) and

other strains in the blood and CSF. As observed for samples in the blood and CSF (Table 2), capsule

genes were also significantly present in the hyper-invasive serotypes (1,5, and 12F). Moreover, RD10

(previously found to be significantly absent from blood and CSF as described in Table 3) was also absent

from serotypes 1 and 12F. However, RD10 was fully conserved in serotype 5. RD10 was also conserved

in 100% of serotypes 16F and 19F. The summary of the gene present-absent analysis is illustrated in Fig

5.b.

Fig 5. The summary of the gene presence-absence analysis. (a) The number of significant genes present-absent in the hyper-invasive serotypes. The gene presence-absence analysis was applied using Scoary to compare the gene pools of the hyper-invasive serotypes and serotypes 16F and 19F. P-values were corrected by the Bonferroni method, and significant genes had an adjusted p-value of less than 0.05. (b) The significant presence and absence of RDs in samples from blood and CSF is shown as a presence-absence heatmap.

Finally and as mentioned, serotypes 6B and 23F were abundant in both carrier and patient groups, the

intra gene presence/absence statistical test for serotype 6B between the nasopharynx (n=60) and sterile

sites (n=37), and for serotype 23F between the nasopharynx (n=43) and sterile sites (n=50) did not

identify any significant gene. The gene content of these two serotypes in the nasopharynx and sterile

sites was similar.

To highlight genes that might assist pneumococci in crossing the blood-brain barrier, a test was also

conducted between the samples from blood and CSF. The analysis between whole blood (n=368) and

CSF (n=284), serotype 1 samples from blood (n=60) and CSF (n=61), and serotype 5 samples from blood

(n=75) and CSF (n=23) did not identify any significant genes.

Discussion

The *S. pneumoniae* genome is highly diverse, with only a small portion of genes conserved across all

strains. In this species, the pan-genome is open, allowing for an extensive gene repertoire due to the

highly recombinogenic nature of pneumococci. Changes in the *S. pneumoniae* habitat may lead to the

utilization of various gene combinations, enabling organisms to diversify their genome and respond

effectively to environmental stresses. This study found a high genetic diversity, with merely 10.7% of

genes classified as core. These core genes have been conserved across all samples for an extensive

period, at least from 1997 to 2015. Their presence may be crucial for cell survival, making them

potential targets for drug design and vaccine development. Specifically, core genes without any SNPs in

their structure are of particular interest. Notable conserved core genes identified in this study included

SP_1961 (rpoB, DNA-dependent RNA polymerase), SP_0251 (formate acetyltransferase), SP_1891 (amiA,

Oligopeptide binding protein), and SP_0855 (parC, Topoisomerase IV). These conserved core genes play

integral roles in DNA transcription and translation.

Serotypes 1, 5, and 12F exhibited a high prevalence among patients but were rarely found in the carrier

group. This observation indicates their increased invasiveness, likely due to a short duration of

nasopharyngeal colonization. Conversely, serotypes 16F and 19F were significantly more frequent

among carriers, suggesting their dominance in the nasopharynx but with lower invasiveness. Most other

serotypes were common among both carriers and patients. Knowing that pneumococcal virulence

strongly depends on the serotype of isolates, we sought to address why several serotypes were shared

across both nasopharynx and sterile sites. Here, we discuss two possible scenarios that could justify the

ubiquitous presence of some serotypes in both nasopharynx and sterile sites.

The first scenario is related to the colonization of *S. pneumoniae*, which is known as a prerequisite for

virulence[4]. Several samples from the carriage group may be actually the invasive serotypes collected

during their colonization phase. Abundant serotypes such as 6B and 23F that had a similar frequency

amongst carriers and patients need to colonize the upper respiratory tract longer than the hyper-

invasive serotypes before entering the sterile organs. In contrast, the hyper-invasive serotypes 1, 5, and

12F colonize the nasopharynx for a short period and quickly enter the sterile sites.

The second possible scenario relates to the differential gene expression pattern of shared genes in the

ubiquitous serotypes. Although the type-specific *cps* genes were identified in the isolates of both

nasopharynx and sterile sites, the expression pattern of these genes could vary within each strain, which

would contribute to the invasiveness of some strains. Several studies described a cycle of encapsulation

and un-encapsulation amongst pneumococcal strains. Isolates benefit from mutations in the *cps* locus to

either cease or re-start the capsule expression [49]. The lack of a capsule at the epithelial surface

enables the bacterium to expose its surface proteins on the cell wall underneath the capsule and

promote adherence to the host epithelial cells. It has been estimated that 15% of isolates in the upper

respiratory tract are unencapsulated and adhere to the respiratory epithelial cells more efficiently than

encapsulated isolates [50][51]. Lack of capsule also facilitates acquiring virulence and resistance genes

from other isolates. The thick capsule prevents immunoglobulins from interacting with the pathogen

surface proteins during disease. Meanwhile, the negatively charged CPS interferes with the function of

the host phagocytes [52][53]. Taken together, the presence of the *cps* locus in the genome of isolates

assigned to the same serotype does not necessarily reflect the encapsulation of all cells.

Serotypes 1 and 5 can infect all age groups and cause severe IPDs [54]. Serotype 1 is genetically distinct

between different geographical regions [55] and is known as the leading cause of pneumococcal

meningitis in Africa [56][57]. Our findings support previous research showing that serotypes 1, 5, and

12F are the major cause of IPDs in Malawi [58]. Serotype 1 was persistently dominant in pre- and post-

PCV13 eras, serotype 5 was only predominant in pre-PCV3, and serotype 12F emerged after vaccination.

The study also characterized the high genetic distinction of the hyper-invasive serotypes in Malawi by

identifying significant genes that are present or absent in their genome structure compared to other

serotypes. Many of the significant genes present in the nasopharynx and sterile sites were homologous

and had the same function, and there were significant genes with an unknown function (hypothetical

proteins). However, of greatest interest, we did find several significant genes with specific functions that

could explain the difference between the biology of the hyper-invasive serotypes and nasopharyngeal

samples.

Genes within RD8a were significantly absent in the genome of serotypes 1, 5, and 12F. RD8a is known as

a region previously linked to the virulence of serotypes 6B and 14 in the United States [59]. Our study

identified this region's conservation in serotypes 13, 14, 16F, and 19F in Malawi. This observation

prompts the hypothesis that RD8a may be essential for the prolonged colonization of serotypes

contrasting with the quicker colonization of serotypes 1, 5, and 12F. The functions of genes in RD8a

strengthen the assumption to some extent that RD8a may be essential for long nasopharyngeal

colonization. The genes within RD8a, such as neuraminidase, *nanA*, and *nanE*, are involved in the

cleaving of terminal sialic acid residues from mucoglycans and epithelial glycoconjugates. This activity

aids the pathogen in breaching the mucus layer and adhering to the epithelial cells in the nasopharynx.

Additionally, since free carbohydrates are limited in the upper respiratory tract [60], cleaved sialic acid

can be used as the carbon source for metabolism. Moreover, during colonization, secretion of the

pneumococcal toxins elevates the level of sodium ions (Na^+) in the nasopharynx [61], which enables the

sodium-solute symporter in RD8a to import a wide variety of substrates with the sodium ions into the

cell [62]. Most importantly, the *ntp* gene cluster in RD8a encodes the *V-type sodium ATP synthase* that

pumps the extra sodium ions out of the cell [43] and uses the sodium-motive force for oxidative

phosphorylation and ATP synthesis [63]. Oxidative phosphorylation is the final step of aerobic

respiration that requires free oxygen molecules for ATP synthesis. Pneumococci are facultative

anaerobes that can either perform aerobic or anaerobic respiration with or without oxygen. In the upper

respiratory tract, they access atmospheric oxygen molecules that can be used by *ntp* genes to perform

oxidative phosphorylation. However, genes in RD8a may not be beneficial for hyper-invasive serotypes

that supposedly do not stay in the nasopharynx for long. Thus, the level of aerobic ATP synthesis is

presumably higher in serotypes 13, 14, 16F, and 19F in contrast with serotypes 1, 5, and 12F, which lack

RD8a.

Pneumococci can ferment up to 30 types of carbohydrates, imported mainly by two types of membrane

transporters, including ATP-binding cassette (ABC) transporters and PTS transporters [64]. The major

differences between ABC and PTS transporters are: (i) ABC transporters use energy from ATP, but PTS

transporters use energy from phosphoenolpyruvate, and (ii) ABC transporters do not modify the

imported substrate, but PTS transporters phosphorylate the incoming sugar upon transport. Generally,

ABC transporters require more energy than PTS transporters, albeit they can transport longer and more

complicated carbohydrates [65]. Unlike isolates in the nasopharynx, serotypes 1, 5, and 12F have access

to more simple and free host dietary carbohydrates in the blood and the central nervous system. Due to

a potential lower ATP synthesis level in serotypes 1, 5, and 12F (due to the lack of RD8a), they may

prefer to use PTS transporters to uptake sugars such as fructose and lactose, and that is why genes that

encode PTS transporters are significantly more present in the hyper-invasive serotypes (1, 5, and 12F). In

addition to the sugar uptake, PTS transporters regulate several pathways in bacteria, such as gene

expression and communication between cells. Thus, the phenotypic effects of the PTS transporters

should not be limited just to their ability to import carbohydrates [66].

Genes within RD10 were absent from serotypes 1 and 12F. However, they were conserved in serotypes

5, 16F, and 19F. Moreover, the location-based gene presence-absence analysis showed that RD10 was

significantly present in nasopharyngeal samples in comparison to samples collected from sterile sites.

Operon RD10 in pneumococcus shares homology with the general secretion pathway protein B

sceA2/Y2 system components in Streptococcus gordonii, which are involved in secreting the general

secretion pathway protein B linked to infective endocarditis [67]. In the *S. pneumoniae* genome, the

homolog of general secretion pathway protein B is PsrP, which is transported to the bacterial cell

surface by the SecA2/Y2 system encoded by genes in RD10. Research on *Streptococcus gordonii* has

indicated that the presence of SecA2/Y2 facilitates adhesion to both epithelial cells in the nasopharynx

and erythrocytes in the blood. [68][69]. This may explain why SecA/Y2 is significantly present in

nasopharyngeal samples and serotype 5 (abundant in the blood). The presence of the secA2/Y2-like

component should also facilitate the export of pneumolysin, which enhances adhesion to the host cell

and contribute to survival in the blood [70][71].

In conclusion, specific genes present or absent in the hyper-invasive serotypes (1, 5, and 12F) may play a

role in their invasiveness and lower colonization rate. Nonetheless, experimental validation is necessary

to confirm the computational findings from this study. While the serotype is the primary determinant of

the pneumococcal population structure, this research has highlighted the substantial genetic divergence

of serotypes 1, 5, and 12F compared to other serotypes. Their substantial presence in the blood and CSF

accounted for the most pronounced genomic and functional differences observed between the

nasopharynx and sterile sites. The lower frequency of serotypes 1, 5, and 12F among carriers could be

attributed to their shorter colonization duration before entering sterile sites. These invasive serotypes

possess elements of PTS transporters but lack genes from RD8a. Interestingly, RD10 is highly conserved

in serotype 5, while it is absent in serotypes 1 and 12F. Notably, this study demonstrates that isolation

sites do not significantly influence the genomic structure of pneumococcal isolates. Although a few

genes were linked to the virulence of commonly present serotypes in both the nasopharynx and sterile

sites, it is suggested that other high-throughput techniques like gene expression analysis may reveal the

differences between these isolates more comprehensively. In summary, this research sheds light on the

pneumococcal population structure and serotypes in Malawi. The unique cluster of significant genes in

the hyper-invasive serotypes, along with highly conserved core genes, could serve as potential

therapeutic targets.

551 Author contributions

Sample collection, metadata curation, and genome sequencing: Jennifer Cornick, Dean Everett, Anmol
Kiran, Chrispin Chaguza, and Chikondi Peno.

Methodology and data analysis: Arash Iranzadeh, Anmol Kiran, and Arghavan Alisoltani.

Result interpretation: Arash Iranzadeh, Arghavan Alisoltani, Nicola Mulder, and Dean Everett.

Initial manuscript writing: Arash Iranzadeh.

Review of the manuscript: Arash Iranzadeh, Arghavan Alisoltani, Anmol Kiran, Robert F Breiman,
Chrispin Chaguza, Chikondi Peno, Dean B Everett, Nicola Mulder.

All authors have given consent to participate in the study.

560 Acknowledgment

Computations were performed using facilities provided by the University of Cape Town's ICTS High-

Performance Computing team: hpc.uct.ac.za. The authors also acknowledge the Centre for High-

Performance Computing (CHPC), South Africa, for providing computational resources to this research

project. We thank the study participants and all involved staff at the Karonga Prevention Study and the

Malawi-Liverpool-Wellcome Trust Clinical Research Programme. We thank Olivier Koole and Naor Bar-

Zeev for their scientific input.

Funding information

We are grateful for the financial support received from the Wellcome Trust and Institute of Infection and Global Health. The study was supported by Wellcome Trust grant number 079828. We would also like to acknowledge the contribution of Global Pneumococcal Sequencing project (https://www.pneumogen.net/gps/), funded by the Bill and Melinda Gates Foundation, and Pneumococcal African Genomic Consortium (http://www.pagegenomes.org) in the generation of data used in this publication.

Conflicts of interest

The author(s) declare that there are no conflicts of interest.

Ethical approval

Not required as the research delas with bacterial samples.

References

[1] T. D. Swarthout *et al.*, 'High residual carriage of vaccine-serotype Streptococcus pneumoniae after introduction of pneumococcal conjugate vaccine in Malawi', *Nature Communications 2020 11:1*, vol. 11, no. 1, pp. 1–12, May 2020, doi: 10.1038/s41467-020-15786-9.

[2] P. Kamthunzi, 'Impact of PCV13 vaccination in Blantyre, Malawi', *Lancet Glob Health*, vol. 9, no. 7, pp. e893–e894, Jul. 2021, doi: 10.1016/S2214-109X(21)00258-8.

[3] L. Paixão *et al.*, 'Host glycan sugar-specific pathways in streptococcus pneumonia: Galactose as a key sugar in colonisation and infection', *PLoS One*, 2015, doi: 10.1371/journal.pone.0121042.

[4] D. Bogaert, R. de Groot, and P. W. M. Hermans, 'Streptococcus pneumoniae colonisation: The key

to pneumococcal disease', *Lancet Infectious Diseases*. 2004. doi: 10.1016/S1473-3099(04)00938-

7.

[5] E. Backhaus *et al.*, 'Epidemiology of invasive pneumococcal infections: Manifestations, incidence

and case fatality rate correlated to age, gender and risk factors', *BMC Infect Dis*, 2016, doi:

10.1186/s12879-016-1648-2.

[6] A. Kadioglu, J. N. Weiser, J. C. Paton, and P. W. Andrew, 'The role of Streptococcus pneumoniae

virulence factors in host respiratory colonization and disease', *Nature Reviews Microbiology*.

2008. doi: 10.1038/nrmicro1871.

[7] J. N. Weiser, D. M. Ferreira, and J. C. Paton, 'Streptococcus pneumoniae: Transmission,

colonization and invasion', *Nature Reviews Microbiology*. 2018. doi: 10.1038/s41579-018-0001-8.

[8] K. A. Geno *et al.*, 'Pneumococcal capsules and their types: Past, present, and future', *Clin*

Microbiol Rev, 2015, doi: 10.1128/CMR.00024-15.

[9] A. L. Nelson, A. M. Roche, J. M. Gould, K. Chim, A. J. Ratner, and J. N. Weiser, 'Capsule enhances

pneumococcal colonization by limiting mucus-mediated clearance', *Infect Immun*, 2007, doi:

10.1128/IAI.01475-06.

[10] J. Brown, S. Hammerschmidt, and C. Orihuela, *Streptococcus Pneumoniae: Molecular*

Mechanisms of Host-Pathogen Interactions. 2015. doi: 10.1016/C2012-0-00722-3.

[11] F. Ganaie *et al.*, 'A new pneumococcal capsule type, 10D, is the 100th serotype and has a large

cps fragment from an oral streptococcus', *mBio*, 2020, doi: 10.1128/mBio.00937-20.

[12] D. M. Weinberger, R. Malley, and M. Lipsitch, 'Serotype replacement in disease after

pneumococcal vaccination', *The Lancet*. 2011. doi: 10.5455/apd.239006.

[13]   D. R. Feikin *et al.*, 'Serotype-Specific Changes in Invasive Pneumococcal Disease after

609   Pneumococcal Conjugate Vaccine Introduction: A Pooled Analysis of Multiple Surveillance Sites',

610   *PLoS Med*, 2013, doi: 10.1371/journal.pmed.1001517.

611   [14]   E. D. McCollum *et al.*, 'Impact of the 13-valent pneumococcal conjugate vaccine on clinical and

612   hypoxemic childhood pneumonia over three years in central Malawi: An observational study',

613   *PLoS One*, 2017, doi: 10.1371/journal.pone.0168209.

614   [15]   N. Bar-Zeev *et al.*, 'Impact and Effectiveness of 13-Valent Pneumococcal Conjugate Vaccine on

615   Population Incidence of Vaccine and Non-Vaccine Serotype Invasive Pneumococcal Disease in

616   Blantyre, Malawi, 2006-2018: Prospective Observational Time-Series and Case-Control Studies',

617   *SSRN Electronic Journal*, Dec. 2020, doi: 10.2139/SSRN.3745169.

618   [16]   E. Heinsbroek *et al.*, 'Pneumococcal carriage in households in Karonga District, Malawi, before

619   and after introduction of 13-valent pneumococcal conjugate vaccination', *Vaccine*, 2018, doi:

620   10.1016/j.vaccine.2018.10.021.

621   [17]   C. Cillóniz, C. Garcia-Vidal, A. Ceccato, and A. Torres, 'Antimicrobial Resistance Among

622   Streptococcus pneumoniae', in *Antimicrobial Resistance in the 21st Century*, Springer, 2018, pp.

623   13–38.

624   [18]   A. Iranzadeh and N. J. Mulder, 'Bacterial Pan-Genomics', in *Microbial Genomics in Sustainable*

625   *Agroecosystems*, Springer, 2019, pp. 21–38.

626   [19]   A. Embry, E. Hinojosa, and C. J. Orihuela, 'Regions of Diversity 8, 9 and 13 contribute to

627   Streptococcus pneumoniae virulence', *BMC Microbiol*, 2007, doi: 10.1186/1471-2180-7-80.

628   [20]   R. Brückner, M. Nuhn, P. Reichmann, B. Weber, and R. Hakenbeck, 'Mosaic genes and mosaic

629   chromosomes-genomic variation in Streptococcus pneumoniae', *International Journal of Medical*

630   *Microbiology*. 2004. doi: 10.1016/j.ijmm.2004.06.019.

631   [21]   A. W. Kamng'ona *et al.*, 'High multiple carriage and emergence of Streptococcus pneumoniae

632   vaccine serotype variants in Malawian children', *BMC Infect Dis*, vol. 15, no. 1, pp. 1–11, Jun.

633   2015, doi: 10.1186/S12879-015-0980-2/FIGURES/7.

634   [22]   C. Chaguza *et al.*, 'Population genetic structure, antibiotic resistance, capsule switching and

635   evolution of invasive pneumococci before conjugate vaccination in Malawi', *Vaccine*, vol. 35, no.

636   35 Pt B, pp. 4594–4602, Aug. 2017, doi: 10.1016/J.VACCINE.2017.07.009.

637   [23]   S. Andrews, 'FastQC: A quality control tool for high throughput sequence data.', *Babraham*

638   *Bioinformatics*, p. http://www.bioinformatics.babraham.ac.uk/projects/, 2010, doi: citeulike-

639   article-id:11583827.

640   [24]   L. Epping, A. J. van Tonder, R. A. Gladstone, S. D. Bentley, A. J. Page, and J. A. Keane, 'SeroBA:

641   rapid high-throughput serotyping of Streptococcus pneumoniae from whole genome sequence

642   data', *Microb Genom*, vol. 4, no. 7, Jul. 2018, doi: 10.1099/MGEN.0.000186.

643   [25]   S. Gladman and T. Seemann, 'VelvetOptimiser', *Free Software Foundation*. 2008. doi:

644   10.1016/S0925-8574(99)00040-3.

645   [26]   A. Gurevich, V. Saveliev, N. Vyahhi, and G. Tesler, 'QUAST: Quality assessment tool for genome

646   assemblies', *Bioinformatics*, 2013, doi: 10.1093/bioinformatics/btt086.

647   [27]   T. Seemann, 'Prokka: Rapid prokaryotic genome annotation', *Bioinformatics*, vol. 30, no. 14, pp.

648   2068–2069, 2014, doi: 10.1093/bioinformatics/btu153.

[28]   A. J. Page *et al.*, 'Roary: Rapid large-scale prokaryote pan genome analysis', *Bioinformatics*, vol.

31, no. 22, pp. 3691–3693, May 2015, doi: 10.1093/bioinformatics/btv421.

[29]   K. Katoh and D. M. Standley, 'MAFFT multiple sequence alignment software version 7:

Improvements in performance and usability', *Mol Biol Evol*, 2013, doi: 10.1093/molbev/mst010.

[30]   B. Q. Minh *et al.*, 'IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the

Genomic Era', *Mol Biol Evol*, vol. 37, no. 5, pp. 1530–1534, May 2020, doi:

10.1093/MOLBEV/MSAA015.

[31]   I. Letunic and P. Bork, 'Interactive tree of life (iTOL) v3: an online tool for the display and

annotation of phylogenetic and other trees', *Nucleic Acids Res*, 2016, doi: 10.1093/nar/gkw290.

[32]   R. Gaujoux, 'An introduction to NMF package', *BMC Bioinformatics*, 2010, doi: 10.1186/1471-

2105-11-367.

[33]   F. Rohart, B. Gautier, A. Singh, and K. A. Lê Cao, 'mixOmics: An R package for 'omics feature

selection and multiple data integration', *PLoS Comput Biol*, 2017, doi:

10.1371/journal.pcbi.1005752.

[34]   O. Brynildsrud, J. Bohlin, L. Scheffer, and V. Eldholm, 'Rapid scoring of genes in microbial pan-

genome-wide association studies with Scoary', *Genome Biol*, vol. 17, no. 1, p. 238, 2016, doi:

10.1186/s13059-016-1108-8.

[35]   D. Szklarczyk *et al.*, 'STRING v11: Protein-protein association networks with increased coverage,

supporting functional discovery in genome-wide experimental datasets', *Nucleic Acids Res*, 2019,

doi: 10.1093/nar/gky1131.

[36]   H. Ogata, S. Goto, K. Sato, W. Fujibuchi, H. Bono, and M. Kanehisa, 'KEGG: Kyoto encyclopedia of

670   genes and genomes', *Nucleic Acids Research*, vol. 27, no. 1. pp. 29–34, 1999. doi:

671   10.1093/nar/27.1.29.

672   [37]   A. Bateman *et al.*, 'The Pfam protein families database', *Nucleic Acids Res*, vol. 32, no. suppl_1,

673   pp. D138–D141, Jan. 2004, doi: 10.1093/NAR/GKH121.

674   [38]   Q. Wang *et al.*, 'Serotype distribution of Streptococcus pneumoniae isolated from children

675   hospitalized in Beijing children's hospital (2013–2019)', *Vaccine*, vol. 38, no. 49, pp. 7858–7864,

676   2020, doi: 10.1016/j.vaccine.2020.10.005.

677   [39]   M. Alizadeh Chamkhaleh *et al.*, 'Serotype distribution of Streptococcus pneumoniae among

678   healthy carriers and clinical patients: a systematic review from Iran', *European Journal of Clinical*

679   *Microbiology and Infectious Diseases*. 2020. doi: 10.1007/s10096-020-03963-z.

680   [40]   N. L. Hiller *et al.*, 'Comparative genomic analyses of seventeen Streptococcus pneumoniae

681   strains: Insights into the pneumococcal supragenome', *J Bacteriol*, vol. 189, no. 22, pp. 8186–

682   8195, Nov. 2007, doi: 10.1128/JB.00690-07/SUPPL_FILE/SUPPLEMENTARY_TABLE_3.ZIP.

683   [41]   S. Zhang and R. Meyer, 'The relaxosome protein MobC promotes conjugal plasmid mobilization

684   by extending DNA strand separation to the nick site at the origin of transfer', *Mol Microbiol*,

685   1997, doi: 10.1046/j.1365-2958.1997.4861849.x.

686   [42]   C. Obert *et al.*, 'Identification of a candidate Streptococcus pneumoniae core genome and

687   regions of diversity correlated with invasive pneumococcal disease', *Infect Immun*, vol. 74, no. 8,

688   pp. 4766–4777, 2006, doi: 10.1128/IAI.00316-06.

689   [43]   K. Yokoyama and H. Imamura, 'Rotation, structure, and classification of prokaryotic V-ATPase',

690   *Journal of Bioenergetics and Biomembranes*. 2005. doi: 10.1007/s10863-005-9480-1.

691   [44]   J. Reizer, A. Reizer, and M. H. Saier, 'A functional superfamily of sodium/solute symporters',

692   *Biochim Biophys Acta*, vol. 1197, no. 2, pp. 133–166, Jun. 1994, doi: 10.1016/0304-

693   4157(94)90003-5.

694   [45]   P. Gaudu, 'Aerobic respiration metabolism in lactic acid bacteria and uses in biotechnology', doi:

695   10.1146/annurev-food-022811-101255.

696   [46]   C. Manzano *et al.*, 'Sortase-Mediated Pilus Fiber Biogenesis in Streptococcus pneumoniae',

697   *Structure*, 2008, doi: 10.1016/j.str.2008.10.007.

698   [47]   J. Lemieux, S. Woody, and A. Camilli, 'Roles of the sortases of Streptococcus pneumoniae in

699   assembly of the RlrA pilus', *J Bacteriol*, 2008, doi: 10.1128/JB.00379-08.

700   [48]   L. Williams, F. Stapleton, and N. Carnt, 'Microbiology, lens care and maintenance', *Contact*

701   *Lenses*, pp. 65–96, Jan. 2019, doi: 10.1016/B978-0-7020-7168-3.00004-0.

702   [49]   J. D. Langereis and M. I. de Jonge, ' Non-encapsulated Streptococcus pneumoniae , vaccination as

703   a measure to interfere with horizontal gene transfer ', *Virulence*, 2017, doi:

704   10.1080/21505594.2017.1309492.

705   [50]   D. van der Windt, H. J. Bootsma, P. Burghout, C. E. van der Gaast-de Jongh, P. W. M. Hermans,

706   and M. van der Flier, 'Nonencapsulated Streptococcus pneumoniae resists extracellular human

707   neutrophil elastase- and cathepsin G-mediated killing', *FEMS Immunol Med Microbiol*, 2012, doi:

708   10.1111/j.1574-695X.2012.01028.x.

709   [51]   U. M. Talbot, A. W. Paton, and J. C. Paton, 'Uptake of Streptococcus pneumoniae by respiratory

710   epithelial cells', *Infect Immun*, 1996.

711   [52]   C. J. Lee, S. D. Banks, and J. P. Li, 'Virulence, immunity, and vaccine related to streptococcus

712   pneumoniae', *Crit Rev Microbiol*, 1991, doi: 10.3109/10408419109113510.

[53] L. E. Keller, D. A. Robinson, and L. S. McDaniel, ' Nonencapsulated Streptococcus pneumoniae:

Emergence and Pathogenesis ', *mBio*, 2016, doi: 10.1128/mbio.01792-15.

[54] W. P. Hausdorff, 'The roles of pneumococcal serotypes 1 and 5 in paediatric invasive disease',

Vaccine, 2007, doi: 10.1016/j.vaccine.2006.09.009.

[55] J. E. Cornick *et al.*, 'Region-specific diversification of the highly virulent serotype 1 Streptococcus

pneumoniae', *Microb Genom*, vol. 1, no. 2, 2015.

[56] J. Leimkugel *et al.*, 'An outbreak of serotype 1 Streptococcus pneumoniae meningitis in northern

Ghana with features that are characteristic of Neisseria meningitidis meningitis epidemics', *J*

Infect Dis, vol. 192, no. 2, pp. 192–199, 2005.

[57] B. D. Gessner, J. E. Mueller, and S. Yaro, 'African meningitis belt pneumococcal disease

epidemiology indicates a need for an effective serotype 1 containing vaccine, including for older

children and adults', *BMC Infect Dis*, 2010, doi: 10.1186/1471-2334-10-22.

[58] J. E. Cornick *et al.*, 'Invasive streptococcus pneumoniae in children, Malawi, 2004-2006', *Emerg*

Infect Dis, 2011, doi: 10.3201/eid1706.101404.

[59] C. Obert *et al.*, 'Identification of a Candidate Streptococcus pneumoniae core genome and

regions of diversity correlated with invasive pneumococcal disease', *Infect Immun*, vol. 74, no. 8,

pp. 4766–4777, Aug. 2006, doi: 10.1128/IAI.00316-06.

[60] D. M. Walters, V. L. Stirewalt, and S. B. Melville, 'Cloning, sequence, and transcriptional

regulation of the operon encoding a putative N-acetylmannosamine-6-phosphate epimerase

(nanE) and sialic acid lyase (nanA) in Clostridium perfringens', *J Bacteriol*, 1999.

[61] L. E. Bakeeva, K. M. Chumakov, A. L. Drachev, A. L. Metlina, and V. P. Skulachev, 'The sodium

cycle. III. Vibrio alginolyticus resembles Vibrio cholerae and some other vibriones by flagellar

motor and ribosomal 5S-RNA structures', *BBA - Bioenergetics*, 1986, doi: 10.1016/0005-

2728(86)90115-5.

[62]   M. H. Saier, 'Families of transmembrane sugar transport proteins', *Molecular Microbiology*. 2000.

738   doi: 10.1046/j.1365-2958.2000.01759.x.

739   [63]   P. D. Boyer, 'THE ATP SYNTHASE—A SPLENDID MOLECULAR MACHINE', *Annu Rev Biochem*, 2002,

740   doi: 10.1146/annurev.biochem.66.1.717.

741   [64]   A. Bidossi *et al.*, 'A functional genomics approach to establish the complement of carbohydrate

742   transporters in Streptococcus pneumoniae', *PLoS One*, 2012, doi: 10.1371/journal.pone.0033320.

743   [65]   C. M. Buckwalter and S. J. King, 'Pneumococcal carbohydrate transport: Food for thought', *Trends

744   in Microbiology*. 2012. doi: 10.1016/j.tim.2012.08.008.

745   [66]   M. H. Saier, 'The Bacterial Phosphotransferase System: New Frontiers 50 Years after Its

746   Discovery', *Journal of Molecular Microbiology and Biotechnology*. 2015. doi: 10.1159/000381215.

747   [67]   B. A. Bensing, B. W. Gibson, and P. M. Sullam, 'The Streptococcus gordonii Platelet Binding

748   Protein GspB Undergoes Glycosylation Independently of Export', *J Bacteriol*, 2004, doi:

749   10.1128/JB.186.3.638-645.2004.

750   [68]   B. A. Bensing and P. M. Sullam, 'Transport of preproteins by the accessory Sec system requires a

751   specific domain adjacent to the signal peptide', *J Bacteriol*, 2010, doi: 10.1128/JB.00373-10.

752   [69]   M. Yamaguchi *et al.*, 'Streptococcus pneumoniae Invades Erythrocytes and Utilizes Them to

753   Evade Human Innate Immunity', *PLoS One*, 2013, doi: 10.1371/journal.pone.0077282.

[70] M. Bandara *et al.*, 'The accessory Sec system (SecY2A2) in Streptococcus pneumoniae is involved

in export of pneumolysin toxin, adhesion and biofilm formation', *Microbes Infect*, 2017, doi:

10.1016/j.micinf.2017.04.003.

[71] R. Wu and H. Wu, 'A molecular chaperone mediates a two-protein enzyme complex and

glycosylation of serine-rich streptococcal adhesins', *Journal of Biological Chemistry*, 2011,

doi: 10.1074/jbc.M111.239350.

Supporting information

S1 Fig. Characteristics of the 1477 pneumococcal isolates used in the study. (a) The relative frequency

of serotypes in the entire cohort, samples were assigned to 56 serotypes. For each sample, the in-silico

serotyping was accomplished by SeroBA. (b) Frequency of isolates in the pre- and post-PCV13 eras in

Malawi. (c) Frequency of isolates obtained from each specimen source.

S2 Fig. Distribution of the abundant serotypes (frequency > 5%) before and after the vaccination

rollout in Malawi in 2011. Serotype 1 persistently dominated both the pre- and post-vaccination eras.

S3 Fig. The serotype distribution among carriers in Karonga and Blantyre. Distributions were similar,

except for serotype 6B, which was more dominant in Karonga, and serotype 13, which was more

prevalent in Blantyre.

S4 Fig. Serotype distribution among meningitis patients in Lilongwe and Blantyre. Only 3.5% of disease

samples (23 out of 652, i.e., 3.5%) were collected from Lilongwe. Serotypes 1 and 12F were predominant

in both regions; however, a larger dataset from Lilongwe is needed to accurately reflect the true

serotype distribution in this area.

S5 Fig. The pan-genome of 1477 pneumococcal samples isolated in Malawi was obtained from 1997 to

2015. The pan-genome is an open pan-genome, which means the number of total genes increases

unlimitedly when the sample size grows. The dashed line represents the number of total genes, and the

solid line represents the number of conserved genes in the pan-genome.

S6 Fig. The three-dimensional PCA of the gene distribution in the vaccine types. For each serotype and

for downsampling, 10 samples were randomly selected from the nasopharynx, blood, and CSF. The PCA

was conducted using the R package MixOmics. Hyper-invasive serotypes 1, 5, and 12F clustered

separately from other strains.

S7 Fig. Genes in RD10 are absent from serotypes 1 and 12F but conserved in serotype 5, 16F, and 19F.

Genes from RD10 encode the components of the secretory system SecA2/Y2 that transports

glycoproteins to the bacterial cell surface, which are required for binding to the human proteins on the

surface of epithelial cells and erythrocytes.

S8 Fig. RD8a consists of two operons RD8a1 (SP_1315-1324) and RD8a2 (SP_1325-1331). This region is

not detected in the significant invasive serotypes 1, 5, and 12F, but it is present in more than 80% of

serotype 16F and 19F that significantly dominates the nasopharynx. The important biological processes

carried out by these genes are the transport of ions across the membrane and the synthesis of ATP

molecules.

S1 Table. Statistical analysis of serotypes' prevalence across specimen sources.

S2 Table. Gene presence-absence analysis (Invasive vs Nasopharyngeal, serotypes 1, 5, and 12F were

excluded).

S3 Table. Gene presence-absence analysis (Serotype 1 vs 16F & 19F).

S4 Table. Gene presence-absence analysis (Serotype 5 vs 16F & 19F).

S5 Table. Gene presence-absence analysis (Serotype 12F vs 16F & 19F).

Figure captions

Fig 1. The distribution of the 56 pneumococcal serotypes assigned to 1477 samples from Malawi. (a)

The relative frequency of each serotype in the nasopharynx of carriers, the blood of bacteremia

patients, and the CSF of meningitis patients is shown in blue, red, and yellow, respectively (UT: Un-

Typeable). (b) The log-transformed odds ratio of the significantly over- and under-abundant serotypes in

the sterile sites (blood and CSF). Fisher's exact test was applied to identify serotypes with a significant

differential abundance among carriers and patients (nasopharynx and sterile sites) at the significance

level of the Benjamini-Hochberg adjusted p-value < 0.01 (BH: Benjamini-Hochberg).

Fig 2. The pan-genome matrix of 1477 pneumococcal isolates from Malawi. The pan-genome is

visualized as a gene presence-absence heatmap representing the hierarchical unsupervised clustering of

samples based on the distribution of genes in the pan-genome. Each row is a sample, and each column

is a gene. A blue dot denotes the presence of each gene. On the right side of the heatmap, the large blue

block shows core genes present in all samples. The left side of the heatmap represents the accessory

genome along with the clustering bands. In addition to the significant serotypes 1, 5, 12F, 16F, and 19F,

other abundant serotypes, including 6A, 6B, and 23F, as well as serotypes with source-based p-value <

0.05, including 21, 11A, and 15B, are also highlighted on the heatmap.

Fig 3. The phylogenetic population structure of 1477 pneumococcal samples from Malawi. The

phylogenetic tree was built based on the multiple sequence alignment of the core genome using the

maximum likelihood method. Colors on the loops show the serotypes, specimen sources (isolation sites),

and PCV13 eras. In addition to the significant serotypes 1, 5, 12F, 16F, and 19F, other abundant

serotypes, including 6A, 6B, and 23F, as well as serotypes with source-based p-value < 0.05, including 21,

11A, and 15B, are also highlighted on the tree.

Fig 4. The PCA of the gene distribution in the pan-genome of pneumococcal isolates from 1477

Malawians. The PCA of variants (gene presence-absence) in the accessory-genome indicates the

influence of (a) specimen sources (isolation sites), (b) serotypes, (c) PCV13 (vaccination) era, and (d)

geographical locations on the gene presence-absence profile of pneumococcal isolates in Malawi.

Serotypes 1 and 5 were clearly separated from other samples.

Fig 5. The summary of the gene presence-absence analysis. (a) The number of significant genes

present-absent in the hyper-invasive serotypes. The gene presence-absence analysis was applied using

Scoary to compare the gene pools of the hyper-invasive serotypes and serotypes 16F and 19F. P-values

were corrected by the Bonferroni method, and significant genes had an adjusted p-value of less than

0.05. (b) The significant presence and absence of RDs in samples from blood and CSF is shown as a

presence-absence heatmap.

Data summary

832 **S6 Table. Samples IDs on European Nucleotide Archive (ENA)**