1 **Title: A Gamma-adapted recombinant subunit vaccine induces broadly neutralizing**
2 **antibodies against SARS-CoV-2 variants and protects mice from SARS-CoV-2 infection.**
3

Authors:
Lorena M. Coria[a,b]*, Juan Manuel Rodriguez[c,d], Agostina Demaria[a,b], Laura A. Bruno[a,b],
Mayra Rios Medrano[a,b], Celeste Pueblas Castro[a,b], Eliana F. Castro[a,b,e], Sabrina A. Del
Priore[c], , Andres C. Hernando Insua[c,d], Ingrid G. Kaufmann[c], Lucas M. Saposnik[a,b], William
B. Stone[f], Lineia Prado[a,b], Valeria Krum[c], Francisco M. Zurvarra[c,d], Johanna E. Sidabra[c],
Ignacio Drehe[c], Jonathan A. Baqué[c], Mariana Li Causi[c], Analia V. De Nichilo[c,d], Cristian J.
Payes[c], Albert J. Auguste[f,g], Julio C. Vega[h], Diego E. Álvarez[a,b], Juan M. Flo[c], Karina A.
Pasquevich[a,b] and Juliana Cassataro[a,b]*.
12

Author Affiliations:
[a]Instituto de Investigaciones Biotecnológicas, Universidad Nacional de San Martín
(UNSAM) – Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). San
Martín (1650), Buenos Aires, Argentina.
[b]Escuela de Bio y Nanotecnologías (EByN), Universidad Nacional de San Martín. San
Martín (1650), Buenos Aires, Argentina.
[c]Laboratorio Pablo Cassará – Unidad de I+D de Biofármacos. C1440FFX, Ciudad
Autónoma de Buenos Aires, Argentina.
[d]Fundación Pablo Cassará - Unidad de I+D de Biofármacos. C1440FFX, Ciudad Autónoma
de Buenos Aires, Argentina.
[e]Instituto de Virología e Innovaciones Tecnológicas (IVIT), Centro de Investigaciones
en Ciencias Veterinarias y Agronómicas (CICVyA), Instituto Nacional de Tecnología
Agropecuaria (INTA)- Consejo Nacional de Investigaciones Científicas y Técnicas
(CONICET), Buenos Aires, Argentina.
[f]Department of Entomology, College of Agriculture and Life Sciences, Fralin Life
Science Institute, Virginia Polytechnic Institute and State University, Blacksburg, VA,
24061.
[g]Center for Emerging, Zoonotic, and Arthropod-borne Pathogens, Virginia Polytechnic
Institute and State University, Blacksburg, VA, 24061.
[h]Laboratorio Pablo Cassará - I+D+i. C1408GBV, Ciudad Autónoma de Buenos Aires,
Argentina.
34

35 **Corresponding Author:** *Address correspondence to Lorena M Coria
36 lcoria@iib.unsam.edu.ar and to Juliana Cassataro: jucassataro@iib.unsam.edu.ar

38 **Keywords:** RBD-adapted vaccine, COVID-19 subunit vaccine, SARS-CoV-2, heterologous
39 booster.

Abstract

The COVID-19 pandemic continues with the emergence of successive new variants of concern (VOC). One strategy to prevent breakthrough infections is developing safe and effective broad-spectrum vaccines. Here, we present preclinical studies of a RBD recombinant vaccine candidate derived from the Gamma SARS-CoV-2 variant adjuvanted with alum. Gamma RBD-derived antigen elicited better neutralizing antibody and T cell responses than formulation containing ancestral RBD antigen. The Gamma-adapted subunit vaccine elicited a long-lasting antibody response with cross-neutralizing activity against different VOC including the Omicron variant. Additionally, Gamma variant RBD-adapted vaccine elicited robust T cells responses with induction of Th1 and CD8$^+$ T cell responses in spleen and lung. Vaccine-induced immunity protected K18-hACE2 mice from intranasal challenge with SARS-CoV-2 increasing survival, reducing body weight loss and viral burden in the lungs and brain. Importantly, the subunit vaccine demonstrated a potent effect as heterologous booster of different vaccine platforms including the non-replicating adenovirus vaccine ChAdOx1-S, the mRNA vaccine BNT162b2 and the inactivated SARS-CoV-2 vaccine BBIBP-CorV, increasing cross-reactive antibody responses. Our study indicates that the adjuvanted Gamma RBD vaccine is highly immunogenic and a broad-spectrum vaccine candidate to combat SARS-CoV-2 variants including Omicron.

Introduction

Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has infected over 630 million people worldwide and resulted in more than 6,6 million deaths (WHO. World Health Organization, Weekly Operational Update on COVID-19). The vaccines authorized for emergency use or fully approved are safe and highly effective against severe disease[1-5]. However, it has been reported that vaccine-induced protection against symptomatic SARS-CoV-2 infection wanes over time[6,7]. Additionally, as COVID-19 pandemic progresses several variants of concern (VOCs) have emerged, including B.1.351 (Beta), B.1.617.2 (Delta) and P.1 (Gamma), B.1.1.529 (Omicron BA.1) and its subvariants (BA.2.12.1, BA.4/BA.5, BQ.1 and XBB and its descendants). Omicron lineages have been reported to be more resistant to neutralization by vaccine-elicited antibodies and, in some cases are more transmissible than previous VOCs[8-12]

Wanning of vaccine-induced immunity and antibody evasive virus variants create the need for new vaccine strategies that can induce more potent, durable, and broader immune responses to enhance protection against SARS-CoV-2. Additional booster doses of current vaccines have been implemented around the world. Current approved vaccines are mainly directed against the spike protein of the prototype Wuhan-1 SARS-CoV-2 strain therefore their efficacy against certain VOCs is limted[11]. Therefore, vaccines are being adapted to variants as a strategy to improve vaccine effectiveness. While current vaccines may be effective at reducing severe disease to existing VOCs, variant-specific antigens, whether in a mono- or multivalent-vaccine, may be required to induce optimal immune responses and reduce infection against emerging variants.

Most of the variant-adapted vaccines boosters were bivalent or monovalent

containing beta and ancestral spike antigens besides Omicron[13-16]. Interestingly, the

Beta variant originated in South Africa, while the Gamma variant originated in Brazil a

few months later. Until now, there are few studies using a Gamma variant adapted

vaccine [17-19].

In this work, we describe the design, formulation, and immunogenicity of a RBD

vaccine candidate comparing ancestral, and Gamma derived RBD versions. Our results

show that Gamma-adapted RBD vaccine is more immunogenic than ancestral RBD

vaccine in terms of inducing broader neutralizing antibodies (and higher antigen (Ag)

specific cellular immune responses. Therefore, we selected RBD Gamma as antigen to

develop a recombinant subunit vaccine that can be used as primary or booster vaccine.

Here we describe the rational design, generation, and preclinical evaluation of the

Gamma RBD-based vaccine candidate using aluminum hydroxide as adjuvant.

Immunogenicity was evaluated after a primary two-dose vaccine schedule and after a

heterologous booster of different anti-SARS-CoV-2 vaccine platforms including the non-

replicating adenovirus vaccine ChAdOx1-S (Oxford AstraZeneca), the mRNA vaccine

BNT162b2 (Pfizer) and the inactivated SARS-CoV-2 vaccine BBIBP-CorV (Sinopharm).

Results

Design and production of RBD antigens

To compare immune responses elicited with ancestral or Gamma derived RBD,

two formulations were developed using RBD homodimers from the ancestral (Wuhan-

Hu-1) and Gamma variants. Antigens comprise single-chain dimeric repeats of the RBD

protein from amino acids 319R to 537K. The Gamma RBD antigen includes the mutations

described in the Gamma SARS-CoV-2 variant of spike in this region: K417T; E484K and

N501Y (P.1/501Y.V3). High-productivity clones were generated under the genetic

background of a CHO-S (Chinese Hamster Ovary) cell line in high-density suspension cell

cultures. Then, proteins were purified, and antigen purity confirmed by SDS-PAGE and

Western Blot (**Fig. 1A-B**). Comparison of the binding affinity of Gamma and ancestral

RBD to hACE2 receptor was performed by ligand-receptor binding ELISA (**Fig. 1C**). Both

antigens bind to hACE2 at comparable levels with a slight greater binding affinity of

Gamma RBD. Purified antigens were then absorbed to aluminum hydroxide (alum) to

generate the final formulations.

Gamma RBD antigen vaccine formulation improved antibody and cellular immune
responses in comparison with ancestral RBD antigen.

Immunogenicity of vaccine formulations was assessed in BALB/c mice after i.m.

immunization in a two-dose vaccine schedule using 10µg of each antigen absorbed in

aluminum hydroxide (**Fig. 2A**). RBD (Gamma and ancestral)-specific IgG was evaluated

by ELISA in sera of immunized animals. Both formulations induced high levels of IgG

antibodies against the ancestral and Gamma RBD antigens (**Fig. 2B**). Anti-RBD IgG

responses elicited by the formulation containing the Gamma antigen were significatively

higher than that induced by the ancestral RBD-based vaccine on days 14 and 42 after

prime immunization (**Fig. 2C**).

Neutralizing antibody activity was evaluated by a live virus assay using ancestral

or Gamma SARS-CoV-2 variants. Immunization with the formulation containing

ancestral RBD antigen induced higher neutralization titers against ancestral SARS-CoV-2

than against Gamma variant (**Fig. 2D**, $p < 0.05$ NT50 Gamma vs ancestral, time points

from 14 to 35 days post prime immunization). In contrast, vaccination with Gamma RBD

plus Alum elicited high neutralizing antibody (Ab) titers against both viruses. In addition,

cellular immune responses were analyzed in splenocytes from immunized animals by

intracellular flow cytometry. Gamma RBD-based vaccine induced higher frequency of

IFN-γ and TNF-α producing CD4$^+$ and CD8$^+$ T cells (p<0.05, medium alone vs RBD peptides

stimulation) compared to the frequency induced by the formulation with the ancestral

RBD antigen (**Fig. 2E**). These results demonstrated that vaccine containing Gamma RBD

antigen induces a broader neutralizing response and a superior Ag-specific T cellular

immune response than RBD derived from the ancestral SARS-CoV-2. Thus, we decided

to move forward with the formulation containing the Gamma RBD antigen.

Gamma RBD adjuvanted formulation induces long-lasting antibody responses with

broad neutralizing activity against different virus variants of concern

Binding antibody responses were evaluated over longer periods after mice

immunization revealing that anti-RBD IgG titers remained high up to 253 days (more

than 8 months) post prime immunization (**Fig. 3B**). Additionally, neutralization capacity

of antibodies elicited after immunization was evaluated using different VOCs in a live

virus PRNT assay at this time point. Immunization with Gamma RBD adsorbed to Alum

elicited high neutralizing Ab titers against Gamma (GMT 254.8) and ancestral (GMT

312.8) virus. Interestingly, elicited antibodies showed cross-reactivity against Alpha

(GMT 67.82), Delta (GMT 85.49), Omicron BA.1 (GMT 135.5) and BA.5 (GMT 33.12) VOCs

(**Fig. 3B**). Of note, there were no significant differences between these variants

indicating that Gamma RBD vaccination elicits broadly long-lasting neutralizing Ab

responses.

Gamma RBD-adapted vaccine elicits Ag-specific mixed Th1-Th2 immune responses

Figure 2 shows that Gamma RBD adjuvanted formulation induces Ag-specific T

cell responses in splenocytes. The cell-mediated immune responses in spleen and lung

generated after vaccination was further evaluated. Gamma RBD plus alum vaccination

induced Ag-specific IFN-y and IL-5 secretion in spleen and lung cells (**Fig. 4A**) indicating

that a mixed Th1/Th2 is generated at local and systemic level.

In the spleen, antigen-specific IFN-y- and TNF-α-producing CD4$^+$ T cells and IFN-y

producing CD8$^+$ T cells were detected upon Ag stimulation of cells from vaccine

immunized group (**Fig. 4B**). In the lung, immunization with Gamma RBD + Alum elicited

RBD-specific IFN-y producing CD4$^+$ T cells (**Fig. 4C**). Production of cytokines by CD8$^+$ T

cells was not detected in the lungs (data not shown).

A large proportion of antigen-specific cytokine producing CD4$^+$ T and CD8$^+$ T cells

in spleens were double positive IFN-y$^+$ TNF- α^+ and single positive IFN-y$^+$ Th1 cells (**Fig.**

4D-E). Analysis of total CD4$^+$ and CD8$^+$ T cell population that produce cytokines revealed

that induced multifunctionality is greater in the population of CD8$^+$ T cells (**Fig. 4E**).

These results suggest that the Gamma RBD-based vaccine promotes a specific T

cell response biased to a multifunctional Th1 profile and effector CD8$^+$ T cells able to

produce IFN-y in spleen. Interestingly, a Th1 response with production of IFN-y was

observed in the lungs.

Gamma RBD-based vaccine protects mice after intranasal SARS-CoV-2 challenge

To determine vaccine ability to protect against SARS-CoV-2 infection, a lethal K-

18-hACE2 transgenic mice model was used[20,21]. Immunized mice were challenged

intranasally with 2x10^5 PFU of SARS-CoV-2 and monitored daily for body weight changes

and survival, a subset of mice was euthanized on day 5 post-infection (p.i.) to measure

viral loads in the lungs and brains.

Body weight started to decrease in infected placebo mice after 3 days p.i.

reaching a 20% body weight decrease on day 9 p.i. whereas mice immunized with the

vaccine did not show significant weight loss during the experiment (**Fig. 5A**). Moreover,

80% of infected placebo mice died approximately 6 days p.i., 83% of vaccine immunized

mice where alive at this time point. (**Fig. 5B**). After 9 days, all mice in the placebo group

succumb to infection whereas a 66% of the immunized mice survived (P = 0.0208

vaccinated vs placebo).

Five days post infection, a subset of animals was euthanized to assess viral loads

in the lungs and brains. High infectious SARS-CoV-2 titers were observed in the lungs

and brains of the placebo group of mice while low virus titers were detected in brains of

animals vaccinated with Gamma RBD + Alum (**Fig. 5C**). Thus, the Gamma RBD-based

vaccine induced protection against experimental intranasal challenge with SARS-CoV-2

in a mouse model of severe disease.

A booster dose of Gamma RBD-based vaccine induces higher neutralizing Ab titers

than homologous booster vaccination in mice previously primed with different vaccine

platforms.

Booster vaccination has become a strategy to prevent SARS-CoV-2 breakthrough

infections. Immunogenicity of the Gamma RBD adjuvanted vaccine as a heterologous

booster for different primary vaccine platforms was studied in comparison with

homologous regimens (i.e., three dose of the same vaccine). To this end, each group of

mice was immunized with one of the following vaccine platforms in a two-dose scheme

primary vaccination: i) non-replicating adenovirus vaccine ChAdOx1-S (Oxford

AstraZeneca), ii) the mRNA vaccines BNT162b2 (Pfizer) and iii) the inactivated SARS-CoV-

2 vaccine BBIBP-CorV (Sinopharm). Three months later, half of the animals received a

homologous boost and the other half a heterologous boost with the Gamma RBD-based

vaccine (**Fig. 6A**).

All vaccine platforms induced RBD-specific IgG after two dose primary

vaccination (**Fig 6B**). Interestingly, three months after primary vaccination the Gamma

RBD-based vaccine candidate elicited the highest Ab titers (day 98 post prime

immunization, **Fig. 6B**). Three weeks after the third dose, in the group that received a

primary regimen with the inactivated BBIBP-CorV- vaccine the increase in RBD-specific

IgG titers was higher after homologous boosting with Gamma-adapted vaccine than

with the homologous vaccine (18.4× vs. 4×-fold-increase respectively, $p<0.05$). Similarly,

primary vaccination with the non-replicating adenovirus vaccine group ChAdOx1-S and

booster with Gamma RBD vaccination showed a higher fold-increase in anti-RBD IgG

titers than homologous ChAdOx1-S boosting (13.9× vs. 0.65×-fold-increase respectively,

$p<0.001$. **Fig. 6C**). In the group that received mRNA vaccine as primary regimen the

increment in RBD-specific IgG titers was similar after homologous or heterologous

booster (6× vs. 4× fold rise respectively, $p>0.05$. **Fig. 6C**).

Analysis of the neutralizing activity of sera from mice before (pre) and after

booster (post) immunization against ancestral SARS-CoV-2 revealed that the fold-

increase in neutralizing Ab titer were significantly higher after heterologous booster

with the Gamma subunit vaccine than homologous booster with BBIBP-CorV and

ChAdOx1-S vaccines (**Fig. 6D,** $p<0.05$). Heterologous boosting with the Gamma subunit

vaccine induced a significantly higher increase of the neutralizing Abs titers against the

Gamma VOC than homologous boosting in all vaccine platforms used to prime mice (**Fig.**

6D). The best neutralization activity (against ancestral and Gamma variants) was

observed in the group that received primary immunization with the mRNA vaccine

BNT162b2 and boosted with the Gamma RBD-adapted vaccine (**Fig 6D**).

Finally, cross-neutralization activity was assessed against Omicron subvariants

BA.1 and BA.5 three weeks after the booster dose. In all groups, antibody neutralization

capacity was reduced against Omicron subvariants compared to neutralizing Ab titers

against the ancestral and Gamma variant. However, a degree of cross-reactive

antibodies against BA.1 and BA.5 was detected, and the groups boosted with the subunit

vaccine outperformed groups boosted with homologous vaccines (adenoviral,

inactivated and mRNA vaccines, **Fig. 6 D-E**). In particular, in the group that received

mRNA as primary vaccination, the neutralizing Ab titers against Omicron BA.1 and BA.5

were significantly higher after a heterologous boost with Gamma RBD-adapted subunit

vaccine than with a homologous boost vaccination **(Fig. 6D-E)**.

These results suggest that the Gamma RBD-adapted recombinant vaccine

candidate could serve as an effective booster of different primary vaccine platforms

inducing a broad cross-reactive immunity.

Discussion

The rapid emergence of SARS-CoV-2 variants has challenged the global

development of vaccines. The ability of VOCs to escape neutralization by antibodies

elicited by infection, vaccination, or therapeutic application has been widely

studied[22,23]. Developing variant adapted vaccines that can elicit broad and strong

immune responses against these predominant circulating variant(s) is proposed as one

of the best strategies to address variants immune evasion.

In this study, we described the immunogenicity of a recombinant adapted

subunit vaccine formulated with aluminum hydroxide to be used as a primary or booster

vaccine. Ancestral and Gamma variant derived RBD antigens were compared in their

capacity to induce neutralizing Ab responses against SARS-CoV-2. It was reported that

sera from individuals vaccinated with an ancestral RBD antigen plus alum vaccine

formulation showed a reduction in neutralizing antibody titers of 1.7x and 2.1x fold

against Gamma and Beta variants of SARS-CoV-2 respectively[24]. In agreement with these

results ancestral RBD could not induce good neutralizing activity against the Gamma

SARS-CoV-2 variant whereas we demonstrated for the first time that the vaccine

formulation containing Gamma RBD was able to induce neutralizing Abs against

ancestral and Gamma SARS-CoV-2 variants.

Beta and Gamma variants, which originated in South Africa and Brazil,

respectively, carry the K417N/T, E484K and N501Y amino acid changes in the spike RBD.

Mutations in the Beta variant decreased the neutralization sensitivity of convalescent

and vaccinated sera by 6x-fold, while mutations in Gamma decreased neutralizing

activity by 3x-fold [25]. Mono and bivalent candidates containing different combinations

of VOCs based antigens have been assessed in humans [13,26]. Omicron-adapted mRNA

vaccines as boosters have been implemented in several countries. Recently, a Beta-

adapted subunit vaccine has been approved for use in Europe. Until present, there are

no clinical studies with a Gamma adapted vaccine.

The Gamma variant booster vaccine candidate tested here showed a broad cross-

reactive neutralizing antibody response against several VOCs including Omicron in mice.

This could be related to the notion that K417N/T and E484K substitutions shared by Beta

and Gamma viruses simulate a structure that enhances the spectrum of neutralizing

activity. In agreement with this hypothesis and our results, a study evaluating the

immunogenicity of an S-protein sequence containing these three substitutions in

addition to D614G (B.1) demonstrated that the variant vaccine is capable of eliciting high

neutralizing Ab titers that cross-reacted with five VOCs in NHP[27]. Moreover, in preclinical

and clinical trials Beta derived vaccines induced broadly neutralizing antibodies that can

cross-react against different VOC [27,28] in a similar way than the Gamma derived vaccine

presented in this work.

A recombinant vaccine containing RBD-dimers derived from ancestral SARS-CoV-

2 and aluminum hydroxide as adjuvant proved to be safe and immunogenic in humans[29].

Structural analysis of these RBD-dimers indicates that are arranged in a way where the

receptor binding motif (RBM) is correctly presented and the major antigenic sites of RBD

are exposed[30]. This supports the versatility of the RBD-dimers as a module to adapt

different variants for the induction of broader immune responses. Hence, it has been

reported that chimeric RBD-dimers elicit broader responses to variants compared with

homodimers[30]. Increasing evidence suggests that vaccine protection against COVID-19

might be waning over time, and newly emerging variants, such as Delta and Omicron,

might evade vaccine-induced immune protection. Our vaccine induced a long-lasting

antibody response with neutralizing activity against different VOCs up to 253 days after

the last immunization. Adjuvanted recombinant COVID-19 vaccine formulations have

reported   durability   of   the   neutralizing-antibody   responses   for   7   months   post-

immunization in NHP[31]. This study together with previous reports suggest that

recombinant variant-specific vaccines may be needed for improved and long-lasting

protection against known and emerging VOCs.

Main correlates of protection against COVID-19 are binding and neutralizing

antibody titers[32]. However, cellular immunity is associated with positive outcomes[33].

Here, we have provided a comprehensive assessment of the T cell response to a Gamma-

adapted recombinant subunit vaccine. Gamma RBD + alum vaccination elicited a robust

Th1-dominated T cell response and RBD-specific CD8$^+$ T cell response. Analysis of

polyfunctionality on vaccine elicited T cells suggests that the major proportion of cells

produce one or two cytokines (IFN-γ and TNF-α). Polyfunctional Th1 responses were

reported after human vaccination with adenoviral ChAdOx1-S and the mRNA BNT162b2

vaccine[34,35]. Protection studies in transgenic mice presented here confirm that the

immunity induce by the Gamma-derived RBD vaccine candidate can protect mice from

a lethal nasal challenge of ancestral SARS-CoV-2 virus.

Updated immunization strategies were implemented as booster shots across the

globe. In this work, we showed that vaccination with Gamma RBD plus alum as booster

of adenoviral, inactivated or mRNA-based vaccines induced a higher binding and

neutralizing antibody titer increase against different VOCs than vaccinations with

homologous vaccines. In general, heterologous boosters showed higher vaccine

effectiveness than homologous boosters for all outcomes in humans, providing

additional support for a mix-and-match approach[36,37]. Our vaccine not only increased

GMT values of neutralizing Ab against ancestral and Gamma virus variants but also

against Omicron lineages (BA.1 and BA.5).

The Gamma-RBD vaccine candidate reported in this article has demonstrated to

be safe in non-clinical safety studies. Both the antigen and the formulated vaccine are

being produced under GMP at manufacturing plants of Laboratorio Pablo Cassará S.R.L.

located in Buenos Aires. Stability studies showed that the vaccine can be stored and

handled in refrigerated conditions (2 – 8 °C) and does not need freezing temperatures.

The development of a vaccine does not require stringent cold-chain transport and

storage facilitates its availability at local and global supply. This gamma RBD plus alum

vaccine was named ARVAC Cecilia Grierson and is currently in a Phase 1 clinical trial

(NCT05656508).

This work highlighted the value of a Gamma variant-adapted vaccine to induce

higher and broader immune responses against SARS-CoV-2 at booster vaccination and

set the precedent for its further clinical development.

Methods

Antigen expression and purification

Constructs from Wuhan (ancestral) and Gamma SARS-CoV-2 variants of the antigen

consisted of a single-chain dimer of the receptor binding domain (RBD), comprising

amino acids 319R to 537K of the Spike protein from ancestral and Gamma virus variant.

The signal peptide sequence from the SARS-CoV-2 Spike protein was used for protein

secretion. The constructs were codon-optimized for CHO cell expression and

synthesized by GenScript (Hong Kong Limited, Japan). All constructs were cloned into

the pJC3UMCS-4 expression vector (Pablo Cassara Foundation) comprising a CMV

promoter and a cis-acting sequence for minimized host gene silencing. Linearized forms

of the vectors were transfected into CHO-S cells by lipofection. After 15-20 days of G418

antibiotic selection, an end point dilution procedure was applied for clone isolation.

Clone candidates were selected by specific and volumetric productivity assessed by

ELISA. Recombinant protein relative abundance was confirmed by SDS-PAGE and

Coomasie Brilliant Blue G-250 staining.

Pilot batches of antigen and vaccines were manufactured by Laboratorio Pablo Cassará,

Argentina. Controlled cell substrates were used to inoculate a 2-L bioreactor with 1.4 L

as working volume. An alternate tangential flow system (ATF, Repligen) was used for

perfusion. The harvest after 20 - 25 days was used as starting material for a downstream

process based on three chromatographic steps: first an affinity mix-mode capture

chromatography step followed by two different ionic chromatography to eliminate

residual host cell DNA and host cell proteins and other process-related impurities. TFF

(Tangential flow filtration) was used to adjust pH, conductivity, and protein

concentrations of the batches.

SDS-PAGE and western blot analysis

RBD samples were run under reducing and non-reducing conditions by SDS-PAGE (10%

polyacrylamide). The samples were visualized by staining with Coomassie Brilliant Blue

G-250. Bands were then transferred to a nitrocellulose membrane (Bio-Rad Inc.

Hercules, CA, USA), blocked with Non-fat milk 5% in phosphate buffered saline (PBS)-

Tween 0.05%, and incubated with anti-RBD rabbit polyclonal serum (1/1000 dilution).

An anti-rabbit IgG horseradish-peroxidase conjugated (1/2000 dilution) was used as a

secondary antibody (Agilent DAKO, Santa Clara, CA, USA) and 4-Chloro-1-Naphthol as

substrate for colorimetric detection.

ACE2 binding to RBD analyzed by titration ELISA

ACE2 binding to the produced RBD variant antigens was analyzed by ELISA. Briefly, 0.1ug

of RBD per well was immobilized on a High binding Microplatte (Greiner Bio-One GmbH,

Austria.) and incubated ON at 4°C. Next, plates were blocked with 5% non-fat milk in

PBS 0.05% Tween20 for 2 h at 37°C and then washed with PBS and 0.05% Tween20.

Plates were then incubated with hACE2 (R&D, Minneapolis MN, USA) for 2 h at 37°C and

another washing step was performed. Finally, plates were incubated with anti-hACE2

(R&D Systems) diluted in 1% non-fat milk PBS 0.05% Tween 20 for 2 h at 37°C. Detection

was performed with a secondary Ab conjugated to HRP (Agilent DAKO, Santa Clara, CA,

USA) and visualized with tetramethylbenzidine (TMB) substrate (BD biosciences,

Franklin Lakes, New Jersey, USA).

Adjuvant and formulations

Pilot batches of vaccines were manufactured using aluminum hydroxide as adjuvant.

The antigen was adsorbed onto the adjuvant by mixing. Free antigen was controlled to

be less than 10% of total antigen in the vaccine.

Ethics statement

All experimental protocols with animals were conducted in strict accordance with

international ethical standards for animal experimentation (Helsinki Declaration and its

amendments, Amsterdam Protocol of welfare and animal protection and National

Institutes of Health, NIH USA, guidelines.). The protocols performed were also approved

by the Institutional Committee for the use and care of experimental animals (CICUAE)

from National University of San Martin (UNSAM) (01/2020).

Animals and immunizations

Eight-week-old female BALB/c mice were obtained from IIB-UNSAM animal facility.

Animals were intramuscularly (i.m) inoculated at day 0 and 14 with i) Gamma RBD (10μg)

+ Alum (n=5), ii) ancestral RBD (10μg) + Alum (n=5), iii) placebo (n=4). In other

experiments, i) Gamma RBD (10μg) + Alum (n=5) and ii) placebo (n=5). Blood samples

were collected weekly to measure total and neutralizing antibody titers and at day 42

post prime immunization animals were sacrificed and spleens and lungs harvested.

Determination of antibody levels in serum

RBD-specific antibody responses (IgG) were evaluated by indirect ELISA as described

previously[38]. Briefly, plates were coated with 0.1 μg/well of RBD (derived from ancestral

or Gamma variant) in phosphate buffered saline (PBS) overnight at 4 °C. Plates were

washed and then incubated with diluted sera and then plates were washed and

incubated with HRP conjugated anti-mouse IgG (SIGMA, St. Louis, MO, USA). Results

were read at 450 nm to collect end point ELISA data. End-point cut-off values for serum

titer determination were calculated as the mean specific optical density (OD) plus 3

standard deviations (SD) from sera of saline immunized mice and titers were established

as the reciprocal of the last dilution yielding an OD higher than the cut-off.

Viruses

Ancestral SARS-CoV-2 (B.1, GISAID Accession ID EPI_ISL_16290469), Alpha (B.1.1.7

GISAID Accession ID EPI_ISL_15806335), Gamma (P.1, GISAID Accession ID

EPI_ISL_15807444) and Omicron BA.5 (GISAID Accession ID EPI_ISL_16297058) were

isolated from nasopharyngeal specimens at the Instituto de Investigaciones

Biotecnológicas (IIB, UNSAM) and adapted in Vero E6 cultures. Delta SARS-CoV-2

(GISAID Accession ID: EPI_ISL_11014871) and Omicron BA.1 (GISAID Accession ID

EPI_ISL_10633761) were isolated at Instituto de Investigaciones Biomédicas en

Retrovirus y SIDA (INBIRS, UBA-CONICET) from nasopharyngeal swabs of patients.

Studies using SARS-CoV-2 were done in a Biosafety level 3 laboratory at IIB, UNSAM, and

the protocol was approved by the INBIRS Institutional Biosafety Committee.

SARS-CoV-2 neutralization assay

Serum samples were heat-inactivated at 56°C for 30 min. Serial dilutions were

performed and then incubated for 1 h at 37ºC in the presence of 300 $TCID_{50}$ of SARS-

CoV-2 in DMEM 2% FBS. One hundred µl of the mixtures were then added onto Vero

cells monolayers. After 72 h at 37ºC and 5% CO_2, cells were fixed with PFA 4% (4°C

overnight) and stained with crystal violet solution in methanol. The cytopathic effect

(CPE) of the virus on the cell monolayer was assessed by surface scanning at 585 nm in

a microplate reader (FilterMAx F5 Microplate reader, Molecular Devices, San Jose, CA,

USA). Average optical density (OD) of each well was used for the calculation of the

percentage of neutralization of viral CPE for each sample as: Neutralization% = 100 x (DO_{sample}

$_{sample}$ - $DO_{virus\ control}$) / ($DO_{cell\ control}$ - $DO_{virus\ control}$). Non-linear curves of Neutralization

(%) vs. Log 1/sera dilution were fitted to obtain the titer corresponding to the 50% of

neutralization (NT50).

Determination of T cell immune responses

Four weeks after the second dose, mice were sacrificed to study cellular responses.

Splenocytes were cultured for 5 days in the presence of RBD antigen (ancestral or

Gamma) or complete medium. Then, supernatants were collected, and cytokines (IFN-y

and IL-5) were measured by ELISA (Biolegend. San Diego, CA). For intracellular cytokine

determination: splenocytes were cultured (4x10^6 cells/well) in the presence of stimulus

medium (complete medium supplemented with anti-CD28 and anti-CD49d) or Ag stimuli

(stimulus medium + RBD-peptides + RBD protein) for 18 h. Next, brefeldin A was added

for 5 h to the samples. Cells were then washed, fixed, permeabilized, stained, and

analyzed by flow cytometry. The cells were stained with Viability dye (Zombie Acqua),

anti-mouse-CD8a Alexa Fluor 488, anti-mouse-CD4 Alexa Fluor 647, anti-IL-4 Brilliant

Violet 421, anti-TNFα PeCy7 and anti-IFN-γ PE (Biolegend. San Diego, CA). BD Fortessa

LSR-X Cytometer with DIVA Software were used for analysis.

SARS-CoV-2 challenge of vaccinated K18-hACE2 mice

Four-week-old K18-hACE2 mice from Jackson Laboratory were used for evaluating

vaccine efficacy. Mice were separated into two groups: i) control (n=8) Placebo (PBS)

and Gamma RBD +Alum (n=9). Mice in each group included males and females. They

were i.m immunized on day 0 and 14 as described for immunogenicity assays. Four

weeks post second vaccination, mice were challenged intranasally (i.n.) with 10^5 PFU of

SARS-CoV-2 strain WA1/2010 split evenly in each nare. Mice were monitored daily for

weight loss and signs of disease for two weeks post-challenge. Three mice per group

were euthanized on day 5 post-challenge to evaluate organ viral loads, by plaque assay

on Vero E6 cells as previously described[38].

Statistical analysis

Statistical analysis and plotting were performed using GraphPad Prism 8 software

(GraphPad Software, San Diego, CA). In experiments with more than two groups, data

were analyzed using one-way ANOVA with Kruskal Wallis or Bonferroni post-test. When

necessary, a logarithmic transformation was applied prior to the analysis to obtain data

with a normal distribution. In experiments with two groups, an unpaired t test or Mann–

Whitney U test were used. A p value <0.05 was considered significant. When bars were

plotted, results were expressed as means ± SEM for each group. Survival curves were

analyzed using Long-rank test.

Data availability

The authors declare that the data supporting the findings of this study are available from

the corresponding author upon reasonable request.

Acknowledgements

The authors are grateful to the technical staff at the I+D Biofármacos Laboratorio Pablo

Cassará and to Fundación Pablo Cassará for their significant assistance in the

preparation and control of the antigen pilot lots and the regulatory documentation:

Cintia Parsza, Brenda Heinrich, Melisa Gambone, Horacio Descoins, Christian Cortez,

María Victoria Román and Romina Albarracín. The authors would also like to thank

Danielle Porier, Krisangel López, and Manette Tanelus for technical assistance with the

SARS-CoV-2 challenge studies. We thank Ministerio de Salud de la Provincia de Buenos

Aires for providing mRNA, adenoviral and inactivated vaccines.

Contributions

L.M.C. was responsible of overall experiment design, conducted humoral and cellular

experiments, collected data, performed data analysis, and wrote the manuscript. J.C.

was responsible for overall experimental design and conceptualization, wrote the

manuscript and received funding support. J.M.R. generated high expression vectors,

designed the antigen and was responsible of development of the vaccine formulation

and downstream development. Coordination of antigen and vaccine production. K.A.P.

and J.M.F. participated of experimental design, data analysis and reviewed the

manuscript. D.E.A. designed and supervised neutralization studies and analysis. A.D.

conducted and analyzed humoral studies and conducted cellular studies. C.P.C., L.A.B,

L.M.S and L.P. conducted humoral and cellular studies. M.R.M. and E.F.C. performed

virus neutralization studies and data analysis. S.A.DP. generated high expression vector

and optimization of high productive clones. A.C.HI. Responsible of upstream and

downstream development of antigens. I.G.K. coordinated overall analytics procedures,

developed validation and analytics methodologies, performed structural

characterization of antigen and vaccine formulation. A.V.DN. generated cell banks,

performed stability, clones purity and identity studies. V.K. and I.D. generated,

characterized, and screened cell clones. F.M.Z., J.A.B. and C.J.P performed upstream and

downstream development of antigen. J.E.S. and M.LC. participated in the development

of validation and analytics methodologies, performed structural characterization of

antigen and vaccine formulation and performed animals studies. J.C.V. participated in

the development of vaccine formulation, collaborates in the overall process of vaccine

development. A.J.A. design, supervise and conduct animal challenge studies and data

analysis. W.B.S. performed animal challenge studies and data analysis. All authors

revised the manuscript.

Competing interest

J.M.R., A.C.HI., F.M.Z. are salaried employees of Fundacion Pablo Cassara. S.A.DP., I.G.K.,

A.V.DN., V.K., I.D., C.J.P., J.A.B., J.E.S., M.LC., J.M.F. and J.C.V. are salaried employees of

Laboratorio Pablo Cassara. L.M.C., J.C., K.A.P., D.E.A., A.D., C.P.C., L.A.B, L.M.S., L.P.,

M.R.M., E.F.C. A.J.A. and W.B.S. declare no competing interests relevant to this article.

Funding

This work was supported by Laboratorio Pablo Cassará and grants from Agencia Nacional de Promoción de la Investigación, el Desarrollo Tecnológico y la Innovación (AGENCIA I+D+i) and Ministerio de Ciencia, Tecnología e Innovación (FONARSEC 0001) to J.C. and from the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under Award Number R01AI153433 to A.J.A.

References

1   Baden, L. R. *et al.* Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *New England Journal of Medicine* **384**, 403-416, doi:10.1056/nejmoa2035389 (2021).

2   Bos, R. *et al.* Ad26 vector-based COVID-19 vaccine encoding a prefusion-stabilized SARS-CoV-2 Spike immunogen induces potent humoral and cellular immune responses. *npj Vaccines* **5**, doi:10.1038/s41541-020-00243-x (2020).

3   Polack, F. P. *et al.* Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *New England Journal of Medicine* **383**, 2603-2615, doi:10.1056/nejmoa2034577 (2020).

4   Voysey, M. *et al.* Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *The Lancet* **397**, 99-111, doi:10.1016/s0140-6736(20)32661-1 (2021).

5   Al Kaabi, N. *et al.* Effect of 2 Inactivated SARS-CoV-2 Vaccines on Symptomatic COVID-19 Infection in Adults. *JAMA* **326**, 35, doi:10.1001/jama.2021.8565 (2021).

6   Chemaitelly, H. *et al.* Waning of BNT162b2 Vaccine Protection against SARS-CoV-

2 Infection in Qatar. *New England Journal of Medicine* **385**, e83,

doi:10.1056/nejmoa2114114 (2021).

7   Goldberg, Y. *et al.* Waning Immunity after the BNT162b2 Vaccine in Israel. *New*

England Journal of Medicine **385**, e85, doi:10.1056/nejmoa2114228 (2021).

8   Wang, Q. *et al.* Antibody evasion by SARS-CoV-2 Omicron subvariants BA.2.12.1,

BA.4 and BA.5. *Nature* **608**, 603-608, doi:10.1038/s41586-022-05053-w (2022).

9   Khan, K. *et al.* Omicron BA.4/BA.5 escape neutralizing immunity elicited by BA.1

infection. *Nature Communications* **13**, doi:10.1038/s41467-022-32396-9 (2022).

10   Tuekprakhon, A. *et al.* Antibody escape of SARS-CoV-2 Omicron BA.4 and BA.5

from vaccine and BA.1 serum. *Cell* **185**, 2422-2433.e2413,

doi:10.1016/j.cell.2022.06.005 (2022).

11   Tseng, H. F. *et al.* Effectiveness of mRNA-1273 against SARS-CoV-2 Omicron and

Delta variants. *Nature Medicine* **28**, 1063-1071, doi:10.1038/s41591-022-01753-

y (2022).

12   Suryawanshi, R. K. *et al.* Limited cross-variant immunity from SARS-CoV-2

Omicron without vaccination. *Nature* **607**, 351-355, doi:10.1038/s41586-022-

04865-0 (2022).

13   Chalkias, S. *et al.* A Bivalent Omicron-Containing Booster Vaccine against Covid-

19. *New England Journal of Medicine* **387**, 1279-1291,

doi:10.1056/nejmoa2208343 (2022).

14   Akache, B. *et al.* Immunogenicity of SARS-CoV-2 spike antigens derived from Beta

& Delta variants of concern. *npj Vaccines* **7**, doi:10.1038/s41541-022-

00540-7 (2022).

15   Sridhar, S. *et al.* The potential of Beta variant containing COVID booster vaccines

for chasing Omicron in 2022. *Nature Communications* **13**, doi:10.1038/s41467-

022-33549-6 (2022).

16   Launay, O. *et al.* Immunogenicity and Safety of Beta-Adjuvanted Recombinant

Booster Vaccine. *New England Journal of Medicine* **387**, 374-376,

doi:10.1056/nejmc2206711 (2022).

17   Sir Karakus, G. *et al. Preclinical efficacy and safety analysis of gamma-irradiated*

inactivated SARS-CoV-2 vaccine candidates (Cold Spring Harbor Laboratory,

2020).

18   Turan, R. D. *et al.* Gamma-irradiated SARS-CoV-2 vaccine candidate, OZG-

38.61.3, confers protection from SARS-CoV-2 challenge in human ACEII-

transgenic mice. *Scientific Reports* **11**, doi:10.1038/s41598-021-95086-4 (2021).

19   Khorattanakulchai, N. *et al.* Receptor binding domain proteins of SARS-CoV-2

variants produced in <i>Nicotiana benthamiana</i> elicit neutralizing antibodies

against variants of concern. *Journal of Medical Virology*, doi:10.1002/jmv.27881

(2022).

20   Winkler, E. S. *et al.* SARS-CoV-2 infection of human ACE2-transgenic mice causes

severe lung inflammation and impaired function. *Nature Immunology* **21**, 1327-

1335, doi:10.1038/s41590-020-0778-2 (2020).

21   Jiang, R.-D. *et al.* Pathogenesis of SARS-CoV-2 in Transgenic Mice Expressing

Human Angiotensin-Converting Enzyme 2. *Cell* **182**, 50-58.e58,

doi:10.1016/j.cell.2020.05.027 (2020).

22   Harvey, W. T. *et al.* SARS-CoV-2 variants, spike mutations and immune escape.

Nat Rev Microbiol **19**, 409-424, doi:10.1038/s41579-021-00573-0 (2021).

23 Tao, K. *et al.* The biological and clinical significance of emerging SARS-CoV-2

variants. *Nat Rev Genet* **22**, 757-773, doi:10.1038/s41576-021-00408-x (2021).

24 Zhao, X. *et al.* Neutralisation of ZF2001-elicited antisera to SARS-CoV-2 variants.

Lancet Microbe **2**, e494, doi:10.1016/S2666-5247(21)00217-2 (2021).

25 Brinkkemper, M. *et al.* A third SARS-CoV-2 spike vaccination improves

neutralization of variants-of-concern. *NPJ Vaccines* **6**, 146, doi:10.1038/s41541-

021-00411-7 (2021).

26 Branche, A. R. *et al. SARS-CoV-2 Variant Vaccine Boosters Trial: Preliminary*

Analyses (Cold Spring Harbor Laboratory, 2022).

27 Wang, Z. *et al.* A potent, broadly protective vaccine against SARS-CoV-2 variants

of concern. *NPJ Vaccines* **7**, 144, doi:10.1038/s41541-022-00571-0 (2022).

28 Chalkias, S. *et al.* Safety, immunogenicity and antibody persistence of a bivalent

Beta-containing booster vaccine against COVID-19: a phase 2/3 trial. *Nat Med*

28, 2388-2397, doi:10.1038/s41591-022-02031-7 (2022).

29 Dai, L. *et al.* Efficacy and Safety of the RBD-Dimer-Based Covid-19 Vaccine ZF2001

in Adults. *N Engl J Med* **386**, 2097-2111, doi:10.1056/NEJMoa2202261 (2022).

30 Xu, K. *et al.* Protective prototype-Beta and Delta-Omicron chimeric RBD-dimer

vaccines against SARS-CoV-2. *Cell* **185**, 2265-2278 e2214,

doi:10.1016/j.cell.2022.04.029 (2022).

31 Grigoryan, L. *et al.* Adjuvanting a subunit SARS-CoV-2 vaccine with clinically

relevant adjuvants induces durable protection in mice. *NPJ Vaccines* **7**, 55,

doi:10.1038/s41541-022-00472-2 (2022).

32 Gilbert, P. B. *et al.* A Covid-19 Milestone Attained - A Correlate of Protection for

Vaccines. *N Engl J Med*, doi:10.1056/NEJMp2211314 (2022).

33   Moss, P. The T cell immune response against SARS-CoV-2. *Nature Immunology*

23, 186-193, doi:10.1038/s41590-021-01122-w (2022).

34   Guerrera, G. *et al.* BNT162b2 vaccination induces durable SARS-CoV-2-specific T

cells with a stem cell memory phenotype. *Sci Immunol* **6**, eabl5344,

doi:10.1126/sciimmunol.abl5344 (2021).

35   Swanson, P. A., 2nd *et al.* AZD1222/ChAdOx1 nCoV-19 vaccination induces a

polyfunctional spike protein-specific T(H)1 response with a diverse TCR

repertoire. *Sci Transl Med* **13**, eabj7211, doi:10.1126/scitranslmed.abj7211

(2021).

36   Jara, A. *et al.* Effectiveness of homologous and heterologous booster doses for

an inactivated SARS-CoV-2 vaccine: a large-scale prospective cohort study.

Lancet Glob Health **10**, e798-e806, doi:10.1016/S2214-109X(22)00112-7 (2022).

37   Rouco, S. O. *et al.* Heterologous booster response after inactivated virus BBIBP-

CorV vaccination in older people. *Lancet Infect Dis* **22**, 1118-1119,

doi:10.1016/S1473-3099(22)00427-3 (2022).

38   Coria, L. M. *et al.* A Novel Bacterial Protease Inhibitor Adjuvant in RBD-Based

COVID-19 Vaccine Formulations Containing Alum Increases Neutralizing

Antibodies, Specific Germinal Center B Cells and Confers Protection Against

SARS-CoV-2 Infection in Mice. *Front Immunol* **13**, 844837,

doi:10.3389/fimmu.2022.844837 (2022).

Figure 1

Figure 1. Characterization of RBD antigens. A. Non-reduced (nr) and reduced (r) SDS-PAGE migration profiles of the pooled samples (Gamma and ancestral RBD) are shown. **B**. **Western blot.** Non-reduced and reduced RBD antigens were detected using a rabbit polyclonal anti-RBD antibody. **C**. Binding affinity of Gamma and ancestral RBD proteins to immobilized human ACE2 receptor by ligand-receptor binding ELISA. Experiments were performed in duplicates and mean ± SD values are shown.

Figure 2

Figure 2. **Gamma RBD-adapted vaccine induce antibodies with a broad neutralizing activity and Ag-specific T cell responses**. **A**. Immunization protocol scheme. BALB/c mice were immunized at day 0 and day 14 via i.m. with: ancestral RBD + Alum (n=5), Gamma RBD + Alum (n=5). Serum samples were obtained at indicated time points for ELISA and neutralization assays. **B**. Kinetics of RBD-specific IgG endpoint titers in sera of immunized animals by ELISA. Points are means ± SEM. **C**. RBD-specific IgG titers in sera of immunized animals at day 14 and 42 post prime immunization. *p<0.05. **p<0.01. Mann Whitney test. **D**. Neutralizing-antibody titers in sera was determined by authentic PRNT SARS-CoV-2 assay for each group of vaccinated mice at different time points. The black lines represent the geometric mean (GMT) of all data points. Titers correspond to the 50% of virus neutralization (NT50). **E**. Mice were sacrificed 42 days after the first immunization to obtain spleens and lungs and T cell response was evaluated by intracellular flow cytometry. Splenocytes were stimulated with complete medium or RBD-peptides pool for 18 h and then brefeldin A was added for 5 h. Afterward, cells were harvested and stained with specific Abs anti-CD8, and anti-CD4, fixed, permeabilized, and stained intracellularly with anti–IFN-γ and anti-TNF-α. Results are presented as percentage of IFN-γ or TNF-α – producing T lymphocytes. Bars are means ± SEM. *p < 0.05 vs. medium. T test.

Figure 3

Figure 3. **Gamma RBD plus alum vaccinated mice induce a long-lasting antibody response with broaden neutralization activity**. BALB/c mice were immunized at day 0 and day 14 via i.m. with: Gamma RBD + Alum (n=5). **A**. Kinetics of RBD-specific (ancestral or Gamma) IgG endpoint titers in sera of immunized animals by ELISA until 253 days post prime immunization (dpp). Points are means ± SEM. **B.** Neutralizing-antibody titers against ancestral SARS-CoV-2 and different VOC (Alpha (B.1.1.7), Gamma (P.1) Delta (B.1.617.2) and Omicron (BA.1 and BA.5) were evaluated 253dpp in Gamma RBD +Alum immunized group. Neutralization titer was defined as the serum dilution that reduces 50% the cytopathic effect (NT50).

Figure 4. Gamma RBD formulated with alum induces T cell responses in mice. Cellular response was evaluated in spleen and lung 28 days after last immunization. **A**. Frequency of secreted IFN-γ and IL-5 following splenocytes or lung cells stimulation with medium or recombinant RBD were determined by ELISA. Bars are means ± SEM of pg/ml of IFN-γ and IL-5 after subtracting the amount in medium stimulated cells. *p < 0.05, **p < 0.01. T test. **Intracellular flow cytometry analysis of cytokine secreting T cells**. Splenocytes (**B**) and lung cells (**C**) were stimulated with complete medium or a peptide pool derived from RBD and then brefeldin A was added. Afterward, cells were harvested and stained with specific Abs anti-CD8, and anti-CD4, fixed, permeabilized, and stained intracellularly with anti–IFN-γ, TNF-α and anti-IL-2. Results are presented as percentage of cytokine-producing T cells. Bars are means ± SEM. *p < 0.05 vs. medium. T test. **D**. **Assessment of polyfunctional T cells**. Frequency of CD4+ and CD8+ T cells population from splenocytes that produce one, two or three cytokines were plotted for each group of mice. **E**. Pie graphs indicate the total proportion of RBD specific cytokine production in splenocytes from the Gamma RBD + Alum immunized mice.

Figure 5

Figure 5. **Vaccination with Gamma RBD + Alum protects K18-hACE2 transgenic mice against intranasal SARS-CoV-2 infection**. Mice received PBS (Control) (n=8) or Gamma RBD + Alum (n=9) administered via i.m. route at day 0 and 14. Four weeks following immunization, K18-hACE2 mice were intranasally infected with 2×10^5 PFU of SARS-CoV-2. **A**. Weight changes in mice were monitored daily until day 14 after infection. Points are means ± SEM of percentage of original weight. **B**. Mice survival were monitored daily until day 14. Each dot represents the percentage of mice alive at that time point. Survival curves were analyzed with Log-rank (Mantel-Cox) test. P=0.0208 vs placebo. **C**. Five days after infection lungs and brains (n=3) were obtained from groups of mice and SARS-CoV-2 virus was titrated. Bars represent the mean ± SEM. **p < 0.01 and ***p < 0.001. T test.

Figure 6

Figure 6. Heterologous boosting with Gamma RBD-adapted vaccine formulation of different primary vaccine platforms. **A**. Immunization protocol scheme. BALB/c mice were immunized at day 0 and day 21 via i.m. with: Gamma RBD + Alum (n=5), ChAdOx1-S (n=10), BNT162b2 (n=10) and BBIBP-CorV (n=10). At day 100 (M3) half of mice received homologous third doses (BNT162b2, BBIBP-CorV or ChAdOx1-S accordingly to table in A) and the other half heterologous booster shots with Gamma RBD + Alum. **B**. RBD-specific IgG titers in sera of immunized animals at day 98 post prime immunization. Bars are GMT ± 95% CI. *p<0.05. **p<0.01. Kruskal Wallis. Bars are GMT ± 95% CI. Numbers in bold are fold-increase in GMT values. **C**. Comparison of RBD-specific IgG titers pre and post-boost immunization. Numbers in bold are GMT fold-increase values. Bars are GMT ± 95% CI *p<0.05. ***p<0.001. Kruskal Wallis. **D**. Neutralizing-antibody titers against ancestral and Gamma SARS-CoV-2 were evaluated pre (98dpp) and post boost immunization (121 dpp) of mice. Neutralization titer was defined as the serum dilution that reduces 50% the cytopathic effect (NT50). Bars are GMT ± SD. Numbers in bold are fold-increase GMT values for each group. *p<0.05. ***p<0.001. Kruskal Wallis test. **E**. Neutralizing activity of antibodies against omicron (BA.1 and BA.5) were evaluated 21 days post boost immunization of mice. Neutralization titer was defined as the serum dilution that reduces 50% the cytopathic effect (NT50). Bars are GMT ± SD and numbers are GMT values for each group of mice. *p<0.05. ***p<0.001. Kruskal Wallis test.