Multi-Omics Atlas-Assisted Discovery of Transcription Factors for Selective T Cell State Programming

H. Kay Chung[1,2,3*#], Cong Liu[4*], Anamika Battu[2], Alexander N. Jambor[4], Brandon M. Pratt[2,9], Fucong Xie[2], Brian P. Riesenberg[2], Eduardo Casillas[1], Ming Sun[1], Elisa Landoni[2], Yanpei Li[2], Qidang Ye[2], Daniel Joo[2], Jarred Green[2], Zaid Syed[2], Nolan J. Brown[2], Mattew Smith[2], Shixin Ma[1], Shirong Tan[1], Brent Chick[5], Victoria Tripple[1], Z. Audrey Wang[2,4], Jun Wang[4], Bryan Mcdonald[1], Peixiang He[6,7], Qiyuan Yang[1], Timothy Chen[1], Siva Karthik Varanasi[1], Michael LaPorte[1], Thomas H. Mann[1], Dan Chen[1], Filipe Hoffmann[1], Josephine Ho[5], Jennifer Modliszewski[2], April Williams[8], Yusha Liu[2], Zhen Wang[11], Jieyuan Liu[11], Yiming Gao[12], Zhiting Hu[11], Ukrae H. Cho[2,3], Longwei Liu[6,7], Yingxiao Wang[6], Diana C. Hargreaves[5], Gianpietro Dotti[2,10], Barbara Savoldo[2,10], Jessica E. Thaxton[2,3], J. Justin Milner[2,10], Susan M. Kaech[1#], Wei Wang[4#]

[1]NOMIS Center for Immunobiology and Microbial Pathogenesis, Salk Institute for Biological Studies, La Jolla, CA, USA
[2]Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA
[3]Department of Cell Biology & Physiology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA
[4] Department of Chemistry and Biochemistry, University of California San Diego, La Jolla, CA, USA.
[5] Molecular and Cell Biology laboratory, Salk Institute for Biological Studies, La Jolla, CA, USA
[6]Alfred E. Mann Department of Biomedical Engineering, University of Southern California, Los Angeles, CA, USA
[7]Department of Bioengineering, Institute of Engineering in Medicine, University of California, La Jolla, CA, USA
[8]Razavi Newman Integrative Genomics and Bioinformatics Core, Salk Institute for Biological Studies, La Jolla, CA, USA
[9]Department of Pharmacology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA.
[10]Department of Microbiology and Immunology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA.
[11]Halıcıoğlu Data Science Institute, University of California San Diego, La Jolla, CA, USA.
[12]Department of Electrical & Computer Engineering, Texas A&M University
* Equal contribution, # Co-corresponding

CD8+ T cells differentiate into diverse states that shape immune outcomes in cancer and chronic infection. To systematically define the transcription factors (TFs) driving these states, we built a comprehensive atlas integrating transcriptional and epigenetic data across nine CD8+ T cell states and inferred TF activity profiles. Our analysis catalogued TF activity fingerprints of each state, uncovering new regulatory mechanisms that govern differentiation of selective cell states. Leveraging this platform, we focused on two transcriptionally similar but functionally opposing states critical in tumor and viral contexts: terminally exhausted T cells (TEX_term), which are dysfunctional, and tissue-resident memory T cells (T_RM), which are protective. Global TF community analysis revealed distinct state-specific biological pathways and TF-driven networks that govern protective versus dysfunctional T cell states. Through _in vivo_ CRISPR screening integrated with single-cell RNA sequencing (_in vivo_ Perturb-seq), we delineated TF-driven networks that govern protective versus dysfunctional T cell states. We identified HIC1 and GFI1 as shared regulators of TEX_term and T_RM differentiation and KLF6 as a unique regulator of T_RM. Importantly, we discovered novel TEX_term single-state TFs, including ZSCAN20 and JDP2 with no prior known function in T cells. Targeted deletion of these TFs enhanced tumor control and synergized with immune checkpoint blockade. Consistently, their depletion in human T cells reduces the expression of inhibitory receptors and improves effector function. By decoupling exhaustion-selective from protective T_RM programs, our platform enables more precise engineering of T cell states, advancing rational design of effective immunotherapies.

Introduction

Cell states can be described as the range of cellular phenotypes arising from a defined cell type's interaction with its environment. Within the immune system, T cells possess multiple differentiation states, particularly as naive T cells differentiate into diverse states with different functionality and trafficking patterns in various immune environments, such as tumors and virus infections[1–4]. As transcription factors (TFs) govern cell state differentiation[5,6], understanding the mechanisms by which TFs influence cell states is vital for programming beneficial cell states for therapeutic applications[7,8]. One promising clinical application of cell state engineering is the enhancement of CD8$^+$ T cell-based therapies, including the adoptive transfer of tumor-infiltrating lymphocytes (TILs) or chimeric antigen receptor-T (CAR-T) cells. However, identifying the crucial TFs that control different CD8$^+$ T cell states is challenging due to the significant heterogeneity among these states and the similarities in their transcriptomes and gene regulatory networks, even when these cell states are functionally distinct.

The transcriptional similarity between cell states with differing functionalities is exemplified by two key CD8$^+$ T cell states involved in controlling solid tumors and viruses: the protective tissue-resident memory (T_{RM}) cell state and the dysfunctional terminally exhausted (TEX_{term}) cell state. Many studies have shown that TILs with T_{RM} characteristics correlate with better survival in solid tumor patients[9–13]. Conversely, during persistent antigen stimulation scenarios such as chronic virus infection (e.g. HIV) or cancer, T cells progressively express diverse inhibitory receptors such as PD1, lose memory potential, and eventually effector functions. This process is known as T cell exhaustion (TEX)[14–17], and cells in this trajectory eventually adopt the TEX_{term} cell state. TEX_{term} cells express higher amounts of a diverse set of inhibitory receptors (e.g., TIM3, and CD101), lack effector and proliferation capacity, and do not respond very effectively to immune checkpoint blockade (ICB) such as anti-PD1 monoclonal antibody (mAb) blockade[18–20]. High TEX_{term} marker expression often indicates poor prognosis in solid tumors, though some markers also correlate with ICB response, highlighting their complex role in tumor immunity[21–24]. Despite their distinct functional impact on cancer outcomes, TEX_{term} and T_{RM} cells both preferentially reside in tissues[1,3] and display remarkable similarities in their transcriptional profiles, including key regulatory TFs such as BLIMP1[14,25–27], BHLHE40[28,29], and NR4A2[9,30,31](Fig. 1a,b, Extended Data Fig. 1a-c). These two cell states even exhibit highly correlated open chromatin regions (Extended Data Fig. 1d), complicating the precise identification of TFs whose disruption may selectively inhibit TEX_{term} while preserving T_{RM} cell development. Given that many TFs are commonly expressed

2

across different CD8[+] T cell states and differentiation trajectories, a sophisticated and precise

bioinformatics approach is crucial to pinpoint the bona fide cell-state-specifying TFs essential for T cell

programming.

We hypothesized that the challenges of identifying key TFs in control of selective CD8[+] T cell

differentiation states can be effectively addressed through a systematic comparison of TF activity across

the T cell differentiation landscape. Two considerations were postulated for accurate TF activity

prediction: first, TF activity is not always proportional to the TF expression level, as it is influenced by

post-translational modifications, co-factors, and target accessibility[32]. Second, the effect of a TF on

target genes propagates over the genetic network, and thus, the global impact of the TF activity needs

to be assessed at the systems level. To address these considerations, a multi-omics atlas-assisted platform

was developed, integrating transcriptomic and chromatin data of nine major CD8[+] T cell states. This

platform analyzes transcriptional networks and genome-wide, "global" influencers in each cell state and

identifies cell-state "*selective*" or "*shared*" TFs. Our platform also enables integrative analysis of TF

communities and TF target genes (referred to as "regulatees"), providing insight into how the milieu of

TFs in each cell state collectively guides specific differentiation states.

Taiji-based multiomics atlas identifies cell-state–selective and multi-state TFs in CD8[+] T cells

Elucidating the TFs and their regulatory networks that govern each cell state is crucial for accurately

programming specific cellular states of interest. Our initial objective was to create a comprehensive

catalogue of TF activity across diverse CD8[+] T cell states by integrating our TF activity analysis

pipeline, Taiji[33–35], with comparative statistical analysis. In Taiji, the gene regulatory network (GRN) is

a weighted, directed network that models regulatory interactions between TFs and their target genes. In

this GRN, each node corresponds to a gene, and its weight is proportional to the gene's expression level.

Each edge represents a regulatory interaction and is weighted based on a combination of factors: the

TF's predicted binding affinity to the target gene, the chromatin accessibility at the target gene's locus,

and the expression levels of both the TF and the target gene (Fig. 1c)[33,34]. To determine the global

influence of each TF within the network, Taiji applies a personalized PageRank algorithm, which assigns

an "importance" score to each node based on both the quantity and quality of incoming connections.

This approach yields a measure of TF activity that reflects the influence of each TF in the broader

regulatory landscape, accounting for upstream regulators, downstream targets, and feedback loops

through iterative computation.

With Taiji, we have identified TFs involved in pan-immune lineage commitment, including NK cells,

dendritic cells, B cells, and γδ T cells[35]. While earlier studies provided foundational insights into cell

differentiation, a more refined analysis is needed to achieve higher resolution to contextualize TF roles

within CD8$^+$ T cell type. Therefore, leveraging the improved statistical filtering in our updated version

of Taiji, we aimed to quantify the global influence of TFs across all CD8$^+$ T cell states.

To begin, we analyzed ATAC-seq and RNA-seq datasets from 121 CD8$^+$ T cell samples spanning nine

distinct states, using both previously published and newly generated datasets from well-characterized

acute and chronic lymphocytic choriomeningitis virus (LCMV) infections[3,9,20,27,36–40] (Extended Data

Fig. 2; Extended Data Table 1). In acute LCMV-Armstrong infections, CD8$^+$ T cells differentiate into

memory precursor (MP), terminal effector (TE), effector memory (T$_{EM}$), central memory (T$_{CM}$), and

T$_{RM}$ states. In chronic LCMV-Clone 13 infections, they adopt heterogeneous exhaustion cell states,

including progenitors of exhaustion (TEX$_{prog}$), effector-like exhaustion (TEX$_{eff}$), and TEX$_{term}$.

Next, we conducted an unbiased comparative analysis using statistical filtering to understand the

specificity of TF activity across the nine major CD8$^+$ T cell states (Extended Data Fig. 3a, Extended

Data Table 2). This identified a total of 255 TFs of which 136 TFs were classified as predominantly

'**single-state**' TFs, with each cell state selectively containing ~12-19 unique TFs (Fig. 1d, Extended

Data Fig. 3b). This category included many novel TFs such as *Hoxa7* in Naïve, *Snai1* in T$_{RM}$, *Hey1* in

TEX$_{prog}$, *Sox8* in TEX$_{eff}$, and *Zscan20* in TEX$_{term}$ cells. In contrast, 119 TFs were classified as key

regulators in more than one cell state, termed '**multi-state**' TFs (Fig. 1e), such as *Tcf7* and *Tbx21*. *Tcf7*

is a known driver of Naive-, MP-, and TEX$_{prog}$ states, all of which are multipotent and have high

proliferative capacity[3,20]. Multi-state TFs like *Vax2*, *Batf*, *Irf8*, and *Stat1* were more enriched within the

set of exhaustion-associated cell states (TEX$_{prog}$, TEX$_{eff}$, and TEX$_{term}$). This analysis of cell-state

selectivity also shows that T$_{RM}$ shares the most TFs with TEX$_{term}$ compared to other cell states (e.g.,

Egr2, *Crem*, *Prdm1*, Extended Data Fig. 3c) in agreement with their high degree of transcriptional and

epigenetic similarity (Fig. 1b; Extended Data Fig. 1b-d).

While Taiji provides a statistically grounded approach for inferring TF activity, there is no absolute
threshold for defining cell state specificity, and some misclassification is expected, particularly for TFs
with overlapping functions or modest differences in activity. Still, Taiji is useful to highlight TFs with
activity patterns enriched in a given cell state. For instance, although *Eomes* is classified as TEX_{term}
single-state TF here, it also functions in effector, T_{EM}, T_{CM}, and T_{RM} differentiation[41–43], illustrating that
more accurate classifications require further investigation and resolution, as performed herein for several
TFs.

Comparative analysis of TFs in TEX_{term} and T_{RM} identifies candidate TFs for state-selective T cell
programming

Despite the strong transcriptional overlap between TEX_{term} and T_{RM} cells, our Taiji pipeline possesses
the predictive power to generate a catalogue of TFs selectively active in either of these two cell states.
This may aid in developing better antitumor or antiviral immunotherapies wherein one can engineer T
cells away from T cell exhaustion and toward more functional effector cell states without negatively
impacting their ability to form into T_{RM} cells in tissues and tumors. Based on statistical criteria (Extended
Data Fig. 3a), we identified 20 and 34 TFs as single-state TFs of T_{RM} or TEX_{term} cells, respectively, and
30 multi-state TFs active in both cell states (Fig. 1f–h; Extended Data Fig. 3a, blue boxes; Extended
Data Table 3). TEX_{term} single-state TFs included many previously unreported TFs such as *Zscan20*,
Jdp2, Zfp324, Zfp143, Zbtb49, and *Arid3a* (Fig. 1f). T_{RM} single-state TFs included *Fosb, Zfp692, Atf4,*
Pbx4, Junb, and Klf6 (Fig. 1g). Of the TEX_{term} and T_{RM} multi-state TFs, some were well-known to
function in the development of *both* cell states, such as *Nr4a2*[12], *Bhlhe40*[28], *Prdm1*[27,39], whereas others
such as *Hic1*[44] and *Gfi1*[45] were not, identifying them as new multi-state TFs to consider (Fig. 1h). We
also analyzed previously reported TFs such as cJUN, BATF/BATF3, and TFAP4 that were identified
from functional screening of $CD8^+$ T cells[43–48] based on limited phenotypic readouts. These prior screens
tended to identify broadly active, multi-state TFs (see Fig. 1). In contrast, our platform enabled a
computationally guided, multi-state screen that revealed TFs predicted to have greater state-selective
activity (Extended Data Fig. 3d).

To evaluate the predicted TFs governing selective T cell differentiation pathways, we identified dynamic
activity patterns of TF groups, termed "TF waves" (Extended Data Fig. 4). TF waves reveal potential
combinations of TFs that govern developmental trajectories coordinately. Seven TF waves associated

with distinctive biological pathways were identified. For example, the T_{RM} TF wave (Fig. 1i) includes

several members of the AP-1 family (e.g. *Atf3*, *Fosb*, *Fosl2*, and *Jun*), which aligns well with a recent

report of their roles in T_{RM} formation[46]. This wave was uniquely linked to the transforming growth factor

beta (TGFβ) response pathway (Extended Data Fig. 4e). Conversely, the TEX TF wave (including

TEX_{prog}, TEX_{eff}, and TEX_{term}) was characterized by a distinct set of TFs, such as *Irf8*, *Jdp2*, *Nfatc1,* and

Vax2, among others, that correlated with pathways related to PD1 and senescence (Fig. 1j, Extended

Data Fig. 4e).

TF community analysis reveals TF-TF networks and biological processes selectively associated

with T_{RM} or TEX_{term} cell states

To uncover transcriptional programs governing T_{RM} or TEX_{term} differentiation, we constructed TF–TF

association networks capturing functional relationships between TFs (Fig. 2a). Analysis of regulatee

based adjacency matrices revealed shared and distinct patterns of TF collaboration across the two states.

Single-state TFs showed strong intra-state interactions; TEX_{term} single-state TFs like ZSCAN20, JDP2,

ZFP324, and IRF8 formed dense intra-state subnetworks in TEX_{term} cells (Fig. 2b; Extended Data Fig.

5a). Similarly, T_{RM} single-state TFs such as FOSB, SNAI1, and KLF6 interacted majorly within T_{RM}

cells (Extended Data Fig. 5b). Multi-state TFs (e.g., HIC1, PRDM1, FLI1, and GFI1) were active in

both states but partnered with distinct TFs depending on the context, indicating they may operate within

regulatory networks unique to each cell state (Fig. 2c; Extended Data Fig. 5c).

We next grouped the TF–TF association networks into distinct "TF neighbor communities" in T_{RM} and

TEX_{term} cells (Extended Data Table 5), and found each community linked to specific biological

processes (Fig. 2d–f). While multi-state TFs shaped overall community topology, single-state TFs drove

unique interaction patterns specific to T_{RM} or TEX_{term} cells within each community. Pathway analysis

revealed that these communities regulate divergent programs in each state—for instance, T_{RM}

community 3 was associated with cell adhesion and TGFβ response (Fig. 2e, g, h), while TEX_{term}

community 3 was linked to apoptosis (Fig. 2f-h). Community 1 in T_{RM} cells controlled RNA metabolism

(Fig. 2e, g, h), whereas in TEX_{term} cells, this community was tied to catabolism, proteolysis, and

autophagy (Fig. 2f-h).

To assess the functional relevance of state-enriched pathways, we focused on the proteasome pathway,

which emerged as a prominent but previously unrecognized feature of TEX$_{term}$ cells (Fig. 2g, h).

Proteasome gene signatures were also enriched in TEX$_{term}$-like CD8$^+$ T cells from NSCLC patients[47]

and murine MCA-205 TILs (Extended Data Fig. 6a,b). Consistently, proteasome activity—measured by

a validated fluorescent probe[48]—was highest in TEX$_{term}$ cells from chronic LCMV (Fig. 2i) and in

tumor-specific TILs (Fig. 2j, k) relative to bystander OT-1 cells (Fig. 2k). To test if high proteasome

activity correlates with dysfunction, we sorted OT-1 cells by proteasome activity probe intensity and

adoptively transferred them into B16F10-OVA tumor-bearing mice. Proteasome^{high} cells showed

reduced tumor control compared to proteasome^{low} cells (Fig. 2l), a trend also seen in endogenous TILs

(Extended Data Fig. 6c). These findings help validate the TF-TF association network and pathway

predictions and identify the ubiquitin-proteasome pathway as a functional hallmark of TEX$_{term}$ cells.

Systematic validation of TEX$_{term}$-driving TFs via *in vivo* CRISPR screening coupled with scRNA-

seq

The Taiji pipeline enabled comparative analysis of TF activity and curated sets of single-state TFs

specific to T$_{RM}$ *vs.* TEX$_{term}$ cells (Fig. 1f-h). To evaluate its accuracy, Perturb-seq, which combines *in*

vivo CRISPR screening with single-cell RNA sequencing (scRNA-seq), was first conducted in separate

infectious environments that promote the differentiation of TEX$_{term}$ and T$_{RM}$ cells, namely chronic and

acute LCMV infection (Fig. 3a and 4a). Our Perturb-seq guide RNA (gRNA) library targeted 19 TFs,

including seven multi-state TFs predicted to impair both TEX$_{term}$ and T$_{RM}$ differentiation and 12 TEX$_{term}$

single-state TFs expected to selectively affect TEX$_{term}$ differentiation. The TEX$_{term}$-TFs included one

known TF, *Nfatc1*, and 11 others that had high specificity scores but were not previously linked to

TEX$_{term}$ differentiation (gray boxes Fig. 1f,h). The multi-state TFs included two positive controls

(*Nr4a2*, *Prdm1*) and five multi-state TFs (*Nfil3*, *Hic1*, *Gfi1*, *Ikzf3*, *Stat3*) not previously tested. To ensure

comprehensive screening, four gRNAs per target were expressed in two dual-gRNA retroviral (RV)

vectors (Extended Data Fig. 7a), along with two control vectors expressing scramble gRNAs

(gScramble). This resulted in a library of 40 dual-gRNA vectors, with 72 gRNAs targeting TFs and 4

gScramble controls (Extended Data Table 6).

To initiate the screening of TEX$_{term}$ cells, Cas9$^+$P14 CD8$^+$ T cells were transduced with this RV-gRNA

library and adoptively transferred into mice infected with **LCMV-Clone 13**, a model of **chronic**

infection and CD8$^+$ T cell exhaustion (*note, recipient mice also expressed Cas9 to prevent rejection of*
donor cells). Droplet-based sequencing was performed 18+ days post-transfer to assess sgRNA and
transcriptomes of each spleen-derived donor Cas9$^+$ P14 CD8$^+$ T (Fig. 3a), analyzing 17,257 cells with
unique gRNA expression.

To determine which TFs impaired TEX$_{term}$ cell differentiation, we first employed uniform manifold
approximation and projection (UMAP) to visualize single-cell transcriptomes. Four primary clusters
were identified consisting of TEX$_{prog}$, TEX$_{eff}$, and TEX$_{term}$ cells and those in cell cycle (Fig. 3b;
Extended Data Fig. 7b,c). All clusters expressed *Tox* and *Pdcd1*, key exhaustion markers, and TEX$_{prog}$
cells were identified by *Tcf7*, *Slamf6*, and *Sell* expression. TEX$_{eff}$ cells expressed effector markers,
including *Cx3cr1*, *Klrd1*, *Klrk1*, *Klf2*, and *Zeb2*[2], while TEX$_{term}$ cells exhibited high expression of
inhibitory receptors and well-established exhaustion markers like *Cd101*, *Cd7*, *Cd38*, *Cd39*, *Cxcr6*, and
Nr4a2. The cell cycle cluster was noted for its expression of *Birc5, Mki67, Stmn1*, and *Tuba1b*.

Next, we evaluated the impact of individual TF depletion by analyzing the distribution of gRNA$^+$ cells
across exhaustion states (Fig. 3c,d). CRISPR KO of the majority of the 19 TEX$_{term}$ driving TFs led to a
reduction in TEX$_{term}$ cell frequency. Notably, KOs of multi-state TFs such as *Hic1*, *Stat3*, *Prdm1*, and
Ikzf3 (which encodes AIOLOS) significantly reduced TEX$_{term}$ differentiation by 92% to 89%. Depletion
of novel TEX$_{term}$ single-state TFs—including *Zfp324*, *Zscan20*, and *Jdp2*—significantly prevented
TEX$_{term}$ differentiation by 78%, 54%, and 43%, respectively (Fig. 3d, bold). Other novel candidates
such as *Etv5*, *Arid3a*, *Zfp410*, *Foxd2*, and *Prdm4* also reduced TEX$_{term}$ representation by 25–40%,
though some did not reach statistical significance. This Perturb-seq analysis highlights the platform's
ability to identify TFs that regulate the TEX$_{term}$ state, with most tested TFs influencing exhaustion to
varying degrees.

To understand how TEX$_{term}$ driving TF KOs affect CD8$^+$ T cell exhaustion, we used flow cytometry and
scRNA-seq to analyze TF KO cells during LCMV-Clone 13 infection (Fig. 3e-i; Extended Data Fig. 8).
We tested six TF KOs, including known control (*Prdm1*) and five newly identified TFs (*Zscan20*, *Jdp2*,
Zfp324, *Stat3*, *Hic1*) that impaired TEX$_{term}$ state differentiation in Perturb-seq. Disrupting these TFs
reduced TEX$_{term}$ cell (PD1$^+$ CX3CR1$^-$ SLAMF6$^-$) frequency by ~50% (Fig. 3e, f) and decreased
expression of inhibitory receptors such as CD101, CD39, and CD38 (Extended Data Fig. 8a,b). All 19

8

TEX$_{term}$-TF KOs exhibited a marked decrease in TEX$_{term}$-signature genes[49], including *Cd7*, *Cxcr6*,

Nr4a2, and *Entpd1*(Fig. 3g).

Lastly, the TEX$_{term}$-driving TF KOs were grouped based on their effects on TEX$_{prog}$ (PD1$^+$ CX3CR1$^-$

SLAMF6$^+$, Fig. 3h) or TEX$_{eff}$ (PD1$^+$ CX3CR1$^+$, Fig. 3i) state differentiation. Loss of *Prdm1* and *Stat3*

markedly increased the frequency of TEX$_{prog}$ cells and upregulated TEX$_{prog}$ signature genes (Fig. 3g, h)

whereas loss of *Hic1*, *Zscan20*, *Zfp324*, or *Jdp2* primarily expanded the TEX$_{eff}$ cell population and

effector signature genes (Fig. 3g, i, j). Deletion of the *Zscan20* and *Jdp2* significantly enhanced effector

cytokine production (e.g., IFNγ and TNFα) and reduced viral loads in recipient mice (Fig. 3k, l).

Depletion of TEX$_{term}$ single-state TFs does not disrupt T$_{RM}$ cell differentiation

A major goal of this work was to identify TFs that selectively repress TEX$_{term}$ cell differentiation without

affecting T$_{RM}$ differentiation, thereby enabling more precise programming of CD8$^+$ T cell states. Since

nearly all the predicted TEX$_{term}$ single-state TFs impaired TEX$_{term}$ differentiation to some degree (Fig.

3), the next step was to evaluate their effects on T$_{RM}$ differentiation to confirm their selective activity.

We used the same Perturb-seq library as before, but this time included only the eight TEX$_{term}$ single-

state TFs and seven multi-state TFs that impaired TEX$_{term}$ state development by more than 25% in

chronic LCMV infection (Fig. 3d). To assess their impact on memory CD8$^+$ T cell development, we

isolated RV-transduced Cas9$^+$ P14 CD8$^+$ T cells from the spleen and small intestine (SI) of mice 18 days

after **acute LCMV-Armstrong** infection. We then analyzed 15,211 cells using scRNA-seq to determine

how these perturbations impacted the formation of intestinal T$_{RM}$ cells, as well as circulating splenic

T$_{CM}$ and T$_{EM}$ cells (Fig. 4a).

The UMAP analysis identified four primary clusters containing cells with features of T$_{CM}$ (*Il7r*, *Tcf7*,

Sell and *S1pr1*), T$_{EM}$ (*Cx3cr1*, *Klrg1,* and *Klf2*), and T$_{RM}$ cells (*Cd69, Cd160, and Itgae* (encoding

CD103)) as well as a small T$_{RM}$ cell population with lower *Itgae*, but higher *Ifng* and *Irf1* expression

designated T$_{RM}$ *Itgae*^{low} (Fig. 4b,c; Extended Data Fig. 9a,b)[27,50]. Examination of the gRNA$^+$ cells

revealed that none of the eight TEX$_{term}$ single-state TF KOs (*Zfp324*, *Irf8*, *Zfp410*, *Nfatc1*, *Zscan20*,

Jdp2, *Arid3a*, and *Etv5)* negatively affected T$_{RM}$ formation significantly (Fig. 4d,e, **bold** gene symbols).

In fact, KO of *Etv5* tended to increase the frequency of T$_{RM}$ cells. To evaluate the specificity of the

TEX$_{term}$ single-state TFs, we also examined the expression of the T$_{RM}$ gene signatures[9,39] in the entire

population of gRNA$^+$ cells for each TF tested (Fig. 4f). With the exception of *Etv5* and *Arid3a*, which increased T$_{RM}$-signature gene expression (Fig. 4f, Extended Data Fig. 9c), perturbation of the TEX$_{term}$ single-state TFs did not substantially alter T$_{RM}$-signature gene expression. The platform also predicted novel multi-state TFs, including *Hic1* and *Gfi1*. Interestingly, disruption of these multi-state TFs significantly reduced T$_{RM}$ cell frequency (Fig. 4e) and T$_{RM}$-signature gene expression (Fig. 4f, Extended Data Fig. 9c), mirroring the effects of *Prdm1*, a known multi-state TF for T$_{RM}$ and TEX$_{term}$ cells.

To further validate the Perturb-seq data, we individually depleted the TEX$_{term}$ single-state TFs *Zscan20* and *Jdp2* and multi-state TF *Prdm1* in Cas9$^+$ P14 CD8$^+$ T cells, adoptively transferred them into LCMV-Armstrong infected animals, and assessed their differentiation into T$_{CM}$, T$_{EM}$, and T$_{RM}$ cells using flow cytometry (Fig. 4g,h). Deletion of *Zscan20* and *Jdp2* did not alter the formation of any memory cell subtypes, while perturbation of *Prdm1* significantly reduced T$_{RM}$ and increased T$_{CM}$ formation, as expected. Altogether, this multi-omics pipeline predicted TEX$_{term}$ single-state TFs that drive TEX$_{term}$ differentiation without affecting with T$_{RM}$ cell formation and multi-state TFs that influence both cell states. These results demonstrate the accuracy and predictive power of our approach for pinpointing single-state and multi-state TFs.

Overexpression of T$_{RM}$-selective TF, *Klf6*, improves T$_{RM}$ differentiation without accelerating T cell exhaustion

To further demonstrate the utility of our cell-state selective TF identification pipeline in discovering novel T$_{RM}$-associated TFs, we evaluated *Klf6*, which was identified through our Taiji analysis as a T$_{RM}$ single-state TF (Fig. 1g). We hypothesized that overexpressing *Klf6* (*Klf6*-OE) could enhance T$_{RM}$ formation during acute viral infection, but would not worsen terminal exhaustion in chronic infection. Our results confirmed this hypothesis. When empty-vector control and *Klf6*-OE P14 CD8$^+$ T cells were co-transferred, *Klf6*-OE cells robustly outcompeted control cells, resulting in a 15-fold enrichment in the SI compared to controls (Fig. 4i). Furthermore, there were ~42 times more CD69$^+$ CD103$^+$double-positive cells in *Klf6*-OE than in control donor cells, indicating *Klf6*-OE dramatically increased T$_{RM}$ development in the SI (Fig. 4j). Importantly, *Klf6*-OE did not increase terminal exhaustion during chronic infection (Extended Data Fig. 10). This work not only identified KLF6 as a new T$_{RM}$-driving TF but also confirmed its selectivity.

10

Disruption of novel TEX$_{term}$ single-state TFs enhances tumor control

This platform predicted cell-state-selective TF activity and identified TEX$_{term}$ single-state TFs as targets for engineering T cells that resist exhaustion yet retain effector and memory functions—offering new strategies to improve immunotherapy efficacy. Given that T$_{RM}$ cells are associated with better clinical outcomes in solid tumors[9–12,51], we hypothesized that KO of exhaustion-selective TFs like *Zscan20* could be more effective than targeting T$_{RM}$ and TEX$_{term}$ multi-state TFs like *Hic1*. Utilizing an adoptive cell therapy model, TF gRNA RV-transduced Cas9$^+$ P14 CD8$^+$ T cells were transferred into mice with established melanoma tumors expressing GP33-41 (Fig. 5a). Unlike depletion of the multi-state TF *Hic1*, depleting the TEX$_{term}$ single-state TF, *Zscan20,* resulted in improved tumor control (Fig. 5b,c). Moreover, *Zscan20* gRNA$^+$ cells more readily formed TEX$_{prog}$ cells rather than TIM3$^+$ CD39$^+$ TEX$_{term}$ cells (Fig. 5d,e). To control for inter-mouse variability in antigen load, we co-transferred *Zscan20* or *Hic1* KO with control P14 CD8$^+$ T cells into the same B16-GP33 tumor-bearing mice (Fig. 5f–j; Extended Data Fig. 11). Both KOs significantly increased the frequency of PD1$^+$ SLAMF6$^+$ TIM3$^-$ and decreased the frequency of TIM3$^+$ exhausted cells and TEX$_{term}$ cell state (PD1$^+$ SLAMF6$^-$ CX3CR1$^-$) compared to controls (Extended Data Fig. 11), consistent with their predicted activity in TEX$_{term}$ cells (Fig. 1f, h). However, *Zscan20* KO robustly enhanced effector marker expression (CX3CR1), granzyme B and cytokine production in TILs, whereas *Hic1* KO did not appear to improve effector function to the same degree (Fig. 5g-j). Thus, despite their similar effects on suppressing TEX$_{term}$ cell differentiation in tumors, differences in their ability to promote functional effector-like states may underlie the differential tumor control observed (Fig. 5b,c). Given that *Hic1* functions as a multi-state TF while *Zscan20* as a single-state TF, these findings support the general rationale for targeting state-specific TFs to enable more selective programming of T cell differentiation.

Cross-species conservation of cell state–selectivity TFs in human tumors

To evaluate the relevance of our murine findings in human T cells—particularly for applications in immunotherapy—we conducted cross-species validation using publicly available single-cell multi-omics and scRNA-seq datasets from human tumor-infiltrating CD8$^+$ T cells (Fig. 6a; Extended Data Fig. 12a; Extended Data Table 1). Leveraging the Taiji TF analysis platform, we mapped murine TEX$_{term}$- and T$_{RM}$-associated TFs onto a curated human pan-cancer CD8$^+$ T cell atlas encompassing six tumor types[52–59](GBM, HNSCC, BCC, HCC, RCC, and ccRCC). Human pan-cancer CD8$^+$ T cells were clustered into heterogeneous cell states, including T$_{RM}$ and TEX$_{term}$ clusters (Fig. 6a; Extended Data Fig.

12b). This Taiji analysis across tumor types revealed strong cross-species conservation: TEX$_{term}$ single-state TFs such as *JDP2*, *ZNF410*, and *FOXD2* consistently exhibited higher activity in TEX$_{term}$ clusters compared to T$_{RM}$-like cells (Fig. 6b). Among the 34 murine TEX$_{term}$-specific TFs, 19 showed conserved activity patterns in human TEX$_{term}$ cells. Similarly, T$_{RM}$-specific TFs—such as *NR4A1*, *KLF6*, and *FOSB*—displayed enriched activity in the human T$_{RM}$ cluster (Extended Data Fig. 12c). Furthermore, 22 of the 30 murine TFs that were active in both TEX$_{term}$ and T$_{RM}$ states showed enriched activity profiles in both cell states in human datasets (Extended Data Fig. 12d). Although a few TFs—such as ZSCAN20—could not be assessed in the Taiji analysis due to missing DNA-binding motifs, we complemented our analysis with comparative RNA expression profiling across 15 tumor types (Fig. 6a,c). Notably, 24 of 34 murine TEX$_{term}$ single-state TFs, including *ZSCAN20* and *JDP2*, showed elevated expression in human TEX cells, further supporting their functional relevance across species.

Given the above correlations between mice and human T cells, we perturbed *ZSCAN20* and *JDP2* to assess the relevance of TEX$_{term}$ single-state TFs in human T cells (Extended Data Fig. 13). Following repeated CD3/CD28 stimulation over 18 days to simulate chronic activation (Fig. 6d), *ZSCAN20*- or *JDP2*-deficient CD8$^+$ T cells exhibited increased expression of CCR7 (naïve/stem cell memory/T$_{CM}$ marker) and decreased levels of inhibitory receptors, including LAG3, PD1, and TIM3 (Fig. 6e, f). These KO cells also produced higher levels of effector cytokines and Granzyme B (Fig. 6g, h), indicating that ZSCAN20 and JDP2 contributed to the regulation of exhaustion-associated features in human CD8$^+$ T cells.

Synergistic tumor control by combining ICB therapy with *Zscan20* and *Jdp2* perturbation highlights clinical relevance

Tumors with high TEX$_{term}$ cell infiltration often exhibit poor responses to immune checkpoint blockade therapy[60]. We hypothesized that targeting TEX$_{term}$ single-state TFs could enhance ICB efficacy. Among the TEX$_{term}$-associated TFs, *Zscan20* and *Jdp2* were prioritized for their conservation and functional relevance in human T cells (Fig. 6b-h). To test synergy with ICB, treatment began one day after adoptive transfer of TF-depleted P14 CD8$^+$ T cells (Fig. 6i). The combination of *Zscan20* or *Jdp2* KO with anti-PD-1 therapy significantly reduced tumor burden (Fig. 6j–m) and improved survival (Extended Data Fig. 14). These findings suggest that selectively disrupting TEX$_{term}$ single-state TF represents a promising strategy to enhance T cell therapy by minimizing dysfunctional states while preserving

385   beneficial T cell phenotypes. Overall, our cross-species multi-omics and functional perturbation
386   approach underscores the translational potential of Taiji-identified TFs for improving adoptive T cell
387   therapies.

389   Discussion

Our study introduces a powerful platform for identifying TFs that are pivotal in guiding the
differentiation of specific CD8$^+$ T cell states during viral infections and tumor progression. This platform
has transformative potential for engineering desirable cell states or preventing detrimental ones, offering
a powerful strategy to enhance T cell–based therapies, including adoptive cell therapy. By constructing
a comprehensive transcriptional and epigenetic atlas from 121 experiments spanning nine distinct CD8$^+$
T cell states, we developed a detailed map of TF activity. This allowed us to predict TF activities within
each state, essentially creating a unique TF fingerprint for each cellular context. Additionally, to
facilitate intuitive and interactive exploration of TF activity and networks analyzed from our multi-
omics atlas, we developed an integrated conversational interface, called TaijiChat–a web-based chatbot
platform enabling natural language querying of both the study's datasets and related literature (Extended
Data Note).

Focusing on two critical cell states TEX$_{term}$ (dysfunctional terminally exhausted) and T$_{RM}$ (functional,
tissue-resident memory) T cells, we examined similarities and differences of TF activity and their
networks in both states and engineered T cells to resist exhaustion while retaining functionality of T$_{RM}$
cells. Using *in vivo* Perturb-seq, we validated TF activity for TEX$_{term}$ and T$_{RM}$ cells in both acute and
chronic infection models. While recent CRISPR screenings in CD8$^+$ T cells have identified important
TFs in cytotoxicity, memory formation[61–64], cell enrichment[65], and exhaustion[66], a systematic and
context-dependent understanding of TF roles across multiple contexts has been lacking. Our study
addresses this gap by generating an accurate catalogue of CD8$^+$ T cell state-defining TFs, enabling cost-
effective validation of predicted TF activity and selectivity via Perturb-seq. Furthermore, our study
offers broader and new insight into context-dependent TF regulation. Previously, differential TF
cooperation in different contexts was reported[30,64,67–69]. We extend this by analyzing global TF
associations across cell states, revealing how TF communities regulate T cell-specific pathways,
including protein catabolism in T cell exhaustion, which aligns with prior research on protein
homeostasis[48,70–72]. These TF networks reveal how various cellular processes are differentially

13

controlled between T_{RM} and TEX_{term} cells, providing rationale for their different functional capabilities within tissues.

One of the key outcomes of this study was the identification of novel transcription factors with state-specific functions, including ZSCAN20 and JDP2 as TEX_{term} single-state TFs, KLF6 as a T_{RM} single-state TF, and newly uncovered roles for multi-state TFs such as HIC1 and GFI1. Perturbing TEX_{term} single-state TFs not only prevented T cell exhaustion but also preserved the ability of these cells to differentiate into effector and memory states. This led to significant improvements in tumor control.

To evaluate the clinical importance of the newly discovered TFs and the catalog of TFs with TEX_{term} and T_{RM} selectivity, we confirmed cross-species conservation of a significant number of TFs using Taiji analysis of a human pan-cancer multi-omics atlas, along with comparative expression analysis across pan-cancer single-cell RNA-seq datasets. Furthermore, we demonstrated enhanced human T cell function following perturbation of the TEX_{term} single-state TFs *ZSCAN20* and *JDP2*. Depletion of these two TFs demonstrates synergistic effects with ICB therapy, resulting in significant tumor regression. These findings highlight a promising strategy for enhancing antitumor immunity through precise cell-state programming.

Our TF analysis platform addresses key challenges in T cell engineering by enabling the prevention of exhaustion while preserving the beneficial properties of T_{RM} cells in solid tumors and chronic infection. This framework offers a rational approach to T cell programming through optimized engineering of 'TF recipes' that control cell-state differentiation with more precision, robustness and durability. For example, future strategies may involve testing combinatorial TF recipes that combine over expression of T_{RM}-promoting TFs (e.g. *KLF6*) with depletion of TEX_{term} single-state TFs. One could compare this to depletion of multi-state TFs to further test the paradigm presented here. Such combinatorial TF recipes may be further refined or expanded using AI-based models to predict modified cell states, such as scLong[73], ultimately advancing the design of durable and effective T cell therapies.

In summary, while our study primarily focuses on $CD8^+$ TEX_{term} and T_{RM} cell differentiation to identify potential strategies for boosting antitumor immunity and combating chronic infections, the pipeline we developed for identifying single-state TFs and 'TF recipes' can be adapted to program other cell types, ultimately improving the effectiveness of cell therapy across a range of applications.

14

References

1. Chung, H. K., McDonald, B. & Kaech, S. M. The architectural design of CD8+ T cell responses in acute and chronic infection: Parallel structures with divergent fates. *J. Exp. Med.* **218**, (2021).
2. Kasmani, M. Y. *et al.* Clonal lineage tracing reveals mechanisms skewing CD8+ T cell fate decisions in chronic infection. *J. Exp. Med.* **220**, (2023).
3. Beltra, J.-C. *et al.* Developmental Relationships of Four Exhausted CD8 T Cell Subsets Reveals Underlying Transcriptional and Epigenetic Landscape Control Mechanisms. *Immunity* **52**, 825–841.e8 (2020).
4. Giles, J. R. *et al.* Shared and distinct biological circuits in effector, memory and exhausted CD8 T cells revealed by temporal single-cell transcriptomics and epigenetics. *Nature Immunology* vol. 23 1600–1613 Preprint at https://doi.org/10.1038/s41590-022-01338-4 (2022).
5. Cui, W., Liu, Y., Weinstein, J. S., Craft, J. & Kaech, S. M. An interleukin-21-interleukin-10-STAT3 pathway is critical for functional maturation of memory CD8+ T cells. *Immunity* **35**, 792–805 (2011).
6. Kaech, S. M. & Cui, W. Transcriptional control of effector and memory CD8+ T cell differentiation. *Nat. Rev. Immunol.* **12**, 749–761 (2012).
7. Flynn, J. K. & Gorry, P. R. Stem memory T cells (TSCM)-their role in cancer and HIV immunotherapies. *Clin Transl Immunology* **3**, e20 (2014).
8. Zhang, Z. *et al.* T Cell Dysfunction and Exhaustion in Cancer. *Front Cell Dev Biol* **8**, 17 (2020).
9. Milner, J. J. *et al.* Runx3 programs CD8 T cell residency in non-lymphoid tissues and tumours. *Nature* **552**, 253–257 (2017).
10. Ganesan, A.-P. *et al.* Tissue-resident memory features are linked to the magnitude of cytotoxic T cell responses in human lung cancer. *Nat. Immunol.* **18**, 940–950 (2017).
11. Djenidi, F. *et al.* CD8 CD103 Tumor–Infiltrating Lymphocytes Are Tumor-Specific Tissue-Resident Memory T Cells and a Prognostic Factor for Survival in Lung Cancer Patients. *The Journal of Immunology* vol. 194 3475–3486 Preprint at https://doi.org/10.4049/jimmunol.1402711 (2015).
12. Corgnac, S., Boutet, M., Kfoury, M., Naltet, C. & Mami-Chouaib, F. The Emerging Role of CD8+ Tissue Resident Memory T (TRM) Cells in Antitumor Immunity: A Unique Functional Contribution of the CD103 Integrin. *Front. Immunol.* **9**, 1904 (2018).
13. Komdeur, F. L. *et al.* CD103+ tumor-infiltrating lymphocytes are tumor-reactive intraepithelial CD8+ T cells associated with prognostic benefit and therapy response in cervical cancer. *Oncoimmunology* **6**, e1338230 (2017).
14. Pauken, K. E. *et al.* Epigenetic stability of exhausted T cells limits durability of reinvigoration by PD-1 blockade. *Science* **354**, 1160–1165 (2016).
15. Sen, D. R. *et al.* The epigenetic landscape of T cell exhaustion. *Science* vol. 354 1165–1169 Preprint at https://doi.org/10.1126/science.aae0491 (2016).
16. Wherry, E. J., Blattman, J. N., Murali-Krishna, K., van der Most, R. & Ahmed, R. Viral persistence alters CD8 T-cell immunodominance and tissue distribution and results in distinct stages of functional impairment. *J. Virol.* **77**, 4911–4927 (2003).
17. Wherry, E. J. *et al.* Molecular signature of CD8+ T cell exhaustion during chronic viral infection. *Immunity* **27**, 670–684 (2007).
18. Im, S. J. *et al.* Defining CD8+ T cells that provide the proliferative burst after PD-1 therapy. *Nature* **537**, 417–421 (2016).
19. Miller, B. C. *et al.* Subsets of exhausted CD8 T cells differentially mediate tumor control and respond to checkpoint blockade. *Nat. Immunol.* **20**, 326–336 (2019).
20. Hudson, W. H. *et al.* Proliferating Transitory T Cells with an Effector-like Transcriptional Signature Emerge from PD-1 Stem-like CD8 T Cells during Chronic Infection. *Immunity* **51**, 1043–1058.e4 (2019).
21. Pu, F. *et al.* TIM-3 expression and its association with overall survival in primary osteosarcoma. *Oncol. Lett.* **18**, 5294–5300 (2019).
22. Zang, K. *et al.* TIM-3 as a Prognostic Marker and a Potential Immunotherapy Target in Human Malignant Tumors: A Meta-Analysis and Bioinformatics Validation. *Front. Oncol.* **11**, 579351 (2021).
23. Liu, F. *et al.* CTLA-4 correlates with immune and clinical characteristics of glioma. *Cancer Cell Int.* **20**, 7 (2020).

24. Zhang, M. *et al.* Prognostic Values of CD38+CD101+PD1+CD8+ T Cells in Pancreatic Cancer. *Immunol. Invest.*
(2019) doi:10.1080/08820139.2019.1566356.
25. Shin, H. *et al.* A role for the transcriptional repressor Blimp-1 in CD8(+) T cell exhaustion during chronic viral
infection. *Immunity* **31**, 309–320 (2009).
26. Behr, F. M. *et al.* Blimp-1 Rather Than Hobit Drives the Formation of Tissue-Resident Memory CD8 T Cells in the
Lungs. *Front. Immunol.* **10**, 400 (2019).
27. Milner, J. J. *et al.* Heterogenous Populations of Tissue-Resident CD8+ T Cells Are Generated in Response to
Infection and Malignancy. *Immunity* **52**, 808–824.e7 (2020).
28. Li, C. *et al.* The Transcription Factor Bhlhe40 Programs Mitochondrial Regulation of Resident CD8 T Cell Fitness
and Functionality. *Immunity* **51**, 491–507.e7 (2019).
29. Wu, J. E. *et al.* In vitro modeling of CD8 T cell exhaustion enables CRISPR screening to reveal a role for BHLHE40.
Sci Immunol **8**, eade3369 (2023).
30. Chen, J. *et al.* NR4A transcription factors limit CAR T cell function in solid tumours. *Nature* **567**, 530–534 (2019).
31. Mackay, L. K. *et al.* The developmental pathway for CD103(+)CD8+ tissue-resident memory T cells of skin. *Nat.*
Immunol. **14**, 1294–1301 (2013).
32. Schacht, T., Oswald, M., Eils, R., Eichmüller, S. B. & König, R. Estimating the activity of transcription factors by the
effect on their target genes. *Bioinformatics* **30**, i401–7 (2014).
33. Zhang, K., Wang, M., Zhao, Y. & Wang, W. Taiji: System-level identification of key transcription factors reveals
transcriptional waves in mouse embryonic development. *Sci Adv* **5**, eaav3262 (2019).
34. Wang, J., Liu, C., Chen, Y. & Wang, W. Taiji-reprogram: a framework to uncover cell-type specific regulators and
predict cellular reprogramming cocktails. *NAR Genom Bioinform* **3**, lqab100 (2021).
35. Liu, C. *et al.* Systems-level identification of key transcription factors in immune cell specification. *PLoS Comput.*
Biol. **18**, e1010116 (2022).
36. Milner, J. J. *et al.* Delineation of a molecularly distinct terminally differentiated memory CD8 T cell population. *Proc.*
Natl. Acad. Sci. U. S. A. **117**, 25667–25678 (2020).
37. Guan, T. *et al.* ZEB1, ZEB2, and the miR-200 family form a counterregulatory network to regulate CD8 T cell fates. *J*
Exp Med **215**, 1153–1168 (2018).
38. Scott-Browne, J. P. *et al.* Dynamic Changes in Chromatin Accessibility Occur in CD8 T Cells Responding to Viral
Infection. *Immunity* **45**, 1327–1340 (2016).
39. Mackay, L. K. *et al.* Hobit and Blimp1 instruct a universal transcriptional program of tissue residency in lymphocytes.
Science **352**, 459–463 (2016).
40. Renkema, K. R. *et al.* KLRG1 Memory CD8 T Cells Combine Properties of Short-Lived Effectors and Long-Lived
Memory. *J Immunol* **205**, 1059–1069 (2020).
41. Intlekofer, A. M. *et al.* Effector and memory CD8+ T cell fate coupled by T-bet and eomesodermin. *Nat Immunol* **6**,
1236–1244 (2005).
42. Reiser, J., Sadashivaiah, K., Furusawa, A., Banerjee, A. & Singh, N. Eomesodermin driven IL-10 production in
effector CD8 T cells promotes a memory phenotype. *Cell Immunol* **335**, 93–102 (2019).
43. Li, G. *et al.* T-Bet and Eomes Regulate the Balance between the Effector/Central Memory T Cells versus Memory
Stem Like T Cells. *PLoS One* **8**, e67401 (2013).
44. Crowl, J. T. *et al.* Tissue-resident memory CD8 T cells possess unique transcriptional, epigenetic and functional
adaptations to different tissue environments. *Nature Immunology* vol. 23 1121–1131 Preprint at
https://doi.org/10.1038/s41590-022-01229-8 (2022).
45. Chandele, A. *et al.* Formation of IL-7Ralphahigh and IL-7Ralphalow CD8 T cells during infection is regulated by the
opposing functions of GABPalpha and Gfi-1. *J. Immunol.* **180**, 5309–5319 (2008).
46. Smith, N. P. *et al.* Resident memory T cell development is associated with AP-1 transcription factor upregulation
across anatomical niches. *bioRxiv* (2023) doi:10.1101/2023.09.29.560006.
47. Guo, X. *et al.* Global characterization of T cells in non-small-cell lung cancer by single-cell sequencing. *Nat Med* **24**,
978–985 (2018).
48. Riesenberg, B. P. *et al.* Stress-Mediated Attenuation of Translation Undermines T-cell Activity in Cancer. *Cancer*
Res. **82**, 4386–4399 (2022).
49. Daniel, B. *et al.* Divergent clonal differentiation trajectories of T cell exhaustion. *Nat. Immunol.* **23**, 1614–1627

(2022).

50. Lin, Y. H. *et al.* Small intestine and colon tissue-resident memory CD8+ T cells exhibit molecular heterogeneity and
differential dependence on Eomes. *Immunity* **56**, 207–223.e8 (2023).

51. Komdeur, F. L. *et al.* CD103 tumor-infiltrating lymphocytes are tumor-reactive intraepithelial CD8 T cells associated
with prognostic benefit and therapy response in cervical cancer. *OncoImmunology* vol. 6 e1338230 Preprint at
https://doi.org/10.1080/2162402x.2017.1338230 (2017).

52. Long, Z. *et al.* Single-cell multiomics analysis reveals regulatory programs in clear cell renal cell carcinoma. *Cell
Discov* **8**, 68 (2022).

53. Terekhanova, N. V. *et al.* Epigenetic regulation during cancer transitions across 11 tumour types. *Nature* **623**, 432–
441 (2023).

54. Riegel, D. *et al.* Integrated single-cell profiling dissects cell-state-specific enhancer landscapes of human tumor-
infiltrating CD8 T cells. *Mol Cell* **83**, 622–636.e10 (2023).

55. Yost, K. E. *et al.* Clonal replacement of tumor-specific T cells following PD-1 blockade. *Nat Med* **25**, 1251–1259
(2019).

56. Ma, L. *et al.* Tumor Cell Biodiversity Drives Microenvironmental Reprogramming in Liver Cancer. *Cancer Cell* **36**,
418–430.e6 (2019).

57. Young, M. D. *et al.* Single-cell transcriptomes from human kidneys reveal the cellular identity of renal tumors.
Science **361**, 594–599 (2018).

58. Cillo, A. R. *et al.* Immune Landscape of Viral- and Carcinogen-Driven Head and Neck Cancer. *Immunity* **52**, 183–
199.e9 (2020).

59. Zheng, L. *et al.* Pan-cancer single-cell landscape of tumor-infiltrating T cells. *Science* **374**, abe6474 (2021).

60. Miller, B. C. *et al.* Author Correction: Subsets of exhausted CD8+ T cells differentially mediate tumor control and
respond to checkpoint blockade. *Nat. Immunol.* **20**, 1556 (2019).

61. Cortez, J. T. *et al.* CRISPR screen in regulatory T cells reveals modulators of Foxp3. *Nature* **582**, 416–420 (2020).

62. Schmidt, R. *et al.* CRISPR activation and interference screens decode stimulation responses in primary human T cells.
Science **375**, eabj4008 (2022).

63. Shifrut, E. *et al.* Genome-wide CRISPR Screens in Primary Human T Cells Reveal Key Regulators of Immune
Function. *Cell* **175**, 1958–1971.e15 (2018).

64. McCutcheon, S. R. *et al.* Transcriptional and epigenetic regulators of human CD8+ T cell function identified through
orthogonal CRISPR screens. *Nat. Genet.* **55**, 2211–2223 (2023).

65. Chen, Z. *et al.* In vivo CD8+ T cell CRISPR screening reveals control by Fli1 in infection and cancer. *Cell* **184**, 1262–
1280.e22 (2021).

66. Zhou, P. *et al.* Single-cell CRISPR screens in vivo map T cell fate regulomes in cancer. *Nature* **624**, 154–163 (2023).

67. Lynn, R. C. *et al.* c-Jun overexpression in CAR T cells induces exhaustion resistance. *Nature* **576**, 293–300 (2019).

68. Seo, H. *et al.* BATF and IRF4 cooperate to counter exhaustion in tumor-infiltrating CAR T cells. *Nat. Immunol.* **22**,
983–995 (2021).

69. Rubin, A. J. *et al.* Coupled Single-Cell CRISPR Screening and Epigenomic Profiling Reveals Causal Gene Regulatory
Networks. *Cell* **176**, 361–376.e17 (2019).

70. Van Der Byl, W. *et al.* The CD8+ T cell tolerance checkpoint triggers a distinct differentiation state defined by protein
translation defects. *Immunity* (2024) doi:10.1016/j.immuni.2024.04.026.

71. Widjaja, C. E. *et al.* Proteasome activity regulates CD8+ T lymphocyte metabolism and fate specification. *J. Clin.
Invest.* **127**, 3609–3623 (2017).

72. Chang, J. T. *et al.* Asymmetric proteasome segregation as a mechanism for unequal partitioning of the transcription
factor T-bet during T lymphocyte division. *Immunity* **34**, 492–504 (2011).

73. Xie, P. *et al.* scLong: A Billion-Parameter Foundation Model for Capturing Long-Range Gene Context in Single-Cell
Transcriptomics. (2024) doi:10.21203/rs.3.rs-5420259/v1.

Figure Legends

Fig. 1| Transcription and epigenomic atlas of CD8$^+$ T cell differentiation states and cell state-selective TF identification pipeline. a, The diagram outlines several CD8$^+$ T cell state trajectories in acute and chronic infection/tumor where various memory and exhaustion states are acquired. Parallel differentiation of T$_{RM}$ and TEX$_{term}$ cells and their similarity in anatomical location. **b,** Pearson correlation matrix of batch effect corrected RNA-seq. Color and size are both proportional to correlation values, with red being the highest correlation. **c,** Pipeline of our integrative analysis. The matched RNA-seq and ATAC-seq data[3,9,20,27,36–40] were used as input for Taiji algorithm to construct a regulatory network and output PageRank score matrix representing the activity of TF. The downstream analysis included the identification of single-state TFs, as well as the construction of transcriptional waves and TF networks. **d-h,** TFs in rows (z- normalized), samples in columns, and the colors of the cell in the matrix indicate the normalized PageRank scores. Each column represents a different data set. **d,** PageRank scores of 136 single-state TFs. **e,** 173 multi-state TFs. **f-h,** Bubble plots of PageRank scores and expression of TFs that are selectively active in TEX$_{term}$ state **(f),** T$_{RM}$ state **(g),** and multi-state TFs that are active in both TEX$_{term}$ and T$_{RM}$ cells **(h)**. TEX$_{term}$ single-state TFs are in bold. Color represents the normalized PageRank score with red displaying high values, and circle size represents the logarithm of mRNA expression across 5 different data sets. TFs are ordered based on the p-value. TFs validated *in vivo* are highlighted in gray. **i-j,** Transcription factor waves associated with exhaustion **(i)** or T$_{RM}$ state **(j)**. Sample numbers for each cell state and the statistics used to curate the TF catalogue are provided in Extended Data Fig. 2a and Extended Data Fig. 3a, respectively.

Fig. 2| Global analysis of TF networks in TEX$_{term}$ and T$_{RM}$ cell states. a, TF network analysis in T$_{RM}$ and TEX$_{term}$ cells encompasses both TF-TF association (blue, Fig. 2b-g). **b-c,** TF-TF association network plots focus on TEX$_{term}$ single-state TFs like ZSCAN20 **(b)** and multi-state TFs like HIC1 **(c)** showing predicted TF-TF network in T$_{RM}$ (green) or TEX$_{term}$ (brown) cells. **d-f,** TF-TF association clustering generates five TF communities in T$_{RM}$ and TEX$_{term}$ cells. The overall community topology is shaped by shared TFs (gray) in both T$_{RM}$ and TEX$_{term}$ cells **(d)** while the TF-TF associations exclusive in T$_{RM}$ **(e)** and TEX$_{term}$ **(f)** cells are shown as green and brown lines, respectively. **g,** Summary of the shared and unique pathways enriched in each community in T$_{RM}$ and TEX$_{term}$ cells. The thickness of the lines reflects p-value. (Gene sets: Extended Data Table. 8). **h**, GSEA of indicated pathways in TEX$_{term}$

vs. T_{RM} cells using batch-effect corrected LCMV bulk-RNA-seq[3,9,20,27,36–40] and human pan-cancer

scRNA-seq data sets[47,59,74]. **i-k,** Flow cytometry analysis of proteasome activity. TEX_{term} cells show the

highest proteasome activity in LCMV-Clone 13 (chronic strain) infected mice **(i),** or MCA-205 tumors

(j). k, Dual transfer of GP33-41 antigen-specific (P14) CD8$^+$ T cells and bystander (OT-1) CD8$^+$ T cells

in B16-GP33 tumor bearing mice. **l,** Evaluation of B16F1-OVA tumor growth by OVA antigen-specific

OT-1 cells with differential proteasome activity. C57BL/6 mice bearing tumors were infused with

proteasome^{high} or proteasome^{low} sorted OT-1 cells, which were pre-stimulated for 7 days with B16F1-

OVA tumor cells. Tumor growth was monitored every two days. Data are presented as the mean ±s.e.m.

Ordinary one-way ANOVA **(i-k),** and two-way ANOVA Tukey's multiple comparison test **(l)** were

performed. ****P<0.0001, ***P<0.001, **P<0.01, *P<0.05.

Fig. 3| *In vivo* Perturb-seq validation of TEX_{term}-driving TFs. a, Schematic of the *in vivo* Perturb-

seq strategy. Dual Cas9$^+$ and P14$^+$ TCR transgenic CD8$^+$ T cells that recognize LCMV epitope GP33-

41 were transduced with retroviruses (RVs) expressing the gRNA library, adoptively transferred into

mice previously infected one day earlier with LCMV-Clone 13, and analyzed 18-23 days later by

scRNA-seq. **b**, UMAP showing TEX_{prog}, TEX_{eff}, TEX_{term}, and cell cycle clusters. Feature plots and

heatmap of the cluster marker mRNA expression are in Extended Data Fig. 7b,c. **c**, Kernel density of

each TF gRNA-expressing cells. **d**, Cluster distribution of cells for each TF gRNA$^+$ cells. TEX_{term} single-

state TFs are indicated in bold. Data represent five pooled replicates from three independent

experiments. Values are shown as mean ± s.e.m. Statistical analysis was performed using two-way

ANOVA followed by Fisher's LSD test, with comparisons made relative to the gScramble control.

Statistical results for TEX_{term} and TEX_{prog} populations are shown; additional statistical comparisons are

provided in Extended Data Table 7. ****P < 0.0001, ***P < 0.001, **P < 0.01, *P < 0.05. **e**, Individual

phenotyping of the TF KOs in LCMV-Clone 13 infected mice based on flow cytometry and

representative flow plots. **f**, Quantification of the frequency of TEX_{term} (PD1$^+$ SLAMF6$^-$ CX3CR1$^-$)

subsets within P14 CD8$^+$ T cells. **g**, TEX_{term}, TEX_{prog}, and TEX_{eff} gene signature analysis in each gRNA

KO compared to gScramble. Gene signatures were generated from ref. [49] (Extended Data Table 8). **h**,

Quantification of the frequency of TEX_{prog} (PD1$^+$ SLAMF6$^+$ CX3CR1$^-$) cells. **i**, Quantification of the

frequency of TEX_{eff} (PD1$^+$ CX3CR1$^+$) cells. **j**, GSEA of an effector-associated gene set between each

TF KOs *vs*. control. **k-l**, P14 CD8$^+$ T cells lacking the indicated TFs were compared to gScramble

controls for IFNγ and TNFα production **(k)** or virus titers **(l)** in LCMV-Clone 13 infected mice. Statistic

19

analysis for **f, h, i, k, l**: Ordinary one-way ANOVA with Dunnett's multiple comparison test versus
gScramble (n ≥ 6 from ≥ 3 biological replicates). Data are presented as the mean ± s.e.m. ****P<0.0001,
***P<0.001, **P<0.01, *P<0.05.

Fig. 4| Functional validation of TFs with distinct roles in TEX$_{term}$ and T$_{RM}$ differentiation. a,
Schematic of *in vivo* Perturb-seq screening in memory CD8$^+$ T cells during acute LCMV-Armstrong
infection. The donor cells were then analyzed for T$_{RM}$ cells in the small intestine (SI) and T$_{CM}$ and T$_{EM}$
cells in the spleen. **b,** UMAP showing T$_{CM}$, T$_{EM}$, T$_{RM}$ and T$_{RM}$- *Itgae^{low}* clusters. **c,** Differential clustering
of cells from different tissue origins. **d,** Kernel density plot of gRNA$^+$ cells on UMAP. **e,** Cluster
distribution of cells with each gRNA. TEX$_{term}$ single-state TFs are in bold letters. Data represent three
replicates. Values are shown as mean ± s.e.m. Statistical analysis was performed using two-way
ANOVA followed by Fisher's LSD test, with comparisons made relative to the gScramble control.
Statistical results for T$_{RM}$ populations are shown; additional statistical comparisons are provided in
Extended Data Table 5. ****P < 0.0001, ***P < 0.001, **P < 0.01, *P < 0.05. **f,** Normalized expression
score of T$_{RM}$ "up and down" gene-signatures[9,39] in cells with each gRNA KO compared to gScramble
cells. **g-h,** Individual TF KO phenotyping. **g,** The ratio of gRNA KO cells in SI:spleen. **h,** The frequency
of splenic T$_{CM}$ (CD62L$^+$ KLRG1$^-$) and T$_{EM}$ (CD62L$^-$ KLRG1$^+$) cells. **i-j,** P14 CD8$^+$ T cells were
transduced with RVs expressing the T$_{RM}$ single-state TF *Klf6* or empty-control vectors and co-
transferred at ~1:1 ratios into mice infected with acute LCMV-Armstrong one day earlier. **i,**
Representative plots of donor P14 cells pre- and post-transfer, with the latter isolated from the SI. **j,**
Quantification of donor P14 cells expressing T$_{RM}$ markers (CD69$^+$ CD103$^+$) in the SI. Data are presented
as the mean ± s.e.m. ***P<0.001, ** P <0.01, * P <0.05. Ordinary one-way ANOVA with Dunnett's
multiple comparison test versus gScramble (n ≥ 5 from ≥ 2 biological replicates, **g-h**). Co-transfer
experiments (**i–j**) were analyzed using paired t-tests (n ≥ 6 from ≥ 2 biological replicates).

Fig. 5 | Distinct roles of TEX$_{term}$ single-state TF *Zscan20* and multi-state TF *Hic1* in tumor control.
a–c, Experimental design (**a**) for adoptive transfer of P14 CD8$^+$ T cells with CRISPR-mediated KO of
either the TEX$_{term}$ single-state TF (*Zscan20*) or a multi-state TF (*Hic1*, active in both TEX$_{term}$ and T$_{RM}$)
into B16-GP33 melanoma-bearing mice. Tumor volumes were measured twice per week (**b**) and weights
were assessed at day 23 (**c**). **d,** Representative flow cytometry plots of PD1$^+$ P14 CD8$^+$ T cells stained

for SLAMF6 and TIM3, with quantification of TEX_{prog} (PD1$^+$ SLAMF6$^+$ TIM3$^-$) and exhausted (PD1$^+$

SLAMF6$^-$ TIM3$^+$) subsets. **e**, Frequency of PD1$^+$ CD39 $^+$ double-positive cells. **f-j**, Co-transfer

experiments in which Cas9$^+$ P14 CD8$^+$ T cells expressing gRNA targeting *Zscan20* or *Hic1* were mixed

with control (scramble gRNA) and transferred into B16-GP33 tumor-bearing mice. Experimental design

(**f**). Representative flow cytometry plots of effector marker CX3CR1 and the cytotoxic cytokine

Granzyme B (GzmB) in TF KO and control cells (**g**). Quantification of CX3CR1$^+$ (**h**), GzmB$^+$ (**i**), and

IFNγ$^+$ TNFα$^+$ (**j**) population in *Zscan20* KO vs. control cells (left), and *Hic1* KO vs. control cells (right).

Statistical analysis was performed using two-way ANOVA with Tukey's multiple comparisons test (**b**)

and one-way ANOVA with Dunnett's test versus gScramble (**c**). Data are from n ≥ 9 from ≥2 biological

replicates (**b-e**). Co-transfer experiments (**g–j**) were analyzed using paired t-tests (n ≥ 6 from ≥2

biological replicates). Data are presented as mean ± s.e.m. ****P < 0.0001, ***P < 0.001, **P < 0.01,

*P < 0.05.

Fig. 6 | Targeting conserved TEX_{term} single-state TFs enhances human T cell effector function and

immunotherapy responses

a, Human pan-cancer single-cell multi-omics and scRNA-seq datasets[52–59] were curated to evaluate TF

mRNA expression across diverse T cell states and to infer TF activity using the scTaiji algorithm. **b,** For

each sample, paired scRNA-seq and scATAC-seq data were used to construct a regulatory network and

compute PageRank-based TF activity scores with scTaiji. Shown are normalized PageRank scores of

TEX_{term} single-state TFs identified in this study (Fig. 1f) with known binding motifs conserved in the

human genome. **c,** mRNA expression of TEX_{term} single-state TFs across TEX_{term} and T_{RM} cell states in

human pan-cancer datasets. TFs with conserved cell-state-selectivity in human and mice are highlighted

in bold (**b,c**). **d,** Human PBMCs with *ZSCAN20* or *JDP2* KO were stimulated with anti-CD3/CD28

antibodies to induce chronic activation and mimic T cell exhaustion. Cells were expanded every 2–3

days at a density of 1 × 10^6 cells/mL. **e,** KO and control human CD8$^+$ T cells were analyzed for

CCR7(central memory/stem-like marker) and LAG3 (exhaustion marker) by flow cytometry as shown

in representative flow plots and cumulative bar graphs. **f,** TF KO and control human CD8$^+$ T cells were

analyzed for inhibitory receptors PD1$^+$ TIM3$^+$ double positive cells by flow cytometry as shown in

representative flow plots and cumulative bar graphs. **g,** Frequency of IFNγ$^+$ TNFα$^+$ double-positive cells

and and IL-2 producing cells. **i,** Polyfunctionality analysis based on the number of cytokines produced

per cell. **i,** Experimental design for testing the synergy between TEX_{term} single-state TF KO and anti–

PD-1 treatment. Cas9$^+$ P14 CD8$^+$ T cells (with or without TF KO) were adoptively transferred into B16-

GP33 tumor-bearing mice on day 7 post-implantation, followed by anti–PD-1 or isotype IgG2a

treatment. **j–k,** Tumor growth curves (**j**) and tumor weights (**k**) of *Zscan20* KO vs. control. **l–m,** Tumor

growth curves (**l**) and tumor weights (**m**) of *Jdp2* KO vs. control. Data are presented as mean ± s.e.m. n

≥ 6 from ≥2 biological replicates. One-way ANOVA with Dunnett's test versus gScramble (**e-g**); two-

way ANOVA with Dunnett's multiple comparisons test versus gScramble (**h**); two-way ANOVA with

Tukey's multiple comparisons test (**j,l**); one-way ANOVA with Tukey's test (**k, m**) ****$P < 0.0001$,

***$P < 0.001$, **$P < 0.01$, *$P < 0.05$.

Methods

Dataset acquisition for murine CD8$^+$ T cell state multiomics atlas: CD8$^+$ T cell samples were collected from ten datasets, including those generated in this study (Extended Data Fig. 2a). In total, we analyzed 121 experiments, comprising 52 ATAC-seq and 69 RNA-seq datasets, which were integrated to generate paired samples and served as input for the Taiji pipeline. The samples encompassed nine distinct CD8$^+$ T cell subtypes: naive cells (Naive), terminal effector cells (TE), memory precursor cells (MP), tissue-resident memory cells (T$_{RM}$), effector memory cells (T$_{EM}$), central memory cells (T$_{CM}$), progenitor exhausted cells (TEX$_{prog}$), intermediate exhausted cells (TEX$_{eff}$) and terminal exhausted cells (TEX$_{term}$). Cell states were defined based on established surface marker combinations and LCMV (lymphocytic choriomeningitis virus)-specific tetramers, including IL7R, KLRG1, PD1, SLAMF6, CD101, Tim3, CD69, CD103, H2-Db LCMV gp 33-41 and H2-Db LCMV gp 276-286 or congenic markers for P14 (T cell receptor (TCR) specific for the LCMV GP33–41 peptide CD8$^+$ T cells), in the context of either acute (LCMV-Armstrong) or chronic (LCMV-Clone 13) infection models. A complete summary of dataset sources, accession numbers, infection conditions, and corresponding cell state definitions (sorting gates) is provided in Extended Data Table 1 and Extended Data Fig. 2a.

TF regulatory networks construction and visualization: To perform integrative analysis of RNA-seq and ATAC-seq data, we developed Taiji v2.0. which allows visualization of several downstream analysis-TF wave, TF-TF association, TF community analysis. Epitensor was used for the prediction of chromatin interactions. Putative TF binding motifs were curated from the latest CIS-BP database[75]. In this analysis, 695 TFs were identified as having binding sites centered around ATAC-seq peak summits. The average number of nodes (genes) and edges (interactions) of the genetic regulatory networks across CD8$^+$ T cell states were 15,845 and 1,325,694, respectively, including 695 (4.38%) TF nodes. On average, each TF regulated 1,907 genes, and each gene was regulated by 22 TFs.

Identification of single-state/multi-state TFs: We first identified universal TFs with mean PageRank across 9 cell states ranked as top 10% and Coefficients of variation (CV) less than 0.5. In total, 54 universal TFs were identified and shown in Extended Data Table 1. The remaining 641 TFs were candidate TFs for single-state TFs. To identify single-state TFs, we divided the samples into two groups: target group and background group. Target group included all the samples belonging to the cell state of interest and the background group comprised the remaining samples. We then performed the normality

23

test using Shapiro-Wilk's method to determine whether the two groups were normally distributed, and

we found that the PageRank scores of most samples (90%) follow a log-normal distribution. Based on

the log-normality assumption, an unpaired t-test was used to calculate the P-value. A P-value cutoff of

0.05 and log2 fold change cutoff of 0.5 were used for calling lineage-specific TFs In total, 255 specific

TFs were identified (Extended Data Table 2). Depending on whether the TF appeared in multiple cell

states, they could be further divided into multi-state TFs (Extended Data Table 2, Fig 1e) and 136 single-

state exclusive TFs (Extended Data Table 2, Fig. 1d). Out of 255 single-state TFs, 84 TFs appear in

TEX_{term} or T_{RM}. To identify the truly distinctive TFs between TEX_{term} and T_{RM}, we performed a second

round of unpaired t-test only between TEX_{term} and T_{RM} (Extended Data Table 3). The same cut-offs, i.e.

P-value of 0.05 and log2 fold change of 0.5, were applied to select TEX_{term} single-taskers and T_{RM} single-

taskers. Thirty out of 84 TFs that did not pass the cut-off were identified as TEX_{term} and T_{RM} multi-

taskers. The full workflow was summarized in Extended Data Fig. 3a.

Identification of transcriptional waves: Combined with prior knowledge on the T cell differentiation

path, TF waves are combinations of TFs which are particularly active in certain differentiation stages,

revealing possible mechanisms of how TF activities are coordinated during differentiation. To be more

specific, we clustered the TFs based on the normalized PageRank scores across samples. First, we

performed principal component analysis (PCA) for dimensionality reduction of the TF score matrix. We

retained the first 10 principal components for further clustering analysis, which explained more than

70% variance (Extended Data Fig. 4b; left panel). We used the k-means algorithm for clustering

analysis. To find the optimal number of clusters and similarity metric, we performed the Silhouette

analysis to evaluate the clustering quality using five distance metrics: Euclidean distance, Manhattan

distance, Kendall correlation, Pearson correlation, and Spearman correlation (Extended Data Fig. 4b;

right panel). Pearson correlation was the most appropriate distance metric since the average Silhouette

width was the highest among all five distance metrics. Based on these analyses, we identified 7 distinct

dynamic patterns of TF activity during immune cell development. We further performed functional

enrichment analysis to identify GO terms for these clusters.

TF-TF collaboration network analysis and visualization: To build the TF-TF association networks,

we first defined a set of relevant TFs for each context (TEX_{term} and T_{RM}) by combining cell state-

important and single-state TFs, resulting in 159 TFs for TEX_{term} and 170 for T_{RM}. The analysis was

based on a TF-regulatee network derived from Taiji, where we first consolidated sample networks by

averaging the edge weights for each TF-regulatee pair. To reduce noise, regulatees with low variation

across all TFs (standard deviation ≤ 1) were removed. Subsequently, a TF-TF correlation matrix was

generated by calculating the Spearman's correlation of edge weights for each TF pair across their

common regulatees. From this matrix, we constructed a graphical model using the R package "huge"[76],

which employs the Graphical Lasso algorithm and a shrunken ECDF (empirical cumulative distribution

function) estimator. An edge between two TFs was established if their correlation was deemed

significant by the model, controlled by a lasso penalty parameter (lambda) of 0.052. This value was

chosen as it represents a local minimum on the sparsity-lambda curve, resulting in approximately 15%

of TF-TF pairs being connected. To validate this method, we estimated the false discovery rate by

generating a null model through random shuffling of the TF-regulatee edge weights. Applying our

algorithm to this null data identified zero interactions, confirming that our approach has a very low false

discovery rate.

TF community construction and visualization: Following the construction of the TF-TF association

networks, we identified functionally related TF communities within each network. We applied the

Leiden algorithm[77], using modularity as the objective function and setting the resolution parameter to

0.9, as this value achieved the highest clustering modularity in our analysis. This procedure identified

five distinct communities for each context (TEX$_{term}$ and T$_{RM}$). The final networks, with their detected

communities, were visualized using the Fruchterman-Reingold layout algorithm[78] to spatially represent

the TF-TF association structure.

Pathway enrichment analysis: The enriched functional terms in this study were analyzed by the R

package clusterProfiler 4.0.5. We used GO, KEGG, MSigDb databases for annotation. For gene set

enrichment analysis (GSEA), the genes were first ranked by the mean edge weight in corresponding

samples and H, C5, C6, and C7 collections from MSigDb were used for annotation. A cutoff of P-value

＜0.05 was used to select the significantly enriched GO terms and KEGG pathways.

Heuristic score calculation via the integration of TF regulatory network and Perturb-seq:

We reasoned that (1) log2FC change in expression due to TF KO, and (2) TF-gene regulatory edge

weights could be combined to provide heuristic scores for a TF's regulatory effect on a target gene. For

each guide RNA KO, Seurat's FindMarkers() function was used to quantify log2FC of gene's expression

with respect to the gScramble condition. Heuristic scores were calculated for each TF-gene pair by

multiplying the gene log2FC with the corresponding edge weight from the Taiji analysis. Regulatees of

a TF were annotated as high-confidence if the magnitude log2FC of the regulate exceeded 0.58

(corresponding to a fold-change of 1.5 or its reciprocal) and if the edge weight belonged to the upper

quantile of all edge weights attributed to the TF. The sign of the log2FC was used to determine if the

TF activated or repressed each target gene.

Human TF activity and cell-state selectivity analysis:

Taiji-based analysis of human multiomic datasets. Transcription factor (TF) activity was inferred using

the Taiji pipeline applied to matched single-cell RNA-seq (scRNA-seq) and ATAC-seq (scATAC-seq)

datasets from various human cancers, including ccRCC (n=2; PRJNA768891, GSE240822), GBM

(GSE240822), BCC (GSE123814, EGAS00001006141), HNSCC (GSE139324, EGAS00001006141),

HCC (GSE125449, EGAS00001006141), and RCC (PMID: 30093597; EGAS00001006141). Cell types

were annotated using canonical marker gene expression and categorized into six major CD8+ T cell

states: T_Ex (exhausted), T_ExProg (progenitor exhausted), T_Eff (effector), T_RM (tissue-resident

memory), T_CM/Naive (central memory/naive), and proliferating. PageRank scores derived from Taiji

were log-transformed, averaged within each T cell category, and then standardized using z-score

normalization. Results were visualized with a focus on TF activity in T_RM and T_Ex populations.

TF expression comparison across human CD8+ T cell states. To assess TF expression across diverse T

cell states, raw count matrices from a published pan-cancer CD8+ T cell atlas (GSE156728) were

reprocessed using Seurat's standard workflow. The dataset encompassed T cells from 11 tumor types,

including BC, BCL, CHOL, ESCA, FTC, MM, OV, PACA, RC, THCA, and UCEC. Cell type

annotations provided by the original study were retained and mapped to the following broad categories:

T_RM, T_Ex, T_EM (effector memory), T_m (memory), T_n (naive), and T_k (cycling). Seurat's

AverageExpression function was used to compute average logCPM expression for each TF in each T

cell category, followed by z-score normalization. Data visualization emphasized comparisons between

T_RM and T_Ex subsets.

Mice and infections: C57BL/6/J, OT-1 (C57BL/6-Tg(TcraTcrb)1100Mjb/J), B6.Cg-*Rag2*$^{tm1.1Cgn}$/J, CD45.1 (B6.SJL-*PtprcaPepcb*/BoyJ) mice were purchased from Jackson Laboratories. P14 mice (Pircher et al., 1987) mice have been previously described. Cas9 P14 mice were generated by crossing P14 mice with B6(C)-Gt(ROSA)26Sorem1.1(CAG-cas9*,-EGFP)Rsky/J (Jackson Laboratories). Animals were housed in specific-pathogen-free facilities at the Salk Institute and University and at the University of North Carolina at Chapel Hill. All animal experiments were approved by the Institutional Animal Care and Use Committee (IACUC). Mice were infected with $2x10^5$ PFU LCMV-Armstrong by intraperitoneal injection or $2x10^6$ PFU LCMV-Clone13 by retro-orbital injection under anesthesia.

Viral titers: LCMV Fluorescence Focus Unit titration was performed seeding Vero cells at a density of 30k cells/100μl in a 96 well flat bottom plate in DMEM + 10% FBS + 2% HEPES + 1% Pen/Strep. On the next day, tissues were homogenized on ice, spun down at 1000g for 5 min at 4°C and supernatants or serum were diluted in 10-fold steps. Diluted samples were added to Vero cells and incubated at 37°C, 5% CO_2 for ~20h. Subsequently, inocula were aspirated and wells were incubated with 4% PFA for 30 min at RT before washing with PBS. VL-4 antibody (BioXCell) was conjugated using the Invitrogen AF488 conjugation kit and added to the wells in dilution buffer containing 3% BSA and 0.3% Triton (Thermo Fisher Scientific) in PBS. Cells were incubated at 4°C overnight before washing with PBS and counting foci under the microscope. FFU was calculated per this formula: FFU/ml=Number of plaques/ (dilution× volume of diluted virus added to the plate).

Cell isolation: Spleens were mechanically dissociated with 1mL syringe plungers over a 70um nylon strainer. Spleens were incubated in ammonium chloride potassium (ACK) buffer for 5 minutes. For isolation of small intestinal IEL, Peyer's patches were first removed by dissection. Intestines were longitudinally cut and then cut into 1cm pieces and washed in PBS. Pieces were incubated in 30mL HBSS with 10% FBS, 10mM HEPES, and 1mM dithioerythritol with vigorous shaking at 37C for 30 minutes. Supernatants were collected, washed, and further isolated using a 40/67% discontinuous percoll density centrifugation for 20 minutes at room temp with no brakes.

Cell lines and *in vitro* cultures: B16-GP33 melanoma cell line were cultured in DMEM (Invitrogen) with 10% fetal bovine serum, 1% penicillin-streptomycin and 250 μg/ml G418 (Invitrogen #10131027). MCA-205 (Sigma) tumor line was maintained in RPMI supplemented with 10% FBS, 300 mg/L L-

glutamine, 100 units/mL penicillin, 100 µg/mL streptomycin, 1mM sodium pyruvate, 100µM NEAA,

1mM HEPES, 55µM 2-mercaptoethanol, and 0.2% Plasmocin mycoplasma prophylactic (InvivoGen).

All the tumor cell lines were used for experiments when in the exponential growth phase. For *in vitro* T

cell culture, splenocytes were activated in RPMI 1640 medium (Invitrogen) containing 10% fetal bovine

serum and 1% penicillin-streptomycin, 2mM L-glutamine, 0.1 mg/ml GP33, beta-mercaptoethanol 50

mM and 10 U/ml IL-2.

Tumor engraftment and treatment of tumor-bearing mice (Fig. 5, 6): 3×10^5 B16-GP33 (Fig. 5),

5×10^5 B16-GP33 tumor cells (Fig. 6) were injected subcutaneously in 100 µl PBS. Around $0.5\text{-}1\text{x}10^6$

Cas9$^+$ P14 T cells with CD45.1 markers were transferred to tumor on day 7 without pre-radiation of

tumor-bearing mice. Tumors were measured every 2-3 days post-tumor engraftment for indicated

treatments and calculated. Tumor volume was calculated by volume = (length × width^2)/2. For antibody-

based treatment, tumor-bearing mice were treated with anti-PD1 antibody (200 µg per injection, clone

RMP1-14, BioXcell) twice per week from day 7 post-tumor implantation. Tumor growth was measured

twice per week with calipers. Survival events were recorded each time a mouse reached the endpoint

(tumor volume ≥1,500 mm^3). Tumor weights were measured on day 23 for Fig. 5 and on day 25 for Fig.

6. All experiments were conducted according to the Salk Institute Animal Care and Use Committee.

Tumor digestion and cell isolation for Fig. 5: Tumors were minced into small pieces in RPMI

containing 2% FBS, DNase I (0.5 µg/ml, Sigma-Aldrich), and collagenase (0.5 mg/ml, Sigma-Aldrich)

and kept for digestion for 30 min at 37˚C, followed by filtration with 70 µm cell strainers (VWR).

Filtered cells were incubated with ACK lysis buffer (Invitrogen) to lyse red blood cells, mixed with

excessive RPMI 1640 medium (Invitrogen) containing 10% fetal bovine serum and 1% penicillin-

streptomycin, and centrifuged at 400g for 5 min to obtain a single-cell suspension.

Proteasome activity analysis: For experiments involving the Proteasome Activity Probe (R&D

systems), cells of interest were incubated with the probe at concentration of 2.5mM for 2 hours at 37C

in PBS. Samples were washed and then stained with Zombie NIR viability dye (Biolegend) in PBS at

4C for 15 minutes. Samples were then stained with some variation of the following antibodies for 30

minutes in FACS Buffer on ice: CD45-BV510 (BD Biosciences), CD45.2-BV510 (Biolegend), CD45.1-

PE-Cy7 (Invitrogen), CD4-APC Fire 810 (Biolegend), CD11b-Alexa Fluor 532 (Invitrogen), CD8-

Spark NIR 685 (Biolegend), CD44-Brilliant Violet 785 (Biolegend), CD62L-BV421 (BD Biosciences),

PD1-BB700 (BD Biosciences), TIM-3-BV711 (Biolegend), LAG-3-APC-eFluor 780 (Invitrogen),

SlamF6-APC (Invitrogen), CD39-Superbright 436 (Invitrogen), CX3CR1-PE/Fire 640 (Biolegend),

CD69-PE-Cy5 (Biolegend), GITR-BV650 (BD Biosciences), CD27-BV750 (BD Biosciences). Samples

were collected on a Cytek Northern Lights and analyzed via Cytek SpectraFlo software.

Tumor experiment for proteasome assay (Fig. 2): MCA-205 fibrosarcomas (2.5×10^5) were

established via subcutaneous injection into the right flank of C57BL/6 mice. After 12-14 days of tumor

growth, spleens, draining lymph nodes, and tumors from groups of mice were harvested and tumors

were processed using the Mouse Tumor Dissociation Kit and gentleMACS dissociator (Miltenyi Biotec)

according to manufacturer's protocol. For purification experiments, samples were pre-enriched using

the EasySep Mouse CD8$^+$ T Cell Isolation Kit (Stemcell Technologies) according to manufacturer's

protocol and stained with Live-or-Dye PE Fixable Viability Stain (Biotium) and CD8a-APC (Invitrogen)

and live CD8$^+$ cells were sorted using the FACSAria II cell sorter. CD8$^+$ spleen and pooled TIL samples

were washed in PBS and frozen for RNA-seq analysis. For adoptive cellular therapy experiments, B16-

GP33 melanomas were established subcutaneously by injecting 5.0×10^5 cells into the right flank of

CD45.1 mice and tumor-bearing hosts were irradiated with 5 Gy 24 hours prior to T cell transfer. In

contrast, mice used in the experiments shown in Fig. 5 and Fig. 6 were not irradiated prior to T cell

transfer. After 7 days of tumor growth, 1.5×10^6 CD45.2 OT-1 T cells and 1.5×10^6 CD45.1/CD45.2 P14

T cells were infused in 100 μL PBS via tail vein into tumor bearing mice. Tumors were harvested 14

days after adoptive cell transfer and CD8 TILs were analyzed for proteasome activity. All experiments

were conducted in accordance with the guidelines of the University of North Carolina at Chapel Hill

Animal Care and Use Committee.

Proteasome^{high}/Proteasome^{low} T cell adoptive transfer experiment (Fig. 2 and Extended Data Fig.

6c): For the adoptive transfer experiment involving proteasome^{high} and proteasome^{low} tumor-specific

OT-1 T cells (Fig. 2l), whole splenocytes from OT-1 mice were activated with 1 μg/mL OVA_257–264

peptide and expanded for 7 days in the presence of 200 U/mL rhIL-2 (NCI). On day 7, OT-1 cells were

FACS-sorted based on proteasome activity to isolate proteasome^{high} and proteasome^{low} OT-1

populations. A total of 2.5×10^5 sorted OT-1 cells were injected into C57BL/6 mice bearing B16F1-

OVA melanomas. Tumors were established by subcutaneous injection of 3×10^5 B16F1-OVA cells into

the right flank 7 days prior to T cell transfer. Recipient mice were preconditioned with 5 Gy total body

irradiation 24 hours before adoptive transfer. Tumor growth was measured every other day with calipers.

For Extended Data Fig. 6c, MCA-205 fibrosarcomas (2.5×10^5) were established via subcutaneous

injection into the right flank of C57BL/6 mice. After 14 days of tumor growth, Live CD45$^+$ CD8$^+$ CD44$^+$

PD1$^+$ T cells were sorted from tumors based on proteasome activity (high vs low) using the FACSAria

II cell sorter. 2.5×10^4 cells were then injected into the 2 day MCA-205 bearing RAG2-/- hosts (n=5 per

group) and tumor growth was monitored every other day starting on day 4. All experiments were

conducted in accordance with the guidelines of the University of North Carolina at Chapel Hill Animal

Care and Use Committee.

Retrovirus transduction and adoptive transfer: For overexpression of the gRNA retrovirus vector,

293T cells were transfected with the Eco-helper and MSCV gRNA vectors. 48 h and 72 h later,

supernatant containing retroviral particles was ready for transduction. Donor P14 splenocytes were *in*

vitro activated by 0.1 mg/ml GP33 and 10 U/ml IL-2 at 37°C for 24h, then spin-transduced (1500 g)

with fresh RV supernatant from 293T cells for 90 min at 30°C in the presence of 5 µg/ml polybrene.

CRISPR-Cas9/RNP nucleofection: Naïve CD8+ T cells were enriched from spleen using the EasySep

Mouse CD8+ T cell Isolation Kit (STEMCELL Technologies). sgRNAs targeting *ZSCAN20, JDP2,*

Etv5, Prdm1, Hic1 genes or the mouse or human genome nontargeting scramble (control) were obtained

from Synthego, IDT, and GeneScript (Extended Data Table 5). Cas9 RNP was prepared immediately

before experiments by incubating 1 µl sgRNA (stock: 3 nmol in 10 µl water), 0.6 µl Cas9 nuclease (IDT,

stock: 62 µM) and 3.4 µl RNase-free water at room temperature for 10min. Nucleofection of naïve

CD8+ T cells was performed using Lonza (Allendale, NJ) P3 primary cell kit and program DN100 with

4D-Nucleofector (Lonza Bioscience) for mouse and EO115 for human stimulated T cells . Each

nucleofection reaction consisted of approximately $5\text{-}10 \times 10^6$ cells in 20 µl of nucleofection reagent and

mixed with 5 µl of RNP: Cas9 complex. After electroporation, 100 µl of T cell culture media was added

to the well to transfer the cells to 1.5 mL Eppendorf tube. The cells were rested at 37°C for 3 min. For

in vivo adoptive transfer, cells were resuspended in PBS at desired concentration and adoptively

transferred into recipient mice.

CRISPR gene editing validation via Sanger sequencing: The genomic DNA (gDNA) was isolated from both the KO-induced CD8+ T cells and the control cells using the Quick-DNA MicroPrep Kit (Zymo). The gDNA concentrations were quantified using the NanoDrop One spectrophotometer (ThermoFisher Scientific). Following isolation, PCR amplification was performed with 2x Phusion Plus Green PCR Master Mix (ThermoFisher Scientific) and the respective validation primers under the following conditions: (98 °C for 5 mins; 35 x of 98 °C for 10 secs, 69 °C for 20 secs, 72 °C for 20-30 secs/kb; 72 °C for 2 mins; hold at 10 °C). The PCR products were run on a 2% agarose gel with SYBR Safe DNA Gel Stain (Invitrogen), and the appropriate bands on the gel were extracted and purified with the Gel DNA Recovery Kit (Zymo). Concentrations of the purified amplicon samples were measured and then sent off for sequencing along with the designated sequencing primers designed from Benchling's Primer3 tool. The samples with the KOs were compared to the wildtype controls using EditCo's Ice Analysis software, providing the indel percentages, KO score, and the indel distributions used to assess the editing efficiency. Indel% ranges from 56% to 97%, and the KO score throughout experiments ranges from 32 to 74.

Flow cytometry, cell sorting, and antibodies: Both single cell suspensions were incubated with Fc receptor-blocking anti-CD16/32 (BioLegend) on ice for 10 min before staining. Cell suspensions were first stained with Red Dead Cell Stain Kit (ThermoFisher) for 10 min on ice. Surface proteins were then stained in FACS buffer (PBS containing 2% FBS and 0.1% sodium azide) for 30 min at 4˚C. To detect cytokine production *ex-vivo*, cell suspensions were re-suspended in RPMI 1640 containing 10% FBS, stimulated by 50 ng/ml PMA and 3 μM Ionomycin in the presence 2.5 μg/ml Brefeldin A (BioLegend #420601) for 4 h at 37°C. Cells were processed for surface marker staining as described above. For intracellular cytokine staining, cells were fixed in BD Cytofix/Cytoperm (BD #554714) for 30 min at 4 °C, then washed with 1× Permeabilization buffer (Invitrogen #00-8333-56). For transcription factor staining, cells were fixed in Foxp3 / Transcription Factor Fixation/Permeabilization buffer (Invitrogen #00-5521-00) for 30 min at 4 °C, then washed with 1× Permeabilization buffer. Cells were then stained with intraceulluar antibodies for 30 min at 4 °C. Samples were processed on LSR-II flow cytometer (BD Biosciences) and data were analyzed with FlowJo V10 (TreeStar). Cells were sorted either on FACSAria™ III sorter or Fusion sorter (BD Biosciences). The following antibodies against mouse proteins were used: anti-CD8a (53-6.7), anti-PD1 (29F.1A12), anti-CX3CR1 (SA011F11), anti-SLAMF6 (13G3), anti-CD38 (90), anti-CD39 (24DMS1), anti-CD101 (Moushi101), anti-KRLG1

(2F1), anti-CD69 (H1.2F3), anti-CD103 (M290), anti-CD62L (MEL-14), anti-Tim3 (RMT3-23), anti-Ly5.1 (A20), anti-Ly5.2 (104), anti-IFN-γ (XMG1.2), anti-TNF-α (MP6-XT22). The following antibodies against human proteins were used: anti-CD8a (RPA-T8), anti-CD4 (SK3), anti-CD45RA (H100), anti-CD45RO (UCHL1), anti-CCR7 (G043H7), anti-CD62L (DREG-56), anti-CD69 (FN50), anti-CD103 (Ber-ACT8), anti-CXCR6 (K041E5), anti-PD1 (EH12.2H7), anti-CD38 (HIT2), anti-CD39 (A1), anti-LAG3 (11C3C65), anti-Tim3 (F38-2E2), anti-TIGIT (A15153G), anti-IFN-γ (4S.B3), anti-TNF-α (MAb11), anti-IL 2 (JES6-5H4), anti-GZMB (QA16A02), anti-G4S Linker (E7O2V). These antibodies were purchased from Invitrogen, Biolegend, Cell Signaling, or eBiosciences.

***In vivo* individual TF KO phenotyping:** To assess the functional impact of individual TF KOs in CD8$^+$ T cells, we used Cas9-expressing P14 donor cells (LCMV-specific TCR transgenic mice, CD45.1 congenic) transduced with GFP-expressing retroviral vectors encoding individual gRNAs. Transductions were performed on the day of adoptive transfer without prior sorting. Without sorting, transduced donor cells (0.5-1 x 10^5) were immediately transferred into congenically distinct Cas9-expressing wild-type recipient mice (CD45.2) infected one day prior with either LCMV-Clone 13 or Armstrong strains. At ≥ day 20 post-infection, spleens from Clone 13 model and spleens and small intestines from Armstrong model were harvested. Single-cell suspensions were prepared and analyzed by flow cytometry. Live, single cells were first gated on CD8$^+$ cells, followed by gating on CD45.1$^+$ P14 donor CD8$^+$ T cells. Successfully transduced (gRNA$^+$) cells were identified by GFP expression, which ranged from 10% to 70% of P14 CD8$^+$ T cells across experiments. Due to variability in the number of GFP$^+$ donor P14 CD8$^+$ T cells obtained from different experiments, all phenotypic analyses were performed within the GFP$^+$ (gRNA$^+$) CD45.1$^+$ CD8$^+$ population. PD-1 positive and negative cells, exhaustion subsets (TEX$_{term}$: PD-1$^+$ SLAMF6$^-$ CX3CR1$^-$, TEX$_{prog}$:PD-1$^+$ SLAMF6$^+$ CX3CR1$^-$, TEX$_{eff}$: PD-1$^+$ CX3CR1$^+$) or expression of phenotypic markers was reported as a percentage within the gRNA$^+$ (GFP$^+$) P14 CD8$^+$ T cell population to ensure consistency across samples.

Co-transfers of control and TF KO/overexpression P14 CD8$^+$ T cells in infection or tumor models: Naive CD8$^+$ T cells were isolated from the spleens and lymph nodes of Cas9-expressing LCMV TCR transgenic (Cas9 P14) or P14 mice using the EasySep™ Mouse CD8$^+$ T Cell Isolation Kit (STEMCELL Technologies). Purified P14 cells were activated for ~24 hours on plates coated with goat anti-hamster

IgG (ThermoFisher), followed by 1 μg/mL hamster anti-mouse CD3 and 1 μg/mL hamster anti-mouse

CD28 antibodies (ThermoFisher), in complete T cell medium (RPMI-1640 supplemented with 10% FBS

(HyClone), 55 μM 2-mercaptoethanol, 100 IU/mL penicillin-streptomycin, and 1% HEPES). After

activation, cells were transduced with retroviruses encoding Klf6 overexpression or gRNAs targeting

Hic1 or Zscan20, and cultured with 20 IU/mL IL-2, 2.5 ng/mL IL-7, and 2.5 ng/mL IL-15 (PeproTech).

At 48 hours post-transduction, reporter expression was confirmed by flow cytometry. Donor cell mixes

were prepared using control vs. Klf6-overexpressing cells (Fig. 4), or gScramble vs. gHic1/gZscan20

cells (Fig. 5). For LCMV infection studies, 1.5×10^5 transduced P14 CD8$^+$ T cells were transferred into

recipient mice, followed by infection with either 2×10^5 PFU LCMV-Armstrong (acute infection,

intraperitoneal) or 2×10^6 PFU LCMV-Clone 13 (persistent infection, intravenous). For tumor studies,

5×10^5 to 1×10^6 transduced T cells (gScramble vs. gTF) were transferred on day 7 after B16-GP33

tumor implantation. All experiments were conducted according to the University of North Carolina at

Chapel Hill Animal Care and Use Committee.

1042

1043 **Perturb seq screening using the retroviral transcriptional factor library:**

1044 D*ual-guide direct-capture retroviral sgRNA vector.* To generate dual-guide sgRNA vector (MSCV-

1045 hU6-mU6-SV40-EGFP), we replaced the hU6 RNA scaffold region of the previously described

1046 retroviral sgRNA vector MG-guide[79] with an additional scaffold[80] and murine U6 promoter.

1047

Dual-guide direct-capture retroviral library construction. For the curated gene list containing 21 TFs,

a total of four gRNA sequences distributed on two individual constructs were designed for each gene.

To construct the library, a customized double-strand DNA fragment pool containing 80 oligonucleotides

targeting those 19 TFs and 4 scramble gRNAs (each oligonucleotide contains two guides targeting the

same gene) (Extended Data Table 5) was ordered from IDT. The dual-guide library was generated using

an In-Fusion (Takara) reaction. In brief, the gRNA containing DNA fragment pool was combined in

MG-guide vector linearized with BpiI (Thermo Fisher). The construct was then transformed into Stellar

competent cells (Takara) and amplified, and the resulting intermediate, individual construct was

assessed for quality using Sanger sequencing. Then, individual dual-gRNA vectors were combined. For

quality control, sgRNA skewing was measured using the MAGeCKFlute[81] to monitor how closely

sgRNAs are represented in a library.

In vivo screening. Retrovirus was generated by co-transfecting HEK293 cells with the dual-guide,

direct-capture retroviral TF library and the packaging plasmid pCL-Eco. Supernatants were collected at

48 and 72 hours post-transfection, then stored at $-80\,°C$. Cas9-expressing P14 CD8$^+$ T cells were

transduced with the viral supernatant to achieve a transduction efficiency of 20–30%. To ensure

sufficient representation of control cells in downstream analysis, 50% of the viral mixture consisted of

retrovirus encoding a non-targeting control gRNA vector. For in vivo experiments, 5×10^4 transduced

P14 cells were intravenously transferred into Cas9-expressing, puromycin-resistant C57BL/6 recipient

mice infected one day prior with either LCMV-Clone 13 or LCMV-Armstrong strain. A total of 25

LCMV-Clone 13 infected mice were used for five biological replicates, and 10 LCMV-Armstrong-

infected mice were used for three biological replicates. Each biological replicate was labeled using

hashtag antibodies (BioLegend, TotalSeq-C) to enable sample demultiplexing and statistical analysis.

At ≥18 days post-infection, donor-derived P14 CD8$^+$ T cells were sorted and pooled for Perturb-seq

analysis. Preliminary tests indicated that T cells expressing gRNA *in vivo* exhibit a greater tendency for

gRNA silencing over extended periods compared to *ex vivo* cultured cells, despite initial successful KOs.

To mitigate gRNA barcode silencing, we harvested tissue between days 18 and 23. Sorted EGFP$^+$ P14

CD8$^+$ T cells were resuspended and diluted in 10% FBS RPMI at a concentration of 1×10^6 cells per

ml. Both the gene expression library and the CRISPR screening library were prepared using a Chromium

Next GEM Single Cell 5′ kit with Feature Barcode technology for CRISPR Screening (10x Genomics).

In brief, the single-cell suspensions were loaded onto the Chromium Controller according to their

respective cell counts to generate 10,000 single-cell gel beads in emulsion per sample. Each sample was

loaded into four separate channels. Chromium Next GEM Single Cell 5' Kit v2, PN-1000263, Chromium

5' Feature Barcode Kit, PN-1000541, 5' CRISPR Kit, PN-1000451, Chromium Next GEM Chip K

Single Cell Kit, PN-1000287, Dual Index Kit TT Set A, PN-1000215, Dual Index Kit TN Set A, PN-

1000250 (10x Genomics) in total were used for each reaction. The resulting libraries were quantified

and quality checked using TapeStation (Agilent). Samples were diluted and loaded onto a NovaSeq

(Illumina) using 100 cycle kit to obtain a minimum of 20,000 paired-end reads (26x91 bp) per cell for

the gene expression library and 5,000 paired-end reads per cell for the CRISPR screening library,

yielding an average of 42639, 36739, and 53413 reads aligned from cells from *in vivo* LCMV-Clone 13,

in vivo LCMV-Armstrong infection, and *in vitro* donor respectively.

Data analysis. Alignments and count aggregation of gene expression and sgRNA reads were completed using Cell Ranger (v.7.0.1). Gene expression and sgRNA reads were aligned using the cell Ranger multi count command with default settings. Gene expression reads were aligned to the mouse genome (mm10 from ENSEMBL GRCm38 loaded from 10x Genomics). The median average of 4, 2, and 33 unique molecular identifiers (UMIs) were detected from cells from *in vivo* LCMV-Clone 13 and LCMV-Armstrong infection, and *in vitro* donor respectively. Droplets with sgRNA UMI passing of default Cell Ranger CRISPR analysis Protospacer UMI threshold were used in further analysis. The filtered feature matrices were imported into Seurat (v.4.3.0)[82] to create assays for a Seurat object containing both gene expression and CRISPR guide capture matrices. Cells detected with sgRNAs targeting two or more genes were then removed to avoid interference from multi-sgRNA-transduced cells. Low-quality cells with <200 detected genes, >10% mitochondrial reads, and <5% ribosomal reads were discarded. A total of 17,257 cells (Clone 13) and 15,211 cells (Armstrong) were passed quality filtering and were used for downstream analysis. Count data was normalized by a global-scaling normalization method and linear transformed.[83] Cluster-specific genes were identified using the FindAllMarkers function of Seurat. We utilized Nebulosa[84] to recover signals from sparse features in single-cell data and made gRNA density plots with scCustomize[85] based on kernel density estimation. In each biological replicate (Clone 13 n=5 and Armstrong =3), the percent cluster distribution of cells with each TF gRNA vector was calculated. Among two gRNA vectors per target TF, the gRNA vector with higher TEX_{term} reduction was shown in Fig. 3d and utilized for Perturb-seq in LCMV-Armstrong infection (Extended Data Table 5). Two-way ANOVA with Fisher's LSD test was performed to determine statistical significance. DEG was calculated using the MAST model[86], the results were then used as inputs for GSEA to evaluate the effect on selected pathways. Genes with p-value < 0.05 were considered as DEG.

UMAP plots were generated by calculating UMAP embeddings using Seurat and then plotting them as scatter plots using ggplot2. Kernel density calculations for each gRNA were performed on UMAP embeddings using the MASS package via the kde2d function. The kernel density results were plotted as a raster layer with ggplot2 over the UMAP scatter plots. Finally, density contour lines were added using ggplot2's built-in 2D kernel density contour geom (geom_density_2d).

ATAC-Seq library preparation and sequencing: ATAC-seq was performed as previously described[87]. Briefly, 5,000-50,000 viable cells were washed with cold PBS, collected by centrifugation, then lysed in resuspension buffer (RSB) (10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 3 mM MgCl2)

supplemented with 0.1% NP40, 0.1% Tween-20, and 0.01% digitonin. Samples were incubated on ice

for 3 min, then washed out with 1 ml RSB containing 0.1% Tween-20. Nuclei were pelleted by

centrifugation at 500g for 10 min at 4°C then resuspended in 50 ul transposition mix (25ul 2x TD buffer,

2.5 ul transposase (100 nM final), 16.5 ul PBS, 0.5 ul 1% digitonin, 0.5 ul 10% Tween-20, 5 ul H2O)

and incubated at 37°C for 30 min in a thermomixer with 1000 RPM mixing. DNA was purified using a

Qiagen MinElute PCR cleanup kit, then PCR amplified using indexed oligos. The optimal number of

amplification cycles for each sample was determined by qPCR. Libraries were size selected using

AmpureXP beads and sequenced using an Illumina NextSeq500 for 75bp paired-end reads.

ATAC-Seq analysis: Paired-end 42-bp, or paired-end 75-bp reads were aligned to the M. musculus

mm10 genome using BWA[88,89] with parameters "bwa mem -M -k 32". ATAC-seq peaks were called

using MACS2[90] program using parameters "callpeaks -qvalue 5.0e-2 –shift -100 –extsize 200".

Differentially accessible regions were identified using DESeq2[91] . Batch effect was removed using

limma[92]. Heatmap visualization of ATAC-seq data was performed using pheatmap.

1131

Single-cell RNA sequencing metadata analysis: Analysis was primarily performed in R (v 3.6.1) using

the package Seurat (v 3.1)[82,93], with the package tidyverse (v 1.2.1)[94] used to organize data and the

package ggplot2 (v 3.2.1) used to generate figures. scRNA-seq data from GSE10898, GSE99254,

GSE98638, GSE199565, and GSE181785 were filtered to keep cells with a low percentage of

mitochondrial genes in the transcriptome (< 5%) and between 200 and 3000 unique genes to exclude

poor quality reads and doublets. Cell cycle scores were regressed when scaling gene expression values

and T cell receptor genes were regressed during the clustering process, which was performed with the

Louvain algorithm within Seurat and visualized with UMAP. To quantify the gene expression patterns,

we used Seurat's module score feature to score each cluster based on its per cell expression of TFs.

1141

To obtain Extended Data Fig. 6a, raw single-cell count data and cell annotation data was downloaded

from NCBI GEO (GSE99254)[47]. Count data was normalized and transformed by derivation of the

residuals from a regularized negative binomial regression model for each gene (SCT normalization

method in Seurat[87], v4.1.1), with 5000 variable features retained for downstream dimensionality

reduction techniques. Integration of data was performed on the patient level with Canonical Correlation

Analysis as the dimension reduction technique[95]. Principal component analysis and UMAP dimension

reduction were performed, with the first 50 PCs utilized in UMAP generation. Cells were clustered

utilizing the Louvain algorithm with multi-level refinement. The data was subset to CD8+ T cells, which

were identified utilizing the labels provided by Guo et al[66]. Cell type labels were confirmed via 1)

SingleR[96] (v1.8.1) annotation utilizing the ImmGen[97] database obtained via celda(1.10), 2) cluster

marker identification, and 3) cell type annotation with the ProjecTILs T cell atlas[7] (v2.2.1). After sub-

setting to CD8$^+$ T cells, cells were again normalized utilizing SCT normalization, with 3000 variable

features retained for dimension reduction. Due to the low number of cells on the per-patient level,

HArmstrongony[98](v1.0) was utilized to integrate the data at the patient level, rather than Seurat. PCA

and UMAP dimensionality reduction were performed as above.

1157

1158 **Statistical analyses:** Statistical tests for flow cytometry data were performed using Graphpad Prism 10.

1159 *p-values* were calculated using either two-tailed unpaired Student's t-tests, one-way ANOVA or two-

1160 way ANOVA as indicated in each figure. Linear regressions were performed using the ordinary least

1161 squares method in R (v 3.6.1). All data were presented as the mean ± s.e.m. The *P* values were

1162 represented as follows: ****$P < 0.0001$, ***$P < 0.001$, **$P < 0.01$, and *$P < 0.05$.

1163

Data availability

ATAC-sequencing data from this paper will be deposited in the GEO database (GSE279498). Taiji v2.0

output of this study (TF activity atlas, TF-TF interaction maps, and TF activity on genome browser

view) will be available at our CD8$^+$ T cell TF atlas portal (https://wangweilab.shinyapps.io/Tcellstates/)

and interactive interface for TF atlas exploration (https://huggingface.co/spaces/taijichat/test-taijichat).

1. Login: Hugging Face account at https://huggingface.co/login using ID: taijichat-reviewers, password:
HelloTaij1Chat#3425
2. Access TaijiChat: After logging in, navigate to the TaijiChat testing page
(https://huggingface.co/spaces/taijichat/test-taijichat)
3. Initial Loading: Upon your first visit, TaijiChat will automatically load necessary packages and build the
system from scratch. This initial setup may take a few minutes. You will know TaijiChat is ready when a
green "running" indicator appears in the top-left corner of the webpage.
4. Testing Chat Functionality: Once TaijiChat is running, select the "Chat" tab on the website. You may then
begin entering questions to test the system. Note that the first query might respond slowly due to system
initialization.
5. Accuracy and Limitations: TaijiChat is powered by state-of-the-art Large Language Models (LLMs), which
can still occasionally make mistakes. Always verify critical scientific information against original research
papers and authoritative sources.

All other raw data are available from the corresponding author upon request.

1183
1184   **Code availability**

1185   All scripts and Taiji v2.0 package will be uploaded at GitHub

1186   (https://github.com/Wang-lab-UCSD/Taiji2).

1187

Methods references

74. Zheng, C. *et al.* Landscape of Infiltrating T Cells in Liver Cancer Revealed by Single-Cell Sequencing. *Cell* **169**, 1342–1356.e16 (2017).

75. Weirauch, M. T. *et al.* Determination and Inference of Eukaryotic Transcription Factor Sequence Specificity. *Cell* vol. 158 1431–1443 Preprint at https://doi.org/10.1016/j.cell.2014.08.009 (2014).

76. Zhao, T., Liu, H., Roeder, K., Lafferty, J. & Wasserman, L. The huge Package for High-dimensional Undirected Graph Estimation in R. (2020) doi:10.48550/arXiv.2006.14781.

77. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-connected communities. *Sci. Rep.* **9**, 5233 (2019).

78. Schönfeld, M. & Pfeffer, J. Fruchterman/Reingold (1991): Graph Drawing by Force-Directed Placement. *Schlüsselwerke der Netzwerkforschung* 217–220 Preprint at https://doi.org/10.1007/978-3-658-21742-6_49 (2019).

79. Bailis, W. *et al.* Author Correction: Distinct modes of mitochondrial metabolism uncouple T cell differentiation and function. *Nature* **573**, E2 (2019).

80. Cong, L. *et al.* Multiplex genome engineering using CRISPR/Cas systems. *Science* **339**, 819–823 (2013).

81. Wang, B. *et al.* Integrative analysis of pooled CRISPR genetic screens using MAGeCKFlute. *Nat. Protoc.* **14**, 756–780 (2019).

82. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* vol. 177 1888–1902.e21 Preprint at https://doi.org/10.1016/j.cell.2019.05.031 (2019).

83. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data using regularized negative binomial regression. *Genome Biol.* **20**, 296 (2019).

84. Alquicira-Hernandez, J. & Powell, J. E. Nebulosa recovers single-cell gene expression signals by kernel density estimation. *Bioinformatics* **37**, 2485–2487 (2021).

85. Marsh, S. E. scCustomize: custom visualizations & functions for streamlined analyses of single cell sequencing. *Preprint at https://doi. org/10.5281/zenodo*.

86. Finak, G. *et al.* MAST: a flexible statistical framework for assessing transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome Biol.* **16**, 278 (2015).

87. McDonald, B. *et al.* Canonical BAF complex activity shapes the enhancer landscape that licenses CD8 T cell effector and memory fates. *Immunity* **56**, 1303–1319.e5 (2023).

88. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, (2009).

89. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **26**, (2010).

90. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, 1–9 (2008).

91. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

92. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

93. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat. Biotechnol.* **36**, 411–420 (2018).

94. Wickham, H. *et al.* Welcome to the Tidyverse. *Journal of Open Source Software* vol. 4 1686 Preprint at https://doi.org/10.21105/joss.01686 (2019).

95. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573–3587.e29 (2021).

96. Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nat. Immunol.* **20**, 163–172 (2019).

97. Heng, T. S. P., Painter, M. W. & Immunological Genome Project Consortium. The Immunological Genome Project: networks of gene expression in immune cells. *Nat. Immunol.* **9**, 1091–1094 (2008).

98. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat. Methods* **16**, 1289–1296 (2019).

99. Zhang, L. *et al.* Single-Cell Analyses Inform Mechanisms of Myeloid-Targeted Therapies in Colon Cancer. *Cell* **181**, 442–459.e29 (2020).

Acknowledgments

We thank members of the Chung, Kaech and Wang laboratory for their advice and assistance with experiments. We appreciate UNC Lineberger Preclinical Research Unit at the University of North Carolina at Chapel Hill assisted animal experiments which is supported in part by an NCI Center Core Support Grant (CA16086) to the UNC Lineberger Comprehensive Cancer Center. This work was supported by NIH grants to S.M.K. R37AI066232, R01AI123864, R21AI151986, and R01CA240909, to W.W. R01AI150282, R01HG009626 (in part) to H.K.C. K01EB034321, to J.J.M. R01AI177864 to J.E.T. R01CA248359 and R01CA244361, to D.C.H. AI151123, to Y.W. EB029122 and GM140929, and to L.L. K01EB035649. H.K.C. is a Damon Runyon Fellow supported by Damon Runyon Cancer Research Foundation grant [DRG-2374-19].

Author contributions

H.K.C. designed and performed experiments. A.B., B.M.R., F.X., Y.L., K.Y., D.J., M.S., J.G., S.M., E.C., B.M. D.C., B.C., Q.Y., F.H., T.H.M., S.K.V., V.T., U.H.C., G.D., B.S., Y.W., L.L., P.H., and J.H. assist with experiments. H.K.C., C.L., A.J., N.J.B., M.S. analyzed the sequencing and bioinformatics data. J.W., B.M., M.S., Y.H., L.H., U.H.C., J.M., contribute to bioinformatics data analysis. Z.W., A.W. contributed to TaijiChat and portal generation. J.H. and D.H. provided resources and training for ATAC-sequencing. B.P.R. and J.E.T. performed proteasome activity experiments. H.K.C., S.M.K., W.W., Y.W., L.L., B.S., G.P., J.E.T., J.J.M., D.C.H, provided scientific input and acquired funding. H.K.C, and C.L. wrote the original manuscript. H.K.C., W.W., and S.M.K. reviewed and edited the paper. H.K.C., S.M.K., and W.W. conceptualized, designed, and supervised the study.

Competing interests

S.M.K. is SAB member for Pfizer, EvolveImmune Therapeutics, Arvinas and Affini-T, advisor for Barer Institute of Raphael Holdings and Academic Editor at Journal of Experimental Medicine. The remaining authors declare no conflict of interest.

Additional information

Supplementary Information is available for this paper. Correspondence and requests for materials should be addressed to H. Kay Chung (hkchung@unc.edu), Susan M. Kaech (skaech@salk.edu) and Wei Wang (wei-wang@ucsd.edu).

 Extended Data Figure Legends

Extended Data Fig. 1 | Parallel differentiation of T_{RM} and TEX_{term} and their transactional and

epigenetic similarity. a, UMAP of scRNA-seq data of T cells from blood, tumor, and adjacent normal

tissues of CRC[99], NSCLC[47], and HCC[74] patients. Unbiased clustering identified multiple T cell states

consistent with those observed murine LCMV infection and tumors. **b,** T_{RM} marker genes show higher

expression in TEX_{term} cluster from Pan-cancer scRNA-seq in **a**. **c,** Both T_{RM} and TEX_{term} clusters

upregulate exhaustion-[17] and T_{RM}[39]-associated gene signatures. **d,** Pearson correlation matrix of batch-

corrected ATAC-seq datasets[3,9,37,38]. Color and size are both proportional to correlation strength.

1276

1277 **Extended Data Figure 2 | Composition of multi-omic atlas**. **a,** A total of 121 experiments across

1278 multiple data sets[3,9,20,27,36–40] were utilized to generate an epigenetic and transcriptional atlas of CD8[+] T

1279 cells under chronic and acute antigen exposure. **b,** Batch-effect corrected ATAC-seq signal at the *Ifng*

1280 locus across nine CD8[+] T cell states.

1281

Extended Data Fig. 3 | Cataloging key TFs across CD8[+] T cell states. a, Logic flow of the unbiased

PageRank comparison to classify single-state/multi-state TFs. **b,** Number of TFs catalogued in each cell

state. **c,** Venn Diagrams showing overlap of TFs with the T_{RM} cell state. **d,** TF activity score (normalized

PageRank) of previously reported TEX_{term}-preventing TFs (TFAP4, JUN, BATF3, and BATF), newly

identified TEX_{term} single-state TFs (JDP2, ZFP324, ZBTB49, ZFP143, ZSCAN20), and NFATC1, a

known TEX_{term}-associated TF.

1288

1289 **Extended Data Fig. 4 | TF wave analysis. a,** Schematic of the analysis pipeline. **b,** Selection of

1290 algorithms and parameters for TF wave analysis. The Pearson correlation was chosen for the distance

1291 metric, with k=7 chosen as the optimal cluster number. **c,** Seven TF waves. Circles represent specific

1292 cell states. Red color indicates normalized PageRank scores. **d,** List of TF members in each wave. **e,**

1293 Heatmap of biological pathways enriched in each TF wave. Red-blue color scale indicates the p-value.

1294

1295 **Extended Data Fig. 5 | TF–TF association network in TEX_{term} and T_{RM} cell states. a–c,** TF–TF

1296 associations of (**a**) TEX_{term} single-state TFs (JDP2, ZFP324, and IRF8); (**b**) T_{RM} single-state TFs (FOSB,

1297 SNAI1, and KLF6); and (**c**) multi-state TFs shared by TEX_{term} and T_{RM} (PRDM1, FLI1, and GFI1). Line

color indicates state specificity: T_{RM} (green) or TEX_{term} (brown) state. Line thickness represents interaction intensity.

Extended Data Fig. 6 | TF network analysis reveals proteasome pathway enrichment in TEX_{term} state with diminished tumor control function. a, Pseudotime analysis of $CD8^+$ TIL scRNA-seq data from NSCLC patients (n = 14), showing a positive correlation between proteasome gene scores (KEGG: M10680) and T cell exhaustion. **b,** Gene set enrichment analysis (GO:0043161, Proteasome-mediated ubiquitin-dependent protein catabolic process) of RNA-seq from $CD8^+$ splenocytes (black) and TILs (purple), n = 3 each. **c,** Tumor growth of $Rag2^{-/-}$ mice bearing MCA-205 sarcomas infused with proteasome^{high} or proteasome^{low} $CD8^+$ TILs isolated from C57BL/6 mice bearing MCA-205 tumors. n = 5 per group. Two-way ANOVA Tukey's multiple comparison test were performed. ****$P<0.0001$, ***$P<0.001$, **$P<0.01$, *$P<0.05$.

Extended Data Fig. 7 | *in vivo* **Perturb-seq with dual-guide RNA in LCMV chronic infection. a,** Retroviral vector design for dual-gRNA delivery. **b,** Feature plot of differentiation markers (Pan-exhaustion: red, TEX_{prog}: pink, TEX_{eff}: blue, Cell cycle: purple, TEX_{term}: brown). **c,** Heatmap of marker gene expression across cell state clusters identified by Seurat's Find Markers() function.

Extended Data Fig. 8 | Individual and co-transfer of TF KO P14 $CD8^+$ T cells identifies novel TFs involved in TEX_{term} differentiation during chronic viral infection. a–b, Quantification of $Cas9^+$ P14 $CD8^+$ T cells transduced with TF-targeting gRNAs isolated from LCMV-Clone 13-infected mice. TEX_{term} single-state TFs are in bold. **a,** Frequency of $PD1^+$ $SLAMF6^-$ $CD101^+$ cells. **b,** Frequency of $CD38^+$ $CD39^+$ double-positive cells. Statistical analysis: Ordinary one-way ANOVA with Dunnett's multiple comparisons test versus gScramble (n ≥ 5 from ≥2 biological replicates). Data are presented as mean ± s.e.m. ****$P< 0.0001$; ***$P < 0.001$; **$P < 0.01$; *$P < 0.05$.

Extended Data Fig. 9 | Single-cell transcriptomic profiling of $CD8^+$ T cells from Perturb-seq in acute LCMV infection. a, Feature plots of differentiation marker genes (T_{CM}: yellow, T_{EM}: blue, T_{RM}-*Itgae*^{low}: dark blue, T_{RM}: green). **b,** Heatmap of differentially expressed genes between T_{RM}-*Itgae*^{low} and T_{RM} clusters. **c,** Volcano plots of differentially expressed genes in $Cas9^+$ P14 $CD8^+$ T cells expressing *gEtv5*, *gArid3a*, g*Hic1*, or *gGfi1*.

1329

Extended Data Fig. 10 | T$_{RM}$ single-state TF, *Klf6* overexpression does not accelerate T cell

exhaustion. a, Experimental setup: *Klf6*-RV or control-RV transduced P14 CD8$^+$ T cells co-transferred

into mice infected with chronic LCMV-Clone 13. **b-e,** Quantification of the frequency of (**b**) PD1$^+$, (**c**)

TEX$_{prog}$ (PD1$^+$ SLAMF6$^+$ CX3CR1$^-$), (**d**) TEX$_{eff}$ (PD1$^+$ SLAMF6$^-$ CX3CR1$^+$) and (**e**) TEX$_{term}$ (PD1$^+$

SLAMF6$^-$ CX3CR1$^-$) populations. Paired t-tests (n $\geq$ 6 from $\geq$2 biological replicates). Data are presented

as mean $\pm$ s.e.m. ****P <0.0001, ***P < 0.001, **P < 0.01, *P < 0.05.

1336

Extended Data Fig. 11 | Depletion of TEX$_{term}$-single state TF, *Zscan20* and TEX$_{term}$ and T$_{RM}$ multi-

state TF, *Hic1* reduces T cell exhaustion. a, Co-transfer of Cas9$^+$ P14 CD8$^+$ T cells transduced with

RV-g*Zscan20* or RV-g*Hic1*, mixed with gRNA control RV transduced cells into B16-GP33 tumor-

bearing mice. **b,** Representative flow plots of PD1$^+$ P14 CD8$^+$ T cells stained for SLAMF6 and CX3CR1.

c-e, Quantification of (**c**) TEX$_{prog}$ (PD1$^+$ SLAMF6$^+$ TIM3$^-$), (**d**) TEX$_{term}$ (PD1$^+$ SLAMF6$^-$ CX3CR1$^-$),

and (**e**) non-progenitor, exhausted (PD1$^+$ SLAMF6$^-$ TIM3$^+$) popultation. Paired t-tests (n $\geq$ 6 from $\geq$2

biological replicates). Data are presented as mean $\pm$ s.e.m. ****P < 0.0001, ***P < 0.001, **P < 0.01,

*P < 0.05.

1345

Extended Data Fig. 12 | Conservation of single- and multi-state TFs in human pan-cancer TEX$_{term}$

and T$_{RM}$ cell states. a, Human pan-cancer datasets utilized in this study[52–59]. **b,** Cluster-specific marker

mRNA expression across integrated pan-cancer single-cell multi-omics datasets. **c,** TF activity analysis

using Taiji on matched matched scRNA-seq and scATAC-seq datasets from various human cancers,

including ccRCC, GBM, BCC, HNSCC, HCC, and RCC. CD8$^+$ T cell states were annotated using

canonical marker gene expression. PageRank scores derived from Taiji were log-transformed, averaged

per state, and z-score normalized. Results were visualized with a focus on TF activity T$_{RM}$ and TEX$_{term}$

cell states. **d,** mRNA expression of TFs were compared across human CD8$^+$ T cell states. The dataset

encompassed T cells from 15 tumor types. Cell type annotations provided by the original study were

retained and mapped to the following broad categories of clusters: T$_{RM}$, TEX$_{term}$, T$_{EM}$, Memory, Naive,

and cell cycle. Seurat's AverageExpression function was used, followed by z-score normalization. Data

visualization emphasized comparisons between T$_{RM}$ and TEX$_{term}$ cell states. Cell-state-selectivity

conserved TFs in humans and mice are highlighted in bold (**c, d**).

1359

1360 **Extended Data Fig. 13 | CRISPR KO validation of *ZSCAN20* and *JDP2* in human PBMCs.** Indel

1361 frequencies, representative Sanger sequencing traces, and indel distributions for (**a**) *ZSCAN20* and (**b**)

1362 *JDP2* KOs.

1363

1364 **Extended Data Fig. 14 | Tumor control of TEX$_{term}$ single-state TF deficient T cells. a,** Survival of

1365 B16-GP33 bearing mice receiving *Zscan20* KO or control P14 CD8$^+$ T cells, followed by anti-PD1 or

1366 isotype IgG2a treatment. **b,** Survival of tumor-bearing mice with *Jdp2* KO P14 CD8$^+$ T cell transfer.

1367

1368 **Extended Data Tables**

1377

Extended Data Note

Development of TaijiChat: An Integrated Conversational Interface for Multi-Omics Data Exploration

Overview of the Multi-Agent Architecture and Design Principles of TaijiChat: To facilitate intuitive and interactive exploration of our Taiji multi-omics atlas, we developed TaijiChat, a web-based chatbot platform enabling natural language querying of both the study's datasets and associated literature. Unlike generic large language models (LLMs), TaijiChat provides domain-specific responses grounded in the content of our manuscript, supplementary materials, and underlying transcriptomic and epigenomic datasets. This tight integration with structured outputs—such as TF activity scores, cell clustering results, and motif enrichment profiles—enables users to dynamically engage in context-aware dialogues with the data. Users can directly query TF activity (e.g., "Which TFs are enriched in TRM cells in GBM?"), request visualizations (e.g., UMAP plots or violin plots of selected genes), or retrieve ranked TF lists by specific cell states or tumor contexts.

In addition to data interrogation, TaijiChat incorporates a literature-aware component. Each generated response is cross-referenced with relevant supporting or contrasting studies from databases such as PubMed and GEO, allowing users to validate findings, assess consistency with prior work, and develop novel hypotheses. For example, querying the role of KLF6 in TRM differentiation yields not only a summary of its activity across tumor-infiltrating lymphocytes (TILs) and associated figures from our atlas, but also citations to independent literature corroborating its specific function in TRM cells.

To systematically enable this advanced functionality, we designed TaijiChat as a sophisticated multi-agent artificial intelligence system, integrating natural language processing, safety-verified code generation, and domain-specific expertise. The system employs a manager-orchestrated architecture consisting of specialized agents responsible for planning, validation, and secure code execution. Specifically, this architecture embodies four core design principles critical for scientific computing applications:

1. Safety-first architecture: Ensures all generated analytical code undergoes mandatory validation through multiple security layers prior to execution.

1406   2.   Separation of concerns: Assigns distinct, specialized roles to individual agents, facilitating

1407   modularity, scalability, and maintainability.

1408

1409   3.   Intelligent orchestration: Provides centralized coordination of complex multi-step workflows,

1410   ensuring coherent interactions across system components.

1411

4.   Domain expertise integration: Directly incorporates scientific methodologies and domain-

specific knowledge into the reasoning and analysis processes, enabling contextually appropriate and

scientifically rigorous responses.

To evaluate TaijiChat's effectiveness, we curated a benchmark set of 21 expert-driven immunology

questions, comparing responses from TaijiChat to those from a general-purpose LLM lacking direct data

integration. TaijiChat demonstrated accurate, evidence-supported responses in all tested scenarios,

highlighting its utility as a domain-specific scientific assistant.

Collectively, TaijiChat represents a platform for hypothesis generation and data interpretation,

effectively combining the intuitive accessibility of conversational AI with rigorous, integrated multi-

omics data analysis and automated literature synthesis.

1422

1.1 Agent Architecture and Coordination: The system consists of four specialized agents operating under centralized coordination: the Manager Agent, Generation Agent, Supervisor Agent, and Executor Agent.

The Manager Agent serves as the central orchestrator in a multi-agent architecture designed for scientific computing and code generation tasks. The system employs a hierarchical workflow where the Manager coordinates three specialized sub-agents: a GenerationAgent that transforms natural language queries into executable Python code and structured execution plans, a SupervisorAgent that performs safety validation and code review to prevent potentially harmful operations, and an ExecutorAgent that executes approved code in a controlled environment. The Manager implements a sophisticated query processing pipeline with built-in retry logic (up to three attempts) to handle generation failures, maintains conversation history for context-aware responses, and supports multi-step computational workflows where intermediate results inform subsequent processing steps. This design ensures robust, safe, and efficient processing of complex scientific queries while maintaining clear separation of concerns across the agent hierarchy.RetryClaude can make mistakes.

The Generation Agent functions as the intelligent planner, employing a structured 13-step reasoning process for query understanding and solution development. This agent integrates domain-specific knowledge of transcription factor biology and experimental methodologies, enabling scientifically appropriate interpretations of natural language queries. The reasoning framework encompasses query decomposition, conversation history integration, multi-modal content evaluation, scientific domain knowledge activation, tool capability assessment, and comprehensive execution plan synthesis.

The Supervisor Agent implements a critical safety validation layer through multi-stage code review processes. This agent performs static code analysis, security pattern detection, and tool usage verification before any code execution. The validation process examines code structure, prevents dangerous system operations, validates tool usage patterns, and provides detailed feedback for rejected code with specific remediation suggestions.

The Executor Agent provides secure code execution within a sandboxed Python environment. This agent implements restricted global scope access, limiting available modules to approved scientific computing libraries and custom tools. All code execution occurs in isolation with comprehensive output capture

1454 and sanitized error reporting, preventing system exposure while maintaining informative debugging

1455 capabilities.

1456

1457

1.2 Safety and Security Architecture: The system implements a three-layer security model designed to

prevent unsafe operations while maintaining scientific functionality. Layer 1 provides generation-time

safety through security-aware code templates and built-in constraint knowledge within the Generation

Agent. Layer 2 implements mandatory pre-execution review through the Supervisor Agent's pattern-

based security analysis and tool compliance verification. Layer 3 enforces execution isolation through

the Executor Agent's sandboxed environment with restricted resource access and output sanitization.

Error recovery mechanisms include intelligent retry strategies with supervisor feedback integration,

progressive query simplification for complex requests, and maintenance of conversation context across

retry attempts. The system limits retry attempts to three per query while incorporating learned patterns

from previous failures to improve subsequent attempts.

1468

1469 *1.3 Tools Integration and Performance Optimization:* The system interfaces with a comprehensive tools

1470 module providing specialized functions for scientific data analysis. Data processing capabilities include

1471 Excel file processing with proper formatting, temporal pattern analysis, correlation matrix access, and

1472 intelligent data subsetting. Literature integration tools provide multi-source academic database searches

across PubMed, ArXiv, and Semantic Scholar, with automated content extraction and deduplication algorithms. Multi-modal analysis capabilities support AI-powered image analysis, PDF document processing, and file format detection. Performance optimization employs a multi-level caching strategy including function-level LRU caching, data-level timeout caching, schema caching for persistent structures, and session-based query-response caching. Resource efficiency measures include lazy loading of datasets, batch processing for large operations, automatic memory management, and intelligent API rate limiting for external services

1.4 Technical Implementation and Performance: The system utilizes OpenAI GPT-4o models for both text and vision processing, operating within a Python 3.x runtime environment with scientific computing libraries including pandas and openpyxl. External API integration provides access to major academic databases while maintaining security through input validation and sandboxed execution.

1.5 Architectural Advantages for Scientific Applications: This multi-agent architecture provides several key advantages for scientific computing applications. The multi-layer validation system ensures research integrity through multiple checkpoints, preventing unsafe operations while maintaining detailed audit trails of all agent interactions. The modular design enables easy maintenance and extensibility, allowing individual agent updates without system-wide modifications and seamless integration of new analytical capabilities. Scientific rigor is emphasized through reproducible operations with consistent outputs, transparent reasoning traces for scientific validation, comprehensive academic literature integration, and built-in domain expertise for methodologically appropriate analyses. The system's fail-safe design defaults to safe states during error conditions while providing robust error recovery with intelligent retry mechanisms.

Figure 1

Fig. 1| Transcription and epigenomic atlas of CD8+ T cell differentiation states and cell state-selective TF identification pipeline. a, The diagram outlines several CD8+ T cell state trajectories in acute and chronic infection/tumor where various memory and exhaustion states are acquired. Parallel differentiation of T_RM and TEX_term cells and their similarity in anatomical location. **b,** Pearson correlation matrix of batch effect corrected RNA-seq. Color and size are both proportional to correlation values, with red being the highest correlation. **c,** Pipeline of our integrative analysis. The matched RNA-seq and ATAC-seq data[3,9,20,27,36–40] were used as input for Taiji algorithm to construct a regulatory network and output PageRank score matrix representing the activity of TF. The downstream analysis included the identification of single-state TFs, as well as the construction of transcriptional waves and TF networks. **d-h,** TFs in rows (z- normalized), samples in columns, and the colors of the cell in the matrix indicate the normalized PageRank scores. Each column represents a different data set. **d,** PageRank scores of 136 single-state TFs. **e,** 173 multi-state TFs. **f-h,** Bubble plots of PageRank scores and expression of TFs that are selectively active in TEX_term state **(f),** T_RM state **(g),** and multi-state TFs that are active in both TEX_term and T_RM cells **(h).** TEX_term single-state TFs are in bold. Color represents the normalized PageRank score with red displaying high values, and circle size represents the logarithm of mRNA expression across 5 different data sets. TFs are ordered based on the p-value. TFs validated *in vivo* are highlighted in gray. **i-j,** Transcription factor waves associated with exhaustion **(i)** or T_RM state **(j).** Sample numbers for each cell state and the statistics used to curate the TF catalogue are provided in Extended Data Fig. 2a and Extended Data Fig. 3a, respectively.

Figure 2

Fig. 2| Global analysis of TF networks in TEX_term and T_RM cell states. a, TF network analysis in T_RM and TEX_term cells encompasses both TF-TF association (blue, Fig. 2b-g). **b-c,** TF-TF association network plots focus on TEX_term single-state TFs like ZSCAN20 **(b)** and multi-state TFs like HIC1 **(c)** showing predicted TF-TF network in T_RM (green) or TEX_term (brown) cells. **d-f,** TF-TF association clustering generates five TF communities in T_RM and TEX_term cells. The overall community topology is shaped by shared TFs (gray) in both T_RM and TEX_term cells **(d)** while the TF-TF associations exclusive in T_RM **(e)** and TEX_term **(f)** cells are shown as green and brown lines, respectively. **g,** Summary of the shared and unique pathways enriched in each community in T_RM and TEX_term cells. The thickness of the lines reflects p-value. (Gene sets: Extended Data Table. 8). **h,** GSEA of indicated pathways in TEX_term vs. T_RM cells using batch-effect corrected LCMV bulk-RNA-seq[3,9,20,27,36–40] and human pan-cancer scRNA-seq data sets[47,59,74]. **i-k,** Flow cytometry analysis of proteasome activity. TEX_term cells show the highest proteasome activity in LCMV-Clone 13 (chronic strain) infected mice **(i),** or MCA-205 tumors **(j). k,** Dual transfer of GP33-41 antigen-specific (P14) CD8+ T cells and bystander (OT-1) CD8+ T cells in B16-GP33 tumor bearing mice. **l,** Evaluation of B16F1-OVA tumor growth by OVA antigen-specific OT-1 cells with differential proteasome activity. C57BL/6 mice bearing tumors were infused with proteasome^high or proteasome^low sorted OT-1 cells, which were pre-stimulated for 7 days with B16F1-OVA tumor cells. Tumor growth was monitored every two days. Data are presented as the mean ±s.e.m. Ordinary one-way ANOVA **(i-k),** and two-way ANOVA Tukey's multiple comparison test **(l)** were performed. ****$P<0.0001$, ***$P<0.001$, **$P<0.01$, *$P<0.05$.

Figure 3

Fig. 3| *In vivo* Perturb-seq validation of TEX_{term}-driving TFs. a, Schematic of the *in vivo* Perturb-seq strategy. Dual Cas9+ and P14+ TCR transgenic CD8+ T cells that recognize LCMV epitope GP33-41 were transduced with retroviruses (RVs) expressing the gRNA library, adoptively transferred into mice previously infected one day earlier with LCMV-Clone 13, and analyzed 18-23 days later by scRNA-seq. **b**, UMAP showing TEX_{prog}, TEX_{eff}, TEX_{term}, and cell cycle clusters. Feature plots and heatmap of the cluster marker mRNA expression are in Extended Data Fig. 7b,c. **c**, Kernel density of each TF gRNA-expressing cells. **d**, Cluster distribution of cells for each TF gRNA+ cells. TEX_{term} single-state TFs are indicated in bold. Data represent five pooled replicates from three independent experiments. Values are shown as mean ± s.e.m. Statistical analysis was performed using two-way ANOVA followed by Fisher's LSD test, with comparisons made relative to the gScramble control. Statistical results for TEX_{term} and TEX_{prog} populations are shown; additional statistical comparisons are provided in Extended Data Table 7. ****$P < 0.0001$, ***$P < 0.001$, **$P <$

0.01, *$P < 0.05$. **e**, Individual phenotyping of the TF KOs in LCMV-Clone 13 infected mice based on flow cytometry and representative flow plots. **f**, Quantification of the frequency of TEX_{term} (PD1+ SLAMF6- CX3CR1-) subsets within P14 CD8+ T cells. **g**, TEX_{term}, TEX_{prog}, and TEX_{eff} gene signature analysis in each gRNA KO compared to gScramble. Gene signatures were generated from ref. 49 (Extended Data Table 8). **h**, Quantification of the frequency of TEX_{prog} (PD1+ SLAMF6+ CX3CR1-) cells. **i**, Quantification of the frequency of TEX_{eff} (PD1+ CX3CR1+) cells. **j**, GSEA of an effector-associated gene set between each TF KOs *vs.* control. **k-l**, P14 CD8+ T cells lacking the indicated TFs were compared to gScramble controls for IFNγ and TNFα production **(k)** or virus titers **(l)** in LCMV-Clone 13 infected mice. Statistic analysis for **f, h, i, k, l**: Ordinary one-way ANOVA with Dunnett's multiple comparison test versus gScramble (n ≥ 6 from ≥ 3 biological replicates). Data are presented as the mean ± s.e.m. ****$P<0.0001$, ***$P<0.001$, **$P<0.01$, *$P<0.05$.

Figure 4

Fig. 4| Functional validation of TFs with distinct roles in TEX_term and T_RM differentiation. a, Schematic of *in vivo* Perturb-seq screening in memory CD8+ T cells during acute LCMV-Armstrong infection. The donor cells were then analyzed for T_RM cells in the small intestine (SI) and T_CM and T_EM cells in the spleen. **b,** UMAP showing T_CM, T_EM, T_RM and T_RM- *Itgae^low* clusters. **c,** Differential clustering of cells from different tissue origins. **d,** Kernel density plot of gRNA+ cells on UMAP. **e,** Cluster distribution of cells with each gRNA. TEX_term single-state TFs are in bold letters. Data represent three replicates. Values are shown as mean ± s.e.m. Statistical analysis was performed using two-way ANOVA followed by Fisher's LSD test, with comparisons made relative to the gScramble control. Statistical results for T_RM populations are shown; additional statistical comparisons are provided in Extended Data Table 5. ****P < 0.0001, ***P < 0.001, **P < 0.01, *P < 0.05. **f,** Normalized expression score of T_RM "up and down" gene-signatures[9,39] in cells with each gRNA KO compared to gScramble cells. **g-h,** Individual TF KO phenotyping. **g,** The ratio of gRNA KO cells in SI:spleen. **h,** The frequency of splenic T_CM (CD62L+ KLRG1-) and T_EM (CD62L- KLRG1+) cells. **i-j,** P14 CD8+ T cells were transduced with RVs expressing the T_RM single-state TF *Klf6* or empty-control vectors and co-transferred at ~1:1 ratios into mice infected with acute LCMV-Armstrong one day earlier. **i,** Representative plots of donor P14 cells pre- and post-transfer, with the latter isolated from the SI. **j,** Quantification of donor P14 cells expressing T_RM markers (CD69+ CD103+) in the SI. Data are presented as the mean ± s.e.m. ***P<0.001, ** P <0.01, * P <0.05. Ordinary one-way ANOVA with Dunnett's multiple comparison test versus gScramble (n ≥ 5 from ≥ 2 biological replicates, **g-h**). Co-transfer experiments (**i–j**) were analyzed using paired t-tests (n ≥ 6 from ≥ 2 biological replicates).

Figure 5

Fig. 5 | Distinct roles of TEX_term single-state TF *Zscan20* and multi-state TF *Hic1* in tumor control. a–c, Experimental design (**a**) for adoptive transfer of P14 CD8+ T cells with CRISPR-mediated KO of either the TEX_term single-state TF (*Zscan20*) or a multi-state TF (*Hic1*, active in both TEX_term and T_RM) into B16-GP33 melanoma-bearing mice. Tumor volumes were measured twice per week (**b**) and weights were assessed at day 23 (**c**). **d**, Representative flow cytometry plots of PD1+ P14 CD8+ T cells stained for SLAMF6 and TIM3, with quantification of TEX_prog (PD1+ SLAMF6+ TIM3−) and exhausted (PD1+ SLAMF6− TIM3+) subsets. **e**, Frequency of PD1+ CD39+ double-positive cells. **f-j**, Co-transfer experiments in which Cas9+ P14 CD8+ T cells expressing gRNA targeting *Zscan20* or *Hic1* were mixed with control (scramble gRNA) and transferred into B16-GP33 tumor-bearing mice. Experimental design (**f**). Representative flow cytometry plots of effector marker CX3CR1 and the cytotoxic cytokine Granzyme B (GzmB) in TF KO and control cells (**g**). Quantification of CX3CR1+ (**h**), GzmB+ (**i**), and IFNγ+ TNFα+ (**j**) population in *Zscan20* KO vs. control cells (left), and *Hic1* KO vs. control cells (right). Statistical analysis was performed using two-way ANOVA with Tukey's multiple comparisons test (**b**) and one-way ANOVA with Dunnett's test versus gScramble (**c**). Data are from n ≥ 9 from ≥2 biological replicates (**b-e**). Co-transfer experiments (**g–j**) were analyzed using paired t-tests (n ≥ 6 from ≥2 biological replicates). Data are presented as mean ± s.e.m. ****P < 0.0001, ***P < 0.001, **P < 0.01, *P < 0.05.

Figure 6

Fig. 6 | Targeting conserved TEX_term single-state TFs enhances human T cell effector function and immunotherapy responses

a, Human pan-cancer single-cell multi-omics and scRNA-seq datasets[52–59] were curated to evaluate TF mRNA expression across diverse T cell states and to infer TF activity using the scTaiji algorithm. b, For each sample, paired scRNA-seq and scATAC-seq data were used to construct a regulatory network and compute PageRank-based TF activity scores with scTaiji. Shown are normalized PageRank scores of TEX_term single-state TFs identified in this study (Fig. 1f) with known binding motifs conserved in the human genome. c, mRNA expression of TEX_term single-state TFs across TEX_term and T_RM cell states in human pan-cancer datasets. TFs with conserved cell-state-selectivity in human and mice are highlighted in bold (b,c). d, Human PBMCs with *ZSCAN20* or *JDP2* KO were stimulated with anti-CD3/CD28 antibodies to induce chronic activation and mimic T cell exhaustion. Cells were expanded every 2–3 days at a density of 1×10^6 cells/mL. e, KO and control human CD8+ T cells were analyzed for CCR7(central memory/stem-like marker) and LAG3 (exhaustion marker) by flow cytometry as shown in representative flow plots and cumulative bar graphs. f, TF KO and control human CD8+ T cells were analyzed for inhibitory receptors PD1+ TIM3+ double positive cells by flow cytometry as shown in representative flow plots and cumulative bar graphs. g, Frequency of IFNγ+ TNFα+ double-positive cells and and IL-2 producing cells. i, Polyfunctionality analysis based on the number of cytokines produced per cell. i, Experimental design for testing the synergy between TEX_term single-state TF KO and anti–PD-1 treatment. Cas9+ P14 CD8+ T cells (with or without TF KO) were adoptively transferred into B16-GP33 tumor-bearing mice on day 7 post-implantation, followed by anti–PD-1 or isotype IgG2a treatment. j–k, Tumor growth curves (j) and tumor weights (k) of *Zscan20* KO vs. control. l–m, Tumor growth curves (l) and tumor weights (m) of *Jdp2* KO vs. control. Data are presented as mean ± s.e.m. n ≥ 6 from ≥2 biological replicates. One-way ANOVA with Dunnett's test versus gScramble (e–g); two-way ANOVA with Dunnett's multiple comparisons test versus gScramble (h); two-way ANOVA with Tukey's multiple comparisons test (j,l); one-way ANOVA with Tukey's test (k, m) ****$P < 0.0001$, ***$P < 0.001$, **$P < 0.01$, *$P < 0.05$.