Control of atypical PKCι membrane dissociation by tyrosine phosphorylation within a PB1-C1 interdomain interface

Mathias Cobbaut[1*], Neil Q. McDonald[1,2*], Peter J. Parker[3,4*]

[1] Signalling and Structural Biology Laboratory, The Francis Crick Institute, 1 Midland Road, London, NW1 1AT, UK

[2] Institute of Structural and Molecular Biology, Department of Biological Sciences, Birkbeck College, London, WC1E 7HX, UK.

[3] Protein Phosphorylation Laboratory, The Francis Crick Institute, 1 Midland Road, London, NW1 1AT, UK

[4] School of Cancer and Pharmaceutical Sciences, King's College London, Guy's Campus, London, SE1 1UL, UK.

* for correspondence: Mathias Cobbaut, mathias.cobbaut@crick.ac.uk; Neil Q. McDonald, neil.mcdonald@crick.ac.uk; Peter J. Parker, peter.parker@crick.ac.uk

Abstract

Atypical PKCs are cell polarity kinases that operate at the plasma membrane where they function within multiple molecular complexes to contribute to the establishment and maintenance of polarity. In contrast to the classical and novel PKCs, atypical PKCs do not respond to diacylglycerol cues to bind the membrane compartment. Until recently it was not clear how aPKCs are recruited; whether aPKCs can directly interact with membranes or whether they are dependent on other protein interactors to do so. Two recent studies identified the pseudo-substrate region and the C1 domain as direct membrane interaction modules, however their relative importance and coupling are unknown. We combined molecular modelling and functional assays to show that the regulatory module of aPKCι, comprising the PB1 pseudo-substrate and C1 domains, forms a cooperative and spatially continuous invariant membrane interaction platform. Furthermore, we show the coordinated orientation of membrane-binding elements within the regulatory module requires a key PB1-C1 interfacial β-strand (BSL). We show this element contains a highly conserved Tyr residue that can be phosphorylated and that negatively regulates the integrity of the regulatory module, leading to membrane release. We thus expose a novel regulatory mechanism of aPKCι membrane binding and release during cell polarization.

Introduction

Protein kinase C (PKC) isozymes are subject to tight regulation of their catalytic activities rendering them responsive to an array of cellular signals in specific compartments consistent with other AGC kinase superfamily members (1, 2). For classical and novel PKC isoforms, changes in lipid signalling that induce formation of diacylglycerol (DAG) drive activation and therefore PKC responses to various G-protein coupled receptors and receptor tyrosine kinases linked to PLC activity (1). The atypical PKCs (iota and zeta) on the other hand do not contain DAG-responsive elements in their regulatory amino-termini, but instead contain a PB1 domain tethered via a pseudo-substrate linker region to a DAG-unresponsive C1 domain (3). The pseudo-substrate (PS) motif is thought to repress catalytic activity by competitively blocking access to substrates (4), while the PB1 domain mediates protein-protein interactions with activating/regulatory proteins such as p62 and Par6, which in turn are also modular proteins allowing for multivalent protein interactions (5, 6).

Many aPKC functions are intricately linked with its recruitment to the plasma membrane. Indeed, the polarity complexes that define apical and basolateral membrane identity bind membranes specific to these compartments (7). The manner in which aPKC is recruited into and excluded from these membrane-associated polarity complexes is still not entirely clear. Since there is an interdependency with other proteins such as Par3 and Cdc42, membrane association was long considered to be indirect occurring through association with these and other partner proteins (8–11). However two recent studies have shown at that aPKCι can associate directly with cell membranes through its regulatory module, defined hereafter as aPKCι^{RM} (12, 13). The membrane-binding determinants within the aPKCι^{RM} have been identified as the PS region, which contains a stretch of positively charged Arg residues, and the C1 domain. It is unclear hitherto what the relative contribution of each region is. For example, in HEK293 cells, substitutions in the PS region render the protein unable to interact with the plasma membrane, whereas in *Drosophila* neuroblasts the C1 domain has been identified as the dominant membrane interaction module (12, 13).

Here, we approach the question of how multiple aPKC membrane-binding determinants coordinate and how they are coupled. We use Alphafold *in silico* predicted models for the aPKC regulatory module combined with biochemical and cellular assays to validate a continuous surface harbouring known membrane-interacting determinants. We show that the PB1 and C1 domains are coupled and

63 oriented though interfacial contacts also aligning the PS motif and C1 domain for membrane

64 interaction. We identify a highly conserved phospho-Tyr residue within the PB1-C1 interface and show

65 that phosphorylation of this residue modulates membrane-binding capabilities of the regulatory

66 module. We propose that this phosphorylation influences the ability of aPKC to associate with and

67 disassociate from membranes in a regulated manner.

Experimental procedures

In silico/molecular modelling

70 To predict the tertiary structure of the regulatory domain we used Alphafold Colab

71 (https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.ipynb) (14) with

72 residues 22-194 of human aPKCι as input. The top scoring model coordinates were used for

73 subsequent prediction of membrane binding motifs using the Membrane Optimal Docking Algorithm

74 (MODA) (15) and ConSurf (16).

Cell lines and reagents

HEK293T cells were grown in Dulbecco's modified eagle medium (DMEM) supplemented with 10%

(v/v) Fetal Bovine serum (ThermoFisher scientific), 100U/ml Penicillin and 100μg/ml Streptomycin

(ThermoFisher scientific). HEK293FS cells were grown in Freestyle 293 Expression Medium

(ThermoFisher Scientific). Cells were obtained from the institute's central catalogue, where they are

routinely checked for mycoplasma. The phospho-Tyr136 antibody was made in-house using an

immunogenic peptide spanning residues 125-141 of haPKCι with modifications at Tyr-136 (phosphate)

and Cys-137 (S-methylcarboxyamide). Anti-thiophosphate ester antibody was from Abcam

(Cambridge, UK). Anti-FLAG M2 antibody, and FLAG-agarose resins were purchased from Sigma (St.

Louis, MO, USA). Streptactin II agarose was from GE healthcare (Little Chalfont, UK). Myc-trap

agarose was from Chromotek (Planegg, Germany) Anti-Myc, secondary HRP-linked goat anti-Rabbit

and Horse anti-Mouse antibodies were from Cell Signaling Technologies (Beverly, MA, USA).

Polyethyleneimine (PEI) was from Polysciences Inc. (Warrington, PA, USA). Mutagenesis and cloning

were done using In-Fusion (Takara, Shiga, JP). Primers and plasmids used are listed in Table S4.

Peptides used were made in-house.

Protein expression and purification

For expression of Strep-tagged Par6 co-expressed with Myc-tagged fl-aPKCι adherent HEK293T cells were transiently transfected using polyethylene-imine (PEI) at a 1:3 (m/m) plasmid/PEI ratio. Forty-eight hours post-transfection, cells were lysed in 50 mM Tris, pH 7.4, 150 mM NaCl, 1% Triton, 0.5mM TCEP supplemented with phosphatase inhibitors (Phosphostop, Roche, Germany), and protease inhibitors (cOmplete, Roche, Germany). Cell lysates were incubated with Streptactin II (GE healthcare) affinity beads for 2 hours at 4°C while rotating. Next, the beads were washed twice with Lysis buffer supplemented with NaCl to 500mM, once with Lysis buffer and once with elution buffer (50mM Tris pH7.4, 150mM NaCl, 0.5mM TCEP). The protein was eluted in elution buffer supplemented with 2.5mM D-desthiobiotin. For the purification of 2xStrep-tagged fl-aPKCι the same protocol was followed but HEK293FS cells were used and transfected for 96h instead of 48h. In addition, a size exclusion chromatography step was performed on a Superdex-200 Increase column and peak fractions were pooled, dialyzed against elution buffer and concentrated. Protein quantity and purity was assessed using SDS-PAGE and A280.

In vitro **kinase assay**

HEK293T cells expressing myc-tagged fl- aPKCι WT and mutants were lysed in lysis buffer (50 mM Tris, pH 7.4, 150 mM NaCl, 1% Triton, 0.5mM TCEP supplemented with phosphatase inhibitors (Phosphostop, Roche, Germany), and protease inhibitors (cOmplete, Roche, Germany)) and immunoprecipitated using Myc-Trap agarose (Chromotek). Beads were washed once in lysis buffer containing 500mM NaCl and twice in TBS. Equal amounts of the bead suspension were then added to 1.5µg myelin basic protein (MBP) and 250µM ATP-γS in assay buffer (1x final; 50mM Tris, 10mM MgCl$_2$). Reactions were incubated at 30°C for 30 min and subsequently added with 2.5 mM p-Nitrobenzyl mesylate (PNBM) for 1.5h at room temperature. Incorporation of S-labeled phosphates was followed via western blotting with an anti-thiophosphate ester antibody.

Differential scanning fluorometry

Protein at 250µg/ml was incubated with SPYRO-orange protein dye (1:400) in 50mM Tris, 150mM NaCl, 0.5mM TCEP with or without 1mM ATP and 10mM MgCl$_2$. The reactions were preincubated at room temperature for 5 min before being distributed into a MicroAmp 96-well qPCR plate. Samples were heated from 25-98°C in a QuantStudio Flex 12 RT-PCR machine (ThermoFisher Scientific). Measurements of fluorescence emission at 570nm were followed for each 0.3°C increment. Raw

curve data was exported and derivative curves were plotted. The T_m of protein in each well was

determined as the temperature at the inflection point of the melting curve using the maximal point of

the derivative.

Immunofluorescence

Cells were grown on 13 mm glass coverslips, transfected with constructs as indicated and 24h post

transfection cultures were treated with 500nM nocodazole for 16h before fixation or left untreated.

Cells were fixed with 4% PFA and permeabilized with PBS+0.1% Triton X-100. Coverslips were then

blocked in 3% BSA in PBS and incubated with FLAG M2 antibody (1:500). Secondary antibody was

1:1500 goat anti-mouse 488 (Thermo Fisher, A11001). Coverslips were mounted using Prolong gold

with DAPI (Thermo Fisher Scientific) and imaged. All the images were acquired using an inverted laser

scanning confocal microscope (Carl Zeiss LSM 880) using a 63x Plan-APOCHROMAT DIC oil-

immersion objective. The same exposure settings were used to acquire all images (excluding DAPI

channel). Images shown in figures were processed in ZEN Blue edition (Zeiss) where a gold LUT

pseudo-colour transformation was applied to the 488-emission channel. All images were batch-

processed to adjust brightness/contrast. Quantification of membrane and cytoplasmic intensities was

done on the raw images in ImageJ. The whole membrane region was traced with the freehand tool

and the average pixel intensity was recorded. In the same way the average intensity for a trace inside

in a sub-membrane region of the cytoplasm was recorded and the fraction of recorded intensities was

plotted.

Cell fractionation

To assess phosphorylation of Tyr-136 in the membrane and cytosolic fractions, HEK293T cells

expressing FLAG-tagged regulatory domain were fractionated as in [17]. Briefly, cells were lysed in

Homogenization buffer (20 mM Tris, pH 7.4, 10mM EDTA, supplemented with phosphatase inhibitors

(Phosphostop, Roche, Germany), and protease inhibitors (cOmplete, Roche, Germany)) and

subjected to sonication. The suspension was then centrifuged at 100.000xg for 20' at 4°C and the

supernatant (Soluble fraction) was collected. The pellet (Particulate fraction) was resuspended in

homogenization buffer containing 1% Triton X-100, sonicated and again centrifuged at 100.000g for

20' at 4°C. The supernatant was collected and FLAG-IP was performed on both supernatants to

isolate the regulatory domain for Western analysis with the pY136 antibody.

Reduction-alkylation for immunodetection with pTyr-136 antibody

Western blots analysing levels of Tyr-136 phosphorylation were performed according to standard procedures but with the subsequent additional steps. After transfer the nitrocellulose membrane was incubated overnight in PBS +10mM DDT. It then was washed twice in PBS and incubated with PBS +10mM iodoacetamide (Sigma) for 1h at room temperature. The membrane was washed twice with PBS and immunoblotting was continued according to standard procedures.

Lipid overlay assay

aPKCι-Par6 WT and mutant complexes were purified as described above using 2xStrep-Par6 as bait. The protein complexes were analysed via silver staining and incubated with lipid coated membranes (PIP-Strips cat. P-6001) (Echelon Bioscience). The PIP strips were subsequently probed with Myc-antibody (1:1000) and HRP-tagged secondary antibody (1:3000). Membranes were processed and exposed in parallel in an Imagequant LAS 4000 (GE Healthcare).

Results

A continuous membrane-binding surface is oriented within a compact PB1-PS-C1 aPKCι regulatory module

In light of the recent discrepancies in the relative contributions of PS and C1 domain to aPKC membrane interaction, we took a holistic approach to understand whether the regulatory module acts as a functional unit contributing two coupled membrane-binding determinants. We used AlphaFold Colab to predict the aPKCι regulatory module (aPKCι^{RM}) spanning the PB1 domain, the PS motif and the C1 domain [14]. The top five structural models predict a compact, globular shape with buried inter-domain contacts suggesting a coupled functional unit (Fig S1A). The models predict that the PB1 and C1 domains are crucially bridged together via an interdomain β-strand which we term beta-strand linker (BSL) (Fig 1A), that is part of a continuous sheet with the β-strands in the PB1 and C1 domains with a parallel orientation to the PB1 and antiparallel to the C1 domain. All of the models show very low predicted alignment errors (PAE), indicative of high confidence in the continuous fold (Fig S1B). Using the highest scoring AF Colab model as input, we set out to predict membrane binding residues via the Membrane Optimal Docking Algorithm (MODA) package [15]. Both the PS and two regions in the C1 domain (termed loop 1 and loop 2) have high membrane-association prediction scores (Fig 1B,C, Table S1). Interestingly, both regions form a continuous surface containing both positively

charged and hydrophobic residues (Fig 1C,D), potentially contributing to a single composite

membrane interaction site. The importance of these residues is further underscored by the fact that

most residues constituting the platform are highly conserved as shown via ConSurf analysis (Fig S1E,

Table S2), while there is less stringency for amino acid composition on the opposite side of the domain

(Fig S1F).

To probe the intrinsic capabilities of aPKCι^{RM} in membrane binding, we expressed FLAG-tagged

aPKCι^{RM} in HEK293T cells. We observed a striking localization pattern that was cell cycle-dependent.

In interphase cells the regulatory module is predominantly localized in the nucleus (Fig 1E), whereas

in mitotic cells, aPKCι^{RM} is uniformly associated with the plasma membrane (Fig 1E, asterisks). This

observation is in line with those from *Drosophila* neuroblasts where over-expressed aPKCι^{RM}

constructs display similar behaviour in interphase and at mitosis [13]. This property allowed us to

assess the membrane binding requirements of aPKCι^{RM} in a physiological context by arresting cells in

mitosis with nocodazole (Fig 1F). To exploit this "mitotic trap" assay, we introduced site-specific

mutations targeting residues with the highest relative prediction scores for membrane interaction

(Table S1). While the WT regulatory module has a strong (~4-fold) enrichment at the plasma

membrane in nocodazole treated cells (Fig 2A), we observed a complete loss of membrane binding

when substituting the highly predictive ^{160}RIWGL^{164} sequence in the C1 domain (loop 2) with

^{160}AIAGD^{164} (Fig 2B,F). Similarly, a complete loss of membrane binding was observed with

^{126}RRGARR^{131} to ^{126}AAGAAA^{131} substitutions in the PS region (termed PS 4R.A, Fig 2C,F).

Substitution of Arg-147 or Arg-150/151 to Ala in the C1 domain (loop 1) had a smaller impact reducing

membrane binding by about 1.5-fold (Fig 2D,E,F). This indicates that both regions in the C1 domain

and the PS motif contribute to allow optimal membrane binding in these mitotic cells. Based upon the

model, it is likely these motifs are organized in a continuous surface, the integrity of which likely

dictates the capacity to bind membranes.

Tyr-136 phosphorylation within an interfacial motif uncouples membrane-binding determinants

but does not impact catalytic activity

One of the more salient features of the regulatory module is the interfacial BSL motif positioning the

PB1 and C1 domain, orienting the membrane-binding determinants within the aPKCι^{RM}. The BSL

shows overall high conservation centred on a centrally positioned, highly conserved Tyr residue 136

(Fig S1G, Table S2). Inspection of Phosphosite indicated that this residue is frequently found to be

phosphorylated in more than 60 high-throughput studies in a variety of conditions and cell types,

alongside the familiar activation loop and turn motif priming sites (Fig 3A) [18]. Consistent with this, it

is also one of few aPKCι phospho-residues identified in the MaxQuant Database (MaxQB) (Table S3)

[19].

As Tyr-136 lies at the core of the regulatory module within the interfacial BSL motif and proximal to the

PS (Fig 3B), we considered whether tyrosine phosphorylation could influence the full-length kinase

conformation, stability and activation state. First, we probed whether phosphorylation affects the

interaction between the PS region and the kinase domain. By definition, the PS is proposed to bind the

catalytic domain at the substrate-binding pocket to be released upon aPKCι activation. We wondered

whether in a linearized context, Tyr phosphorylation proximal to the PS could influence this behaviour.

To test this, we immunoprecipitated a Myc-tagged kinase domain (aPKCι^{KD}) from cell lysates using a

biotin-labeled extended pseudo-substrate peptide (ePSP) containing either a non-phosphorylated Tyr

or pTyr residue or Glu or Phe substitutions to mimic either phosphorylated or non-phosphorylated

states respectively (Fig 3C). We saw specific binding of the peptide to the core kinase domain and

observed loss of kinase domain interaction with the peptide when the Tyr residue is either

phosphorylated or substituted with Glu. Conversely a non-phosphorylated or Phe substituted peptide

effectively captured the aPKCι^{KD}. To confirm whether the phosphorylation of this residue reduces the

interaction between aPKCι kinase domain and its regulatory module, we assessed the protection of

the kinase core by co-expressing the isolated kinase domain with either wild type (WT) or mutant

forms of aPKCι^{RM} (Fig 3D). While the WT aPKCι^{RM} protein product protects the aPKCι^{KD}, a Y136E

substitution causes a reduction in the levels of aPKCι^{KD} protein, as does a previously described

pseudo-substrate disengaged A129E mutant [4]. These results indicate that the kinase domain-

regulatory module interaction is weakened by phosphorylation of Tyr-136.

To probe whether this effect is due to loss of direct binding of the PS region to the kinase domain or

the aPKCι^{RM} conformation, we purified WT, Y136E and A129E mutant full-length aPKCι (aPKCι^{FL}) from

HEK293FS cells and subjected them to differential scanning fluorometry (DSF) (Fig 3E). In the apo-

form, the melting temperature (T_m) for the WT protein is 53.3°C, similar to the A129E mutant (T_m =

53.7°C). The Y136E mutant is less thermostable with a T_m of 51.3°C in the apo-form. Addition of

ATP.Mg^{2+} stabilizes both WT and mutant kinases, however comparatively the A129E mutant is

stabilized to a lesser degree than WT (ΔT_m= 6.7°C vs WT ΔT_m= 10.2°C). This suggests that

occupation of the active site with ATP, while stabilizing the kinase core, may result in a less

thermostable open conformation, possibly because of phosphate-driven charge repulsion with the Glu

substitution. The Y136E mutant is less thermostable than WT in both the presence and absence of

ATP, indicating that decreased thermostability is caused by mutation-driven alterations in the

regulatory module.

To test the impact of Tyr-136 phosphorylation on aPKCι catalytic activity, we followed the

phosphorylation of MBP *in vitro* after immunoprecipitation of WT and mutant forms of aPKCι^{FL}. To

monitor the incorporation of new phosphate groups on MBP we made use of γ-thiophosphate ATP (Fig

3F). MBP was readily phosphorylated by the conformationally "open" A126E mutant but showed only a

marginal nonsignificant increase in phosphorylation upon incubation with a Y136E mutant compared

to WT. This indicates that in the 'tethered' state of the full-length protein, the impact of the weakened

regulatory module/catalytic domain interaction consequent to Tyr-136 phosphorylation is modest

compared to the pseudo-substrate A129E mutation, while there is a strong effect on the overall

aPKCι^{RM} conformation.

Tyr-136 phosphorylation disrupts aPKCι^{RM} membrane binding in cells

Since the above results indicate a change in regulatory module conformation with little impact on

activity, we tested whether Tyr-136 phosphorylation had a direct influence on the aPKCι^{RM} membrane

binding properties. We assessed membrane occupation in mitotic cells as above, expressing the WT

regulatory module (Fig 4A) or mutants in nocodazole treated cells; the PS 4R.A mutant was used as a

control for loss of membrane binding. Substitution of Tyr-136 with Glu resulted in a significant 1.5-fold

reduction of membrane intensity (Fig 4B,F), whereas a Tyr-Phe substitution had a negligible impact on

membrane binding (Fig 4C,F), consistent with Tyr phosphorylation being disruptive for membrane

interaction. Interestingly a substitution of A129 to Glu had only a small but significant impact on

membrane binding capabilities, despite its negative charge central to the PS region (Fig 4D,F),

whereas substitutions of the PS Arg residues resulted in a complete loss of membrane binding (Fig

4E,F).

To probe Tyr-136 phosphorylation in the regulatory module of aPKC we developed a site-specific and

phospho-specific rabbit antibody. Initial attempts resulted in poor and nonspecific immunoreactivity,

but using a derivatized phosphopeptide with S-methylcarboxyamide coupled to the adjacent Cys residue C-terminal to the phosphorylated Tyr residue, we were able to detect Tyr phosphorylated Myc-aPKC in pervanadate treated cells after immunoprecipation (Fig S2A). The alkylation greatly improved epitope recognition, specific to nitrocellulose membranes following alkylation of proteins with iodoacetamide, confirming the double-modified epitope recognition (Fig S2A,B). Interestingly, using this antibody, we could show significant levels of Tyr-136 phosphorylation in regulatory module constructs in nocodazole treated cells (Fig 4G). The Y136 phosphorylation of the 'free' aPKCι^{RM} in mitotic cells, suggested that this was not due to the aPKCι kinase domain exhibiting Tyr specificity due to proximity (*in casu* the PS motif), as seen in other Ser/Thr kinases as recently observed in the PKDs (21). Indeed, we could detect no phosphorylation of Tyr-136 upon incubation of aPKCι^{FL} with ATP.Mg^{2+} (data not shown). Exploring candidate upstream kinases, we pulled down proteins from a cell extract using a biotinylated ePSP peptide containing either a Y, F or pY residue corresponding to Tyr-136 and then spiked the precipitated protein mixture with ATP.Mg^{2+}, allowing any associated kinase to (auto)phosphorylate. Western blot analysis with a phospho-Tyr antibody showed tyrosine kinase (autophosphorylation) activity around ~60kDa and ~170kDa in protein mixtures associating with the Y and F but not the product pY peptide (Fig S3A). In a subsequent screen with a panel of Tyr kinase inhibitors we found responsiveness of aPKCι^{RM} Tyr-136 phosphorylation in nocodazole treated conditions to the Src family kinase (SFK) inhibitors PP2 and dasatinib (Fig S3B). This implicates SFKs in targeting this site and is consistent with the finding of a ~60KDa tyrosine phosphorylated protein(s) captured on the site-affinity matrix.

The fact that Tyr-136 is phosphorylated in nocodazole treated cells led us to ask the question whether the phosphorylated species segregates to the membrane or cytosolic compartment in mitotic cells (Fig 4H). To assess any preferential localization of the phosphorylated species we fractionated nocodazole-treated cells by ultracentrifugation (100,000xg) to separate membrane (particulate) and cytosolic (soluble) components. We could detect no phosphorylation of the regulatory module in the particulate fraction indicating that this species does not associate with the membrane fraction, while phosphorylated protein was detected in the cytosol (Fig 4I). This further supports our observation with the Glu phospho-mimetic mutant, showing that tyrosine phosphorylation impacts the protein cycling between membrane and cytosolic fractions.

To further assess direct lipid binding in the context of the full-length protein, including the effects of

increased dynamics between regulatory and kinase domain in the mutants, we purified the aPKC-Par6

complex from HEK293T cells using Par6 as bait (Fig 5A). The binary complex is thought to be in a

more open conformation as Par6 is proposed to displace the PS region, potentially exposing the

aPKCι[RM] for lipid binding (4). These binary protein complexes were then used in a lipid overlay assay

with PIP-strips (Fig 5B). The WT complex interacts preferentially with monophosphorylated inositol

headgroups, (PI3P, PI4P, PI5P) but also with doubly phosphorylated headgroups (PI(3,4)P_2,

PI(3,5)P_2,PI(4,5)P_2) with no strong preference for the position of the phosphate (Fig 5B). The complex

also binds phosphatidylserine while we observed no binding to Sphingosine-1-phosphate (S1P) which

was previously described as a lipid activator for aPKCs (20). Substituting Tyr-136 with Glu results in

reduced binding to all phospholipids, whereas a Phe mutant displays WT characteristics (Fig 5B).

Interestingly a PS-disengaged mutant (A129E) results in increased lipid binding despite the glutamic

substitution. This is indicative of the fact that this mutation can further drive the Par6-bound open

conformation, allowing for enhanced lipid binding. Consistent with the mitotic trap assay the Ala to Glu

substitution only seems to have a small effect on membrane binding.

Discussion

Cell polarity - the spatially asymmetric organization of a cell and its components - is an essential

biological process, orchestrated by key regulatory inputs. One central component of this process

conserved in higher eukaryotes is atypical protein kinase Cι. Together with its partner Par6 this kinase

is involved in regulating a variety of protein complexes, prominently those including Par3, Lgl1/2, Crb,

but also Dlg, Par-1, FARP2, Yurt, and others (22–28). While there are functional data on the

interdependency of polarity components, molecular insight into the regulation of these assemblies and

their distribution is still lacking. For example, it is not clear how the aPKCs are recruited to membranes

and membrane-associated protein assemblies. Several studies have shown a dependency on Par-3

(8–11), however, Par3 independent mechanisms have also been described (29). The correct lipid

environment plays a crucial role in recruiting and tethering the right polarity components, with anionic

phospholipids and prominently PI(4,5)P_2 playing a dominant role in the apical domain (30, 30, 31).

Several polarity proteins respond to these lipids via basic-hydrophobic (BH) motifs (32, 33).

Interestingly, in a recent study the pseudo-substrate of aPKCs was identified as such a lipid-

responsive BH domain in cells (12). This aligns with the long-standing notion that the PKC pseudo-

substrate region interacts with negatively charged lipids (34). In addition the response to PIP3

downstream of insulin has also been mapped previously to the PS region (35). However a subsequent

study by the Prehoda group showed that in dividing drosophila neuroblasts membrane targeting was

dependent on the C1 domain, also via an anionic lipid-binding mechanism (13).

Here we have investigated membrane binding by aPKCι^{RM} making use of recent advances in AI by

Alphafold to derive a structural model for multi-domain modules of proteins. While attempts at

purification of aPKCι^{RM} for structure determination were unsuccessful, AlphaFold Colab predictions

indicated it adopts a single compact unit comprising the PB1 and C1 domains, in which they are

tethered via an interdomain β-strand proximal to the pseudo-substrate embedded in a continuous

sheet spanning across both domains. Based on this predicted model we used the MODA prediction

algorithm which identified 3 regions containing putative membrane-binding residues. Interestingly,

together these predicted membrane-binding regions form a highly conserved continuous platform in

the aPKCι^{RM} model. Aside from the PS sequence ^{126}RRGARRWRK^{134} we identify ^{160}RIWGLGRQ^{167} in

the C1 domain as a second sequence strongly contributing to membrane binding. Mutagenesis in

either of these sequences completely disrupts membrane binding in the mitotic cell assay, indicating

that neither alone provides sufficient affinity and an intact platform is required to establish the avidity

effect for optimal membrane binding. Additionally, Arg-147 and Arg-150/151 in the C1 domain also

contribute albeit to a lesser degree.

The organization of the membrane binding platform is crucially driven by the interfacial BSL motif, with

at its core an evolutionarily conserved phospho-Tyr residue that allows for regulatory input.

Interestingly we observed this dynamic behaviour in mitotic cells where we found that aPKCι Tyr-136

was phosphorylated only in the cytoplasm. Whether the released protein retains an active, open

conformation impacting cytosolic targets has yet to be determined, although our evidence indicates

that a tyrosine phosphorylation mimetic alone is not substantially activating despite weakening the

inhibitory interaction of the regulatory module on the catalytic domain (Table 1). Based on the effects

on thermostability, we propose that the PB1 and C1 domains become discontinuous when Tyr-136 in

the BSL is phosphorylated, compromising the overall conformation and membrane binding (Table1,

Fig 5C, left hand side). This distinguishes the BSL from the pseudo-substrate, which, besides being

part of the aPKC membrane interaction module, regulates kinase activity by direct interaction with the

kinase domain evidenced by a loss of kinase domain stabilization and increased catalytic activity upon

A129E substitution (Table 1). The pseudo-substrate is therefore the main regulatory unit influencing

catalysis (Fig 5C, right hand side).

Construct	WT	Y136E	A129E
Kinase domain stabilization *(co-expression assay)*			
Catalytic activity for MBP *(in vitro kinase assay)*			
aPKC^RM membrane binding *(mitotic trap assay)*			
Thermostability (+ATP.Mg^2+) *(DSF assay)*			

Table 1. effects measured in the respective assays. Colour-coded based on the average

measurements obtained in the assays, cells conditionally formatted from red (low) to green (high).

The observation that there is a pTyr residue central to the regulation of membrane binding implicates

cross-talk with Tyr kinase activities contributing to a polarized distribution of the aPKCs. The role for

Tyr kinase activity in junction formation and cell polarity was clear early on (36–38), however, few

molecular links have been established. We show for Tyr-136 that SFKs are potential upstream

regulators. Interestingly, Src has long been known to associate with aPKCs, docking to their PXXP

rich motifs N-terminal to the pseudo-substrate site; an in vitro phosphorylation reaction with Src using

aPKC-derived peptides also included a peptide containing pTyr-136, further supporting the fact that

SFKs can act upstream (39, 40). The cell-cycle dependency of Tyr-136 phosphorylation we observed

raises intriguing questions for further exploration regarding the Src-aPKC connection in asymmetrical

cell division. v-Src transformation of the zebrafish embryo enveloping layer epithelium causes mitotic-

dependent extrusion, partially by affecting cell polarity through Cdc42 and aPKC (41 42). Src also

promotes epithelial polarity switching in colorectal cancer spheroids, promoting a switch from apical-in

(central lumen) to apical-out oriented metastatic cell clusters (43). Other Src family kinases such as

Fyn and Lyn have also been involved in the regulation of cell polarity, opening up further possibilities

of connecting PTK and polarity pathways (44, 45).

Recruitment and retainment of aPKC at membranes involves a variety of factors that are either PKC

intrinsic or extrinsic through membrane-anchored proteins. This complex mesh allows for a variety of

inputs and affinity thresholds. We show here the mechanism by which aPKCs can interact with

membranes through a continuous interaction platform spanning the whole regulatory region,

orchestrated by an interdomain β-strand that can be regulated via tyrosine phosphorylation. The

hierarchy and local mechanisms by which aPKCs are recruited and retained at the correct membrane

locus, as well as a further understanding of tyrosine kinase crosstalk with cell polarity, are intriguing

questions for future exploration.

Acknowledgements

384 We thank the Crick Microscopy, Structural Biology and Peptide Synthesis science technology
385 platforms for reagents and expertise. We thank Christopher Earl and David C. Briggs for crystallization
386 efforts and lab members in the Parker and McDonald labs for constructive discussions. The study was
387 funded by the Francis Crick Institute, which receives its core funding [FC001115 and FC001130] from
388 Cancer Research UK, UK Medical Research Council and the Wellcome Trust.

Competing interest statement

391 The authors declare that they have no known competing financial interests or personal relationships

392 that could have appeared to influence the work reported in this paper.

References

1. Parker, P. J., Brown, S. J., Calleja, V., Chakravarty, P., Cobbaut, M., Linch, M., Marshall, J. J. T., Martini, S., McDonald, N. Q., Soliman, T., and Watson, L. (2021) Equivocal, explicit and emergent actions of PKC isoforms in cancer. *Nat Rev Cancer*. **21**, 51–63

2. Leroux, A. E., Schulze, J. O., and Biondi, R. M. (2018) AGC kinases, mechanisms of regulation and innovative drug development. *Seminars in Cancer Biology*. **48**, 1–17

3. Parker, P. J., Justilien, V., Riou, P., Linch, M., and Fields, A. P. (2014) Atypical Protein Kinase Cι as a human oncogene and therapeutic target. *Biochem Pharmacol*. **88**, 1–11

4. Graybill, C., Wee, B., Atwood, S. X., and Prehoda, K. E. (2012) Partitioning-defective Protein 6 (Par-6) Activates Atypical Protein Kinase C (aPKC) by Pseudosubstrate Displacement. *J Biol Chem*. **287**, 21003–21011

5. Hirano, Y., Yoshinaga, S., Takeya, R., Suzuki, N. N., Horiuchi, M., Kohjima, M., Sumimoto, H., and Inagaki, F. (2005) Structure of a Cell Polarity Regulator, a Complex between Atypical PKC and Par6 PB1 Domains*. *Journal of Biological Chemistry*. **280**, 9653–9661

6. Hirano, Y., Yoshinaga, S., Ogura, K., Yokochi, M., Noda, Y., Sumimoto, H., and Inagaki, F. (2004) Solution Structure of Atypical Protein Kinase C PB1 Domain and Its Mode of Interaction with ZIP/p62 and MEK5*. *Journal of Biological Chemistry*. **279**, 31883–31890

7. Buckley, C. E., and St Johnston, D. (2022) Apical–basal polarity and the control of epithelial form and function. *Nat Rev Mol Cell Biol*. 10.1038/s41580-022-00465-y

8. Wodarz, A., Ramrath, A., Grimm, A., and Knust, E. (2000) Drosophila Atypical Protein Kinase C Associates with Bazooka and Controls Polarity of Epithelia and Neuroblasts. *J Cell Biol*. **150**, 1361–1374

9. Franz, A., and Riechmann, V. (2010) Stepwise polarisation of the Drosophila follicular epithelium. *Developmental Biology*. **338**, 136–147

10. Wang, S.-C., Low, T. Y. F., Nishimura, Y., Gole, L., Yu, W., and Motegi, F. (2017) Cortical forces and CDC-42 control clustering of PAR proteins for *Caenorhabditis elegans* embryonic polarization. *Nature Cell Biology*. **19**, 988–995

11. Martin-Belmonte, F., Gassama, A., Datta, A., Yu, W., Rescher, U., Gerke, V., and Mostov, K. (2007) PTEN-mediated apical segregation of phosphoinositides controls epithelial morphogenesis through Cdc42. *Cell*. **128**, 383–397

12. Dong, W., Lu, J., Zhang, X., Wu, Y., Lettieri, K., Hammond, G. R., and Hong, Y. (2020) A polybasic domain in aPKC mediates Par6-dependent control of membrane targeting and kinase activity. *J Cell Biol*. 10.1083/jcb.201903031

13. Jones, K. A., Drummond, M. L., and Prehoda, K. E. (2022) *Cooperative regulation of C1-domain membrane recruitment polarizes atypical Protein Kinase C*, 10.1101/2022.01.01.474707

14. Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M. (2022) ColabFold: Making Protein folding accessible to all. *Nature Methods*. 10.1038/s41592-022-01488-1

15. Kufareva, I., Lenoir, M., Dancea, F., Sridhar, P., Raush, E., Bissig, C., Gruenberg, J., Abagyan, R., and Overduin, M. (2014) Discovery of novel membrane binding structures and functions. *Biochem Cell Biol*. **92**, 555–563

16. Ashkenazy, H., Abadi, S., Martz, E., Chay, O., Mayrose, I., Pupko, T., and Ben-Tal, N. (2016) ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation in macromolecules. *Nucleic Acids Research*. **44**, W344–W350

17. Wescott, G. G., Manring, C. M., and Terrian, D. M. (1999) Translocation Assays of Protein Kinase C Activation. in *Neurodegeneration Methods and Protocols* (Harry, J., and Tilson, H. A. eds), pp. 125–132, Methods in Molecular Medicine™, Humana Press, Totowa, NJ, 10.1385/0-89603-612-X:125

18. Hornbeck, P. V., Zhang, B., Murray, B., Kornhauser, J. M., Latham, V., and Skrzypek, E. (2015) PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. *Nucleic Acids Res*. **43**, D512-520

19. Schaab, C., Geiger, T., Stoehr, G., Cox, J., and Mann, M. (2012) Analysis of high accuracy, quantitative proteomics data in the MaxQB database. *Mol Cell Proteomics*. **11**, M111.014068

20. Kajimoto, T., Caliman, A. D., Tobias, I. S., Okada, T., Pilo, C. A., Van, A.-A. N., Andrew McCammon, J., Nakamura, S.-I., and Newton, A. C. (2019) Activation of atypical protein kinase C by sphingosine 1-phosphate revealed by an aPKC-specific activity reporter. *Sci Signal*. 10.1126/scisignal.aat6662

21. Cobbaut, M., Derua, R., Parker, P. J., Waelkens, E., Janssens, V., and Van Lint, J. (2018) Protein kinase D displays intrinsic Tyr autophosphorylation activity: insights into mechanism and regulation. *FEBS Letters*. **592**, 2432–2443

22. Elbediwy, A., Zhang, Y., Cobbaut, M., Riou, P., Tan, R. S., Roberts, S. K., Tynan, C., George, R., Kjaer, S.,
Martin-Fernandez, M. L., Thompson, B. J., McDonald, N. Q., and Parker, P. J. (2019) The Rho family GEF
FARP2 is activated by aPKCι to control tight junction formation and polarity. *J Cell Sci*. 10.1242/jcs.223743
23. Golub, O., Wee, B., Newman, R. A., Paterson, N. M., and Prehoda, K. E. (2017) Activation of Discs large by
aPKC aligns the mitotic spindle to the polarity axis during asymmetric cell division. *eLife*. **6**, e32137
24. Soriano, E. V., Ivanova, M. E., Fletcher, G., Riou, P., Knowles, P. P., Barnouin, K., Purkiss, A., Kostelecky, B.,
Saiu, P., Linch, M., Elbediwy, A., Kjær, S., O'Reilly, N., Snijders, A. P., Parker, P. J., Thompson, B. J., and
McDonald, N. Q. (2016) aPKC Inhibition by Par3 CR3 Flanking Regions Controls Substrate Access and
Underpins Apical-Junctional Polarization. *Dev. Cell*. **38**, 384–398
25. Yamanaka, T., Horikoshi, Y., Sugiyama, Y., Ishiyama, C., Suzuki, A., Hirose, T., Iwamatsu, A., Shinohara, A.,
and Ohno, S. (2003) Mammalian Lgl Forms a Protein Complex with PAR-6 and aPKC Independently of
PAR-3 to Regulate Epithelial Cell Polarity. *Current Biology*. **13**, 734–743
26. Sotillos, S., Díaz-Meco, M. T., Caminero, E., Moscat, J., and Campuzano, S. (2004) DaPKC-dependent
phosphorylation of Crumbs is required for epithelial cell polarity in Drosophila. *J Cell Biol*. **166**, 549–557
27. Hurov, J. B., Watkins, J. L., and Piwnica-Worms, H. (2004) Atypical PKC Phosphorylates PAR-1 Kinases to
Regulate Localization and Activity. *Current Biology*. **14**, 736–741
28. Gamblin, C. L., Parent-Prévost, F., Jacquet, K., Biehler, C., Jetté, A., and Laprise, P. (2018) Oligomerization
of the FERM-FA protein Yurt controls epithelial cell polarity. *J. Cell Biol*. **217**, 3853–3862
29. Shahab, J., Tiwari, M. D., Honemann-Capito, M., Krahn, M. P., and Wodarz, A. (2015) Bazooka/PAR3 is
dispensable for polarity in Drosophila follicular epithelial cells. *Biology Open*. **4**, 528–541
30. Kanemaru, K., Shimozawa, M., Kitamata, M., Furuishi, R., Kayano, H., Sukawa, Y., Chiba, Y., Fukuyama, T.,
Hasegawa, J., Nakanishi, H., Kishimoto, T., Tsujita, K., Tanaka, K., Itoh, T., Sasaki, J., Sasaki, T., Fukami, K.,
and Nakamura, Y. (2022) Plasma membrane phosphatidylinositol (4,5)-bisphosphate is critical for
determination of epithelial characteristics. *Nat Commun*. **13**, 2347
31. Lu, J., Dong, W., Hammond, G. R., and Hong, Y. (2022) Hypoxia controls plasma membrane targeting of
polarity proteins by dynamic turnover of PI4P and PI(4,5)P2. *eLife*. **11**, e79582
32. Bailey, M. J., and Prehoda, K. E. (2015) Establishment of Par-Polarized Cortical Domains via
Phosphoregulated Membrane Motifs. *Dev Cell*. **35**, 199–210
33. Brzeska, H., Guag, J., Remmert, K., Chacko, S., and Korn, E. D. (2010) An Experimentally Based Computer
Search Identifies Unstructured Membrane-binding Sites in Proteins. *J Biol Chem*. **285**, 5738–5747
34. Mosior, M., and McLaughlin, S. (1991) Peptides that mimic the pseudosubstrate region of protein kinase
C bind to acidic lipids in membranes. *Biophys J*. **60**, 149–159
35. Ivey, R. A., Sajan, M. P., and Farese, R. V. (2014) Requirements for Pseudosubstrate Arginine Residues
during Autoinhibition and Phosphatidylinositol 3,4,5-(PO4)3-dependent Activation of Atypical PKC. *J Biol
Chem*. **289**, 25021–25030
36. Meyer, T. N., Schwesinger, C., Ye, J., Denker, B. M., and Nigam, S. K. (2001) Reassembly of the tight
junction after oxidative stress depends on tyrosine kinase activity. *J Biol Chem*. **276**, 22048–22055
37. Tsukamoto, T., and Nigam, S. K. (1999) Role of tyrosine phosphorylation in the reassembly of occludin
and other tight junction proteins. *Am J Physiol*. **276**, F737-750
38. Yap, A. S., Stevenson, B. R., Cooper, V., and Manley, S. W. (1997) Protein tyrosine phosphorylation
influences adhesive junction assembly and follicular organization of cultured thyroid epithelial cells.
Endocrinology. **138**, 2315–2324
39. Wooten, M. W., Vandenplas, M. L., Seibenhener, M. L., Geetha, T., and Diaz-Meco, M. T. (2001) Nerve
Growth Factor Stimulates Multisite Tyrosine Phosphorylation and Activation of the Atypical Protein
Kinase C's via a src Kinase Pathway. *Mol Cell Biol*. **21**, 8414–8427
40. Suzuki, A., Akimoto, K., and Ohno, S. (2003) Protein Kinase C λ/ι (PKCλ/ι): A PKC Isotype Essential for the
Development of Multicellular Organisms. *The Journal of Biochemistry*. **133**, 9–16
41. Anton, K. A., Kajita, M., Narumi, R., Fujita, Y., and Tada, M. (2018) Src-transformed cells hijack mitosis to
extrude from the epithelium. *Nat Commun*. **9**, 4695
42. Wang, Y., Du, D., Fang, L., Yang, G., Zhang, C., Zeng, R., Ullrich, A., Lottspeich, F., and Chen, Z. (2006)
Tyrosine phosphorylated Par3 regulates epithelial tight junction assembly promoted by EGFR signaling.
The EMBO Journal. **25**, 5058–5070
43. Okuyama, H., Kondo, J., Sato, Y., Endo, H., Nakajima, A., Piulats, J. M., Tomita, Y., Fujiwara, T., Itoh, Y.,
Mizoguchi, A., Ohue, M., and Inoue, M. (2016) Dynamic Change of Polarity in Primary Cultured Spheroids
of Human Colorectal Adenocarcinoma and Its Role in Metastasis. *The American Journal of Pathology*.
186, 899–911

44.   Morinaga, T., Yanase, S., Okamoto, A., Yamaguchi, N., and Yamaguchi, N. (2017) Recruitment of Lyn from
endomembranes to the plasma membrane through calcium-dependent cell-cell interactions upon
polarization of inducible Lyn-expressing MDCK cells. *Sci Rep*. **7**, 493
45.   Luo, J., Mcginnis, L. K., and Kinsey, W. H. (2009) Fyn Kinase Activity Is Required for Normal Organization
and Functional Polarity of the Mouse Oocyte Cortex. *Mol Reprod Dev*. **76**, 819–831
509
510

Figure legends

Figure 1. A. Representation of the aPKCι regulatory module as predicted by Alphafold Colab (best scoring model). The individual domains and regions are indicated and coloured; from N-C PB1 domain (cyan), Pseudo-substrate (PS) region (dark purple), Beta-Strand Linker (BSL) (light purple), C1 domain (blue). B. Per residue prediction score of the MODA prediction with above-threshold scoring regions indicated. C. Cartoon diagram (omitting secondary structure features) of predicted membrane interacting regions from the MODA prediction displayed in the AF Colab model. D. Mesh representation of panel C showing a continuous sidechain surface E. Localization of FLAG-tagged aPKCι regulatory module in HEK293T cells; red asterisks indicate mitotic cells. F. Localization of FLAG-tagged aPKCι regulatory module in nocodazole-treated (500nM, 16h) HEK293T cells.

Figure 2. A. Localization of WT FLAG-tagged regulatory module in nocodazole-treated (500nM, 16h) HEK293T cells. B-D. Expression of the indicated mutant FLAG-tagged regulatory modules in nocodazole-treated (500nM, 16h) HEK293T cells. Mutated residues are depicted on the protein model in red. E. Quantification of membrane/cytoplasmic pixel intensities measured in >45 cells imaged via confocal microscopy. Representative experiment of 3 biological replicates. Differences analysed via one-way ANOVA (****$P \leq 0.0001$)

Figure 3. A. Identification of main phosphorylation sites on aPKCι identified in the Phosphosite database B. visualization of Tyr-136 central to the BSL region of the regulatory module, colour coded as in Fig. 1A C. Peptide-pulldown of Myc-tagged aPKCι kinase domain with extended pseudosubstrate peptides (Biotin-Ahx-SIYRRGARRWRKLYCANGHT-CONH2) containing the indicated modifications at Y^{14}, i.e. pY, Y/E and Y/F. D. Co-expression of FLAG-tagged regulatory module and Myc-tagged kinase domain of aPKC in HEK293T cells. Equal amounts of DNA were transfected and stabilization of Myc-tagged kinase domain by different mutants of the regulatory module was followed. Quantification of Myc-kinase domain levels relative to WT aPKCι^{RM} co-expression shown in the right panel (n=3 biological replicates, average +/- SEM, analysed via one-sample t-test (v.a.v. a value of 1): ns $P >$ 0.05, * $P \leq 0.05$, *** $P \leq 0.001$). E. DSF assay measuring thermal stability of WT and mutant fl-aPKCι in absence or presence of ATP.Mg^{2+}. Plotted is the T_m derived from the inflection point of the melting curve (using the derivative maximum). F. *In vitro* kinase assay (IVKA) with immunoprecipitated WT or mutant kinases on MBP. Incorporation of new phosphates into MBP was followed with ATP-γS.

Quantification of MBP thiophosphate levels shown on the right (n=3 biological replicates, average +/-

SEM, analysed via unpaired t-test: ns P > 0.05)

Figure 4. A-E. Expression of the indicated mutant FLAG-tagged regulatory modules in nocodazole-

treated (500nM, 16h) HEK293T cells. F. Quantification of membrane/cytoplasmic pixel intensities

measured in >45 cells imaged via confocal microscopy. Representative experiment of 3 biological

replicates. Differences analysed via one-way ANOVA (ns P > 0.05, ****P ≤ 0.0001) G. Western blot

analysis of phospho-Tyr-136 levels in the in untreated (mixed pop.) cells or cells treated with 500nM

Nocodazole for 16h (Noc.). H. Representation of the observed localization of regulatory module

species and possible mapping of its phospho-form. I. Fractionation of cells by ultracentrifugation and

probing for Tyr-136 phosphorylation of the regulatory module (representative experiment of n=2

biological replicates).

Figure 5. A. Silver-stained SDS-PAGE gel of the Myc-aPKC-2xStrep-Par6 binary complexes purified

through 2xStrep-Par6. B. Lipid overlay assay using the protein complexes defined in A. Levels of

aPKCι bound to the PIP-strip membrane were quantified using anti-Myc tag antibody. C. schematic

representation of the impact of mutants and phospho-states of aPKCι. The left panel displays the

anticipated consequence of Tyr-136 phosphorylation in the BSL as mimicked by a Y136E substitution,

disrupting the continuous membrane binding platform constituted by the PS and C1 domain. The right

panel displays the impact of a pseudo-substrate disengaged conformation, mimicked by an A129E

substitution which mainly results in relieved auto-inhibition and increased catalysis.

Figure 1

Figure 2

Figure 3

Figure 4

A

WT Y136E Y136F A129E

aPKC

Par6

B

WT Y136E Y136F A129E

Myc

C

PB1 C1 KD → Y136E → PB1 C1 KD

KD C1 PB1 → A129E → KD C1 PB1

Figure 5