Phylogenetic comparative approach reveals evolutionary conservatism, ancestral composition, and integration of vertebrate gut microbiota

Benoît Perez-Lamarque [1,2] * (ORCID: 0000-0001-7112-7197)
Guilhem Sommeria-Klein [3] (ORCID: 0000-0002-5331-3639)
Loréna Duret [1] (ORCID: 0000-0001-7031-4900)
Hélène Morlon [1] (ORCID: 0000-0002-3195-7521)

[1] Institut de biologie de l'École normale supérieure (IBENS), École normale supérieure, CNRS, INSERM, Université PSL, 46 rue d'Ulm, 75 005 Paris, France

[2] Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'histoire naturelle, CNRS, Sorbonne Université, EPHE, UA, CP39, 57 rue Cuvier 75 005 Paris, France

[3] Department of Computing, University of Turku, Yliopistonmäki, 20014 Turku, Finland

* corresponding author: benoit.perez@ens.psl.eu; ORCID: 0000-0001-7112-7197; +33144323535, 46 rue d'Ulm, 75 005 Paris, France

Author Contributions: BPL, GSK, LD, and HM designed the study. BPL and GSK implemented the model. BPL performed the simulations and the empirical applications. BPL, GSK, and HM wrote the manuscript.

Competing Interest Statement: The authors declare no conflict of interest.

Classification: Biological Sciences; Evolution

Keywords: gut microbiome, interactome, phylosymbiosis, holobiont evolution, phylogenetic signal, comparative methods.

Abstract:

Host-associated microbial communities can play a central role in the adaptation of their hosts to different environments during evolutionary diversifications. Yet, little is known about the deep-time evolution of these microbiota: How conserved is their composition? What was the composition of ancestral microbiota? Do microbial taxa covary in abundance over millions of years? Multivariate phylogenetic models of trait evolution are key to answering similar questions for complex host phenotypes, yet they are not directly applicable to the relative abundances of microbial taxa. Here, we extend these models to the case of compositional data, thereby providing a powerful approach for estimating phylosymbiosis (*i.e.* the extent to which closely related host species harbor similar microbiota), ancestral microbiota composition, and integration (*i.e.* evolutionary covariations in bacterial abundances). We then apply our model to gut bacteria data for >200 mammal and >300 bird species. We find a significant phylosymbiosis in both mammals and birds that is not entirely explained by diet and biogeography, indicating that other evolutionary-conserved traits shape microbiota composition. We identify main shifts in microbiota composition during the evolution of the two groups and infer an ancestral mammalian microbiota consistent with an insectivorous diet. We also find remarkably consistent evolutionary covariations among bacterial orders in mammals and birds, suggesting strong constraints in gut microbiota evolution. Our results highlight the importance of host evolutionary history in shaping the gut microbiota composition of two major vertebrate groups and provide a phylogenetic comparative model for microbiota evolution broadly applicable across microbiota-host systems.

Significance statement:

Little is known about how host-associated microbial communities evolve as their hosts diversify. We have developed a novel approach to infer deep-time evolutionary trends in microbiota composition from the phylogenies of present-day hosts and their associated microbiota. Our approach allows testing for phylosymbiosis, i.e. whether the microbiota is evolutionary conserved, reconstructing ancestral microbiota composition, and identifying microbial taxa that vary in a concerted way. Applying our approach to the gut microbiota of mammals and birds, we find that the microbiota of both vertebrate groups is evolutionary conserved, and infer an ancestral mammalian microbiota consistent with an insectivorous diet. We observe conserved covariations in abundances among bacterial orders that suggest previously unsuspected constraints in vertebrate gut microbiota over millions of years.

Introduction:

Host-associated microbial communities, referred to as the *microbiota*, often play central roles in the biology of the hosts and their interactions with the environment. As host clades diversify, the microbiota can furthermore play a key role in the adaptation of their hosts to different ecological conditions. This raises important questions on the evolution of the microbiota as hosts diversify. First, how much is microbiota composition conserved over host evolutionary timescales? While the microbiota can be quite labile within and between host species (1–5), more closely related host species often tend to have more similar microbiota, a pattern referred to as *phylosymbiosis* (6, 7). In animals, levels of phylosymbiosis appear to be heterogeneous across tissues (e.g. gut or skin microbiota) and lineages (6, 8–10).

The presence of a phylogenetic signal in microbiota composition across hosts could potentially be used to reconstruct ancestral microbiota composition. Ancestral reconstructions could be particularly useful to detect events during host diversification associated with major shifts in microbiota composition or to verify hypotheses on ancestral diets. A phylogenetic signal in microbiota composition may also inform on potential long-term evolutionary covariations in abundances between microbial taxa. Positive or negative covariations may arise from direct interactions between microbial taxa, such as cross-feeding, trophic relationships, or competition (11–13), or from (anti)correlated responses to variations in the environment (*e.g.* similar or opposite responses to decreased pH). We refer to these covariations as *microbiota integration* by analogy with the often observed *phenotypic integration* between traits in complex phenotypes (14). Such covariations would indicate constraints in the evolution of microbiota composition.

Phylogenetic comparative methods offer a rich toolbox for quantifying phylogenetic signal, reconstructing ancestral states, and detecting integration in multidimensional phenotypes (15). These methods rely on modeling the evolution of a set of phenotypic traits across evolutionarily related species through a multivariate stochastic process, such as the Brownian motion process, running along the species' phylogenetic tree (16, 17). The multivariate Brownian process models the gradual evolution of traits through the accumulation of stochastic changes drawn from a multivariate normal distribution with a variance-covariance matrix that reflects the magnitude of the changes for each trait (the variance terms) and the covariation in the changes between trait pairs (the covariance terms). This process is relevant to represent long-term variations in the abundances of the different microbial taxa that constitute the microbiota, as such variations are an emerging outcome of: (i) the stochastic accumulation of changes in the numerous host traits that can influence the microbiota, including both extrinsic (e.g. biogeography, habitat) and intrinsic (e.g. diet, antimicrobial excretions) traits (6, 12, 18) and (ii) interactions between microbial taxa (11). Indeed, the Brownian motion process has already been used to model variations in microbial abundances over host evolutionary time (19, 20). However, the process is

not directly applicable to compositional data made of relative microbial abundances as
it does not constrain its components to sum to 1, and absolute abundances are
unfortunately typically not provided by mainstream metabarcoding technics used to
characterize microbiota composition. Thus, current phylogenetic comparative methods
cannot directly be used in the context of microbiota evolution without transgressing
several model assumptions (21).
123
Here, we develop an approach to apply the multivariate Brownian motion
process to compositional data. We also include a widely-used tree transformation (22)
that quantifies phylosymbiosis by evaluating how much host phylogeny contributes to
explaining interspecific variation in present-day microbiota composition.
Phylosymbiosis is typically assessed using correlative approaches such as Mantel
tests (6), which are known to suffer from frequent false negatives, while process-based
approaches such as ours tend to be more powerful (21, 23, 24). We apply our new
approach to the gut bacterial microbiota of mammals and birds. The gut microbiota is
key to the functioning of their hosts, contributing to their nutrition, their protection, and
their development (25). Strong phylosymbiosis in gut bacterial microbiota has been
reported for mammals, including primates and rodents (26–28), while it is thought to
be absent for birds, with some exceptions in a few young clades (8, 29, 30). We revisit
this dichotomy here, on the premise that previous analyses may have not been
powerful enough to detect phylosymbiosis in birds (21). We analyse potential drivers
of phylosymbiotic patterns, including geography, diet, and flying ability, we estimate
the ancestral microbiota composition of mammals and birds, and we investigate
patterns of microbiota integration.

Results & Discussion:

We developed a method to infer the dynamics of microbiota composition during host diversification from host-microbiota data (*i.e.* a fixed, bifurcating host phylogeny and microbiota relative abundances for each extant host species) using the multivariate Brownian motion process (Figure 1 and Material & Methods). We assume that all microbial taxa are present in all hosts, potentially in very low (undetectable) abundances and that they were already present in the most recent common ancestor of all host species. These assumptions are met if we consider a taxonomic level in the definition of microbial taxa that is high enough given the host clade, such as bacterial orders in the vertebrate gut microbiota. We assume that, from ancestral values at the root X_0, the log-absolute abundances of the different microbial taxa change on the host phylogeny following a multivariate Brownian motion model with variance-covariance matrix R (Figure 1a). In addition, we account for variation linked to present-day factors by applying the widely-used Pagel's λ transformation to the host phylogeny (Figure 1b). Unlike the traditional case of the multivariate Brownian motion process applied to phenotypic data, where the phenotype is directly measured at present, in the case of the microbiota, relative rather than absolute abundances are measured. To handle this difficulty, we treat total microbial abundances in each host as latent variables, and sample from the joint posterior distribution of these latent variables and our parameters of interest: Pagel's λ, which provides us with an estimate of phylosymbiosis, the R matrix which reflects microbiota integration, and Z_0, which indicates the relative microbial abundances in the ancestral microbiota.

We tested this inference method on data simulated from our model and found that we can accurately estimate the ancestral bacterial relative abundances Z_0 (with a tendency for homogenization) and the variance-covariance matrix R between microbial taxa, provided that the number of host species (n) and bacterial taxa (p) are large enough ($n \geq 50$ and $p \geq 5$, see Supplementary Results 1). Similarly, the level of phylosymbiosis λ is accurately estimated for $p \geq 5$, and its significance is correctly inferred for $n \geq 50$ (see Supplementary Results 1). As expected, this approach provides a more powerful way to detect phylosymbiosis than Mantel tests (Table S1).

173
174

Figure 1: A comparative phylogenetic model for the dynamics of microbiota
composition during host diversification: **(a)** We model fluctuations in the abundances of
microbial taxa along a host phylogeny with a multivariate Brownian motion parametrized by
the ancestral abundances (X_0) and the variance-covariance matrix (R). The variance terms (on
the diagonal) reflect the magnitude of the changes, while the covariance terms reflect positive
or negative covariations in abundances between pairs of microbial taxa. The relative ancestral
abundances (Z_0) and the variance-covariance matrix R are estimated by adjusting the model
to the host-microbiota data (host phylogeny and microbiota relative abundances for each host).
(b) Following the widely-used Pagel's λ transformation, we extend the relative length of the
terminal branches of the host phylogenetic tree by a scaling factor 1-λ. λ is comprised between
0 and 1 and is co-estimated during inference. λ close to 1 indicates that closely related hosts
tend to have similar microbiota due to shared evolutionary history (strong phylosymbiosis),
while λ close to 0 indicates that microbiota composition is determined by present-day
processes with little influence of host evolutionary history (weak or absent phylosymbiosis).
The significance of phylosymbiosis is assessed with permutations.

We applied our model to the gut bacterial microbiota of 215 mammal species and 323 bird species from (8) and found a pervasive signal of phylosymbiosis. We focused on the 14 most abundant bacterial orders, corresponding in abundance to 84% and 82% of the total gut bacterial microbiota of mammals and birds, respectively. We found a markedly higher level of phylosymbiosis in mammals ($\lambda \simeq 0.65$) than in birds ($\lambda \simeq 0.31$; Table S2, Figure S1), consistent with previous literature and our finding that microbiota composition is more species-specific in mammals than in birds (Table S3). Indeed, bird microbiota is generally more sensitive to short-term environmental changes such as anthropogenic perturbations or parasite infections (30). By explicitly modeling the non-phylogenetic component of microbiota composition using a Pagel's λ transformation, we detected a low but significant level of phylosymbiosis in the gut microbiota of birds (Table S2), contrary to previous conclusions (8, 30) that relied on Mantel tests. λ values are higher at the level of bacterial phyla (Table S2; Figure S1), suggesting that microbiota composition is more evolutionarily conserved at higher taxonomic levels. Testing model performance on data simulated directly on the mammal and bird phylogenetic trees, we found a low type-I error rate and a high statistical power, suggesting that the phylosymbiosis we detected in birds is not due to false detection by our method, but rather to a higher power than previously used methods (Table S4). Phylosymbiosis is not linked to an effect of captivity nor the spurious concatenation of different studies either (Supplementary Results 2). Phylosymbiosis is particularly strong in Primates, Passeriformes, and Cetartiodactyla, lower but significant in Columbiformes, Chiroptera, and Carnivora, and non-significant in Rodentia, Charadriiformes, and Anseriformes (Table S2). Non-significant phylosymbiosis in these orders is likely due to an insufficient number of sampled species ($n < 25$, see Supplementary Results 1). It appears that vertebrate orders with mainly herbivorous diets have stronger phylosymbiosis, although this would need to be tested more robustly with a better species coverage (Table S2).

Our results suggest that phylosymbiosis is only partially explained by evolutionary conservatism in flying ability, diet, or biogeography. First, excluding flying mammals (Chiroptera) or non-flying birds did not impact our estimates of phylosymbiosis (Table S2). Second, permutation tests shuffling the microbiota of host species having the same diet, biogeography, flying ability, or combination of these traits resulted in much lower λ values (Figures 2 & S2). In mammals, λ values resulting from such shuffling are still significant (Figure 2), suggesting that the evolutionary conservatism of flying ability, diet, and biogeography contributes to phylosymbiosis without fully explaining it (18). In birds, shuffling often resulted in non-significant λ values (Figure 2), indicating a weak or absent contribution of diet or biogeography in the observed phylosymbiosis. Similarly, the conservatism of these traits is not sufficient to explain the phylosymbiosis measured in some of the larger mammal and bird clades, such as Primates, Cetartiodactyla, and Passeriformes (Figure S3). Thus, we suspect that other evolutionary-conserved physiological, immunological, or ecological traits act

232 as host filters (11, 18) and contribute to phylosymbiosis in the gut microbiota of
233 mammals and birds (9, 31).
234
Our ancestral reconstructions of the microbiota of early mammals and birds
suggest that Proteobacteria and Firmicutes were much more abundant in the ancestral
gut microbiota of birds than mammals (Figures 3, 4, S4 & S5). As it is common in
phylogenetic ancestral reconstruction, the uncertainty is quite high (Figure S6); it is
larger in mammals than in birds because of the long branches that separate marsupials
and eutherians at the origin of all mammals. Comparing the ancestral microbiota
composition of mammals to that of the extant wild mammal species, we found the
highest similarity with invertebrate feeders, such as the insectivorous armadillos
(*Zaedyus pichiy*, Figures 5 & S7). This result is robust to uncertainty in our estimate of
ancestral microbiota composition (Figure S6b) and when including species sampled in
captivity (Figure S8). This is consistent with the current hypothesis, based on the fossil
record and ancestral diet reconstruction, of a generalist insectivorous diet in early
mammals (32, 33). We found the gut microbiota composition of modern birds to be
only weakly structured by diet compared to that of mammals, making the inferred
ancestral microbiota composition of birds less informative in this respect (no strong
clustering in the PCA plots; PermANOVA testing the effect of diet: $R^2 \sim 0.03$, $p < 0.001$
in birds *versus* $R^2 \sim 0.22$, $p < 0.001$ in mammals; Figures 5, S7 & S8; Table S5).

252
253
Figure 2: Phylogenetically-conserved diets, biogeographies, or flying abilities partially
contribute to phylosymbiosis in the gut microbiota of mammals, but not birds. For both
mammals and birds, we compared the estimated level of phylosymbiosis (mean λ value in
orange) to levels of phylosymbiosis (λ values) estimated when shuffling the species that have
the same diet (green boxplot), biogeography (blue boxplot), flying ability (flying or non-flying;
purple boxplot), or combination of the latter traits (in red). For each shuffling strategy, we
performed 100 randomizations. Combining all traits strongly constrains the possible
permutations, which may consequently retain a phylogenetic signal in the shuffling and lead to
high λ values although the traits are actually not strongly contributing to phylosymbiosis.

263

264

265 **Figure 3: Ancestral reconstruction of mammalian gut microbiota**: Phylogenetic tree of the
266 sampled mammal species and associated relative abundances of the 14 most abundant
267 bacterial orders (bar charts on the right). Pie charts at the root and nodes of the tree represent
268 estimated ancestral microbiota compositions (mean of the posterior distribution of Z_0 at the
269 root and generalized least squares estimates at other internal nodes). Compositions are not
270 represented at the most recent nodes for the sake of clarity.

Figure 4: Ancestral reconstruction of avian gut microbiota: Phylogenetic tree of the sampled birds, and associated relative abundances of the 14 most abundant bacterial orders (bar charts on the right). Pie charts at the root and nodes of the tree represent estimated ancestral microbiota compositions (mean of the posterior distribution of Z_0 at the root and generalized least squares estimates at other internal nodes). Compositions are not represented at the most recent nodes for the sake of clarity.

We detected significant changes in microbiota composition in the ancestors of some mammal and bird orders (Figures 3, 4, & S9). In mammals, the largest shift in microbiota composition occurred in the ancestor of Chiroptera, with an increased proportion of Enterobacteriales (Proteobacteria), Mycoplasmatales (Tenericutes), and to a lesser extent Actinomycetales (Actinobacteria), as well as a decreased proportion of Bacteroidales (Bacteroidetes), and in Firmicutes, Clostridiales were replaced by Bacillales and Lactobacillales (Figure 3; Table S6). Other shifts occurred in the ancestor of Carnivora, with an increased proportion of Fusobacteriales (Fusobacteria), and in the ancestors of Primates and Cingulata, with an increased proportion of some Firmicutes orders (*e.g.* Erysipelotrichales; Figures 3 & S9). In addition, Proteobacteria (especially Enterobacteriales and Pseudomonadales) almost disappeared in the ancestral microbiota of Ungulata and Simiiformes (New and Old World monkeys; Table S6). In birds, we found a shift in microbiota composition in the ancestor of Passeriformes, with more Bacillales and Enterobacteriales, and to a lesser extent Pseudomonadales, and a quasi-disappearance of Bacteroidales (Figures 4 & S5; Table S6). The ancestors of Anseriformes and Charadriiforms were characterized by a larger proportion of Bacteroidales, as well as a large proportion of Fusobacteriales, often absent or present in low abundances in other bird gut microbiota. Finally, the relative abundance of Actinomycetales increased in Columbiformes (Table S6). We found similar estimates of ancestral gut microbiota composition when running separate inferences for the different mammal and bird orders (Figure S9). Some of these compositional shifts might be linked to the ecological changes that these lineages experienced, such as the acquisition of flight for bats or carnivorous diets for Carnivora and Charadriiforms (8, 34).

Figure 5: Projection of the estimated ancestral gut microbiota of mammals and birds onto the space of present-day gut microbiota. Top panels: projection of bacterial orders contributing to the two principal components (PC). Colors represent the contribution of the taxa to the principal components. Percentages indicate the explained variance of each PC. Only the 9 most abundant orders are represented for the sake of clarity. Bottom panels: Projection of the extant and ancestral microbiota compositions. Extant microbiota of species sampled in the wild are colored according to the species' diet. Ancestral microbiota compositions of mammals and birds are represented in blue. On each PCA plot, we indicated the three extant species with microbiota compositions closest to the ancestral microbiota composition. The ancestral gut microbiota of mammals is closest to the gut microbiota of present-day invertebrate feeders; the gut microbiota of birds does not strongly reflect diet.

Far from varying as uncorrelated units during the evolutionary history of mammals and birds, we found significant covariances between many microbial taxa, both positive and negative (Figure 6a), suggesting strong constraints in the evolution of the microbiota. These patterns of microbiota integration are strikingly similar in mammals and birds (Figure 6b), indicating that they are conserved over long evolutionary times. Our simulation analyses on the mammal and bird trees suggest that these results are not artefactual, since we recover significant covariances only when we include them in the simulations (Table S7, Supplementary Results 1). Similar covariances were obtained when performing separate inferences on the different mammal and bird orders (Figure S10), which both confirms our results and suggests that the model assumption of a constant variance-covariance matrix across the host phylogenetic tree is reasonable. Combined with the high bacterial variability in time, across individuals, and across host species at low taxonomic levels, these consistent patterns at the level of bacterial orders on large time scales suggest that there is a certain level of functional redundancy among bacteria taxa within orders in the vertebrate gut microbiota.

Based on their covariances, bacterial orders can be clustered into two main subsets within which taxa tend to covary in a concerted way, while taxa from different subsets tend to be anti-correlated (Figure 6). The first subset ("subset 1") is formed in particular by the orders Clostridiales, Bacteroidales, and Fusobacteriales, and the second subset ("subset 2") is mainly composed of the orders Enterobacteriales, Lactobacillales, Pseudomonades, Actinomycetales, and Bacillales. Although some host species have a microbiota composed of an even mixture of these two bacterial subsets, one subset generally prevails. The first subset is dominant in the microbiota of most mammals (excluding Chiroptera), the ancestors of birds, and some extant bird lineages (*e.g.* Anseriformes, Columbiformes, or Accipitriformes); the second subset predominates in the microbiota of Chiroptera and other bird lineages, including Passeriformes (Figure S11). This result suggests the existence of two main "enterotypes" in the vertebrate gut microbiota at the level of bacterial orders, by analogy with the enterotypes observed in human gut microbiota at the bacterial genus level (35, 36), which are conserved over millions of years.

Figure 6: Estimated variances and covariances between the main bacterial taxa tend to be similar in the gut microbiota of mammals and birds. (a) For each variance-covariance matrix between bacterial taxa estimated using our model of host microbiota evolution, we represented negative covariances in red and positive covariances in blue, while variances are represented in shades of green. Non-significant covariances are represented in white. Grey rectangles correspond to subsets of bacterial orders that tend to covary positively. **(b)** Correlation between covariances between the main bacterial taxa estimated in the gut microbiota of mammals or birds. The red line indicates the corresponding linear model, while the grey line corresponds to y=x.

We can only speculate on the processes underlying positive or negative
covariances between bacterial orders: we cannot distinguish from our analyses
whether they indicate direct interactions between bacterial taxa (*e.g.* cross-feeding or
competition) or indirect interactions mediated by similar/opposed microbial responses
to changes in the gut environment. For instance, the frequent and strong negative
covariations observed between the abundant Enterobacteriales (Proteobacteria) and
the major bacterial orders Clostridiales (Firmicutes) and Bacteroidales (Bacteroidetes)
may result from direct competitions (37), host immunological controls over
Proteobacteria (38), and/or be mediated by the oxygen concentration in the gut, as
Proteobacteria are facultative anaerobes, while other phyla are obligate anaerobes
(39). The strongest positive covariations we inferred between Actinomycetales,
Pseudomonadales, and Rhizobiales, which are the most abundant bacterial orders in
plant tissues (40), may reflect a plant-based diet, which would lead to a concomitant
increase of plant-associated bacteria in the gut microbiota of herbivorous vertebrates
(41). Some of the covariations we detected (e.g. the negative covariation between
Lactobacillales and Bacteroidales) have also been observed in human microbiome
data using co-occurrence network analyses (13), suggesting that at least some
covariations between microbial taxa that occur over short timescales within host
species are conserved over macroevolutionary timescales.
378
To test the adequacy of our model to the data, we simulated microbiota under
our model using the parameters estimated on mammal and bird data. We found that
simulated microbiota have compositions similar to those observed in extant mammals
and birds (Figure S12), which indicates that, despite its simple assumptions, our
multivariate Brownian motion model generates realistic gut microbiota (19, 21).
Nevertheless, the gut microbiota composition of mammals and birds appears more
constrained than the sets of compositions we can simulate using multivariate Brownian
motions (Figure S12). This may be linked to constraints that are not accounted for by
our model, such as the potential existence of carrying capacities for some bacterial
orders, or non-constant or non-homogeneous variance-covariance matrices (*e.g.*
effects of host traits such as diet or gut pH on covariation). Extensions of our
multivariate Brownian motion approach could accommodate such constraints, but this
will highly complexify inferences. Meanwhile, our phylogenetic comparative model for
microbiota evolution is general enough to be broadly applicable across other host-
microbiota systems.
394
Besides modeling assumptions, our results may be influenced by the inherent
biases of metabarcoding data. Bacterial relative abundances characterized using
metabarcoding techniques are a distortion of the actual relative abundances (42, 43),
since metabarcoding is sensitive to the number of rRNA copies in the bacterial
genomes, primer biases, and the quality and completeness of the reference database
for taxonomic assignation (at the bacterial order/phylum level in our case). These
issues are unlikely to artefactually generate phylosymbiosis or covariations across
bacterial taxa because we expect such biases to be homogeneous across host

species; nevertheless, they are likely to affect our ancestral reconstructions of
microbiota compositions.
405
Our approach to quantifying phylosymbiosis characterizes microbiota
composition in terms of the relative abundances of higher bacterial taxa (orders or
phyla). This characterization hides variations in the presence/absence of bacterial taxa
at lower taxonomic levels (*e.g.* genus or species). Indeed, distinct mammal or bird
species are known to host different bacterial species (8), and this may not translate
into abundance variations at higher taxonomic levels if the different bacterial species
belong to the same higher taxa. Besides the widely-used Mantel tests, such variations
could be accounted for by stochastic processes modeling the evolution of
presence/absence on host phylogenies (44), although we are not aware that these
approaches have been used to detect bacterial phylosymbiosis. Yet another level of
variation in microbiota composition that can contribute to phylosymbiosis arises
through genetic differentiation below the bacterial species level: if a bacterial species
is vertically transmitted during host diversification, we expect bacterial strains from
closely related host species to be more genetically similar (26, 45, 46). This latter
process can be specifically tested thanks to cophylogenetic methods that consider the
evolution of each microbial species separately (47, 48). The above-mentioned
methods are complementary, as they focus on different levels of variations in
microbiota composition, and on the distinct processes that simultaneously generate
phylosymbiosis (6, 18).
425
Phylosymbiosis is a widespread pattern that has fascinated microbial ecologists
and evolutionary biologists since its discovery, spurring debates on the main processes
underlying the pattern. Drawing upon phylogenetic comparative methods, we have
developed a new approach to studying phylosymbiosis. Our results on simulations and
birds suggest that phylosymbiosis may be even more prevalent than currently
recognized, but sometimes undetected with correlative approaches. We have shown
that conservatisms in diet, biogeography, and flying ability are not enough to explain
phylosymbiosis, calling for an investigation of the role of other host ecological traits, as
well as physiological and immunological traits. One of our most striking results, in the
face of the well-known high variability of the gut microbiota, is its high level of
integration, with conserved covariations between bacterial orders over millions of
years. The same two subsets of bacterial orders tend to covary in a concerted way in
both mammals and birds, leading to the existence of two main "enterotypes" in
vertebrate gut microbiota. Deciphering the host traits and microbial interactions at the
origin of such constraints will help clarify the processes that shape the composition of
the vertebrate gut microbiota.

Material & Methods:

A multivariate Brownian motion model for variations in microbiota composition over host evolutionary time:

We denote by p the total number of microbial taxa detected across the microbiota of the n sampled host species. Standard metabarcoding techniques only measure the relative abundance of each microbial taxon j in each extant host species i, which we denote by $Z_{ij} = X_{ij}/Y_i$, where X_{ij} is the unmeasured absolute abundance of microbial taxon j in host i and $Y_i = \sum_j X_{ij}$ is the unmeasured total microbial abundance in the microbiota of host i. We assume that the logarithms of microbial absolute abundances $\log X_{ij}$ vary along the host phylogenetic tree according to a multivariate Brownian motion starting from the ancestral abundances at the root, denoted by X_{0j} (Figure 1). Indeed, taking the logarithm of the abundances yields values on the real axis that are amenable to be modeled with a Brownian motion, similar to continuous phenotypic traits. This model implies a log-normal distribution of abundances, as is commonly observed in microbial communities (49), and it can easily accommodate undetected microbial taxa in some hosts by assigning them very low unobserved relative abundances. To make the model identifiable, we express the total abundances Y_i relative to the unknown total abundance at the root Y_0, and we only infer $\tilde{Y}_i = Y_i/Y_0$. Each microbial taxon i is characterized by a certain variance and pairs of microbial taxa can affect each other through a covariance term, so that their changes in abundance over time can be positively or negatively correlated. All variance and covariance values are assumed to be constant along the host phylogeny and are summarized by the invertible variance-covariance matrix *R* (Figure 1a).

Model inference:

To infer the model parameters, we sampled from their joint posterior distribution $P(\log Z_0, R, \lambda, \log \tilde{Y}_1, \dots, \log \tilde{Y}_n | Z_{11}, \dots, Z_{ij}, \dots, Z_{np}, C)$ using a No U-turn Hamiltonian Monte Carlo sampler, a computationally efficient Markov Chain Monte Carlo algorithm for continuous variables (Supplementary Methods 1). We implemented it in the probabilistic programming language Stan and we ran and compiled it through the RStan interface (50, 51). Inferences were performed with 4 independent chains and a minimum of 4,000 iterations per chain including a warmup of 2,000 iterations. We checked the convergence of the chains using the Gelman statistics and effective sample sizes (ESS). We extracted the mean posterior value of each parameter and its associated 95% credible interval across posterior samples.

We considered a covariance to be significant if 0 was not included in its 95% credible interval. We could not use the same approach for λ, because it only takes positive values. Furthermore, model selection using Bayes factors led to many false negatives on simulated data (Supplementary Methods 2 & Results 1). Therefore, we

assessed the significance of λ using permutations. We shuffled at random the extant
host species to break the phylogenetic structure and ran again model inference of the
randomized dataset. We performed 100 replications and compared the distribution of
λ values thus obtained to the original λ estimate: if the original λ was greater than at
least 95% of the λ values obtained through permutations, we considered that there
was a significant impact of host evolution on microbiota evolution.
491
492   **Simulations:**
493
We evaluated our approach using simulations. We simulated the evolution of a
microbiota along a host phylogeny using Multivariate Brownian motions for log-
abundances. We simulated phylogenies with n = 20, 50, 100, or 250 extant host
species using a pure birth model (*pbtree* function in the phytools R-package (52)). We
considered microbiota with p = 3, 5, 10, or 15 microbial taxa and uniformly sampled
the logarithms of their ancestral abundances at the root of the host phylogeny between
-4 and 0 before normalizing them so that $\sum_j Z_{0j} = 1$. We generated random positive
definite variance-covariance matrices R following (53) and (54) with eigenvalues of 1/4.
Finally, we applied Pagel's λ transformations with λ = 1, 0.75, 0.5, 0.25, or 0. For each
combination of n, p, and λ values, we performed 100 independent simulations, leading
to a total of 8,000 simulations. We verified that our approach correctly estimates the
parameters λ, Z_0, and R, and detects phylosymbiosis (significant λ) and covariations
(significant R components) when they are simulated. We compared the performances
of our approach for detecting phylosymbiosis to that of Mantel tests (24).
508
We also evaluated our inference approach using data simulated on the
phylogenetic tree of mammals or birds, and using conditions and parameters matching
the empirical data. We performed simulations with 7 taxa (corresponding to the 7
bacterial phyla in the data, see below) and 14 taxa (corresponding to the 14 bacterial
orders in the data). We used values of λ = 1, 0.75, 0.5, 0.25, or 0, and values for the
other model parameters similar to those estimated from the empirical data (Figure
S13). We performed 100 simulations per condition (thus reaching a total of 2,000
simulations).
517
518   **Empirical application:**
519
We downloaded the dataset of (8) that gathered the gut microbiota of 2,677
mammal individuals from >200 species and 1,630 bird individuals from >300 species,
characterized by metabarcoding using the V4 region of the 16S rRNA gene. Only
studies using the standard protocol of the Earth Microbiome Project (55) were included
(see (8) for details), making samples comparable across different studies (42). Song
et al. converted bacterial reads into amplicon sequence variants (ASV), assigned each
ASV taxonomically using the Greengenes database (8, 56), and rarefied ASV tables
at 10,000 reads per sample. We complemented their dataset with the consensus

phylogenetic trees of (57) and (58) for mammals and birds, respectively. We only kept
the species having their microbiota compositions characterized by at least 2 microbiota
samples (Table S2). We checked that gut microbiota from the same host species were
more similar than gut microbiota from different species using PermANOVA (59). Then,
we obtained the microbiota composition of each host species by averaging the
samples per species and extracted the relative abundances of the main bacterial
orders and phyla per host species. We verified that similar results were obtained when
repeating our analyses by randomly sampling one individual per host species (Figure
S14). We only considered the 14 most abundant bacterial orders, *i.e.* those that each
represented more than 1% of the total bacterial abundance, and the 7 most abundant
bacterial phyla (Figure S15). We also repeated all analyses using only the 9 (resp. 5)
most abundant orders (resp. phyla). We did not apply our model at lower taxonomic
levels as the quality of the taxonomic assignation then decreased sharply.
541
Our multivariate Brownian motion model of microbiota does not explicitly
consider losses of bacterial taxa from the microbiota through time. Yet, some bacterial
taxa can be absent or undetected in the gut microbiota of mammals and birds. We
assumed that the absence of a particular taxon came from a very low abundance,
below the detection threshold: we thus arbitrarily set the relative abundances of absent
taxa to 0.001%. Setting the minimal relative abundances of absent taxa to 0.01%
reduced the estimated variance of the rare taxa but did not affect other estimates
(Figure S16).
550
551   We applied the model separately on all mammals and all birds, getting estimates
552 of Pagel's λ, the ancestral microbiota composition Z_0, and the variance-covariance
553 matrix R for each vertebrate class.
554

Effect of host traits on phylosymbiosis:

We gathered data on host species traits from (8) for diet, biogeography, and
flying ability. We assigned a dominant diet to each host species as either "plants",
"fruits", "invertebrates" or "meat" following the EltonTraits database (60). We assigned
a biogeography to each species by picking the biographic realm (Afrotropical,
Antarctic, Australasian, Nearctic, Neotropic, Oriental, or Palearctic) where the highest
number of wild individuals were sampled, or if not available, where the highest number
of captive individuals were sampled (this was the case for 48% of the mammalian
species and 18% of the avian ones). We treated flying ability as binary (yes/no). First,
we assessed the influence of flight on the gut microbiota by performing inferences on
non-flying mammal species only (*i.e.* excluding bats) and on flying bird species only.
Similarly, we investigated the effect of captivity on our inferences by replicating them
using only the gut microbiota of wild or captive individuals. Second, we tested whether
the evolutionary conservatism of diet, biogeography, or flying ability may explain
phylosymbiosis in mammals and birds by performing permutations. We shuffled host
species having the same diet, biogeography, and/or flying ability and re-ran the

inferences on these randomized datasets. For each tested trait, we performed 100 independent randomizations. Finally, we verified that phylosymbiosis did not artefactually arise from the concatenation of the separate studies composing this dataset by randomizing the species that came from the same study.

Comparison between ancestral and present-day microbiota composition:

We compared the estimated ancestral microbiota composition Z_0 of all mammals or birds to that of extant species using principal component analysis (PCA) after applying a centered log-ratio transform to the abundances (61). Given Z_0, we also jointly estimated the ancestral abundances at each node of the host phylogenetic tree using generalized least squares following (54, 62, 63). We additionally performed separate model inference for all orders of mammals (Carnivora, Cetartiodactyla, Chiroptera, Primates, and Rodentia; Table S2) and birds (Anseriformes, Charadriiformes, Columbiformes, and Passeriformes) represented by at least 15 species, and compared the ancestral microbiota composition obtained with separate and joint inferences.

Integration analyses:

We identified the significantly positive or negative covariances between bacterial orders. In addition, to characterize potential subsets of bacterial taxa that tend to vary in a concerted way, we clustered taxa using the *cluster_fast_greedy* function in the R-package igraph (64), based on the estimated variance-covariance matrix *R*, modified to retain information of only positive covariances (negative ones were set to 0).

Model adequacy:

To assess whether our model for the evolution of the gut microbiota of mammals and birds yields realistic microbiota compositions, we simulated the process of microbiota evolution on the mammal or bird phylogenies using the parameters estimated for mammals and birds ($\log Z_0$, R, and λ). Next, we compared the simulated microbiota compositions to the empirical microbiota compositions of the extant mammal or bird species using principal component analysis (PCA). We performed 20 independent simulations for each of our model inferences.

Data availability:

Raw data and processed data from (8) used to perform the empirical applications are available in Qiita (https://qiita.ucsd.edu/study/description/11166).

Our phylogenetic comparative method, referred to as ABDOMEN (A Brownian moDel Of Microbiota EvolutioN), is available on GitHub with a tutorial: https://github.com/BPerezLamarque/ABDOMEN.

Acknowledgments:

The authors acknowledge Julien Clavel, Jonathan Drury, and members of the BioDiv team at IBENS for helpful discussions. They also acknowledge the Hubert Curien Alliance program for funding workshops that initiated the project. GSK acknowledges support from the Academy of Finland (decision 340314) and the Sakari Alhopuro foundation (grant 20210172). This work was performed using HPC resources from GENCI-IDRIS (Grants 2021-A0100312405 and 2022- AD010313735).

References:

1. K. R. Amato, *et al.*, Evolutionary trends in host physiology outweigh dietary niche in structuring primate gut microbiomes. *ISME J.* **13**, 576–587 (2019).
2. L. A. David, *et al.*, Diet rapidly and reproducibly alters the human gut microbiome. *Nature* **505**, 559–563 (2014).
3. R. E. Ley, C. A. Lozupone, M. Hamady, R. Knight, J. I. Gordon, Worlds within worlds: evolution of the vertebrate gut microbiota. *Nat. Rev. Microbiol.* **6**, 776–788 (2008).
4. S. M. Hird, C. Sánchez, B. C. Carstens, R. T. Brumfield, Comparative Gut Microbiota of 59 Neotropical Bird Species. *Front. Microbiol.* **6**, 1403 (2015).
5. S. Hacquard, *et al.*, Microbiota and host nutrition across plant and animal kingdoms. *Cell Host Microbe* **17**, 603–616 (2015).
6. S. J. Lim, S. R. Bordenstein, An introduction to phylosymbiosis. *Proc. R. Soc. B Biol. Sci.* **287**, 20192900 (2020).
7. A. W. Brooks, K. D. Kohl, R. M. Brucker, E. J. van Opstal, S. R. Bordenstein, Phylosymbiosis: relationships and functional effects of microbial communities across host evolutionary history. *PLOS Biol.* **14**, e2000225 (2016).
8. S. J. Song, *et al.*, Comparative analyses of vertebrate gut microbiomes reveal convergence between birds and bats. *MBio* **11**, 1–14 (2020).
9. F. Mazel, *et al.*, Is host filtering the main driver of phylosymbiosis across the Tree of Life? *mSystems* **3**, 1–15 (2018).
10. B. Perez-Lamarque, H. Krehenwinkel, R. G. Gillespie, H. Morlon, Limited evidence for microbial transmission in the phylosymbiosis between Hawaiian spiders and their microbiota. *mSystems* **7**, e01104-21 (2022).
11. K. R. Foster, J. Schluter, K. Z. Coyte, S. Rakoff-Nahoum, The evolution of the host microbiome as an ecosystem on a leash. *Nature* **548**, 43–51 (2017).
12. K. D. Kohl, Ecological and evolutionary mechanisms underlying patterns of phylosymbiosis in host-associated microbial communities. *Philos. Trans. R.*

654  *Soc. B Biol. Sci.* **375**, 20190251 (2020).

655 13. K. Faust, *et al.*, Microbial co-occurrence relationships in the human
656  microbiome. *PLoS Comput. Biol.* **8**, e1002606 (2012).

657 14. M. Pigliucci, Phenotypic integration: Studying the ecology and evolution of
658  complex phenotypes. *Ecol. Lett.* **6**, 265–272 (2003).

659 15. J. Clavel, G. Escarguel, G. Merceron, mvMORPH: An R package for fitting
660  multivariate evolutionary models to morphometric data. *Methods Ecol. Evol.* **6**,
661  1311–1319 (2015).

662 16. L. J. Revell, L. J. Harmon, D. C. Collar, Phylogenetic signal, evolutionary
663  process, and rate. *Syst. Biol.* **57**, 591–601 (2008).

664 17. L. J. Harmon, *Phylogenetic Comparative Methods* (2017).

665 18. N. A. Moran, H. Ochman, T. J. Hammer, Evolutionary and ecological
666  consequences of gut microbial communities. *Annu. Rev. Ecol. Evol. Syst.* **50**,
667  451–475 (2019).

668 19. M. del M. Labrador, J. Doña, D. Serrano, R. Jovani, Quantitative interspecific
669  approach to the stylosphere: Patterns of bacteria and fungi abundance on
670  passerine bird feathers. *Microb. Ecol.* **81**, 1088–1097 (2021).

671 20. D. C. Capunitan, O. Johnson, R. S. Terrill, S. M. Hird, Evolutionary signal in the
672  gut microbiomes of 74 bird species from Equatorial Guinea. *Mol. Ecol.* **29**,
673  829–847 (2020).

674 21. S. M. Hird, Microbiomes, community ecology, and the comparative method.
675  *mSystems* **4**, 1–5 (2019).

676 22. M. Pagel, Inferring the historical patterns of biological evolution. *Nature* **401**,
677  877–884 (1999).

678 23. L. J. Harmon, R. E. Glor, Poor statistical performance of the Mantel test in
679  phylogenetic comparative analyses. *Evolution (N. Y).* **64**, 2173–2178 (2010).

680 24. B. Perez-Lamarque, *et al.*, Do closely related species interact with similar
681  partners? Testing for phylogenetic signal in bipartite interaction networks. *Peer
682  Community J.* **2**, e59 (2022).

683 25. M. McFall-Ngai, *et al.*, Animals in a bacterial world, a new imperative for the life
684  sciences. *Proc. Natl. Acad. Sci.* **110**, 3229–3236 (2013).

685 26. M. Groussin, *et al.*, Unraveling the processes shaping mammalian gut
686  microbiomes over evolutionary time. *Nat. Commun.* **8**, 14319 (2017).

687 27. H. Ochman, *et al.*, Evolutionary relationships of wild hominids recapitulated by
688  gut microbial communities. *PLoS Biol.* **8**, 3–10 (2010).

689 28. K. D. Kohl, M. D. Dearing, S. R. Bordenstein, Microbial communities exhibit
690  host species distinguishability and phylosymbiosis along the length of the
691  gastrointestinal tract. *Mol. Ecol.* **27**, 1874–1883 (2018).

692 29. B. K. Trevelline, J. Sosa, B. K. Hartup, K. D. Kohl, A bird's-eye view of
693  phylosymbiosis: weak signatures of phylosymbiosis among all 15 species of
694  cranes. *Proc. R. Soc. B Biol. Sci.* **287**, 20192988 (2020).

695 30. K. H. Bodawatta, S. M. Hird, K. Grond, M. Poulsen, K. A. Jønsson, Avian gut
696  microbiomes taking flight. *Trends Microbiol.* **30**, 268–280 (2022).

697 31. J. K. Goodrich, E. R. Davenport, J. L. Waters, A. G. Clark, R. E. Ley, Cross-
698  species comparisons of host genetic associations with the microbiome.
699  *Science (80-.).* **352**, 532–535 (2016).

700 32. D. M. Grossnickle, S. M. Smith, G. P. Wilson, Untangling the multiple
701  ecological radiations of early mammals. *Trends Ecol. Evol.* **34**, 936–949
702  (2019).

703 33. P. G. Gill, *et al.*, Dietary specializations and diversity in feeding ecology of the

earliest stem mammals. *Nature* **512**, 303–305 (2014).

34.   A. H. Nishida, H. Ochman, Rates of gut microbiome divergence in mammals.
Mol. Ecol. **27**, 1884–1897 (2018).

35.   M. Arumugam, *et al.*, Enterotypes of the human gut microbiome. *Nature* **473**,
174–180 (2011).

36.   A. H. Nishida, H. Ochman, A great-ape view of the gut microbiome. *Nat. Rev.*
Genet. **20**, 195–206 (2019).

37.   N. G. Shealy, W. Yoo, M. X. Byndloss, Colonization resistance: metabolic
warfare as a strategy against pathogenic Enterobacteriaceae. *Curr. Opin.*
Microbiol. **64**, 82–90 (2021).

38.   J. Mirpuri, *et al.*, Proteobacteria-specific IgA regulates maturation of the
intestinal microbiota. *Gut Microbes* **5**, 28–39 (2013).

39.   N. R. Shin, T. W. Whon, J. W. Bae, Proteobacteria: Microbial signature of
dysbiosis in gut microbiota. *Trends Biotechnol.* **33**, 496–503 (2015).

40.   M. R. Wagner, *et al.*, Host genotype and age shape the leaf and root
microbiomes of a wild perennial plant. *Nat. Commun.* **7**, 12151 (2016).

41.   H. Dion-Phénix, *et al.*, Bacterial microbiota similarity between predators and
prey in a blue tit trophic network. *ISME J.* **15**, 1098–1107 (2021).

42.   R. Knight, *et al.*, Best practices for analysing microbiomes. *Nat. Rev. Microbiol.*
16, 410–422 (2018).

43.   A. Lavrinienko, T. Jernfors, J. J. Koskimäki, A. M. Pirttilä, P. C. Watts, Does
intraspecific variation in rDNA copy number affect analysis of microbial
communities? *Trends Microbiol.* **29**, 19–27 (2021).

44.   M. P. Braga, M. J. Landis, S. Nylin, N. Janz, F. Ronquist, Bayesian inference of
ancestral host-parasite interactions under a phylogenetic model of host
repertoire evolution. *Syst. Biol.* **69**, 1149–1162 (2020).

45.   B. Perez-Lamarque, H. Morlon, Characterizing symbiont inheritance during
host–microbiota evolution: Application to the great apes gut microbiota. *Mol.*
Ecol. Resour. **19**, 1659–1671 (2019).

46.   J. G. Sanders, *et al.*, Stability and phylogenetic correlation in gut microbiota:
lessons from ants and apes. *Mol. Ecol.* **23**, 1268–1283 (2014).

47.   W. Dismukes, M. P. Braga, D. H. Hembry, T. A. Heath, M. J. Landis,
Cophylogenetic methods to untangle the evolutionary history of ecological
interactions. *Annu. Rev. Ecol. Evol. Syst.* **53**, 1–24 (2022).

48.   B. Perez-Lamarque, H. Morlon, Comparing different computational approaches
for detecting long-term vertical transmission in host-associated microbiota. *Mol.*
Ecol. (2022) https:/doi.org/10.1111/mec.16681.

49.   C. Quince, T. P. Curtis, W. T. Sloan, The rational exploration of microbial
diversity. *ISME J.* **2**, 997–1006 (2008).

50.   Stan Development Team, RStan: the R interface to Stan (2022).

51.   R Core Team, R: A language and environment for statistical computing. (2022).

52.   L. J. Revell, phytools: An R package for phylogenetic comparative biology (and
other things). *Methods Ecol. Evol.* **3**, 217–223 (2012).

53.   J. C. Uyeda, D. S. Caetano, M. W. Pennell, Comparative analysis of principal
components can be misleading. *Syst. Biol.* **64**, 677–689 (2015).

54.   J. Clavel, L. Aristide, H. Morlon, A penalized likelihood framework for high-
dimensional phylogenetic comparative methods and an application to New-
World monkeys brain evolution. *Syst. Biol.* **68**, 93–116 (2019).

55.   L. R. Thompson, *et al.*, A communal catalogue reveals Earth's multiscale
microbial diversity. *Nature* **551**, 457–463 (2017).

56. T. Z. DeSantis, *et al.*, Greengenes, a chimera-checked 16S rRNA gene
database and workbench compatible with ARB. *Appl. Environ. Microbiol.* **72**,
5069–5072 (2006).
57. N. S. Upham, J. A. Esselstyn, W. Jetz, Inferring the mammal tree: Species-
level sets of phylogenies for questions in ecology, evolution, and conservation.
PLoS Biol. **17**, e3000494 (2019).
58. W. Jetz, G. H. Thomas, J. B. Joy, K. Hartmann, A. O. Mooers, The global
diversity of birds in space and time. *Nature* **491**, 444–448 (2012).
59. J. Oksanen, *et al.*, vegan: Community Ecology Package, R package version
2.4-0. *R Packag. version 2.2-1* (2016).
60. H. Wilman, *et al.*, EltonTraits 1.0: Species-level foraging attributes of the
world's birds and mammals. *Ecology* **95**, 2027–2027 (2014).
61. J. Aitchison, Principal component analysis of compositional data. *Biometrika*
70, 57 (1983).
62. E. P. Martins, T. F. Hansen, Phylogenies and the comparative method: A
general approach to incorporating phylogenetic information into the analysis of
interspecific data. *Am. Nat.* **149**, 646–667 (1997).
63. C. W. Cunningham, K. E. Omland, T. H. Oakley, Reconstructing ancestral
character states: a critical reappraisal. *Trends Ecol. Evol.* **13**, 361–366 (1998).
64. G. Csardi, T. Nepusz, The igraph software package for complex network
research. *InterJournal Complex Syst.* **Complex Sy**, 1695 (2006).
775