Inhibition of Type I PRMTs Reforms Muscle Stem Cell Identity Enhancing their Therapeutic Capacity

Claudia Dominici[1,4], Oscar Villareal[1], Junio Dort[6], Emilie Heckel[6], Yu Chang Wang[7], Ioannis Ragoussis[7], Jean-Sebastien Joyal[6], Nicolas A. Dumont[6], Stéphane Richard[1,2,3,4,5]*

[1]Segal Cancer Center, Lady Davis Institute for Medical Research and [2]Gerald Bronfman Department of Oncology and Departments of [3]Medicine, [4]Human Genetics and [5]Biochemistry, McGill University, Montréal, Québec, Canada H3T 1E2

[6]CHU Sainte-Justine Research Center, Université de Montréal, Montréal, Québec, Canada H3T 1C5

[7]Genome Quebec Innovation Centre, Montreal, Quebec, Canada.

Running title: Type I PRMT repression improves muscle stem cell self-renewal and muscle regeneration

Keywords: muscle stem cell, PRMT1, muscle regeneration, small molecule inhibitor

*Corresponding author: stephane.richard@mcgill.ca

Dominici et al.,

Abstract

In skeletal muscle, muscle stem cells (MuSC) are the main cells responsible for regeneration upon injury. In diseased skeletal muscle, it would be therapeutically advantageous to replace defective MuSCs, or rejuvenate them with drugs to enhance their self-renewal and ensure long-term regenerative potential. One limitation of the replacement approach has been the inability to efficiently expand MuSCs *ex vivo,* while maintaining their stemness and engraftment abilities. Herein, we show that inhibition of type I protein arginine methyltransferases (PRMTs) with MS023 increases the proliferative capacity of *ex vivo* cultured MuSCs. Single cell RNA sequencing (scRNAseq) of *ex vivo* cultured MuSCs revealed the emergence of novel subpopulations in MS023-treated cells which are defined by elevated Pax7 expression and markers of MuSC quiescence, both features of enhanced self-renewal. Furthermore, the scRNAseq identified MS023-specific subpopulations to be metabolically reprogrammed with upregulated glycolysis and oxidative phosphorylation (OxPhos). Transplantation of MuSCs treated with MS023 had a better ability to repopulate the MuSC niche and contributed more robustly to muscle regeneration following injury. Interestingly, the preclinical mouse model of Duchenne muscular dystrophy had increased bilateral grip strength 10 days after a single intraperitoneal dose of MS023. These findings suggest that type I PRMT inhibition metabolically reprograms MuSCs resulting in improved self-renewal and muscle regeneration fitness.

Dominici et al.,

Introduction

Muscle stem cells (MuSCs) reside on the periphery of muscle fibers in skeletal muscle and are activated following injury to drive muscle regeneration (Lepper *et al*, 2011; Relaix & Zammit, 2012). Major challenges arise when attempting to manipulate MuSCs *ex vivo*. Purified MuSCs spontaneously differentiate upon *ex vivo* expansion (Fu *et al*, 2015a; Sacco *et al*, 2008), limiting the potential of CRISPR-based applications and stem cell engraftment therapy. It is known that *ex vivo* cultured MuSCs are vastly inferior at engrafting and contributing to muscle regeneration compared to freshly isolated MuSCs, indicating the importance of retaining stemness for MuSC engraftment and muscle regeneration (Montarras *et al*, 2005; Sakai *et al*, 2017). Cell-intrinsic defects in patients with Duchenne muscular dystrophy (DMD) prevent MuSCs from maintaining a proper balance between stem cell self-renewal and differentiation, ultimately leading to depletion of regeneration-competent MuSCs (Bentzinger *et al*, 2014; Dumont *et al*, 2015; Wang *et al*, 2013).

MuSCs have metabolic flexibility that allows them to adapt to demands presented by their changing environments (Relaix *et al*, 2021). Indeed, metabolic signatures are strong contributors to the heterogeneity observed in isolated MuSCs (Cho & Doles, 2017). Several findings have indicated that the metabolic state of MuSCs is inextricably tied to their function and fate. Quiescent MuSCs predominantly rely on fatty acid oxidation (FAO) and oxidative phosphorylation (OxPhos) (Machado *et al*, 2017; Ryall *et al*, 2015). While glycolysis remains low in quiescent cells, it is upregulated as MuSCs begin to proliferate following activation, representing a metabolic shift in activated MuSCs (L'Honore *et al*, 2018; Ryall *et al.*, 2015). The dynamic regulation of metabolic state during the transition from MuSC quiescence to proliferation therefore presents a unique avenue to regulate MuSC regenerative capacity.

Dominici et al.,

There is growing evidence to suggest a role for the post-translational modification arginine methylation in muscle cellular metabolism (Blanc & Richard, 2017; Saber & Rudnicki, 2022; vanLieshout & Ljubicic, 2019). Arginine methylation is carried out by 9 protein arginine methyltransferases (PRMTs) (Xu & Richard, 2021). PRMTs are classified according to their catalytic activity: type I enzymes (ex: PRMT1, PRMT2, PRMT3, CARM1, PRMT6 and PRMT8) catalyze arginine asymmetrical dimethylation (ADMA), type II enzymes (ex: PRMT5, PRMT9) catalyze arginine symmetrical dimethylation (SDMA), and the unique type III enzyme PRMT7 catalyzes arginine monomethylation (MMA) (Bedford & Clarke, 2009; Jain & Clarke, 2019). PRMTs methylate many cellular proteins including histones and are known epigenetic regulators (Xu & Richard, 2021).

PRMTs are known regulators of MuSCs. PRMT1-null mouse MuSCs are highly proliferative and can be expanded in culture, but are unable to terminate differentiation, leading to a severe muscle regeneration defect *in vivo* following muscle injury (Blanc *et al*, 2017). CARM1 is a well-known transcriptional coactivator for nuclear receptors and p300/CPB (Bedford & Clarke, 2009) and it is a mediator of skeletal muscle plasticity (Saber & Rudnicki, 2022; vanLieshout & Ljubicic, 2019). CARM1 methylates the myogenic transcription factor, PAX7, to complex with histone methyltransferase complex to allow the expression of myogenic transcription factor Myf5 during asymmetric stem cell divisions (Kawabe *et al*, 2012). CARM1 regulates MuSCs metabolism by methylating AMP-activated protein kinase (AMPK) in MuSCs (Stouth *et al*, 2020). PRMT5 regulates MuSC proliferation by epigenetically downregulating expression of the cell cycle inhibitor p21 and deletion of PRMT5 in MuSCs blocks their expansion *ex vivo* and causes defects in muscle regeneration *in vivo* (Zhang *et al*, 2015). PRMT7 functions in the myogenic process and is required for MuSC self-renewal and muscle

Dominici et al.,

regeneration, as deletion of PRMT7 in MuSCs leads to persistent p21 expression (Blanc *et al*, 2016).

Given the importance in maintaining MuSC stemness and the enhanced MuSC proliferation observed with PRMT1 deficiency (Blanc *et al.*, 2017), we aimed to determine whether the reversible inhibition of type I PRMTs using the MS023 inhibitor (Eram *et al*, 2016) would affect cellular metabolism and fulfil the requirements of generating a stem-like MuSC in culture with subsequent regenerative capabilities. Small molecule inhibitors of epigenetic regulators are a promising avenue to pursue in the treatment of muscle wasting diseases. Indeed, an inhibitor of Setd7-mediated lysine methylation (Judson *et al*, 2018), was shown to have beneficial effects on self-renewal and regeneration.

Herein, we show that MS023, an inhibitor of type I PRMTs, fulfils the requirements of a reversible inhibitor which maintains the stemness of cultured MuSCs, while allowing for engraftment and regenerative capabilities. Although type I PRMTs have well-known indications as epigenetic modulators, we demonstrate that targeting type I PRMTs alters the metabolic profile of dividing MuSCs to favour self-renewal and improve regeneration.

Results

Type I PRMT inhibitor MS023 enhances self-renewal and *in vitro* expansion of MuSCs

We isolated single muscle fibers from 6 to 8 week old C57BL/6 wild type mice, cultured them in vitro with MS023 or vehicle (DMSO) for 48 h, and stained them for the myogenic transcription factor Pax7 (Seale *et al*, 2000; Soleimani *et al*, 2012), and the proliferation marker ki67. The number of Pax7/ki67 double-positive cells increased significantly in the presence of MS023 (Eram *et al.*, 2016) for 48 h (Figure 1A, 1B). We observed a significant increase in the number of clusters with 3 or more cells with MS023 treatment (Figure 1B), as observed for PRMT1-deficient MuSCs (Blanc *et al.*, 2017). Treatment with a specific CARM1 inhibitor TP-064 (Nakayama *et al*, 2018) over the course of 6 days did not increase MuSC cell number, ruling out that MS023 increased proliferation is mediated by CARM1 (Supplemental Figure S1A). These data show that type I PRMT inhibition causes *in vitro* MuSC expansion and is likely by the inhibition of PRMT1. To determine the activation state of these fiber-associated MuSCs, we examined fibers after 48 h in culture for the expression of Pax7 and the myogenic regulator transcription factor, MyoD (Rudnicki *et al*, 1993; Troy *et al*, 2012). Interestingly, MS023 treatment restricted the proportion of committed $Pax7^{neg}/MyoD^{pos}$ cells, while expanding the proportion of activated and cycling cells ($Pax7^{pos}/MyoD^{pos}$, Figure 1C, D).

MuSCs spontaneously differentiate into non-cycling myoblasts and lose their stemness *in vitro* (Montarras *et al.*, 2005; Ryall *et al.*, 2015; Sacco *et al.*, 2008). The enhanced proliferation of MuSCs observed with type I PRMT inhibition on muscle fibers prompted us to examine whether similar effects could be achieved with FACS-purified MuSCs cultured *in vitro*. MuSCs incubated with MS023 for 48 h had reduced ADMA containing proteins, as visualized by immunoblotting (Supplemental Figure S1B), and had increased proliferative capacity of MuSCs,

Dominici et al.,

as determined by ki67 immunofluorescence (Supplemental Figure S1C) and by FACS analysis of

ki67-PE intracellularly-stained MuSCs (Supplemental Figure S1D). These data are consistent

with what was observed in muscle fiber-associated MuSCs (Figure 1A-C). Importantly,

immunofluorescence analysis of myogenic markers in cultured MuSCs revealed that MS023

treatment prevented initiation of differentiation, as indicated by the significantly reduced sub-

population of committed Pax7^{neg}/MyoD^{pos} cells compared to control DMSO culture (13% versus

31%, respectively) (Figure 1E, F). Additionally, there was an increase in the proportion of cells

that have retained their stemness by delaying expression of MyoD (Pax7^{pos}/MyoD^{neg}) in the

MS023-treated samples compared to DMSO (40% versus 9%, Figure 1E, F). Similar proportions

were observed with fiber-associated MuSCs (34% versus 14%, Figure 1C, D). Furthermore, we

observed that the impediment of precocious differentiation observed by treatment with MS023

led to enhanced long-term culture capabilities of MuSCs up to the 14th passage (Supplemental

Figure S1E). Altogether, these findings indicate that treatment with MS023 maintained cultured

MuSCs in a stem-like state.

Transcriptionally distinct MuSC sub-populations emerge under type I PRMT inhibition

To identify subpopulations of MuSCs generated by MS023, we performed single-cell RNA

sequencing (scRNAseq). MuSCs were purified from 8 week-old wild type C57BL/6 mice in

biological duplicates immediately after 1) isolation (termed d0; sample day 0), and 2) culture in

growth medium for 4 days with 0.033% DMSO as control (sample d4) or with 1μM MS023

(sample d4MS023), and 3) grown in growth media for 6 days with 0.033% DMSO removed at

day 4 (sample d6), or 6 days in culture with 1μM MS023 removed at day 4 (sample

d6MS023rd4). Cells were collected at these time points and processed on the 10x Genomics

Dominici et al.,

Chromium platform for scRNAseq. Data filtering and analysis was performed using the Seurat package for scRNAseq analysis (Butler *et al*, 2018). After filtering out low-quality cells, approximately 4,000 cells (average, 3,887 ± 44 cells) from each sample were retained for subsequent analysis (Supplemental Table S2). A Pearson correlation of >0.98 (Supplemental Figure S2A) was obtained between replicates and we chose to include a replicate for each time point for analysis.

Dimensionality reduction was performed using uniform manifold approximation and projection (UMAP) on pooled cells from all samples, resulting in 10 distinct subpopulations of myogenic cells (labelled 0-9, Supplemental Figure S2B). We then visualized the distribution of individual samples within the UMAP embedding (Figure 2C, Supplemental Figure S2C). To determine the identity of the 10 subpopulations, we examined the expression of *Pax7* and the bHLH transcription factors *Myod1, Myf5 and Myog* (Figure 2B).

Two populations of *Pax7* and *Myf5* expressing cells were observed which we called quiescence 1 and 2 (Q1 and Q2), while the two populations that expressed both *Myod1* and *Myog* were called differentiated myoblasts 1 and 2 (DM1 and DM2, Figure 2A). The remaining *Myod1* populations we named myoblast clusters 1 to 5 (M1-M5), and the last one C1 for committed 1 (Figure 2A). The freshly isolated MuSCs of day 0 (d0) segregated into Q1 and Q2 (Supplemental Figure S2D, Supplemental Table S3), which in addition to *Pax7* and *Myf5*, expressed markers of MuSC quiescence (*Fos, Egr1, Jun*), and early activation markers (*MyoD1, Mt1, Mt2*, Supplemental Table S1). Our Q1 and Q2 populations are similar to quiescence and early activation populations reported recently from freshly isolated (De Micheli *et al*, 2020; Dell'Orso *et al*, 2019; van Velthoven *et al*, 2017). The two *Myog* expressing clusters expressed other markers of early differentiation (*Acta2*; DM1), and late differentiation (*Tnnt3, Myl1, Acta1*;

Dominici et al.,

DM2) (Supplemental Table S1). An enrichment for cell cycle and proliferation markers was found throughout the remaining 5 undifferentiated myoblast clusters (*Cdk1*, *Ccnd1*, *Mki67*, *Top2a, Birc5,* and *Cenpa*; M1-M5) (Supplemental Table S1).

The control DMSO day 4 (d4) sample was distributed among 3 subpopulations of myoblasts (M1, M2, and M4), while two additional days in culture day 6 (d6) contained M1, M2, and M4, but also had differentiated clusters DM1 and DM2 (Figure 2C, 2D; Supplemental Figure S2E, Supplemental Table S3). In comparison, day 4 with MS023 (d4MS023) was distributed uniquely among clusters M3 and M5. Day 6 MS023 removed at day 4 (d6MS023rd4) had clusters of day 4 MS023 treatment, as well as DM1 and DM2 (Figure 2C, 2D Supplemental Figure S2E, Supplemental Table S3). Notably, d6MS023rd4 had nearly twice as many cells in DM1 (20.8%) as d6 (11.5%) (Figure 2D), indicating that MS023 inhibition was reversible and cells enter the differentiation program more efficiently. In addition, all samples had in common a small subpopulation of cells (<200 cells) in cluster C1 (Supplemental Table S3).

Analysis of the top 100 enriched genes in each cluster revealed distinct transcriptional profiles (Supplemental Table S1). For example, the d4MS023-occupied cluster M5 was enriched for cell cycle markers (*Cdk1*, *Cdk4*, *Ccnd3*) and other markers of proliferation (*Mki67*, *Birc5*, *Top2a*) (Supplemental Table S1). The other d4MS023-occupied cluster M3 retained some markers of proliferation (*Ccnd1*), however also expressed the cell cycle inhibitor *Cdkn2a* and the structural muscle gene *Myl6*, in addition to a strong enrichment for ribosomal genes (over 20% of the top100 markers belonged to the ribosomal protein family, a feature that is unique to cluster M3). Therefore, clusters M3 and M5 are unique to MS023-treated MuSCs and represent robust proliferation clusters with extensive ribosomal protein expression.

Dominici et al.,

The top 100 enriched genes in each sample were also investigated (Supplemental Table S2, Heatmap in Supplemental Figure S2J). Interestingly, d4MS023 genes were strongly enriched for the reactome pathways of respiratory electron transport, the citric acid cycle, and formation of ATP. In contrast, d4 was enriched for the reactome pathways of RNA metabolism, translation, and mRNA splicing (Supplemental Figure S2H, S2I).

The d4 cluster M2 shared many of the same gene enrichments as cluster M5 (*Cdk1*, *Mki67*, *Birc5*, *Top2a)* (Supplemental Table S1). Conversely, the other d4 cluster M4 down-regulated these genes while maintaining expression of replication-dependent transcripts such as *Ube2c* and *Dut*, indicating increased heterogeneity in cycling cells compared to d4MS023 (Supplemental Table S1). The final d4 cluster M1 resembled the transitional d4MS023 cluster M3 in that it expressed a mix of cell cycle (*Ccnd1*) and differentiation (*Myl6*) markers, however lacked the enrichment of ribosomal genes observed in M3 (Supplemental Table S1).

It was important to note is the stark difference in the pathway enrichment signatures of the transitional clusters M1 (d4) and M3 (d4MS023). The genes which were uniquely enriched in cluster M3 were strongly associated with protein synthesis (e.g., *Rps9, Eef1g, Uba52*) (Supplemental Table S1, Supplemental Figure S2H, S2I). In contrast, the enriched genes unique to cluster M1 were associated with cellular response to toxic substance and cellular detoxification (*Anxa1, Txn1, Hmox1, Prdx5*) (Supplemental Figure S2H, S2I).

We further investigated the distribution of expression levels for *Pax7, Myf5, Myod1,* and *Myog* across clusters and samples (Supplemental Figure S2F, S2G). We observed a unique expression pattern for *Pax7* in d4MS023 and d6MS023rd4 samples (Supplemental Figure S2G). In d4 MuSCs, cells were divided into two populations, Pax7-negative (expression level 0) and Pax7-positive (standard distribution around expression level 1). In d6 myoblasts, Pax7-positive

cells maintain the same level of *Pax7* expression, but a higher proportion of cells became Pax7-negative, consistent with canonical *Pax7* expression during MuSC differentiation (Yin *et al*, 2013). Meanwhile, with MS023 treatment, three subpopulations emerged, a Pax7-negative (expression level 0) and two Pax7-positive (standard distribution around expression level 1) populations. The Pax7-positive populations in d4MS023 and d4MS023rd4 had higher *Pax7* expression levels than their untreated counterparts and were more comparable with day 0 (quiescent) MuSC *Pax7* expression levels (Supplemental Figure S2G).

Together, these findings suggest that the MuSC subpopulations generated by MS023 treatment possess several unique features. Day 4 MS023 cells had proliferating signatures, less heterogeneity and were localized mainly within one cluster (cluster M5), unlike day 4 DMSO proliferating cells which were divided into two clusters (clusters M2 and M4), each with different expression profiles of cell cycle regulators. Cluster M5 had a striking enrichment for genes involved in cellular metabolism that was unique to M5. Furthermore, the day 4 MS023 transitional cluster M3 had a strong enrichment for genes involved in mRNA translation which was not observed in the day 4 DMSO transitional cluster M1. Analysis of expression level patterns in d4MS023 cells indicated a subpopulation of MuSCs with elevated *Pax7* expression compared to day 4 DMSO control cells. Therefore, through repression of type I PRMTs with MS023, we have reprogramed MuSCs to acquire a unique and previously uncharacterized identity.

Dominici et al.,

Trajectory analysis reveals temporally distinct expression patterns in MuSCs treated with MS023 with unique enrichment for energy metabolism

Using the Monocle v2.16.0 R package (Trapnell *et al*, 2014), we conducted trajectory analysis on 3 groups of samples to obtain hierarchical links between clusters: (1) all samples; (2) d0, d4, d6; (3) d0, d4MS023, d6MS023rd4. Monocle analysis placed each cell from the 9 clusters (Q1/Q2, M1-5, and DM1/2) onto a pseudotime axis based on differential gene expression patterns. The trajectory analysis of all samples pooled produced a pseudotime axis with a single branchpoint and 3 cell states which we labelled Q for quiescence, M for myoblast, and DM for differentiating myoblast (Figure 3A). Cells from the Q1 and Q2 clusters were found at the beginning of pseudotime, cells from the M1-5 clusters populated the area around the branchpoint, and cells from DM1/2 mainly populated the upward-oriented branch, following an overall trajectory of Q1/2→M1-5→DM1/2 (Figure S3A).

Day 0 cells populated the beginning of pseudotime, with d4 and d4MS023 cells appearing shortly after. d6 and d6MS023rd4 cells had a similar distribution as d4 and d4MS023, with the exception of having increased density around the terminus of the upward-oriented branch (Supplemental Figure S3A). Interestingly, d4MS023 and d6MS023rd4 cells emerge later along the pseudotime axis than d4 and d6 cells and appeared closer to the branchpoint (Supplemental Figure S3A).

We examined the genes which were significantly differentially regulated across the branchpoint of the pooled sample trajectory to confirm whether this split is representative of the molecular events which determine whether MuSCs will commit to differentiation or continue to proliferate. As expected, genes involved in the myogenic process were highly upregulated along the branch that diverges upwards along the trajectory (eg. *Tnnt1, Tnnt2, Acta1, Myog*), while

genes involved in cell cycle and proliferation were upregulated along the branch that diverges downwards along the trajectory (eg. *Mki67, Cdca8, Top2a, Birc5*) (Supplemental Figure S3B, S3C).

Trajectory analysis was also performed on a subset of samples (d0→d4→d6; DMSO trajectory and d0→d4MS023→d4MS023rd4; MS023 trajectory) to identify treatment-specific alterations in the molecular events which drive fate-determining decisions (Figure 3B, 3C, 3E, 3F). Both the DMSO and MS023 trajectories retained a single branch point indicating that the overall lineage patterns were conserved.

Closer inspection of genes which were significantly differentially regulated across the branchpoint in the DMSO and MS023 trajectories revealed a unique signature in the MS023-trajectory. The list of genes which changed significantly across the branchpoint for both trajectories (7094 for the DMSO trajectory, 7616 for the MS023 trajectory) were filtered for genes which were unique to each condition. The DMSO trajectory had 1,517 unique genes, while the MS023 trajectory had 2037 unique genes (Supplemental Figure S3D). Interestingly, the MS023 trajectory-unique genes revealed an enrichment for oxidative phosphorylation and mitochondrial ATP synthesis (Figure 3F, 3G), consistent with cluster marker analysis for MS023-specific clusters (Supplemental Figure S2H, S2I: *electron transport, ATP synthesis*). Expression of select genes involved in the mitochondrial electron transport chain (*Ndua2, Ndufa4, Uqcr10, Cox4I1, Cox5b, Cox8a, Atp5c1, AtpIf1, Atp5j2*) were further examined for expression level within the scRNAseq UMAP plot, and it was confirmed that they were elevated in MS023-specific clusters (Figure 4A). Moreover, RT-qPCR analysis of a subset of these genes confirmed they were upregulated in d4MS023 MuSCs compared to d4 MuSCs (Supplemental Figure S4A). Therefore, these pathways both define the unique identity of MS023-treated

Dominici et al.,

MuSCs, and serve as driving forces in their fate-determination. MS023-treated proliferating

MuSCs retain oxidative phosphorylation and have elevated glycolysis

The transition of MuSC from quiescence to proliferation is accompanied by a sharp

reduction in oxidative metabolism and a shift towards glycolysis (L'Honore *et al.*, 2018; Ryall *et*

al., 2015). Therefore, the retention of a transcriptional signature indicating high levels of

mitochondrial oxidative phosphorylation in MS023-treated cells was unexpected. We sought

validate these findings in *ex vivo* cultured MuSCs by performing metabolic analysis with

Seahorse Extracellular Flux Analyzer (Agilent). We isolated MuSCs from C57BL/6 wild type

mice and seeded them onto matrigel-coated 96-well plates and treated with 1μM MS023 or

DMSO for 48 hours (n= 3 mice per treatment condition). Oxygen consumption rate (OCR) was

first measured to characterize mitochondrial bioenergetics (Figure 4B). As predicted, MS023-

treated MuSCs displayed increased OCR measurements at nearly all timepoints (Figure 4B, *p*=

0.000001 - 0.02), with significantly elevated basal respiration (*p*= 0.006) and maximal

respiration (*p*= 0.04), and a trend of increased ATP production (approaching statistical

significance, *p*=0.1). To confirm an increase in mitochondrial biogenesis, d4 and d4MS034

MuSCs were stained with MitoTracker CMXRos (ThermoFisher Scientific). The average

intensity of the MitoTracker signal was significantly increased in d4MS023 cells (Figure 4C, 4D,

p< 0.0001)

We next determined whether markers of glycolysis were also differentially regulated in

MS023-treated MuSCs. Indeed, several of the marker genes for MS023-specific clusters were

components of glycolysis which displayed increased expression compared to clusters of DMSO-

treated cells (*Hk1, Eno1, Pfkm, Gapdh, Pkm, Slc2a1*; Supplemental Figure S4B). We next

performed extracellular acidification rate (ECAR) analysis as an indicator of glycolysis again

Dominici et al.,

using the Seahorse Extracellular Flux Analyzer. MS023-treated cells were strikingly more glycolytic than DMSO-treated cells (Supplemental Figure S4C). Glycolysis, glycolytic capacity, and glycolytic reserve were all significantly increased ($p= 0.02$, $p= 0.004$ and $p= 0.001$, respectively) (Supplemental Figure S4C). As further indicators of glycolysis, the growth media supernatant of d4 and d4MS023 cultured cells was analyzed using a Nova Biomedical Bioprofile 400 analyzer (Cambridge Scientific). As expected, d4MS023 cells had significantly higher levels of glucose uptake (measured as grams/liter per 10,000 cells, $p= 0.004$). Additionally, the glycolytic byproduct lactate was present in higher concentrations d4MS023 media and was undetectable in the d4 media (Supplemental Figure S4D). Collectively, these findings indicate that MS023-treated MuSCs utilize both OxPhos and glycolysis at elevated levels compared to untreated control cells to boost cellular metabolism, suggesting a possible mechanism through which enhanced proliferation is supported.

***In vitro* expanded MS023-treated MuSCs retain the potential to become mature myoblasts**

We next wanted to validate that removal of MS023 created conditions which are permissive to terminal differentiation. Primary MuSCs were cultured under four conditions before switching to differentiation media: (1) for four days with DMSO, (2) washout for additional two days without DMSO (DMSOwo), (3) for four days with MS023, and (4) washout for additional two days without MS023 (MS023wo). Cells treated for four days were switched to differentiation media with the continued presence of DMSO or MS023. Washout cells were switched to differentiation media without DMSO or MS023. Expectedly, cells differentiated in the presence of MS023 had an impaired ability to form multinucleated myotubes and had a poor fusion index of 19%

Dominici et al.,

compared to the 77% fusion index observed with DMSO treated MuSCs (Figure 5A, 5B).

Interestingly, MS023wo cells had completely restored their ability to terminate differentiation,

with their fusion index of 76% being comparable to that of DMSOwo cells (78%) (Figure 5C,

5D). Together, these experiments define MS023 as a reversible inhibitor of type I PRMT activity

which enhances MuSC proliferation and self-renewal while still allowing for full differentiation

capabilities upon removal of the compound.

Type I PRMT inhibitor MS023 enhances MuSC engraftment

We next sought to confirm that MS023 washout conditions create a pool of MuSCs that are

regeneration-competent upon engraftment into an injured muscle *in vivo*. Primary MuSCs were

isolated from a β-actin-GFP reporter mouse (C57BL/6-Tg(CAG-EGFP)10sb/J, Jackson

Laboratory 003291, FACS gating strategy in Supplemental Figure S5B, S5C), and treated with

MS023 or DMSO *in vitro* for 6 days to allow for expansion. Following culture, 15,000 GFP$^+$

myoblasts from each group were injected into the tibialis anterior (TA) muscle of wild type mice.

One day after stem cell injection, the TA was injured with cardiotoxin (CTX). After 3 weeks,

mice were sacrificed and GFP$^+$ differentiated muscle fibers were quantified following cross-

sectional staining of the injured TA muscle (n=3 mice per condition) (Supplemental Figure

S5A). Interestingly, the mice which received MS023-treated MuSCs had more GFP$^+$ mature

muscle fibers than the mice which received DMSO-treated MuSCs (88 fibers/section ± 32 for

MS023 versus 31 fibers/section ± 14 for DMSO, *p*= 0.02) (Figure 5E, 5F). These data indicate

that MS023-treated MuSCs were able to regain their ability to differentiate *in vivo* following

Type I PRMT de-repression and contribute more robustly to muscle regeneration following

injury. We next wanted to determine whether MS023-treated engrafted cells were able to

Dominici et al.,

repopulate the niche. Therefore, we repeated the engraftment experiment with MS023 and

DMSO-treated GFP$^+$ MuSCs, and this time re-isolated MuSCs three weeks after CTX injection.

Strikingly, MS023-treated MuSC engraftment yielded a 6.8-fold increase in the number of GFP$^+$

following muscle regeneration (p=0.004) (Figure 5G, 5H).

MS023 injection increases muscle strength in *mdx* mice

Patients with DMD lack functional dystrophin, a structural protein required to connect mature

myofibers to the extracellular matrix. The resulting muscle tissue is vulnerable to injury and is

caught in constant cycles of muscle degeneration and regeneration (Serrano *et al*, 2011). The

Dmd^{mdx} mice harbor a spontaneous mutation in the gene encoding dystrophin and are therefore

commonly used as a model for DMD (Bulfield *et al*, 1984; Ryder-Cook *et al*, 1988).

To determine whether MS023 could provide any therapeutic benefits in a dystrophic

context, we delivered 80 mg/kg of MS023 or vehicle via intraperitoneal injection to *mdx* mice

once a day for 3 days. Tail pieces were collected 48 h after the final injection and whole tissue

lysate was analyzed for ADMA- and SDMA-containing proteins using immunoblotting. We

observed reduced ADMA proteins and a subsequent increase in SDMA proteins in MS023-

treated mice compared to vehicle-treated mice (Supplemental Figure S6A). At 10 days following

the final injection, 2-paw and 4-paw grip strength measurements were taken. Additionally,

endurance was tested by placing mice on a wire grid and inverting them so that the amount of

time spent hanging could be recorded (Figure 6A for schematic). We observed a 50% increase in

2-paw grip strength (p= 0.0113) and a 45% increase (p= 0.0089) in 4-paw grip strength in the

MS023-treated mice versus those that received vehicle at the 10-day time point (Figure 6B). The

hanging test to measure endurance showed a positive trend towards increased endurance at the

Dominici et al.,

10-day time point in MS023 versus vehicle-treated mice, however this was not statistically

significant ($p= 0.1$) (Supplemental Figure S6B). A final measurement was performed at the 32-

day endpoint, during which mice were placed under terminal anesthesia and the extensor

digitorum longus (EDL) hindlimb muscle was isolated, incubated ex vivo, and used to generate a

force/frequency curve. Interestingly, this experiment revealed that MS023-treated mice

responded with a ~30% increase in force compared to their vehicle-treated counterparts ($p=$

0.00727) (Figure 6C). Moreover, we observed a reduction in the variation of the minimum Feret

diameter measurements of fibers from TA cross sections of MS023-treated mice ($p= 0.04$)

(Figure 6D, 6E). These findings indicate that transient repression of PRMT1 with MS023 can

offer a therapeutic avenue to explore for enhancing MuSC proliferation while still maintaining

the ability to differentiate to mature muscle fibers, thus providing improved muscle strength.

Dominici et al.,

Discussion

In the present manuscript, we show that inhibition of type I PRMTs promoted the *in vitro* proliferation of Pax7-positive MuSCs, as visualized by ki67 staining. scRNAseq analysis of MS023 treated MuSCs identified new transitional clusters. MS023-specific subpopulations harboured unique transcriptional signatures of stemness and energy metabolism. The MS023 treated MuSCs exhibited elevation of both glycolysis and OxPhos. Engrafted MuSCs treated with MS023 were better at regenerating injured muscle and at repopulating the niche. A single treatment of the *mdx* mouse model of DMD with MS023 resulted in enhanced grip strength and force generation. Therefore, repression of type I PRMTs reforms the metabolic state of cultured MuSCs to enhance their capacity for self-renewal *ex vivo* and maintain their ability to robustly participate in muscle regeneration. These finding demonstrate that type I PRMT inhibitors may serve as viable treatment options for skeletal muscle diseases.

The ability to tailor energy metabolism to the specific demands of cell proliferation and differentiation is known as metabolic plasticity, and is required for maintaining tissue homeostasis (Folmes *et al*, 2012). The transition from quiescence to proliferation is regulated by a shift from OxPhos to glycolysis (Pala *et al*, 2018; Ryall *et al.*, 2015). During MuSC quiescence, high levels of NAD+ are generated by oxidative metabolism which in turn activates Sirtuin 1 (SIRT1) to deacetylate and repress expression of differentiation genes required for the myogenic program (Ryall *et al.*, 2015). The shift from OxPhos to glycolysis upon exit from quiescence results in reduced NAD+ levels and subsequent activation of differentiation genes. Furthermore, it has been shown that entry to the process of terminal differentiation is mediated by the activation of pyruvate dehydrogenase (PDH), marking a switch from glycolysis to OxPhos (Hori *et al*, 2019). Another key player in metabolic plasticity in MuSCs is AMP kinase, a major

Dominici et al.,

energy sensor in the cell, regulating energy balance and metabolic state (Hardie *et al*, 2016; Herzig & Shaw, 2018). Null alleles of AMPK have contributed to the understanding of its role in MuSCs (Fu *et al*, 2015b; Theret *et al*, 2017). Activated AMPK stimulates autophagy, mitophagy, and mitochondrial biogenesis, thus maintaining mitochondrial homeostasis (Herzig & Shaw, 2018). Our data shows that in inhibition of type I PRMTs in MuSCs stimulates expression of autophagic markers (*Atg5, Lamp1, Lamp2, Sqstm1*) downstream targets of AMPK, however it remains to be determined whether AMPK is involved in MS023 treated cells.

Treatment of MuSCs with MS023 resulted in metabolic reprogramming of MuSCs, supporting a role for PRMT1 as a metabolic coactivator. PRMT1 has been shown to interact with AMPK in skeletal muscle during the early stages of muscle atrophy (Stouth *et al*, 2018). Moreover, CARM1 directly methylates AMPK to regulate its activity during muscle response to denervation (Stouth *et al.*, 2020). PRMT1 has also been shown to play a signaling role to regulate thermogenesis. PRMT1 acts through the transcriptional co-activator PGC-1α in human and mouse adipocytes to regulate thermogenic fat activation, and PRMT1-deficient mice were unable to induce the thermogenic program following cold exposure (Qiao *et al*, 2019). Methylation of PGC-1α by PRMT1 was also shown to stimulate mitochondrial biogenesis and may regulate nuclear-encoded mitochondrial genes(Teyssier *et al*, 2005). Additionally, PRMT1-mediated methylation positively regulates the insulin receptor (IR)-phosphatidylinositol 3-kinase (PI3-K) pathway which is required for glucose transport in skeletal muscle (Iwasaki & Yada, 2007). Overexpression of PRMT1 in the liver under hypoxic conditions causes hypermethylation of FoxO1 and increased translocation to the nucleus, resulting in reduced glucose uptake (Bayen *et al*, 2018). Therefore, these findings indicate that PRMT1 methylates key regulators of cellular metabolism and highlight a crosstalk between metabolic effectors and type I PRMTs.

Dominici et al.,

scRNAseq analysis of MS023 treated MuSCs identified new transitional clusters which harboured unique transcriptional signatures of stemness and energy metabolism. DMSO clusters M1/M2/M4 and MS023 clusters M3/M5 were both enriched for proliferation markers (*Cdk1, Mki67*), and M3/M5 also uniquely harboured elevated metabolic genes including components of the ETC (*Atp5k, Cox5a, Ndufa2*) and components of glycolysis (*Eno1, Gapdh*). The simultaneous elevation of both glycolytic and OxPhos components in MS023 subpopulations likely contributes to the enhanced proliferation phenotype. MuSC subpopulations have been characterized at the single-cell level during muscle regeneration, revealing regulatory mechanisms that guide the transition from quiescence to activation, proliferation, and differentiation. A general progressive elevation of metabolic enzyme expression has been noted from low levels in MuSCs isolated from uninjured muscle, to slightly elevated levels in MuSCs isolated from acutely injured muscle (60h after injury), to the highest levels in cultured proliferating MuSCs (Dell'Orso *et al.*, 2019). A more in-depth analysis of MuSC subpopulations throughout muscle regeneration investigated additional time points after injury (2, 5, and 7 days) and generated a hierarchical continuum model of MuSCs throughout the regeneration process, with a focus on ligand-receptor cell communication networks (De Micheli *et al.*, 2020). Additionally, analysis of young and aged MuSCs revealed that, while they have similar transcriptional signatures, aged MuSCs display delayed activation state dynamics when exiting quiescence, thus preventing them from entering the differentiation program as efficiently as young MuSCs (Kimmel *et al*, 2020). It has been previously reported that the quiescent MuSCs captured by scRNAseq exist in two groups, one 'quiescent' and one 'early activation', wherein the early activation group highly expresses Fos and Jun while down-regulating *Hox* genes (van den Brink *et al*, 2017). Our Q1 and Q2 clusters are in alignment with these two states (Q1 -

quiescence; Q2 - early activation), and bare transcriptional similarities to our MS023-specific clusters M3 and M5, such as upregulated *Fos* and *Jun*, as well as high levels of *Pax7*. Notably, capture of rare transient MuSC states remains difficult due to poor sampling or limited cell number. To overcome these issues, one study evaluated 365,000 single cells/nuclei from over 100 combined datasets, allowing for the characterization of short-lived MuSC subpopulations that emerge throughout the myogenic process (McKellar *et al*, 2021). We show that MS023 treatment generates unique subpopulations of self-renewing MuSCs that have not been previously identified, suggesting that transient cell states may be also amplified through treatment with epigenetic inhibitors, thus facilitating a more in-depth understanding of their function and contributions to muscle regeneration.

PRMTs are known as epigenetic regulators by methylating histones directly and by serving as coactivators/repressors (Xu & Richard, 2021). We now show that type I PRMT inhibition induces metabolic reprogramming favouring proliferation and self-renewal, however, the mechanism whether epigenetic, regulating of mRNA splicing or a signaling cascade remains to be defined. It has been shown that inhibition of the epigenetic modulator *Setd7* enhances the proliferation of cultured MuSCs, while retaining their stemness by preventing transport of β-catenin into the nucleus, thereby failing to activate the differentiation program (Judson *et al.*, 2018). Taken together, these findings suggest that inhibiting methyltransferases can affect MuSC fate.

We report heightened engraftment capabilities of MS023-treated MuSCs accompanied by high Pax7 expression levels. It has been shown that freshly isolated MuSCs retain high Pax7 expression levels and are superior at engrafting compared to cultured MuSCs (Montarras *et al.*, 2005; Sacco *et al.*, 2008). Generating stable Pax7 expression in myogenic precursor cells with

lentiviral delivery of transgenes is a strategy for enhancing engraftment potential following *ex vivo* expansion of MuSCs (Kim *et al*, 2021), although this approach may be hampered by safety concerns. Therefore, we have identified an additional strategy for increasing Pax7 expression and improving engraftment efficiency and self-renewal of *ex vivo* expanded MuSCs through inhibition of type I PRMTs. Furthermore, our findings show that injection of MS023 in the dystrophic mouse model *mdx* led to enhanced muscle strength with effects lasting up to 30 days, suggesting that the transient increase in MuSC proliferation and self-renewal afforded by MS023 treatment has lasting effects that could benefit patients with muscle wasting disease.

In sum, we report that inhibition of type I PRMTs elevates cellular metabolism in MuSCs, stimulates mitochondrial biogenesis and increases OxPhos, generating a more stem-like MuSC with enhanced self-renewal and engraftment capabilities.

Dominici et al.,

Materials and Methods

Mice

C57BL/6J (Jackson Laboratory 000664) were the wild type mice used for MuSC FACS
purification for scRNAseq, muscle fiber isolation, and for all other *in vitro* experiments.
β-Actin GFP mice (C57BL/6-Tg(CAG-EGFP)10sb/J, Jackson Laboratory 003291) were used to
obtain GFP[Pos] MuSCs for further implantation experiments. Mice lacking dystrophin expression
(C57BL/10ScSn-*Dmd[mdx]*/J, Jackson Laboratory 001801) were used for intraperitoneal MS023
injection and muscle physiology assessment. The experiments with the *Dmd[mdx]* mice were
performed at Sainte-Justine and were approved by the CHU Sainte-Justine Research Ethics
Committee and performed in compliance with the Comité Institutionnel des Bonnes Pratiques
Animales en Recherche (CIBPAR; approval number 2020-2668) in accordance with the
Canadian Council on Animal Care guidelines. All other mouse husbandry and experiments were
conducted in accordance with the Institutional Animal Care and Use Committee (IACUC) of
McGill University. All animal procedures conducted at McGill were approved by the Animal
Welfare Committee of McGill University (protocol #3506).

Primary MuSC isolation

Skeletal muscle tissue was isolated from the abdominal and diaphragm muscles of wild type
mice and muscle stem cells were isolated as previously described using fluorescence activated
cell sorting (FACS) (Pasut *et al*, 2012). Briefly, dissected muscles were minced with dissection
scissors and digested with collagenase/dispase solution (2.4U/ml collagenase D, 2.4U/ml
Dispase II in Ham's F10 media) at 37°C for 1h. Digested tissue was triturated and filtered
through a 40 μM cell strainer. Cells were pelleted for 18 min at 1800 rpm and resuspended in 2%

Dominici et al.,

BSA/PBS. Cells were stained for 15 min at room temperature with ITGA7-Alexa647 (R&D systems) for positive selection, and PE-CD45 (Invitrogen), PE-CD11b (Invitrogen, PE-CD31 (BD Pharmigen), and PE-ScaI (BD Pharmigen) for negative selection. Hoescht was used to gate the living cells. Cells were washed once with 2% BSA/PBS and pelleted prior to final resuspension and one last filter through a 40 μM cell strainer. ITGA7+/CD45-/CDCD11b-/Sca1-/Hoechst+ cells were sorted into full myoblast growth media using the FACSAriaIII cell sorter (BD Biosciences).

Myoblast growth and differentiation culture

Purified MuSCs were seeded onto collagen-coated plates and expanded in growth media (Ham's F10 media with 20% FBS, 2.5 ng/mL human recombinant bFGF, and 1% Penicillin/Streptomycin) at 37°C and 5% CO_2. Media was replenished every two days. To differentiate myoblasts into myotubes, myoblasts were grown to 90% confluency in growth media, washed twice with 1X PBS, and switched to differentiation media (DMEM, 1% FBS, and 1% Penicillin/Streptomycin). For inhibitor treatment, cells were incubated with 0.033% DMSO or 1 μM MS023 (Sigma) in 0.033% DMSO and media was replenished every second day for the duration of the indicated treatment time.

Western blot

Proteins from total cell or tissue lysate (50 mM HEPES [pH 7.4], 150 mM NaCl, and 1% Triton X-100) were resolved on SDS 8-15% polyacrylamide gels and transferred onto nitrocellulose membranes using the Trans-Blot turbo transfer system (BioRad). Membranes were blotted with

the primary antibodies against ADMA (Millipore), SDMA (Millipore), and β-Actin (Sigma) overnight at 4°C. Following 3 washes in TBST, membranes were incubated with HRP-conjugated secondary antibodies (Sigma) for 45 min and visualised on X-ray films with Western Lightning Plus ECL (Perkin Elmer).

Cultured myoblast immunofluorescence

Myoblasts expanded in growth media were seeded onto collagen-coated glass coverslips (VWR) in 6-well plates. Coverslips were transferred to a 12-well plate containing 4% paraformaldehyde (PFA) and cells were fixed for 15 min at room temperature. Cells were then permeabilized with 0.2% Triton X-100, 0.125M glycine in PBS for 12 min at room temperature (RT). Incubation with blocking buffer (2% BSA, 5% horse serum and 0.1% Triton X-100) proceeded for 1 h at RT. Primary antibodies were diluted in blocking buffer to detect Pax7 (Developmental Studies Hybridoma Bank, 1:100), MyoD (Santa Cruz Biotechnology, 1:200), or ki67 (abcam, 1:1000). After 16h incubation at 4°C, cells were washed 3 times with 1X PBS for 10 min. Secondary antibody (AlexaFluor anti-mouse or anti-rabbit 488nm or 568nm) was used at a dilution of 1:400 in blocking buffer for 45 min in the dark at RT. Cells were washed 3 times for 10 min with 1X PBS. Finally, the cover slips were transferred to a microscope slide and mounted with ProLong Gold Antifade Mountant with DAPI (ThermoFisher Scientific). For MitoTracker staining, primary MuSCs were seeded onto 8-well Lumox chamber slides (Sarstedt) and cultured for the indicated time points. Cells were then incubated for 30 min with a final concentration of 100 nM MitoTracker Red CMXRos (ThermoFisher Scientific) at 37°C and 5% CO_2. Cells were fixed with ice-cold 100% methanol at -20°C for 15 min and washed 3 times for 10 min with 1X PBS prior to mounting with ProLong Gold Antifade Mountant with DAPI. All cells were visualized

Dominici et al.,

on a Zeiss Axio Imager M1 microscope (Carl Zeiss, Thornwood NY), and resulting images were analyzed using Zeiss' ZEN Digital imaging suite software.

Muscle fiber isolation and culture

Wild type mice were sacrificed and their extensor digitorum longus (EDL) muscles were dissected using standard dissection techniques. Isolated muscles were incubated with 0.4% collagenase (Sigma) in DMEM for 30 min at 37°C and 5% CO_2. Whole muscle was then triturated with a plastic disposable Bohr pipette to dissociate individual fibers from the whole muscle as described previously(Gallot *et al*, 2016). To mimic activating conditions, fibers were cultured in fiber growth media (DMEM plus 20% FBS, 1% chick embryo extract, 2.5 ng/mL bFGF, 1% penicillin/streptomycin) at 37°C and 5% CO_2. For quiescent satellite cell analysis, fibers were fixed immediately following dissociation using 4% PFA prepared fresh in 1X PBS.

Muscle fiber immunofluorescence

Cultured muscle fibers from wild type mice were fixed in 4% paraformaldehyde PFA and washed twice with 1X PBS. Fibers were then permeabilized with 0.2% Triton X-100, 0.125M glycine in PBS for 15 min at RT. Blocking followed for one hour at room temperature with 2% BSA, 5% horse serum and 0.1% Triton X-100. Primary antibodies were diluted in blocking buffer to detect Pax7 (Developmental Studies Hybridoma Bank, 1:100), MyoD (Santa Cruz Biotechnology, 1:200), ki67 (abcam, 1:1000). After 16 h incubation at 4°C, fibers were washed 3 times with 1X PBS for 10 min. Secondary antibody (AlexaFluor anti-mouse or anti-rabbit 488nm or 568nm) was used at a dilution of of 1:400 in blocking buffer for 45 min in the dark at RT. Fibers were washed 3 times for 10 min with 1X PBS. Finally, the fibers were transferred to a

Dominici et al.,

microscope slide outlined using an ImmEdge hydrophic barrier pen and mounted with ProLong

Gold Antifade Mountant with DAPI (ThermoFisher Scientific). Fiber-associated satellite cells

were then visualized on a Zeiss Axio Imager M1 microscope (Carl Zeiss, Thornwood NY), and

resulting images were analyzed using Zeiss' ZEN Digital imaging suite software.

scRNAseq sample preparation and computational analysis

Each biological replicate for the scRNAseq corresponded to one wild type C57BL/6 mouse. The

purified MuSCs were cultured in Ham's F10 (Gibco) with 20% FBS (HyClone), 2.5 ng/mL

human recombinant bFGF (Gibco), 1% Penicillin/Streptomycin (Wisent Inc.) with 0.033%

DMSO or 1 μM MS023 in 0.033% DMSO. The medium was changed on day 2 and day 4 of

culturing. The purified MuSCs from each replicate were stained for viability with calcein-AM

and ethidium-homodimer1 (P/ N L3224 Thermo Fisher Scientific). scRNA libraries were

generated using the GemCode Single- Cell Instrument (10x Genomics, Pleasanton, CA, USA)

and Single Cell 3' Library & Gel Bead Kit v2 and Chip Kit (P/N 120236 P/N 120237 10x

Genomics). The sequencing ready libraries were purified with SPRIselect, quality controlled for

sized distribution and yield (LabChip GX Perkin Elmer), and quantified using qPCR (KAPA

Biosystems Library Quantification Kit for Illumina platforms P/N KK4824) as previously

described(Couturier *et al*, 2020). Libraries were subsequently shipped and sequenced using

Illumina NovaSeq6000 at IGM Genomics Center, UCSD, San Diego, CA. Cell barcodes and

UMI (unique molecular identifiers) barcodes were demultiplexed and paired-end reads were first

aligned to the mouse genome (mm10) using the Cell Ranger software v3.1.0 (10X Genomics,

https://support.10xgenomics.com/single-cell-gene-expression/software/pipelines/latest/what-is-

cell-ranger). Pre-processing was then carried out with the Seurat v3.2.0 R package (Butler et al.,

Dominici et al.,

2018). Genes detected in less than 3 cells as well as cells containing less than 200 genes detected were removed. Cells were further filtered out for each sample based on the distribution of genes detected as well as the percentage of mitochondrial counts to balance the number of cells per sample to ~4000, and the raw count matrices of all samples were merged. Read counts for each cell were then normalized by the cell total, multiplied by 10000 and natural-log transformed. The expression of the 2000 genes with highest cell-to-cell variation was standardized and the heterogeneity associated with mitochondrial contamination was regressed out. Principal component analysis was performed on the scaled data and the top 10 principal components were used to construct a K-nearest neighbor cell graph. Clustering of cells was carried out through the Louvain algorithm with the granularity parameter set to 0.4 and visualized with the Uniform Manifold Approximation and Projection (UMAP) (Becht et al., 2019) dimensional reduction technique using the Scanpy v1.5.2 python module (Wolf et al., 2018). Cluster biomarkers were identified using the Wilcoxon Rank Sum test. Cell trajectories across pseudotime were analyzed by the Monocle v2.16.0 R package (Trapnell et al., 2014). Cell progress was defined by differentially expressed genes based on the clusters identified by Seurat. The dimensionality of the data was reduced through the Discriminative Dimensionality Reduction with Trees (DDRTree) algorithm to two dimensions and the cells were ordered along the trajectory according to pseudotime. Genes with branch-dependent expression were identified through the branched expression analysis modeling (BEAM) test. RNA velocity analysis was performed by the scVelo v0.2.2 python module (Bergen et al., 2020). Spliced and unspliced mRNAs were first distinguished through the velocyto v0.17.17 python module (La Manno et al., 2018). Velocities representing the direction and speed of cell motion were then computed and projected onto the UMAP embedding.

Dominici et al.,

Oxygen consumption rate and extracellular acidification measurements

Oxygen consumption rates (OCR) and extracellular acidification rate (ECAR) were measured using a Seahorse XFe96 Flux Analyzer and analyzed with Wave 2.6.0 software (Agilent Technologies). FACS sorted muscle stem cells were seeded in a matrigel-coated 96-well plate and cultured with MS023 or DMSO for 48 h. Cells were incubated in MitoStress test assay media (DMEM 5030 media, 10mM glucose, 1mM sodium pyruvate, 2mM glutamine, PH7.4) or GlycoStress test assay media (DMEM 5030 media, 2mM glutamine, pH 7.4) 1 h prior to OCR and ECAR measurements, respectively. Oligomycin (3mM, Sigma-Aldrich), carbonyl cyanide-p-trifluoromethoxy-phenylhydrazone (FCCP; 1mM, Abcam) and rotenone-antimycin A (RAA; 2.5mM, Sigma-Aldrich) were added to cells to measure OCR parameters while glucose (10mM, Sigma-Aldrich), Oligomycin (1mM, Sigma-Aldrich) and 2-DG (100mM) were used to measure ECAR parameters. Following the assay, cells were stained with crystal violet and nuclei were counted in each well for normalization.

MuSC engraftment and muscle histology

Primary MuSCs were FACS-purified from donor GFP mice and cultured with MS023 or DMSO for 5 days. Cells were then trypsinized, resuspended in 1X PBS, and pelleted for 10 min at 1,800 RPM. Cell pellets were resuspended in 1ml of 1X PBS and counted using a haemocytometer. 15,000 cells from each condition were then injected into the TA muscle of wild type mice. Recipient mice were also injected with 50 μl of 10 μM cardiotoxin (CTX) 24 h prior to MuSC engraftment to induce muscle regeneration. Recipient mice were sacrificed 3 weeks after CTX injection, and the TA muscles were dissected and fixed in 2% PFA at 4°C for 16 h. Fixed TA

Dominici et al.,

muscles were then embedded in OCT and frozen prior to cryosectioning onto a glass microscope slide. Resulting tissue sections were permeabilized for 12 min with 0.2% Triton X-100, 0.125M glycine in PBS at RT. Blocking followed with M.O.M blocking reagent (Vector Laboratories) for 1 h at RT. Primary antibodies were diluted in blocking reagent to detect Pax7 (Developmental Studies Hybridoma Bank, 1:10) and laminin (Sigma, 1:50). After 16h incubation at 4°C, sections were washed 3 times with 1X PBS for 10 min. Secondary antibody (AlexaFluor anti-mouse or anti-rabbit 488nm or 568nm) was used at a dilution of 1:400 in blocking reagent for 45 min in the dark at RT. Sections were then washed 3 times for 10 min with 1X PBS. Tissue sections were mounted with ProLong Gold Antifade Mountant with DAPI (ThermoFisher Scientific) and covered with a coverslip. Visualization was performed on a Zeiss Axio Imager M1 microscope (Carl Zeiss, Thornwood NY), and resulting images were analyzed using Zeiss' ZEN Digital imaging suite software.

MS023 injection in mdx mice

12-week old *mdx* mice received an intraperitoneal injection of 80 mg/kg MS023 (Sigma) dissolved in 50 µl N-methyl-2-pyrrolidone, 200 µl Captisol, 200 µl polyethylene glycol 400 and 550 µl PBS once per day for three consecutive days. As control, 12-week old mice received vehicle (50 µl N-methyl-2-pyrrolidone, 200 µl Captisol, 200 µl polyethylene glycol 400 in 550 µl PBS) alone. Tail pieces were collected 48h after the last injection and lysed for western blot analysis with anti-ADMA and anti-SDMA and β-actin antibodies. Grip strength and hanging test measurements were taken at 10 days after the final injection. 29 days after the final injection, mice were anesthetized and sacrificed for endpoint *ex vivo* force measurements.

Dominici et al.,

Grip strength measurements and hanging test

Grip strength was measured for the front two paws and for all four limbs using a Bioseb Grip

Strength Test instrument. Each mouse was placed on the rod (2 paws) or mesh (4 paws) and

pulled from the tail three times with 1 min rest between trials. The highest measurement was

retained for subsequent analysis. To assess endurance, mice were placed on a plastic mesh taped

to a hollow cylinder. The cylinder was slowly inverted over soft bedding to allow mice to grip

the mesh, and the amount of time each mouse remined suspended before falling onto the bedding

was recorded 3 times with 2 min rest between trials. The longest time measurement was retained

for analysis.

Ex vivo force measurements of mdx mice

Mice were anesthetized with pentobarbital sodium (50 mg/kg). Proximal and distal tendons of

the EDL were attached with silk suture (3.0) and the muscle was carefully isolated and placed in

a buffered physiological solution (Krebs-Ringer supplemented with glucose and bubbling

carbogen gas) maintained at 25°C (Dort *et al*, 2021). The distal tendon of the EDL muscle was

attached to the electrode, and the proximal tendon was attached to the force lever arm (300C-LR

dual-mode lever; Aurora Scientific, Canada). Muscle length was adjusted to find the optimal

muscle length (L_0) at which it generates the maximum isometric twitch tension. The EDL muscle

was allowed to equilibrate in the contractile bath for 10 min equilibration prior the contractile

measurement. Muscle was stimulated at different frequencies (10, 25, 50, 80, 100, 150 Hz), with

2 min rest between each stimulation, to obtain a force-frequency curve. Thereafter, muscle length

and weight were measured. Specific muscle force (N/cm^2) was determined using the following

Dominici et al.,

formula: (force (N) x fiber length (0.44 x L_0 for the EDL muscle) x muscle density (1.06 g/cm^3))/muscle mass (g).

DECLARATION OF INTERESTS

The authors declare no competing interests.

ACKNOWLEDGEMENTS

The authors would like to thank Chris Young and Mathew Duguay at the LDI Flow Cytometry Facility for their assistance with FACS experiments. We thank Drs. Vahab Soleimani, Vladimer Ljubicic and Etienne Audet-Walsh for critically reading the manuscript and helpful suggestions and discussions. The research was funded by Canadian Institute of Health Research (CIHR) FDN-154303 to S.R. C.D. held a studentship from the fonds de la recherche en santé du Québec (FRQS).

AUTHOR CONTRIBUTIONS

C.D. conceptualized the project, conducted the experiments, and wrote the paper. S.R. conceptualized the project, secured funding, supervised the research, and wrote the paper. O.V. performed bioinformatics analysis of scRNAseq data. N.A.D. critically reviewed the paper and supervised the *mdx* injection experiments. J.D performed the *mdx* injections and C.D and J.D recorded strength measurements. J.J supervised the Seahorse experiments, and E.H. and C.D performed the Seahorse experiments and analysis. Y.C. performed the library preparations of the scRNAseq samples under supervision of J.R.

Dominici et al.,

Figure Legends

Figure 1. Enhanced self-renewal of MS023-treated MuSCs.

(A) Muscle fibers cultured for 48 h with MS023 or DMSO. Pax7 identifies MuSCs, and ki67 identifies proliferating cells. Scale bar represents 50 μM. (B) Quantification of total MuSCs per fiber. 30 fibers were counted per condition, horizontal bar represents average number of MuSCs per fiber (n=3 mice per condition). (C) Muscle fibers cultured for 48 h with MS023 or DMSO. 30 fibers counted per condition (n=3 mice per condition). Scale bar represents 50 μM. (D) Quantification of Pax7/MyoD expressing MuSCs from (C). (E) Pax7/MyoD immunofluorescence staining of cultured myoblasts treated with MS023 or DMSO for 48 h. Scale bar represents 50 μM. (F) Quantification of Pax7/MyoD expressing cells from (E).

Figure 2. Single-cell graph-based clustering analysis.

(A) UMAP embedding representation for all cells. (B) Gene expression density on the UMAP embedding plot of myogenic markers Pax7, Myf5, MyoD, and Myogenin. (C) Distribution of cells from each sample within the UMAP embedding representation. (D) Proportion of cells from each sample belonging to each of the 10 identified clusters.

Figure 3. Pseudotime trajectory of pooled DMSO- and MS023-treated MuSCs.

(A) Trajectory of all cells depicting three cell states. Pseudotime starts at the left endpoint of the plot. Q: Quiescence. M: Proliferating myoblast. DM: Differentiating myoblast. (B-D) Monocle trajectory of day-0 and DMSO-treated samples (B), heatmap of top differentially regulated genes across pseudotime (C) and GO enrichment analyses of DMSO-unique branchpoint genes (D). (E-G) Monocle trajectory of day-0 and MS023-treated samples (E), heatmap of top differentially

Dominici et al.,

regulated genes across pseudotime (F) and GO enrichment analyses of MS023-unique

branchpoint genes (G).

Figure 4. MS023-treated proliferating MuSCs display elevated oxidative phosphorylation.

(A) UMAP plots showing expression of components of the electron transport chain localized to

MS023 clusters. (B) Seahorse XF96 analysis of oxygen consumption rate (OCR) in DMSO and

MS023-treated freshly isolated primary MuSCs and quantification of basal respiration, maximal

respiration, and ATP-linked production. Error bars represent mean ± SEM from 7 individual

wells per condition. (C) MitoTracker staining (red) of freshly isolated MuSCs treated with

DMSO (upper panel) or MS023 (lower panel) and counterstained with DAPI to visualize nuclei.

Minimum 100 cells quantified per condition from 3 independent experiments. Scale bar

represents 20 μM. (D) Quantification of MitoTracker signal intensity of cells from (C), error

bars represent mean ± SEM from 3 individual replicates, > 100 nuclei quantified per condition (p

<0.0001).

Figure 5. MS023 treatment is reversible and allows expanded MuSCs to differentiate *ex*

***vivo* and *in vivo*.**

(A) Immunofluorescence of MyHC counterstained with DAPI in primary myotubes that were

differentiated in the presence of DMSO or MS023 following 4 days of treatment with DMSO or

MS023. (B) Fusion index calculated as the ratio of nuclei within multinucleated myotubes to

total nuclei from >200 nuclei per condition, error bars represent mean ± SEM from 3 individual

replicates. (C) Immunofluorescence of MyHC counterstained with DAPI in primary myotubes

that were differentiated in the absence of DMSO or MS023 following 4 days of treatment with

Dominici et al.,

DMSO or MS023 and 2 days of washout. Scale bar represents 50 µM (D) Fusion index

calculated the same as 6B. (E) Immunofluorescence of GFP$^+$ myofibers following transplantation

of 15,000 DMSO or MS023-treated MuSCs. Scale bar represents 50 µM. (F) Quantification of

GFP$^+$ myofibers. Error bars represent mean ± SEM from 3 individual biological replicates. (G)

Representative FACS plots of re-isolated MuSCs following engraftment and muscle

regeneration. P3 represents the population of purified GFP+ MuSCs. (H) Fold enrichment of

MS023-treated MuSC re-isolation compared to DMSO. Error bars represent mean ± SEM from 3

individual biological replicates.

Figure 6. *In vivo* **administration of MS023 to dystrophic mice improves muscle strength.**

(A) Experimental schema. (B) Normalized grip strength measurements taken from two forelimbs

and all 4 limbs of mice treated with vehicle or MS023 10 days after the last injection. Error bars

represent mean ± SEM from 6 biological replicates. (C) Force/frequency curve generated for

mice treated with vehicle or MS023. (D) Representative cross-sectional area of TA muscles

isolated from mice treated with vehicle or MS023 and immunostained with anti-laminin

antibodies to visualize myofibers. Scale bar represents 50 µM. (E) Quantification of the variance

coefficient of TA muscle minimum fiber Feret measurement from mice treated with vehicle or

MS023. Error bars represent mean ± SEM from 3 biological replicates, >200 fibers measured per

mouse)

Dominici et al.,

Supplemental Figure S1

(A) Primary MuSCs were isolated from WT mice and seeded on collagen-coated 6 well plates.

Upon seeding, cells were treated with 1μM MS023, 1 μM TP064, or DMSO for 6 days. Cells

were trypsinized and counted at days 2, 4 and 6 using a haemocytometer.

(B) Western blot of total asymmetric dimethyl arginine (ADMA) from whole cell lysate of

primary myoblasts treated with MS023 (1μM) or DMSO. (C) Ki67 immunofluorescence of

MS023-treated primary myoblasts. Error bars represent mean ± SEM from 3 individual

experiments, minimum of 200 cells per condition counted. Scale bar represents 50 μM. (D)

Ki67-PE immunostained MS023-treated primary myoblasts analyzed with FACS, foldchange of

mean intensity calculated compared to DMSO-treated cells. Error bars represent mean ± SEM

from 3 individual replicates. (E) (left) Immunofluorescence of 14th passage MuSCs with DMSO

or MS023 treatment, Pax7 (green), MyoD (purple), counterstained with DAPI. Scale bar

represents 50 μM. (right) Pie chart representing quantification of Pax7 and MyoD expression

from cells in left panel.

Supplemental Figure S2

(A) Reproducibility of average gene expression across biological replicates represented as

Pearson correlation coefficients. (B) UMAP plot labelled with clusters 0-9. (C) Distribution of

individual samples mapped onto the UMAP embedding representation for all cells. (D)

Distribution of cells from d0 across the clusters represented in a pie chart. Cluster colour legend

indicated on the right. (E) Distribution of individual clusters mapped onto the UMAP embedding

representation for all cells. (F) Expression RidgePlots of key myogenic markers Pax7, Myf5,

MyoD (Myod1) and Myog separated by cluster. (G) Expression RidgePlots of key myogenic

Dominici et al.,

markers Pax7, Myf5, MyoD (Myod1) and Myog separated by sample. (H) GO enrichment terms of top 100 marker genes unique to cluster M1 (DMSO specific, top panel), M3 (MS023 specific, middle panel), and genes that are common between clusters M1 (DMSO) and M3 (MS023) (lower panel). (I) GO enrichment terms of top 100 marker genes unique to d4 cells (top panel) and d4MS023 cells (lower panel). (J) Heatmap of the top 100 enriched genes in each cluster.

Supplemental Figure S3

(A) Cell trajectories for each sample and cluster, with the pseudotime starting at the left endpoint of the plot. (B) Heatmap of significantly differentially expressed genes across the branchpoint of the pooled sample trajectory. (C) Distributions in the kinetic trends of each lineage for the expression for selected genes (D) Venn diagram of overlapping differentially regulated genes across the branch point of the DMSO and MS023 trajectories with a q value <0.1.

Supplemental Figure S4

(A) Fold change of selected OxPhos genes in response to treatment with MS023. Bulk RNA was isolated from freshly purified primary myoblasts. Error bars represent mean ± SEM from 3 biological replicates. (B) UMAP plots showing expression of components of glycolysis localized to MS023 clusters. (C) Seahorse XF96 analysis of extracellular acidification rate (ECAR) in DMSO and MS023-treated freshly isolated primary MuSCs and quantification of glycolysis, glycolytic capacity, and glycolytic reserve. Error bars represent mean ± SEM from 7 individual wells per condition. (D) Measurement of glucose uptake (left) and lactate production (right) as quantified from the media of d4 (DMSO) and d4MS023 (MS023) MuSCs. Error bars represent mean ± SEM from 3 independent experiments.

Dominici et al.,

Supplemental Figure S5

(A) Experimental outline for MuSC engraftment. (B) FACS plot of primary purified MuSCs

represented as P4 (0.6% of the total population). (C) Purity of sorted cells (99%)

Supplemental Figure S6

(A) Whole tissue lysate from homogenized tail pieces of mice treated with Vehicle or MS023

probed with antibodies against total asymmetric dimethyl arginine (ADMA), symmetric

dimethyl arginine (SDMA), and actin for loading control. (B) Hanging test measurements taken

from mice treated with vehicle or MS023 at 10d post-injection. Error bars represent mean ± SEM

from 6 biological replicates.

References

Bayen S, Saini S, Gaur P, Duraisamy AJ, Kumar Sharma A, Pal K, Vats P, Singh SB (2018)
    PRMT1 promotes hyperglycemia in a FoxO1-dependent manner, affecting glucose
    metabolism, during hypobaric hypoxia exposure, in rat model. *Endocrine* 59: 151-163
Bedford MT, Clarke SG (2009) Protein arginine methylation in mammals: who, what, and why.
    Mol Cell 33: 1-13
Bentzinger CF, von Maltzahn J, Dumont NA, Stark DA, Wang YX, Nhan K, Frenette J,
    Cornelison DD, Rudnicki MA (2014) Wnt7a stimulates myogenic stem cell motility and
    engraftment resulting in improved muscle strength. *J Cell Biol* 205: 97-111
Blanc RS, Richard S (2017) Regenerating muscle with arginine methylation. *Transcription* 8:
    175-178
Blanc RS, Vogel G, Chen T, Crist C, Richard S (2016) PRMT7 Preserves Satellite Cell
    Regenerative Capacity. *Cell Rep* 14: 1528-1539
Blanc RS, Vogel G, Li X, Yu Z, Li S, Richard S (2017) Arginine Methylation by PRMT1
    Regulates Muscle Stem Cell Fate. *Mol Cell Biol* 37
Bulfield G, Siller WG, Wight PA, Moore KJ (1984) X chromosome-linked muscular dystrophy
    (mdx) in the mouse. *Proc Natl Acad Sci U S A* 81: 1189-1192
Butler A, Hoffman P, Smibert P, Papalexi E, Satija R (2018) Integrating single-cell
    transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol*
    36: 411-420

Cho DS, Doles JD (2017) Single cell transcriptome analysis of muscle satellite cells reveals widespread transcriptional heterogeneity. *Gene* 636: 54-63

Couturier CP, Ayyadhury S, Le PU, Nadaf J, Monlong J, Riva G, Allache R, Baig S, Yan X, Bourgey M *et al* (2020) Single-cell RNA-seq reveals that glioblastoma recapitulates a normal neurodevelopmental hierarchy. *Nature Communications* 11: 3406

De Micheli AJ, Laurilliard EJ, Heinke CL, Ravichandran H, Fraczek P, Soueid-Baumgarten S, De Vlaminck I, Elemento O, Cosgrove BD (2020) Single-Cell Analysis of the Muscle Stem Cell Hierarchy Identifies Heterotypic Communication Signals Involved in Skeletal Muscle Regeneration. *Cell Rep* 30: 3583-3595.e3585

Dell'Orso S, Juan AH, Ko KD, Naz F, Perovanovic J, Gutierrez-Cruz G, Feng X, Sartorelli V (2019) Single cell analysis of adult mouse skeletal muscle stem cells in homeostatic and regenerative conditions. *Development* 146

Dort J, Orfi Z, Fabre P, Molina T, Conte TC, Greffard K, Pellerito O, Bilodeau JF, Dumont NA (2021) Resolvin-D2 targets myogenic cells and improves muscle regeneration in Duchenne muscular dystrophy. *Nat Commun* 12: 6264

Dumont NA, Wang YX, von Maltzahn J, Pasut A, Bentzinger CF, Brun CE, Rudnicki MA (2015) Dystrophin expression in muscle stem cells regulates their polarity and asymmetric division. *Nat Med* 21: 1455-1463

Eram MS, Shen Y, Szewczyk M, Wu H, Senisterra G, Li F, Butler KV, Kaniskan HU, Speed BA, Dela Sena C *et al* (2016) A Potent, Selective, and Cell-Active Inhibitor of Human Type I Protein Arginine Methyltransferases. *ACS Chem Biol* 11: 772-781

Folmes CD, Dzeja PP, Nelson TJ, Terzic A (2012) Metabolic plasticity in stem cell homeostasis and differentiation. *Cell Stem Cell* 11: 596-606

Fu X, Xiao J, Wei Y, Li S, Liu Y, Yin J, Sun K, Sun H, Wang H, Zhang Z *et al* (2015a) Combination of inflammation-related cytokines promotes long-term muscle stem cell expansion. *Cell Res* 25: 1082-1083

Fu X, Zhu MJ, Dodson MV, Du M (2015b) AMP-activated protein kinase stimulates Warburg-like glycolysis and activation of satellite cells during muscle regeneration. *J Biol Chem* 290: 26445-26456

Gallot YS, Hindi SM, Mann AK, Kumar A (2016) Isolation, Culture, and Staining of Single Myofibers. *Bio Protoc* 6

Hardie DG, Schaffer BE, Brunet A (2016) AMPK: An Energy-Sensing Pathway with Multiple Inputs and Outputs. *Trends Cell Biol* 26: 190-201

Herzig S, Shaw RJ (2018) AMPK: guardian of metabolism and mitochondrial homeostasis. *Nat Rev Mol Cell Biol* 19: 121-135

Hori S, Hiramuki Y, Nishimura D, Sato F, Sehara-Fujisawa A (2019) PDH-mediated metabolic flow is critical for skeletal muscle stem cell differentiation and myotube formation during regeneration in mice. *FASEB J* 33: 8094-8109

Iwasaki H, Yada T (2007) Protein arginine methylation regulates insulin signaling in L6 skeletal muscle cells. *Biochem Biophys Res Commun* 364: 1015-1021

Jain K, Clarke SG (2019) PRMT7 as a unique member of the protein arginine methyltransferase family: A review. *Arch Biochem Biophys* 665: 36-45

Judson RN, Quarta M, Oudhoff MJ, Soliman H, Yi L, Chang CK, Loi G, Vander Werff R, Cait A, Hamer M *et al* (2018) Inhibition of Methyltransferase Setd7 Allows the In Vitro Expansion of Myogenic Stem Cells with Improved Therapeutic Potential. *Cell Stem Cell* 22: 177-190.e177

Kawabe Y, Wang YX, McKinnell IW, Bedford MT, Rudnicki MA (2012) Carm1 regulates Pax7 transcriptional activity through MLL1/2 recruitment during asymmetric satellite stem cell divisions. *Cell Stem Cell* 11: 333-345

Kim H, Selvaraj S, Kiley J, Azzag K, Garay BI, Perlingeiro RCR (2021) Genomic Safe Harbor Expression of PAX7 for the Generation of Engraftable Myogenic Progenitors. *Stem Cell Reports* 16: 10-19

Kimmel JC, Hwang AB, Scaramozza A, Marshall WF, Brack AS (2020) Aging induces aberrant state transition kinetics in murine muscle stem cells. *Development* 147

L'Honore A, Commere PH, Negroni E, Pallafacchina G, Friguet B, Drouin J, Buckingham M, Montarras D (2018) The role of Pitx2 and Pitx3 in muscle stem cells gives new insights into P38alpha MAP kinase and redox regulation of muscle regeneration. *Elife* 7

Lepper C, Partridge TA, Fan CM (2011) An absolute requirement for Pax7-positive satellite cells in acute injury-induced skeletal muscle regeneration. *Development* 138: 3639-3646

Machado L, Esteves de Lima J, Fabre O, Proux C, Legendre R, Szegedi A, Varet H, Ingerslev LR, Barres R, Relaix F *et al* (2017) In Situ Fixation Redefines Quiescence and Early Activation of Skeletal Muscle Stem Cells. *Cell Rep* 21: 1982-1993

McKellar DW, Walter LD, Song LT, Mantri M, Wang MFZ, De Vlaminck I, Cosgrove BD (2021) Large-scale integration of single-cell transcriptomic data captures transitional progenitor states in mouse skeletal muscle regeneration. *Commun Biol* 4: 1280

Montarras D, Morgan J, Collins C, Relaix F, Zaffran S, Cumano A, Partridge T, Buckingham M (2005) Direct isolation of satellite cells for skeletal muscle regeneration. *Science* 309: 2064-2067

Nakayama K, Szewczyk MM, Dela Sena C, Wu H, Dong A, Zeng H, Li F, de Freitas RF, Eram MS, Schapira M *et al* (2018) TP-064, a potent and selective small molecule inhibitor of PRMT4 for multiple myeloma. *Oncotarget* 9: 18480-18493

Pala F, Di Girolamo D, Mella S, Yennek S, Chatre L, Ricchetti M, Tajbakhsh S (2018) Distinct metabolic states govern skeletal muscle stem cell fates during prenatal and postnatal myogenesis. *J Cell Sci* 131

Pasut A, Oleynik P, Rudnicki MA (2012) Isolation of muscle stem cells by fluorescence activated cell sorting cytometry. *Methods Mol Biol* 798: 53-64

Qiao X, Kim DI, Jun H, Ma Y, Knights AJ, Park MJ, Zhu K, Lipinski JH, Liao J, Li Y *et al* (2019) Protein Arginine Methyltransferase 1 Interacts With PGC1alpha and Modulates Thermogenic Fat Activation. *Endocrinology* 160: 2773-2786

Relaix F, Bencze M, Borok MJ, Der Vartanian A, Gattazzo F, Mademtzoglou D, Perez-Diaz S, Prola A, Reyes-Fernandez PC, Rotini A *et al* (2021) Perspectives on skeletal muscle stem cells. *Nat Commun* 12: 692

Relaix F, Zammit PS (2012) Satellite cells are essential for skeletal muscle regeneration: the cell on the edge returns centre stage. *Development* 139: 2845-2856

Rudnicki MA, Schnegelsberg PN, Stead RH, Braun T, Arnold HH, Jaenisch R (1993) MyoD or Myf-5 is required for the formation of skeletal muscle. *Cell* 75: 1351-1359

Ryall JG, Dell'Orso S, Derfoul A, Juan A, Zare H, Feng X, Clermont D, Koulnis M, Gutierrez-Cruz G, Fulco M *et al* (2015) The NAD(+)-dependent SIRT1 deacetylase translates a metabolic switch into regulatory epigenetics in skeletal muscle stem cells. *Cell Stem Cell* 16: 171-183

Dominici et al.,

Ryder-Cook AS, Sicinski P, Thomas K, Davies KE, Worton RG, Barnard EA, Darlison MG, Barnard PJ (1988) Localization of the mdx mutation within the mouse dystrophin gene. *Embo j* 7: 3017-3021

Saber J, Rudnicki MA (2022) Carm1 and the Epigenetic Control of Stem Cell Function. *Stem Cells Transl Med*

Sacco A, Doyonnas R, Kraft P, Vitorovic S, Blau HM (2008) Self-renewal and expansion of single transplanted muscle stem cells. *Nature* 456: 502-506

Sakai H, Fukuda S, Nakamura M, Uezumi A, Noguchi YT, Sato T, Morita M, Yamada H, Tsuchida K, Tajbakhsh S *et al* (2017) Notch ligands regulate the muscle stem-like state ex vivo but are not sufficient for retaining regenerative capacity. *PLoS One* 12: e0177516

Seale P, Sabourin LA, Girgis-Gabardo A, Mansouri A, Gruss P, Rudnicki MA (2000) Pax7 is required for the specification of myogenic satellite cells. *Cell* 102: 777-786

Serrano AL, Mann CJ, Vidal B, Ardite E, Perdiguero E, Muñoz-Cánoves P (2011) Cellular and molecular mechanisms regulating fibrosis in skeletal muscle repair and disease. *Curr Top Dev Biol* 96: 167-201

Soleimani VD, Punch VG, Kawabe Y, Jones AE, Palidwor GA, Porter CJ, Cross JW, Carvajal JJ, Kockx CE, van IWF *et al* (2012) Transcriptional dominance of Pax7 in adult myogenesis is due to high-affinity recognition of homeodomain motifs. *Dev Cell* 22: 1208-1220

Stouth DW, Manta A, Ljubicic V (2018) Protein arginine methyltransferase expression, localization, and activity during disuse-induced skeletal muscle plasticity. *Am J Physiol Cell Physiol* 314: C177-C190

Stouth DW, vanLieshout TL, Ng SY, Webb EK, Manta A, Moll Z, Ljubicic V (2020) CARM1 Regulates AMPK Signaling in Skeletal Muscle. *iScience* 23: 101755

Teyssier C, Ma H, Emter R, Kralli A, Stallcup MR (2005) Activation of nuclear receptor coactivator PGC-1alpha by arginine methylation. *Genes Dev* 19: 1466-1473

Theret M, Gsaier L, Schaffer B, Juban G, Ben Larbi S, Weiss-Gayet M, Bultot L, Collodet C, Foretz M, Desplanches D *et al* (2017) AMPKalpha1-LDH pathway regulates muscle stem cell self-renewal by controlling metabolic homeostasis. *EMBO J* 36: 1946-1962

Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, Lennon NJ, Livak KJ, Mikkelsen TS, Rinn JL (2014) The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat Biotechnol* 32: 381-386

Troy A, Cadwallader AB, Fedorov Y, Tyner K, Tanaka KK, Olwin BB (2012) Coordination of satellite cell activation and self-renewal by Par-complex-dependent asymmetric activation of p38α/β MAPK. *Cell Stem Cell* 11: 541-553

van den Brink SC, Sage F, Vertesy A, Spanjaard B, Peterson-Maduro J, Baron CS, Robin C, van Oudenaarden A (2017) Single-cell sequencing reveals dissociation-induced gene expression in tissue subpopulations. *Nat Methods* 14: 935-936

van Velthoven CTJ, de Morree A, Egner IM, Brett JO, Rando TA (2017) Transcriptional Profiling of Quiescent Muscle Stem Cells In Vivo. *Cell Rep* 21: 1994-2004

vanLieshout TL, Ljubicic V (2019) The emergence of protein arginine methyltransferases in skeletal muscle and metabolic disease. *Am J Physiol Endocrinol Metab* 317: E1070-E1080

Wang YX, Bentzinger CF, Rudnicki MA (2013) Treating muscular dystrophy by stimulating intrinsic repair. *Regen Med* 8: 237-240

Dominici et al.,

Xu J, Richard S (2021) Cellular pathways influenced by protein arginine methylation: Implications for cancer. *Mol Cell* 81: 4357-4368

Yin H, Price F, Rudnicki MA (2013) Satellite cells and the muscle stem cell niche. *Physiol Rev* 93: 23-67

Zhang T, Gunther S, Looso M, Kunne C, Kruger M, Kim J, Zhou Y, Braun T (2015) Prmt5 is a regulator of muscle stem cell expansion in adult mice. *Nat Commun* 6: 7140

Dominici et al., Figure 1

Dominici et al., Figure 3

A State ● Q/M ● DM ● M — Pseudotime 0 10 20

B DMSO trajectory

E MS023 trajectory

C

F

D DMSO-unique branchpoint genes

GO Term	p-value
mRNA splicing, via spliceosome (GO:0000398)	1.47E-07
mRNA processing (GO:0006397)	5.91E-07
protein modification by small protein removal (GO:0070646)	2.53E-06
RNA splicing, via transesterification reactions with bulged adenosine as nu	3.20E-06
ubiquitin-dependent protein catabolic process (GO:0006511)	3.86E-06
protein deubiquitination (GO:0016579)	4.22E-06
mitochondrial gene expression (GO:0140053)	1.25E-05
proteasome-mediated ubiquitin-dependent protein catabolic process (GO:	1.66E-05
protein polyubiquitination (GO:0000209)	1.90E-05

G MS023-unique branchpoint genes

GO Term	p-value
translation (GO:0006412)	6.80E-27
ribosome biogenesis (GO:0042254)	2.00E-16
respiratory electron transport chain (GO:0022904)	9.97E-13
mitochondrial ATP synthesis coupled electron transport (GO:0042775)	6.57E-12
NADH dehydrogenase complex assembly (GO:0010257)	5.22E-10
mitochondrial respiratory chain complex I biogenesis (GO:0097031)	5.22E-10
mitochondrial respiratory chain complex I assembly (GO:0032981)	5.22E-10
mitochondrial respiratory chain complex assembly (GO:0033108)	1.62E-09

Dominici et al., Figure 4

A

B

C

D

Dominici et al., Figure 5

Dominici et al., Figure 6

Supplemental figures

Supplemental figure S1

A

B

C

D

E

Supplemental figure S2

Supplemental figure S2 (cont'd)

F

G

H

M1
(DMSO)
unique

Biological Process (GO)			
GO-term	description	count in gene set	false discovery rate
GO:0097237	cellular response to toxic substance	9 of 190	2.78e-05
GO:1990748	cellular detoxification	6 of 83	0.00034
GO:0009636	response to toxic substance	11 of 471	0.00034
GO:2000812	regulation of barbed-end actin filament capping	3 of 5	0.00072
GO:0098869	cellular oxidant detoxification	5 of 79	0.0021

M3
(MS023)
unique

Biological Process (GO)			
GO-term	description	count in gene set	false discovery rate
GO:0006412	translation	28 of 313	5.15e-29
GO:0006518	peptide metabolic process	28 of 440	9.75e-26
GO:1901566	organonitrogen compound biosynthetic process	32 of 1122	7.94e-20
GO:0002181	cytoplasmic translation	11 of 55	5.45e-14
GO:0044271	cellular nitrogen compound biosynthetic process	36 of 2781	5.23e-12

M1
M3
overlap

Biological Process (GO)			
GO-term	description	count in gene set	false discovery rate
GO:0042127	regulation of cell population proliferation	12 of 1594	0.00035
GO:0065008	regulation of biological quality	15 of 3420	0.0020
GO:0042981	regulation of apoptotic process	10 of 1476	0.0020
GO:0033599	regulation of mammary gland epithelial cell proliferation	3 of 18	0.0020
GO:0033554	cellular response to stress	10 of 1370	0.0020

I

d4
unique

Reactome Pathways			
pathway	description	count in gene set	false discovery rate
MMU-8953854	Metabolism of RNA	13 of 448	7.48e-06
MMU-72766	Translation	7 of 148	0.00031
MMU-72172	mRNA Splicing	6 of 163	0.0034
MMU-72163	mRNA Splicing - Major Pathway	6 of 156	0.0034
MMU-499943	Interconversion of nucleotide di- and triphosphates	3 of 24	0.0073

d4MS023
unique

Reactome Pathways			
pathway	description	count in gene set	false discovery rate
MMU-163200	Respiratory electron transport, ATP synthesis by chemiosm...	10 of 78	6.30e-10
MMU-1428517	The citric acid (TCA) cycle and respiratory electron transport	10 of 129	3.15e-08
MMU-8949613	Cristae formation	5 of 17	2.14e-06
MMU-163210	Formation of ATP by chemiosmotic coupling	5 of 17	2.14e-06
MMU-156827	L13a-mediated translational silencing of Ceruloplasmin exp...	6 of 50	5.41e-06

Supplemental figure S2 (cont'd)

Supplemental figure S3

Supplemental figure S4

A

Fold change normalized to Mlec/Parn

DMSO (black), MS023 (teal)

Nd1 Atp5o Ndufa2* Cox5b* Uqcr11

B

Hk1 Eno1 Pfkm Gapdh Pkm Slc2a1

C

ECAR (Glycolysis)

ECAR (mpH/min/Norm. Unit) vs Time (minutes)

DMSO MS023

Glycolysis *p*=0.02
DMSO MS023

Glycolytic Capacity *p*=0.004
DMSO MS023

Glycolytic Reserve *p*=0.001
DMSO MS023

D

Glucose uptake *p* = 0.0042
DMSO MS023

Lactate production
Undet.
DMSO MS023

Supplemental figure S5

A

Day 0
Isolate
MuSCs

Culture

Day 5
Inj. CTX in
recipient TA

Day 6
Inj. 15K
MuSCs

3 weeks
post-CTX
Inj.

β-Actin-GFP

DMSO MS023

Sacrifice

B

Tube: Sample_001

Population	#Events	%Parent	%Total
All Events	159,388	####	100.0
Cells	4,848	3.0	3.0
FSC Singlets	3,377	69.7	2.1
SSC Singlets	3,036	89.9	1.9
P1	2,095	69.0	1.3
P2	1,216	58.0	0.8
P4	1,000	82.2	0.6

C

Tube: Sample_002 Purity

Population	#Events	%Parent	%Total
All Events	248	####	100.0
Cells	105	42.3	42.3
FSC Singlets	105	100.0	42.3
SSC Singlets	104	99.0	41.9
P1	103	99.0	41.5
P2	102	99.0	41.1
P4	101	99.0	40.7

Supplemental figure S6

A

B

Hanging test 10d

ns

■ Vehicle
□ MS023

Hanging test 20d

■ Vehicle
□ MS023