Tet-dependent 5-hydroxymethyl-Cytosine modification of mRNA regulates the axon guidance genes *robo2* and *slit* in *Drosophila*

Badri Nath Singh*[1], Hiep Tran*[1], Joseph Kramer[3], Elmira Kirishenko[1], Neha Changela[1], Fei Wang[1], Yaping Feng[1], Dibyendu Kumar[1,] Min Tu[1], Shun Liang[4], Jie Lan[4,5], Martin Bizet[4], François Fuks [4], and Ruth Steward[1,2,6]

*equal contribution

[1] Waksman Institute, Rutgers University, Piscataway, NJ 08854, [2]Department of Molecular Biology and Biochemistry, Cancer Institute of New Jersey, Rutgers University

[3]Department of Pathology and Laboratory Medicine, Rutgers Biomedical and Health Sciences, Rutgers University, New Brunswick

[4]Laboratory of Cancer Epigenetics, Faculty of Medicine, ULB Cancer Research Center (U-CRC), Université Libre de Bruxelles (ULB), Brussels, Belgium.

[5]Present address, Institute for Genetics, Justus-Liebig University Giessen, 35392 Giessen, Germany

[6]Corresponding author: E-mail, steward@waksman.rutgers.edu

Abstract

Modifications of mRNA, especially methylation of adenosine, have recently drawn much attention. The much rarer modification, 5-hydroxymethylation of cytosine (5hmC), is not well understood and is the subject of this study. Vertebrate Tet proteins are 5-methylcytosine (5mC) hydroxylases enzymes catalyzing the transition of 5mC to 5hmC in DNA and have recently been shown to have the same function in messenger RNAs in both vertebrates and in Drosophila. The *Tet* gene is essential in Drosophila because Tet knock-out animals do not reach adulthood. We describe the identification of Tet-target genes in the embryo and larval brain by determining Tet DNA-binding sites throughout the genome and by mapping the Tet-dependent 5hmrC modifications transcriptome-wide. 5hmrC-modified sites can be found along the entire transcript and are preferentially located at the promoter where they overlap with histone H3K4me3 peaks. The identified mRNAs are frequently involved in neuron and axon development and Tet knock-out led to a reduction of 5hmrC marks on specific mRNAs. Among the Tet-target genes were the *robo2* receptor and its *slit* ligand that function in axon guidance in Drosophila and in vertebrates. *Tet* knock-out embryos show overlapping phenotypes with *robo2* and are sensitized to reduced levels of *slit*. Both Robo2 and Slit protein levels were markedly reduced in Tet KO larval brains. Our results establish a role for Tet-dependent 5hmrC in facilitating the translation of modified mRNAs, primarily in developing nerve cells.

Main

The regulatory function of epigenetic mechanisms such as modifications of specific DNA bases or amino acids in histone tails have been investigated for many years. These processes are overlayed upon the genetic code and have profound effects on transcription and overall gene expression. The importance of similar modifications of RNA bases has become apparent and its pervasiveness has engendered the nascent field of epitranscriptomics[1]. Approximately 150 modifications of all four nucleosides have been detected in total RNAs [2]. These modifications are mostly associated with the more abundant ribosomal and transfer RNAs but are also present in a subset of messenger RNA. The mRNA modifications provide a critical layer of regulation of the transcriptome in both Drosophila and vertebrates, and influence gene expression through the control of mRNA biogenesis [3]. Cytosine bases convey epigenetic information in both DNA and mRNA. 5-methylcytosine (5mrC) is abundant in RNA and present in cytoplasmic and mitochondrial ribosomal RNA, t-RNA, non-coding RNA, and mRNA [4]. In contrast, 5hmrC is detected, in mRNA and is much less abundant [5].

In Drosophila DNA, 5mC is present at low levels and so far, no function has been documented for it. Also, the methyltransferases that catalyze the C-methylation in vertebrates are not present in Drosophila, with the exception of DNMT2, which primarily modifies several tRNAs and viral transcripts [6]. However, both 5mrC and 5hmrC are present in *Drosophila* RNA. The 5hmrC modification appears to be specific to mRNA and is controlled, at least in part, by the Drosophila Tet (Ten-Eleven-Translocation) protein [5]. Tet proteins were first identified as DNA-modifying enzymes that function as 5-methylcytosine (5mC) hydroxylases, catalyzing the transition of 5mC to 5hmC in vertebrate DNA [7].

The three vertebrate *TET* genes (*TET1, 2* and *3)* function as epigenetic regulators of gene expression. The transition of 5mC to 5hmC leads to the elimination of the methyl mark on DNA and activates the transcription of target genes [7]. Mammalian TET proteins, TET3 in particular, catalyze the same reaction on RNA, converting 5mrC to 5hmrC in tissue culture and mouse embryonic stem cells (ESCs) [8]. Vertebrate TET1 and TET3 isoforms have an N-terminal DNA binding domain (CxxC) and a C-terminal metal-binding catalytic domain (HxD), while TET2 lacks the N-terminal domain [9]. *Drosophila* has only one *Tet* gene, that encodes the two major protein forms from two distinct promoters [10]. The larger protein (Tet-L) includes the DNA binding and catalytic domains, while the smaller form (Tet-S) has only the catalytic domain. Both DNA binding and catalytic domains of *Drosophila* Tet are about 50% homologous to those of TET1 and 3, and the specific amino acids within the catalytic domain responsible for metal ion-binding are identical in *Drosophila* and vertebrates [11].

Complete loss-of function of *Tet* (*Tet^null^*) leads to lethality in the late pupal stage, with partial loss-of-function alleles surviving as adults for varying amounts of time [10]. All mutant animals show abnormal locomotion and knock-down of Tet in neurons that control the circadian rhythm results in perturbation of that rhythm, indicating that Tet is likely essential in diverse neuronal cells. The neuronal phenotypes agree well with the expression of the Tet gene, first expressed in three-hour old embryos,

and throughout embryogenesis and larval development the protein is found primarily in nerve cells [10,12].

While the role of Tet in vertebrate DNA modification and its consequences have been reported

in much detail, little is known about the function of Tet and 5hmrC in mRNA. Tet2 regulates pathogen

induced myelopoiesis as well as endogenous retroviruses by controlling the 5hmrC mark on mRNAs [13].

The 5hmrC mark is present in mRNA of mouse Embryonic Stem Cells (ESC), where Tet proteins

control the 5 hydroxymethylation of key-pluripotency transcripts [14].

While Tet function in RNA modification has been analyzed in tissue culture cells in *Drosophila*

and mouse, we report our work on identifying genes that are regulated by Tet in *Drosophila* embryos

and nerve tissue. These Tet-target genes were identified through genome- and transcriptome-wide

experiments, namely ChIP-seq, hmeRIP-seq, and RNA-seq. Two of these target genes, *robo2* and *slit*,

are known for their requirement in axon guidance in both vertebrates and Drosophila and we chose

them for further analyses. We found that *Tet* mutant animals show overlapping phenotypes with *robo2*

and *slit* in the developing nervous system, and that *slit* dominantly enhances defects in CNS

development in *Tet^null* embryos. Further, we found that Tet activity and 5hmrC modification of the mRNA

encoding *robo2* and *slit* is required for the proper expression of these pathfinding genes since loss of

Tet results in reduced protein expression.

97 Results

98 Tet functions as a 5-methylcytosine hydroxylase and modifies polyA$^+$ RNA.

Previously we have shown by dot blot analysis in S2 Drosophila cells and larval brains that the

5hmrC modification was primarily found on polyA$^+$ RNA and was strongly reduced in Tet knock-down

(KD) cells as well as in larval brains from complete loss-of-function animals (*Tet^null*) [5]. We have

confirmed and quantified these results using ultra-performance liquid chromatography tandem mass

spectrometry (UHPLC-MS/MS). Measurements of 5mrC and 5hmrC abundance in S2 cells indicate that

5hmrC was strongly enriched in polyA$^+$ RNA whereas 5mrC was underrepresented in that fraction as

compared to total RNA (Fig. 1A and B). Thus, our results are consistent with the observation that 5mrC

is associated with rRNA, tRNA and polyA$^+$ RNA, while 5hmrC is primarily found in mRNA. We then

examined changes of 5hmrC and 5mrC in polyA$^+$ RNA isolated from normal and *Tet^null* larval brains. We

found that 5hmrC was decreased about 5-fold in the mutant brains as compared to control (Fig. 1C).

Moreover, 5mrC was observed to increase almost 3-fold in the absence of Tet function. (Figure 1D).

Similar results were found in wildtype (wt) and Tet KD embryos (Fig. S1A and B). These results confirm

and extend our previous antibody-based analyses [5] and indicate that Tet is responsible for much of the

conversion of 5mrC to 5hmrC in *Drosophila* mRNA.

114 Tet binds DNA preferentially at the transcription start site of target genes in *Drosophila*.

Members of the Tet protein family are known DNA and RNA binding proteins. Moreover, in

vertebrates Tet proteins have been shown to bind DNA at promoter regions to regulate gene
expression through active DNA demethylation [14,15]. We sought to identify the genes that are regulated
by *Drosophila* Tet. We began our experiments by determining if *Drosophila* Tet also binds DNA and
mapping the binding sites. We performed ChIP-seq experiments and mapped Tet-binding peaks
genome wide using a Tet-GFP fusion protein in two samples from different stages of development: 3rd
instar larval brain and imaginal discs (larval brain fraction, LBF) and 0-12h embryos. Samples were
normalized to input chromatin. As negative control we used chromatin from LBF and 0-12 h embryos
lacking GFP however it did not produce enough material for library preparation and sequencing (see
methods).

Bioinformatic analysis of the LBF ChIP-seq results identified 3413 Tet binding peaks distributed
on 2240 genes. Example of Tet binding peak profile is shown in Fig. 2A. Tet preferentially occupies
promoter regions (Fig. 2B) and shows the strongest binding to promoter regions. (Fig. 2C). In murine
ESCs a GC rich DNA motif has been shown to be enriched in Tet1 bound loci [15]. In LBF we identified a
highly conserved CG-rich sequence as one of the highest-ranking motives within Tet bound regions
using MEME-ChIP Motif Analysis (Fig. 2D and Fig. S2A).

A Tet-binding profile in a composite model across the protein coding regions illustrates that Tet
binding is highest near the promoter and gradually decreases until it undergoes a notable drop at the
transcription termination sites (TTS). This closely mirrors the profile observed for H3K4me3, an
epigenetic mark associated with actively transcribing regions frequently found at transcription start sites
[16] (Fig. 2E). While 36% of all Tet peaks co-localize with this chromatin modification (H3K4me3, Fig. 2F),
40% of the Tet binding sites that are localized to the promoter region co-localized with the H3K4me3
mark (Fig. 2G).

In embryo samples, we detected 5180 Tet-binding peaks associated with 2578 genes. Example
of Tet binding peak profile is shown in Fig. 3A. Tet is enriched throughout the gene body and intronic
regions (Fig. 3B) however the strength of binding is strongest at promoters and (Fig 3C). A Tet-binding
profile across the protein coding regions is similar to that observed in LBF (Fig. 3E). Analysis of the
DNA sequences bound by Tet protein in embryos uncovered a highest ranking binding motif that shows
significant similarity to the larval Tet consensus sequence (Fig. 3D and S2) and, as with the larval ChIP
samples, we observe Tet occupancy to be correlated with H3K4me3 binding sites, primarily associated
with promoters (Fig 3E): 42% of all embryonic Tet peaks co-localized with H3K4me3 chromatin
modification marks (Fig. 3F) and 51% of the promoter binding sites overlapped with H3K4me3 mark
(Fig. 3G). In both embryos and LBF Tet binds to approximately the same number of target genes and
30% of Tet's targets are identical in both tissues (Fig 3H).

Our results indicate that Tet binding sites are distributed throughout the physical map of the
genome (Fig. S2B). To confirm these results and show that the Tet-DNA binding domain is sufficient to
target Tet to DNA we constructed transgenic flies carrying a Myc-tagged DNA-binding domain of Tet

(CxxC) under the control of the heat shock promoter (hsp70-GAL4::UAS-TetCxxCRFPmyc). We

expressed the Tet DNA-binding domain by exposing larvae to heat shock and stained salivary glands

with anti-Myc and anti-H3K4me3 antibody. Tet showed many bands distributed on all arms of the

chromosomes, but virtually no staining of the chromocenter which contains very few genes nor of

ribosomal RNA in the nucleolus. H3K4me3 is also present in a distinct binding pattern on all

chromosome, but in contrast to Tet is abundant in the chromocenter and the nucleolus. As indicated by

Chip-seq, HA-Tet and H3K4me3 staining overlapped significantly on giant chromosomes (FigS2C).

These staining results are in agreement with our observation that Tet binds to genes on all

chromosomes of *Drosophila* (Fig. S2).

Our Chip-seq experiments were done in embryos and LBF, two tissues at diverse stages of fly

development, but in which Tet protein is highly expressed. The distribution and characteristics of

binding sites; In both tissues we identified about 2500 genes genome-wide that showed significant Tet-

binding. Tet binding characteristics were similar in both tissues in that the most significant Tet-binding

peaks, showing strongest binding, were preferentially located at promoters. Thus, it appears that only

part of the Tet targets are fixed while others show stage-specific variations throughout development.

Identification of Tet-target mRNAs by hMeRIP-seq in fly tissues

We next determined how many of the genes with Tet-binding peaks also showed 5hmrC

modifications of their RNA. To do this we mapped Tet-dependent 5hmrC modifications on RNAs

transcriptome-wide in the same tissues we used for our Chip-seq analysis. We first performed hMeRIP-

seq on total RNA using basically the same approach we used previously in S2 cells [5]. RNAs isolated

from wt 0-12 h embryos and from wt and *Tet^null* Larval Brain Fraction (LBF) was treated with anti-5hmC

anti-body or immunoglobulin as negative control, and followed by Next Generation Sequencing (NGS,

see methods).

In the embryo we identified 1815 peaks on 1402 mRNAs. A representative 5hmrC peak profile is

shown in Fig. 4A. The 5hmrC modification is preferentially associated with gene bodies and a

comparison to the expected distribution of peaks shows that the modification is not random (Fig. 4B).

Moreover, as the presence of the 5hmrC modification is not proportional to the abundance of the mRNA

the modification appears to function broadly within the transcriptome and is not a regulatory modality

restricted either to rare or hyperabundant transcripts (Fig. 4C). The 5hmrC-associated sequences

identified from these experiments revealed a specific UC-rich motif present within these mRNAs that

closely resembles the motif observed in S2 cells and mammalian ESCs (Fig. 4E and Fig. S3) [5,14].

In mRNA from the wild type LBF, we detected 3711 peaks on 1775 transcripts. A representative

profile of 5hmrC enriched peaks in wt and *Tet^null* is shown in Fig. 5A. In wt the peaks were distributed

across the gene body (Fig. 5B) and 5hmrC marks were found to decorate mRNAs independent of their

abundance (Fig. 5D). Analysis of the peak sequences indicated the modifications were primarily

associated with a UC-rich motif highly related to that identified in embryonic samples (Fig. 5F). In

mRNA from *Tet^null* larvae we identified 5,374 peaks in 1710 mRNA. Comparison of mRNAs identified in

both the wt and *Tet^null* samples indicate that the distribution of 5hmrC peaks is similar both in the

presence and absence of Tet function. However, In the *Tet^null* samples, 45% of the transcripts identified

had at least one peak that showed a reduction of >1.4-fold in the 5hmrC modification relative to wild-

type (Fig. 5C) and the reduction was most pronounced on intronic and coding region peaks (45% and

46%) compared to the peaks found in the UTRs (5', 19%, and 3', 16%). Thus, within a given mRNA

transcript some peaks were affected in *Tet^null* LBF, while others remained unchanged. These results

suggest preference of Tet to modify specific regions of transcripts.

In addition, 37% of the modified mRNA in embryos were also identified in the LBF, while 30% of

the larval modified mRNAs were also present in the embryonic fraction (Fig. S4C). Taken together

these results suggest that Tet targets a distinct cohort of mRNAs in embryos and larval brains and

controls specific 5hmrC modifications along transcripts.

RNA levels in wild type and *Tet^null* larval brains

Our results so far indicate that Tet binds to the promoter of a subset of possibly actively transcribed

genes and controls the 5hmrC modification of their mRNAs. The modification may have an effect on the

stability, processing, and/or translation of the transcripts. To determine if there is a link between 5hmrC

modification and mature mRNA levels, we performed NGS of RNA isolated from wildtype and *Tet^null*

LBF. We found that out of 9000 total transcripts the levels of 445 were significantly increased and 115

were decreased in *Tet^null* LBF (Fig. 6A). When we compared these mRNAs with the 5hmrC-modified

mRNAs present in LBF, we found that 1716 or ~20% of the total transcripts were modified, but only 15

or 3 % of the RNAs that were upregulated in *Tet^null* and 13 or 11 % of the decreased mRNAs were

modified (Fig. 6A, B). This result indicates that the levels of the vast majority of 5hmrC modified

mRNAs do not change levels in *Tet^null* LBF. Thus, the 5hmrC modification of the mRNAs does not

appear to control the steady state level of transcripts. It is therefore likely that the change in levels of

the mRNAs observed in *Tet^null* brains represent a secondary effect.

Cellular function of genes controlled by Tet

Tet protein is detected in embryos from blastoderm stage onwards and is most strongly expressed

in neuronal tissues and also in cardiac and muscle precursor cells. In third instar larvae, the gene is

strongly expressed in the brain and neuronal cells in imaginal discs [10]. It was therefore important to

assess if our molecular analyses would agree with this expression pattern and if target genes are

associated with neuronal functions. We performed Gene Ontology (GO) analyses of the genes

identified via ChIP-seq as well as of the genes encoding the 5hmrC-modified mRNAs that were

identified in our hMeRIP-seq analyses in the embryo and the LBF (Fig. S5 A-D). The genes identified

in both embryonic and larval samples through both ChIP-seq and hMeRIP-seq all show enrichment for

genes involved in axon guidance. When we looked at the GO terms of transcripts that showed a

reduction of the 5hmrC modification in Tet^{null} samples, axon guidance genes were highly represented, in

fact, GO terms of transcripts showing > 1.4 times reduction of the modification in Tet^{null} samples

identified mostly genes associated with neuronal functions (see highlighted genes in Fig. 6C).

It is striking that in our two very different experimental approaches, ChIP-seq and hMeRIP-seq we

identified genes with overlapping functions (Fig. S5 A-D). The importance of our results is also

underlined by the observation that of the transcripts that show > 1.4 times reduction of 5hmrC levels in

Tet^{null} samples, 40% were derived from genes that also have at least one Tet DNA-binding site (Fig.

6C). In LBF samples, 43% of all the transcripts that show 5hmrC modification are derived from genes

that have been shown to bind Tet (Fig. S4A). In embryo samples, 29% of all the transcripts that showed

5hmrC modification are derived from genes that have been shown to bind Tet (Fig. S4B). Further, 29%

of modified transcripts in embryos and 37% of modified transcripts in LBF show 5hmrC marks at both

developmental stages (Fig. S4C). An example of the experimental IGV tracks of all our results for a

gene in the larval CNS and the embryo are shown in Fig.6D and Fig.S6A, respectively.

These analyses confirm that Tet-dependent 5hmC is often found on mRNAs derived from genes

that show Tet binding. Notably, close to 50% of transcripts that show a reduction in the 5hmrC mark in

Tet^{null} tissues are derived from Tet-target genes. However, the levels of these mRNAs are generally

unaffected by the loss of Tet suggesting that the 5hmrC modification does not affect steady state level

of mRNAs but other aspects of mRNA function such as translation or localization.

Tet target genes

We used the results above to identify Tet-target genes and sought to determine whether the

phenotypic effects of the loss of Tet's activity were derived from its inability to regulate target mRNAs

[5,10]. We looked for genes that are 1. active in the nervous system where Tet is enriched and 2. showed

Tet protein binding to DNA, and 3. whose mRNA showed a reduction in 5hmrC in Tet^{null} animals. Axon

guidance genes as a group frequently showed Tet-DNA-binding and 5hmrC mRNA modification by Tet

(Figure 6D). Among the genes that fulfilled the three criteria were two well-studied genes that function

in axon guidance, *robo2* and *slit* (Fig. S7). The Slit/Robo signaling pathway is required for axonal

pathfinding and the bilateral organization of the CNS in both vertebrates and invertebrates [17]. Robo

proteins are transmembrane receptors on axonal growth cones for the secreted Slit ligands. Glial cells

present at the midline secrete Slit and signaling between Robo and Slit is essential to inhibit midline

crossing of axons through commissures via repulsion [18]. Slit also has previously been implicated as a

target of Tet activity in midline glia [12]. We examined axonal pathfinding in the embryonic, ventral nerve

cord (VNC) and reasoned that if Tet impinges upon the levels of Robo2 and/or Slit, we should observe

midline defects in Tet^{null} animals like those seen in *robo2* or *slit* mutant embryos. Gross CNS

commissural structure is maintained in *Tet^{null}* embryos (Fig. 7B', HRP), however, examination of

neuronal subpopulations within the longitudinal neuropils indicates frequent pathfinding defects. A well

described subpopulation, Fas2+ neurons, exhibit extensive midline crossing of growth cones in these

Tet^{null} embryos (Fig. 7B, arrows; Table S1). Additionally, the most lateral of the Fas2+ longitudinal tracks

are often incomplete or absent (Fig. 7B, 46%-arrowheads). A second subpopulation of neurons

expressing Connectin also appears to be altered in *Tet^{null}* VNCs and fails to populate one of the

longitudinal tracks compared to wild type (Fig S7B; arrows). These phenotypes are strikingly similar to

the axonal pathfinding defects seen in *robo2* embryos with Tet's effects being slightly more severe (Fig.

7B and C and table S1) [18]. We sought to determine whether the reduction of Tet-mediated 5hmrC

deposition on the *robo2* or *slit* mRNAs resulted in mRNA species with reduced activity or potential for

expression. Thus, we examined genetic interactions between Tet and the Slit/Robo signaling pathway

in *Tet^{null}* embryos lacking one copy of *robo2* or *slit*. We additionally examined Robo1, a gene that is also

involved in midline repulsion but is not 5hmrC modified. Decreasing the dose of Robo2 or Robo1 in a

Tet^{null} background has little effect on Fas2+ axonal pathfinding in comparison to *Tet^{null}* alone (table S1).

The failure to see an effect with Robo2 may stem from the observation that the levels of midline

crossing in *Tet^{null}* embryos exceeds that seen for *robo2*^{null} embryos (Table S1 and [19]. However, reducing

the gene dose of Slit by half enhances the midline crossing of Fas2+ neurons in *Tet^{null}* embryos (Table

S1; Figure 7; 48% vs 32% *Tet^{null}*), whereas heterozygous *slit* embryos show midline crossing in < 1% of

segments (Fig 7D). Moreover, *Tet^{null}* mutant animals appear to be sensitized towards midline crossing

in general when lacking full *slit* function. Notably, the commissures (red arrowheads, 7E') are poorly

defined likely due too many axons inappropriately transiting the midline.

Given that these *robo2* or *slit* encode mRNAs that carry the 5hmrC mark and a reduction of that

mark in the *Tet^{null}* background, we expected Tet to potentially control their protein levels (Fig. S7).

Indeed, both proteins were clearly reduced in brain extracts from *Tet^{null}* larvae relative to wt (Fig. 7F and

7F'). These results support the idea that the function of Tet-dependent 5hmrC modification is to control

high levels of translation of specific target mRNAs and that in the context of embryonic axonal

pathfinding Tet provides an additional, novel layer of regulation of the medically important Slit/Robo

pathway.

Based on all our results we suggest the model shown in Figure 7G, we propose that Tet binds,

possibly as a complex to DNA binding sites mediated by its DNA-binding domain. The Tet binding sites

are preferentially located at promoter regions of genes that also show H3K4me3, generally accepted as

a mark of active transcription. We further postulate that Tet binds nascent mRNA in cooperation with

associated proteins (RNA-binding proteins, and with a so far unidentified RNA methyltransferase) to set

the 5hmC mark. The 5hmrC marked mRNAs are then exported from the nucleus and recognized by a

reader protein that will control the efficient loading of the modified mRNAs onto polysomes, where the

mRNAs are proficiently translated.

While several aspects of this model need to be investigated our results provide a consistent

framework of how Tet and Tet-dependent RNA modifications may function in controlling gene

expression. Recently, mutations in human Tet3 have been shown to cause neurodevelopmental delays.

It will be interesting to investigated if 5hmC RNA modification is deficient in the affected patients [20].

301   Discussion

In our previous study we investigated if Tet proteins, that are well known as 5-methylcytosine

(5mC) hydroxylases catalyzing the change from 5mC to 5hmC in DNA, can have a similar function in

RNA [5]. For these molecular studies we mainly used *Drosophila* S2 cells as source material. In the

present study we used animal sources, embryos, and larval brain tissues to investigate the function of

Tet in modifying mRNA *in vivo*. We also wanted to delineate the molecular and cellular processes for

which the modification is required, and to identify in vivo targets of the Tet protein.

Our results demonstrate that Tet protein binds to distinct genes, functions in modifying mRNAs,

and that this modification modulates translational output of the mRNAs. We used our molecular results

to identify Tet target genes. We selected genes that, 1. contain promoter proximal Tet-binding site(s)

that overlap with H3K4me3 modifications, 2. whose mRNA showed 5hmrC modifications that were

reduced in *Tet^null* neuronal tissues, and 3. whose mRNA levels displayed negligible changes in *Tet^null*

neuronal tissues.

We found that these target genes were most often associated with axonal growth and

pathfinding. Two such genes, *robo2* and *slit*, were selected because they fulfill the conditions outlined

above and are members of a conserved set of cell-signaling molecules responsible for controlling the

activity of axonal growth cones of the developing CNS in vertebrates and invertebrates [21]. Robo

receptors interact with a Slit ligand to specify overall axonal growth cone repulsion. Phenotypic analysis

of the developing CNS in Tet-deficient animals indicates a specific requirement for Tet in the proper

patterning of the CNS; *Tet^null* embryos showed a similar CNS phenotype to Robo2 or Slit deficient

animals. Indeed, in the absence of Tet and the ensuing reduction of the 5hmrC modification of the

mRNAs, levels of the corresponding proteins, Robo2 and Slit, are reduced, resulting in aberrant axonal

pathfinding and other defects in nervous system patterning [10,12].

325   Tet controls the 5hmrC modification on mRNA

In mass spectrometry experiments we determined that 5hmrC is highly enriched in polyA$^+$ RNA

confirming our previous dot blot results. This modification is significantly rarer than other well-studied

mRNA modifications, such as 5mrC or 6mA (Fig. 1) [5,22]. Coupled with our transcriptomic analyses,

these observations show that the modification resides on a subset of mRNAs. Because Tet is

expressed in Drosophila almost exclusively in nerve cells, we determined the levels of 5mrC and

5hmrC in wild type 0-12 h embryos (where Tet is highly expressed) and in larval brains. We found that

5mrC levels are about two orders of magnitude higher than 5hmrC levels (~$2x10^5$ 5mrC and ~$2x10^7$

5hmrC in larval brains), and therefore detecting 5hmrC is not trivial.

334   The presence of 5hmrC is notably reduced (~ 5 fold) in *Tet^null* samples. Our results are

335   consistent with the *Drosophila* Tet enzyme being responsible for this 5hmrC modification (Fig. 1 and

336   S1). However, since we detect ~20% of the wild type 5hmrC levels in mutant tissues that lack Tet

337   completely, we assume that an additional hydroxymethyltransferase(s) that can modify 5mrC do exist in

338   the *Drosophila* genome. The existence of additional enzyme(s) contributing to mRNA

339   hydroxymethylation has also been postulated in mouse ESCs [14].

340   Our mass spectrometry findings and the results from our hMeRIP-seq experiments on larval

brain fractions (LBF) and embryos are consistent with what has been previously reported for *Drosophila*

tissue culture cells and for ESCs (Fig. 1,4,5 and S1, S3) [14]. We identified ~3000 5hmrC peaks in ~1500

transcripts in S2 cells [5]. In ESCs the number of peaks was 1633 in 795 transcripts [14]. In our *in vivo*

experiments we identified 1815 peaks in 1402 transcripts in embryos, and 3711 peaks on 1776

transcripts in LBF. Of the modified transcripts in embryos 37% were also identified as modified

transcripts in the LBF. In all samples the modification peaks centered around a UC-rich consensus

motif (Fig. S3). The consistency of the mapping results of the 5hmrC modifications in *Drosophila* tissue

culture cells, embryos, larval brain fraction, and ESCs underlines the probable conserved function of

Tet across the species.

350   The distribution of the 5hmrC peaks on transcripts derived from LBF are found at similar levels

in all parts of the transcripts, the UTRs, the coding region, and introns. However, in *Tet^null* LBF

significantly more peaks are reduced in the CDS and introns than in the UTRs (Fig. 5C). This

observation suggests that Tet may target coding sequences and introns specifically. We do not yet

understand if modifications in different parts of the transcripts have diverse functions and if they may be

controlled by additional enzyme(s).

Drosophila Tet's DNA binding activity

358   We found that in both embryos and in LBFs, Tet recognizes a DNA motif similar to the motif

bound by Tet1 in vertebrate ESCs (Fig. S2D) [15,23]. A majority of these peaks are associated with coding

regions and are frequently found at the promoter. Almost 50% of the peaks overlap with the H3K4me3

mark, an indication that the genes are actively transcribed. The distribution of Tet-binding peaks and

the overlap with the H3K4me3 mark agree well with the localization of the Tet-DNA-binding domain on

salivary gland chromosomes confirming that the binding sites are found almost exclusively in

euchromatin and are distributed on all 4 chromosomes (Fig. S2A).

365   We propose that the selection of target RNAs modified by Tet is at least in part facilitated by

Tet's DNA-binding of specific genes. The concurrence of Tet-DNA binding peaks on genes that also

showed Tet-dependent 5hmrC modifications of their mRNA is consistent with this idea. The majority of

the genes that show Tet binding and modified mRNAs are divergent in both tissues indicating that in

addition to a conserved function of Tet in different neuronal cells, Tet also has a tissue-specific or

possibly even cell-type-specific function.

Identifying Tet target mRNAs

Tet is highly expressed in nervous tissues and the loss of Tet function leads to abnormal

neuronal functions such as defects in larval locomotion or abnormalities in the circadian rhythm

(PLOS). Our immunoprecipitation of 5hmrC-modified RNAs identified 1775 genes in larval brain

fractions. 45 % (798) showed a significant decrease in the overall 5hmrC peaks in a Tet^{null} background.

Of the genes with reduced 5hmrC marks, 44% showed Tet-DNA binding. Notably, the mRNAs in which

the reduction of the 5hmrC mark was seen were mostly associated with genes that function in different

aspects of nerve cell development. First among them are axon outgrowth genes that were also

identified in the GO-term analysis as abundant gene categories associated with Tet binding sites and

mRNAs carrying the 5hmrC mark (Fig. 6D, S5).

Our initial examination of the developing embryonic ventral nerve cord (VNC) in *Tet* mutants

identified subtle defects in CNS patterning. We then examined subsets of VNC neurons using

antibodies to Fas2 and Connectin (Fig 7B, B' and S7B, B') guided by our molecular results. Overall

commissural structure is maintained in Tet^{null} embryos, however neurons expressing Fas2 show a

failure of the midline to repel axon crossing effectively. And so, we looked among the Tet mRNA targets

with known functions in axon guidance and found that both *slit* and *robo2* mRNAs were represented.

Both genes have Tet-binding sites near the TSS, their mRNA is modified, and the modification is

reduced in Tet^{null} LBF, while their mRNA levels are not significantly changed (Fig. S7). Comparison of

the CNS in Tet^{null} and $robo2^{null}$ embryos identified a set of overlapping phenotypes with high frequency

midline crossing defects of Fas2+ neurons, as well as discontinuities in the most lateral, longitudinal

Fas2 and Connectin axonal tracts (for description of embryonic nerve cord see [18]. Notably, these tracts

correspond to neurons which express the Robo2 protein [24,25]. Fas2+ neurons of *slit/+* embryos show no

phenotype but removing one copy of *slit* in Tet^{null}/Tet^{null} embryos had a strong enhancing effect of the

axon guidance phenotype, consistent with a close interaction of the two genes (Fig. 7, Table 1).

The overlapping phenotypes of Tet, *robo2 and slit,* together with the molecular data that

identified Robo2 and Slit as Tet targets, prompted us to investigate if Robo2 and Slit protein expression

was affected by the loss of Tet. Indeed, in Western blots from Tet^{null} larval brain extracts both Robo2

and Slit protein levels were strongly reduced (Fig. 7F, F'), indicating that Tet's profound consequences

on VNC patterning occurs, at least in part through the control of expression of the Robo2 and Slit

proteins. As Robo2 and slit mRNA levels are not changed in Tet^{null} LBF (Fig. S9), we suggest that the

Tet-dependent 5hmrC modification positively controls the level of translation of the two mRNAs. While

we have not investigated the protein levels of additional Tet-targets, we expect that Tet controls protein

levels through the 5hmC modification of many target mRNAs.

Which step in RNA processing leading to mRNA translation is affected in Tet^{null} animals will

have to be elaborated. Based on our previous results, that showed 5hmrC modified RNAs found on

polysomes, at least one possibility is that the 5hmrC modification facilitates the loading of the mRNAs

on ribosomes [5].

Tet and the 5hmrC modification function in mRNA processing of specific neuronal mRNAs

guaranteeing that the modified mRNAs are translated efficiently, and at levels necessary for normal

neuronal function, thus adding an additional level of control of gene expression.

Materials and Methods

Drosophila Genetics

All flies were reared at 25°C and kept on standard medium. The mutant Tet alleles are described in [5,10]; the wild-type allele used in all experiments is w^{1118}. Stocks utilized to examine genetic interactions with Tet were sli²/CyO, and robo1⁴/CyO (Bloomington Drosophila Stock center), and robo2^x123/CyO [26]. The material used for all whole-genome analysis was either hand dissected third instar larval brains, or, because some experiments necessitated a large input, dissected anterior parts of larvae including the 3 anterior abdominal segments that contain the brain besides other tissues such as imaginal discs, salivary glands, mouth parts and epidermis. Because Tet is highly expressed in the brain and the nerve cell in discs, but not in the other tissues, we call this the Larval Brain Fraction, LBF. Brains and larvae from wt and Tet-GFP third instar larvae were dissected in cold-PBS supplemented with protease inhibitor, snap frozen on dry ice, and stored at -80°C.

Immunohistochemistry and Imaging

The following antibodies were used for immunolabelling of late stage embryos and chromosomal preparations: mouse anti-Fas2 (Developmental Studies Hybridoma Bank, DSHB), rabbit anti-HRP (Jackson Immunoresearch), mouse anti-Connectin (DSHB), rabbit anti-dsRED (Invitrogen), rabbit and mouse anti-GFP (Invitrogen), mouse anti-H3K4me3 (Invitrogen). Secondary antibodies were purchased from Invitrogen. DNA was labeled with DAPI (Invitrogen). Embryos were collected and fixed via a formaldehyde/MeOH method [10]. Polytene chromosome preparations and staining were performed as in Karachentsev et al. [27]. Images of the ventral nerve cord were obtained using a Leica SP8 using a 40x Objective. Fas2 and HRP labeled embryos were imaged and typically contained 8-10 hemisegments. Hemisegments were examined for midline crossing and in some instances the presence or integrity of the most lateral Fas2+ longitudinal track. Similar imaging and analysis were performed on Connectin/HRP labeled embryos.

LC-MS/MS for 5mC and 5hmC detection and quantification

Mass spectrometry analysis was performed as described previously [28]. Briefly, 3 µL of 10× buffer (500 mM Tris-HCl, 100 mM NaCl, 10 mM $MgCl_2$, 10 mM $ZnSO_4$, pH 7.0), 2 µL (180 units) of S1 nuclease, 2 µL (0.001 units) of venom phosphodiesterase I and 1 µL (30 units) of CAIP were added to 1 µg of mRNA from *Drosophila* wild type and Tet-deficient larval brains, respectively (in 22 µL of H_2O). The mixture (30 µL) was incubated at 37°C for 4 h. The resulting solution was three times extracted with chloroform. The upper aqueous phase was collected and passed through a solid phase extraction cartridge filled with 50 mg of sorbent of graphitized carbon black to remove the salts. The eluate was then dried with nitrogen at 37°C for subsequent chemical labeling and LC-ESI-MS/MS analysis by an AB 3200 QTRAP mass spectrometer (Applied Biosystems, Foster City, CA, USA).

Embryo and Larval Tet ChIP-seq

0-12h embryos were collected, processed, and chromatin was prepared according to Yad et al. [29], except lysates were sonicated on a Covaris S2 sonication device (intensity 8, duty cycle 20%, cycle burst 200) for 30 minutes at 4˚C to reach fragments ranging from 150–500 bp and then centrifuged at 20,000g at 4˚C for 1 minute. Supernatants were collected and centrifuged again for 15 minute to remove debris. Chromatin samples were then snap frozen in dry ice and stored at -80˚C until immunoprecipitation in triplicets. All buffers contained cOmplete EDTA-free protease inhibitor cocktail (Roche).

For the larval brain fraction (LBF), 300 frozen larval heads were thawed on ice and 1 ml of NU-1 buffer (5 mM HEPES-KOH pH 7.9, 5 mM MgCl$_2$, 0.1 mM EDTA pH 8.0, 0.5 mM EGTA pH 8.0, 350 mM sucrose, 1mM DTT). 1% formaldehyde was added to NU-1 buffer before use. Samples were homogenized immediately at room temperature using Dounce with a loose pestle 30 times without foaming for 15 minutes. Samples were filtered first through BD Falcon Cell Strainer 70 μm (Cat No.352350) followed by 50 μm Filcon (Cat No. 340603). Samples were quenched with freshly prepared 125 mM glycine incubated for 5 minutes at room temperature on a shaker and transferred to ice for 5 minutes. Samples were centrifuged at 4000 g at 4˚C for 5 minutes. The pellet was washed twice with 1 ml cold PBS and resuspended in 350 μl chilled sonication buffer (50mM HEPES-KOH pH 7.9, 140 mM NaCl, 1mM EDTA pH 8.0, 1% Triton X-100, 0.1% sodium deoxycholate, 1% SDS) and incubated for 20 minutes at 4˚C. Lysates were sonicated as described above and chromatin was stored at - 80˚C until immunoprecipitation.

Chromatin Immunoprecipitation

Chromatin samples were thawed on ice and pre-cleared for 15 minutes by rotation in 25 μl of pre-washed binding control magnetic agarose beads (Chromotek). Chromatin was diluted ten-fold in sonication buffer without SDS. 1% of the diluted lysate was recovered and used as input. Diluted chromatin was incubated with 25 μl of pre-washed GFP-Trap MA beads (Chromotek) and rotated at 4˚C overnight. Lysates were washed on magnetic stand with 1 ml each low salt RIPA buffer (140 mM NaCl, 1mM EDTA pH 8.0, 1% Triton X-100, 0.1% sodium deoxycholate, 10mM Tris-HCl pH 8.0) (5 times), high salt RIPA buffer (500 mM NaCl, 1mM EDTA pH 8.0, 1% Triton X-100, 0.1% sodium deoxycholate, 10mM Tris-HCl pH 8.0) (2 times), LiCl buffer (250mM LiCl, 1mM EDTA pH 8.0, 0.5% IGEPAL CA-630, 0.5% sodium deoxycholate, 10mM Tris-HCl pH 8.0) (1 time), TE buffer (10mM Tris-HCl pH 8.0, 1mM EDTA pH 8.0) (1 time). All buffers contained cOmplete EDTA-free protease inhibitor cocktail (Roche).

ChIP DNA was eluted by shaking 2 hours at 37˚C with 100 μl of elution buffer (1% SDS, 50mM NaHCO$_3$, 10μg/ml RNaseA), then 4 hours with 0.2μg/ml proteinase K. Beads were concentrated on

486 magnet and elute was recovered. Samples were de-crosslinked overnight at 65˚C. Inputs were
487 processed like ChIP samples. DNA was purified by phenol/chloroform/isoamyl alcohol followed by SPRI
488 select beads (Beckman Coulter) and DNA concentration was measured with Qubit fluorometer (Thermo
489 Fisher).
490
Embryo Tet ChIP-seq library preparation and sequencing
NGS Libraries were made from eluted DNA using the NEBNext Ultra II DNA Library Prep kit
(New England Biolabs) according to the manufacturer's protocol. Briefly, 20 ng of DNA fragments were
end-repaired and the blunt, phosphorylated ends were treated with Klenow DNA polymerase and dATP
to yield a 3′ A base overhang for ligation of Illumina adapters. After adapter ligation, DNA was PCR
amplified with indexed primer for 12 cycles. Libraries were size-selected using Ampure XP beads
(Beckman Coulter) to remove adapter dimers. DNA was quantified by fluorometry with the Qubit 2.0
(Thermo Scientific) and DNA integrity was assessed with a Fragment Analyzer (Agilent). The libraries
were pooled and sequenced on the NextSeq 500 platform using 75 bp single end sequencing
according to manufacturer's protocol using Reagent v.2.5 at the Waksman Institute Genomics Core.
Coverage ranged from 30 million to 60 million tags per ChIP-seq sample.
502
503 **Larva Tet ChIP-seq library preparation and sequencing**
504   ACCEL-NGS® 1S plus DNA library kit was used to prepare indexed libraries from IP and input
505 DNA. Libraries were pooled respecting equimolarity. Sequencing was performed on Illumina MISeq
506 sequencer in 150 bp paired-end reads.
507
Embryo Tet ChIP-seq data analysis
Raw reads were trimmed using cutadapt v2.0 [30] to remove adapter and low-quality reads. The
processed reads were mapped to the *Drosophila melanogaster* BDGP6 (dm6) reference genome from
Ensembl release 88 using the BWA version 0.7.5-r404 for Chip-seq [31]. For analysis, only unique reads
with mapping quality >20 were accepted. Further, redundant reads with identical coordinates were
filtered out. Aligned reads were processed by Model-based Analysis of ChIP-seq (MACS2) [32] using
Input ChIP DNA as control. For peak calling the MACS2 'callpeak' function was used (-p 1e-2 -g
1.2e+08 -B --nomodel –ext size 147 –SPMR) for each replicate vs. control input. Peaks were selected
using the following criteria: p-value <10e-5, fold enrichment over control greater than 10 and a minimal
number of reads higher than 50. Bedtools (version v2.24.0) [33] was used to identify overlapping peaks in
replicates. A sliding window of 50, 100, 150, 200, 250 and 300 bp around the peak summit (base
position of maximum enrichment) was used to determine best range for overlapping peaks. The
number of overlapping peaks saturated around window size of 250 bp. Thus, for downstream analysis,
windows size of 250 bp was used to identify overlapping peaks in replicates. The Integrated Genomics

Viewer (IGV) [34] was used for visualization of ChIP-seq data sets. For visualization in IGV, bigwig peak

files were generated using "bdgcmp" function in MACS2 with option "-m logFE -p 0.00001". Peaks were

annotated using the "annotatePeaks.pl" feature of HomerTools [35] with default settings and gtf was

obtained from of Ensembl dm6 release 88. De novo motif discovery was carried out on all intersecting

peaks of Tet ChIP-seq. DNA sequences (FASTA) were generated from chromosome coordinates

produced by peak detection and windowing using the BEDTools. De novo motif analysis was performed

using MEME-ChIP [36]. Gene ontology (GO) analysis was done using Database for Annotation,

Visualization and Integrated Discovery (DAVID) [37,38]. Binding profile within gene body was generated

using deepTools2 with computeMatrix and plotProfile functions [39].

H3K4me3 ChIP-seq public datasets and analysis

Embryo and larva H3K4me3 ChIP-seq data were obtained from the modENCODE project

(GEO: GSE16013) [40]. The analysis was carried out from raw data following the same approach

described for Tet ChIP-seq. The overlapping of Tet-ChIP seq peaks and H3K4me3 was computed

using BEDTools [33].

Larva Tet ChIP-seq data analysis

Tet-Chip sequencing data were pre-processed using the following steps: the raw sequencing

data were first analysed with FastQC (Andrews, 2010,

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Low-complexity reads were removed with

the AfterQC tool [41] with default parameters and Trimmomatic [42] with default parameters was used to

remove adapter sequences. The resulting fastq data were again analysed with FastQC to ensure that

no further processing was needed. Pre-processed reads were then mapped against the *Drosophila*

reference genome (BDGP6.28) with the bowtie2 algorithm [43] using the ensembl reference

transcriptome (version 100). Tet-binding peak regions were identified by applying the MACS2 peak-

calling tool [32] to immunoprecipitated (IP) samples, using their input counterpart to estimate background

noise (q-value < 0.05). It is worth noting that the "expected genome size" MACS2 parameter was set as

the *Drosophila* genome length excluding 'N' bases (*i.e.* 142 573 024 bp), and summit positions were

identified using the MACS2 "-call-summits" option. To avoid identifying extremely large peak regions,

the peaks were resized to 100 bp on both sides of the identified summit. Binding profile within gene

body was generated using deepTools2 with computeMatrix and plotProfile functions [39].

HydroxyMethylated RNA Immunoprecipitation sequencing (hMeRIP-seq)

0-12h embryos were collected, immediately frozen on dry ice, and stored at -80°C until RNA

purification. The larval brain fraction (LBF), was dissected, immediately frozen on dry ice and stored at -

80°C until RNA isolation. The RNA immunoprecipitation was performed essentially as described in

Dominissi et al [44]. Briefly, total RNA was isolated using RNeasy Maxi Kit (Qiagen). For each sample 1

mg of total RNA (1 µg/µl) was divided into batches of 45µg and incubated at 94°C in fragmentation

buffer (100 mM Tris-HCl pH7.0, 100 mM $ZnCl_2$) for 40 seconds. Fragmented RNA batches were

pooled, and ethanol precipitated at -80˚C overnight. RNA samples were washed with 75% ethanol and

resuspended in RNase-free water. Fragmentation efficiency was checked on a Bioanalyzer RNA chip

(Agilent). RNA fragments were denatured by heating at 70°C for 5 minutes, then chilled on ice for 5

minutes. For immunoprecipitation, RNA samples were incubated overnight at 4°C with 12.5 µg of anti-

5-hmC antibody (Diagenode rat monoclonal MAb-633HMC) or without antibody as negative control in

IP buffer (750 mM NaCl, 50 mM Tris-HCl pH7.4, 0.5% IGEPAL CA-630, RNasin 400 U/ml and RVC 2

mM). 60 µl of equilibrated Dynabeads Protein G (Life Technologies) were added to the samples and

incubated at 4°C for 2.5 hours. The magnetic stand beads were washed with 1 ml IP buffer for 5

minutes three times. To elute immunoprecipitated RNA, 1 ml TriPure Reagent (Roche) was added,

mixed thoroughly and centrifuged at room temperature for 5 minutes. Aqueous phase was recovered,

and equal amount of chloroform was added, vortexed and aqueous phase was collected after

centrifugation and ethanol precipitated at -80˚C overnight. RNA was resuspended in nuclease free

water and used for library preparation. All buffers contained cOmplete EDTA-free protease inhibitor

cocktail (Roche).

hMeRIP-seq library preparation and sequencing

Library preparation was done with the TruSeq ChIP Sample Prep Kit (Illumina) after reverse

transcription of pulled-down RNA and synthesis of a second strand (NEB) by Next mRNA second

strand synthesis module (NEB)). Briefly, 5 to 10 ng dsDNA was subjected to 5' and 3' protruding end

repair. Then, non-templated adenines were added to the 3' ends of the blunted DNA fragments. This

last step allows ligation of Illumina multiplex adapters. The DNA fragments were then size selected in

order to remove all unligated adapters and to sequence 200-300-bp fragments. 18 cycles of PCR were

carried out to amplify the library. DNA was quantified by fluorometry with the Qubit 2.0 and DNA

integrity was assessed with a 2100 bioanalyzer (Agilent). 6 pM of DNA library spiked with .5% PhiX

viral DNA was clustered on cBot (Illumina) and then sequenced on a HiScanSQ module (Illumina).

hMeRIP-seq data analysis

The processed reads were mapped to the reference genome Drosophila melanogaster BDGP6

(dm6) from Ensembl by using Hisat2 (version 2.1.0) for RNA seq and hMeRIP seq [45]. To analyze gene

expression, HTSeq framework, version 0.5.3p9, was used to count the aligned reads in genes [46]. Mode

"union" and mapping quality cut-off 20 were used for our analysis. Count-table was normalized so that

all samples have the same level of total mapped reads. DEseq2 was used to identify differentially

expressed genes [47]. Cufflinks v2.2.1 was applied to calculate the rpkm values [48,49]. A gene was

considered as significantly changed when fold change >=2 or <= -2 and adjusted p value < 0.05.

"SplitNCigarReads" funciton in GATK (version 3.3-0) (https://gatk.broadinstitute.org/) were used to split

reads that contain Ns in their cigar string (e.g., spanning splicing events in hMeRIP-seq data). "rmdup"

function of samtools (version 1.3.1) were used to remove a duplicate mapping of reads. Then the same

peak calling procedure as ChIP seq data analysis was performed to call peaks of hMeRIP-seq data.

The peaks of hMeRIP-seq were selected using P-value < 10e-5. Peaks of hMeRIP-seq were

considered as reduced when the normalized hMeRIP-seq signal in control samples was at least 1.4-

fold change higher than the signal in Tet depleted samples. The fold change and P-value were

calculated using "limma" package in R [50].

Western blot

One hundred third instar larval brains from wild type or *Tet^null* were dissected and immediately

frozen on dry ice. Total protein was isolated from these brains using RIPA buffer and 75 ug of the total

protein was loaded to each well. Slit antibody (DSHB, C555.6D, Spyros Artavanis-Tsakonas) was used

at 1: 200 dilution and Robo2 antibody [51] was used at 1: 1000 dilution. The western blot signals were

detected using IRDye 800CW Infrared Dyes conjugated secondary antibody in LICOR Odyssey CLx

imaging system. Signals were quantified using LICOR Image Studio Lite software. See Figure S8 for

unprocessed western blot exposure.

References

Roundtree, I. A., Evans, M. E., Pan, T. & He, C. Dynamic RNA Modifications in Gene Expression Regulation. *Cell* **169**, 1187-1200 (2017). https://doi.org:10.1016/j.cell.2017.05.045

Boccaletto, P. *et al.* MODOMICS: a database of RNA modification pathways. 2021 update. *Nucleic Acids Res* **50**, D231-D235 (2022). https://doi.org:10.1093/nar/gkab1083

Schaefer, M. R. The Regulation of RNA Modification Systems: The Next Frontier in Epitranscriptomics? *Genes (Basel)* **12** (2021). https://doi.org:10.3390/genes12030345

Gao, Y. & Fang, J. RNA 5-methylcytosine modification and its emerging role as an epitranscriptomic mark. *RNA Biol* **18**, 117-127 (2021). https://doi.org:10.1080/15476286.2021.1950993

Delatte, B. *et al.* RNA biochemistry. Transcriptome-wide distribution and function of RNA hydroxymethylcytosine. *Science* **351**, 282-285 (2016). https://doi.org:10.1126/science.aac5253

Durdevic, Z. *et al.* Efficient RNA virus control in Drosophila requires the RNA methyltransferase Dnmt2. *EMBO Rep* **14**, 269-275 (2013). https://doi.org:10.1038/embor.2013.3

Tahiliani, M. *et al.* Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* **324**, 930-935 (2009). https://doi.org:10.1126/science.1170116

Fu, L. *et al.* Tet-mediated formation of 5-hydroxymethylcytosine in RNA. *J Am Chem Soc* **136**, 11582-11585 (2014). https://doi.org:10.1021/ja505305z

Tan, L. & Shi, Y. G. Tet family proteins and 5-hydroxymethylcytosine in development and disease. *Development* **139**, 1895-1902 (2012). https://doi.org:10.1242/dev.070771

Wang, F. *et al.* Tet protein function during Drosophila development. *PLoS One* **13**, e0190367 (2018). https://doi.org:10.1371/journal.pone.0190367

Dunwell, T. L., McGuffin, L. J., Dunwell, J. M. & Pfeifer, G. P. The mysterious presence of a 5-methylcytosine oxidase in the Drosophila genome: possible explanations. *Cell Cycle* **12**, 3357-3365 (2013). https://doi.org:10.4161/cc.26540

Ismail, J. N., Badini, S., Frey, F., Abou-Kheir, W. & Shirinian, M. Drosophila Tet Is Expressed in Midline Glia and Is Required for Proper Axonal Development. *Front Cell Neurosci* **13**, 252 (2019). https://doi.org:10.3389/fncel.2019.00252

Shen, Q. *et al.* Tet2 promotes pathogen infection-induced myelopoiesis through mRNA oxidation. *Nature* **554**, 123-127 (2018). https://doi.org:10.1038/nature25434

Lan, J. *et al.* Functional role of Tet-mediated RNA hydroxymethylcytosine in mouse ES cells and during differentiation. *Nat Commun* **11**, 4956 (2020). https://doi.org:10.1038/s41467-020-18729-6

Wu, H. *et al.* Dual functions of Tet1 in transcriptional regulation in mouse embryonic stem cells. *Nature* **473**, 389-393 (2011). https://doi.org:10.1038/nature09934

16   Jones, P. A. & Liang, G. Rethinking how DNA methylation patterns are maintained. *Nat Rev Genet*
10, 805-811 (2009). https://doi.org:10.1038/nrg2651

17   Blockus, H. & Chedotal, A. Slit-Robo signaling. *Development* **143**, 3037-3044 (2016).
https://doi.org:10.1242/dev.132829

18   Simpson, J. H., Kidd, T., Bland, K. S. & Goodman, C. S. Short-range and long-range guidance by
slit and its Robo receptors. Robo and Robo2 play distinct roles in midline guidance. *Neuron* **28**,
753-766 (2000). https://doi.org:10.1016/s0896-6273(00)00151-3

19   Evans, T. A., Santiago, C., Arbeille, E. & Bashaw, G. J. Robo2 acts in trans to inhibit Slit-Robo1
repulsion in pre-crossing commissural axons. *Elife* **4**, e08407 (2015).
https://doi.org:10.7554/eLife.08407

20   Beck, D. B. *et al.* Delineation of a Human Mendelian Disorder of the DNA Demethylation
Machinery: TET3 Deficiency. *Am J Hum Genet* **106**, 234-245 (2020).
https://doi.org:10.1016/j.ajhg.2019.12.007

21   Gorla, M. & Bashaw, G. J. Molecular mechanisms regulating axon responsivenss at the midline.
Dev Biol **466**, 12-21 (2020). https://doi.org:10.1016/j.ydbio.2020.08.006

22   Dominissini, D. *et al.* Topology of the human and mouse m6A RNA methylomes revealed by m6A-
seq. *Nature* **485**, 201-206 (2012). https://doi.org:10.1038/nature11112

23   Yao, B. *et al.* Active N(6)-Methyladenine Demethylation by DMAD Regulates Gene Expression by
Coordinating with Polycomb Protein in Neurons. *Mol Cell* **71**, 848-857 e846 (2018).
https://doi.org:10.1016/j.molcel.2018.07.005

24   Simpson, J. H., Bland, K. S., Fetter, R. D. & Goodman, C. S. Short-range and long-range guidance
by Slit and its Robo receptors: a combinatorial code of Robo receptors controls lateral position. *Cell*
103, 1019-1032 (2000). https://doi.org:10.1016/s0092-8674(00)00206-3

25   Spitzweck, B., Brankatschk, M. & Dickson, B. J. Distinct protein domains and expression patterns
confer divergent axon guidance functions for Drosophila Robo receptors. *Cell* **140**, 409-420 (2010).
https://doi.org:10.1016/j.cell.2010.01.002

26   Santiago-Martinez, E., Soplop, N. H. & Kramer, S. G. Lateral positioning at the dorsal midline: Slit
and Roundabout receptors guide Drosophila heart cell migration. *Proc Natl Acad Sci U S A* **103**,
12441-12446 (2006). https://doi.org:10.1073/pnas.0605284103

27   Karachentsev, D., Sarma, K., Reinberg, D. & Steward, R. PR-Set7-dependent methylation of
histone H4 Lys 20 functions in repression of gene expression and is essential for mitosis. *Genes*
Dev **19**, 431-435 (2005). https://doi.org:10.1101/gad.1263005

28   Huang, W. *et al.* Formation and determination of the oxidation products of 5-methylcytosine in
RNA. *Chem Sci* **7**, 5495-5502 (2016). https://doi.org:10.1039/c6sc01589a

29   Ghavi-Helm, Y., Zhao, B. & Furlong, E. E. Chromatin Immunoprecipitation for Analyzing

Transcription Factor Binding and Histone Modifications in Drosophila. *Methods Mol Biol* **1478**, 263-

277 (2016). https://doi.org:10.1007/978-1-4939-6371-3_16

30   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *2011* **17**,

3 (2011). https://doi.org:10.14806/ej.17.1.200

31   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics **25**, 1754-1760 (2009). https://doi.org:10.1093/bioinformatics/btp324

32   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

https://doi.org:10.1186/gb-2008-9-9-r137

33   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics **26**, 841-842 (2010). https://doi.org:10.1093/bioinformatics/btq033

34   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24-26 (2011).

https://doi.org:10.1038/nbt.1754

35   Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-

regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-589 (2010).

https://doi.org:10.1016/j.molcel.2010.05.004

36   Machanick, P. & Bailey, T. L. MEME-ChIP: motif analysis of large DNA datasets. *Bioinformatics* **27**,

1696-1697 (2011). https://doi.org:10.1093/bioinformatics/btr189

37   Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large gene

lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57 (2009).

https://doi.org:10.1038/nprot.2008.211

38   Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional

annotation of gene lists (2021 update). *Nucleic Acids Res* **50**, W216-221 (2022).

https://doi.org:10.1093/nar/gkac194

39   Ramirez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis.

Nucleic Acids Res **44**, W160-165 (2016). https://doi.org:10.1093/nar/gkw257

40   Negre, N. *et al.* A cis-regulatory map of the Drosophila genome. *Nature* **471**, 527-531 (2011).

https://doi.org:10.1038/nature09990

41   Chen, S. *et al.* AfterQC: automatic filtering, trimming, error removing and quality control for fastq

data. *BMC Bioinformatics* **18**, 80 (2017). https://doi.org:10.1186/s12859-017-1469-3

42   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics **30**, 2114-2120 (2014). https://doi.org:10.1093/bioinformatics/btu170

43   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-

359 (2012). https://doi.org:10.1038/nmeth.1923

44   Dominissini, D., Moshitch-Moshkovitz, S., Salmon-Divon, M., Amariglio, N. & Rechavi, G.

Transcriptome-wide mapping of N(6)-methyladenosine by m(6)A-seq based on immunocapturing

and massively parallel sequencing. *Nat Protoc* **8**, 176-189 (2013).
https://doi.org:10.1038/nprot.2012.148
45   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907-915 (2019).
https://doi.org:10.1038/s41587-019-0201-4
46   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-throughput
sequencing data. *Bioinformatics* **31**, 166-169 (2015). https://doi.org:10.1093/bioinformatics/btu638
47   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* **15**, 550 (2014). https://doi.org:10.1186/s13059-014-0550-8
48   Trapnell, C. *et al.* Transcript assembly and quantification by RNA-Seq reveals unannotated
transcripts and isoform switching during cell differentiation. *Nat Biotechnol* **28**, 511-515 (2010).
https://doi.org:10.1038/nbt.1621
49   Trapnell, C. *et al.* Differential analysis of gene regulation at transcript resolution with RNA-seq. *Nat*
Biotechnol **31**, 46-53 (2013). https://doi.org:10.1038/nbt.2450
50   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res* **43**, e47 (2015). https://doi.org:10.1093/nar/gkv007
51   Rajagopalan, S., Vivancos, V., Nicolas, E. & Dickson, B. J. Selecting a longitudinal pathway: Robo
receptors specify the lateral position of axons in the Drosophila CNS. *Cell* **103**, 1033-1045 (2000).
https://doi.org:10.1016/s0092-8674(00)00207-5
737
738

Acknowledgements

We thank Cordelia Rauskolb, Bryce Nickels, and Michael Verzi for helpful comments on the manuscript, Premal Shah and John Fervante for discussions and suggestion, and Benjamin Rogers-Boehme for help with Figures. We also thank Barry Dickson for anti-Robo2 antibodies and Le Nguyen for expert fly food preparation and stock maintenance. Stocks obtained from the Bloomington Drosophila Stock Center (NIHP40OD018537) were used in this study. This work was supported by NIH grant, (R01 GM118404) to RS. HT is supported by Vietnam Education Foundation (VEF) and Charles and Johanna Busch Pre-doctoral Fellowships.

Figures and Legends

752 **Figure 1. 5hmrC is found in PolyA+ RNA and is controlled by Tet as measured by mass spectometry. A.**

753 5hmrC in total and polyA+ RNA isolated from S2 cells; **B.** 5mrC in total and polyA+ RNA isolated from S2 cells; **C.**

754 5hmrC in total RNA isolated from wild-type and *Tet^null* larval brain; **D.** 5mrC in total RNA isolated from wild-type

755 and *Tet^null* larval brain.

Figure 2. Genome-wide Tet protein binding sites in Drosophila larval brain fractions, Tet-ChIP-seq

analysis: A. Representative gene showing Tet binding peak at the promoter. Arrow indicates promoter

orientation; **B.** Genome wide distribution of Tet occupancy in larva brain fraction. The genomic regions (3'UTR,

5'UTR, exons, intergenic, introns, promoter-TSS transcription start sites, and TTS, transcription termination sites)

were defined based on RefSeq gene (dm6) annotations; **C.** Strength of Tet enrichment on fly genome counted as

peak score across the gene body plotted from 3413 peaks; **D.** Genome wide distribution of Tet binding sites

displayed as enriched sequence motif among 3413 peaks identified by de novo motif discovery in this study; **E.**

Binding profile of LBF Tet (red) and H3K4me3 (green) within the gene body ± 5kb; **F.** 36% of Tet occupied genes

on various genomic regions overlapped with the H3K4me3 mark; **G.** Promoter-associated Tet binding peaks on

40% of genes overlap with H3K4me3 marks.

Figure 3. Genome-wide Tet protein binding sites in Drosophila 0-12 hour embryos, Tet-ChIP-seq analysis:
A. Representative gene showing Tet binding peak at the promoter. Arrow indicates promoter orientation; **B.**
Genome wide distribution of embryo Tet ChIP-seq peaks in different genomic regions; **C.** Strength of Tet
enrichment on different genomic regions counted as peak score plotted from 5180 peaks; **D.** Enriched sequence
motif among 5180 embryo Tet ChIP-seq peaks identified by de novo motif discovery in this study; **E.** Binding
profile of embryo Tet (red) and H3K4me3 (green) within the gene body ± 5kb; **F.** 42% of Tet bound genes in
embryo have H3K9me3 modification; **G.** 51% of genes that show binding of Tet to the promoter that overlap with
H3K4me3; **H.** 27% of Tet bound genes in embryo also have Tet binding peaks in larva brain fraction.

Figure 4. **Transcriptome-wide distribution of 5hmrC in Drosophila 0-12 h embryo mRNA, hMePIP-seq: A**.

Example of gene showing 5hmrC peak distribution. Arrow indicates promoter orientation; **B.** Distribution of 5hmrC

peaks on embryonic transcripts and comparison of actual and predicted peaks according to the type of structural

element within the transcript; **C.** Distribution of all expressed (gray) or 5hmrC enriched (green) transcripts,

showing the number of mRNAs as a function of their expression levels in wt embryo; **E.** Sequence motif identified

in within 1815 5hmrC peaks.

Figure 5. Transcriptome-wide distribution of 5hmrC in LBF mRNA, hMePIP-seq: A. Example of gene
showing 5hmrC peak distribution. Arrow indicates promoter orientation; **B.** Distribution of 5hmrC peaks on wt LBF
transcripts and comparison of actual and predicted peaks according to the type of structural element within the
transcript; **C.** Distribution of 5hmrC peaks reduced by >1.4 times in *Tet*^null compared to the peaks found in the wt
LBF; note that peaks in the protein coding sequences and introns are significantly more reduced in *Tet*^null than are
the peaks in the 5' and 3' UTR; **D.** Distribution of all expressed (gray) or 5hmrC enriched (green) transcripts,
showing the number of mRNAs as a function of their expression levels in wt LBF; **E.** Distribution of all expressed
(gray) or hmrC enriched (green) transcripts, showing the number of mRNAs as a function of their expression
levels in *Tet*^null LBF; **F.** Sequence motif identified within 3711 5hmrC peaks.

Figure 6. The 5hmrC modified mRNAs. A. Volcano plot of mRNAs that are increased (green) or decreased
(red) relative to wildtype levels in *Tet^null* LBF preparations; **B.** Proportion of modified mRNAs in all 9000 wild type
transcripts, and in the decreased and increased portions of mRNAs from *Tet^null* LBF; note the low level of modified
transcripts in these two groups of mRNAs; **C.** Percent of transcripts that show a reduction of 5hmrC modification
of at least 1.4 times compared to wt LBF transcripts; **D.** Percent of transcript that show a >1.4 times reduction of
5hmrC modification that also show Tet binding to the corresponding gene; **E.** GO term analysis of transcripts that
show a >1.4 times reduction in 5hmrC modification; **F.** IGV tracks of a representative gene showing the
distribution of indicated peaks along the gene body. ChIP-seq, hMeRIP-seq and RNA-seq data are shown in
reads per million with the y-axis. Genomic regions with statistically significant enrichment were measured by -
log10 (peak P values); $P<10^{-8}$) are indicated. Effect of Tet depletion on 5hmC levels are also represented. Y axis
scale is indicated above each track. Blue arrows show reduction in 5hmrC peaks.

Figure 7. Tet regulates the expression of members of the Slit/Robo signaling pathway. Stage 16/17

embryonic ventral nerve cords immunolabelling a subpopulation of CNS neurons with Fas2 (**A-E)** and the general

neuronal cell surface marker, HRP. (**A'-E')**. **A, A'.** wild-type; **B, B'.** *Tet^null^/Tet^null^*; **C, C'.** *robo2^x123^/robo2^x123^;* **D, D'.**

*slı²/+***; E, E'.** *slı²/+*; T*et^null^/Tet^null^.* Examples of midline crossing are indicated by white arrows and malformed lateral

Fas2 tracks are noted with white arrowheads. Red arrows in **E'** highlight commissural malformations present in

*Tet^null^/Tet^null^* embryos with reduced slit dosage.

814