AedG, a new weapon in *Acinetobacter's* fight against antibiotics

Varsha Naidu[a,b], Bhumika Shah[b,c], Claire Maher[a,b], Ian T. Paulsen[b,c], Karl A. Hassan[a,b*]

Affiliations

[a]School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW 2308,

Australia

[b]ARC Centre of Excellence in Synthetic Biology, Macquarie University, North Ryde, NSW,

Australia

[c]School of Natural Sciences, Faculty of Science and Engineering, Macquarie University, North

Ryde, NSW, Australia

*Corresponding Author

School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW,

Australia. Phone: +6124921 7236. Email: karl.hassan@newcastle.edu.au

Abstract

A novel multidrug efflux pump, AdeG from the Drug:H$^+$ antiporter 2 family was discovered in

Acinetobacter multiresistance plasmids. Here, we profiled the antimicrobial resistance

potential and examined the distribution of this gene. Putative homologs of this novel efflux

pump were found in many *Acinetobacter* species and other Gram-negative species. The

pump conferred resistance to at least eight diverse antimicrobials. These results show that

the novel pump is a new determinant in the *Acinetobacter* resistance arsenal.

Introduction

The success of pathogenic *Acinetobacter* species in clinical settings is at least partly due to their drug resistance capabilities. Multidrug efflux pumps are significant contributors of resistance in *Acinetobacter*, as a single pump will frequently confer resistance to a broad range of antimicrobials. All *Acinetobacter* species encode efflux pumps that may confer resistance to antibiotics (1). Many of these pumps are found in the core genomes of the species, if not the genus, suggesting that they have been vertically inherited in these lineages for a considerable time (1, 2, 3). Others appear to be associated with mobile genetic elements, such as plasmids and transposons and have a scattered distribution in sub-sets of phylogenetically distant strains, suggesting horizontal transfer between strains (4, 5). For example, some *Acinetobacter* strains encode the tetracycline efflux pump, TetB, that was originally characterised in *Escherichia coli* on the transposon Tn*10* and is found in other Gram-negative species (6, 7).

We recently identified genes encoding a putative multidrug efflux pump from the Drug:H$^+$ antiporter 2 (DHA2) family of the Major Facilitator Superfamily (MFS) of transport proteins in several *Acinetobacter* plasmids (8). Here we demonstrate that the novel DHA2 family efflux pump can mediate resistance to a range of antimicrobials, and genes encoding homologs of the pump are widespread, but always associated with an *adeAB(C)* locus in *Acinetobacter* species. We call the novel efflux pump AedG, as <u>A</u>cinetobacter <u>e</u>fflux pump from the <u>D</u>HA2 family member G (9).

Methods

Bacterial growth conditions, Cloning, PCR and Complementation.

Bacterial strains were grown at 37 °C with shaking (200 rpm) in Mueller Hinton (MH) broth

or LB broth or on LB plates supplemented with 100 mg/L ampicillin, and IPTG where

necessary (see below). The coding region of *aedG* together with an RGSH6 epitope tag at its

3⬚ end was synthesised and cloned into the pTTQ18 plasmid (10), followed by next-

generation sequence verification by Twist Bioscience. The recombinant plasmid carring the

amvA gene (pTTQ18*amvA*) was obtained from our previous work (9). The empty plasmid

(pTTQ18) and recombinant plasmids (pTTQ18*aedG* and pTTQ18*amvA*) were transformed into

chemically competent *E. coli* BL21 cells. Transformants were selected on LB plates

containing 100 mg/L ampicillin and colony PCR was used to confirm positive insertion of

plasmids.

Antimicrobial susceptibility testing

Antimicrobial susceptibility assays were conducted in MH broth supplemented with 0.05

mM IPTG using the broth microdilution method as described (11). Assays for each

antimicrobial was performed with three biological replicates and four technical replicates,

resulting in twelve replicates in total. Cell growth was measured spectrophotometrically

using the PHERAstar FSX (BMG LABTECH, Germany).

Whole Cell transport assay

E. coli BL21 cells carrying either pTTQ18 (empty vector), pTTQ18*aedG,* or pTTQ18*amvA* were

grown in replicates of three in LB broth supplemented with 100 mg/L ampicillin to OD_{600} =

0.6, followed by induction of gene expression with IPTG (0.2 mM) for 1h. The cells were

washed three times in transport assay buffer (20 mM HEPES-NaOH (pH 7), 145 mM NaCl, 5

mM KCl) and resuspended in the same buffer to an OD_{600} = 1. Acriflavine (2 μM) or

ethidium (10 μM) and CCCP (10 μM) were added, and after 30 minutes incubation at 37 °C,

the cells were washed three times in transport assay buffer and resuspended in the same

buffer. Transport reaction was initiated at 37 °C by adding 1% glucose and a change in

fluorescence was measured using the PTI QuantaMaster 8000 (HORIBA Scientific) with an

excitation and emission wavelengths of 530 nm and 610 nm, respectively.

Distribution of AdeG and its phylogenetic relationship to other DHA2 family proteins

The *adeG* loci from *Acinetobacter* plasmids and their surrounding genes were aligned and

visualised using clicker v0.0.25 (12). The amino acid sequences of AedG homologs from

different lineages were aligned to the sequences of other representative DHA2 family

proteins using MUSCLE (13), and phylogeny inferred using MrBayes v3.2.6 (14).

Results and discussion

<u>AedG homologs are distributed widely across the *Acinetobacter* genus</u>

Genes encoding AedG were previously identified on plasmids carried by *A. baumannii, A.*

bereziniae, A. defluvii, A. johnsonii, A. nosocomialis, A. seifertii, A. ursingii and *Acinetobacter*

strains with no species designation (8). These genes were nearly identical, as only one

nucleotide difference was present between sub-sets of genes, producing proteins with

either a tyrosine or histidine at the predicted cytoplasmic boundary of TM helix 13 in the

proteins expressed. The genes were each located downstream of a gene cluster comprised

of *adeABC* and *adeRS* (Figure 1A), which encode a tripartite RND efflux system, AdeABC, and

its two-component transcriptional regulator, AdeRS (refs). The 3' ends of *adeC* in the *aedG*

linked clusters were formed by an inverted repeat sequence that was also found adjacent to

a nearby IS*18* (Figure 1A) (8). The locations of these repeat sequences suggested that the

intervening sequence, comprised of the IS*18* transposase gene, *adeRS* and *adeABC*, may be

mobilised by the IS*18* transposase. These loci also carried a partial *adeS* gene (3' end)

downstream of the *aedG* homologs, that could be related to an alternative organisation of

these genes in an ancestral locus (Figure 1A).

We sought to find additional homologs of *aedG* and determine whether they are also

associated with IS*18*/*adeRS*/*adeABC*. We initially queried the NCBI nr/nt database using

tblastn with the AedG sequence encoded in *A. nosocomialis* AC1530 plasmid pAC1530 and

found close to 300 high scoring hits from strains designated in 23 *Acinetobacter* species and

Acinetobacter strains with no species designation. Surveys of the local genetic environments

of the high scoring hits showed one of two general arrangements. Either *aedG* was located

downstream of the IS*18*/*adeRS*/*adeABC* cluster, as in pAC1530, or *aedG* was located

between homologs of *adeAB* and *adeRS* gene pairs and *adeC* and the IS*18* transposase gene

were absent, as in *A. baumannii* ABF9692 plasmid pABF9692 (Figure 1B). It is not unusual

for *adeC* to be absent from *adeAB* loci in *Acinetobacter* genomes, and AdeAB can function

with AdeK as an alternative outer membrane protein (8, 15). Insertion elements were also

seen in some of the loci, such as that in *A. lwoffii* strain AL_065 pAL_065-2, where insertion

sequence elements appear to have disrupted *adeS* (Figure 1B).

To identify additional homologs of AedG, we queried the NCBI experimental clustered

protein database using the amino acid sequence of AedG encoded by pAC1530. This search

identified 69 clusters whose representative members had above 65 % amino acid sequence

identity with the query AedG sequence across at least 95 % of its length. Proteins in the top

scoring cluster were encoded by *Acinetobacter* species. The local genetic environments of

these homologs resembled those of *aedG* in pAC1530 or pABF9692. Other high scoring

clusters included *Acinetobacter* proteins encoded by *aedG* homologs in an alternative

genetic environment. For example, the genetic environment of *aedG* homologs in the

Candidatus Acinetobacter avistercoris isolate 5402 metagenome assembled genome (MAG)

and an *A. bereziniae* GD03393 plasmid resembled that in pABF9692 but included *adeC*

(Figures 1B and S1). Of note, the *A. bereziniae* GD03393 plasmid harboured a second *aedG*

homolog similar to pAC1530 (Figure S1).

The proteins in lower scoring clusters were encoded by species outside the *Acinetobacter*

genus. The genes encoding these proteins were frequently adjacent to divergently

transcribed *tetR* family regulator genes that may control expression of the *aedG* homologs.

Some non-*Acinetobacter* aedG homologs were adjacent to RND efflux pump genes. For

example, the *aedG* homolog in the MAG of Proteobacteria bacterium isolate CTSoil_107

(likely *Pleomorphomonas* based on GyrB and RpoD sequence similarity) was adjacent to

distant homologs of *adeRS/adeAB* (Figure S2), and the *aedG* homolog from *Pelagibacterium*

sp. SCN 63-126 ABS74_C0016 was encoded adjacently to a novel RND efflux system (Figure

1B). These results demonstrate a tendency for genes encoding AedG to be co-localised with

those encoding RND efflux systems, specifically AdeAB(C) in *Acinetobacter* species. AedG

and AdeAB(C) may cooperate in the efflux of substrates from the cytoplasm to outside the

cell.

A phylogenetic analysis was performed to examine the relationships of AedG homologs with

other members of the DHA2 family. This analysis showed that AedG homologs are most

closely related to the AmvA (also known as AedF) (9, 16) and SmvA (17) efflux pumps from

Acinetobacter and *Salmonella*, respectively (Figure 1C). Pairwise amino acid sequence

similarity scores confirmed the relatedness of AedG homologs and AmvA/SmvA (Figure S3).

<u>AedG confers resistance to multiple antibiotics, biocides and dyes</u>

Based on the proximity of *aedG* genes to homologs of *adeAB(C)* in *Acinetobacter* species

and the similarity of AedG variants to drug transporters in the DHA2 family, it seemed likely

that AedG pumps would mediate resistance to one or more antimicrobials. We profiled the

antimicrobial resistance potential of the AdeG pump encoded in pAC1530 to eight

antimicrobials and compared resistance levels to those conferred by the multidrug efflux

pump AmvA in the same expression system (9, 16). The *aedG* homolog from pAC1530 was

synthesised and cloned into the pTTQ18 expression plasmid (Amp^r) behind the *tac* promoter

(10) with a C-terminal RGSH6 tag and sequence verified (Twist Bioscience). The *amvA* gene

from *A. baumannii* ATCC 17978 (also known as AedF) was cloned previously (9). AedG

conferred increased resistance to all eight antimicrobials tested when expressed in *E. coli*

BL21 cells and promoted higher levels of resistance to tetracycline, DAPI, ethanol and

ethidium bromide than AmvA (Figure 2A). These results suggest that AedG can recognise a

broad spectrum of antimicrobials similar to AmvA and is a new multidrug resistance protein

in *Acinetobacter*.

<u>AedG functions via an active efflux mechanism</u>

The ability of AedG to confer resistance to the nucleic acid intercalating dye ethidium

bromide allowed us to test efflux using real-time fluorometric transport assays. *E. coli* BL21

cells expressing AedG, AmvA or no additional protein (negative control) were loaded with 10

µM ethidium bromide in the absence of energy. The cells were washed, and re-energised

and ethidium efflux monitored by fluorescence changes. Both AmvA and AedG promoted

the active export of efflux ethidium bromide, as seen in fluorescence decrease over time

relative to control cells (Figure 2B). This result confirms efflux as the mechanism of

resistance mediated by AedG and that AedG is a new multidrug efflux pump.

Conclusions

The genes encoding AedG are part of the *Acinetobacter* accessory genome. This is in

contrast to AmvA, its closest relative, which is encoded in the core genome of *A. baumannii*

and many other *Acinetobacter* species. AmvA expression is not readily responsive to

antimicrobial treatment, and recent work showed that it may have a physiological function

in polyamine efflux (18). The organisation of *aedG* genes in *Acinetobacter*, downstream of

adeAB(C) and in proximity to *adeRS* homologs (Figures 1A and 1B), suggests that these

genes may be controlled by AdeRS and co-expressed with *adeAB(C)*. As such AedG

homologs may play a more significant role in antimicrobial resistance than AmvA in isolates

that carry them.

Figure 1. (A) Organisation of the local genetic environment of the *aedG* homolog in *A.*

nosocomialis AC1530 plasmid pAC1530 and related plasmids, including the locations of

surrounding genes and inverted repeat (IR) sequences that flank the IS*18* transposase gene

and *adeC*. (B) Alignment of loci carrying an *aedG* homolog in selected bacterial isolates.

Genes highlighted by colour and labels are found in more than one of the loci shown, or are

features noted in the text. This figure was made using clinker v0.0.25 (12). (C) The amino

acid sequences of AedG homologs from the lineages represented in (B) were aligned to the

sequences of other representative DHA2 family proteins using MUSCLE (13), and phylogeny

inferred using MrBayes v3.2.6 (14). The node labels show posterior probability values.

Figure 2. (A) *E. coli* BL21 cells carrying empty pTTQ18-*aedG* or pTTQ18-*amvA* were examined

in minimum inhibitory concentration (MIC) assays for resistance to a broad range of

antimicrobials, including the antibiotics tetracycline (TET) and erythromycin (ERY); the

biocides benzalkonium chloride (BAC), chlorhexidine (CHX), and ethanol (EtOH); and the

antimicrobial dyes 4⍰,6-diamidino-2-phenylindole (DAPI), acriflavine (ACR) and ethidium

bromide (EtBr). Resistance assays were performed as previously described (9). (B) Ethidium

transport from *E. coli* BL21 cells carrying either pTTQ18, pTTQ18-*amvA*, or pTTQ18-*aedG*.

Cells were loaded with 10 μM ethidium, transport was initiated by adding glucose and efflux

monitored fluorescently (Ex:530nm; Em:610nm) in a PTI QuantaMaster 8000 (HORIBA

Scientific) as described previously (9).

Figure S1. Local genetic organisation of the *adeG* loci in *A. bereziniae* GD03393 plasmid. One

gene cluster shows high sequence similarity to the *aedG* locus in *A. nosocomialis* AC1530

pAC1530 and an adjacent cluster shows high similarity to the *aedG* locus in the *Candidatus*

Acinetobacter avistercoris 5402 metagenome assembled genome. This figure was made

using clinker v0.0.25 (12).

Figure S2. (A) The phylogenetic tree of AdeB homologs from representative *Acinetobacter*

lineages (Figure 1B). The MAG of Proteobacteria bacterium isolate CTSoil_107, the *A.*

baumannii ATCC 17978 chromosome, and the AdeJ sequence from the *A. baumannii* ATCC

17978 chromosome were aligned using MUSCLE (13). The phylogeny of these sequences

was inferred using MrBayes v3.2.6 (14). The node labels show posterior probability values.

(B) Pairwise amino acid sequence similarity (bottom left) and identity (top right) scores for

the proteins shown in (A). These values were determined using MatGat version 2.0 (19) with

a BOLSUM 50 scoring matrix.

	AedG pAL_065_2	AedG pABF9692	AedG Can. A. avistercoris MAG	AedG pAC1530	AedG Pelagibacterium MAG	AedG Proteobacterium MAG	AmvA	SmvA	PqrB	CmgA	QacA	QepA	LfrA	SgvT1	RifP	Ptr
AedG pAL_065-2		98	88.1	97	72.6	66.6	42.7	43.5	36.2	33.8	34.2	35.5	32.9	30.9	31.2	30.9
AedG pABF9692	99.2		89.1	99	72.4	67.2	42.9	43.5	36.4	33.5	34.4	36.2	32.8	30.6	31.6	31
AedG Can. A. avistercoris MAG	93.5	94.2		88.9	70.1	66.6	44.9	43.7	35.8	34.4	33	35.7	33.5	32.5	32.2	30.9
AedG pAC1530	98.6	99.4	94.2		72.4	67	42.7	43.5	35.8	33.6	33.8	36	32.2	30.4	31.2	31
AedG Pelagibacterium MAG	83.9	83.5	83.3	83.7		71.7	43	45.3	38	32.5	32.1	38.3	33.9	32	34.7	34.6
AedG Proteobacteria MAG	82.6	82.4	81.2	82.6	84.4		41.5	43.1	37.2	34.7	31.8	36.7	35.7	32	32.2	32
AmvA	66.5	66.1	67.5	66.3	65.3	62.6		55.2	33	32	28.7	30.7	33.6	27.9	31.2	31.6
SmvA	67.1	67.5	67.7	67.7	67.1	63.4	76.4		37.4	34.2	32.3	35.5	32.4	31.7	34.9	31.5
PqrB	57.1	57.1	54.9	56.9	56.5	57.1	59.4	59.2		39.7	36	45.8	47.8	40	40.4	38.1
CmgA	58.5	58.3	58.3	58.1	56.2	57.5	59.3	56.6	59.8		33	35.8	33.3	32.5	32.5	34.7
QacA	57.2	57.2	56.8	56.8	54.5	56	53.5	54.1	60.3	55.3		33.1	33.5	29.4	30.3	31.5
QepA	55.6	55.8	57.3	55.8	55.2	55.4	54.4	56.4	64.8	55.6	59.9		42.4	31.3	35.7	35.2
LfrA	54.4	54.2	53.2	53.8	53.4	55.7	58.5	54.8	69.2	54.6	57.8	63.4		34	33	31.4
SgvT1	52.6	52.5	55.3	53.4	53.2	52.6	52.1	51.3	59.4	53.8	52.8	53.6	53.2		50.3	48.2
RifP	52.9	53.1	53.3	52.9	54.2	53.3	54.4	53.6	59	51.5	55	57.1	53.8	68.5		62.5
Ptr	53.6	54.4	56.7	54.2	57.2	55.9	55.7	53.7	56.3	56.3	54.3	56.4	55.4	66.8	77.8	

Figure S3. Pairwise amino acid sequence similarity (bottom left) and identity (top right)

scores for members of the DHA2 family of efflux proteins. These values were determined

using MatGat version 2.0 (19) with a BOLSUM 50 scoring matrix.

Conflicts of Interest

None

Funding information

This work was supported by an Australian Research Council Future Fellowship to KAH

(FT180100123) and a Project Grant from the National Health and Medical Research Council

of Australia to ITP and KAH (GNT1120298).

Ethical Approval

N/A

References

1.   Brzoska AJ, Hassan KA, de Leon EJ, Paulsen IT, Lewis PJ. Single-Step Selection of Drug
Resistant Acinetobacter baylyi ADP1 Mutants Reveals a Functional Redundancy in the
Recruitment of Multidrug Efflux Systems. PLoS One. 2013;8(2):e56090.
2.   Hassan KA, Maher C, Elbourne LD, Henderson PJ, Paulsen IT. Increasing the PACE of
characterising novel transporters by functional genomics. Curr Opin Microbiol. 2021;64:1-8.
3.   Henderson PJF, Maher C, Elbourne LDH, Eijkelkamp BA, Paulsen IT, Hassan KA.
Physiological Functions of Bacterial "Multidrug" Efflux Pumps. Chem Rev. 2021;121(9):5417-
78.
4.   Bi D, Zheng J, Xie R, Zhu Y, Wei R, Ou HY, et al. Comparative Analysis of AbaR-Type
Genomic Islands Reveals Distinct Patterns of Genetic Features in Elements with Different
Backbones. mSphere. 2020;5(3).
5.   Guardabassi L, Dijkshoorn L, Collard JM, Olsen JE, Dalsgaard A. Distribution and in-
vitro transfer of tetracycline resistance determinants in clinical and aquatic Acinetobacter
strains. J Med Microbiol. 2000;49(10):929-36.
6.   Levy SB, McMurry L. Plasmid-determined tetracycline resistance involves new
transport systems for tetracycline. Nature. 1978;276(5683):90-2.
7.   Li L, Hassan KA, Brown MH, Paulsen IT. Rapid multiplexed phenotypic screening
identifies drug resistance functions for three novel efflux pumps in Acinetobacter
baumannii. J Antimicrob Chemother. 2016;71(5):1223-32.

8.   Naidu V, Bartczak A, Brzoska AJ, Lewis P, Eijkelkamp BA, Paulsen IT, et al. Evolution
of RND efflux pumps in the development of a successful pathogen. Drug Resist Updat.
2023;66:100911.
9.   Hassan KA, Brzoska AJ, Wilson NL, Eijkelkamp BA, Brown MH, Paulsen IT. Roles of
DHA2 family transporters in drug resistance and iron homeostasis in Acinetobacter spp. J
Mol Microbiol Biotechnol. 2011;20(2):116-24.
10.   Stark MJ. Multicopy expression vectors carrying the lac repressor gene for regulated
high-level expression of genes in Escherichia coli. Gene. 1987;51(2-3):255-67.
11.   Wiegand I, Hilpert K, Hancock RE. Agar and broth dilution methods to determine the
minimal inhibitory concentration (MIC) of antimicrobial substances. Nat Protoc.
2008;3(2):163-75.
12.   Gilchrist CLM, Chooi YH. Clinker & clustermap.js: Automatic generation of gene
cluster comparison figures. Bioinformatics. 2021.
13.   Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. BMC Bioinformatics. 2004;5:113.
14.   Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogenetic trees.
Bioinformatics. 2001;17(8):754-5.
15.   Sugawara E, Nikaido H. Properties of AdeABC and AdeIJK efflux systems of
Acinetobacter baumannii compared with those of the AcrAB-TolC system of Escherichia coli.
Antimicrob Agents Chemother. 2014;58(12):7250-7.
16.   Rajamohan G, Srinivasan VB, Gebreyes WA. Molecular and functional
characterization of a novel efflux pump, AmvA, mediating antimicrobial and disinfectant
resistance in Acinetobacter baumannii. J Antimicrob Chemother. 2010;65(9):1919-25.
17.   Santiviago CA, Fuentes JA, Bueno SM, Trombert AN, Hildago AA, Socias LT, et al. The
Salmonella enterica sv. Typhimurium smvA, yddG and ompD (porin) genes are required for
the efficient efflux of methyl viologen. Mol Microbiol. 2002;46(3):687-98.
18.   Short FL, Liu Q, Shah B, Clift HE, Naidu V, Li L, et al. The Acinetobacter baumannii
disinfectant resistance protein, AmvA, is a spermidine and spermine efflux pump. Commun
Biol. 2021;4(1):1114.
19.   Campanella JJ, Bitincka L, Smalley J. MatGAT: an application that generates
similarity/identity matrices using protein or DNA sequences. BMC bioinformatics. 2003;4:29.