AadT, a new weapon in *Acinetobacter's* fight against antibiotics

Varsha Naidu[a,b], Bhumika Shah[b,c], Claire Maher[a,b], Ian T. Paulsen[b,c], Karl A. Hassan[a,b*]

Affiliations

[a]School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW 2308,

Australia

[b]ARC Centre of Excellence in Synthetic Biology, Macquarie University, North Ryde, NSW,

Australia

[c]School of Natural Sciences, Faculty of Science and Engineering, Macquarie University, North

Ryde, NSW, Australia

*Corresponding Author

School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW,

Australia. Phone: +6124921 7236. Email: karl.hassan@newcastle.edu.au

Abstract

A novel multidrug efflux pump, AadT from the Drug:H$^+$ antiporter 2 family, was discovered in

Acinetobacter multidrug resistance plasmids. Here, we profiled the antimicrobial resistance

potential and examined the distribution of this gene. Putative homologs of this efflux pump

were encoded in many *Acinetobacter* species and other Gram-negative species, and were

genetically associated with novel variants of *adeAB(C)*, which encodes a major tripartite efflux

pump in *Acinetobacter*. The AadT pump conferred decreased susceptibility to at least eight

diverse antimicrobials, including antibiotics erythromycin, tetracycline; biocides

chlorhexidine; and dyes ethidium bromide and DAPI. These results show that AadT is a new

determinant in the *Acinetobacter* resistance arsenal and may cooperate with variants of

AdeAB(C)*.

Introduction

The success of pathogenic *Acinetobacter* species in clinical settings is at least partly due to their drug resistance capabilities. Multidrug efflux pumps are significant contributors of resistance in *Acinetobacter*, as a single pump will frequently confer resistance to a broad range of antimicrobials. All *Acinetobacter* species encode efflux pumps that may confer resistance to antibiotics (1). Many of these pumps are encoded in the core genomes of the species, if not the genus, suggesting that the genes have been vertically inherited in these lineages since speciation (1-3). Others appear to be associated with mobile genetic elements, such as plasmids and transposons and have a scattered distribution in sub-sets of phylogenetically distant strains, suggesting horizontal transfer between strains (4, 5). For example, some *Acinetobacter* strains encode the tetracycline efflux pump, TetB, that was originally characterised in *Escherichia coli* on the transposon Tn*10* and is found in other Gram-negative species (6, 7).

We recently identified genes encoding a putative multidrug efflux pump from the Drug:H$^+$ antiporter 2 (DHA2) family of the Major Facilitator Superfamily (MFS) of transport proteins in several *Acinetobacter* plasmids (8). Here we demonstrate that the novel DHA2 family efflux pump can decrease susceptibility to a range of antimicrobials, and that genes encoding homologs of the pump are widespread, but always associated with a variant of the *adeAB(C)* locus in *Acinetobacter* species. We call the novel efflux pump AadT, as the <u>A</u>cinetobacter <u>a</u>ntimicrobial <u>d</u>rug <u>t</u>ransporter (9).

Methods

<u>Bacterial growth conditions, Cloning, PCR and Complementation</u>

Bacterial strains were grown at 37 °C with shaking (200 rpm) in Mueller Hinton (MH) broth or

LB broth or on LB plates supplemented with 100 mg/L ampicillin, and IPTG where necessary

(see below). The coding region of *aadT* together with an RGSH6 epitope tag at its 3' end was

synthesised and cloned into the pTTQ18 plasmid (10), followed by next-generation sequence

verification by Twist Bioscience. The recombinant plasmid carring the *amvA* gene

(pTTQ18$_{amvA}$) was obtained from our previous work (9). The empty plasmid (pTTQ18) and

recombinant plasmids (pTTQ18$_{aadt}$ and pTTQ18$_{amvA}$) were transformed into chemically

competent *E. coli* BL21 cells. Transformants were selected on LB plates containing 100 mg/L

ampicillin and colony PCR was used to confirm positive insertion of plasmids.

<u>Antimicrobial susceptibility testing</u>

Antimicrobial susceptibility assays were conducted in cation adjusted MH broth

supplemented with 0.05 mM IPTG using the broth microdilution method as described (11).

Assays for each antimicrobial were performed with three biological replicates and four

technical replicates, resulting in twelve replicates in total. Cell growth was measured

spectrophotometrically as OD_{600} endpoint readings using the PHERAstar FSX (BMG LABTECH,

Germany).

<u>Whole Cell transport assay</u>

E. coli BL21 cells carrying either pTTQ18 (empty vector), pTTQ18$_{aadt,}$ or pTTQ18$_{amvA}$ were

grown in biological replicates of three in LB broth supplemented with 100 mg/L ampicillin to

OD_{600} = 0.6, followed by induction of gene expression with IPTG (0.2 mM) for 1h. The cells

were washed three times in transport assay buffer (20 mM HEPES-NaOH (pH 7), 145 mM NaCl,

5 mM KCl) and resuspended in the same buffer to an OD_{600} = 1. Ethidium bromide (10 μM)

and CCCP (10 μM) were added, and after 30 minutes incubation at 37 °C, the cells were

washed three times in transport assay buffer and resuspended in the same buffer. Transport

reaction was initiated at 37 °C by adding 1 % glucose and fluorescence was measured over

time using the PTI QuantaMaster 8000 (HORIBA Scientific) with an excitation and emission

wavelengths of 530 nm and 610 nm, respectively.

<u>Distribution of AadT and its phylogenetic relationship to other DHA2 family proteins</u>

Variants of *aadT* were identified in NCBI databases using NCBI blast searches (see detailed

descriptions in the text). Representative *aadT* loci and their surrounding genes were aligned

and visualised using clicker v0.0.25 (12). The amino acid sequences of representative AadT

homologs from different lineages were aligned to the sequences of other representative

DHA2 family proteins using MUSCLE (13), and phylogeny inferred using MrBayes v3.2.6 (14).

The pairwise sequence identity and similarity of the representative proteins was determined

using MatGat (15). The same approaches were used to infer the phylogeny and sequence

similarities of representative homologs of adeB.

Results and discussion

<u>AadT homologs are distributed widely across the *Acinetobacter* genus</u>

Genes encoding AadT were previously identified on plasmids carried by *A. baumannii, A.*

bereziniae, A. defluvii, A. johnsonii, A. nosocomialis, A. seifertii, A. ursingii and *Acinetobacter*

strains with no species designation (8). These genes were nearly identical, as only one

nucleotide difference was present between sub-sets of genes, producing proteins with either

a tyrosine or histidine at the predicted cytoplasmic boundary of TM helix 13. The genes were

each located downstream of a gene cluster comprised of variants of *adeABC* and *adeRS*

(Figure 1A), which encode a tripartite RND efflux system, AdeABC, and its two-component

transcriptional regulator, AdeRS (16). The 3' ends of *adeC* in the *aadT* linked clusters were

formed by an inverted repeat sequence that was also found adjacent to a nearby IS*18* (Figure

1A) (8). The locations of the repeat sequences suggested that the intervening sequence,

comprising the IS*18* transposase gene, *adeRS* and *adeABC*, may be mobilised by the IS*18*

transposase. These loci also carried a partial *adeS* gene (3' end) downstream of the *aadT*

homologs, that could be related to an alternative organisation of these genes in an ancestral

locus (Figure 1A).

We sought to find additional homologs of *aadT* and determine whether they are also

associated with IS*18/adeRS/adeABC*. We initially queried the NCBI nr/nt database using

tblastn with the AadT amino acid sequence encoded in the *A. nosocomialis* AC1530 plasmid

pAC1530 and found close to 300 high scoring hits from strains designated in 23 *Acinetobacter*

species and *Acinetobacter* strains with no species designation. Surveys of the local genetic

environments of the high scoring hits showed one of two general arrangements. Either *aadT*

was located downstream of the IS*18/adeRS/adeABC* cluster, as in pAC1530, or *aadT* was

located between homologs of *adeAB* and *adeRS* gene pairs and *adeC* and the IS*18* transposase

gene were absent, as in *A. baumannii* ABF9692 plasmid pABF9692 (Figure 1B). It is not unusual

for *adeC* to be absent from *adeAB* loci in *Acinetobacter* genomes, and AdeAB can function

with AdeK as an alternative outer membrane protein (8, 17). Insertion elements were also

seen in some of the loci, such as that in *A. lwoffii* strain AL_065 pAL_065-2, where insertion

sequence elements appear to have disrupted *adeS* (Figure 1B).

To identify additional homologs of AadT, we queried the NCBI experimental clustered protein

database using the amino acid sequence of AadT encoded by pAC1530. This search identified

69 clusters whose representative members had above 65 % amino acid sequence identity

with the query AadT sequence across at least 95 % of its length. Proteins in the top scoring

cluster were encoded by *Acinetobacter* species. The local genetic environments of these

homologs resembled those of *aadT* in pAC1530 or pABF9692. Other high scoring clusters

included *Acinetobacter* proteins encoded by *aadT* homologs in an alternative genetic

environment. For example, the genetic environment of *aadT* homologs in the *Candidatus*

Acinetobacter avistercoris isolate 5402 metagenome assembled genome (MAG) and an *A.*

bereziniae GD03393 plasmid resembled that in pABF9692 but included *adeC* (Figures 1B and

S1). Of note, the *A. bereziniae* GD03393 plasmid harboured a second *aadT* homolog similar

to pAC1530 (Figure S1).

The proteins in lower scoring clusters were encoded by species outside the *Acinetobacter*

genus. The genes encoding these proteins were frequently adjacent to divergently

transcribed *tetR* family regulator genes that may control expression of the *aadT* homologs.

Some non-*Acinetobacter aadT* homologs were adjacent to RND efflux pump genes. For

example, the *aadT* homolog in the MAG of Proteobacteria bacterium isolate CTSoil_107 (likely

Pleomorphomonas based on GyrB and RpoD sequence similarity) was adjacent to putative

distant homologs of *adeRS*/*adeAB* (Figure 1), and the *aadT* homolog from *Pelagibacterium* sp.

SCN 63-126 ABS74_C0016 was encoded adjacently to a novel RND efflux system (Figure 1B).

These results demonstrate a tendency for genes encoding AadT to be co-localised with those

encoding RND efflux systems, specifically variants of AdeAB(C) in *Acinetobacter* species. AadT

and AdeAB(C) may cooperate in the efflux of substrates from the cytoplasm to outside the

cell, with AadT moving substrates from the cytoplasm to the periplasm and AdeAB(C) moving

substrates from the periplasm out of the cell (18).

A phylogenetic analysis was performed to examine the relationships of AadT homologs with

other members of the DHA2 family. This analysis showed that AadT homologs are most

closely related to the AmvA (also known as AedF) (9, 19) and SmvA (20) efflux pumps from

Acinetobacter and *Salmonella*, respectively (Figure 1C). Pairwise amino acid sequence

similarity scores confirmed the relatedness of AadT homologs and AmvA/SmvA (Figure S2).

Phylogenetic and pairwise amino acid sequence similarity studies were conducted to examine

the relationships of AdeB homologs encoded adjacently to *aadT* variants (Figure 1B). The

results (Figure 1D and 1E) showed that the AdeB pump encoded in pAC1530 is most similar

to the chromosomally encoded AdeB pump from *A. baumannii* ATCC 17978 (95 % amino acid

sequence similarity; Figure 1E). The AdeB pumps encoded in pAL_065-1, pABF9692 and the

MAG of *Candidatus Acinetobacter avistercoris* 5402 showed greater than 98 % amino acid

sequence similarity to each other, but were less than 88 % similar to the chromosomal AdeB

or AdeB encoded on pAC1530 (Figures 1D and 1E). The putative AdeB homolog encoded by

the Proteobacteria CTSoli_107 isolate showed low similarity to all other AdeB homologs

(Figure 1E). These results indicate that at least two distct putative *adeAB(C)* loci have been

acquired on different *Acinetobacter* plasmids and formed an association with *aadT*. As above,

the *A. bereziniae* GD03393 plasmid carries both of these loci (Figure S1).

AadT confers resistance to multiple antibiotics, biocides and dyes

Based on the proximity of *aadT* genes to homologs of *adeAB(C)* in *Acinetobacter* species and

the similarity of AadT variants to drug transporters in the DHA2 family, it seemed likely that

AadT pumps would mediate resistance to one or more antimicrobials. We profiled the

antimicrobial resistance potential of the AadT pump encoded in pAC1530 to eight

antimicrobials and compared resistance levels to those conferred by the multidrug efflux

pump AmvA in the same expression system (9, 19). The *aadT* homolog from pAC1530 was

synthesised and cloned into the pTTQ18 expression plasmid (Amp^r) behind the *tac* promoter

(10) with a C-terminal RGSH6 tag and sequence verified (Twist Bioscience). The *amvA* gene

from *A. baumannii* ATCC 17978 (also known as AedF) was cloned previously (9). AadT reduced

susceptibility to all eight antimicrobials tested when expressed in *E. coli* BL21 cells and

promoted higher levels of tolerance to tetracycline, DAPI, ethanol and ethidium bromide than

AmvA (Figure 2A). These results suggest that AadT can recognise a broad spectrum of

antimicrobials similar to AmvA and is a new multidrug transport protein in *Acinetobacter*.

<u>AadT functions via an active efflux mechanism</u>

The ability of AadT to confer tolerance to the nucleic acid intercalating dye ethidium bromide

allowed us to test efflux using real-time fluorometric transport assays. *E. coli* BL21 cells

expressing AadT, AmvA or no additional protein (negative control) were loaded with 10 µM

ethidium bromide in the absence of energy. The cells were washed, and re-energised and

ethidium efflux monitored by fluorescence changes. Both AmvA and AadT promoted the

active export of efflux ethidium bromide, as seen in fluorescence decrease over time relative

to control cells (Figure 2B). This result confirms efflux as the mechanism of tolerance

mediated by AadT and that AadT is a new multidrug efflux pump.

Conclusions

The genes encoding AadT are part of the accessory genomes of more than 20 *Acinetobacter*

species. This is in contrast to AmvA, its closest known relative in *Acinetobacter*, which is

encoded in the core genome of *A. baumannii* and other *Acinetobacter* species. AmvA

expression in *Acinetobacter* is not readily responsive to antimicrobial treatment, and recent

results from our groups showed that it may have a physiological function in polyamine efflux

(21). The organisation of *aadT* genes in *Acinetobacter*, downstream of *adeAB(C)* variants and

in proximity to *adeRS* homologs (Figures 1A and 1B), suggests that *aadT* genes may be

controlled by AdeRS and co-expressed with *adeAB(C)*. As such AadT homologs may play a

more significant role in antimicrobial resistance than AmvA in the isolates that carry them.

That *aadT* has formed a genetic association with several novel RND efflux pump genes,

mostly homologs of *adeAB(C)*, suggests a selective advantage to this arrangement, which as

mentioned above, may relate to cooperative transport activities of these pumps (18, 22).

The discovery of genes encoding novel variants of AdeAB(C) and AdeRS in *Acinetobacter*

plasmids is significant. These genes were found in hundreds of *Acinetobacter* sequences. The

variant AdeAB(C) pumps may have different substrate preferences to the chromosomally

encoded pumps. Details of these substrate preferences will be of significant interest in

determining their importance to antibiotic resistance in pathogenic *Acinetobacter.* The role

of the plasmid encoded AdeRS system is also of significant interest. This system seems likely

to control expression of the plasmid encoded *adeAB(C)* genes and as mentioned above may

control *aadT* gene expression. It is also possible that the plasmid encoded AdeRS systems

could promote expression of chromosomal *adeAB(C)* homologs, which are tightly regulated

in many strains.

Figure 1. (A) Organisation of the local genetic environment of the *aadT* homolog in *A.*
nosocomialis AC1530 plasmid pAC1530 and related plasmids, including the locations of
surrounding genes and inverted repeat (IR) sequences that flank the IS*18* transposase gene
and *adeC*. (B) Alignment of loci carrying an *aadT* homolog in selected bacterial isolates. Genes
highlighted by colour and labels are found in more than one of the loci shown, or are features
noted in the text. This figure was made using clinker v0.0.25 (12). (C) The amino acid
sequences of AadT homologs from the lineages represented in (B) were aligned to the
sequences of other representative DHA2 family proteins using MUSCLE (13), and phylogeny
inferred using MrBayes v3.2.6 (14). The node labels show posterior probability values. (D) The
putative AdeB homologs depicted in panel (B), and the AdeJ sequence from the *A. baumannii*
ATCC 17978 chromosome were aligned using MUSCLE (13) and the phylogeny of these
sequences was inferred using MrBayes v3.2.6 (14) and is shown in the tree. The node labels

show posterior probability values. (E) Pairwise amino acid sequence similarity (bottom left)

and identity (top right) scores for the proteins used in the phylogenetic analysis (D). These

values were determined using MatGat version 2.0 (15) with a BOLSUM 50 scoring matrix.

Figure 2. (A) *E. coli* BL21 cells carrying empty pTTQ18-*aadT* or pTTQ18-*amvA* were examined

in minimum inhibitory concentration (MIC) assays for resistance to a broad range of

antimicrobials, including the antibiotics tetracycline (TET) and erythromycin (ERY); the

biocides benzalkonium chloride (BAC), chlorhexidine (CHX), and ethanol (EtOH); and the

antimicrobial dyes 4',6-diamidino-2-phenylindole (DAPI), acriflavine (ACR) and ethidium

bromide (EtBr). The OD$_{600}$ measurements were taken as endpoint readings after 24h growth.

Resistance assays were performed as previously described (9). (B) Ethidium bromide

transport assays in *E. coli* BL21 cells carrying either pTTQ18, pTTQ18-*amvA*, or pTTQ18-*aadT*.

Cells were loaded with 10 µM ethidium, transport was initiated by adding glucose and efflux

monitored fluorescently (Ex:450nm; Em:509nm) in a PTI QuantaMaster 8000 (HORIBA

Scientific) as described previously (9). The total area under the curve (AUC) was calculcated

after transport initiation by glucose for each strain BL21 pTTQ18 (150); pTTQ18-*amvA* (72);

and pTTQ18-*aadT* (120). The grey shaded area represents the standard error of three

biological replicates per strain.

249 Figure S1. Local genetic organisation of the *aadT* loci in *A. bereziniae* GD03393 plasmid. One

250 gene cluster shows high sequence similarity to the *aadT* locus in *A. nosocomialis* AC1530

251 pAC1530 and an adjacent cluster shows high similarity to the *aadT* locus in the *Candidatus*

252 *Acinetobacter avistercoris* 5402 metagenome assembled genome. This figure was made using

253 clinker v0.0.25 (12).

Figure S2. Pairwise amino acid sequence similarity (bottom left) and identity (top right)

scores for members of the DHA2 family of efflux proteins. These values were determined

using MatGat version 2.0 (15) with a BOLSUM 50 scoring matrix.

Conflicts of Interest

<u>None</u>

Funding information

This work was supported by an Australian Research Council Future Fellowship to KAH

(FT180100123) and a Project Grant from the National Health and Medical Research Council

of Australia to ITP and KAH (GNT1120298).

Ethical Approval

N/A

References

1.   Brzoska AJ, Hassan KA, de Leon EJ, Paulsen IT, Lewis PJ. Single-Step Selection of Drug
Resistant Acinetobacter baylyi ADP1 Mutants Reveals a Functional Redundancy in the
Recruitment of Multidrug Efflux Systems. PLoS One. 2013;8(2):e56090.
2.   Hassan KA, Maher C, Elbourne LD, Henderson PJ, Paulsen IT. Increasing the PACE of
characterising novel transporters by functional genomics. Curr Opin Microbiol. 2021;64:1-8.
3.   Henderson PJF, Maher C, Elbourne LDH, Eijkelkamp BA, Paulsen IT, Hassan KA.
Physiological Functions of Bacterial "Multidrug" Efflux Pumps. Chem Rev. 2021;121(9):5417-
78.
4.   Bi D, Zheng J, Xie R, Zhu Y, Wei R, Ou HY, et al. Comparative Analysis of AbaR-Type
Genomic Islands Reveals Distinct Patterns of Genetic Features in Elements with Different
Backbones. mSphere. 2020;5(3).
5.   Guardabassi L, Dijkshoorn L, Collard JM, Olsen JE, Dalsgaard A. Distribution and in-
vitro transfer of tetracycline resistance determinants in clinical and aquatic Acinetobacter
strains. J Med Microbiol. 2000;49(10):929-36.
6.   Levy SB, McMurry L. Plasmid-determined tetracycline resistance involves new
transport systems for tetracycline. Nature. 1978;276(5683):90-2.
7.   Li L, Hassan KA, Brown MH, Paulsen IT. Rapid multiplexed phenotypic screening
identifies drug resistance functions for three novel efflux pumps in Acinetobacter
baumannii. J Antimicrob Chemother. 2016;71(5):1223-32.

8.   Naidu V, Bartczak A, Brzoska AJ, Lewis P, Eijkelkamp BA, Paulsen IT, et al. Evolution
of RND efflux pumps in the development of a successful pathogen. Drug Resist Updat.
2023;66:100911.
9.   Hassan KA, Brzoska AJ, Wilson NL, Eijkelkamp BA, Brown MH, Paulsen IT. Roles of
DHA2 family transporters in drug resistance and iron homeostasis in Acinetobacter spp. J
Mol Microbiol Biotechnol. 2011;20(2):116-24.
10.   Stark MJ. Multicopy expression vectors carrying the lac repressor gene for regulated
high-level expression of genes in Escherichia coli. Gene. 1987;51(2-3):255-67.
11.   Wiegand I, Hilpert K, Hancock RE. Agar and broth dilution methods to determine the
minimal inhibitory concentration (MIC) of antimicrobial substances. Nat Protoc.
2008;3(2):163-75.
12.   Gilchrist CLM, Chooi YH. Clinker & clustermap.js: Automatic generation of gene
cluster comparison figures. Bioinformatics. 2021.
13.   Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. BMC Bioinformatics. 2004;5:113.
14.   Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogenetic trees.
Bioinformatics. 2001;17(8):754-5.
15.   Campanella JJ, Bitincka L, Smalley J. MatGAT: an application that generates
similarity/identity matrices using protein or DNA sequences. BMC Bioinformatics. 2003;4:29.
16.   Marchand I, Damier-Piolle L, Courvalin P, Lambert T. Expression of the RND-type
efflux pump AdeABC in Acinetobacter baumannii is regulated by the AdeRS two-component
system. Antimicrob Agents Chemother. 2004;48(9):3298-304.
17.   Sugawara E, Nikaido H. Properties of AdeABC and AdeIJK efflux systems of
Acinetobacter baumannii compared with those of the AcrAB-TolC system of Escherichia coli.
Antimicrob Agents Chemother. 2014;58(12):7250-7.
18.   Tal N, Schuldiner S. A coordinated network of transporters with overlapping
specificities provides a robust survival strategy. P Natl Acad Sci USA. 2009;106(22):9051-6.
19.   Rajamohan G, Srinivasan VB, Gebreyes WA. Molecular and functional
characterization of a novel efflux pump, AmvA, mediating antimicrobial and disinfectant
resistance in Acinetobacter baumannii. J Antimicrob Chemother. 2010;65(9):1919-25.
20.   Santiviago CA, Fuentes JA, Bueno SM, Trombert AN, Hildago AA, Socias LT, et al. The
Salmonella enterica sv. Typhimurium smvA, yddG and ompD (porin) genes are required for
the efficient efflux of methyl viologen. Mol Microbiol. 2002;46(3):687-98.
21.   Short FL, Liu Q, Shah B, Clift HE, Naidu V, Li L, et al. The Acinetobacter baumannii
disinfectant resistance protein, AmvA, is a spermidine and spermine efflux pump. Commun
Biol. 2021;4(1):1114.
22.   Lee A, Mao WM, Warren MS, Mistry A, Hoshino K, Okumura R, et al. Interplay
between efflux pumps may provide either additive or multiplicative effects on drug
resistance. J Bacteriol. 2000;182(11):3142-50.