AedG, a new weapon in *Acinetobacter*'s fight against antimicrobials

Varsha Naidu[a,b], Bhumika Shah[b,c], Claire Maher[a,b], Ian T. Paulsen[b,c], Karl A. Hassan[a,b*]

[a]School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW 2308,

Australia

[b]ARC Centre of Excellence in Synthetic Biology, Macquarie University, North Ryde, NSW,

Australia

[c]School of Natural Sciences, Faculty of Science and Engineering, Macquarie University, North

Ryde, NSW, Australia

*Corresponding Author

School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW,

Australia. Phone: +6124921 7236. Email: karl.hassan@newcastle.edu.au

Abstract

A novel multidrug efflux pump, AdeG from the Drug:H$^+$ antiporter 2 family was discovered in *Acinetobacter* plasmids. Here, we profiled the antimicrobial resistance potential and examined the distribution of this gene. Close and distant homologs of the efflux pump were found in several *Acinetobacter* species and other Gram-negative species, respectively. The pump conferred resistance to at least eight diverse antimicrobials. These results show that the novel pump is a new determinant in the *Acinetobacter* resistance arsenal*.*

 Introduction

The success of pathogenic *Acinetobacter* species in clinical settings is at least partly due to

their drug resistance capabilities. Multidrug efflux pumps are significant contributors of

resistance in *Acinetobacter*, as a single pump will frequently confer resistance to a broad

range of antimicrobials. All *Acinetobacter* species encode efflux pumps that may confer

resistance to antibiotics (1). Many of these pumps are found in the core genomes of the

species, if not the genus, suggesting that they have been vertically inherited in these lineages

for a considerable time (1-3). Others appear to be associated with mobile genetic elements,

such as plasmids and transposons and have a scattered distribution in sub-sets of

phylogenetically distant strains, suggesting horizontal transfer between strains (4, 5). For

example, some *Acinetobacter* strains encode the tetracycline efflux pump, TetB, that was

originally characterised in *Escherichia coli* on the transposon Tn*10* and is found in other Gram-

negative species (6, 7).

We recently identified genes encoding a putative multidrug efflux pump from the Drug:H$^+$

antiporter 2 (DHA2) family of the Major Facilitator Superfamily (MFS) of transport proteins in

several *Acinetobacter* plasmids (8). Here we demonstrate that the novel DHA2 family efflux

pump can mediate resistance to a range of antimicrobials, and genes encoding homologs of

the pump are widespread, but always associated with an *adeAB(C)* locus in *Acinetobacter*

species. We call the novel efflux pump AedG, as <u>A</u>cinetobacter <u>e</u>fflux pump from the <u>DHA2</u>

family member G (9).

<u>AedG homologs are distributed widely across the *Acinetobacter* genus</u>

Genes encoding AedG were previously identified on plasmids carried by *A. baumannii, A.*

bereziniae, A. defluvii, A. johnsonii, A. nosocomialis, A. seifertii, A. ursingii and *Acinetobacter*

strains with no species designation (8). These genes were nearly identical; only one nucleotide

difference was present between sub-sets of genes, producing proteins with either a tyrosine

or histidine at the predicted cytoplasmic boundary of TM helix 13 in the proteins expressed.

The genes were each located downstream of a gene cluster comprised of *adeABC* and *adeRS*

(Figure 1A), which encode a tripartite RND efflux system, AdeABC, and its two-component

transcriptional regulator, AdeRS (refs). The 3' ends of *adeC* in the *aedG* linked clusters were

formed by an inverted repeat sequence that was also found adjacent to a nearby IS*18* (Figure

1A) (8). The locations of these repeat sequences suggested that the intervening sequence,

comprised of the IS*18* transposase gene, *adeRS* and *adeABC*, may be mobilised by the IS*18*

transposase. These loci also carried a partial *adeS* gene (3' end) downstream of the *aedG*

homologs, that could be related to an alternative organisation of these genes in an ancestral

locus (Figure 1A).

We sought to find additional homologs of *aedG* and determine whether they are also

associated with IS*18/adeRS/adeABC*. We initially queried the NCBI nr/nt database using

tblastn with the AedG sequence encoded in *A. nosocomialis* AC1530 plasmid pAC1530 and

found close to 300 high scoring hits from strains designated in 23 *Acinetobacter* species and

Acinetobacter strains with no species designation. Surveys of the local genetic environments

of the high scoring hits showed one of two general arrangements. Either *aedG* was located

downstream of the IS*18/adeRS/adeABC* cluster, as in pAC1530, or *aedG* was located between

homologs of *adeAB* and *adeRS* gene pairs and *adeC* and the IS*18* transposase gene were

absent, as in *A. baumannii* ABF9692 plasmid pABF9692 (Figure 1B). It is not unusual for *adeC*

to be absent from *adeAB* loci in *Acinetobacter* genomes, and AdeAB can function with AdeK

as an alternative outer membrane protein (8, 10). Insertion elements were also seen in some

of the loci, such as that in *A. lwoffii* strain AL_065 pAL_065-2, where insertion sequence

elements appear to have disrupted *adeS* (Figure 1B).

To identify additional homologs of AedG, we queried the NCBI experimental clustered protein

database using the amino acid sequence of AedG encoded by pAC1530. This search identified

69 clusters whose representative members had above 65 % amino acid sequence identity

with the query AedG sequence across at least 95 % of its length. Proteins in the top scoring

cluster were encoded by *Acinetobacter* species. The local genetic environments of these

homologs resembled those of *aedG* in pAC1530 or pABF9692. Other high scoring clusters

included *Acinetobacter* proteins encoded by *aedG* homologs in an alternative genetic

environment. For example, the genetic environment of *aedG* homologs in the *Candidatus*

Acinetobacter avistercoris isolate 5402 metagenome assembled genome (MAG) and an *A.*

bereziniae GD03393 plasmid resembled that in pABF9692 but included *adeC* (Figures 1B and

S1). Of note, the *A. bereziniae* GD03393 plasmid harboured a second *aedG* homolog similar

to pAC1530 (Figure S1).

The proteins in lower scoring clusters were encoded by species outside the *Acinetobacter*

genus. The genes encoding these proteins were frequently adjacent to divergently

transcribed *tetR* family regulator genes that may control expression of the *aedG* homologs.

Some non-*Acinetobacter* aedG homologs were adjacent to RND efflux pump genes. For

example, the *aedG* homolog in the MAG of Proteobacteria bacterium isolate CTSoil_107

(likely *Pleomorphomonas* based on GyrB and RpoD sequence similarity) was adjacent to

distant homologs of *adeRS*/*adeAB* (Figure S2), and the *aedG* homolog from *Pelagibacterium*

sp. SCN 63-126 ABS74_C0016 was encoded adjacently to a novel RND efflux system (Figure

1B). These results demonstrate a tendency for genes encoding AedG to be co-localised with

those encoding RND efflux systems, specifically AdeAB(C) in *Acinetobacter* species. AedG and

AdeAB(C) may cooperate in the efflux of substrates from the cytoplasm to outside the cell.

A phylogenetic analysis was performed to examine the relationships of AedG homologs with

other members of the DHA2 family. This analysis showed that AedG homologs are most

closely related to the AmvA (also known as AedF) (9, 11) and SmvA (12) efflux pumps from

Acinetobacter and *Salmonella*, respectively (Figure 1C). Pairwise amino acid sequence

similarity scores confirmed the relatedness of AedG homologs and AmvA/SmvA (Figure S3).

<u>AedG confers resistance to multiple antibiotics, biocides and dyes</u>

Based on the proximity of *aedG* genes to homologs of *adeAB(C)* in *Acinetobacter* species and

the similarity of AedG variants to drug transporters in the DHA2 family, it seemed likely that

AedG pumps would mediate resistance to one or more antimicrobials. We profiled the

antimicrobial resistance potential of the AdeG pump encoded in pAC1530 to eight

antimicrobials and compared resistance levels to those conferred by the multidrug efflux

pump AmvA in the same expression system (9, 11). The *aedG* homolog from pAC1530 was

synthesised and cloned into the pTTQ18 expression plasmid (Amp^r) behind the *tac* promoter

(13) with a C-terminal RGSH6 tag and sequence verified (Twist Bioscience). The *amvA* gene

from *A. baumannii* ATCC 17978 (also known as AedF) was cloned previously (9). AedG

conferred increased resistance to all eight antimicrobials tested when expressed in *E. coli*

BL21 cells and promoted higher levels of resistance to tetracycline, DAPI, ethanol and

ethidium bromide than AmvA (Figure 2A). These results suggest that AedG can recognise a

broad spectrum of antimicrobials similar to AmvA and is a new multidrug resistance protein

in *Acinetobacter*.

<u>AedG functions via an active efflux mechanism</u>

The ability of AedG to confer resistance to the nucleic acid intercalating dye ethidium bromide

allowed us to test efflux using real-time fluorometric transport assays. *E. coli* BL21 cells

expressing AedG, AmvA or no additional protein (negative control) were loaded with 10 µM

ethidium bromide in the absence of energy. The cells were washed, and re-energised and

ethidium efflux monitored by fluorescence changes. Both AmvA and AedG promoted the

active export of efflux ethidium bromide, as seen in fluorescence decrease over time relative

to control cells (Figure 2B). This result confirms efflux as the mechanism of resistance

mediated by AedG and that AedG is a new multidrug efflux pump.

Conclusions

The genes encoding AedG are part of the *Acinetobacter* accessory genome. This is in contrast

to AmvA, its closest relative, which is encoded in the core genome of *A. baumannii* and many

other *Acinetobacter* species. AmvA expression is not readily responsive to antimicrobial

treatment, and recent work showed that it may have a physiological function in polyamine

efflux (14). The organisation of *aedG* genes in *Acinetobacter*, downstream of *adeAB(C)* and in

proximity to *adeRS* homologs (Figures 1A and 1B), suggests that these genes may be

controlled by AdeRS and co-expressed with *adeAB(C)*. As such AedG homologs may play a

more significant role in antimicrobial resistance than AmvA in isolates that carry them.

Acknowledgements

This work was supported by an Australian Research Council Future Fellowship to KAH

(FT180100123) and a Project Grant from the National Health and Medical Research Council

of Australia to ITP and KAH (GNT1120298).

Figure 1. (A) Organisation of the local genetic environment of the *aedG* homolog in *A.*

nosocomialis AC1530 plasmid pAC1530 and related plasmids, including the locations of

surrounding genes and inverted repeat (IR) sequences that flank the IS*18* transposase gene

and *adeC*. (B) Alignment of loci carrying an *aedG* homolog in selected bacterial isolates.

Genes highlighted by colour and labels are found in more than one of the loci shown, or are

features noted in the text. This figure was made using clinker v0.0.25 (14). (C) The amino

acid sequences of AedG homologs from the lineages represented in (B) were aligned to the

sequences of other representative DHA2 family proteins using MUSCLE (15), and phylogeny

inferred using MrBayes v3.2.6 (16). The node labels show posterior probability values.

Figure 2. (A) *E. coli* BL21 cells carrying empty pTTQ18-*aedG* or pTTQ18-*amvA* were examined

in minimum inhibitory concentration (MIC) assays for resistance to a broad range of

antimicrobials, including the antibiotics tetracycline (TET) and erythromycin (ERY); the

biocides benzalkonium chloride (BAC), chlorhexidine (CHX), and ethanol (EtOH); and the

antimicrobial dyes 4',6-diamidino-2-phenylindole (DAPI), acriflavine (ACR) and ethidium

bromide (EtBr). Resistance assays were performed as previously described (9). (B) Ethidium

transport from *E. coli* BL21 cells carrying either pTTQ18, pTTQ18-*amvA*, or pTTQ18-*aedG*. Cells

were loaded with 10 µM ethidium, transport was initiated by adding glucose and efflux

monitored fluorescently (Ex:530nm; Em:610nm) in a PTI QuantaMaster 8000 (HORIBA

Scientific) as described previously (9).

Figure S1. Local genetic organisation of the *adeG* loci in *A. bereziniae* GD03393 plasmid. One gene cluster shows high sequence similarity to the *aedG* locus in *A. nosocomialis* AC1530 pAC1530 and an adjacent cluster shows high similarity to the *aedG* locus in the *Candidatus* Acinetobacter avistercoris 5402 metagenome assembled genome. This figure was made using clinker v0.0.25 (14).

Figure S2. (A) The phylogenetic tree of AdeB homologs from representative *Acinetobacter*

lineages (Figure 1B). The MAG of Proteobacteria bacterium isolate CTSoil_107, the *A.*

baumannii ATCC 17978 chromosome, and the AdeJ sequence from the *A. baumannii* ATCC

17978 chromosome were aligned using MUSCLE (15). The phylogeny of these sequences was

inferred using MrBayes v3.2.6 (16). The node labels show posterior probability values. (B)

Pairwise amino acid sequence similarity (bottom left) and identity (top right) scores for the

proteins shown in (A). These values were determined using MatGat version 2.0 (17) with a

BOLSUM 50 scoring matrix.

Figure S3. Pairwise amino acid sequence similarity (bottom left) and identity (top right)

scores for members of the DHA2 family of efflux proteins. These values were determined

using MatGat version 2.0 (17) with a BOLSUM 50 scoring matrix.

References

1.   Brzoska AJ, Hassan KA, de Leon EJ, Paulsen IT, Lewis PJ. Single-step selection of drug

resistant Acinetobacter baylyi ADP1 mutants reveals a functional redundancy in the

recruitment of multidrug efflux systems. Plos One. 2013;8(2):e56090.

2.   Hassan KA, Maher C, Elbourne LD, Henderson PJ, Paulsen IT. Increasing the PACE of

characterising novel transporters by functional genomics. Curr Opin Microbiol. 2021;64:1-8.

3.   Henderson PJF, Maher C, Elbourne LDH, Eijkelkamp BA, Paulsen IT, Hassan KA.

Physiological Functions of Bacterial "Multidrug" Efflux Pumps. Chem Rev. 2021;121(9):5417-

78.

4.   Bi D, Zheng J, Xie R, Zhu Y, Wei R, Ou HY, et al. Comparative Analysis of AbaR-Type

Genomic Islands Reveals Distinct Patterns of Genetic Features in Elements with Different

Backbones. mSphere. 2020;5(3).

5.   Guardabassi L, Dijkshoorn L, Collard JM, Olsen JE, Dalsgaard A. Distribution and in-

vitro transfer of tetracycline resistance determinants in clinical and aquatic Acinetobacter

strains. J Med Microbiol. 2000;49(10):929-36.

6.   Levy SB, McMurry L. Plasmid-determined tetracycline resistance involves new

transport systems for tetracycline. Nature. 1978;276(5683):90-2.

7.   Li L, Hassan KA, Brown MH, Paulsen IT. Rapid multiplexed phenotypic screening

identifies drug resistance functions for three novel efflux pumps in Acinetobacter

baumannii. J Antimicrob Chemother. 2016;71(5):1223-32.

8.   Varsha Naidu AB, Anthony J. Brzoska, Peter Lewis, Bart A. Eijkelkamp, Ian T. Paulsen,

Liam D.H. Elbourne, Karl A. Hassan. Evolution of RND efflux pumps in the development of a

successful pathogen. Drug Resistance Updates. 2023;66: 100911.

9.   Hassan KA, Brzoska AJ, Wilson NL, Eijkelkamp BA, Brown MH, Paulsen IT. Roles of

DHA2 family transporters in drug resistance and iron homeostasis in Acinetobacter spp. J

Mol Microbiol Biotechnol. 2011;20(2):116-24.

10.   Sugawara E, Nikaido H. Properties of AdeABC and AdeIJK efflux systems of

Acinetobacter baumannii compared with those of the AcrAB-TolC system of Escherichia coli.

Antimicrob Agents Chemother. 2014;58(12):7250-7.

11.   Rajamohan G, Srinivasan VB, Gebreyes WA. Molecular and functional

characterization of a novel efflux pump, AmvA, mediating antimicrobial and disinfectant

resistance in Acinetobacter baumannii. J Antimicrob Chemother. 2010;65(9):1919-25.

12.   Santiviago CA, Fuentes JA, Bueno SM, Trombert AN, Hildago AA, Socias LT, et al. The

Salmonella enterica sv. Typhimurium smvA, yddG and ompD (porin) genes are required for

the efficient efflux of methyl viologen. Mol Microbiol. 2002;46(3):687-98.

13.   Stark MJ. Multicopy expression vectors carrying the lac repressor gene for regulated

high-level expression of genes in Escherichia coli. Gene. 1987;51(2-3):255-67.

14.   Gilchrist CLM, Chooi YH. Clinker & clustermap.js: Automatic generation of gene

cluster comparison figures. Bioinformatics. 2021.

15.   Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and

space complexity. BMC Bioinformatics. 2004;5:113.

16.   Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogenetic trees.

Bioinformatics. 2001;17(8):754-5.

17.   Campanella JJ, Bitincka L, Smalley J. MatGAT: an application that generates

similarity/identity matrices using protein or DNA sequences. BMC Bioinformatics. 2003;4:29.