The gut metagenome harbors metabolic and antibiotic resistance signatures of

moderate-to-severe asthma

Naomi G. Wilson[1], Ariel Hernandez-Leyva[1], Drew J. Schwartz[2], Leonard B. Bacharier[3],

Andrew L. Kau[1,*]

[1] Division of Allergy and Immunology, Department of Medicine and Center for Women's

Infectious Disease Research, Washington University School of Medicine, St. Louis, MO,

63110, USA

[2] Division of Infectious Diseases, Department of Pediatrics and Center for Women's

Infectious Disease Research, Washington University School of Medicine, St. Louis, MO,

63110, USA

[3] Division of Allergy, Immunology and Pulmonary Medicine, Department of Pediatrics,

Monroe Carell Jr Children's Hospital at Vanderbilt University Medical Center, Nashville,

TN 37232, USA

* Correspondence: akau@wustl.edu

Abstract

Asthma is a common allergic airway disease that develops in association with the

human microbiome early in life. Both the composition and function of the infant gut

microbiota have been linked to asthma risk, but functional alterations in the gut

microbiota of older patients with established asthma remain an important knowledge

gap. Here, we performed whole metagenomic shotgun sequencing of 95 stool samples

from 59 healthy and 36 subjects with moderate-to-severe asthma to characterize the

metagenomes of gut microbiota in children and adults 6 years and older. Mapping of

functional orthologs revealed that asthma contributes to 2.9% of the variation in

metagenomic content even when accounting for other important clinical demographics.

Differential abundance analysis showed an enrichment of long-chain fatty acid (LCFA)

metabolism pathways which have been previously implicated in airway smooth muscle

and immune responses in asthma. We also observed increased richness of antibiotic

resistance genes (ARGs) in people with asthma. One differentially abundant ARG was a

macrolide resistance marker, *ermF*, which significantly co-occurred with the *Bacteroides*

fragilis toxin, suggesting a possible relationship between enterotoxigenic *B. fragilis*,

antibiotic resistance, and asthma. Lastly, we found multiple virulence factor (VF) and

ARG pairs that co-occurred in both cohorts suggesting that virulence and antibiotic

resistance traits are co-selected and maintained in the fecal microbiota of people with

asthma. Overall, our results show functional alterations via LCFA biosynthetic genes

and increases in antibiotic resistance genes in the gut microbiota of subjects with

moderate-to-severe asthma and could have implications for asthma management and

treatment.

Importance:

Asthma is an airway disease that affects the everyday lives of millions of people

and accounts for approximately 1.5 million emergency room visits yearly in the US(1)

Both antibiotic usage and gut microbiota dysbiosis have been linked to the development

of asthma, however, little is known about the specific gut microbial functions associated

with asthma, particularly in older populations. In this study, we characterize the gut

microbiota of school-aged children and adults with moderate-to-severe asthma to

uncover asthma-associated microbial functions that may contribute to disease features.

We find that people with asthma have an increase in gut microbial genes associated

with long-chain fatty acid metabolism as well as an accumulation of antibiotic resistance

genes, both of which may have practical consequences for monitoring and treatment of

asthma.

Introduction

Asthma is a common respiratory disease characterized by symptoms of airway

obstruction including wheeze, cough, and shortness of breath. In most cases, asthma

onsets in early childhood with the development of sensitization to environmental

allergens. Ongoing environmental exposures lead to airway inflammation and ultimately

result in asthma symptoms manifesting within the first few years of life. Recent findings

support the notion that asthma develops in association with the human gut microbiome

composition early in life(2, 3). This finding is supported by 16S rRNA sequencing

surveys demonstrating that alterations in the gut microbiota precede asthma

development within the first few months of life(2, 4).

Early childhood gut microbial communities have been proposed to contribute to

asthma by several mechanisms. Epoxide hydrolases encoded by enterococci and other

gut bacteria produce the lipokine 12,13-diHOME that predisposes towards atopic

sensitization and asthma(4, 5). Similarly, short-chain fatty acids (SCFAs), produced by

the metabolism of dietary fibers by diverse members of the gut microbiota, are thought

to protect from asthma through their effect on the host G-protein coupled receptor

GPR41, shaping immune cell differentiation in the lungs, and ameliorating allergic

airway inflammation(2, 6–9).

In addition to microbially-encoded metabolic features, carriage of antibiotic

resistance genes (ARGs) within the gut microbiota, termed the resistome, has been

associated with asthma risk. In infants, microbial signatures associated with the

development of asthma are also associated with increased richness of ARGs in the gut

microbiome(10). These differences in ARG carriage were found to be driven primarily by

E. coli, which is a common colonizer in the first days of life(10). These findings are

important in understanding the origins of asthma since antibiotic exposure correlates

both to the number of ARGs within the gut microbiome(11) and the later development of

asthma and other allergic diseases(12–14). This association between antibiotic

exposure and asthma is supported by animal models that found antibiotic treatment

worsens allergic airway inflammation (AAI)(15–17).

While there is an abundance of data supporting the idea that asthma

susceptibility is associated with features of the gut microbiota in early childhood, the

potential effect of gut microbial functions on asthma later in life remains an important

knowledge gap. Since asthma often begins in infancy when the gut microbiota

composition is highly unstable, disease-causing microbial functions may not persist into

older children and adults. Nevertheless, the gut microbiota in older individuals could

underlie the variable manifestations of asthma(18) and may hold valuable prognostic

and therapeutic significance.

Asthma-associated differences in later childhood and adult gut microbial

communities have already been noted in several reports. Studies in preschool-aged

children have noted distinct taxonomic composition of gut microbial communities in

asthmatic subjects compared to healthy controls(3). These differences are reported to

include reductions in *Akkermansia muciniphila*(19), *Faecalibacterium prausnitzii*(20) as

well as *Roseburia* species(21). Functional characterization of microbial communities by

whole metagenomic sequencing from an older population of asthmatic women(20) has

shown that pathways related to lipid and amino acid metabolism, as well as

carbohydrate utilization were enriched in asthmatics. In contrast, microbial pathways

involved in the production of SCFAs, like butyrate, were enriched in the healthy cohort

of the same study(20). These findings are supported by a complementary study

designed to test the effect of probiotic supplementation on asthma that found an

association of improved asthma symptoms with SCFA biosynthesis as well as

tryptophan metabolism pathways in the adult gut microbiota(22).

Here, we describe an analysis of whole metagenomic sequencing data from a

cohort of 36 subjects with physician-diagnosed, moderate-severe asthma along with a

matched cohort of 59 healthy controls. This study tests the hypothesis that the gut

metagenome harbors signatures of asthma later in life. Our results identify global

differences in metagenomic functions between healthy and asthmatic subjects and

reveal an enrichment in long-chain fatty acid biosynthetic pathways. We also find an

increased richness of ARGs in asthmatics and co-occurrence of ARGs with known

bacterial virulence factors, suggesting a potential relationship between antibiotic

exposure and pathogen colonization in asthmatics.

Results

Whole metagenomic shotgun sequencing of fecal samples from adults and

children with asthma and healthy controls

We performed whole metagenomic sequencing on fecal samples from asthmatic

subjects and healthy controls taking part in the Microbiome & Asthma Research Study

(MARS), which we have previously described(23, 24). MARS participants were recruited

from the St. Louis, Missouri area and included pediatric (6-10 years) and adult (18-40

years) age groups. All asthma cohort patients had a physician diagnosis of moderate-to-

severe asthma, and history of allergic sensitization as evidenced by positive skin testing

or serum specific-IgE to one or more common aeroallergens. In total, we analyzed 95

patient stool samples including 17 adults and 19 school-aged participants with asthma,

and 40 adults and 19 school-aged participants without asthma.

NovaSeq S4 sequencing of our libraries yielded 1.69 billion paired-end reads

translating to a total of approximately 500 Gigabases (Gb). After filtering for read quality,

dropping host contaminants, and trimming adaptor content, we achieved 1.23 billion

paired-end reads and an average 3.4 Gb per stool sample with a range of 0.4-9.9

Gb/sample (Figure S1A). Neither host contamination nor sequencing depth differed

between asthma and healthy cohorts (t-test p=0.2 and 0.7, Table S1). All samples

achieved an estimated average metagenomic coverage of at 89% (range of 61-98%)

with the annotation-free redundancy-based metagenome coverage estimator,

Nonpareil(25) (Figure S1B). Further, estimated metagenome coverage was not different

between the asthma and healthy cohorts, although we noted coverage was slightly

reduced in the pediatric cohort (Figure SB, Table S1). We employed the read-based

annotation pipeline, HUMAnN(26) to determine the abundance of genes and functional

pathways in the stool metagenomes. We found that the most abundant functional

pathways (Figure S1C) across all MARS participants are involved in essential

processes of gut microbes such as starch degradation and glycolysis, demonstrating

that our sequencing captured core functions of the gut metagenome, as expected.

Taken together, we concluded that our sequencing is of sufficient depth and quality to

be used for further analyses.

Gut taxonomic composition differs between people with and without asthma

We first leveraged the clade marker annotation tool, MetaPhlAn(26), to analyze

the taxonomic composition of the study participants. We found dominate genera typical

in gut microbiota communities including *Bacteroides* (phylum Bacteroidota) and

Faecalibacterium (phylum Bacillota) (Figure S1D). Simpson alpha diversity was slightly

higher in the asthma cohort even when taking read depth and age group into account

(Figure S1E). Bray-Curtis dissimilarity (Figure 1F) was shifted between the asthma and

healthy cohorts (p<0.0004, R^2=0.029) even when accounting for other covariates

including age (p<0.001, R^2=0.032), race (p=0.0006, R^2=0.026), recent antibiotic usage

(p=0.9, R^2=0.006), read depth (p=0.2, R^2=0.013), obesity (p=0.7, R^2=0.008), sex (p=0.4,

$R^2=0.011$), and tobacco exposure (p=0.2, $R^2=0.012$) by sequential PERMANOVA

(Figure 1G). There was also no significant interaction between asthma status and age

group (p=0.8, $R^2=0.007$), or between asthma status and recent antibiotic usage (p=0.6,

$R^2=0.009$) (Figure S1G). To determine differentially abundant taxa, we tested the fixed

effect of asthma along with the random effects of age group and race in a general linear

model(27) and found *Eubacterium rectale* and *Prevotella copri* were enriched in the

healthy cohort (Figure S1H, Table S2). All of these findings are consistent with 16S

rRNA sequencing performed in a previous study(23) which lent us further confidence

that our sequencing data was suitable for functional profiling.

Fatty acid metabolism pathways are enriched in the gut metagenomes of people

with asthma

Given that our samples had adequate coverage to capture expected taxonomic

shifts, we started interrogating the differences in metagenomic functions of the gut

microbiota attributable to asthma status. The alpha diversity of genes (UniRef90

clusters) was neither different between the asthma and healthy cohorts nor between the

pediatric and adult cohorts, suggesting that our gene profiling reached a similar total

number of genes in both cohorts (Figure 1A). Using PERMANOVA, we noted that, even

while accounting for significant covariates of age (p<0.001, $R^2=0.029$), race (p<0.001,

$R^2=0.024$), and read depth (p=0.03, $R^2=0.015$), asthma status also significantly

impacted gut microbiome functional composition (p=0.008, $R^2=0.017$; Figure 1B, C). We

note that age group's interaction term with asthma did not significantly contribute to the

variance in beta diversity, suggesting that the influence of asthma and age on beta

diversity is non-overlapping. These findings support the idea that the gut metagenomic

content of people with asthma is different than that of healthy individuals, even when

accounting for other clinical sources of interpersonal gut microbiome variation.

We next considered which metagenomic functions and metabolic pathways may

be involved in the differences between asthma and healthy cohorts. We first examined a

list of specific metagenomic functions previously implicated in asthma, including genes

related to histamine production, 12-13 diHOME biosynthesis, and tryptophan

metabolism, but we were unable to identify a difference between cohorts (Figure S2A).

To identify pathways that differed between asthma and healthy subjects, we performed

a Wilcoxon Rank Sum test with a false discovery rate q<0.2 on the relative abundance

of all pathways annotated by the MetaCyc database that were above 10% prevalence

within the population. Using these criteria, we found seven pathways that were enriched

in asthma and one that was enriched in the healthy cohort out of 312 total pathways

(Figure 1D). To determine if these findings were robust to other analysis methods, we

performed additional differential abundance approaches on the 312 MetaCyc pathways,

including a Wilcoxon test on centered log-transformed counts and ALDEX2, both of

which demonstrated that these pathways differed between healthy and asthmatic

cohorts (See Table S3). All differentially abundant pathways enriched in patients with

asthma were involved in fatty acid synthesis, and included the production of oleate,

palmitoleate, (5Z)-dodecenoate, 8-amino-7-oxononanoate, biotin, and octanoyl acyl-

carrier protein, as well as saturated fatty acid elongation. In the healthy cohort, only a

single L-lysine biosynthesis pathway was enriched.

Using taxonomically tiered functional mapping, we determined which taxa were

driving the observed differences in asthma-associated pathways. For the L-lysine

biosynthesis III pathway which was more abundant in healthy subject, we found that it

primarily originated from *Blautia obeum*, Figure S2B). In the case of the asthma-

enriched pathways, we found that *Bacteroides vulgatus* and *Alistipes finegoldii* account

for the largest fraction of complete fatty acid biosynthesis pathways (Figure 1E, Figures

S3C). However, the differential abundance of these asthma-associated pathways was

probably not due solely to an enrichment of *B. vulgatus* or *A. finegoldii* in asthma stool

since neither species was differentially abundant (maaslin2 q-value=0.58 and 0.25,

respectively; See Table S2). Further, the majority of mapped pathways were not

attributable to any single species and these unmapped pathway counts made up more

of the overall pathway richness than *B. vulgatus* (Wilcoxon q values < 0.05 for all seven

pathways; see "Community" stratification in Figure S2C). Taken together, these findings

indicate that the differences may be either driven by community-level effort (i.e. distinct

steps of the pathway are encoded across more than one species), or that current

databases are insufficiently granular to identify the key taxa responsible for these

differences.

We reviewed the enzymatic steps of each of the eight pathways represented in

Figure 1D and found that, of the 78 total reactions in these pathways, only 11 reactions

were shared between 2 pathways (Figure S3). The 8-amino-7-oxononanoate

biosynthesis I pathway consists of the first 11 reactions of the larger biotin biosynthesis

pathway and the latter only has four additional reaction steps past synthesizing 8-

amino-7-oxonanoate to produce biotin. Additionally, the (5Z)-dodecenoate pathway can

feed directly into the palmitoleate biosynthesis pathway, and that the octanoyl acyl

carrier protein pathway shares an upstream substrate (acetoacetyl-acyl carrier protein)

with the saturated fatty acid elongation pathway (Figure S3). Together, our findings

indicate that long chain fatty acid biosynthesis is differentially abundant in the asthma

gut metagenome via related but largely non-redundant pathways.

Given the association between obesity with fatty acid metabolism(28) as well as

asthma(29–31), we next wanted to determine whether obesity (which we define here as

a BMI greater than 30 in adults or a BMI-for-age percentile of greater than 95% in

children) confounds the association of microbial fatty acid metabolism with asthma. We

compared the abundance of the differentially abundant fatty acid pathways between all

non-obese and obese patients and found no significant difference (Figure 1F). Within

the asthma cohort, there was similarly no statistically significant difference between the

asthmatic obese and asthmatic non-obese patients, suggesting that obesity is not a

confounder for the difference we observed in fatty acid metabolism. To determine

whether fatty acid metabolism is related to the intensity of asthma symptoms and their

effect on everyday life activities, we utilized a validated survey of asthma control (The

Asthma Control Test; ACT)(32). None of the fatty acid pathways were differentially

abundant between well-controlled and poorly-controlled asthmatics (Figure 1F). We

tested if age group affects the differentially abundant metabolic pathways and found that

these pathways were not differentially abundant between age groups alone (Figure 1F).

We also tested the impact of asthma and age as independent variables to differentially

abundant metabolic pathways using a Two-way ANOVA. We found that, even while

taking age into account, these pathways are differentially abundant between asthma

and healthy cohorts, but are not different by age or an interaction between asthma and

age (Figure 1G, 2-Way ANOVA). Given that the effect of asthma status on differentially

abundant metagenomic functions was distinct from that of age, we primarily focused our

subsequent analyses on the asthma and healthy cohorts overall, combining age groups.

Richness of antibiotic resistance genes is increased in the gut metagenomes of

people with asthma

Since people with asthma tend to be prescribed antibiotics frequently(33) and

oral antibiotic exposure is a risk factor for the acquisition of ARGs in the gut(11), we

wanted to determine if the members of our asthma cohort were more likely to have

received antibiotics. To test this, we counted how many subjects had taken a course of

antibiotics within one year of their participation in the study. As part of the study design,

participants could not take antibiotics in the month prior to fecal donation. We found that

a greater proportion of the asthma cohort received antibiotics in the past year compared

to that of healthy participants (42% of asthma cohort versus 15% of the healthy cohort,

Fisher's test, p=0.011, Figure 2A). This finding represents evidence of increased

antibiotic exposure amongst subjects with asthma in our study.

We next sought to characterize the gut antibiotic resistome in the asthma and

healthy cohorts. To test if the increased antibiotic exposure in the asthma cohort was

reflected in the gut resistome, we utilized the ShortBRED pipeline(34) to detect reads

mapped to the Comprehensive Antibiotic Resistance Database (CARD)(35). We first

asked whether there were more ARGs in our asthma cohort by summarizing our dataset

into richness (Total number of unique ARGs detected per sample) and load (Total sum

of ARG RPKM per sample). We found that ARG richness was higher in people with

asthma even when accounting for differences due to age (p=0.03) and sequencing

depth (p=0.09 while ARG load was not different between asthma and healthy cohorts

(p=0.4) when accounting for age (p<0.001) and read depth (0.002) (Figure 2B). We note

that *E coli* was not differentially abundant between asthma and healthy cohorts (p=0.52,

Table S2), so the richness increase we observe in the asthma cohort is not due solely to

an increase in *E. coli* relative abundance. These results suggest that there are a higher

number of unique ARGs, or a higher diversity, in asthma compared to healthy controls.

From our 95 stool samples, we detected 71 unique ARGs, comprising 32

antimicrobial resistance families, 29 drug classes, and 7 mechanisms of resistance, with

26 ARGs (37% of the total) conferring multi-drug resistance (Figure 2C). Similar to

previous studies of gut resistomes, we found that tetracycline resistance markers were

the most commonly detected ARGs and inactivation is the most common mechanism of

resistance followed by efflux pumps(10) (Figure 2C). Using the abundance data of each

detected ARG, we determined that asthma (p=0.005, R^2=0.028) and age (p<0.001,

R^2=0.053) were the strongest factors contributing to the variance in ARG beta diversity

even when accounting for important technical and demographic covariates (Figure 3A

and 3B). We next wanted to ascertain to what degree the resistome profile was

determined by microbial composition. We used a Procrustes analysis(36) to compare

compositional data generated from MetaPhlAn(26) to the antibiotic resistome profile

derived from ShortBRED and found that the microbiome composition correlated to the

resistome profile (Figure 3C, PROTEST corr = 0.627, p-value < 0.0001), indicating that

ARG profiles are directly related to bacterial species composition.

Macrolide resistance markers are differentially abundant in asthma

To determine ARGs that are differentially abundant between asthmatic and

healthy gut metagenomes, we applied negative binomial tests to the abundance of all

ARGs detected in at least 7 samples. This prevalence cutoff was chose because it is

the minimum number of samples needed to detect a difference using a negative

binomial distribution. We found that genes encoding resistance to macrolides (*ermF,*

ermB and *ermA*), vancomycin (*vanRO*), tetracycline (*tet(45)*), as well as multi-drug

efflux pumps (*smeB, mdtO*, and *oqxA*) were enriched in the asthma cohort (Figure 4A,

Table S4). Prominent amongst these was the 23S rRNA methyltransferase *ermF*, which

is typically encoded by *Bacteroides* species and confers resistance to macrolides.

Next, we explored the genomic context of *ermF* by assembling metagenomic

sequencing reads into contigs with metaSPAdes(37) and annotating open reading

frames with Prokka(38) and BLAST. We detected full-length *ermF* with 98% or higher

identity in 53 out of 95 samples. Out of 53 contigs, the vast majority originated from

members of the Bacteroidota, 75.4% originated from the *Bacteroides* genus and 60.3%

of them were likely from *B. fragilis* based on the top BLAST homology. Of the contigs

that encoded *ermF,* 68% occurred on scaffolds with at least one other open reading

frame within ten kilobases (Figure 4B). We found that many *ermF genes* are co-located

with genes associated with mobile genetic elements such as transposases, mobilization

genes, and toxin/antitoxin systems, as well as with other ARGs like *btgA* which encodes

clindamycin resistance (Figure 4B,C). This indicates that *ermF* occurs in multiple

different genomic contexts within our cohort and suggests that its presence is not strictly

due to propagation of a single *B. fragilis* strain.

People with asthma have a distinct set of co-existing pairs of antibiotic resistance

genes and virulence factors in the gut metagenome

In our prior work on this same cohort of patients, we found that, compared to

healthy subjects, a greater portion of asthma subjects were colonized with *B. fragilis*

strains harboring the virulence factor *B. fragilis* toxin (*bft*), which we showed has the

potential to shape inflammation in the lung(23). Given that our resistome analysis

pointed to an enrichment of a *B. fragilis* ARG, we wanted to test whether the *ermF* gene

is co-selected with *bft.* When only taking metagenomes encoding *ermF* into account, we

observed an enrichment of *bft* prevalence in the asthma cohort (Figure 4D, p=0.009). In

contrast, among metagenomes with no detectable *ermF,* there is no enrichment of *bft* in

the asthma cohort (Figure 4D, p=0.39). When reviewing the entire MARS population, we

found no statistically significant co-occurrence of *ermF* with *bft* (Figure 4E; one-tailed

Fishers test p>0.05) and this was consistent with healthy gut metagenomes from the

Human Microbiome Project (Figure 4E, Fisher's test p=0.56). However, in our MARS

samples, we did not find any instances where *bft* and *ermF* occurred on the same

scaffold, so it remains unclear whether these two genes are encoded within the same *B.*

fragilis strain or within two separate strains. Nevertheless, these results suggest that the

environment supporting the gut microbiota of asthmatic individuals presents

opportunities or niches for *ermF* and *bft* to co-occur.

To explore the possibility that virulence traits and ARGs are linked in the gut

microbiota, we characterized virulence factor (VF) content of all samples using the

Virulence Factor Database(39) and compared these data to the antibiotic resistome

profiles. We did not find the same overall shift in the virulence factor beta diversity

between asthma and healthy that we observed with the resistomes (Figure S4A-C), but

we did find differentially abundant VFs belonging to capsule and peritrichous flagella VF

families (Table S5, q values<0.2). Further, we found that microbiota composition is

highly correlated with virulence factor profile (Figure S4D, Protest correlation

coefficient=0.61, p<0.0001). Given that microbiota composition strongly affects both VF

and ARG content, we used a partial correlation between VF and ARG richness to test

our hypothesis while removing the effect of total metagenomic content. We found a

positive partial correlation between VF and ARG richness in both the asthma and

healthy cohorts (Figure 5A). Similarly, virulence factor and resistome beta diversity

profiles were also positively correlated (Figure 5B, Protest correlation coefficient=0.574,

p=1e-4). Together, our results suggest that these two microbial features, virulence and

antibiotic resistance, are closely linked within the gut metagenome.

We next performed a co-occurrence analysis to uncover other linked virulence

and antibiotic resistance traits that could be important in gut ecology. We found

numerous co-occurring VF-ARG pairs in MARS gut metagenomes (Figure 5C, p<0.05).

Several of these positively co-occurring pairs were shared between the two cohorts

(yellow), suggesting that these relationships are not dependent on asthma status. In

contrast, many pairs specifically co-occur in one cohort and may indicate microbial

interactions important in asthma but not healthy gut metagenomes (Figure 5C). In

summary, we found that VF and ARG presence is linked in the gut metagenome and

that people with asthma have a distinct set of co-occurring functions compared to

healthy people.

While our co-occurrence analysis between VFs and ARGs demonstrated multiple

examples of virulence and antibiotic resistance traits found in the same gut

metagenome, this analysis does not indicate if these genes are present in a single

organism. To obtain a more granular view of VF-ARG co-occurrence, we limited our

analysis to look for VF-ARG pairs that could be encoded by the same species. This

analysis showed that the asthma cohort had a greater number of ARGs (p=0.007 and

0.01) and VFs (p=0.005 and 0.09) annotated as coming from *Klebsiella pneumoniae*

and *Escherichia coli*, respectively (Figure S5A). Individual co-occurrences attributable to

each of these species are summarized in Figure S5B and show that *cepA*, encoding a

beta-lactamase, and *chuU*, a VF involved in iron acquisition, are both putatively

encoded by *E. coli* and co-occur in asthmatics, suggesting that the metagenome-wide

co-occurrence of CepA and Chu families observed in Figure 5C may be due to

enrichment within one or more *E. coli* strains harboring these VF/ARG pairs. Together,

our co-occurrence analyses show that there appear to be multiple co-occurring VFs and

ARGs, similar to *B. fragilis*-encoded *bft* and *ermF*, in the gut metagenome and within

putative individual species that could be important for asthma. The cohort-specific co-

occurring VF-ARG pairs found here could serve as candidates for future studies of

asthma gut microbiome ecology.

Discussion

In this study, we present an exploratory analysis of fecal whole metagenomic

sequencing contrasting subjects with moderate-to-severe asthma to a group of healthy

controls to identify disease-associated microbial genes with the strongest likelihood of

affecting disease. Our sequencing and subsequent analyses revealed that the

functional content of individuals with asthma differed significantly from that of healthy

controls. We found an enrichment of functions associated with saturated and mono-

unsaturated fatty acids, including oleate, palmitoleate, 5(Z)-dodecenoate, biotin, 8-

amino-oxononanoate, saturated fatty acid elongation, and octanoyl acyl carrier protein

pathways. Currently, the functional significance of gut bacterial synthesis of these long-

chain fatty acids (LCFA) to asthma has not been well defined. Excess LCFAs, usually

studied in the context of dietary fat intake, have been associated with metabolic

diseases including diabetes, obesity, and atherosclerosis risk(30) but is also linked to

asthma risk in adults(29–31, 40). Increasing recognition that obesity predisposes to

asthma has motivated investigation of the impact of fatty acids on airway biology and

has shown that LCFA signaling through free fatty acid receptor 1 (FFAR1, also called

GPR40) induces airway smooth muscle cell contraction and proliferation, both of which

are important components of asthma pathophysiology(30, 41). Notably, a study that

sequenced airway microbes in children with cystic fibrosis implicated a similar list of

LCFA production pathways during exacerbations, suggesting that microbially produced

LCFAs may influence airway physiology(42). To our knowledge, the potential for gut

microbes to contribute to the amount of free fatty acids available to the lung has not yet

been defined, however, LCFAs are readily absorbed into the circulation(43) and could

plausibly reach the airways. Further, previous studies have shown the effect of SCFA

(e.g. acetate, butyrate, propionate) produced by gut microbes to directly alter lung

inflammation via GPR41 (FFAR3)(8, 9). While our study did not find a direct enrichment

of SCFA production pathways in the healthy cohort as has been previously

reported(20), we did observe that lysine biosynthesis was enriched. Since lysine may

serve as a precursor to the SCFA butyrate(44), SCFAs may still be more abundant in

our healthy cohort but may be subject to transcriptional regulation that would not be

detected by metagenomic DNA sequencing. Together, our metabolic pathway analyses

of the gut metagenome demonstrate a positive association between LCFAs produced

by gut microbes and asthma, in contrast to the negatively associated SCFAs.

In addition to metabolic alterations, analysis of the gut resistome demonstrated

that subjects with asthma had a distinct ARG composition. In a recently published

prospective gut metagenomic study of infants, asthma-associated taxonomic signatures

were associated with a higher number of ARGs(10). These differences in the resistome

were largely driven by a single species of bacteria, *E. coli*, and reveals that acquisition

of ARGs in subjects with asthma may begin in early childhood and could affect asthma

development. In our study of older subjects with established asthma, we similarly found

a higher richness of ARGs that is associated with asthma in both school-aged children

and adults, supporting the idea that increased ARG carriage may persist in people with

asthma throughout life. Based on our resistome annotation, however, ARGs in our

cohort were likely from a diverse assemblage of bacteria in contrast to what was

observed in infants. This is likely due to differences in gut dynamics between age

groups. The infant microbiome is heavily shaped by limited available niches in the

developing gut, which favor transient, facultative anaerobes like *E. coli* (10)*,* whereas

the gut resistome in older subjects reflects selective pressures experienced over a

lifetime. One important consequence of increased richness of ARGs in people with

asthma is that it may promote persistence of some bacterial strains(45, 46) and

contribute to the taxonomic differences in the gut microbiota between asthma and

healthy people(3, 23).

While asthma was among the important factors accounting for a significant

amount of the variance in ARG beta diversity, we found that recent antibiotic exposure

(within the past year) was not. Notably, no participant in our cohort received a course of

antibiotics in the month prior to fecal sampling since this could have confounded our

analyses on asthma-associated microbial community changes. Previous studies have

shown that the gut microbiota recovers in approximately a month after perturbation from

antibiotics in healthy adults(47). We interpret these findings to mean that recent

exposure (within 1 - 12 months) to antibiotics does not drastically change the resistome,

whereas repeated exposures over time may be more important for driving the

population-wide shifts we observed in our cohort(46).

Of the ARGs found to be enriched within asthmatic resistomes, the ARG *ermF*,

encoding resistance to macrolide antibiotics, was especially prominent amongst the

asthmatic cohort. While we did not collect data on the antibiotic drug classes, number of

courses and their duration, or the reason for prescription of antibiotics, our subjects

received, it is likely that our asthma population has been exposed to macrolides.

Macrolide antibiotics, including clarithromycin and azithromycin, are commonly

prescribed for upper and lower airway infections which disproportionately affect people

with asthma(48). This class of antibiotics, particularly azithromycin, have been a focus

of special concern for driving antibiotic resistance due to their frequent usage and

pharmacological properties(49–51). Nevertheless, azithromycin has been noted to have

beneficial effects in asthma, and some(52), but not all(53), studies suggest that

azithromycin may prevent exacerbations in asthmatics. Given the interest in

azithromycin as a treatment modality in asthma, there will be an urgent need for

additional studies to determine the robustness of the association between asthma and

macrolide ARG accumulation in the gut to inform parameters for antibiotic selection and

prescription in people with asthma.

Additional exploration of the gut metagenomes revealed potential co-selection in

people with asthma for *B. fragilis* genes *ermF* and *bft* (*B. fragilis* toxin), the latter of

which is more prevalent in fecal samples from the asthma compared to healthy

cohort(23). Untargeted analysis of gut resistomes revealed multiple examples of

virulence factor and ARG co-occurrence as well as positive correlations between ARG

and VF richness in people with and without asthma**.** Our findings are consistent with

previous reports that found correlations between VFs and ARG richness and VF-ARG

cooccurrence relationships in both gut metagenomes(54) and human-associated

bacterial genomes(55). Our findings also add to these studies by demonstrating that,

while the correlation between VF and ARG richness does not appear to be any stronger

in the asthma cohort after taking gene richness into account, the two MARS cohorts do

not have identical sets of statistically significant co-occurring VF-ARG pairs. These data

suggest that people with asthma may be experiencing different selection pressures from

that of healthy people, leading to accumulation of a distinct set of virulence and

antibiotic determinants. Given that antibiotics induce gut inflammation through the

disruption of the gut microbiota(56), and strains encoding virulence factors such as *bft*

are known to thrive in an inflammatory environment(57), one plausible model for the

apparent accumulation of distinct VF-ARG pairs is that antibiotic treatment not only

selects for ARGs(11, 46), but simultaneously selects for VFs. Together with evidence

that virulence determinants, such as *bft,* are associated with airway inflammation(23),

our model implies that heightened antibiotic treatment may contribute to the

manifestations of asthma via co-selection for VFs and ARGs. Considering that prenatal

and early life antibiotic exposure is linked to asthma risk(13, 56), this model could be

used to test whether the initial events driving VF and ARG co-occurrence start with the

first vertical transmission events in very early life.

Our study has several limitations that constrain the scope of our claims. First,

MARS is an exploratory, cross-sectional study with only a moderate number of subjects

recruited from a single site, which is less ideal for identifying disease-associated

microbiome differences(58). As a result, our study had limited statistical power to detect

less prevalent or abundant functions. Second, our study focused on school-aged and

older subjects with moderate-to-severe asthma, and thus our findings may not be

applicable to other younger populations or those with less severe disease. These

population differences may explain why we were unable to identify statistically

significant differences in microbial metabolic pathways identified from other studies

including bile acid metabolism(2), epoxide hydrolases(5), histamine metabolism(59, 60),

or tryptophan metabolism(61, 62) (Figure S2A). Third, the factors driving the shift in gut

bacterial metabolism to LCFA biosynthesis and whether gut microbiome enrichment of

this pathway is sufficient to change the hosts' LCFA profile is not known. Collecting

blood to interrogate host metabolism as well as dietary information at the time of fecal

sample collection would have helped to disentangle the effects of diet on host and gut

microbiota metabolism. Fourth, a record of the frequency and class of antibiotics

administered to our participants would have allowed us to confirm whether macrolide

administration associates with the enrichment of *ermF* in our asthma cohort and

whether a higher diversity of antibiotic usage correlates with ARG richness. It is likely

that antibiotic exposures accumulated throughout life contribute to the resistome, and a

complete catalog of exposures is critical to determine patterns of antibiotic prescription

most likely to account for the ARG associations to asthma found in this study. Fifth, as

with all metagenomic sequencing studies, we are limited by annotation bias in existing

databases. This is a concern for our virulence factor and antibiotic resistance profiling

especially, where we rely on the database to predict source species for ARGs and VFs.

We also recognize that the databases we used for these two analyses are biased

towards well-studied human pathogens rather than commensals or opportunistic

pathogens. However, we note that other investigators have reported similar co-

occurrence of ARGs and VFs(54, 55), and co-selection of these features is biologically

plausible.

Despite these constraints on the scope of our study, we provide evidence that

there is an increased production of LCFA and an increased richness of ARGs encoded

by the gut microbiota in people with asthma. These findings could have applications in

the care of patients with asthma. If LCFA pathways are shown to play a causal role in

airway inflammation in future studies, microbiota-directed therapeutics in the form of

dietary interventions or probiotics, could be developed to modify gut microbial

metabolism to protect against asthma. Additionally, our resistome findings add to the

growing concern over antibiotic resistance in patients with asthma by suggesting that

antibiotic administration may also contribute to gut carriage of virulence factors that can

alter airway inflammation. Ultimately, our study shows that the gut microbiota of school-

aged and older subjects with moderate-to-severe asthma harbor important functional

alterations that could serve as a foundation for future studies investigating how gut

microbial functions affect pulmonary diseases.

Author Contributions

N.G.W. and A.L.K. conceptualized the work. L.B.B. and A.L.K. planned the clinical

study. N.G.W., A.H-L., D.J.S., and A.L.K. analyzed the data and drafted the manuscript.

All authors interpreted the data and contributed to revising the manuscript.

Acknowledgements

We would like to thank our clinical study coordinators Tarisa Mantia, Caitlin

O'Shaughnessy, and Shannon Rook; the physicians of Washington University Pediatric

and Adolescent Ambulatory Research Consortium, especially Dr. Jane Garbutt; the

Volunteer for Health registry; and the MARS participants and their families. We would

also like to acknowledge the Center for Genome Sciences Sequencing Core and

Genome Technology Access Center at the McDonnell Genome Institute for sequencing

services. The authors also thank Dr. Leyao Wang and members of the Kau lab for

critical reading of the manuscript. A.L.K. is funded by the AAAAI Foundation Faculty

Development Award and the NIH K08 AI113184. A.H-L received funding through the

NIH T32 GM007200 and F30 DK127584.

Declaration of Interests

The authors have no competing interest to declare.

Methods

MARS Study Population

The Microbiome and Asthma Research Study (MARS) consisted of 104 subjects

from the St Louis, MO USA area that are either healthy or had physician-diagnosed

moderate-to-severe asthma. This study included an adult cohort (ages 18-40 years) and

pediatric cohort (ages 6-10 years). As described in previous manuscripts(23, 24), 9

patients were disqualified or did not donate stool samples. The remaining 95 patients

donated stool samples either at home or at the recruitment visit and were evaluated with

a clinical questionnaire to gather relevant metadata. Stool samples were kept at -20°C

and delivered within 24 hours to the study site, Kau Lab at Washington University

School of Medicine, where they were stored at -80°C for no more than three years until

processing for DNA isolation. This study was approved by the Washington University

Institutional Review Board (IRB# 201412035). Written informed consent documents

were obtained from all MARS subjects or their legal guardians. All recruitment, follow

up, and sample acquisition occurred between November 2015 and December 2017.

Fecal DNA Isolation

Frozen human stool samples were pulverized in liquid nitrogen using a pestle

and mortar. We then homogenized the stool in a mixture of phenol, chloroform, and

isoamyl alcohol with a bead beater using sterilized zirconium and steel beads as

previously described(63) to extract crude DNA. We purified the fecal DNA with a 96-well

QIAGEN PCR Clean up kit and quantitated by measuring the absorbance at 260/280

nm. Sample DNA concentrations were normalized to 0.5 ng/mL. Neither depletion of

human DNA sequence nor enrichment of microbial or viral DNA was performed. No

experimental quantification like a spike-in were used.

Whole Metagenomic Sequencing of Fecal Communities

To generate fecal metagenomic sequencing data, we adapter-ligated libraries by

tagmentation using an adaptation of the Nextera Library Prep kit (Illumina, cat. No. FC-

121-1030/1031)(64). Individual libraries were then purified with AMPure XP SPRI

beads, quantitated using Quant-iT (Invitrogen, cat. Q33130), and then combined in an

equimolar ratio. We confirmed that each library was adequately represented in the

combined library by preliminary sequencing on a MiSeq instrument at the Washington

University in St. Louis Center for Genome Sciences to assess the evenness of the

library. Once the quality of the library was assured, we sequenced the combined library

on a NovaSeq 6000 S4 with 2x150 bp chemistry to achieve an average of 3.4 Giga-

base-pairs (Gb) per sample. NovaSeq services and data demultiplexing were performed

by the Genome Technology Access Center at the McDonnell Genome Institute (St

Louis, MO). All samples were tagmented simultaneously and sequenced on the same

run to avoid batch effects.

Processing of sequencing data

Metagenomic raw demultiplexed reads were then processed to (1) remove

spurious human sequences (human reference database was hg37dec_v0.1.1), (2)

remove low quality sequences, and (3) trim remaining adapter content using Kneaddata

v. 0.10.0 (huttenhower.sph.harvard.edu/kneaddata) bypassing the tandem repeat finder

step ("- -bypass-trf"). FastQC (fastqc v0.11.7) and MultiQC (multiqc v1.2) with default

settings were used to create quality reports and visualize processing steps. See Figure

S1A and Table S1 for number of reads dropped per processing step. After trimming and

filtering, no samples had adaptor content, overrepresented sequences, or an average

sequence quality score below Phred 24. Estimated metagenome coverage was

calculated with Nonpareil(25, 65) (version 3.4.1) via the online querying tool at

http://enve-omics.ce.gatech.edu/nonpareil/submit.

Read-based metagenome profiling

To obtain functional information about the metagenomic contents of fecal

samples, we processed samples using HUMAnN(26) v3.0.0 on filtered reads with

default parameters. The marker gene database used by HUMAnN to identify taxonomic

identities was ChocoPhlAn v201901b and the protein database used by HUMAnN to

identify functions was the UniRef90 full database v201901b. Alpha diversity analysis of

Uniref90 genes and two-sample tests of KEGG orthologs were performed on respective

genes that were present (>0 copies per million) in at least 16 out of 95 samples, which

was the lowest prevalence cutoff that would allow for Bonferroni corrected Wilcoxon p-

values below 0.0001. HUMAnN was used to determine the abundance of metagenomic

pathways by mapping UniRef90 genes to the MetaCyc database. We performed

differential abundance analysis using the Wilcoxon 2-sample tests on pathways that had

a minimum of 10% prevalence.

To identify antibiotic resistance genes present in the fecal metagenomes of

MARS stools, we used ShortBRED-identify(34) (v0.9.4) with the Comprehensive

Antibiotic Resistance Database(35) (downloaded 2021-07-05 16:10:04.04555) and

Virulence Factor Database(39) (downloaded Fri Jul 16 10:06:01 2021). ShortBRED-

Quantify was run on the filtered reads with default parameters. ARGs or VFs that had

an abundance greater than zero in less than 7 out of 95 samples were excluded from

downstream analyses. This prevalence cutoff was determined using the binomial

distribution to maintain a 95% confidence that enrichment was not due to random

chance (using stats::binom in R). In the analyses that compared virulence factor profiles

to antibiotic resistance gene profiles, any gene with the same name was excluded from

the list of antibiotic resistance and considered a virulence factor only, to prevent

spurious results due to co-correlations. Only one gene matched this criterion: *ugd*

(UDP-glucose 6-dehydrogenase).

Microbial composition was determined with MetaPhlAn 3.0(26) which is included

in the HUMAnN pipeline described(26). MaasLin(27) (Maaslin2_1.5.1) was used in R to

find taxa of any taxonomic level that correlated with asthma by setting asthma as a fixed

effect and setting age group and race as random effects.

For PERMANOVA analyses, BMI class refers to two stratifications: Non-obese

(underweight, healthy, or overweight) and obese determined for adults by BMI cutoffs

and for pediatric patients by BMI-for-age percentile as defined by the Centers for

Disease Control and Prevention (see

cdc.gov/healthyweight/assessing/bmi/childrens_bmi/about_childrens_bmi.html). Race

was reported by the subject and split into the two categories of Caucasian and non-

Caucasian.

Metagenome Assemblies

Filtered reads were assembled into contigs using spades(37) (v3.14.0) with the

"meta" flag and k-mers lengths as follows: -k 21,33,55,77. The resulting scaffolds

achieved an average N50 of 3525 +/- 178 bp, an average L50 of 7192 +/- 372 and an

average total length of 136.8 +/- 4.5 Mbp as measured by QUAST (v 4.5) (66, 67) (see

Table S1). Determination of *ermF* location was performed by aligning the 801-bp coding

sequence of *ermF* from CARD(35) to all scaffolds. Scaffolds containing BLAST hits with

98% identity or higher to the full-length CARD *ermF* sequence were further annotated

by Prokka (v1.14.5) to find open reading frames and annotate them. Manual BLAST

was used to annotate "hypothetical protein" open reading frames for the contexts of

ermF hits.

Statistics and Reproducibility

R version 3.6.3 was used for all analyses downstream of HUMAnN and

ShortBRED, and for data visualization. Wilcoxon tests with false discovery rate multiple

testing correction or Type II ANOVAs were used to determine statistically significant

differences with the car::Anova package in R. PERMANOVAs were performed in R

using the vegan::adonis package with default settings and 100,000 iterations. The

following symbols were used to designate significance: * $p < 0.05$, ** $p < 0.01$, *** $p <$

0.001 and the following for q values (FDR-adjusted p-values): * $q < 0.2$, ** $q < 0.05$.

Data Availability

All sequencing data available at European Nucleotide Archive

(https://www.ebi.ac.uk/ena/browser/home) under project accession number

PRJEB56741. Demographic data can be found in this manuscript (Table S1) and

previous articles about the MARS study(23, 24). A STORMS (Strengthening The

Organizing and Reporting of Microbiome Studies) checklist(68) is available at doi:

10.5281/zenodo.7492635.

References

1.   Asthma-related emergency department visits 2010–2018 | CDC.

https://www.cdc.gov/asthma/asthma_stats/asthma-ed-visits_2010-2018.html.

Retrieved 28 December 2022.

2.   Arrieta M-C, Stiemsma LT, Dimitriu PA, Thorson L, Russell S, Yurist-Doutsch S,

Kuzeljevic B, Gold MJ, Britton HM, Lefebvre DL, Subbarao P, Mandhane P,

Becker A, Mcnagny KM, Sears MR, Kollmann T, Mohn WW, Turvey SE, Finlay

BB, Brett Finlay B. 2015. Early infancy microbial and metabolic alterations affect

risk of childhood asthma. Sci Transl Med 7:307–152.

3.   Hufnagl K, Pali-Schöll I, Roth-Walter F, Jensen-Jarolim E. 2020. Dysbiosis of the

gut and lung microbiome has a role in asthma. Semin Immunopathol 42:75–93.

4.   Fujimura KE, Sitarik AR, Havstad S, Lin DL, Levan S, Fadrosh D, Panzer AR,

Lamere B, Rackaityte E, Lukacs NW, Wegienka G, Boushey HA, Ownby DR,

Zoratti EM, Levin AM, Johnson CC, Lynch S V. 2016. Neonatal gut microbiota

associates with childhood multisensitized atopy and T cell differentiation. Nat Med

22:1187–1191.

5.   Levan SR, Stamnes KA, Lin DL, Panzer AR, Fukui E, McCauley KK, Fujimura KE,

McKean M, Ownby DR, Zoratti EM, Boushey HA, Cabana MD, Johnson CC,

Lynch S V. 2019. Elevated faecal 12,13-diHOME concentration in neonates at

high risk for asthma is produced by gut bacteria and impedes immune tolerance.

Nat Microbiol 4:1851–1861.

6.   Roduit C, Frei R, Ferstl R, Loeliger S, Westermann P, Rhyner C, Schiavi E, Barcik

W, Rodriguez-Perez N, Wawrzyniak M, Chassard C, Lacroix C, Schmausser-

Hechfellner E, Depner M, von Mutius E, Braun-Fahrländer C, Karvonen A,

Kirjavainen P, Pekkanen J, Dalphin J-C, Riedler J, Akdis C, Lauener R, O'Mahony

L. 2018. High levels of Butyrate and Propionate in early life are associated with

protection against atopy. Allergy 1–11.

7.   Cait A, Hughes MR, Antignano F, Cait J, Dimitriu PA, Maas KR, Reynolds LA,

Hacker L, Mohr J, Finlay BB, Zaph C, McNagny KM, Mohn WW. 2018.

Microbiome-driven allergic lung inflammation is ameliorated by short-chain fatty

acids. Mucosal Immunol 11:785–795.

8.   Trompette A, Gollwitzer ES, Yadava K, Sichelstiel AK, Sprenger N, Ngom-Bru C,

Blanchard C, Junt T, Nicod LP, Harris NL, Marsland BJ. 2014. Gut microbiota

metabolism of dietary fiber influences allergic airway disease and hematopoiesis.

Nat Med 20:159–166.

9.   Zaiss MM, Rapin A, Lebon L, Dubey LK, Mosconi I, Sarter K, Piersigilli A, Menin

L, Walker AW, Rougemont J, Paerewijck O, Geldhof P, McCoy KD, Macpherson

AJ, Croese J, Giacomin PR, Loukas A, Junt T, Marsland BJ, Harris NL. 2015. The

Intestinal Microbiota Contributes to the Ability of Helminths to Modulate Allergic

Inflammation. Immunity 43:998–1010.

10. Li X, Stokholm J, Brejnrod A, Vestergaard GA, Russel J, Trivedi U, Thorsen J,

Gupta S, Hjelmsø MH, Shah SA, Rasmussen MA, Bisgaard H, Sørensen SJ.

2021. The infant gut resistome associates with E. coli, environmental exposures,

gut microbiome maturity, and asthma-associated bacterial composition. Cell Host

Microbe 1–13.

11. Ramirez J, Guarner F, Bustos Fernandez L, Maruy A, Sdepanian VL, Cohen H.

712  2020. Antibiotics as Major Disruptors of Gut Microbiota. Front Cell Infect Microbiol

713  10:1–10.

714 12. Kozyrskyj AL, Ernst P, Becker AB. 2007. Increased risk of childhood asthma from

715  antibiotic use in early life. Chest 131:1753–1759.

716 13. McKeever TM, Lewis SA, Smith C, Collins J, Heatlie H, Frischer M, Hubbard R.

717  2002. Early exposure to infections and antibiotics and the incidence of allergic

718  disease: A birth cohort study with the West Midlands General Practice Research

719  Database. J Allergy Clin Immunol 109:43–50.

720 14. Hoskin-Parr L, Teyhan A, Blocker A, Henderson AJW. 2013. Antibiotic exposure

721  in the first two years of life and development of asthma and other allergic

722  diseases by 7.5 yr: A dose-dependent relationship. Pediatr Allergy Immunol

723  24:762–771.

724 15. Russell SL, Gold MJ, Willing BP, Thorson L, McNagny KM, Finlay BB. 2013.

725  Perinatal antibiotic treatment affects murine microbiota, immune responses and

726  allergic asthma. Gut Microbes 4:158–164.

727 16. Yang X, Feng H, Zhan X, Zhang C, Cui R, Zhong L, Ying S, Chen Z. 2019. Early-

728  life vancomycin treatment promotes airway inflammation and impairs microbiome

729  homeostasis. Aging (Albany NY) 11:2071–2081.

730 17. Borbet TC, Pawline MB, Zhang X, Wipperman MF, Reuter S, Maher T, Li J, Iizumi

731  T, Gao Z, Daniele M, Taube C, Koralov SB, Müller A, Blaser MJ. 2022. Influence

732  of the early-life gut microbiota on the immune responses to an inhaled allergen.

733  Mucosal Immunol 15:1000–1011.

734 18. Wenzel SE. 2012. Asthma phenotypes: The evolution from clinical to molecular

approaches. Nat Med 18:716–725.

19. Michalovich D, Rodriguez-Perez N, Smolinska S, Pirozynski M, Mayhew D, Uddin

S, Van Horn S, Sokolowska M, Altunbulakli C, Eljaszewicz A, Pugin B, Barcik W,

Kurnik-Lucka M, Saunders KA, Simpson KD, Schmid-Grendelmeier P, Ferstl R,

Frei R, Sievi N, Kohler M, Gajdanowicz P, Graversen KB, Lindholm Bøgh K, Jutel

M, Brown JR, Akdis CA, Hessel EM, O'Mahony L. 2019. Obesity and disease

severity magnify disturbed microbiome-immune interactions in asthma patients.

Nat Commun 10.

20. Wang Q, Li F, Liang B, Liang Y, Chen S, Mo X, Ju Y, Zhao H, Jia H, Spector TD,

Xie H, Guo R. 2018. A metagenome-wide association study of gut microbiota in

asthma in UK adults. BMC Microbiol 18:1–7.

21. Chiu CY, Chan YL, Tsai MH, Wang CJ, Chiang MH, Chiu CC. 2019. Gut microbial

dysbiosis is associated with allergen-specific IgE responses in young children with

airway allergies. World Allergy Organ J 12:100021.

22. Liu A, Ma T, Xu N, Jin H, Zhao F, Kwok L-Y, Zhang H, Zhang S, Sun Z. 2021.

Adjunctive Probiotics Alleviates Asthmatic Symptoms via Modulating the Gut

Microbiome and Serum Metabolome. Microbiol Spectr 9:1–17.

23. Wilson NG, Hernandez-Leyva A, Rosen AL, Jaeger N, McDonough RT, Santiago-

Borges J, Lint MA, Rosen TR, Tomera CP, Bacharier LB, Swamidass SJ, Kau AL.

2022. The asthma gut microbiota influences lung inflammation in gnotobiotic

mice. bioRxiv 2022.08.09.502549.

24. Jaeger N, McDonough RT, Rosen AL, Hernandez-Leyva A, Wilson NG, Lint MA,

Russler-Germain EV, Chai JN, Bacharier LB, Hsieh C-S, Kau AL. 2020. Airway

Microbiota-Host Interactions Regulate Secretory Leukocyte Protease Inhibitor

Levels and Influence Allergic Airway Inflammation. Cell Rep 33.

25. Rodriguez-R LM, Konstantinidis KT. 2014. Nonpareil: a redundancy-based

approach to assess the level of coverage in metagenomic datasets.

Bioinformatics 30:629–635.

26. Beghini F, McIver LJ, Blanco-Míguez A, Dubois L, Asnicar F, Maharjan S, Mailyan

A, Manghi P, Scholz M, Thomas AM, Valles-Colomer M, Weingart G, Zhang Y,

Zolfo M, Huttenhower C, Franzosa EA, Segata N. 2021. Integrating taxonomic,

functional, and strain-level profiling of diverse microbial communities with

biobakery 3. Elife 10:1–42.

27. Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, Tickle TL,

Weingart G, Ren B, Schwager EH, Chatterjee S, Thompson KN, Wilkinson JE,

Subramanian A, Lu Y, Waldron L, Paulson JN, Franzosa EA, Bravo HC,

Huttenhower C. 2021. Multivariable association discovery in population-scale

meta-omics studies. PLoS Comput Biol 17.

28. Brayner B, Kaur G, Keske MA, Perez-Cornago A, Piernas C, Livingstone KM.

2021. Dietary Patterns Characterized by Fat Type in Association with Obesity and

Type 2 Diabetes: A Longitudinal Study of UK Biobank Participants. J Nutr

151:3570–3578.

29. Wendell SG, Baffi C, Holguin F. 2014. Fatty acids, inflammation, and asthma. J

Allergy Clin Immunol 133:1255–1264.

30. Mizuta K, Matoba A, Shibata S, Masaki E, Emala CW. 2019. Obesity-induced

asthma: Role of free fatty acid receptors. Jpn Dent Sci Rev 55:103–107.

31.   Scott HA, Gibson PG, Garg ML, Wood LG. 2011. Airway inflammation is

augmented by obesity and fatty acids in asthma. Eur Respir J 38:594–602.

32.   Schatz M, Sorkness CA, Li JT, Marcus P, Murray JJ, Nathan RA, Kosinski M,

Pendergraft TB, Jhingran P. 2006. Asthma Control Test: Reliability, validity, and

responsiveness in patients not previously followed by asthma specialists. J

Allergy Clin Immunol 117:549–556.

33.   Snyder BM, Patterson MF, Gebretsadik T, Cacho F, Ding T, Turi KN, Abreo A,

Wu P, Hartert T V. 2021. Association between asthma status and prenatal

antibiotic prescription fills among women in a Medicaid population. J Asthma

59:2100–2107.

34.   Kaminski J, Gibson MK, Franzosa EA, Segata N, Dantas G, Huttenhower C.

2012. High-Specificity Targeted Funcitonal Profiling in Microbial Communities with

ShortBRED. Nature 486:207–214.

35.   Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, Edalatmand A,

Huynh W, Nguyen A-L V, Cheng AA, Liu S, Min SY, Miroshnichenko A, Tran H-K,

Werfalli RE, Nasir JA, Oloni M, Speicher DJ, Florescu A, Singh B, Faltyn M,

Hernandez-Koutoucheva A, Sharma AN, Bordeleau E, Pawlowski AC, Zubyk HL,

Dooley D, Griffiths E, Maguire F, Winsor GL, Beiko RG, Brinkman FSL, Hsiao

WWL, Domselaar G V, McArthur AG. 2019. CARD 2020: antibiotic resistome

surveillance with the comprehensive antibiotic resistance database. Nucleic Acids

Res 48:517–525.

36.   Mardia K, Kent J, Bibby J. 1979. Mardia, Kent, Bibby - 1979 - Multivariate

Analysis.pdf 521.

37.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, Pyshkin A V., Sirotkin A V., Vyahhi N, Tesler

G, Alekseyev MA, Pevzner PA. 2012. SPAdes: A new genome assembly

algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–

477.

38.   Seemann T. 2014. Prokka: Rapid prokaryotic genome annotation. Bioinformatics

30:2068–2069.

39.   L C, J Y, J Y, Z Y, L S, Y S, Q J. 2005. VFDB: a reference database for bacterial

virulence factors. Nucleic Acids Res 33.

40.   Nagel G, Linseisen J. 2005. Dietary intake of fatty acids, antioxidants and

selected food groups and asthma in adults'. Eur J Clin Nutr 59:8–15.

41.   Mizuta K, Zhang Y, Mizuta F, Hoshijima H, Shiga T, Masaki E, Emala CW. 2015.

Novel identification of the free fatty acid receptor FFAR1 that promotes

contraction in airway smooth muscle. Am J Physiol - Lung Cell Mol Physiol

309:L970–L982.

42.   Felton E, Burrell A, Chaney H, Sami I, Koumbourlis AC, Freishtat RJ, Crandall

KA, Hahn A. 2021. Inflammation in children with cystic fibrosis: contribution of

bacterial production of long-chain fatty acids. Pediatr Res 90:99–108.

43.   Niot I, Poirier H, Tran TTT, Besnard P. 2009. Intestinal absorption of long-chain

fatty acids: Evidence and uncertainties. Prog Lipid Res 48:101–115.

44.   Vital M, Howe AC, Tiedje JM. 2014. Revealing the bacterial butyrate synthesis

pathways by analyzing (meta)genomic data. MBio 5:1–11.

45.   Yassour M, Vatanen T, Siljander H, Hämäläinen AM, Härkönen T, Ryhänen SJ,

Franzosa EA, Vlamakis H, Huttenhower C, Gevers D, Lander ES, Knip M, Xavier

RJ. 2016. Natural history of the infant gut microbiome and impact of antibiotic

treatment on bacterial strain diversity and stability. Sci Transl Med 8.

46. Schwartz DJ, Langdon AE, Dantas G. 2020. Understanding the impact of

antibiotic perturbation on the human microbiome. Genome Med 12:1–12.

47. Palleja A, Mikkelsen KH, Forslund SK, Kashani A, Allin KH, Nielsen T, Hansen

TH, Liang S, Feng Q, Zhang C, Pyl PT, Coelho LP, Yang H, Wang J, Typas A,

Nielsen MF, Nielsen HB, Bork P, Wang J, Vilsbøll T, Hansen T, Knop FK,

Arumugam M, Pedersen O. 2018. Recovery of gut microbiota of healthy adults

following antibiotic exposure. Nat Microbiol 3:1255–1265.

48. Juhn YJ. 2014. Risks for infection in patients with asthma (or other atopic

conditions): Is asthma more than a chronic airway disease? J Allergy Clin

Immunol 134:247-257.e3.

49. Malhotra-Kumar S, Lammens C, Coenen S, Van Herck K, Goossens H. 2007.

Effect of azithromycin and clarithromycin therapy on pharyngeal carriage of

macrolide-resistant streptococci in healthy volunteers: a randomised, double-

blind, placebo-controlled study. Lancet 369:482–490.

50. Doan T, Hinterwirth A, Worden L, Arzika AM, Maliki R, Abdou A, Kane S, Zhong

L, Cummings SL, Sakar S, Chen C, Cook C, Lebas E, Chow ED, Nachamkin I,

Porco TC, Keenan JD, Lietman TM. 2019. Gut microbiome alteration in MORDOR

I: a community-randomized trial of mass azithromycin distribution. Nat Med

25:1370–1376.

51. Doan T, Arzika AM, Hinterwirth A, Maliki R, Zhong L, Cummings S, Sarkar S,

Chen C, Porco TC, Keenan JD, Lietman TM. 2019. Macrolide Resistance in

MORDOR I — A Cluster-Randomized Trial in Niger. N Engl J Med 380:2271–

2273.

52. Gibson PG, Yang IA, Upham JW, Reynolds PN, Hodge S, James AL, Jenkins C,

Peters MJ, Marks GB, Baraket M, Powell H, Taylor SL, Leong LEX, Rogers GB,

Simpson JL. 2017. Effect of azithromycin on asthma exacerbations and quality of

life in adults with persistent uncontrolled asthma (AMAZES): a randomised,

double-blind, placebo-controlled trial. Lancet 390:659–668.

53. Brusselle GG, VanderStichele C, Jordens P, Deman R, Slabbynck H, Ringoet V,

Verleden G, Demedts IK, Verhamme K, Delporte A, Demeyere B, Claeys G,

Boelens J, Padalko E, Verschakelen J, Maele G Van, Deschepper E, Joos GFP.

2013. Azithromycin for prevention of exacerbations in severe asthma (AZISAST):

a multicentre randomised double-blind placebo-controlled trial. Thorax 68:322–

329.

54. Escudeiro P, Pothier J, Dionisio F, Nogueira T. 2019. Antibiotic Resistance Gene

Diversity and Virulence Gene. mSphere 4:1–13.

55. Pan Y, Zeng J, Li L, Yang J, Tang Z, Xiong W, Li Y, Chen S, Zeng Z. 2020.

Coexistence of Antibiotic Resistance Genes and Virulence Factors Deciphered by

Large-Scale Complete Genome Analysis. mSystems 5.

56. Strati F, Pujolassos M, Burrello C, Giuffrè MR, Lattanzi G, Caprioli F, Troisi J,

Facciotti F. 2021. Antibiotic-associated dysbiosis affects the ability of the gut

microbiota to control intestinal inflammation upon fecal microbiota transplantation

in experimental colitis models. Microbiome 9:1–15.

57. Casterline BW, Hecht AL, Choi VM, Bubeck Wardenburg J. 2017. The

Bacteroides fragilis pathogenicity island links virulence and strain competition. Gut

Microbes 0976:1–10.

58. Walter J, Armet AM, Finlay BB, Shanahan F. 2020. Establishing or Exaggerating

Causality for the Gut Microbiome: Lessons from Human Microbiota-Associated

Rodents. Cell 180:221–232.

59. Barcik W, Pugin B, Westermann P, Perez NR, Ferstl R, Wawrzyniak M,

Smolinska S, Jutel M, Hessel EM, Michalovich D, Akdis CA, Frei R, O'Mahony L.

2016. Histamine-secreting microbes are increased in the gut of adult asthma

patients. J Allergy Clin Immunol 138:1491-1494.e7.

60. Barcik W, Pugin B, Brescó MS, Westermann P, Rinaldi A, Groeger D, Van Elst D,

Sokolowska M, Krawczyk K, Frei R, Ferstl R, Wawrzyniak M, Altunbulakli C, Akdis

CA, O'Mahony L. 2019. Bacterial secretion of histamine within the gut influences

immune responses within the lung. Allergy Eur J Allergy Clin Immunol 74:899–

909.

61. Licari A, Fuchs D, Marseglia G, Ciprandi G. 2019. Tryptophan metabolic pathway

and neopterin in asthmatic children in clinical practice. Ital J Pediatr 45:1–4.

62. Van der Leek AP, Yanishevsky Y, Kozyrskyj AL. 2017. The kynurenine pathway

as a novel link between allergy and the gut microbiome. Front Immunol 8:1374.

63. Kau AL, Planer JD, Liu J, Rao SS, Yatsunenko T, Trehan I, Manary MJ, Liu T-CC,

Stappenbeck TSTS, Maleta KM, Ashorn P, Dewey KG, Houpt ER, Hsieh C-SS,

Gordon JI. 2015. Functional characterization of IgA-targeted bacterial taxa from

undernourished Malawian children that produce diet-dependent enteropathy. Sci

Transl Med 7:1–15.

64.   Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony RK.

2015. Inexpensive multiplexed library preparation for megabase-sized genomes.

PLoS One 10:1–15.

65.   Rodriguez-R LM, Gunturu S, Tiedje JM, Cole JR, Konstantinidis KT. 2018.

Nonpareil 3: Fast Estimation of Metagenomic Coverage and Sequence Diversity.

mSystems 3.

66.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: Quality assessment

tool for genome assemblies. Bioinformatics 29:1072–1075.

67.   Mikheenko A, Valin G, Prjibelski A, Saveliev V, Gurevich A. 2016. Icarus:

visualizer for de novo assembly evaluation. Bioinformatics 32:3321–3323.

68.   Mirzayi C, Renson A, Furlanello C, Sansone SA, Zohra F, Elsafoury S, Geistlinger

L, Kasselman LJ, Eckenrode K, van de Wijgert J, Loughman A, Marques FZ,

MacIntyre DA, Arumugam M, Azhar R, Beghini F, Bergstrom K, Bhatt A, Bisanz

JE, Braun J, Bravo HC, Buck GA, Bushman F, Casero D, Clarke G, Collado MC,

Cotter PD, Cryan JF, Demmer RT, Devkota S, Elinav E, Escobar JS, Fettweis J,

Finn RD, Fodor AA, Forslund S, Franke A, Furlanello C, Gilbert J, Grice E, Haibe-

Kains B, Handley S, Herd P, Holmes S, Jacobs JP, Karstens L, Knight R, Knights

D, Koren O, Kwon DS, Langille M, Lindsay B, McGovern D, McHardy AC,

McWeeney S, Mueller NT, Nezi L, Olm M, Palm N, Pasolli E, Raes J, Redinbo

MR, Rühlemann M, Balfour Sartor R, Schloss PD, Schriml L, Segal E, Shardell M,

Sharpton T, Smirnova E, Sokol H, Sonnenburg JL, Srinivasan S, Thingholm LB,

Turnbaugh PJ, Upadhyay V, Walls RL, Wilmes P, Yamada T, Zeller G, Zhang M,

Zhao N, Zhao L, Bao W, Culhane A, Devanarayan V, Dopazo J, Fan X, Fischer

M, Jones W, Kusko R, Mason CE, Mercer TR, Sansone SA, Scherer A, Shi L,

Thakkar S, Tong W, Wolfinger R, Hunter C, Segata N, Huttenhower C, Dowd JB,

Jones HE, Waldron L. 2021. Reporting guidelines for human microbiome

research: the STORMS checklist. Nat Med 27:1885–1892.

Figure 1: Gut metagenomes from individuals with asthma show increased genes
encoding fatty acid metabolism. A) Stacked violin plots of Uniref90 cluster richness
(unique Uniref90 cluster with CPM>0) grouped by either healthy and asthma cohort
(blue green colors in background) or age (brown colors in foreground. **B)** Non-metric
multidimensional scaling plot of Bray-Curtis Dissimilarity distance between Uniref90
(copies per million) profiles. Axis 1 and 2 of five total are shown of an NMDS with stress
value 0.09. **C)** Sequential PERMANOVA of Bray-Curtis dissimilarities between Uniref90
profiles. Input order of terms to the test is identical to the order of the barplot from top to
bottom. **D)** Relative abundance of MetaCyc pathways that were differentially abundant

given a Wilcoxon q value below 0.2 (p-value after FDR correction). **E)** Stacked bar plot

of differentially abundant fatty acid metabolism pathways mapped to respective taxa by

MetaPhlAn3.0/HUMAnN3.0, averaged within asthma or healthy cohorts. **F)** Heatmap of

MetaCyc pathway abundance ratios between groups in important clinical demographics:

Asthma vs. Healthy, Adult vs. Pediatric, Obese vs Non-Obese, and Well-Controlled

Asthmatics vs. Poorly-Controlled Asthmatics. Asterisk denotes a significant differential

abundance (*q<0.2) according to Wilcoxon tests controlled for multiple comparison

testing within each demographic category. **G)** Differentially abundant MetaCyc

pathways plotted as four cohorts: asthma by age with respective Two-Way ANOVAs.

Only statistically significant p values shown.

946
Figure 2: Gut metagenomes from individuals with asthma harbor an increased
richness of antibiotic resistance genes. A) Table describing short-term antibiotic
usage in the MARS cohorts. **B)** Overlapping violin plots of ARG richness and load by
grouped by either healthy and asthma cohort (blue green colors in background) or age
(brown colors in foreground. **C)** Stacked bar plots of average ARG richness painted by
antimicrobial family (AMR), drug class to which the ARG confers resistance, and ARG
resistance mechanism.

Figure 3: The gut antibiotic resistome is altered in asthma patients. A) Non-metric Multidimensional Scaling (NMDS) plot of antibiotic resistome with units in Bray-Curtis dissimilarity of total-sum scaled RPKM, labeled by asthma and age cohorts. Showing two axes out of five with stress value=0.1. **B)** Effect of demographic categories on antibiotic resistome data in A (sequential PERMANOVA). **C)** Procrustes and PROTEST analysis between MetaPhlAn species-level Bray-Curtis dissimilarity distances and CARD ShortBRED Bray-Curtis dissimilarity distances. Arrows connect the two data points belonging to identical samples.

964
Figure 4: **Resistance gene *ermF* is differentially abundant in diverse genomic**
contexts of gut resistomes belonging to individuals with asthma. A) Boxplots of
antibiotic resistance gene (ARG) abundance by cohort on log-scale. Showing only
ARGs present in at least 7 out of 95 samples and have q-values less than 0.2. A
pseudocount of 0.0015 RPKM (designated as the limit of detection "LOD") was used for
the negative binomial tests. Bolded genes are enriched in the asthma cohort while non-
bolded are enriched in the healthy cohort. **B)** Summary of *ermF* contexts on contigs
from metagenomic assemblies that had at least one detectable open reading frame
flanking the *ermF* within 10 kilobases. **C)** Three representative *ermF* context maps
generated in GeneSpy. **D)** Count tables of *ermF*+ (top) and *ermF*- (bottom) MARS fecal
samples split by *bft* presence and asthma status. Both tables showing two-sided p-
value. **E)** Count tables of metagenomes (top from MARS and bottom from Human
Microbiome Project) split by the presence of *B. fragilis* toxin (*bft*) and *ermF*. Top: one-
sided p-value shown; bottom: two-sided p-value shown.

Figure 5: **Asthma patients have unique sets of virulence factor and antibiotic resistance gene associations. A)** Partial correlations split by asthma status between virulence factor richness and ARG richness after accounting for species richness. **B)** Procrustes and PROTEST analysis between Bray-Curtis dissimilarity distances of virulence factors and CARD resistomes. Arrows connect the two data points belonging to identical samples. **C)** Heatmap of statistically significant (cooccur R package p<0.05) co-occurrence relationships between all VFs and ARGs. Colors indicate direction of co-occurrence and in which cohort(s) the respective effect was detected. Grey squares mark pairs with no statistically significant co-occurrence. White squares were pairs filtered out due to a lack of observed co-occurrence.

 Supplemental Figure and Data Table Captions

Figures:
Figure S1: MARS whole metagenomic shotgun sequencing captures essential
functions and taxonomic shifts of the asthma gut microbiota. A) Summary of select
sequencing statistics from NovaSeq shotgun metagenomic sequencing and subsequent
filtering steps. **B)** Boxplot of redundancy-based estimated metagenome coverage (%)
as calculated by running the forward reads through the Nonpareil tool. Split into asthma
and age group and Two-way Type II ANOVA results shown. **C)** Bar plot of MetaCyc
pathway copies per million (CPM) in all MARS samples annotated by HUMANnN
pipeline, with horizontal length representing mean and bars the standard error. For all
panels: N= 20 healthy children, 39 healthy adults, 19 asthmatic children, 17 asthmatic
adults. **D)** Relative abundance stacked barplots of top abundant bacterial genera split
by age group and asthma cohort. **E)** Simpson alpha diversity boxplots split by asthma
and age group cohorts (2-Way Type II ANOVA). **F)** NMDS of Bray-Curtis Dissimilarity of
species-level relative abundance grouped by age and asthma. **G)** Sequential
PERMANOVA to test effect of demographics on beta diversity. Terms were input into
the test as ordered from top to bottom of barplot. Dotted vertical line represents a p
value of 0.05. Color scale is mapped to the R^2 value. **H)** Arcsine transformed relative
abundance boxplots of differentially abundant species as determined by Maaslin2 with
age group and race modeled as random effects. For all panels: N= 20 healthy children,
39 healthy adults, 19 asthmatic children, 17 asthmatic adults.

1013

Figure S2: KEGG orthologs, KEGG pathways, and differentially abundant
MetaCyc fatty acid pathways. A) Relative abundance of KEGG orthologs previously
implicated in asthma. Copies per million (CPM) are counts normalized by gene size and
read depth, then total-sum-scaled to one million. **B-C)** Stacked bar plots of differentially
abundant pathways mapped to respective taxa including "Community" bin which
accounts for the remaining reads that mapped to the pathway but not to any single
species by MetaPhlAn3.0/HUMAnN3.0, averaged within asthma or healthy cohorts. **B)**
L-lysine biosynthesis III pathway. Only top 13 taxa shown in addition to Community
category. **C)** Seven fatty acid metabolism pathways differentially abundant in the
asthma cohort. Only top 9 taxa shown in addition to Community category. Stars
represent a q value < 0.05 of Wilcoxon tests between the Community pathway richness
and *B. vulgatus*-encoded pathway richness. For all panels: N= 59 healthy, 36 asthmatic
individuals.

1027

1028 **Figure S3: Pathway collage for differentially abundant MetaCyc pathways.** "PWY-
1029 6519: 8-amino-7-oxononanoate biosynthesis I" is completely overlapping with "BIOTIN-
1030 BIOSYNTHESIS-PWY: biotin biosynthesis I" and its steps are highlighted in blue text.
1031 Pathway collage made on MetaCyc browser tool.

1032

1033 **Figure S4: Gut virulence factor ecology shifts with age group but not asthma**
1034 **cohort. A)** Total-sum scaled RPKM Bray-Curtis Dissimilarity Non-metric
1035 Multidimensional Scaling (NMDS) plot labeled by asthma and age cohorts. Showing two
1036 axes out of 5 with stress value=0.09. **B)** Effect of demographic categories on virulence

factor profile in A (by sequential PERMANOVA, input terms ordered from top to bottom of barplot). **C)** Stacked violin plots of virulence factor alpha diversity grouped by eithter healthy and asthma cohort (blue green colors in background) or age (brown colors in foreground). Two-Way ANOVA results shown in table above plot. **D)** Procrustes plot and PROTEST analysis between virulence factor profile Bray-Curtis dissimilarity distances and Metaphlan species relative abundance Bray-Curtis dissimilarity distances. Arrows connect the two data points belonging to identical samples. For all panels: N= 20 healthy children, 39 healthy adults, 19 asthmatic children, 17 asthmatic adults.

Figure S5: Asthma-associated ARG richness and ARG-VF co-occurrence relationships are observed within *K. pneumoniae* and *E. coli*. A) Richness bar plots between antibiotic resistance genes (ARGs) and virulence factors (VFs) grouped by asthma status. **B)** Heatmap of the co-occurrence of each VF/ARG pair colored by the direction in which (positively or negatively co-occurring) and the cohort for which (asthma vs. healthy) the pair had a p-value less than 0.05 via R cooccur function. Blank squares were pairs filtered out due to a lack of observed co-occurrence. SS: secretion systems. (N=36 Healthy, 59 Asthmatic)

Tables:
Table S1: Fecal shotgun metagenomics filtering and assembly summary statistics
Table S2: Maaslin 2 Analysis of Metaphlan Community Composition
Table S3: Comparison of MetaCyc Pathway Differential Abundance Analyses
Table S4: Negative Binomial tests of antibiotic resistance gene abundance (RPKM) between healthy and asthma cohorts
Table S5: Negative Binomial tests of virulence factor abundance (RPKM) between healthy and asthma cohorts