Simultaneous proteome localization and turnover analysis reveals spatiotemporal features of protein homeostasis disruptions

[1]Jordan Currie, [1]Vyshnavi Manda, [1]Sean K. Robinson, [4]Celine Lai, [5]Vertica Agnihotri, [1]Veronica Hidalgo, [1]R. W. Ludwig, [6]Kai Zhang, [1]Jay Pavelka, [6]Zhao V. Wang, [5]June-Wha Rhee, [1,2,3]Maggie P. Y. Lam, [1,3]Edward Lau

1. Department of Medicine
2. Department of Biochemistry and Molecular Genetics
3. Consortium for Fibrosis Research and Translation
   University of Colorado School of Medicine
   Aurora, CO 80045, USA
4. Stanford Cardiovascular Institute
   Stanford University
   Stanford, CA 94305, USA
5. Department of Medicine, Division of Cardiology
   City of Hope Comprehensive Cancer Center
   Durante, CA 91010, USA
6. Department of Diabetes and Cancer Metabolism
   Beckman Research Institute, City of Hope National Medical Center
   Duarte, CA 91010, USA

Correspondence
Edward Lau, PhD: edward.lau@cuanschutz.edu

Abstract

The functions of proteins depend on their spatial and temporal distributions, which are not directly measured by static protein abundance. Under endoplasmic reticulum (ER) stress, the unfolded protein response (UPR) pathway remediates proteostasis in part by altering the turnover kinetics and spatial distribution of proteins. A global view of these spatiotemporal changes has yet to emerge and it is unknown how they affect different cellular compartments and pathways. Here we describe a mass spectrometry-based proteomics strategy and data analysis pipeline, termed Simultaneous Proteome Localization and Turnover (SPLAT), to measure concurrently the changes in protein turnover and subcellular distribution in the same experiment. Investigating two common UPR models of thapsigargin and tunicamycin challenge in human AC16 cells, we find that the changes in protein turnover kinetics during UPR varies across subcellular localizations, with overall slowdown but an acceleration in endoplasmic reticulum and Golgi proteins involved in stress response. In parallel, the spatial proteomics component of the experiment revealed an externalization of amino acid transporters and ion channels under UPR, as well as the migration of RNA-binding proteins toward an endosome co-sedimenting compartment. The SPLAT experimental design classifies heavy and light SILAC labeled proteins separately, allowing the observation of differential localization of new and old

protein pools and capturing a partition of newly synthesized EGFR and ITGAV to the ER under
stress that suggests protein trafficking disruptions. Finally, application of SPLAT toward human
induced pluripotent stem cell derived cardiomyocytes (iPSC-CM) exposed to the cancer drug
carfilzomib, identified a selective disruption of proteostasis in sarcomeric proteins as a potential
mechanism of carfilzomib-mediated cardiotoxicity. Taken together, this study provides a global
view into the spatiotemporal dynamics of human cardiac cells and demonstrates a method for
inferring the coordinations between spatial and temporal proteome regulations in stress and drug
response.

Keywords

48　Unfolded protein response, protein turnover, spatial proteomics, mass spectrometry, stress
49　response, subcellular localization, spatiotemporal dynamics

Introduction

Protein turnover is an important cellular process that maintains the quality and quantity of protein
pools in homeostasis, and involves fine regulations of the rates of synthesis and degradation of
individual proteins. A close relationship exists between turnover kinetics with the spatial
distribution of proteins. Cellular organelles including the cytosol, endoplasmic reticulum (ER),
and mitochondria are equipped with distinct quality control and proteolytic mechanisms that
maintain protein folding and regulate protein degradation in a localization dependent manner
(Lemberg and Strisovsky, 2021; Mårtensson et al., 2019; Tsai et al., 2022). Newly synthesized
proteins need to be properly folded and trafficked to their intended subcellular localization
through subcellular targeting and sorting mechanisms, whose capacity has to be coordinated to
match the rate of protein synthesis (Chartron et al., 2016; Jan et al., 2014; Lakkaraju et al., 2008).
A primary subcellular trafficking mechanism of new proteins is the ER vesicular transport and
secretory pathway through the endomembrane system, which is a rate-limiting step in the
production of membrane and extracellular proteins. A mismatch between temporal synthesis
rate and spatial localization capacity can lead to ER stress and subsequently mislocalization of
newly synthesized proteins (Hetz et al., 2020).
Disruption of protein turnover and homeostasis is broadly implicated in human diseases
including cardiomyopathies, cancer, and neurodegeneration (Hetz et al., 2020; Ren et al., 2021).
In stressed cells, the accumulation of misfolded proteins triggers the unfolded protein response
(UPR), which involves signaling pathways that suppress protein synthesis and promote protein
folding and proteolysis to relieve proteostatic stress. At the same time, UPR activation invokes
a spatial reorganization of the proteome, including but not limited to the transient translocation
of UPR pathway mediators to the nucleus during initial stress response, the retrotranslocation of
misfolded ER proteins to the cytosol for proteasomal clearance under endoplasmic-reticulum-
associated protein degradation (ERAD), and the sequestration of RNA and RNA-binding proteins
into stress granules. Despite ongoing research, how the cellular proteome reorganizes under
proteostatic stress remains incompletely understood, and the full scope of differentially localized
proteins and pathways remains to be elucidated.

We wonder how the spatial and temporal regulations of proteins change in conjunction
under UPR, such as whether proteins with differential spatial distributions are also differentially
turned over. Advances in mass spectrometry methods have allowed the turnover rate and
subcellular localization of proteins to be measured on a large scale. The turnover rate and half-
life of proteins can be measured using stable isotope labeling in cells and in intact animals
followed by mass spectrometry measurements of isotope signatures and kinetics modeling to
derive rate constants (Claydon and Beynon, 2012; Doherty et al., 2009; Hammond et al., 2022;
Schwanhäusser et al., 2011). Quantitative comparison of turnover rates provides a temporal view
into proteostatic regulations and can implicate new pathological signatures and pathways over
steady-state mRNA and protein levels (Andrews et al., 2022; Lam et al., 2014; Lau et al., 2018).
In parallel, spatial proteomics methods have allowed increasing power to discern the subcellular
localization of proteins on a large scale (Christopher et al., 2021; Geladaki et al., 2019; Kennedy
et al., 2020; Mulvey et al., 2021; Orre et al., 2019). In recent work using a differential solubility
fractionation strategy and mass spectrometry, we observed broad substantial rearrangement of
proteins across three subcellular fractions in an acute paraquat challenge model of UPR in the
mouse heart, consistent with protein translocation being an important layer of proteome
regulation under proteostatic stress (Dostal et al., 2020). Nevertheless, an integrated strategy
that can simultaneously measure protein turnover kinetics and spatial information has thus far
not been realized.

Here we extended protein turnover measurements to include subcellular localization
dynamics, by integrating dynamic SILAC labeling with differential ultracentrifugation-based
spatial proteomics profiling strategies. We describe an experimental strategy and computational
analysis pipeline to perform simultaneous proteome localization and turnover (SPLAT)
measurements in baseline and stressed cells. SPLAT builds on prior work in protein turnover
measurements and subcellular localization profiling, by combining dynamic SILAC isotope
labeling, differential ultracentrifugation, isobaric TMT labeling, and kinetic modeling. This
strategy allows for concurrent measurement of changes in the turnover dynamics and subcellular
distributions of whole cell proteomes under perturbation in a single experiment. Applying this
method to human AC16 cardiac cells under thapsigargin- and tunicamycin-induced UPR and to
human induced pluripotent stem cell derived-cardiomyocytes (iPSC-CM) under carfilzomib
induced proteasome inhibition, we delineated prominent changes in the spatial and temporal
distributions of proteins on a proteome scale. The inclusion of spatial information of light and
heavy SILAC labeled proteins moreover allowed disaggregation of the localization and trafficking
of new and old protein pools.

Results

Simultaneous acquisition of turnover and spatial information using a double labeling strategy

We reason that we can use a hyperplexing strategy to simultaneously encode temporal and
spatial protein information through isotope labels in the MS1 and MS2 levels, respectively.

Hence, we designed a workflow that combines dynamic SILAC metabolic labeling in cultured
cells, with TMT labeling of spatially separated fractions to simultaneously measure new protein
synthesis as well as subcellular localization under baseline and perturbation conditions (**Figure
1A**). To determine the rate of protein turnover during control, thapsigargin, and tunicamycin
conditions, a dynamic SILAC strategy was used to measure the rate of appearance of post-
labeling synthesized protein. Briefly, cells were pulsed with a lysine and arginine depleted media
supplemented with heavy labeled lysine and arginine concurrently with drug treatment to label
post-treatment synthesized proteins and derive fractional synthesis rates through kinetic
modeling.
Upon harvesting, the cells were fractionated to resolve subcellular compartments. We
adopted a protein correlation profiling approach. In particular, the LOPIT-DC (Localisation of
Organelle Proteins by Isotope Tagging after Differential ultraCentrifugation) method (Geladaki et
al., 2019) uses sequential ultracentrifugation to enrich different subcellular fractions from the
same samples, which facilitates ease of adoption and reproducibility. Briefly, the cells were lysed
under gentle conditions and then sequentially pelleted through ultracentrifugation steps, which
pellets subcellular fractions based on their sedimentation rate and which is a function of particle
mass, shape, and volume. The ultracentrifugation fractions were each subsequently solubilized,
and the extracted proteins were digested and further labeled with tandem mass tag (TMT)
isobaric stable isotope labels. The acquired mass spectrometry data therefore carries temporal
information in the dynamic SILAC tags and spatial information in the TMT channel intensities
(**Figure 1B**).
To process the double isotope encoded mass spectrometry data, we assembled a
custom computational pipeline comprising database search and post-processing, and
quantification for dynamic SILAC and TMT data (**Figure 1C**). The turnover kinetics information
from the dynamic SILAC data is analyzed using a mass spectrometry software tool we previously
developed, RIANA (Hammond et al., 2022), which integrates the areas-under-curve of mass
isotopomers from peptides over a specified retention time window, then performs kinetic curve-
fitting to a mono-exponential model to measure the fractional synthesis rates (FSR) of each
dynamic SILAC-labeled (K and R containing) peptide. To extract the ultracentrifugation fraction
quantification information for spatial analysis, we developed a new version of the pyTMT tool
which we previously described (Dostal et al., 2020), and used it to perform TMT label
quantification for the Comet/Percolator workflow. To account for specific challenges related to
spatial proteomics data features, we made two new modifications. First, we account for isotope
impurities in TMT tags. Because the TMT data are row normalized in the LOPIT-DC design, we
incorporated correction of isotope contamination of TMT channels based on the batch
contamination data sheet (**Supplemental Table S1**) to account for isotope impurity in fractional
abundance calculation from randomized channels across experiments (**Supplemental
Methods**). Second, we implemented an isoform-aware quantitative rollup of peptide channel
intensities into the protein level for the downstream spatial proteomics analysis. Standard protein
inference invokes parsimony rules that assign peptides to the protein within a protein group with
the highest level of evidence, but razor peptides can conflate spatial information from different
proteins with different localization. Here the TMT-quantified peptides are summed into protein

groups using a more conservative aggregate method, such that the identified peptides that are
assigned to two or more top-level UniProt protein accessions are discarded to avoid
confounding of spatial information in the TMT channels. Moreover, protein groups containing
two or more proteins belonging to the same top-level UniProt accession are removed from
consideration if one of the non-canonical isoforms contain a unique peptide, and are otherwise
rolled up to the canonical protein. Protein isoforms are only included in downstream analysis
through quantified isoform-unique peptides. Following RIANA and pyTMT processing, the
SPLAT pipeline combines the dynamic SILAC and TMT information by peptides and appends a
heavy ("_H") tag to the UniProt accession of all peptides containing dynamic SILAC
modifications for separate localization analyses. The data were then used for temporal kinetics
summaries using the MS1 encoded information and subcellular localization classification from
the MS2 encoded information (**Figure 1D**). By separately analyzing heavy and light peptides, the
subcellular spatial information of the heavy (new) and light (old) subpools of thousands of
proteins can be mapped simultaneously in normal and perturbed cells.

173
Figure 1. Overview of the SPLAT strategy. **A.** Experimental workflow. Control, thapsigargin-treated, and
tunicamycin-treated human A16 cardiomyocytes were labeled with $^{13}C_6{}^{15}N_2$ L-Lysine and $^{13}C_6{}^{15}N_4$ L-

Arginine dynamic SILAC labels. For each condition, 3 biological replicate SPLAT experiments were
performed (n=3). After 16 hours, cells were harvested and mechanically disrupted, followed by differential
ultracentrifugation steps to pellet proteins across cellular compartments. Proteins from the
ultracentrifugation fractions were digested and labeled using tandem mass tag (TMT) followed by mass
spectrometry. **B.** Dynamic SILAC labeling allowed differentiation of pre-existing (unlabeled, i.e., SILAC
light) and post-labeling (heavy lysine or arginine, i.e., +R[10.0083]) synthesized peptides at 16 hours. The
light and heavy peptides were isolated for fragmentation separately to allow the protein sedimentation
profiles containing spatial information to be discerned from TMT channel intensities. **C.** Computational
workflow. Mass spectrometry raw data were converted to mzML format to identify peptides using a
database search engine. The mass spectra and identification output were processed using RIANA (left) to
quantitate the time dependent change in SILAC labeling intensities and determine the protein half-life, and
using pyTMT (right) to extract and correct TMT channel intensities from each light or heavy peptide MS2
spectrum. The TMT data were further processed using pRoloc/Bandle to predict protein subcellular
localization via supervised learning. **D.** Temporal information and spatial information is resolved in MS1
and MS2 levels, respectively. SPLAT allows the subcellular spatial information of the heavy (new) and light
(old) subpools of thousands of proteins to be quantified simultaneously in normal and perturbed cells. HL:
Half-life.

Protein turnover kinetics regulations under unfolded protein response vary by cellular compartments

We applied SPLAT to identify protein spatiotemporal changes in human AC16 cells under UPR
induced by 1 µM thapsigargin for 16 hours. Thapsigargin at the dosage and duration used is a
common and robust model to induce ER stress and integrated stress response in cardiac and
other cell types through the inhibition of sarco/endoplasmic reticulum Ca2+-ATPase (SERCA).
Thapsigargin treatment at 16 hours robustly induced known ER stress markers (Glembotski,
2007) including BiP/HSPA5, HSP90B1, PDIA4 (limma FDR adjusted P < 0.01) (**Figure 2A**). Three
biological replicate SPLAT experiments were carried out for normal and thapsigargin-treated
AC16 cells (n=3 each). We analyzed the spatial fractionation patterns of the proteins following
ultracentrifugation and TMT labeling, and classified the subcellular localization of proteins using
a Bayesian model BANDLE as previously described (Crook et al., 2018). A spatial classification
model is trained separately for each treatment using a basket of canonical organelle markers
(**see Methods**) which showed clear separation in PC1 and PC2 in each condition (**Supplemental
Figure S1**). The ultracentrifugation profiles of each cellular compartment are highly consistent
across treatments and replicates (**Supplemental Figure S2**). To minimize the potential ratio
compression that can result from MS2-based TMT quantification, we employed extensive two-
dimensional fractionation and narrow isolation window, and verified that identified MS2 spectra
had high precursor ion purity (median purity 92–93%) (**Supplemental Figure S3**). We further
performed a direct comparison of MS2 and MS3 based quantification on an identical sample
(control replicate 2) (**Supplemental Figure S4**), which confirmed that MS2-based quantification
produced acceptable spatial resolution, consistent with previous observations (Shin et al., 2020).
In total using MS2-based TMT quantification, we mapped the subcellular profiles of 4360
protein features (i.e., 1,820 new and 2,540 old proteins) in normal AC16 cells across 3 biological
replicate experiments using a stringent two-peptide filter at 1% FDR of protein identification,

with 1946 old proteins and 1,462 new proteins assigned to one of 12 subcellular localization with
>95% confidence after removing outliers (see Methods) (**Figure 2B; Supplemental Data S1**).
The accuracy of the spatial classification is supported by the observation that 69.5% of assigned
proteins in normal AC16 cells contain matching cellular component annotation in Gene Ontology
despite the current incompleteness of annotations (**Figure 2C**) and 71.6% of proteins match
their localization annotation in thapsigargin-treated cells (**Supplemental Figure S5**). From the
associated SILAC data of the proteins with spatial information, we further quantified and
compared the turnover kinetics of 2516 proteins (**Supplemental Data S2**); hence we were able
to acquire proteome-wide spatial and temporal information in matching samples from a single
experiment.
Considering the temporal kinetics data, we observed a proteome-wide decrease in
fractional synthesis rates under thapsigargin challenge compared with normal cells (median
protein half-life 46.9 vs. 19.9 hours; Mann-Whitney test $P < 2.2\text{e}{-}16$) (**Figure 2D**). This slowdown
is consistent with the extensive shutdown in protein translation due to ribosome remodeling
under integrated stress response (Bresson et al., 2020; Pakos-Zebrucka et al., 2016), shown
here by the decreased rate of SILAC incorporation into proteins. Notwithstanding the overall
slowdown, we also observed a wide range of protein turnover rates in both the untreated and
thapsigargin treated conditions that differ by the assigned subcellular compartment
(**Supplemental Figure S6**). Changes in protein kinetics following thapsigargin also varies by
compartment, with ER and Golgi proteins having significantly less slowdown of protein kinetics
compared to protein in other compartments (Mann-Whitney test P: $9.4\text{e}{-}11$ and $9.8\text{e}{-}5$,
respectively; $<< 0.05/13$) (**Figure 2E**). On an individual protein level, out of the 2516 proteins
measured, 1542 showed significant changes in temporal kinetics (Mann-Whitney test, FDR
adjusted P value < 0.1), but the vast majority of these proteins show decreases in turnover as
expected, with only 12 proteins showing significant increased temporal kinetics. Among these
are the induced ER stress markers BiP/HSPA5, HSP90B1, and PDIA4 (**Figure 2F; Supplemental
Data S3**) but also other ER and Golgi proteins that may be involved in stress response (**Figure
2G**). SDF2L1 (stromal-cell derived factor 2 like 1) is recently described to form a complex with
the ER chaperone DNAJB11 to retain it in the ER (Hanafusa et al., 2019). In control cells, we
found that SDF2L1 has a basal turnover rate of 0.027/hr. Upon thapsigargin treatment, its
turnover rate increased to 0.048 /hr (adjusted P: 0.07). DNAJB11 also experienced accelerated
kinetics (1.28-fold in thapsigargin, adjusted P 0.029) hence both proteins may be preferentially
synthesized during UPR. On a proteome level, gene set enrichment analysis (GSEA) of temporal
kinetics changes show a preferential enrichment of proteins in unfolded protein response (FDR
adjusted P: $4.1\text{e}{-}4$), ER to Golgi anterograde transport (FDR adjusted P 0.036) and N-linked
glycosylation (FDR adjusted P: $1.7\text{e}{-}3$) but a negative enrichment of translation-related terms
(**Figure 2H**). Overall, protein kinetic changes are modestly correlated with protein abundance
changes (**Supplemental Figure S7**), suggesting that AC16 cells actively regulate protein
synthesis and degradation kinetics in normal and stressed conditions beyond changes in protein
abundance.

258
Figure 2. Simultaneous measurements of spatial and temporal kinetics under UPR. **A.** Bar charts showing
activation of known ER stress markers upon thapsigargin treatment for 8 hours and 16 hours. X-axis: ER
stress markers; y-axis: expression ratio (n=6 normal AC16; n=3 thapsigargin). *: limma adjusted P < 0.01;
: limma adjusted P < 0.001; *: limma adjusted < 0.0001; error bars: s.d. **B.** PC1 and PC2 of proteins
spatial map showing the localization of confidently allocated proteins in normal and thapsigargin-treated
AC16 cells. Each data point represents a protein; color represents classification of subcellular localization.
C. Distribution of light (unlabeled) protein features in each of the 12 subcellular compartments (n=3); fill
color represents whether the protein is also annotated to the same subcellular compartment in UniProt
Gene Ontology Cellular Component terms. **D.** Histograms of the determined log10 protein turnover rates
in control and thapsigargin treated cells (n=3). Text overlay indicates median half-life. **E.** Boxplot showing
the log2 turnover rate ratios in thapsigargin over normal AC16 cells for proteins that are localized to the
ER (blue) (T) or not (F); or the Golgi (GA; green). P values: Mann-Whitney test. A Bonferroni corrected
threshold of 0.05/13 is considered significant. Center line: median; box limits: interquartile range; whiskers:

1.5x interquartile range. **F.** Example of best fit curves in the first-order kinetic model at the protein level
between normal (gray), and thapsigargin treated (red) AC16 cells showing four known ER stress markers
with elevated turnover (HSPA5, RCN3, HSP90B1, and PDIA4). Because the sampling time point is known,
the measured relative isotope abundance of a peptide (prior to reaching the asymptote) is sufficient to
define the kinetic curve and the parameter of interest (k). **G.** Turnover rate ratio (thapsigargin vs. normal)
of the top proteins with elevated temporal kinetics in UPR within the ER (blue) and Golgi (green); .: Mann-
Whitney test FDR adjusted P value < 0.1; *: < 0.05; ** < 0.01; red dashed line: 1:1 ratio; bars: standard
error. **H.** Gene set enrichment analysis (GSEA) of turnover rate ratios in thapsigargin treatment; proteins
with faster kinetics are significantly enriched in DNA damage response and UPR pathways. Color: FDR
adjusted P values in GSEA; x-axis: GSEA normalized enrichment score (NES). Size: number of proteins in
gene set.

Changes in protein subcellular distribution under ER stress

We next analyzed the spatial proteomics component of the data to find proteins that
change in their subcellular localization following thapsigargin treatment. To do so, we used a
Bayesian statistical model implemented in the BANDLE package to estimate the differential
spatial localization of proteins. In total, we identified 1,306 translocating protein features (687
light and 619 heavy) in thapsigargin under a stringent filter of BANDLE differential localization
probability > 0.95 with an estimated FDR of 0.0018 (0.18%), and further filtered using a bootstrap
differential localization probability of > 0.95. We then further prioritized 330 pairs of differentially
localized proteins where the light and heavy features both show confident differential localization
(**Supplemental Data S2**, **Supplemental Data S4**). The differential localization of these 330
proteins recapitulate previously established relocalization events in cellular stress response,
capturing the migration of caveolae toward the mitochondrion under cellular stress (Fridolfsson
et al., 2012) (**Supplemental Figure S8A)**, and the engagement of EIF3 to ribosomes in EIF3-
dependent translation initiation in integrated stress response (Guan et al., 2017) (**Supplemental
Figure S8B**), thus supporting the confidence of the spatial translocation assignment.
From the results, we discerned three major categories of differential localization
behaviors in ER stress that revealed new insights into proteome-wide features of UPR. First, we
observed the externalization of proteins toward the plasma membrane (**Figure 3A**). The large
neutral amino acid transporter component SLC3A2 is localized to the lysosome fraction in normal
cells (Pr > 0.999) but in thapsigargin-treated cells is localized confidently to the plasma
membrane (BANDLE differential localization probability > 0.999) (**Figure 3B**). Showing similar
behaviors are SLC7A5, the complex interacting partner of SLC3A2; and two other amino acid
transporters SLC1A4, and SLC1A5 (**Figure 3C**); whereas the ion channel proteins SLC30A1,
ATP1B1, ATB1B3 and ATP2B1 also showed confident localization toward the cell surface
(**Figure 3C**). The change in localization of SLC3A2 is corroborated by immunostaining (**Figure
3D**), which shows a decrease in co-localization between immunostaining signals of SLC3A2 and
lysosome marker LAMP2 upon thapsigargin treatment (**Figure 3E**).
Second, thapsigargin treated AC16 cells are associated with an increase in proteins
classified into the peroxisome fraction including proteins whose locations changed from the ER,
Golgi, and the nucleus in normal cells (**Figure 3F**). In mammalian cells, ER and peroxisomes are
spatially adjacent; the peroxisome associated fractions sediment prominently at 5000–9000 × g

(F3 and F4) in the LOPIT-DC protocol (**Supplemental Figure S2**), marked by canonical
peroxisome markers PEX14 and ACOX1 (**Supplemental Data S2**). However, although this
compartment was trained using peroxisome markers, the majority of proteins categorized into
this compartment are not annotated to be in the peroxisome whereas 30 out of 49 (61%) of
proteins in the control cells allocated to this compartment were annotated also as endosome,
including canonical markers EEA1 and VPS35L. We thus refer to this compartment hereafter as
peroxisome/endosome. Moreover, proteins that become differentially localized to this fraction in
thapsigargin include known stress granule proteins UBAP2, USP10, CNOT1, CNOT2, CNOT3,
CNOT7, CNOT10, ZC3H7A, and NUFIP2 (**Figure 3G and Supplemental Figure S9)**, which show
high-confidence translocation to the peroxisome/endosome fraction, and are known RNA
binding proteins that participate in phase separation, consistent with stress granule formation in
UPR. Notably, LMAN1, LMAN2, SCYL2, and SNX1 are RNA-binding proteins that are not
currently established stress granule components and show identical translocation patterns,
nominating them as potential participants in RNA granule related processes in AC16 cells for
further studies (**Supplemental Figure S9**). Other proteins in this fraction include the ER-to-Golgi
transport vesicle proteins GOLT1B, GOSR2, RER1, and NAPA (**Supplemental Figure S9**).
Lastly, we see evidence of proteins from the ER and Golgi targeted to the lysosome (see
Tunicamycin section below). Thus taken together, the spatial proteomics component of the data
reveals a complex network of changes in protein spatial distribution during UPR.

333
Figure 3. SPLAT captures extensive protein translocation in AC16 cells under UPR. **A.** Alluvial plot of
translocation events (> 0.99 BANDLE translocation probability; estimated FDR < 1%) following
thapsigargin treatment showing a cohort of proteins moving from the Golgi apparatus (GA) and lysosome
towards the plasma membrane (PM) (n=3). **B.** Protein spatial map for SLC3A2 (open black circle) in normal
(left) and thapsigargin-treated (right) AC16 cells, showing its colocalization with lysosomal proteins in
normal cells and in PM proteins in thapsigargin-treated cells. Colors represent allocated subcellular

localization. **C.** Ultracentrifugation fraction profile of SLC3A2 and other amino acid transporters SLC7A5,
SLC1A4, SLC1A5 and ion channel proteins SLC30A1, ATP1B1, ATP1B3, and ATP2B1 with similar
migration patterns. X-axis: fraction 1 to 10 of ultracentrifugation. Y-axis: relative channel abundance. Bold
lines represent the protein in question; light lines represent ultracentrifugation profiles of all proteins
classified to a respective localization. Colors correspond to subcellular localization in panel B and for all
AC16 data throughout the manuscript; numbers within boxes correspond to BANDLE allocation probability
to compartment. **D.** Immunofluorescence of SLC3A2 (red) against the lysosome marker LAMP2 (green)
and DAPI (blue). Numbers in cell boundary: colocalization score per cell. Scale bar: 90 μm. **E.**
Colocalization score (Mander's correlation coefficient) between SLC3A2 and LAMP2 decreases
significantly (two-tailed unpaired t-test P value: 3e–8) following thapsigargin treatment, consistent with
movement away from lysosomal fraction (n= 205 normal cells, n = 32 thapsigargin treated cells). Center
line: median; box limits: interquartile range; whiskers: 1.5x interquartile range; points: outliers. **F.** Alluvial
plot showing the migration of ER, GA, and nucleus proteins toward the peroxisome/endosome containing
fraction in thapsigargin treated cells. **G.** Ultracentrifugation fraction profile of stress granule proteins
UBAP2, USP10, CNOT1, CNOT2, ZC3H7A, and NUFIP2. RNA Granule Score: score from RNA Granule
Database (https://rnagranuledb.lunenfeld.ca/). A score of 7 or above is considered a known stress granule
protein. Phi: predicted phase separation participation. Circle denotes a prediction of True within the
database. RBP: Annotated RNA binding protein on the RNA Granule Database. One circle denotes known
RBP in at least one data set; two circles denote known RBP in multiple datasets.

Partition of newly synthesized and pre-existing protein pools

We next considered the interconnectivity of temporal and spatial dynamics, namely whether
some localization changes are contingent upon protein pool lifetime, such as where light (old)
protein does not change in spatial distribution but the heavy (new) proteins display differential
translocation upon UPR. Because the spatial profiles of the light and heavy proteins are acquired
independently, this experimental design allowed us to examine whether old and new proteins
are localized to identical cellular locales. In both normal and thapsigargin-treated cells, we found
that the independently measured spatial profiles of light (pre-existing) proteins and their
corresponding heavy SILAC (newly synthesized) counterparts are highly concordant, with a
normalized spatial distribution distance (see Supplemental Methods) of 0.020 [0.015 – 0.030],
compared to 0.117 [0.080 – 0.155] in random pairs of pre-existing proteins (1,614 light-heavy
pairs, Mann-Whitney P < 2.2e–16) in normal cells, and 0.028 [0.019 – 0.041] and 0.122 [0.081
– 0.161] in thapsigargin-treated cells (1,614 light-heavy pairs, Mann-Whitney P < 2.2e–16)
(**Figure 4A**). This robust agreement provided an additional independent confirmation on the
accuracy of the spatial measurements. Consistently, among heavy-light protein pairs with
confidently assigned subcellular localization, the heavy and light proteins are assigned to the
identical subcellular compartment in 93% and 89% of the cases in normal and thapsigargin-
treated AC16 cells, respectively (**Figure 4B**). We focused on the unusual cases where the spatial
distribution distances between the heavy and light proteins increased noticeably following
thapsigargin treatment, as they may be indicative of localization changes that are dependent
upon time since synthesis. These include two proteins EGFR and ITGAV with an uncommon
increase in heavy-light spatial distances (Z: 12.00 and 2.24, respectively) (**Figure 4C**). Epidermal
growth factor receptor (EGFR/ErbB1/HER1) is a receptor tyrosine kinase with multiple subcellular

localizations and signaling roles, and is implicated in cardiomyocyte survival (Lee et al., 2020).
Following a variety of stressors, EGFR is known to be inactivated by intracellular trafficking,
including being internalized to the early endosome and lysosome following oxidative stress and
hypoxia in cancer cells (Tan et al., 2016). In the spatial proteomics data, the spatial distribution
of EGFR borders the lysosome and plasma membrane fractions, which we interpret as EGFR
having potential multiple pools including a cell surface fraction (**Figure 4D**). In the thapsigargin
treated cells, the light-heavy spatial distance of EGFR increased from 0.014 in normal cells to
0.052, and the dynamic SILAC labeled pool (heavy) becomes internalized toward the ER
(BANDLE differential localization probability: >0.999) but not the pre-existing (light) pool (**Figure**
4E). The data therefore suggests that the internalization of EGFR under thapsigargin is likely to
be due to endomembrane stalling or redistribution upon new protein synthesis, possibly leading
to fewer new EGFR molecules reaching the cell surface. Likewise, in thapsigargin-treated AC16
cells, newly-synthesized ITGAV (integrin subunit alpha V) shows a partition from the plasma
membrane fraction to the ER fraction but not the old/existing protein pool, concomitant with an
increase in spatial distribution distance from 0.017 to 0.057 between old and new proteins
(**Figure 4F–G**). With the function of integrins as cell surface receptors that function in
intracellular-to-extracellular and retrograde communication, the ER localization of newly
synthesized ITGAV, such as due to stress-induced stalling of protein trafficking along the
secretory pathway, could indicate a decrease in integrin signaling function through spatial
regulation rather than protein abundance. To partially corroborate the partial redistribution of
EGFR, we performed immunocytochemistry imaging of EGFR subcellular distribution in AC16
cells with or without thapsigargin (**Figure 4H**). Thapsigargin treatment did not increase cell size
(**Figure 4I**), and whereas there is an increase in immunofluorescence signal of EGFR in
thapsigargin (**Figure 4J**), this signal is distributed preferentially to the interior of the cell such that
there is a significant reduction in the ratio of mean intensity at cell borders over the whole cell in
thapsigargin vs. untreated cells (0.673 vs. 0.746, n=93 and 71 cells, Mann-Whitney P: 1.7e–4)
(**Figure 4K**), consistent with a partial redistribution of EGFR toward an internal pool. Taken
together, these examples demonstrate the SPLAT strategy can be used to distinguish time-
dependent differential localization of proteins such as due to the trafficking of newly synthesized
proteins.
412
413
414

Figure 4. SPLAT reveals protein-lifetime dependent translocation. **A.** Histogram showing the similarity in
light and heavy proteins in normalized fraction abundance profiles in (left) normal and (right) thapsigargin-
treated AC16 cells. X-axis: the spatial distribution distance of two proteins is measured as the average
euclidean distance of all TMT channel relative abundance in the ultracentrifugation fraction profiles across
3 replicates; y-axis: count. Blue: distance for 1,614 quantified light-heavy protein pairs (e.g., unlabeled
EGFR, heavy SILAC-labeled EGFR). Grey: distribution of each corresponding light protein with another
random light protein. P value: Mann-Whitney test. **B.** Proportion of heavy-light protein pairs with
confidently assigned localization that are assigned to the same location (purple) in normal (left; 93%) and

thapsigargin-treated (right; 89%) cells. **C.** Ranked changes in heavy-light pair euclidean distance upon thapsigargin treatment. The difference in heavy-light distances in thapsigargin is adjusted by the average changes in the spatial distance of the light protein with 250 other sampled light proteins to calculate the normalized difference. The majority of proteins show no change (+/- 0.02 in euclidean distance). The positions of EGFR and ITGAV are highlighted. Inset: Z score distribution of all changes. **D.** Spatial map showing the location of the light and heavy EGFR in normal and thapsigargin-treated AC16 cells. Each data point is a light or heavy protein species. Colors correspond to other AC16 experiments in the manuscript. Numbers correspond to euclidean distance in fraction profiles over 3 replicates. **E.** Corresponding fraction profiles; x-axis: ultracentrifugation fraction; y-axis: fractional abundance. Post-labeling synthesized EGFR is differentially distributed in thapsigargin and shows ER retention (blue), whereas the preexisting EGFR pool remains to show a likely cell surface localization (pink) after thapsigargin. **F-G**. As above, for ITGAV. **H.** Confocal imaging of EGFR immunofluorescence supports a partial relocalization of EGFR from the cell surface toward internal membranes following thapsigargin treatment. Numbers: The mean intensity of the labeled EGFR channel of a 3 pixel border at cell boundaries was divided by mean intensity of the whole cell to estimate the ratio of EGFR at the plasma membrane to the cell interior. Blue: DAPI; Green: EGFR; scale bar: 90 μm. **I.** Cell areas; Mann-Whitney P: 0.72. Center line: median; box limits: interquartile range; whiskers: 1.5x interquartile range. **J.** Total EGFR intensity per cell; Mann-Whitney P: 2.2e-05. Center line: median; box limits: interquartile range; whiskers: 1.5x interquartile range. **K.** Edge/total intensity ratios in normal and thapsigargin-treated AC16 cells (n=71 normal cells; n=93 thapsigargin cells; Mann-Whitney P: 1.7e–4). Center line: median; box limits: interquartile range; whiskers: 1.5x interquartile range.

Spatiotemporal proteomics highlights similarities and differences of ER stress induction protocols

We next investigated the protein spatiotemporal features of AC16 cells under the treatment of tunicamycin, another compound commonly used to induce ER stress in cardiac cells (Liu et al., 2012; Toro et al., 2022) by inducing proteostatic stress via inhibition of nascent protein glycosylation. Three biological replicate SPLAT experiments were performed in tunicamycin-treated cells to resolve the temporal kinetics and subcellular localisation of proteins (**Supplemental Figures S1, S2C, S5B, S6, S7B, S10**). Tunicamycin treatment at 1 μg/mL for 16 hours robustly induced the known ER stress response markers BiP/HSPA5, HSP90B1, PDIA4, CALR, CANX, and DNAJB11 (limma FDR adjusted P < 0.10) (**Figure 5A**) demonstrating effective ER stress induction. Overall, tunicamycin treatment led to a lesser slowdown of temporal kinetics than thapsigargin (average protein half-life 32.6 hours) (**Figure 5B; Supplemental Data S5**). As in thapsigargin treatment, the kinetic changes following tunicamycin are different across cellular compartments, with ER and Golgi proteins having relatively faster kinetics than other cellular compartments (Mann-Whitney test P: 1.3e–13 and 6.6e–5; << Bonferroni corrected threshold 0.05/13); whereas the greatest reduction was observed among proteins localized to the lysosome, a compartment closely linked to glycosylation and recycling of glycans (Mann-Whitney test P: 1.8e–10) (**Figure 5C**). Gene set enrichment analysis (GSEA) of turnover rate ratios revealed a significant positive enrichment of UPR proteins (adjusted P 3.5e–3) and DNA repair terms (e.g., processing of DNA double-strand break ends; adjusted P 0.017) and a negative enrichment of translation related terms (**Figure**

5D). Compared to thapsigargin treatment however, no significant enrichment of glycosylation
and vesicle transport related terms were found in tunicamycin. Inspection of individual protein
kinetics changes likewise revealed both similar induction of the ER stress response markers
HSPA5, HSP90B1 and PDIA4 as in thapsigargin treatment, but other stress response genes
PDIA3 and NIBAN1 are not induced in thapsigargin (**Figure 5E**). On the other hand, RCN3
(reticulocalbin 3) is an ER lumen calcium binding protein that regulates collagen production
(Martínez-Martínez et al., 2017) and shows increased temporal kinetics in thapsigargin (**Figure
2F**) but not in tunicamycin (ratio 0.76 over normal; **Supplemental Data S5**), altogether
reflecting potential differences in stress response modality to a different ER stress inducer.
Parallel to the less prominent changes in vesicle transport, tunicamycin treatment also
led to fewer translocating proteins than thapsigargin, with 620 translocating features (including
282 light proteins and 338 heavy proteins) at BANDLE differential localization probability >
0.95, corresponding to an estimated FDR of 0.35%, and thresholded by bootstrapping
differential localization probability > 0.95; from which we highlighted 109 proteins where the
heavy and light versions both showed translocation. The spatial data revealed a high degree of
similarity but also notable differences with thapsigargin-induced ER stress. We found that in
both tunicamycin and thapsigargin treatment, there was evidence of lysosome targeting from
other endomembrane compartments, including: RRBP1, a ribosome-binding protein of the ER,
GANAB, a glucosidase II alpha subunit integral to the proper folding of proteins in the ER,
FKBP11, a peptidyl-prolyl cis/trans isomerase important to the folding of proline-containing
peptides, IKBIP, an interacting protein to the IKBKB nuclear kinase, and MANF, a neurotrophic
factor which has relations to ER stress-related cell death when its expression is lowered
(BANDLE differential localization probability > 0.999) (Sayers et al., 2022) (**Figure 5F**). Among
these proteins was the collagen synthesis enzyme P3H1 in both thapsigargin and tunicamycin.
Interestingly, prior work found no correlation between the protein abundance of collagen
modifying enzymes with the known reduction of collagen synthesis in chondrocytes and
fibroblasts under ER stress (Vonk et al., 2010). The results here suggest that the functional
decline may instead correlate with a change in the subcellular localization of collagen
modifying enzymes in AC16 cells. Tunicamycin treatment also induced old-new protein
partitions in EGFR and ITGAV as observed in thapsigargin (**Supplemental Figure S11**).
Notably, although tunicamycin also induced the translocation of proteins toward the
peroxisome/endosome fraction, different proteins are involved, including the stress response
proteins DNAJB11, DNAJC3, DNAJC10, and PDIA6 as well as other proteins EMC4, EMC8,
VAPA, and VAPB (**Supplemental Figure S12**) which further outlines different modalities of
cellular response toward two different ER stress inducers. The translocating stress response
proteins DNAJB11, DNAJC10, and PDIA6 also showed significant acceleration in temporal
kinetics in tunicamycin (Mann-Whitney test, FDR adjusted P < 0.10; **Supplemental Data S6**)
which is consistent with specific production of the proteins followed by shuttling to subcellular
location for their function during stress response.

Figure 5. Comparison of ER stress induction methods. **A.** Bar charts showing activation of known ER
stress markers upon tunicamycin treatment for 8 hours and 16 hours. X-axis: ER stress markers; y-axis:
expression ratio (n=6 normal AC16; n=3 tunicamycin). *: limma adjusted P < 0.01; **: limma adjusted P <
0.001; ***: limma adjusted P < 0.0001; error bars: s.d.. **B.** Histograms of the determined log10 protein
turnover rates in control and tunicamycin treated cells (n=3). **C.** Boxplot showing the log2 turnover rate
ratios in tunicamycin over normal AC16 cells for proteins that are localized to the ER (T) or not (F); Golgi
apparatus, or the lysosome. P values: two-tailed t-test. Center line: median; box limits: interquartile range;
whiskers: 1.5x interquartile range. A Bonferroni corrected threshold of 0.05/13 is considered significant.
D. Gene set enrichment analysis (GSEA) of turnover rate ratios in tunicamycin treatment. Color: FDR
adjusted P values in GSEA; x-axis: GSEA normalized enrichment score (NES). Size: number of proteins in
the gene set. **E.** Example of best fit curves in the first-order kinetic model at the protein level between

normal (gray), tunicamycin (blue) and thapsigargin (red) treated AC16 cells showing known ER stress
markers with elevated turnover in both ER stress inducers (HSPA5, HSP90B1, and PDIA4) as well as stress
response proteins with elevated turnover only in tunicamycin (PDIA3, DNAJB11, NIBAN1). **F.** Alluvial plot
showing the migration of ER, GA, and peroxisome/endosome proteins toward the lysosome (left). On the
right, the ultracentrifugation fraction profiles of translocating proteins RRBP1, FKBP11, GANAB, MANF,
IKBIP, and P3H1 are shown that are targeted toward the lysosome in both tunicamycin and thapsigargin
treatment (BANDLE differential localization probability > 0.95). Numbers in boxes are the BANDLE
allocation probability in each condition (n=3).

Application of SPLAT to the mechanism of cardiotoxicity in iPSC-CM models

We next assessed the applicability of SPLAT toward a different, non-proliferating cell type
(human iPSC-derived cardiomyocytes [iPSC-CMs]) and its utility for interrogating the mechanism
of cardiotoxicity following proteasome inhibitor treatment (**Figure 6A**). The ubiquitin proteasome
system is responsible for the degradation of over 70% of cellular proteins. Compounds that
inhibit proteasome function, including carfilzomib, are widely used as cancer treatment and have
led to remarkable improvement in the survival of multiple myeloma patients. Mechanistically,
carfilzomib functions by binding to and irreversibly inhibiting the proteasome catalytic subunit
PSMB5 (Cromm and Crews, 2017), leading to the accumulation of unfolded proteins in cancer
cells. Importantly, despite its efficacy as a cancer treatment, carfilzomib also leads to cardiotoxic
adverse effects including heart failure (<20%), arrhythmia (<10%), and hypertension (11-37%) in
a significant number of patients (IBM Watson Health, 2023). This cardiotoxicity has been
modeled in vitro by exposure of 0.01 – 10 µM carfilzomib to human iPSC-CMs (Forghani et al.,
2021), yet the molecular mechanisms of carfilzomib cardiotoxicity remain incompletely
understood. To examine the protein spatiotemporal changes upon carfilzomib-mediated
proteasome inhibition in cardiac cells, we differentiated contractile iPSC-CMs using a small
molecule based protocol, and treated the cells with 0.5 µM carfilzomib. To verify toxicity
modeling, we measured iPSC-CM viability and phenotypes under 0 to 5 µM carfilzomib for 24
and 48 hours. Under the chosen treatment (0.5 µM for 48 hours), iPSC-CMs showed sarcomeric
disarray consistent with prior observations on the cardiotoxic effects of carfilzomib (**Figure 6B**)
but maintained viability (82%) (**Figure 6C**), while showing significant decreases in oxygen
consumption (**Figure 6D**), basal respiration (**Figure 6E**), and maximal respiration (**Figure 6F**),
whereas higher doses are accompanied with drops in viability at 48 hours and an increase in
proton leak (**Figure 6G**). ATP production at the 0.5 µg dose was significantly lower than untreated
cells at both 24 and 48 hours (**Figure 6H**). Therefore cardiotoxicity due to carfilzomib can be
recapitulated in a human iPSC-CM model, consistent with prior work in the literature.
From the untreated and carfilzomib-treated (0.5 µM, 48 hrs) human iPSC-CMs, we
constructed a protein subcellular spatial map that takes into account several features of the
iPSC-CM cell type, including the inclusion of cell junction and desmosome proteins, as well as
a sarcomere protein compartment, that are not apparently recognized as discrete compartments
in the prior spatial maps (**Supplemental Figure S13**). In addition, the 40S and 60S ribosomes
showed clear separation in iPSC-CM unlike in AC16 cells and are hence classified separately.
This separation is consistent with less active protein translation in this cell type. In total, we
mapped the subcellular localization of 5,047 protein features including 2,680 old proteins and

2,367 new proteins using a stringent two-peptide filter at 1% FDR of protein identification,
including 2,010 old proteins and 1,737 new proteins assigned to one of 13 subcellular localization
with >95% confidence after removing outliers (**Figure 6I; Supplemental Data S7**). The iPSC-
CM spatial map achieved similar levels of concordance with known cellular component
annotations as in AC16 cells (70.8% with known annotation matching the assigned compartment
in normal iPSC-CM; 63.0% in carfilzomib-treated cells) (**Supplemental Figure S14**). The iPSC-
CM map has comparable spatial distance between light and heavy protein pairs as in AC16 cells,
and 87.6% heavy-light protein pairs map to the same compartment in the baseline, supporting
that there is sufficient spatial resolution to resolve subcellular compartment differences in this
cell type (**Supplemental Figure S14**).

Figure 6. Applicability in human iPSC-derived cardiomyocytes. A. Schematic of human iPSC
differentiation into cardiomyocytes, carfilzomib treatment, and SPLAT analysis. **B.** Confocal microscopy
images showing sarcomeric disarray in iPSC-CMs upon 48 hrs of 0.5 μM carfilzomib; green: cTNT, red:
alpha-actinin; blue: DAPI; scale bar: 20 μm. **C–H.** Cell viability (%), normalized Seahorse oxygen
consumption rate (OCR; pmol/min), basal respiration, maximal respiration, proton leak, and ATP
production upon 0 – 5 μM carfilzomib for 24 or 48 hrs; .: adjusted P < 0.1; *: adjusted P < 0.05; **: adjusted
P < 0.01, ANOVA with Tukey's HSD post-hoc at 95% confidence level; n=5. Error bars: s.d. for bar charts
in panels C, E, F, G, H; s.em. for the OCR graph in panel D. Colors in panel D: dosage, same as panel C.
O: Oligomycin; AA/R: Antimycin A/Rotenone. **I.** Spatial map with 13 assigned subcellular localizations in
iPSC-CMs at the baseline (top) and upon 0.5 μM carfilzomib treatment (n=2). **J.** Histogram of log10 protein
turnover rates (k), with median half-life of 97.4 hours and 100.0 hours in normal and carfilzomib-treated
iPSC-CM. **K.** Proteasome activity in iPSC-CMs treated with 0 (Ctrl) vs. 0.5 μM carfilzomib (Cfzb) for 48
hrs. P value: two-tailed t-test; n = 3. **L.** Autophagy assay for iPSC-CMs treated with 0 (Ctrl) vs. 0.5 μM
carfilzomib (Cfzb) for 48 hrs, and positive control (Pos); data were normalized to DAPI and normal cells. P
value: two-tailed t-test; n = 10. **M.** log2 Turnover rate ratios between carfilzomib-treated and untreated
iPSC-CM from the spatiotemporal proteomics data. Proteins assigned the proteasome compartment have
significantly increased temporal kinetics; proteins in the lysosome/junction and chromatin/sarcomere
compartments have significantly reduced temporal kinetics. P values: Mann-Whitney; with a threshold of
0.05/14 considered significant. Center line: median; box limits: interquartile range; whiskers: 1.5x
interquartile range.

Among proteins with spatial information, we compared the temporal kinetics of 2,648
proteins. Unexpectedly, there was no overall slowdown of protein temporal kinetics with the
median protein half-lives being 97.4 hours and 100.0 hours in normal and carfilzomib-treated
cells, respectively (**Figure 6J**; **Supplemental Data S8**), suggesting that at 48 hours following
proteasome inhibitor treatment, the observed cellular toxicity is not directly explainable by a drop
of per-protein average in global protein degradation. At 48 hours of carfilzomib treatment in
iPSC-CMs, proteasome chymotrypsin-like activities are partially suppressed but significant
partial activities are also observable (**Figure 6K**); whereas other proteolysis mechanisms may
also compensate for proteasome inhibition, including a suggestive increase in autophagy (P:
0.053) (**Figure 6L**). Inspection of the spatial data revealed that the changes in protein kinetics
upon carfilzomib are localization specific, with a significant reduction in chromatin/sarcomere
protein turnover rate, and significant increase for the proteasome compartment under carfilzomib
treatment (**Figure 6M**).

Notably, on an individual protein level we find that the majority of proteins with increased
protein kinetics belong to subunits of the regulatory 19S complex rather than the core 20S
complex (**Figure 7A**) suggesting possible changes in 26S proteasome activity and target
engagement. In addition to proteasome subunits, the temporal kinetics revealed a robust
induction of chaperons HSP90AA1/HSP90A, HSP90AB1/HSP90B, HSPA4, and BAG3; and
ERAD associated proteins VCP and UFD1 (**Figure 7B, Supplemental Data S8**). Within the
mitochondrion, quality control proteins HSPD1, HSPE1, and CLPB are induced (**Figure 7A**). In
contrast, among proteins that show reduced turnover in carfilzomib treatment are major
sarcomeric proteins MYH6, MYH7, MYBPC3, MYL7; as well as proteins classified to the cell

junction compartment dystrophin (DMD) and utrophin (UTRN), and the desmosome complex
protein desmoplakin (DSP) (**Figure 7A & C**).

We observed an interconnectivity of spatial and temporal changes, with 23 out of 339
pairs of confident translocators also showing significant kinetic changes. BAG3, a muscle
chaperone important for sarcomere turnover (Martin et al., 2021a), shows elevated kinetics
(**Figure 7B**) and a partition away from the soluble cytosol compartment (**Figure 7D**) toward an
expanded compartment in carfilzomib that co-sediments with Golgi markers. Inspection of
existing annotations show that the majority of categorized proteins are not canonical Golgi
proteins but contain cytoplasm and endosome terms; hence it likely represents a less soluble
cytoplasmic fraction consistent with a lower abundance in the last ultracentrifugation step
(**Supplemental Figure S13**, **Supplemental Figure S14, Supplemental Data S9**). This is
consistent with the known dynamic partitioning of BAG3 between the cytosol and myofilament
fractions for its function (Martin et al., 2021b). Secondly, we find that accelerated temporal
kinetics of PA200/PSME4 proteasome activator (**Figure 7B**) in conjunction with a change in
localization from the nuclear compartment in baseline toward the proteasome compartment
upon carfilzomib (**Figure 7E**). The PSME4/PA200 proteasome activator is known to bind with the
20S/26S proteasome complex to stimulate proteolysis and has a putative nuclear localization
signal (Ustrell et al., 2002). The change in localization is therefore consistent with increased
binding with the proteasome complex. In parallel, the proteasome activator PA28/PSME3 also
relocalizes to the proteasome upon carfilzomib (**Supplemental Figure S15**), altogether
suggesting a remodeling of proteasome configuration upon carfilzomib.

Taken together, the spatiotemporal proteomics data here identified major proteostatic
pathways induced in carfilzomib, involving a potential remodeling of the proteasome, induction
of chaperones and ERAD proteins, and mitochondrial protein quality control mechanisms that
may be important for preserving function. On the other hand, a preferential decrease of temporal
kinetics in sarcomere and desmosome proteins suggest that the interruption of protein quality
control and turnover in these important cardiomyocyte components may be principal sites of
lesion in carfilzomib cardiotoxicity. Finally, we assessed the protein-level expression profiles in
the hearts of mice treated with carfilzomib for 2 weeks to model cardiac dysfunction
(Supplemental Methods). Notably, we find differential protein abundance analysis showed that
MHC-β (MYH7) and desmoplakin (DSP) are the 1st and 5th most significantly up-regulated
proteins among 3,379 quantified proteins in the hearts of mice treated with carfilzomib
(**Supplemental Figure S16**), consistent with their accumulation following proteasome inhibition
and suggesting the possibility that similar proteostatic lesions may underlie cardiotoxicity
mechanism in vivo.

Figure 7. Proteostatic pathways and lesions in carfilzomib mediated cardiotoxicity in iPSC-CMs. **A.** Changes in protein turnover rates between carfilzomib vs. normal iPSC-CMs across selected cellular compartments; **: P < 0. 01; *: P < 0.05; .: P < 0.1; Mann-Whitney test FDR adjusted P values. error bars: standard error. **B.** Kinetic curve representations of proteins with accelerated temporal kinetics in carfilzomib including PSMC2 which corresponds to the ratio in panel A, as well as additional ERAD proteins and chaperones; gray: normal iPSC-CM; green: carfilzomib. **C.** Kinetic curve representations of slowdown of protein kinetics in DSP, DMD, MYH6, and MYH7, corresponding to the ratios in panel A. **D–E.** Spatial map (PC1 vs. PC2) and ultracentrifugation fraction profiles of **D.** BAG3 and **E.** PSME4 in normal and carfilzomib-treated human iPSC-CM, showing a likely differential localisation in conjunction with kinetics changes. White-filled circles: light and heavy BAG3 or PSME4 in each plot. The kinetic curves of BAG3 and PSME4 are in panel B. Numbers at arrows correspond to BANDLE differential localization probability (Diff. Loc. Pr.).

Discussion

Advances in spatial proteomics have opened new avenues to discover the subcellular
localization of proteins on a proteome scale. Thus far however, few efforts have linked the spatial
dynamics of the proteome to other dynamic proteome parameters, which hinders a multi-
dimensional view of protein function (Burnum-Johnson et al., 2022; Larance et al., 2013). Co-
labeling of SILAC (MS1) and TMT (MS2) tags have been used to increase the channel capacity
of quantitative proteomics experiments (Dephoure and Gygi, 2012). Here we adopted the
extended labeling capacity to encode spatial and temporal information in the same experiment
(**Figure 1D**). The SPLAT design has the advantages of resolving proteome-wide temporal
kinetics and spatial distributions in the same experiments and identifying their interactions such
as compartment-specific turnover; secondly, it allows separate observations of spatial
distribution of new vs. old protein pools. Despite each of the SILAC labeled pairs (light and heavy)
and their associated TMT profiles being separately quantified by mass spectrometry as
independent ions, the spatial profiles of light and heavy proteins are highly similar under baseline
conditions (**Figure 4A-B**) which provides additional assurance of spatial assignments.

Applying the workflow to human AC16 cardiac cells under thapsigargin and tunicamycin
induced ER stress and the associated UPR, we observed that both ER stress inducing drugs led
to a global suppression in turnover rate, consistent with the reduced translation known to be
caused by ER stress. The temporal kinetics data of individual proteins revealed the coordinated
activation of known and suspected stress mediators through their increased kinetics, particularly
concentrated in the ER and Golgi compartments (**Figure 2E**). At the same time, the spatial
proteomics profiles revealed substantial endomembrane remodeling and hundreds of
translocating proteins under ER stress. A recent work also reported ~75 translocator protein
candidates under acute low-dose thapsigargin (250 nM for 1 hr) in U-2 OS cells, although the
experimental approach was more optimized toward mRNA detection (Villanueva et al., 2022).
The spatial data here add to an emerging view of dynamical protein regulation under cellular
stress, illustrating a differential localization of RNA binding proteins to stress granules, targeting
of specific proteins toward lysosomes, as well as membrane trafficking of ion channels and
amino acid transporters. UPR activation is known to induce the biosynthesis of non-essential or
partly-essential amino acids despite protein synthesis suppression (Gonen et al., 2019); the
recycling of lysosomal lysine and arginine regulates the sensitivity to ER stress (Higuchi-Sanabria
et al., 2020); whereas deprivation of amino acids is known to activate downstream pathways of
UPR including CHOP in vitro (Harding et al., 2003). Moreover, the knockdown of SLC3A2 has
been found to suppress the activation of ER stress response pathways including ATF4/6
induction (Liu et al., 2018), together suggesting amino acid transporters may function in ER stress
response. The data here indicate that these transporters may in turn be regulated by their spatial
localization beyond steady-state abundance. Other changes are found that may be specific to
stressors. Tunicamycin treatment leads to a further decrease in turnover of lysosomal proteins
compared to other organelles. The lysosome plays important roles in recycling of glycans, a
process which may be slowed under ER stress in response to the decrease in glycosylation.

Changes in protein spatial distribution can occur due to a relocation of an existing protein,
where in a protein may respond to a signaling cue such as a post-translational modification

status and subsequently migrate to a subcellular location. Alternatively, an alternate localization
of newly synthesized proteins can also drive spatial redistribution. By comparing the
translocation behaviors of new and old protein pools separately, we were able to observe a
partitioning of new and old protein pools including epidermal growth factor receptor
(EGFR/ErbB1/HER1). ErbB family proteins are required for both normal heart development and
prevention of cardiomyopathies in the adult heart. EGFR is a receptor tyrosine kinase of this
family capable of triggering multiple signaling cascades, and can be activated via both ligand
dependent and ligand independent pathways. Upon ER stress induction, EGFR
immunofluorescence showed a partial translocation away from the plasma membrane. Although
immunofluorescence cannot distinguish between old and new protein pools, this partial
translocation is consistent with the mass spectrometry data showing partial translocation,
involving the newly synthesized heavy protein pool. We hypothesize that this partial translocation
is suggestive of a ligand-independent trafficking of newly synthesized protein, rather than ligand
dependent activation and internalization that is agnostic to protein lifetime. Of note, ligand-
independent activation and internalization of EGFR has been previously induced via both
starvation and tyrosine kinase inhibitor treatment, leading to cellular autophagy (Tan et al., 2015),
hence this partial translocation may carry functional significance to protective cellular response.
We further applied SPLAT to a different, non-proliferating cell type, namely human iPSC-
derived cardiomyocytes, which have gained increasing utility for modeling the cardiotoxic effects
of the cancer drugs, including carfilzomib. In the carfilzomib experiment, the data from SPLAT
revealed a surprising similarity in global turnover rates between control and treatment. These
observations are consistent with a compensatory rescue of proteasome abundance and activity
previously observed in proteasome inhibition by carfilzomib (Demo et al., 2007; Forghani et al.,
2021) or bortezomib (Meiners et al., 2003) in other cell types. Proteasomes are known to be
regulated by negative feedback mechanisms (Meiners et al., 2003; Xie and Varshavsky, 2001;
Xu et al., 2008), which could explain the lack of change in proteome-wide half-life differences
and instead suggest that toxicity may derive from more specific cellular lesions. We identified a
significant reduction in turnover in sarcomere proteins, which may be particularly sensitive to
interruptions in proteasome activity and moreover may account for the bulk of turnover flux in
iPSC-CMs given their high abundance. Finally, the activation and translocation of proteostatic
pathway proteins BAG3 and PA200/PSME4 in cardiac cells may be explored as potential targets
to ameliorate proteostatic disruptions and cardiotoxic effects.

Limitations

Although there is no inherent limit to the number of dynamic SILAC labeling time points that can
be investigated, in our experiments we have used only a single time point per treatment (16 hours
post thapsigargin or tunicamycin; 48 hours post carfilzomib), which was selected based on the
drug treatment models but also needed to be sufficient to capture the median half-life of proteins
in the cell types studied (**Figure 2D; Figure 6J**). Hence the collection time points need to be
optimized for different cell types with distinct intrinsic protein turnover rates. Synthesized
proteins may have further relocalized following the start of dynamic SILAC labeling, hence some
acute translocation responses may be missed. Although the inclusion of earlier time points might

provide insight into early translocation events, reaching this objective may be technically
challenging, as the acquisition of spatial localization information from the heavy SILAC labeled
peptides would be hindered by their low intensity. The double labeling design also requires
independent MS2 acquisition of light and heavy peptides, which can decrease the depth and
data completeness of mass spectrometry-based proteomics analysis. Future work may alleviate
this limitation by modifying the mass spectrometry acquisition methods to automatically trigger
the acquisition of heavy peptides and reduce incomplete light-heavy pairs.

Secondly, SPLAT shares limitations that are common to common spatial proteomics
strategies. The differential ultracentrifugation method employed here requires ~10^7 cells and
cannot resolve some subcellular fractions, e.g., lysosome from cell junction. The number of
classifiable subcellular localizations here is in line with other LOPIT-DC studies, and may be
improved in future work that attempts to couple turnover analysis to gradient-based
sedimentation approaches with higher spatial resolution. Protein correlation profiling based
techniques generally face challenges in recognizing proteins with multiple localizations or partial
translocations. For instance, the multi-functional ERAD protein p97/VCP is known to have
multiple subcellular localizations, but its precise subcellular translocation profile is difficult to
interpret from the TMT data and is unclassified to any compartment. Because translocation may
be sub-stoichiometric, translocated proteins can have lower confidence in classification of
location. Hence, other biologically relevant translocators reside in the data that await exploration.
In the thapsigargin and tunicamycin experiments, upward of 1,000 candidate translocation
patterns were detected with significant differences in localization (99% probability) using
BANDLE, but many proteins presented a challenge to clear interpretation of compartments upon
manual inspection. Progress in this area may require development of spatial separation methods
that combine orthogonal separation principles.

In summary, we describe an experimental workflow and data analysis pipeline that
integrates dynamic time-resolved stable isotope labeling kinetic analysis with differential
ultracentrifugation-based subcellular proteomics to characterize proteome-wide spatial and
temporal changes upon perturbation. This method may be broadly useful for understanding the
function and behaviors of proteins inside the cell, and may provide new insight into the
mechanisms that regulate protein stability and localization in stress, disease, and drug treatment.

Methods

774 Additional methods can be found in the Supplemental Information file.

AC16 Cell culture, metabolic labeling, UPR induction

AC16 cells procured from Millipore between passage number 11 and 16 were cultured in
DMEM/F12 supplemented with 10% FBS and no antibiotics. Cells were maintained at 37°C with
5% CO_2 and 10% O_2. For isotopic labeling, SILAC DMEM/F12 (Thermo Scientific) deficient in
both L-lysine and L-arginine was supplemented with 1% dialyzed FBS and heavy amino acids
$^{13}C_6{}^{15}N_2$ L-Lysine-2HCl and $^{13}C_6{}^{15}N_4$ L-Arginine-HCl (Thermo Scientific) at concentrations of

0.499 mM and 0.699 mM, respectively. Light media was switched to heavy media and cells were
labeled for 16 hours prior to harvest. UPR was induced with 1 µM thapsigargin (SelleckChem) or
1 µg/mL tunicamycin (Sigma) at the same time as isotopic labeling.

Cell harvest, differential centrifugation, and isobaric labeling

Cell harvest and subcellular fractionation was performed based on the LOPIT-DC differential
ultracentrifugation protocol as described in Geladaki et al. (Geladaki et al., 2019). Briefly, AC16
cells were treated, harvested with trypsinization, washed 3× with room temperature PBS, and
resuspended in a detergent free gentle lysis buffer (0.25 M sucrose, 10 mM HEPES pH 7.5, 2
mM magnesium acetate). 1.5 mL of suspension at a time was lysed using an Isobiotec ball
bearing homogenizer with a 16 µM clearance size until ~80% of cell membranes were lysed, as
verified with trypan blue (approximately 15 passes through the chamber). Lysates were spun 3
times each in a 4°C swinging bucket centrifuge 200 × g, 5 min to remove unlysed cells. The
supernatant was retained and used to generate the 9 ultracentrifugation pellets using spin
parameters shown in **Supplemental Table S2.**

The supernatant generated in the final spin was removed and all pellets and the final supernatant
were stored at -80°C until proceeding. Supernatant was thawed on ice and precipitated in 3×
the volume of cold acetone overnight at –20°C.This was used to generate pellet 10 by
centrifuging at 13,000 × g for 10 minutes at 4°C. Excess acetone was removed and the pellet
was allowed to dry briefly before resuspension in a resolubilization buffer of 8 M urea, 50 mM
HEPES pH 8.5, and 0.5% SDS with 1x Halt Protease and Phosphatase Inhibitor Cocktail (Thermo
Scientific). The suspension was sonicated in a Biorupter with settings 20× 30s on 30s off at 4°C.

Pellets from the ultracentrifugation fractions 1 to 9 were resuspended in RIPA buffer with Halt
Protease and Phosphatase Inhibitor Cocktail (Thermo Scientific) with sonication in a Biorupter
with settings 10x 30s on 30s off at 4°C. Insoluble debris was removed from all samples (1-10) by
centrifugation at 14,000 × g, 5 minutes. Protein concentration of all samples was measured with
Rapid Gold BCA (Thermo Scientific). The samples were digested and isobarically tagged using
the iFASP protocol (McDowell et al., 2013). 25 ug protein per sample in 250 uL 8M urea was
loaded onto Pierce Protein Concentrators PES, 10K MWCO prewashed with 100 mM TEAB. The
samples were again washed with 8 M urea to denature proteins and remove SDS. The samples
were washed with 300 uL 100 mM TEAB twice. The samples were then reduced and alkylated
with TCEP and CAA for 30 minutes at 37°C in the dark. CAA and TCEP were removed with
centrifugation and the samples were washed 3x with 100 mM TEAB. Samples were digested
atop the filters overnight at 37°C with mass spectrometry grade trypsin (Promega) at a ratio of
1:50 enzyme:protein. A total of 0.2 mg of TMT-10plex isobaric labels (Thermo Scientific) per
differential centrifugation fraction were equilibrated to room temperature and reconstituted in 20
µL LC-MS grade anhydrous acetonitrile. In each experiment, labels were randomly assigned to
each fraction (**Supplemental Tables S3-S4**) with a random number generator to mitigate
possible batch effect. Isobaric tags were added to peptides still atop the centrifugation filters
and incubated at room temperature for 1 hour with shaking. The reactions were quenched with

1 µL 5% hydroxylamine at room temperature for 30 minutes with shaking. Labeled peptides were
eluted from the filters with centrifugation. To further elute labeled peptides 40 µL 50 mM TEAB
was added and filters were again centrifuged. All 10 labeled fractions per experiment were
combined and mixed well before dividing each experiment into two aliquots. Aliquots were dried
with speed-vac and stored at –80°C.

Liquid chromatography and mass spectrometry

One aliquot per experiment was reconstituted in 50 µL 20 mM ammonium formate pH 10 in LC-
MS grade water (solvent A) for high pH reverse phase liquid chromatography (RPLC)
fractionation. The entire sample was injected into a Jupiter 4 µm Proteo 90 Å LC Column of 150
× 1 mm on a Ultimate 3000 HPLC system. The gradient was run with a flow rate of 0.1 mL/min
as follows: 0–30 min: 0%–40% Solvent B (20 mM ammonium formate pH 10 in 80% LC-MS
grade acetonitrile); 30–40 min: 40%-80% Solvent B; 40–50 min: 80% Solvent B. Fractions were
collected every minute and pooled into a total of 20 peptide fractions, then dried with speed-
vac.
The dried fractions were reconstituted in 10 µL each of pH 2 MS solvent A (0.1% formic
acid) and analyzed with LC-MS/MS on a Q-Exactive HF orbitrap mass spectrometer coupled to
an LC with electrospray ionization source. Peptides were separated with a PepMap RSLC C18
column 75 µm x 15 cm, 3 µm particle size (ThermoScientific) with a 90 minute gradient from 0 to
100% pH 2 MS solvent B (0.1% formic acid in 80% LC-MS grade acetonitrile). Full MS scans
were acquired with a 60,000 resolution. A stepped collision energy of 27, 30 and 32 was used
and MS2 scans were acquired with a 60,000 resolution and an isolation window of 0.7 m/z.

Mass spectrometry data processing and turnover analysis

Mass spectrometry raw data were converted to mzML format using ThermoRawFileParser
v.1.2.0 (Hulstaert et al., 2020) then searched against the UniProt Swiss-Prot human canonical
and isoform protein sequence database (retrieved 2022-10-27) using Comet v.2020_01_rev3
(Eng et al., 2015). The fasta database was further appended with contaminant proteins using
Philosopher v4.4.0 (total 42,402 forward entries). The search settings were as follows: peptide
mass tolerance: 10 ppm; isotope error: 0/1/2/3; number of enzyme termini: 1; allowed missed
cleavages: 2; fragment bin tolerance: 0.02; fragment bin offset: 0; variable modifications: TMT-
10plex tag +229.1629 for TMT experiments, and lysine + 8.0142, arginine + 10.0083 for all SILAC
experiments; fixed modifications: cysteine + 57.0214. The search results were further reranked
and filtered using Percolator v3.0 (The et al., 2016) at a 5% FDR. Following database search, the
mzML files and Percolator PSMs were input to the SPLAT pipeline. The dynamic SILAC data
were analyzed using RIANA v0.7.1 (Hammond et al., 2022) to integrate the peak intensity within
a 25 ppm error of the light (+0), heavy (+8, +10), and double K/R (+16, +18, +20) peptide peaks
over a 20 second retention time window encompassing the first and last MS2 scan where the
peptide is confidently identified. We then calculated the fractional synthesis of all K/R containing
peptides as the intensity of the 0th isotopomer peak (m0) over the sum of applicable light and
heavy isotopomers (e.g., m0/m0+m8 for a peptide with one lysine). RIANA then performs

intensity-weighted least-square curve-fitting using the scipy optimize function to a first-order
exponential rise model to find the best-fit peptide turnover rate. Protein turnover rates are
calculated as the harmonic mean of peptide turnover rates

Subcellular localization classification

Subcellular localization classification and translocation predictions were performed using the
pRoloc (Gatto et al., 2014) and the BANDLE (Crook et al., 2022) packages in R/Bioconductor.
Three replicate batches of AC16 cells per condition each were individually labeled and treated,
fractionated and analyzed by mass spectrometry, and biological replicate data were used for
pRoloc and BANDLE analysis. Briefly, the subcellular localization markers were selected from
the intersecting proteins with a prior data set generated from human U-2 OS osteosarcoma cells
(Geladaki et al., 2019) with further curation to account for cell type specific marker expression
(**Supplemental Data S10**). A random walk algorithm is used to prune the markers to maximize
normalized between-class separation. For differential localization analyses we used the Markov-
chain Monte-Carlo (MCMC) and non-parametric model in BANDLE to find unknown protein
classification and evaluate differential localization probability. MCMC parameters are 9 chains,
10,000 iterations, and 5,000 burn-in, 20 thinning, seed 42; convergence of the Markov chains is
assessed visually by rank plots as recommended (Crook et al., 2019). For additional analysis to
describe baseline protein classification and compare MS2 and MS3 performance, a T-
augmented Gaussian mixture model with a maximum a posteriori (MAP) method in pRoloc was
used.

Human iPSC-derived cardiomyocytes and proteasome inhibition

Human AICS-0052-003 induced pluripotent stem cell (iPSC) (mono-allelic C-terminus mEGFP-
tagged MYL7 WTC; Allen Institute Cell Collection) line was acquired from Coriell Institute and
seeded onto Geltrex (Gibco) coated 6 well plates and maintained in StemFlex (Thermo Scientific)
media at 37 °C, 5% CO2 with daily media changes. At 80% confluency, cells were passaged
using 0.5 mM EDTA before resuspension in StemFlex supplemented with 10 µM Y-27632
(Selleck). Cells were replated into Geltrex coated 12 well plates at a density of 3×10^5 cells/well
and daily media changes of StemFlex continued until the cells reached 80% confluency, day 0
of cardiac differentiation. The iPSCs were differentiated into cardiomyocytes using a small
molecule based GSK-3 inhibition-Wnt inhibition protocol (Burridge et al., 2014). Briefly, on day
0, cell media was replaced with 2 mL/well RPMI supplemented with B-27 minus insulin (Gibco)
and 6 µM CHIR99021 (STEMCELL); on day 2, the media was changed to 2 mL/well RPMI+B-27
minus insulin. On day 3, the media was changed to 2 mL/well RPMI+B27 minus insulin
supplemented with 5 µM IWR-1-Endo (STEMCELL). On day 7, the media was changed to 2 mL
RPMI+B27 with insulin. Differentiation was confirmed via visualization of morphology,
spontaneous contraction of cells, and imaging of the GFP tagged MYL7/MLC-2a. On day 9, the
media was changed to RPMI+B27 with insulin without glucose to select for cardiomyocytes. The
cardiomyocytes were then passaged at low density with 2 µM CHIR-99021 (Maas et al., 2021).
At approximately 75% confluency on passage 2, CHIR supplemented media was removed and

replaced with RPMI B-27 with insulin, and used for experiments on day 25–30 post
differentiation. For isotopic labeling, RPMI (Thermo Scientific) deficient in both L-lysine and L-
arginine was supplemented with B27 with insulin and heavy amino acids $^{13}C_6{}^{15}N_2$ L-Lysine-2HCl
and $^{13}C_6{}^{15}N_4$ L-Arginine-HCl (Thermo Scientific) at concentrations 0.219 mM and 1.149 mM,
respectively. 48 hours after CHIR removal, light media was replaced with this heavy media and
cells were labeled for 48 hours prior to harvest. 0.5 µM carfilzomib (Selleck) was added with
heavy media in the treatment group. Harvesting and ultracentrifugation proceeded as above with
the following exception. Due to the diffuse nature of pellets generated in the iPSC-derived
cardiomyocytes (iPSC-CM) control experiment, the MLA-50 (Beckman) rotor was switched to
the TLA-55 rotor after generation of pellet 5. Consistent force (g) of each spin was maintained
by increasing the smaller rotor's RPM on subsequent spins. This change was repeated during
the iPSC-CM treatment experiment. Proteins from each cellular fraction were digested and
analyzed with mass spectrometry as above.

Data Availability

912   Raw mass spectrometry data have been uploaded to ProteomeXchange under the accession
913   numbers PXD038054, PXD041386, PXD046669, PXD046670, and PXD046671. Source data are
914   provided with this paper.

Code Availability

916   Software code for the SPLAT pipeline is available on GitHub at https://github.com/lau-lab/splat,
917   https://github.com/ed-lau/riana and https://github.com/ed-lau/pytmt.
918

Author Contributions

920   J.C. and E.L. conceptualized the study. J.C., V.M., S.K.R. V.H., C.L., V.A., R.W.L., K.Z., J.P., J-
921   W.R. and M.P.L. performed experiments. J.C., V.M., and E.L. processed the data and interpreted
922   the results. J.C. and E.L. wrote software code and performed analysis. J.C., S.K.R., V.M., and
923   E.L. drafted and revised the manuscript. J-W.R., Z.V.W., M.P.L., and E.L. managed funding. All
924   authors read and approved the final version of the manuscript.

Acknowledgments

926   The authors thank Dr. Christopher Ebmeier (Director, Proteomics and Mass Spectrometry Core
927   Facility, University of Colorado Boulder) for his assistance with TMT-MS3 experiments. This work
928   was supported in part by NIH award K08-HL148540 to J-W.R., NIH awards R01-HL141278 and
929   R01-GM144456 to M.P.L. and NIH awards R35-GM146815 to E.L; and the University of
930   Colorado SOM Translational Research Scholars Program (TSRP) award to E.L..

References

Andrews, B., Murphy, A.E., Stofella, M., Maslen, S., Almeida-Souza, L., Skehel, J.M., Skene, N.G., Sobott, F., Frank, R.A.W., 2022. Multidimensional Dynamics of the Proteome in the Neurodegenerative and Aging Mammalian Brain. Mol. Cell. Proteomics 21, 100192. https://doi.org/10.1016/j.mcpro.2021.100192

Bresson, S., Shchepachev, V., Spanos, C., Turowski, T.W., Rappsilber, J., Tollervey, D., 2020. Stress-Induced Translation Inhibition through Rapid Displacement of Scanning Initiation Factors. Mol. Cell 80, 470-484.e8. https://doi.org/10.1016/j.molcel.2020.09.021

Burnum-Johnson, K.E., Conrads, T.P., Drake, R.R., Herr, A.E., Iyengar, R., Kelly, R.T., Lundberg, E., MacCoss, M.J., Naba, A., Nolan, G.P., Pevzner, P.A., Rodland, K.D., Sechi, S., Slavov, N., Spraggins, J.M., Van Eyk, J.E., Vidal, M., Vogel, C., Walt, D.R., Kelleher, N.L., 2022. New Views of Old Proteins: Clarifying the Enigmatic Proteome. Mol. Cell. Proteomics 21, 100254. https://doi.org/10.1016/j.mcpro.2022.100254

Burridge, P.W., Matsa, E., Shukla, P., Lin, Z.C., Churko, J.M., Ebert, A.D., Lan, F., Diecke, S., Huber, B., Mordwinkin, N.M., Plews, J.R., Abilez, O.J., Cui, B., Gold, J.D., Wu, J.C., 2014. Chemically defined generation of human cardiomyocytes. Nat. Methods 11, 855–860. https://doi.org/10.1038/nmeth.2999

Chartron, J.W., Hunt, K.C.L., Frydman, J., 2016. Cotranslational signal-independent SRP preloading during membrane targeting. Nature 536, 224–228. https://doi.org/10.1038/nature19309

Christopher, J.A., Stadler, C., Martin, C.E., Morgenstern, M., Pan, Y., Betsinger, C.N., Rattray, D.G., Mahdessian, D., Gingras, A.-C., Warscheid, B., Lehtiö, J., Cristea, I.M., Foster, L.J., Emili, A., Lilley, K.S., 2021. Subcellular proteomics. Nat. Rev. Methods Primer 1, 32. https://doi.org/10.1038/s43586-021-00029-y

Claydon, A.J., Beynon, R., 2012. Proteome dynamics: revisiting turnover with a global perspective. Mol. Cell. Proteomics MCP 11, 1551–1565. https://doi.org/10.1074/mcp.O112.022186

Cromm, P.M., Crews, C.M., 2017. The Proteasome in Modern Drug Discovery: Second Life of a Highly Valuable Drug Target. ACS Cent. Sci. 3, 830–838. https://doi.org/10.1021/acscentsci.7b00252

Crook, O.M., Breckels, L.M., Lilley, K.S., Kirk, P.D.W., Gatto, L., 2019. A Bioconductor workflow for the Bayesian analysis of spatial proteomics. F1000Research 8, 446. https://doi.org/10.12688/f1000research.18636.1

Crook, O.M., Davies, C.T.R., Breckels, L.M., Christopher, J.A., Gatto, L., Kirk, P.D.W., Lilley, K.S., 2022. Inferring differential subcellular localisation in comparative spatial proteomics using BANDLE. Nat. Commun. 13, 5948. https://doi.org/10.1038/s41467-022-33570-9

Crook, O.M., Mulvey, C.M., Kirk, P.D.W., Lilley, K.S., Gatto, L., 2018. A Bayesian mixture modelling approach for spatial proteomics. PLOS Comput. Biol. 14, e1006516. https://doi.org/10.1371/journal.pcbi.1006516

Demo, S.D., Kirk, C.J., Aujay, M.A., Buchholz, T.J., Dajee, M., Ho, M.N., Jiang, J., Laidig, G.J., Lewis, E.R., Parlati, F., Shenk, K.D., Smyth, M.S., Sun, C.M., Vallone, M.K., Woo, T.M., Molineaux, C.J., Bennett, M.K., 2007. Antitumor Activity of PR-171, a Novel Irreversible Inhibitor of the Proteasome. Cancer Res. 67, 6383–6391. https://doi.org/10.1158/0008-5472.CAN-06-4086

Dephoure, N., Gygi, S.P., 2012. Hyperplexing: A Method for Higher-Order Multiplexed Quantitative Proteomics Provides a Map of the Dynamic Response to Rapamycin in Yeast. Sci. Signal. 5. https://doi.org/10.1126/scisignal.2002548

Doherty, M.K., Hammond, D.E., Clague, M.J., Gaskell, S.J., Beynon, R.J., 2009. Turnover of
the human proteome: determination of protein intracellular stability by dynamic SILAC.
J. Proteome Res. 8, 104–112. https://doi.org/10.1021/pr800641v
Dostal, V., Wood, S.D., Thomas, C.T., Han, Y., Lau, E., Lam, M.P.Y., 2020. Proteomic
signatures of acute oxidative stress response to paraquat in the mouse heart. Sci. Rep.
10, 18440. https://doi.org/10.1038/s41598-020-75505-8
Eng, J.K., Hoopmann, M.R., Jahan, T.A., Egertson, J.D., Noble, W.S., MacCoss, M.J., 2015. A
deeper look into Comet--implementation and features. J. Am. Soc. Mass Spectrom. 26,
1865–1874. https://doi.org/10.1007/s13361-015-1179-x
Forghani, P., Rashid, A., Sun, F., Liu, R., Li, D., Lee, M.R., Hwang, H., Maxwell, J.T.,
Mandawat, A., Wu, R., Salaita, K., Xu, C., 2021. Carfilzomib Treatment Causes
Molecular and Functional Alterations of Human Induced Pluripotent Stem Cell–Derived
Cardiomyocytes. J. Am. Heart Assoc. 10, e022247.
https://doi.org/10.1161/JAHA.121.022247
Fridolfsson, H.N., Kawaraguchi, Y., Ali, S.S., Panneerselvam, M., Niesman, I.R., Finley, J.C.,
Kellerhals, S.E., Migita, M.Y., Okada, H., Moreno, A.L., Jennings, M., Kidd, M.W.,
Bonds, J.A., Balijepalli, R.C., Ross, R.S., Patel, P.M., Miyanohara, A., Chen, Q.,
Lesnefsky, E.J., Head, B.P., Roth, D.M., Insel, P.A., Patel, H.H., 2012. Mitochondria-
localized caveolin in adaptation to cellular stress and injury. FASEB J. Off. Publ. Fed.
Am. Soc. Exp. Biol. 26, 4637–4649. https://doi.org/10.1096/fj.12-215798
Gatto, L., Breckels, L.M., Wieczorek, S., Burger, T., Lilley, K.S., 2014. Mass-spectrometry-
based spatial proteomics data analysis using pRoloc and pRolocdata. Bioinformatics
30, 1322–1324. https://doi.org/10.1093/bioinformatics/btu013
Geladaki, A., Kočevar Britovšek, N., Breckels, L.M., Smith, T.S., Vennard, O.L., Mulvey, C.M.,
Crook, O.M., Gatto, L., Lilley, K.S., 2019. Combining LOPIT with differential
ultracentrifugation for high-resolution spatial proteomics. Nat. Commun. 10, 331.
https://doi.org/10.1038/s41467-018-08191-w
Glembotski, C.C., 2007. Endoplasmic Reticulum Stress in the Heart. Circ. Res. 101, 975–984.
https://doi.org/10.1161/CIRCRESAHA.107.161273
Gonen, N., Meller, A., Sabath, N., Shalgi, R., 2019. Amino Acid Biosynthesis Regulation during
Endoplasmic Reticulum Stress Is Coupled to Protein Expression Demands. iScience 19,
204–213. https://doi.org/10.1016/j.isci.2019.07.022
Guan, B.-J., van Hoef, V., Jobava, R., Elroy-Stein, O., Valasek, L.S., Cargnello, M., Gao, X.-H.,
Krokowski, D., Merrick, W.C., Kimball, S.R., Komar, A.A., Koromilas, A.E., Wynshaw-
Boris, A., Topisirovic, I., Larsson, O., Hatzoglou, M., 2017. A Unique ISR Program
Determines Cellular Responses to Chronic Stress. Mol. Cell 68, 885-900.e6.
https://doi.org/10.1016/j.molcel.2017.11.007
Hammond, D.E., Simpson, D.M., Franco, C., Wright Muelas, M., Waters, J., Ludwig, R.W.,
Prescott, M.C., Hurst, J.L., Beynon, R.J., Lau, E., 2022. Harmonizing Labeling and
Analytical Strategies to Obtain Protein Turnover Rates in Intact Adult Animals. Mol. Cell.
Proteomics 21, 100252. https://doi.org/10.1016/j.mcpro.2022.100252
Hanafusa, K., Wada, I., Hosokawa, N., 2019. SDF2-like protein 1 (SDF2L1) regulates the
endoplasmic reticulum localization and chaperone activity of ERdj3 protein. J. Biol.
Chem. 294, 19335–19348. https://doi.org/10.1074/jbc.RA119.009603
Harding, H.P., Zhang, Y., Zeng, H., Novoa, I., Lu, P.D., Calfon, M., Sadri, N., Yun, C., Popko,
B., Paules, R., Stojdl, D.F., Bell, J.C., Hettmann, T., Leiden, J.M., Ron, D., 2003. An
Integrated Stress Response Regulates Amino Acid Metabolism and Resistance to
Oxidative Stress. Mol. Cell 11, 619–633. https://doi.org/10.1016/S1097-2765(03)00105-
9

Hetz, C., Zhang, K., Kaufman, R.J., 2020. Mechanisms, regulation and functions of the
unfolded protein response. Nat. Rev. Mol. Cell Biol. 21, 421–438.
https://doi.org/10.1038/s41580-020-0250-z
Higuchi-Sanabria, R., Shen, K., Kelet, N., Frankino, P.A., Durieux, J., Bar-Ziv, R., Sing, C.N.,
Garcia, E.J., Homentcovschi, S., Sanchez, M., Wu, R., Tronnes, S.U., Joe, L., Webster,
B., Ahilon-Jeronimo, A., Monshietehadi, S., Dallarda, S., Pender, C., Pon, L.A., Zoncu,
R., Dillin, A., 2020. Lysosomal recycling of amino acids affects ER quality control. Sci.
Adv. 6, eaaz9805. https://doi.org/10.1126/sciadv.aaz9805
Hulstaert, N., Shofstahl, J., Sachsenberg, T., Walzer, M., Barsnes, H., Martens, L., Perez-
Riverol, Y., 2020. ThermoRawFileParser: Modular, Scalable, and Cross-Platform RAW
File Conversion. J. Proteome Res. 19, 537–542.
https://doi.org/10.1021/acs.jproteome.9b00328
IBM Watson Health, 2023. Micromedex (Electronic Version).
Jan, C.H., Williams, C.C., Weissman, J.S., 2014. Principles of ER cotranslational translocation
revealed by proximity-specific ribosome profiling. Science 346, 1257521.
https://doi.org/10.1126/science.1257521
Kennedy, M.A., Hofstadter, W.A., Cristea, I.M., 2020. TRANSPIRE: A Computational Pipeline to
Elucidate Intracellular Protein Movements from Spatial Proteomics Data Sets. J. Am.
Soc. Mass Spectrom. 31, 1422–1439. https://doi.org/10.1021/jasms.0c00033
Lakkaraju, A.K.K., Mary, C., Scherrer, A., Johnson, A.E., Strub, K., 2008. SRP Keeps
Polypeptides Translocation-Competent by Slowing Translation to Match Limiting ER-
Targeting Sites. Cell 133, 440–451. https://doi.org/10.1016/j.cell.2008.02.049
Lam, M.P.Y., Wang, D., Lau, E., Liem, D.A., Kim, A.K., Ng, D.C.M., Liang, X., Bleakley, B.J., Liu,
C., Tabaraki, J.D., Cadeiras, M., Wang, Y., Deng, M.C., Ping, P., 2014. Protein kinetic
signatures of the remodeling heart following isoproterenol stimulation. J. Clin. Invest.
124, 1734–1744. https://doi.org/10.1172/JCI73787
Larance, M., Ahmad, Y., Kirkwood, K.J., Ly, T., Lamond, A.I., 2013. Global Subcellular
Characterization of Protein Degradation Using Quantitative Proteomics. Mol. Cell.
Proteomics 12, 638–650. https://doi.org/10.1074/mcp.M112.024547
Lau, E., Cao, Q., Lam, M.P.Y., Wang, J., Ng, D.C.M., Bleakley, B.J., Lee, J.M., Liem, D.A.,
Wang, D., Hermjakob, H., Ping, P., 2018. Integrated omics dissection of proteome
dynamics during cardiac remodeling. Nat. Commun. 9, 120.
https://doi.org/10.1038/s41467-017-02467-3
Lee, J.W., Koeppen, M., Seo, S.-W., Bowser, J.L., Yuan, X., Li, J., Sibilia, M., Ambardekar,
A.V., Zhang, X., Eckle, T., Yoo, S.-H., Eltzschig, H.K., 2020. Transcription-independent
Induction of ERBB1 through Hypoxia-inducible Factor 2A Provides Cardioprotection
during Ischemia and Reperfusion. Anesthesiology 132, 763–780.
https://doi.org/10.1097/ALN.0000000000003037
Lemberg, M.K., Strisovsky, K., 2021. Maintenance of organellar protein homeostasis by ER-
associated degradation and related mechanisms. Mol. Cell 81, 2507–2519.
https://doi.org/10.1016/j.molcel.2021.05.004
Liu, C., Li, X., Li, C., Zhang, Z., Gao, X., Jia, Z., Chen, H., Jia, Q., Zhao, X., Liu, J., Liu, B., Xu,
Z., Tian, Y., He, K., 2018. SLC3A2 is a novel endoplasmic reticulum stress-related
signaling protein that regulates the unfolded protein response and apoptosis. PloS One
13, e0208993. https://doi.org/10.1371/journal.pone.0208993
Liu, C.-L., Li, X., Hu, G.-L., Li, R.-J., He, Y.-Y., Zhong, W., Li, S., He, K.-L., Wang, L.-L., 2012.
Salubrinal protects against tunicamycin and hypoxia induced cardiomyocyte apoptosis
via the PERK-eIF2α signaling pathway. J. Geriatr. Cardiol. JGC 9, 258–268.
https://doi.org/10.3724/SP.J.1263.2012.02292

Maas, R.G.C., Lee, S., Harakalova, M., Snijders Blok, C.J.B., Goodyer, W.R., Hjortnaes, J.,
Doevendans, P.A.F.M., Van Laake, L.W., van der Velden, J., Asselbergs, F.W., Wu, J.C.,
Sluijter, J.P.G., Wu, S.M., Buikema, J.W., 2021. Massive expansion and
cryopreservation of functional human induced pluripotent stem cell-derived
cardiomyocytes. STAR Protoc. 2, 100334. https://doi.org/10.1016/j.xpro.2021.100334
Mårtensson, C.U., Priesnitz, C., Song, J., Ellenrieder, L., Doan, K.N., Boos, F., Floerchinger, A.,
Zufall, N., Oeljeklaus, S., Warscheid, B., Becker, T., 2019. Mitochondrial protein
translocation-associated degradation. Nature 569, 679–683.
https://doi.org/10.1038/s41586-019-1227-y
Martin, T.G., Myers, V.D., Dubey, P., Dubey, S., Perez, E., Moravec, C.S., Willis, M.S., Feldman,
A.M., Kirk, J.A., 2021a. Cardiomyocyte contractile impairment in heart failure results
from reduced BAG3-mediated sarcomeric protein turnover. Nat. Commun. 12, 2942.
https://doi.org/10.1038/s41467-021-23272-z
Martin, T.G., Tawfik, S., Moravec, C.S., Pak, T.R., Kirk, J.A., 2021b. BAG3 expression and
sarcomere localization in the human heart are linked to HSF-1 and are differentially
affected by sex and disease. Am. J. Physiol.-Heart Circ. Physiol. 320, H2339–H2350.
https://doi.org/10.1152/ajpheart.00419.2020
Martínez-Martínez, E., Ibarrola, J., Fernández-Celis, A., Santamaria, E., Fernández-Irigoyen, J.,
Rossignol, P., Jaisser, F., López-Andrés, N., 2017. Differential Proteomics Identifies
Reticulocalbin-3 as a Novel Negative Mediator of Collagen Production in Human
Cardiac Fibroblasts. Sci. Rep. 7, 12192. https://doi.org/10.1038/s41598-017-12305-7
McDowell, G.S., Gaun, A., Steen, H., 2013. iFASP: combining isobaric mass tagging with filter-
aided sample preparation. J. Proteome Res. 12, 3809–3812.
https://doi.org/10.1021/pr400032m
Meiners, S., Heyken, D., Weller, A., Ludwig, A., Stangl, K., Kloetzel, P.-M., Krüger, E., 2003.
Inhibition of Proteasome Activity Induces Concerted Expression of Proteasome Genes
and de Novo Formation of Mammalian Proteasomes. J. Biol. Chem. 278, 21517–21525.
https://doi.org/10.1074/jbc.M301032200
Mulvey, C.M., Breckels, L.M., Crook, O.M., Sanders, D.J., Ribeiro, A.L.R., Geladaki, A.,
Christoforou, A., Britovšek, N.K., Hurrell, T., Deery, M.J., Gatto, L., Smith, A.M., Lilley,
K.S., 2021. Spatiotemporal proteomic profiling of the pro-inflammatory response to
lipopolysaccharide in the THP-1 human leukaemia cell line. Nat. Commun. 12, 5773.
https://doi.org/10.1038/s41467-021-26000-9
Orre, L.M., Vesterlund, M., Pan, Y., Arslan, T., Zhu, Y., Fernandez Woodbridge, A., Frings, O.,
Fredlund, E., Lehtiö, J., 2019. SubCellBarCode: Proteome-wide Mapping of Protein
Localization and Relocalization. Mol. Cell 73, 166-182.e7.
https://doi.org/10.1016/j.molcel.2018.11.035
Pakos-Zebrucka, K., Koryga, I., Mnich, K., Ljujic, M., Samali, A., Gorman, A.M., 2016. The
integrated stress response. EMBO Rep. 17, 1374–1395.
https://doi.org/10.15252/embr.201642195
Ren, J., Bi, Y., Sowers, J.R., Hetz, C., Zhang, Y., 2021. Endoplasmic reticulum stress and
unfolded protein response in cardiovascular diseases. Nat. Rev. Cardiol. 18, 499–521.
https://doi.org/10.1038/s41569-021-00511-w
Sayers, E.W., Bolton, E.E., Brister, J.R., Canese, K., Chan, J., Comeau, D.C., Connor, R., Funk,
K., Kelly, C., Kim, S., Madej, T., Marchler-Bauer, A., Lanczycki, C., Lathrop, S., Lu, Z.,
Thibaud-Nissen, F., Murphy, T., Phan, L., Skripchenko, Y., Tse, T., Wang, J., Williams,
R., Trawick, B.W., Pruitt, K.D., Sherry, S.T., 2022. Database resources of the national
center for biotechnology information. Nucleic Acids Res. 50, D20–D26.
https://doi.org/10.1093/nar/gkab1112

Schwanhäusser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W., Selbach, M., 2011. Global quantification of mammalian gene expression control. Nature 473, 337–342. https://doi.org/10.1038/nature10098

Shin, J.J.H., Crook, O.M., Borgeaud, A.C., Cattin-Ortolá, J., Peak-Chew, S.Y., Breckels, L.M., Gillingham, A.K., Chadwick, J., Lilley, K.S., Munro, S., 2020. Spatial proteomics defines the content of trafficking vesicles captured by golgin tethers. Nat. Commun. 11, 5987. https://doi.org/10.1038/s41467-020-19840-4

Tan, X., Lambert, P.F., Rapraeger, A.C., Anderson, R.A., 2016. Stress-Induced EGFR Trafficking: Mechanisms, Functions, and Therapeutic Implications. Trends Cell Biol. 26, 352–366. https://doi.org/10.1016/j.tcb.2015.12.006

Tan, X., Thapa, N., Sun, Y., Anderson, R.A., 2015. A Kinase-Independent Role for EGF Receptor in Autophagy Initiation. Cell 160, 145–160. https://doi.org/10.1016/j.cell.2014.12.006

The, M., MacCoss, M.J., Noble, W.S., Käll, L., 2016. Fast and Accurate Protein False Discovery Rates on Large-Scale Proteomics Data Sets with Percolator 3.0. J. Am. Soc. Mass Spectrom. 27, 1719–1727. https://doi.org/10.1007/s13361-016-1460-7

Toro, R., Pérez-Serra, A., Mangas, A., Campuzano, O., Sarquella-Brugada, G., Quezada-Feijoo, M., Ramos, M., Alcalá, M., Carrera, E., García-Padilla, C., Franco, D., Bonet, F., 2022. miR-16-5p Suppression Protects Human Cardiomyocytes against Endoplasmic Reticulum and Oxidative Stress-Induced Injury. Int. J. Mol. Sci. 23, 1036. https://doi.org/10.3390/ijms23031036

Tsai, P.-L., Cameron, C.J.F., Forni, M.F., Wasko, R.R., Naughton, B.S., Horsley, V., Gerstein, M.B., Schlieker, C., 2022. Dynamic quality control machinery that operates across compartmental borders mediates the degradation of mammalian nuclear membrane proteins. Cell Rep. 41, 111675. https://doi.org/10.1016/j.celrep.2022.111675

Ustrell, V., Hoffman, L., Pratt, G., Rechsteiner, M., 2002. PA200, a nuclear proteasome activator involved in DNA repair. EMBO J. 21, 3516–3525. https://doi.org/10.1093/emboj/cdf333

Villanueva, E., Smith, T., Pizzinga, M., Elzek, M., Queiroz, R.M.L., Harvey, R.F., Breckels, L.M., Crook, O.M., Monti, M., Dezi, V., Willis, A.E., Lilley, K.S., 2022. A system-wide quantitative map of RNA and protein subcellular localisation dynamics (preprint). Cell Biology. https://doi.org/10.1101/2022.01.24.477541

Vonk, L.A., Doulabi, B.Z., Huang, C.-L., Helder, M.N., Everts, V., Bank, R.A., 2010. Endoplasmic reticulum stress inhibits collagen synthesis independent of collagen-modifying enzymes in different chondrocyte populations and dermal fibroblasts. Biochem. Cell Biol. Biochim. Biol. Cell. 88, 539–552. https://doi.org/10.1139/o09-174

Xie, Y., Varshavsky, A., 2001. RPN4 is a ligand, substrate, and transcriptional regulator of the 26S proteasome: a negative feedback circuit. Proc. Natl. Acad. Sci. U. S. A. 98, 3056–3061. https://doi.org/10.1073/pnas.071022298

Xu, H., Ju, D., Jarois, T., Xie, Y., 2008. Diminished feedback regulation of proteasome expression and resistance to proteasome inhibitors in breast cancer cells. Breast Cancer Res. Treat. 107, 267–274. https://doi.org/10.1007/s10549-007-9553-4