Endophytic fungi related to the ash dieback causal agent encode signatures of pathogenicity on European ash

Maryam Rafiqi[1,2]*, Chatchai Kosawang[3], Jessica A. Peers[4], Lukas Jelonek[5], Hélène Yvanne[4], Mark McMullan[4] and Lene R. Nielsen[3]*

[1] Jodrell Laboratory, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK;

[2] Plant Pathology, Agrobiosciences, Mohammed VI Polytechnic University, Ben Guerir, Morocco

[3] Department of Geosciences and Natural Resource Management, University of Copenhagen, Rolighedsvej 23, 1958, Frederiksberg C, Denmark

[4] The Earlham Institute, Norwich Research Park, Norwich, NR4 7UZ, UK

[5] Bioinformatics and Systems Biology, Justus Liebig University Giessen, Giessen, Germany

Abstract

Tree diseases constitute a significant threat to biodiversity worldwide. Pathogen discovery in natural habitats is of vital importance to understanding current and future threats and prioritising efforts towards developing disease management strategies. Ash dieback is a fungal disease of major conservational concern that is infecting common ash trees, Fraxinus excelsior, in Europe. The disease is caused by a non-native fungal pathogen, Hymenoscyphus fraxineus. Other dieback causing-species have not previously been identified in the genus Hymenoscyphus. Here, we discover the pathogenicity potential of two newly identified related species of Asian origin, H. koreanus and H. occultus, and one Europe-native related species, H. albidus. We sequence the genomes of all three Hymenoscyphus species and compare them to that of H. fraxineus. Phylogenetic analysis of core eukaryotic genes identified H. albidus and H. koreanus as sister species, whilst H. occultus diverged prior these and H. fraxineus. All four Hymenoscyphus genomes are of comparable sizes (55-62 Mbp) and GC contents (42–44%) and encode for polymorphic secretomes. Surprisingly, 1,133 predicted secreted proteins are shared between the ash dieback pathogen H. fraxineus and the three related Hymenoscyphus endophytes. Amongst shared secreted proteins are cell death-inducing effector candidates, such as necrosis, and ethylene-inducing peptide 1-like proteins, NLPs, that are upregulated during in planta growth of all Hymenoscyphus species. Indeed, pathogenicity tests showed that all four related Hymenoscyphus species develop pathogenic growth on European ash stems, with native H. albidus being the least virulent. Our results identify the threat Hymenoscypohus species pose to the survival of European ash trees, and highlight the importance of promoting pathogen surveillance in environmental landscapes. Identifying new pathogens and including them in the screening for durable immunity of common ash trees is key to the long-term survival of ash.

Main

Fungal diseases have huge economic and environmental impacts worldwide. Fungal diseases infecting crops are of high research priority because they directly affect food security and the global economy. In contrast, fungal diseases of non-crop plants in natural habitats and urban areas are far less studied and, consequently, less well understood. Several plant diseases affecting environmental landscapes have caused huge permanent damage to biodiversity whilst their pathogenesis was still unknown (Fisher *et al.*, 2012; Rafiqi, 2018). One such disease is ash dieback.

The ash dieback pandemic was first observed in 2002 in Poland but has quickly spread to much of Europe invading ash trees (*Fraxinus excelsior*) in natural and urban habitats (Drenkhan *et al.*, 2017). The ash dieback pandemic is expected to have huge economic and environmental consequences. For example, in the UK alone, the disease is expected to cost around £15 billion (Hill *et al.*, 2019). The substantial loss of *F. excelsior* is likely to negatively affect carbon sequestration and threaten around 1000 other species that are associated with ash, including birds, mammals, invertebrates, vascular plants, lichens, and fungi (Mitchell, 2014). It is anticipated that ash dieback will change the European landscape forever.

The ash dieback disease, also called Chalara dieback of ash, is caused by the ascomycete pathogen *Hymenoscyphus fraxineus* (formally known under asexual, *Chalara fraxinea*, and sexual, *Hymenoscyphus pseudoalbidus*), a commensal endophyte of Asian origin that causes no or minor symptoms when colonising leaves of Asian ash host species (Zheng and Zhuang, 2013; Zhao *et al.*, 2013a; Drenkhan *et al.*, 2017), yet is highly pathogenic on European ash. The genome of *H. fraxineus* is about 62 Mbp and encodes a large and diversified set of host-interacting genes (McMullan *et al.*, 2018). *H. fraxineus* European population was founded by two divergent haploid individuals and has a bottlenecked genetic diversity when compared to the native Asian population of the endophyte (McMullan *et al.*, 2018; Gross, Hosoya and Queloz, 2014). Subsequent introduction of new individuals is expected to increase the pathogen's adaptive potential in Europe, which in turn would further jeopardise the survival of European ash trees. What triggers *H. fraxineus* to switch from a commensal endophyte on Asian ash to a virulent pathogen on European ash is as yet unknown.

It is widely accepted that all plants, including lichens, algae, and nonvascular plants, harbour microorganisms, called endophytes, living within them (Bacon and White, 2000). Typically, endophytes live with no apparent symptoms on plants. However, depending on environmental conditions and interactions with the host, some endophytes have visible effects on their host's immunity and physiology. Therefore, endophytes can be pathogenic or nonpathogenic (Hardoim *et al.*, 2015; Brader *et al.*, 2017; Bacon and Hinton, 1996). Research in this area is scarce, although analysis of the endospheric microbiome of many plant species is increasingly detecting, through amplicon sequencing, potential pathogens that did not cause noticeable symptoms in the host at the sampling time (Pereira *et*

al., 2019; Manzotti *et al.*, 2020). Whether the pathogenicity of these endophytes could be triggered and

how is currently unresolved. Specific ecological drivers, competition for nutrients, as well as host

factors and interactions with other endophytes could all underlie the suppression of disease symptoms

despite pathogen occurrence. Pathogenic fungi infecting plants in agricultural or environmental

landscapes are not clustered in a particular taxonomic category that separates them from harmless

endophytes. Instead, some endophytic species possess pathogenic and nonpathogenic trophic states. For

example, *Sydowia polyspora* is both a foliar commensal endophyte and a preemergent seed pathogen

in *Pinus ponderosa* (Ridout and Newcombe, 2017). Many fungal species include both pathogenic and

non-pathogenic members. For example, *Fusarium oxysporum*, a ubiquitous soil-borne fungus, is

classified in more than 150 forms or *formae speciales* of pathogenic and non-pathogenic strains, each

one with the ability to infect one or a group of plant species (Nirmaladevi *et al.*, 2016). Pathogenic *F.*

oxysporum species often carry dispensable chromosomes that are not required for non-pathogenic

growth but are virulence-associated and carry genes coding for specific host-interacting proteins and

toxins, also referred to as effectors. Horizontal transfer of these chromosomes from pathogenic to a non-

pathogenic recipient lineage of *F. oxysporum* renders the latter pathogenic on the respective host (van

Dam *et al.*, 2017). Such dispensable chromosomes have, however, not yet been identified in *H.*

fraxineus (McMullan et al., 2018).

Forecasting fungal pathogen invasions in environmental landscapes is challenging and requires cross-

disciplinary experimental approaches because of a myriad of variables and the complexity of

interactions between the invading species and biological and physical characteristics of the recipient

ecosystem (Santini *et al.*, 2013; Alpert, Bone and Holzapfel, 2000; Hayes and Barry, 2008). The main

factors include pathogen virulence and residence time, host susceptibility and abundance, as well as

climatic factors (Hayes and Barry, 2008).

Hymenoscyphus is a large genus of the family of Helotiaceae and includes over 150 species (Kirk *et al.*,

2008), with new species increasingly being discovered (Zheng and Zhuang, 2013; Çetinkaya and Uzun,

2020; Gross and Han, 2015). *Hymenoscyphus* species were long classified as saprotrophic decomposers

and non-pathogenic endophytes up until the emergence of *H. fraxineus* as a successful invasive

pathogen of ash trees in Europe. Here, we interrogate if other species of the genus *Hymenoscyphus*

could be pathogenic and cause disease on European ash. We address this question by studying three

Hymenoscyphus species closely related to *H. fraxineus*: *H. albidus*, isolated from European ash growing

in Wales (UK), *H. koreanus,* and *H. occultus*; two newly-discovered species of Asian origin and not

(yet) introduced to Europe; they have been isolated by Gross and Han (2015) from petioles of Korean

ash (*F. chinensis* subsp. *rhynchophylla*) (Gross and Han, 2015; Kosawang *et al.*, 2020).

The fungal secretome refers to the set of functionally diverse families of secreted proteins, many of

which are involved in a range of diverse biological processes that directly affect the outcome of plant-

fungi interactions. Examples include cell wall degrading enzymes, extracellular proteinases, toxins, and
effector proteins that are upregulated during *in planta* growth of the pathogen and implicated in the
suppression of host immune defences (Vincent, Rafiqi and Job, 2019). Whilst identifying and analysing
predicted secretory proteins involved in disease development can't help accurately predict pathogen
invasions, it increasingly contributes to our understanding of pathogenicity and host responses. Here,
we sequence and compare the genomes of three *Hymenoscyphus* species related to *H. fraxineus*. Since
these fungal species can vary in their secretome predominantly by gene gains and losses or by the rapid
evolution of secreted proteins, we compare the full set of proteins predicted to be secreted by these four
fungal species, analyse putative effectors that could be correlated with pathogenicity and, ultimately,
we test the pathogenicity of these *Hymecoscyphus* species on European ash and discuss the threat they
further pose to the survival of European ash trees. The results show that in addition to the primary
causative agent, *H. fraxineus*, the ash dieback disease can be caused by other member species in the
Hymenoscyphus genus, which highlight the importance of promoting pathogen surveillance in
environmental landscapes, identifying current and future threats to biodiversity, locally and at a global
level.

Results and discussion

1- Whole genome sequencing of three endophytic fungal species related to *H. fraxineus*

Genome sequences of three *Hymenoscyphus* species, *H. albidus* (76x coverage), *H. koreanus* (114x
coverage) and *H. occultus* (102x coverage), were generated and compared to the genome of related and
previously published *H. fraxineus* (McMullan *et al.*, 2018). We assessed the completeness of the three
Hymenoscyphus genome assemblies, using KAT and BUSCO. The *k-mer* distribution of such
assemblies is compatible with the distribution of a complete one, and the estimated genome size is close
to the actual size of the genome assembly. Approximately 96% of single copy ortholog genes from the
Sodariomycetes odb9 were completely assembled in all three genomes (Table S1). Among a total of
3725 tested BUSCO genes, only 61, 73, and 78 single-copy orthologs were missing in the *H. albidus*,
H. koreanus, and *H. occultus* genomes, respectively. These results suggest that draft genome sequences
generated in this study were relatively complete in terms of both assembly completeness and gene
content. Genome features of *H. albidus*, *H. koreanus,* and *H. occultus* are detailed in Table 1.

All *Hymenoscyphus* genomes assembled here are of similar size (55-62 Mbp) and GC content (42–
44%) and are comparable to *H. fraxineus*. The total number of coding sequences (CDSs) varied among
the three genomes: 13,657 in *H. albidus*, 15,042 in *H. koreanus* and 14,686 in *H. occultus* (table 1). Of

these CDSs, 7,791 (57.0%) in *H. albidus*, 8,313 (55.2%) in *H. koreanus*, and 8,180 (55.6%) in *H. occultus* harboured recognised protein signatures in Pfam/CDD databases. When compared to other *Hymensoscyphus* species, the predicted number of CDSs in the three genome assemblies were slightly greater than that of *H. fraxineus* (11,097 CDSs) (McMullan *et al.*, 2018), but were in the same range as that of *H. varicosporoides* PMI_453 (14,929 CDSs; https://mycocosm.jgi.doe.gov/Hymvar1/Hymvar1.home.html).

OrthoMCL clustering of CDSs of all four *Hymenoscyphus* species (*H. fraxineus* and the three other related species addressed here) assigned 49,045 CDSs (90% of total CDSs) to 12,426 orthogroups (OGs), 8,665 (69.7%) of which contained CDSs from all four *Hymenoscyphus* species. Transporter, synthase, and cytochrome 450 CDSs were among the largest OGs. We found a total of 71 CDSs, representing 9 OGs specific to *H. fraxineus* (2 CDSs, 2 OGs), *H. albidus* (4 CDSs, 1 OG), and *H. occultus* (63 CDSs, 6 OGs), respectively. No CDSs specific to *H. koreanus* were found. The fact that only a small fraction, less than 1%, of predicted CDSs in all four *Hymenoscyphus* species are species-specific emphasises that these closely-related species likely share many common adaptations to similar environmental conditions. Phylogenetic analysis of core eukaryotic genes (CEGs) showed *H. albidus* and *H. koreanus* as sister species, whilst *H. occultus* was a more distant relative in the clade (Fig. 1). In previous studies, *H. fraxineus* and *H. albidus* were initially thought to be sister species, mostly due to their similar morphologies and high level of synteny (Elfstrand *et al.*, 2021). In the present study we gain additional resolution with the inclusion of the newly discovered related species *H. koreanus* and *H. occultus*. Our expectation is, that with an increase in attention to this genus, additional species will be discovered and further our understanding of the biology of this group. Our finding supports the phylogenetic relatedness of *Hymenoscyphus* species previously described by (Gross and Han, 2015).

Table 1 Genome assembly statistics for *Hymenoscyphus albidus*, *H. koreanus* F52847-4 and *H. occultus* F52847-22 isolates

Characteristic	*H. albidus*	*H. koreanus* F52847-4	*H. occultus* F52847-22
Genome size (Mb)	55.7	57.4	61.6
No. of scaffolds	2002	1517	728
N50/L50 scaffold length (Kb)	265/53.5	225/76.2	115/154.7
Coverage	76x	114x	102x
%GC	44.3	44.1	42.1
No. of CDSs	13657	15042	14686
Average gene length	1520	1576	1572

2- Comparative analysis of the *in silico* secretomes of *H. albidus*, *H. koreanus*, *H. occultus* and *H. fraxineus*

Common gene ontologies

We mined the genomes of *H. albidus*, *H. koreanus* and *H. occultus* for predicted secreted proteins, using a pipeline described in materials and methods. All three fungi code for secretomes of similar sizes: 1446, 1583 and 1676 predicted secreted proteins, respectively (Table S2). Secretomes of these three species were compared to that of related pathogenic *H. fraxineus*. Whilst the four predicted secretomes are highly heterogenous, strikingly a high number of proteins (1133), were shared between the ash dieback pathogen *H. fraxineus* and the three related *Hymenoscyphus* endophytes. 594 proteins were shared with *H. koreanus*, 567 proteins with *H. albidus* and 411 with *H. occultus*, with an overlap of 324 proteins shared amongst all four secretomes (Fig 2). Genes in common between pathogenic species that are involved in plant interaction shed light on a foundation of pathogenicity whereas novel genes may play a role in niche adaptation or disease development. Regardless of their phylogenetic distance, the fact that all four *Hymenoscyphus* species share a significant number of predicted secreted proteins suggests possible common infection strategies among species, which is in agreement with the observed similarity between *H. fraxineus* and *H. albidus* in developing necrotrophic growth on ash leaves (Hietala *et al.*, 2022).

Polymorphism and positive selection

We analysed genetic diversity within coding sequences in all four *Hymenoscyphus* species. Genes coding for secreted proteins were associated with a SNP density that is significantly higher on average than that observed in other genes (Fig 3A). In addition, SNPs observed in secretome genes were associated with an increase in the rate of non-synonymous nucleic acid changes to the rate of synonymous nucleic acid changes (dN/dS) (Fig 3B). The polymorphism observed here in conjunction with high adaptive diversity in secretome genes among all four *Hymenoscyphus* species suggest genes coding for host-interacting proteins, including but not limited to effectors, undergo a faster rate of adaptive evolution than non-secreted genes and may play a key role in the infection process and pathogenesis on ash trees.

Large repertoires of CAZymes

To identify carbohydrate-active enzyme (CAZyme) gene content, protein sequences from the three *Hymensocyphus* species and pathogenic *H. fraxineus* were submitted for CAZymes prediction, using at least two integrated automated dbCAN2 tools (Zhang *et al.*, 2018). Secretomes of all four species encode comparably high numbers (292-307) of CAZymes (Fig 4). Glycoside hydrolases (GH) appear

to be the most highly represented (around 40-50%) of all CAZymes modules, followed by auxiliary

activities (AA), carbohydrate esterases (CE) and carbohydrate-binding modules (CBM), respectively.

CAZymes are plant cell wall degrading enzymes (PCWDs) that play a critical role in disintegrating

components of the host plant cell wall, which provides pathogens with a carbon source and access to

other nutrients in the host apoplast. Plant pathogenic fungi tend to secrete more CAZymes than

saprophytes and symbionts. Within plant pathogens, necrotrophic and hemibiotrophic fungi contain

more CAZymes than biotrophic fungi (Zhao *et al.*, 2013c). The high number of CAZymes in the

predicted secretomes of *Hymenoscyphus* species could, therefore, be attributed to their trophic mode

(Stenlid *et al.*, 2017). Cytological details of how these fungi grow *in planta* are yet unavailable, although

some evidence for the formation of biotrophic fungal structures was observed in ash leaf cells inoculated

with the ash dieback pathogen, *H. fraxineus,* (Mansfield, Galambos and Saville, 2018), suggesting a

hemibiotrophic lifestyle, though details of how long the initial biotrophic phase takes before the

pathogen switches into a necrotophic growth phase has not yet been identified (Mansfield, Brown and

Papp-Rupar, 2019). Large sets of CAZymes are more frequent in necrotrophic pathogens (Zhao *et al.*,

2013b). In agreement with this, a new classification of fungi based exclusively on genome-derived

analysis of CAZymes has grouped *H. fraxineus* with polymertroph pathogens with a narrow host range,

corresponding to necrotrophs (Hane *et al.*, 2020). Indeed a recent comparative study by (Hietala *et al.*,

2022) showed that both *H. fraxineus* and *H. albidus* develop similar necrotophic growth on leaves of

European ash. Detailed cytological studies are further needed to understand the nature of the life style

of *Hymenoscyphus* species on living ash trees and how CAZymes are involved.

Shared Cell-death inducing effectors

Amongst predicted secreted proteins shared between pathogenic *H. fraxineus* and the three additional

Hymenoscyphus species studied here are plant cell death inducing proteins. These include proteins with

predicted nuclease activity. Typically, each *Hymenoscyphus* species is predicted to secrete three to four

S1/P1 nucleases, one Ribonuclease 2-5A and three close homologues of T2 RNAse family protein.

Secreted nuclease effectors are increasingly identified in the secretomes of fungal pathogens (Kumakura

et al., 2021; Rafiqi *et al.*, 2021; Pennington *et al.*, 2019b; Rafiqi *et al.*, 2022). They might either act

intracellularly as a cytotoxin by scavenging host nucleic acids (Balabanova *et al.*, 2012; Pennington *et

al.*, 2019a) or extracellularly to interact with the damage-associated molecular pattern extracellular

RNA and DNA (Kumakura *et al.*, 2021; Park *et al.*, 2019; Widmer, 2018). Other cell-death inducing

effector candidates in the secretomes of this *Hymenoscyphus* group include homologues of necrosis-

inducing toxins, called Nep1-like proteins (NLPs) (Fellbrich *et al.*, 2002; Bailey, 1995). NLPs cause

necrosis in dicotyledonous plants and are suggested to function as virulence factors that accelerate

disease development and pathogen growth in the host (Feng *et al.*, 2014). Reverse transcription

quantitative real-time PCR (RT-qPCR) of a selected set of genes coding for cell death inducing effector
candidates showed upregulation of these genes during *in planta* growth of all *Hymenoscyphus* species
(Fig 5). Predicted S1/P1 nuclease genes were induced 3dpi and 6dpi in the three *Hymenoscyphus* species
as well as in pathogenic *H. fraxineus* (Fig 5A). NLP homologues are 15- and 30-fold induced in both
H. albidus and *H. fraxineus*, respectively (Fig 5C and 5F), suggesting a potential involvement in
virulence. It is particularly surprising to find that *H. albidus*, which has long been regarded as a
nonpathogenic endophyte in European ash trees, is expressing PCD-inducing toxins. This is in line with
recent increasing number of studies showing that *H. albidus* and *H. fraxineus*, the ash dieback primary
causal agent, both develop similar necrotic growth on leaves of common ash trees. Although it causes
shorter lesions on ash stems than those caused by *H. fraxineus*, *H. albidus* has been consistently shown
to induce comparable lesions in leaves of European ash (*F. excelsior*), American green ash (*F.*
pennsylvanica) as well as Manchurian ash (*F. mandshurica*) (Kowalski, Bilanski and Holdenrieder,
2015; Gross and Sieber, 2016; Hietala *et al.*, 2022). Both expressing PCD-inducing effector candidates
and causing necrosis on the host leaf tissue suggest that *H. albidus* is a mild leaf pathogen of ash trees
rather than a nonpathogenic endophyte, as it has been long considered. The fact that *H. albidus* has not
been associated with the recent continental breakout of the ash dieback epidemic in Europe may be
attributed to its relatively low abundance, perhaps due to its low fecundity, in addition to potential
environmental selection pressures that may have acted disadvantageously on its spread and virulence
within its native range (Drenkhan *et al.*, 2016; Hietala *et al.*, 2022) as compared with invasive *H.*
fraxineus. Furthermore, European ash has coexisted with native *H. albidus* for a much longer time and
may, therefore, have evolved to minimise the cost of infection by this pathogen, but was in fact
previously unexposed and immunologically naïve to invasive *H. fraxineus*.

Pathogenicity on European ash

Even with the availability of large genome data of numerous groups of plant-interacting fungi, no study
has, to date, addressed a clear correlation between genomic features or secretome composition of fungi
and their pathogenicity potential. This is mainly due to the complexity of interactions between fungi
and their host plants. As such, many fungal species include pathogenic as well as non-pathogenic
members (Nirmaladevi *et al.*, 2016), and others possess pathogenic and nonpathogenic states (Ridout
and Newcombe, 2017). Therefore, predicting the pathogenicity of fungal microorganisms is as yet not
accurate. Nevertheless, we set out to investigate the pathogenicity of all three *Hymenoscyphus* species,
studied here. Stems of European ash seedlings were inoculated with mycelia of *H. koreanus*, *H. occultus*
and *H. albidus*. *H. fraxineus* was used as a positive control and sterile wood plugs as negative controls.
These latter did not induce the formation of any necrotic lesions on ash stems (Fig 6a-c). However,
inoculations with each of the four *Hymenoscyphus* species resulted in necrotic lesions manifested by

discoloration of the stem tissue to light or dark brown (Fig 6d-o). In our pathogenicity test, lesions

developed by *H. fraxineus*, *H. koreanus* and *H. occultus* were relatively deeper than those formed by

H. albidus (Fig 6m-o), suggesting a slight difference in virulence. Our results demonstrate that all four

related *Hymenoscyphus* species develop pathogenic growth on European ash stems.

Conclusions

In this study, we explored the pathogenicity of three fungal species closely related to the ash dieback

pathogen, *H. fraxineus*. We sequenced the genomes of two new *Hymenoscyphus* species of Asian

origin; *H. koreanus* and *H. occultus* as well as a native European isolate of *H. albidus*, which has long

been considered as a harmless native decomposer and commensal on *F. excelsior* (Hietala and Solheim,

2011). Recently isolated from ash tree collections, the two new Asian species have not previously been

reported as pathogens (Gross and Han, 2015). The genomes of all three sequenced species are of similar

size and GC content and are comparable to *H. fraxineus* genome. Phylogenetic analysis of core

eukaryotic genes identified *H. albidus* and *H. koreanus* as sister species, whilst *H. occultus* was a more

distant relative in the clade (Fig. 1), which is in agreement with previous findings (Gross and Han,

2015). The past two decades have witnessed a surge in newly-discovered phytopathogenic species,

which impacted phylogenetic analyses priorly performed. One such example is the fungal pathogen

causal of sudden oak death, *Phytophthora ramorum*, which was first isolated in 2000 (Werres *et al.*,

2001); fifteen years later, over 50 other Phytophthora species have been discovered and added to the

Tree of Life. Currently, a large part of the inventory of pathogenic forest fungi remains undescribed;

they are more likely to be discovered when they start causing noticeable, sometimes irreparable, damage

to trees and the environmental landscape as a whole.

To identify genes that play a role in *Hymenoscyphus-Fraxinus* interaction, we analysed the secretomes

of *H. albidus*, *H. koreanus*, and *H. occultus*, and we compared them to the previously described

secretome of *H. fraxineus*. Whilst most predicted secreted proteins are highly heterogenous in all four

secretomes, a high number of 1133 proteins were shared between the ash dieback pathogen *H. fraxineus*

from one hand and the other three related *Hymenoscyphus* fungi on the other hand. One particular group

of shared proteins, toxins and cell-death inducing proteins, was not only expressed, but also induced

upon growth of the fungal species on detached leaves of European ash. These closely related species

may, thus, share possible common niche adaptation, infection mechanisms or disease development

strategies on European ash, raising the intriguing possibility that the three *Hymenoscyphus* species

sequenced here may be pathogenic on European ash. Indeed, pathogenicity test using artificial infection

of European ash stems with mycelia of all four *Hymenoscyphus* species produced necrotic lesions under

controlled conditions, confirming the virulence of these fungal species on European ash trees (Fig 6).

Lesions induced by all species were comparable, except those induced by *H. albidus*, which were

slightly smaller in size and reduced in depth. Future possible introduction of Asian *H. koreanus* and *H.*
occultus populations into Europe is likely to jeopardise the survival of European ash, hence the need
for heightened biosecurity awareness to guide conservation strategies. This study is the first to discover
new potential ash dieback pathogens, raising awareness and promoting plant disease surveillance and
investigations in environmental landscapes. As a consequence of human activity, trade, and climate
change, plant pathogens are broadening their geographical ranges and increasingly appearing in new
habitats (Rafiqi *et al.*, 2018). Forests are a particularly vulnerable ecosystem. Thus, the increasing need
for pathogen surveillance and forecast for better detection and response to future outbreaks.

Material and Methods

Fungal isolates, maintenance, and sequencing

Asian isolates: *Hymenoscyphus koreanus* isolate F52847-4 and *H. occultus* isolate F52847-22 were
collected in Korea from petioles of Korean ash and kindly provided by Dr. Andrin Gross (Swiss Federal
Institute for Forest Snow and Landscape Research WSL, Birmensdorfm Switzerland). Fungal isolates
were cultured in potato dextrose broth (VWR chemicals, USA) at room temperature and fungal mycelial
mats were collected for DNA extraction 21 days after inoculation. Fungal DNA was extracted using
E.Z.N.A HP plant DNA Mini kit (Omega Biotek, USA). The DNA was sent for library construction
with TruSeq DNA PCR-free library prep kit with 350-bp insert (Illumina, USA) prior to sequencing
with Illumina HiSeq 2500 platform on rapid run mode generating 2x250 bp (Macrogen Inc, Korea).

European isolates: *H. albidus* was collected near Aberystwyth in Wales, UK, and kindly provided by
Dr. Fiona Corke (IBERS, Plas Gogerddan, Aberystwyth University, United Kingdom) and Dr. Anne
Edwards (John Innes Centre, Norwich Research Park, United Kingdom). Single spore isolates were
grown in liquid culture, harvested, freeze-dried, and the genomic DNA extracted using the MagAttract
HMW DNA Kit (QIAGEN, Germany). Library preparation and sequencing were done at the Earlham
Institute, Norwich. Another isolate of *H. albidus* (ID 5/37/E5) kindly provided by Dr Halvor Solheim,
Norwegian Institute of Bioeconomy Research, Ås (NIBIO) was used for inoculation tests as described
below.

Genome assembly and annotation

Paired-end raw reads were adapter and quality trimmed (Q≥23) using Bbduk in the BBTools suite
version 37.25 (https://jgi.doe.gov/data-and-tools/bbtools/). The draft genomes of the three
Hymenoscyphus species were *de novo* assembled using SPAdes version 3.10.1(Bankevich *et al.*, 2012)
with the following parameters: k-mers 55, 77, 99 and 127, -- careful and –cov-cutoff auto. We used
Blobtools version 1.1.1 (https://github.com/DRL/blobtools) and nucmer (Kurtz *et al.*, 2004) to identify

and dislodge scaffolds of bacterial contaminants and repetitive scaffolds (< 1,000 bp), respectively. We

used the k-mer Analysis Toolkit (KAT) (Mapleson *et al.*, 2017) and BUSCO version 3 (Simão et al.,

2015) against the near-universal single-copy orthologues of Sodariomycetes odb9.

Annotation of the draft genomes of *H. koreanus* F52847-4 and *H. occultus* F52847-22 was achieved

using the FunGAP pipeline version 1.0 (Min, Grigoriev and Choi, 2017), while the annotation of the

draft genome of *H. albidus* was carried out with the funannotate eukaryotic genome annotation pipeline

version 1.2 (https://github.com/nextgenusfs/funannotate). Both FunGAP and funannotate incorporate

genome masking using RepeatModeler/Repeatmasker, *ab initio* gene prediction using either Augustus

and Maker for FunGAP or Augustus and GeneMark for funannotate and evidence-based evaluation of

the predicted gene models. Functional annotation of the predicted coding sequences (CDSs) from the

H. *albidus*, *H. koreanus*, and *H. occultus* genome assemblies depended on InterproScan version

5.25.64.0 (Jones *et al.*, 2014) using PFAM database version 31.0 (Mistry *et al.*, 2021) for *H. koreanus*

and *H. occultus* and PFAM and CDD database version 3.16 (Mistry *et al.*, 2021) for *H. albidus*. The

genome assemblies of *H. koreanus* F52847-1 and *H. occultus* F52847-22 and *H. albidus* were deposited

at the Electronic Research Data Archive at the University of Copenhagen and can be accessed via

https://sid.erda.dk/sharelink/C4VcS7pMHV.

Phylogenetic analysis

To generate a phylogenetic tree, core eukaryotic genes (CEGs) were identified using the Ascomycota

lineage in BUSCO v5.2.1 (Simão *et al.*, 2015). From five *Hymenoscyphus* species, 1560 CEG single-

copy orthogroups were identified. The species investigated were *H. fraxineus*, *H. albidus*, *H. koreanus*,

and *H. occultus*, with *H. varicosporoides* used as an outgroup. Orthologous genes were aligned using

MAFFT v7.271 (Katoh *et al.*, 2002). A phylogenetic tree was generated using RaxML v8.2.12

(Stamatakis, 2014) with the PROTGAMMAGTR model and *H. varicosporoides* as the root. The tree

was visualised using FigTree v1.4.3 (Rambaut, 2012).

Secretome prediction

We used a previously described pipeline (Rafiqi *et al.*, 2022) to predict fungal secretomes of *H. albidus*,

H. occultus, and *H. koreanus*. The pipeline is similar to the one previously used to mine the secretome

of *H. fraxineus* (McMullan *et al.*, 2018), and uses SignalP 4. 1f (Nielsen, 2017) to filter proteins that

contain predicted signal peptides. The set of predicted secreted proteins was further used in the pipeline

to predict transmembrane helices with TMHMM 2.0c (Krogh *et al.*, 2001) and cellular localization

signals with TargetP 1.1b (Emanuelsson *et al.*, 2000). Protein sequences containing predicted

transmembrane helices or mitochondrial targeting signal were removed from the list. The remaining set

of proteins was annotated with Hmmer (Zhang *et al.*, 2018) against PfamScan (Finn *et al.*, 2014a; Finn

et al., 2014b) for domain information, TargetP (Almagro Armenteros *et al.*, 2019) for subcellular

localisation, Predictnls (Cokol, Nair and Rost, 2000) for prediction of nuclear localization signals, T-
Reks (Jorda and Kajava, 2009) for detection of repeats, Disulfinder (database: uniprotkb/swiss-prot) for
prediction of the disulfide bond, and MOTIF search for the search of known motifs. The positions of
the motifs RxLR, [LI]xAR, [RK]CxxCx12}H, [YFW]xC, YxSL[RK], G[IFY][ALST]R, DELD, and
[SG]PC[KR]P were identified with a script based on regular expressions.

Genetic diversity

To investigate genetic diversity within coding sequences, OrthoFinder v.2.2.6 (Emms and Kelly, 2019)
was used to identify orthogroups from four *Hymenoscyphus* species: *H. fraxineus*, *H. albidus*, *H.*
koreanus, and *H. occultus*. Single-copy orthogroups were selected and orthologous genes from the four
species were each combined into a single file. Genes were aligned using GUIDANCE2 in codons with
the MAFFT algorithm (Sela *et al.*, 2015). Due to erroneous sequence lengths, 37 genes were removed,
leaving 7308 genes.

Synonymous (dS) and non-synonymous (dN) mutations were quantified using (Yang, 2007) PAML
v4.9 with the yn00 method on the single-copy orthogroups. The dN/dS ratio was calculated. DnaSP
(Rozas *et al.*, 2003) was run on the orthologous genes to investigate single nucleotide varient density
per gene, which was calculated by dividing the number of SNPs by the net number of sites. DnaSP
failed to run on two genes, leaving 7306 genes.

Stem inoculation

An inoculation experiment was performed on 4/7/2018 to determine if inoculation with the fungi (*H.*
koreanus F52847-4, *H. occultus* F52847-22, and *H. albidus* 5/37/E5) could cause the development of
necrotic lesions on *Fraxinus excelsior*. Two-year-old open-pollinated offspring from clone #35 from
the clonal seed orchard Tuse Næs (N55° 45 57.99 E11° 42 47.48) that had been kept under disease-free
conditions were used for the experiment. The seedlings were inoculated with infected wood plugs.
Before inoculation, sterile wood plugs (3 x 4 x 10 mm^3 in size) had been placed on five week old
cultures allowing the mycelium to colonize the wood plugs. An incision was cut into the stem of the
plant and the infected wood plug was placed in the incision. Parafilm was wrapped around the stem. A
culture of *H. fraxineus* strain 18.3 (Kosawang *et al.*, 2020) was used as a positive control and sterile
wood plugs as negative controls. Three biological replicates were performed for each inoculation type
(i.e. the three species and the positive and negative controls). During the experiment, the plants were
kept in a closed chamber at room temperature and 16 hours of light. The experiment was terminated
after 5 weeks (8/8/2018). Parafilm was removed and photos of the lesions were taken for documentation.

qRT-PCR of selected *Hymenoschyphus* effector candidates

Six, two-week-old mycelia plugs of the three *Hymenoscyphus* species and *H. fraxineus* strain 18.3 were grown on detached leaves of common ash (*Fraxinus excelsior*) clone 27x rinsed with running tap water for 30 mins. The plugs were placed face-down allowing the fungal mycelia to interact with the leaves. Three of the plugs were harvested after 3 and 6 days after inoculation (DAI) for RNA extraction. For the control treatment, the plugs were grown in malt extract broth and were collected accordingly after 3 and 6 DAI. Total RNA was isolated from the plugs using an E.Z.N.A Plant RNA kit (Omega Bio-tek, USA) with on-column DNase I digestion, and cDNA was prepared from 500 ng total RNA using a qScript cDNA synthesis kit (Quanta Biosciences, USA) following the manufacturer's instruction. Three biological replicates of control and treatment were performed for each species of *Hymenoscyphus*.

Gene expression analysis was performed in two technical replicates for each biological replicate on a Mx3005P qPCR system (Stratagene, USA) and FIREPol EvaGreen qPCR Mix Plus (Solis Biodyne, Estonia). Analysis of melting curves was carried out at the end of each run to detect nonspecific amplifications. Relative expression of the effector-encoding genes was calculated in relation to the expression of the beta-tubulin gene according to the $2^{-\Delta\Delta CT}$ method (Livak and Schmittgen, 2001). Gene expression data were analysed statistically using student t-test with a confidential level of 95% implemented in Microsoft Excel version 16.30 (Microsoft Corp., USA).

Data availability

The genome assemblies and annotations of *H. koreanus* F52847-1, *H. occultus* F52847-22 and *H. albidus* were deposited at the Electronic Research Data Archive at the University of Copenhagen and can be accessed via https://sid.erda.dk/sharelink/DljEqJhxMG.

Acknowledgements

We thank Dr Andrin Gross (The Swiss Federal Institute for Forest, Snow and Landscape Research, Switzerland), Dr Fiona Corke (Aberystwyth University, Aberystwyth, UK), Dr Anne Edwards, Professor Allan Downie (John Innes Centre, Norwich, UK) and Dr Halvor Solheim (Norwegian Institute of Bioeconomy Research, Ås (NIBIO)) for providing fungal material. We thank our funders Godfred Birkedal Hartmanns Familiefond (2017), the Danish Council for Independent Research (grant no. 6111-00254) and Royal Botanic Gardens Kew Pilot Study Fund 11231-105. Work at the Earlham Institute (EI; MM, JAP & HY) was supported by Biotechnology and Biological Sciences Research Council (BBSRC), part of UK Research and Innovation, through the Core Capability Grant (BB/CCG1720/1) and (BBS/E/T/000PR9818) WP1 Signatures of Domestication and Adaptation. Sequencing at the EI is supported by BBSRC National Capability in Genomics and Single

398 Cell Analysis (BBS/E/T/000PR9816) by members of the Genomics Pipelines and Core Bioinformatics

399 Groups. HY & <u>JAP</u> were also supported by the BBSRC funded Norwich Research Park Biosciences

400 Doctoral Training Partnership grants BB/M011216/1 & BB/T008717/1 as well as JAP Year in Industry

401 (BBS/E/T/000PR9811).

Authors contribution

404 MR, MM and LRN planned and designed the research. MR, CK, JAP, JL, HY, MM and LRN performed

405 experiments and analysed data. MR, CK, MM and LRN wrote the manuscript. MR and CK contributed

406 equally.

References

Almagro Armenteros, J. J., Salvatore, M. A.-O., Emanuelsson, O. A.-O., Winther, O., von Heijne, G. A.-
O., Elofsson, A. A.-O. and Nielsen, H. A.-O. (2019) 'Detecting sequence signals in targeting peptides
using deep learning. LID - 10.26508/lsa.201900429 [doi] LID - e201900429', (2575-1077 (Electronic)).
Alpert, P., Bone, E. and Holzapfel, C. (2000) 'Invasiveness, invasibility and the role of environmental
stress in the spread of non-native plants', *Perspectives in Plant Ecology, Evolution and Systematics,*
3(1), pp. 52-66.
Bacon, C. W. and Hinton, D. M. (1996) 'Symptomless endophytic colonization of maize by Fusarium
moniliforme', *Canadian Journal of Botany,* 74(8), pp. 1195-1202.
Bacon, C. W. and White, J. (2000) *Microbial endophytes.* CRC press.
Bailey, B. A. (1995) 'Purification of a protein from culture filtrates of *Fusarium oxysporum* that
induces ethylene and necrosis in leaves of *Erythroxylum coca*', *Phytopathology,* 85, pp. 1250–1255.
Balabanova, L. A., Gafurov, Y. M., Pivkin, M. V., Terentyeva, N. A., Likhatskaya, G. N. and Rasskazov,
V. A. (2012) 'An extracellular S1-type nuclease of marine fungus Penicillium melinii', *Mar Biotechnol*
(NY), 14(1), pp. 87-95.
Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M.,
Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler, G.,
Alekseyev, M. A. and Pevzner, P. A. (2012) 'SPAdes: a new genome assembly algorithm and its
applications to single-cell sequencing', *J Comput Biol,* 19(5), pp. 455-77.
Brader, G., Compant, S., Vescio, K., Mitter, B., Trognitz, F., Ma, L.-J. and Sessitsch, A. (2017) 'Ecology
and Genomic Insights into Plant-Pathogenic and Plant-Nonpathogenic Endophytes', *Annual Review*
of Phytopathology, 55(1), pp. 61-83.
Cokol, M., Nair, R. and Rost, B. (2000) 'Finding nuclear localization signals', *EMBO Rep,* 1(5), pp. 411-
5.
Drenkhan, R., Riit, T., Adamson, K. and Hanso, M. (2016) 'The earliest samples of Hymenoscyphus
albidus vs. H. fraxineus in Estonian mycological herbaria', *Mycological Progress,* 15(8), pp. 835-844.
Drenkhan, R., Solheim, H., Bogacheva, A., Riit, T., Adamson, K., Drenkhan, T., Maaten, T. and Hietala,
A. M. (2017) 'Hymenoscyphus fraxineus is a leaf pathogen of local Fraxinus species in the Russian Far
East', *Plant Pathology,* 66(3), pp. 490-500.
Elfstrand, M., Chen, J., Cleary, M., Halecker, S., Ihrmark, K., Karlsson, M., Davydenko, K., Stenlid, J.,
Stadler, M. and Durling, M. B. (2021) 'Comparative analyses of the Hymenoscyphus fraxineus and

Hymenoscyphus albidus genomes reveals potentially adaptive differences in secondary metabolite and transposable element repertoires', *BMC Genomics,* 22(1), pp. 503.

Emanuelsson, O., Nielsen H Fau - Brunak, S., Brunak S Fau - von Heijne, G. and von Heijne, G. (2000) 'Predicting subcellular localization of proteins based on their N-terminal amino acid sequence', (0022-2836 (Print)).

Emms, D. M. and Kelly, S. (2019) 'OrthoFinder: phylogenetic orthology inference for comparative genomics', *Genome Biol,* 20(1), pp. 238.

Fellbrich, G., Romanski, A., Varet, A., Blume, B., Brunner, F., Engelhardt, S., Felix, G., Kemmerling, B., Krzymowska, M. and Nurnberger, T. (2002) 'NPP1, a Phytophthora-associated trigger of plant defense in parsley and Arabidopsis', *Plant J,* 32(3), pp. 375-90.

Feng, B. Z., Zhu, X. P., Fu, L., Lv, R. F., Storey, D., Tooley, P. and Zhang, X. G. (2014) 'Characterization of necrosis-inducing NLP proteins in Phytophthora capsici', *BMC Plant Biol,* 14, pp. 126.

Finn, R. D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R. Y., Eddy, S. R., Heger, A., Hetherington, K., Holm, L., Mistry, J., Sonnhammer, E. L., Tate, J. and Punta, M. (2014a) 'Pfam: the protein families database', *Nucleic Acids Res,* 42(Database issue), pp. D222-30.

Finn, R. D., Bateman A Fau - Clements, J., Clements J Fau - Coggill, P., Coggill P Fau - Eberhardt, R. Y., Eberhardt Ry Fau - Eddy, S. R., Eddy Sr Fau - Heger, A., Heger A Fau - Hetherington, K., Hetherington K Fau - Holm, L., Holm L Fau - Mistry, J., Mistry J Fau - Sonnhammer, E. L. L., Sonnhammer El Fau - Tate, J., Tate J Fau - Punta, M. and Punta, M. (2014b) 'Pfam: the protein families database', (1362-4962 (Electronic)).

Fisher, M. C., Henk, D. A., Briggs, C. J., Brownstein, J. S., Madoff, L. C., McCraw, S. L. and Gurr, S. J. (2012) 'Emerging fungal threats to animal, plant and ecosystem health', *Nature,* 484(7393), pp. 186-194.

Gross, A. and Han, J. G. (2015) 'Hymenoscyphus fraxineus and two new Hymenoscyphus species identified in Korea', *Mycological Progress,* 14(4), pp. 19.

Gross, A., Hosoya, T. and Queloz, V. (2014) 'Population structure of the invasive forest pathogen Hymenoscyphus pseudoalbidus', *Mol Ecol,* 23(12), pp. 2943-60.

Gross, A. and Sieber, T. N. (2016) 'Virulence of Hymenoscyphus albidus and native and introduced Hymenoscyphus fraxineus on Fraxinus excelsior and Fraxinus pennsylvanica', *Plant Pathology,* 65(4), pp. 655-663.

Hane, J. K., Paxman, J., Jones, D. A. B., Oliver, R. P. and de Wit, P. (2020) '"CATAStrophy," a Genome-Informed Trophic Classification of Filamentous Plant Pathogens – How Many Different Types of Filamentous Plant Pathogens Are There?', *Frontiers in Microbiology,* 10.

Hardoim, P. R., van Overbeek, L. S., Berg, G., Pirttilä, A. M., Compant, S., Campisano, A., Döring, M. and Sessitsch, A. (2015) 'The Hidden World within Plants: Ecological and Evolutionary Considerations for Defining Functioning of Microbial Endophytes', *Microbiol Mol Biol Rev,* 79(3), pp. 293-320.

Hayes, K. R. and Barry, S. C. (2008) 'Are there any consistent predictors of invasion success?', *Biological Invasions,* 10(4), pp. 483-506.

Hietala, A. M., Agan, A., Nagy, N. E., Børja, I., Timmermann, V., Drenkhan, R. and Solheim, H. (2022) 'The Native Hymenoscyphus albidus and the Invasive Hymenoscyphus fraxineus Are Similar in Their Necrotrophic Growth Phase in Ash Leaves', *Frontiers in Microbiology,* 13.

Hietala, A. M. and Solheim, H. (2011) 'Hymenoscyphus species associated with European ash1', *EPPO Bulletin,* 41(1), pp. 3-6.

Hill, L., Jones, G., Atkinson, N., Hector, A., Hemery, G. and Brown, N. (2019) 'The £15 billion cost of ash dieback in Britain', *Current Biology,* 29(9), pp. R315-R316.

Jones, P., Binns, D., Chang, H. Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S. Y., Lopez, R. and Hunter, S. (2014) 'InterProScan 5: genome-scale protein function classification', *Bioinformatics,* 30(9), pp. 1236-40.

Jorda, J. and Kajava, A. V. (2009) 'T-REKS: identification of Tandem REpeats in sequences with a K-meanS based algorithm', (1367-4811 (Electronic)).

Katoh, K., Misawa, K., Kuma, K. and Miyata, T. (2002) 'MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform', *Nucleic Acids Res,* 30(14), pp. 3059-66.
Kirk, P. M., Cannon, P. F., David, J. C. and Stalpers, J. A. (2008) 'Ainsworth and Bisby's Dictionary of
the fungi, Wallingford', *International Mycological Institute*.
Kosawang, C., McKinney, L. V., Nielsen, L. R. and Kjær, E. D. (2020) 'Variation in aggressiveness of
Hymenoscyphus fraxineus genotypes amid the ash dieback epidemic', *Plant Pathology,* 69(4), pp.
677-684.
Kowalski, T., Bilanski, P. and Holdenrieder, O. (2015) 'Virulence of Hymenoscyphus albidus and H.
fraxineus on Fraxinus excelsior and F. pennsylvanica', *PLoS One,* 10(10), pp. e0141592.
Krogh, A., Larsson B Fau - von Heijne, G., von Heijne G Fau - Sonnhammer, E. L. and Sonnhammer, E.
L. (2001) 'Predicting transmembrane protein topology with a hidden Markov model: application to
complete genomes', (0022-2836 (Print)).
Kumakura, N., Singkaravanit-Ogawa, S., Gan, P., Tsushima, A., Ishihama, N., Watanabe, S., Seo, M.,
Iwasaki, S., Narusaka, M., Narusaka, Y., Takano, Y. and Shirasu, K. (2021) 'Guanosine-specific single-
stranded ribonuclease effectors of a phytopathogenic fungus potentiate host immune responses',
bioRxiv, pp. 2021.10.13.464185.
Kurtz, S., Phillippy, A., Delcher, A. L., Smoot, M., Shumway, M., Antonescu, C. and Salzberg, S. L.
(2004) 'Versatile and open software for comparing large genomes', *Genome Biol,* 5(2), pp. R12.
Mansfield, J., Brown, I. and Papp-Rupar, M. (2019) 'Life at the edge – the cytology and physiology of
the biotroph to necrotroph transition in Hymenoscyphus fraxineus during lesion formation in ash',
Plant Pathology, 68(5), pp. 908-920.
Mansfield, J. W., Galambos, N. and Saville, R. (2018) 'The use of ascospores of the dieback fungus
Hymenoscyphus fraxineus for infection assays reveals a significant period of biotrophic interaction in
penetrated ash cells', *Plant Pathology,* 67(6), pp. 1354-1361.
Manzotti, A., Bergna, A., Burow, M., Jørgensen, H. J. L., Cernava, T., Berg, G., Collinge, D. B. and
Jensen, B. (2020) 'Insights into the community structure and lifestyle of the fungal root endophytes
of tomato by combining amplicon sequencing and isolation approaches with phytohormone
profiling', *FEMS Microbiology Ecology,* 96(5).
Mapleson, D., Garcia Accinelli, G., Kettleborough, G., Wright, J. and Clavijo, B. J. (2017) 'KAT: a K-mer
analysis toolkit to quality control NGS datasets and genome assemblies', *Bioinformatics,* 33(4), pp.
574-576.
McMullan, M., Rafiqi, M., Kaithakottil, G., Clavijo, B. J., Bilham, L., Orton, E., Percival-Alwyn, L., Ward,
B. J., Edwards, A., Saunders, D. G. O., Garcia Accinelli, G., Wright, J., Verweij, W., Koutsovoulos, G.,
Yoshida, K., Hosoya, T., Williamson, L., Jennings, P., Ioos, R., Husson, C., Hietala, A. M., Vivian-Smith,
A., Solheim, H., MaClean, D., Fosker, C., Hall, N., Brown, J. K. M., Swarbreck, D., Blaxter, M., Downie,
J. A. and Clark, M. D. (2018) 'The ash dieback invasion of Europe was founded by two genetically
divergent individuals', *Nat Ecol Evol,* 2(6), pp. 1000-1008.
Min, B., Grigoriev, I. V. and Choi, I. G. (2017) 'FunGAP: Fungal Genome Annotation Pipeline using
evidence-based gene model evaluation', *Bioinformatics,* 33(18), pp. 2936-2937.
Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E. L. L., Tosatto, S.
C. E., Paladin, L., Raj, S., Richardson, L. J., Finn, R. D. and Bateman, A. (2021) 'Pfam: The protein
families database in 2021', *Nucleic Acids Res,* 49(D1), pp. D412-D419.
Mitchell, R. (2014) *The potential ecological impact of ash dieback in the UK.*
Nguyen, L. T., Schmidt, H. A., von Haeseler, A. and Minh, B. Q. (2015) 'IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies', *Mol Biol Evol,* 32(1), pp. 268-
74.
Nielsen, H. (2017) 'Predicting Secretory Proteins with SignalP', (1940-6029 (Electronic)).
Nirmaladevi, D., Venkataramana, M., Srivastava, R. K., Uppalapati, S. R., Gupta, V. K., Yli-Mattila, T.,
Clement Tsui, K. M., Srinivas, C., Niranjana, S. R. and Chandra, N. S. (2016) 'Molecular phylogeny,
pathogenicity and toxigenicity of Fusarium oxysporum f. sp. lycopersici', *Scientific Reports,* 6(1), pp.
21367.

Park, H.-J., Wang, W., Curlango-Rivera, G., Xiong, Z., Lin, Z., Huskey, D. A., Hawes, M. C., VanEtten, H. D. and Turgeon, B. G. (2019) 'A DNase from a Fungal Phytopathogen Is a Virulence Factor Likely Deployed as Counter Defense against Host-Secreted Extracellular DNA', *mBio,* 10(2), pp. e02805-18.

Pennington, H. G., Jones, R., Kwon, S., Bonciani, G., Thieron, H., Chandler, T., Luong, P., Morgan, S. N., Przydacz, M., Bozkurt, T., Bowden, S., Craze, M., Wallington, E. J., Garnett, J., Kwaaitaal, M., Panstruga, R., Cota, E. and Spanu, P. D. (2019a) 'The fungal ribonuclease-like effector protein CSEP0064/BEC1054 represses plant immunity and interferes with degradation of host ribosomal RNA', *PLoS Pathog,* 15(3), pp. e1007620.

Pennington, H. G., Jones, R., Kwon, S., Bonciani, G., Thieron, H., Chandler, T., Luong, P., Morgan, S. N., Przydacz, M., Bozkurt, T., Bowden, S., Craze, M., Wallington, E. J., Garnett, J., Kwaaitaal, M., Panstruga, R., Cota, E. and Spanu, P. D. (2019b) 'The fungal ribonuclease-like effector protein CSEP0064/BEC1054 represses plant immunity and interferes with degradation of host ribosomal RNA', *PLoS pathogens,* 15(3), pp. e1007620-e1007620.

Pereira, E., Vázquez de Aldana, B. R., San Emeterio, L. and Zabalgogeazcoa, I. (2019) 'A survey of culturable fungal endophytes from Festuca rubra subsp. pruinosa, a grass from marine cliffs, reveals a core microbiome', *Frontiers in microbiology,* 9, pp. 3321.

Rafiqi, M. (2018) 'Plant Killers- State of The World's Fungi 2018', pp. 56–61.

Rafiqi, M., Jelonek, L., Diouf, A. M., Mbaye, A., Rep, M. and Diarra, A. (2021) 'Profile of the in silico secretome of the palm dieback pathogen, a fungus that puts natural oases at risk', *bioRxiv*, pp. 2021.11.18.469146.

Rafiqi, M., Jelonek, L., Diouf, A. M., Mbaye, A., Rep, M. and Diarra, A. (2022) 'Profile of the in silico secretome of the palm dieback pathogen, Fusarium oxysporum f. sp. albedinis, a fungus that puts natural oases at risk', *PLoS One,* 17(5), pp. e0260830.

Rafiqi, M., Saunders, D., McMullan, M., Oliver, R., Bone, R., Fones, H., Gurr, S. a., Vincent, D. and Coker, T. a. B., Richard (2018) *Plant Killers*: Royal Botanic Gardens, Kew.

Rambaut, A. 2012. Figtree 1.4.3. Available at http://tree.bio.ed.ac.uk/software/figtree/.

Ridout, M. and Newcombe, G. (2017) 'Sydowia polyspora is both a Foliar Endophyte and a Preemergent Seed Pathogen in Pinus ponderosa', *Plant Disease,* 102(3), pp. 640-644.

Rozas, J., Sánchez-DelBarrio, J. C., Messeguer, X. and Rozas, R. (2003) 'DnaSP, DNA polymorphism analyses by the coalescent and other methods', *Bioinformatics,* 19(18), pp. 2496-7.

Santini, A., Ghelardini, L., De Pace, C., Desprez-Loustau, M. L., Capretti, P., Chandelier, A., Cech, T., Chira, D., Diamandis, S., Gaitniekis, T., Hantula, J., Holdenrieder, O., Jankovsky, L., Jung, T., Jurc, D., Kirisits, T., Kunca, A., Lygis, V., Malecka, M., Marcais, B., Schmitz, S., Schumacher, J., Solheim, H., Solla, A., Szabò, I., Tsopelas, P., Vannini, A., Vettraino, A. M., Webber, J., Woodward, S. and Stenlid, J. (2013) 'Biogeographical patterns and determinants of invasion by forest pathogens in Europe', *New Phytologist,* 197(1), pp. 238-250.

Sela, I., Ashkenazy, H., Katoh, K. and Pupko, T. (2015) 'GUIDANCE2: accurate detection of unreliable alignment regions accounting for the uncertainty of multiple parameters', *Nucleic Acids Res,* 43(W1), pp. W7-14.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. and Zdobnov, E. M. (2015) 'BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs', *Bioinformatics,* 31(19), pp. 3210-2.

Stamatakis, A. (2014) 'RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies', *Bioinformatics,* 30(9), pp. 1312-1313.

Stenlid, J., Elfstrand, M., Cleary, M., Ihrmark, K., Karlsson, M., Davydenko, K. and Durling, M. B. (2017) 'Genomes of Hymenoscyphus fraxineus and Hymenoscyphus albidus encode surprisingly large cell wall degrading potential, balancing saprotrophic and necrotrophic signatures', *Baltic Forestry,* 23(1), pp. 41-51.

van Dam, P., Fokkens, L., Ayukawa, Y., van der Gragt, M., Ter Horst, A., Brankovics, B., Houterman, P. M., Arie, T. and Rep, M. (2017) 'A mobile pathogenicity chromosome in Fusarium oxysporum for infection of multiple cucurbit species', *Sci Rep,* 7(1), pp. 9042.

Vincent, D., Rafiqi, M. and Job, D. (2019) 'The Multiple Facets of Plant-Fungal Interactions Revealed
Through Plant and Fungal Secretomics', *Front Plant Sci,* 10, pp. 1626.
Werres, S., Marwitz, R., Man In't veld, W. A., De Cock, A. W. A. M., Bonants, P. J. M., De Weerdt, M.,
Themann, K., Ilieva, E. and Baayen, R. P. (2001) 'Phytophthora ramorum sp. nov., a new pathogen on
Rhododendron and Viburnum', *Mycological Research,* 105(10), pp. 1155-1165.
Widmer, H. 'Functional characterisation of a fungal endonuclease effector and regulated host cell
death'.
Yang, Z. (2007) 'PAML 4: phylogenetic analysis by maximum likelihood', *Mol Biol Evol,* 24(8), pp.
1586-91.
Zhang, H., Yohe, T., Huang, L., Entwistle, S., Wu, P., Yang, Z., Busk, P. K., Xu, Y. and Yin, Y. (2018)
'dbCAN2: a meta server for automated carbohydrate-active enzyme annotation', *Nucleic Acids Res,*
46(W1), pp. W95-W101.
Zhao, Y.-J., Hosoya, T., Baral, H.-O., Hosaka, K. and Kakishima, M. (2013a) 'Hymenoscyphus
pseudoalbidus, the correct name for Lambertella albida reported from Japan', *Mycotaxon,* 122(1),
pp. 25-41.
Zhao, Z., Liu, H., Wang, C. and Xu, J.-R. (2013b) 'Comparative analysis of fungal genomes reveals
different plant cell wall degrading capacity in fungi', *BMC Genomics,* 14(1), pp. 274.
Zhao, Z., Liu, H., Wang, C. and Xu, J. R. (2013c) 'Comparative analysis of fungal genomes reveals
different plant cell wall degrading capacity in fungi', *BMC Genomics,* 14, pp. 274.
Zheng, H. and Zhuang, W. (2013) 'Four new species of the genus Hymenoscyphus (fungi) based on
morphology and molecular data', *Science China Life Sciences,* 56(1), pp. 90-100.
Çetinkaya, A. S. and Uzun, Y. (2020) 'Hymenoscyphus caudatus, a new ascomycete record for the
mycobiota of Turkey', *Anatolian Journal of Botany*.

H. albidus

H. koreanus

H. fraxineus

H. occultus

H. varicosporoides

0.03

Relative frequency distribution of CAZymes modules in 4 related *Hymenoscyphus* species

Legend: GH ■ CE ■ GT ■ PL ■ CBM ■ AA ■ GH & CBM ■ CE &CBM