Loss of NR5A1 in Sertoli cells after sex determination changes their cellular identity and induces their death by anoikis

Sirine SOUALI-CRESPO[1], Diana CONDREA[1], Nadège VERNET[1], Betty FÉRET[1], Muriel KLOPFENSTEIN[1], Erwan GRANDGIRARD[1,3], Violaine ALUNNI[1,2], Marie CERCIAT[1,3], Matthieu JUNG[1,3], Chloé MAYERE[4], Serge NEF[4], Manuel MARK[1,5], Frédéric CHALMEL[6*], and Norbert B. GHYSELINCK[1*#]

1. Institut de Génétique et de Biologie Moléculaire et Cellulaire (IGBMC), Département de Génétique Fonctionnelle et Cancer, Centre National de la Recherche Scientifique (CNRS UMR7104), Institut National de la Santé et de la Recherche Médicale (INSERM U1258), Université de Strasbourg (UNISTRA), 1 rue Laurent Fries, BP-10142, F-67404 Illkirch Cedex, France
2. GenomEast Platform, France Génomique consortium, IGBMC, 1 rue Laurent Fries, F-67404 Illkirch Cedex, France
3. Imaging Center, IGBMC, F-67404 Illkirch Cedex, France
4. Department of Genetic Medicine and Development, Faculty of Medicine, University of Geneva, Geneva, Switzerland
5. Service de Biologie de la Reproduction, Hôpitaux Universitaires de Strasbourg (HUS), France
6. Univ Rennes, EHESP, Inserm, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, F-35000 Rennes, France

* These authors contributed equally to this work

Author for correspondence: norbert@igbmc.fr

Tel: +33 388 655 674; Fax: +33 388 653 201

Keywords: anoikis, cell-death, fetal gonad, integrin, Leydig, testis, SF-1, single-cell.

ABSTRACT

NR5A1 is an orphan nuclear receptor crucial for gonadal development in

mammals. In the mouse testis it is expressed both in Sertoli cells (SC) and

Leydig cells (LC). To investigate its role posteriorly to sex determination, we

have generated and analysed mice lacking NR5A1 in SC from embryonic day

(E) 13.5 onwards (*Nr5a1*^{SC–/–} mutants). Ablation of *Nr5a1* impairs the expression

of genes characteristic of SC identity (e.g., *Sox9*, *Amh*), makes SC to

progressively die from E14.5 by a *Trp53*-independent mechanism, and induces

disorganization of the testis cords, which, together, yields germ cells (GC) to

prematurely enter meiosis and die, instead of becoming quiescent. Single-cell

RNA-sequencing experiments revealed that *Nr5a1*-deficient SC acquire a pre-

granulosa cell-like identity, and profoundly modify the landscape of the

adhesion molecules and extracellular matrix they express. We propose

therefore that SC lacking NR5A1 transdifferentiate and die by anoikis. Fetal LC

do not display major changes in their transcriptome, indicating that SC are not

required beyond E14.5 for their emergence or maintenance. In contrast, adult

LC were missing in *Nr5a1*^{SC–/–} postnatal testes. In addition, adult males display

Müllerian duct derivatives (i.e., uterus, vagina), as well as a decreased

anogenital distance and a shorter penis that can be explained by loss of AMH

production and defective HSD17B1- and HSD17B3-mediated synthesis of

testosterone in SC during fetal life. Together, our findings indicate that *Nr5a1*

expressed in SC after the period of sex determination safeguards SC identity,

which maintains proper seminiferous cord organization and prevents GC to

enter meiosis.

INTRODUCTION

Testis and ovary both originate from a bipotential gonad, which is composed of

somatic progenitor cells that are competent to adopt one or the other sex-specific cell

fate, and of primordial germ cells (GC). Sex determination initiates at around

embryonic day (E) 11.5, from when somatic progenitor cells undergo sex-specific cell

differentiation. The specification of the supporting cell lineage [i.e., either Sertoli cells

(SC) in the testis upon SRY expression, or pre-granulosa cells (pGrC) in the ovary

upon WNT/CTNNB1 signalling stabilization] is the first essential step in this process.

Following supporting cell differentiation, the sex-specific fate decision propagates to

the GC and to other somatic lineages, including the steroidogenic cells Leydig cells

(LC) in the testis, which, in turn, drive acquisition of primary and secondary sexual

characters at later stages of development, through hormone secretion [1-3]. In

humans, mutations in genes driving this complex developmental process can give

rise to a group of defects known as disorders of sex development (DSD), the

diagnosis and the management of which requires an understanding of the molecular

mechanisms underpinning cell differentiation [4].

Amongst the genes responsible for DSD is *Nr5a1*, encoding the orphan nuclear

receptor NR5A1 [5], a transcription factor expressed at early stages of development,

when cells are partitioned into adrenal and gonadal primordium [6]. NR5A1 controls

the expression of genes implicated in proliferation, survival and differentiation of

somatic progenitor cells [7]. Therefore, knock-out of *Nr5a1* in the mouse results in

regression of the gonads by E11.5 due to apoptosis of somatic cells, and agenesis of

the adrenal gland [8]. From E11.5, NR5A1 and SRY are expressed in the bipotential

supporting cells of male embryos, where they upregulate SOX9 expression, initiating

thereby pre-SC and ultimately SC differentiation [2,3,9]. Once specified, the SC start

forming cords by enclosing GC as early as E12.5 in the mouse [10], providing them

with a specialized environment that promotes their survival and orchestrates their

differentiation [11]. SC also (i) allow the differentiation of fetal LC (FLC) through

paracrine Desert Hedgehog (DHH) signalling [12], (ii) produce, under the control of

NR5A1 and SOX9, high levels of anti-Müllerian hormone (AMH), which triggers the

regression of the Müllerian duct normally giving rise to the female genitalia [13].

Because gonads are absent in *Nr5a1*-null mice [8], studying the cell-specific

roles of NR5A1 in the developing testis required analysis of mice bearing tissue-

targeted mutations. Here we have analysed the outcome of *Nr5a1* ablation in SC

from E13.5 onwards. We show that loss of NR5A1 from this stage induces some SC

to transdifferentiate into female somatic cells, and to die by a *Trp53*-independent

mechanism related to anoikis.

RESULTS

Deletion of* Nr5a1 *in Sertoli cells

To achieve *Nr5a1* ablation in SC after sex determination, we introduced the

Plekha5[Tg(AMH-cre)1Flor] transgene [14] in mice bearing *lox*P-flanked alleles of *Nr5a1* and

a Cre-dependent reporter transgene [15]. Cre-mediated recombination was assessed

by immunohistochemistry (IHC) using antibodies recognizing the yellow fluorescent

protein (YFP). Surprisingly, YFP was detected in all SC of embryonic day (E) 12.5

gonads (arrows, **Fig.1A,B**), indicating Cre-mediated excision occurred earlier than

anticipated [14]. However, the NR5A1 protein was still detected in SC at E12.5

(**Fig.1A,B**). At E13.5, the nuclei of SC were all NR5A1-positive in the control

(**Fig.1C**), but NR5A1-negative in the mutant testis (**Fig.1D**). Importantly, NR5A1 was

still detected in the nuclei of LC, identified by their expression of the steroid 3 beta-

hydroxy-steroid dehydrogenase type 1 (HSD3B1), in both control and mutant testes

(**Fig.1E,F**). Thus, NR5A1 was lost selectively in SC from E13.5 onwards, generating

males hereafter called *Nr5a1*$^{SC-/-}$ mutants.

Ablation of Nr5a1 _in SC impairs expression of SOX9, SOX8, SOX10 and AMH_

As NR5A1 regulates the expression of *Amh* and *Sox9* genes [9,16], we tested

expression of these two proteins by IHC. Both AMH and SOX9 were detected at

normal levels at E12.5 (**Fig.2A,B,G,H**). Their expression decreased progressively

between E13.5 and E14.5 in *Nr5a1*$^{SC-/-}$ mutant testes (**Fig.2C-F,I-L**). At E14.5, only

few *Nr5a1*-deficient SC were SOX9-positive, some of which with cytoplasmic SOX9

(compare insets, **Fig.2K,L**). The expression of SOX8 and SOX10 was also

decreased in *Nr5a1*$^{SC-/-}$gonads (**Suppl.Fig.1**). Accordingly, the steady state levels of

Amh, *Sox9* and *Sox8* mRNAs was decreased in *Nr5a1*$^{SC-/-}$ whole gonads (**Fig.2S**).

The level of prostaglandin D2 synthase (*Ptgds*) mRNA, another SOX9 target-gene

[17], was also reduced in E14.5 *Nr5a1*$^{SC-/-}$ testes. This finding may explain the

cytoplasmic localization of SOX9 in some SC since prostaglandin D2, the end

product of PTGDS, is required for SOX9 nuclear translocation [18].

In parallel, we analysed the expression of GATA4 and WT1, important for *Amh*

expression and SC differentiation, respectively [19,20]. Both were detected at similar

levels in SC nuclei of control and mutant testes, from E12.5 to E14.5 (**Fig.2M-R;**

Suppl.Fig.1).

**Nr5a1*-deficient SC die through a TRP53-independent mechanism*

Thanks to the expression of the YFP-reporter we quantified the surface

occupied by SC (green pixels) relative to the whole testis surface (**Fig.3A,D**). At

E14.5, this ratio was about 30% lower in mutant than in control testes [28.3 ± 3.8

(n=5) versus 37.9 ± 2.5 (n=5), respectively; p<0.01]. This suggested that *Nr5a1-*

deficient SC had either shut down expression of YFP or had disappeared from the

testis. If YPF expression was shut down, the excised *Nr5a1* allele (L-) should be

present in the testis at birth. If SC were lost, it should no longer be detected. PCR

analysis of genomic DNA showed that the L– allele was not amplified in *Nr5a1*$^{SC–/–}$

mutants (**Suppl.Fig.2A**), indicating that *Nr5a1*-deficient SC were lost from the mutant

testis. In agreement with this possibility, numerous cells displaying features of SC

(i.e., located at the periphery of the testis cords, amongst GATA4-expressing SC)

were TUNEL-positive in E14.5 *Nr5a1*$^{SC–/–}$ testis (arrowheads, **Fig.3E,F**). This

indicates that NR5A1-deficient SC progressively died.

Because TRP53 was not detected in fetal SC at E14.5 (**Fig.3G,H**), we

wondered whether it was functionally involved in SC-death. We introduced

conditional alleles of *Trp53* [21] in *Nr5a1*$^{SC–/–}$ mutants bearing the YFP reporter

transgene. Efficient ablation of *Trp53* was assessed by PCR on genomic DNA

extracted form FACS-purified, YFP-positive, SC (**Suppl.Fig.2B**). E14.5 *Nr5a1*$^{SC–/–}$

and *Nr5a1*$^{SC–/–}$;*Trp53*$^{SC–/–}$ mutants displayed similar reduced surfaces occupied by

SC (**Fig.3I,J**, compare with 3A), out of which most of them had lost expression of

SOX9. At post-natal day 15 (PND15), both *Nr5a1*$^{SC–/–}$ and *Nr5a1*$^{SC–/–}$;*Trp53*$^{SC–/–}$

testes lacked SC-surrounded seminiferous tubules, but contained interstitial cells, out

of which about half of them were HSD3B1-positive LC (**Fig.3K-P**). At adulthood, none

of the interstitial cells expressed the adult LC-specific marker BHMT [22], indicating

this cell-type was unable to emerge (**Fig.3Q-S**). Altogether, this indicates that

NR5A1-deficient SC died, even though they lacked TRP53, yielding a postnatal testis

made of FLC and other interstitial cells.

Nr5a1$^{SC–/–}$ adult males exhibit Müllerian duct retention

To analyse the outcome of the loss of SC and AMH production in *Nr5a1*[SC−/−]

mutants, the male phenotype was investigated at PND60. They all were sterile and

had a shorter anogenital distance (AGD) index [0.30 ± 0.03 mm/g of body weight

(n=16), versus 0.36 ± 0.02 mm/g (n=17) in controls; p<0.001] (**Fig.4A,C**). At autopsy

(n = 8), they displayed normal derivatives of the Wolffian ducts (i.e., epididymis, vas

deferent, seminal vesicle, ampullary glands) and the urogenital sinus (i.e., prostatic

lobes) (**Fig.4E,F**). However, they also displayed Müllerian duct derivatives (**Fig.4F-**

H), namely a vagina, an uterine body and bilateral, uterine horns, which were either

incomplete and truncated (6 out of 8), or complete on one side (2 out of 8).

Histological analyses revealed that the uterine body and vagina displayed a columnar

epithelium (**Fig.4I**) and a stratified, squamous epithelium (**Fig.4J**), as anticipated for

these female organs.

Nonetheless, *Nr5a1*[SC−/−] males displayed 40% lighter seminal vesicles

[4.71 ± 1.66 mg/g of body weight (n=17) versus 7.43 ± 1.22 mg/g (n=16) in controls;

p<0.05] and 10% shorter penis bones [6.7 ± 0.4 mm (n=17) versus 7.5 ± 0.3 mm

(n=16) in controls; p<0.05] (**Fig.4B,D**). As AGD, seminal vesicle growth and penis

bone length vary as a function of androgen exposure [23], we tested blood

testosterone levels. They were comparable at birth [0.25 ± 0.17 ng/ml (n=10) in

mutants, versus 0.22 ± 0.10 ng/ml serum (n=10) in controls; p=0.59], and at PND60

[0.38 ± 0.17 ng/ml (n=17) in controls, versus 0.43 ± 0.23 ng/ml serum (n=16) in

controls; p=0.68], indicating normal testosterone production.

Transcriptomic signatures of cells in control and* Nr5a1 *-deficient gonads.

We then performed single-cell RNA sequencing (scRNA-seq) experiments

using dissociated cell suspensions obtained from 12 control and 16 *Nr5a1*[SC−/−] whole

testes at E14.5. After data processing and quality control, we assembled an atlas

composed of 3,988 and 5,010 control and mutant testicular cells, respectively. On

average, we detected ~13,860 unique molecular indices (UMIs) and ~3,556 genes in

each individual cell. The 8,998 testicular cells were partitioned into 23 cell clusters

(termed C1-C23) and projected into a two-dimensional space (**Fig.5A; Table.S1**).

Known marker genes of distinct cell-types were used to identify each cluster

(**Suppl.Fig.3**).

179   Only a few changes of gene expression were observed between control and

Nr5a1$^{SC-/-}$ testes for endothelial, immune and perivascular cells. In Leydig, interstitial,

and coelomic epithelium cells the expression of 59, 49, and 47 genes was

deregulated (**Table.S2**). Functional analysis revealed that down-regulated genes

were related to ribosome in Leydig (14 genes, *p* value<4.8^{e-15}) and interstitial cells

(12 genes, *p* value<1.3^{e-10}) (**Suppl.Fig.4; Table.S3**). In GC, the expression of 636

genes was significantly deregulated (**Table.S1**), amongst which *Stra8* and *Rec8* were

up, while *Nanos2* and *Piwil4* were down (**Fig.5B**). Processes such as DNA

recombination (33 genes, *p* value<2.0^{e-10}), meiotic cell cycle (32 genes, *p* value<1.1^{e-9})

and oocyte differentiation (8 genes, *p* value<5.6^{e-3}) were identified amongst the

upregulated genes (**Suppl.Fig.4; Table.S3**). In agreement with this finding we

showed that GC were in the S-phase of the meiotic prophase at E14.5 and E15.5 in

Nr5a1$^{SC-/-}$ testes instead of becoming mitotically quiescent, as in the control

situation. Accordingly, many GC of the *Nr5a1*$^{SC-/-}$ testes expressed meiotic proteins,

such as STRA8 or REC8, and died because of premature meiotic initiation

(**Suppl.Fig.5**).

195   Three clusters of SC were identified (C18, C19 and C21), out of which C19 and

C21 were mainly composed of control and NR5A1-deficient SC, respectively

(**Fig.5A**). Almost all SC of C18 were mitotic (**Suppl.Fig.3**). In total, 1,671 genes were

significantly deregulated in SC (**Suppl.Table.S1**), amongst which *Amh*, *Sox9* and

Ptgds were down as anticipated (see above), while *Wnt4*, *Fst* and *Rspo1* were up

(**Fig.5C**). We confirmed deregulated expression of some by IHC (**Fig.5D-K**) and that

of others by RT-qPCR (**Fig.5L**), validating thereby the scRNA-seq experiment.

According to its transcriptomic signature, cluster C20 was assigned the *rete*

testis identity (**Suppl.Fig.3**). This structure is a set of ducts, through which

spermatozoa are transported from the gonad via the efferent duct to the epididymis

[24]. At E14.5, the *rete testis* is identified as epithelial cords between the

mesonephric tubules and the testis cords (**Suppl.Fig.6A**). The *rete testis* cells

express *Nr5a1*, *Pax8* and *Aldh1a3* [25]. As the *Plekha5*$^{Tg(AMH-cre)1Flor}$ transgene is not

functional in the *rete testis* [26], we anticipated that C20 could comprise cells in which

NR5A1 expression was not lost. In agreement, we observed that cells in C20

organized as a continuum, which stemmed from C19 with *Nr5a1*⁺/*Yfp*⁻/*Pax8*⁻

/*Aldh1a3*⁻ cells and extended with *Nr5a1*⁺/*Yfp*⁻/*Pax8*⁺/*Aldh1a3*⁺ cells (*rete*) originating

from both control and *Nr5a1*$^{SC-/-}$ testes. It ended at the junction with C21 with *rete*

testis cells, originating exclusively from the *Nr5a1*$^{SC-/-}$ testes, and in which cre was

active   (*Nr5a1*⁻/*Yfp*⁺/*Pax8*⁺/*Aldh1a3*⁺   cells)   (**Suppl.Fig.6B**).   Accordingly,   IHC

experiments confirmed that *rete testis* cells actually expressed NR5A1, but not YFP,

control and *Nr5a1*$^{SC-/-}$ testes (**Suppl.Fig.6C-J**).

**Nr5a1*-deficient SC change their molecular identity.*

As indicated above, the NR5A1-deficient SC from cluster C21 gained

expression of *Wnt4*, *Fst* and *Rspo1*, which characterize the fetal ovary somatic cells

[4]. This raised the question as to whether SC had changed their identity. To test for

this possibility, we decided to compare our data with a reference atlas of embryonic

and fetal gonad development ranging from E10.5 to E16.5 [25]. After processing this

atlas with the pipeline that we used for our data set, (**Suppl.Fig.7**), we next mapped

our data set to this reference atlas (**Fig.6A**). Most of control and *Nr5a1*$^{SC-/-}$ cell

clusters were predicted to correspond to clusters with identical (or compatible)

identities, as expected (**Fig.6B**). In addition, GC from control testes matched in the

atlas with E13.5 male GC, while those from *Nr5a1*$^{SC-/-}$ testes matched with E13.5

female (i.e., meiotic) GC (**Fig.6C**), as expected from IHC analyses (**Suppl.Fig.5**), and

validating thereby the whole prediction. Interestingly, cluster C20 was predicted to

match the best with *rete testis* cells, a feature fitting well with our analysis by IHC

(see above, **Suppl.Fig.6**). Most importantly, cluster C21 was predicted to correspond

to pGrC in the atlas. Further analysis distinguishing the developmental stages

indicated that they actually matched with E16.5 pGrC (**Fig.6C**). The remaining cells

matched with E13.5 SC. Thus, a fraction of NR5A1-deficient SC lost their identity,

and acquired a "pGrC-like" identity. Accordingly, functional analysis of genes

differentially expressed by NR5A1-deficient SC assigned upregulated genes to WNT

signalling pathway (54 genes, *p* value<4.3e^{-8}), known to drive pGrC differentiation in

the fetal ovary [27] (**Suppl.Fig.4; Table.S3**). Nonetheless, the process of *trans*-

differentiation was not fully achieved because IHC experiments failed to evidence

expression of FOXL2 in the *Nr5a1*$^{SC-/-}$ mutant testis at E15.5 (**Fig.6D,E**).

Ablation of Nr5a1 *in SC induces disorganisation of the testis cords.*

On histological sections at E14.5, the majority of *Nr5a1*$^{SC-/-}$ seminiferous cords

displayed reduced diameters, as well as poorly defined and discontinuous contours

(broken dotted line and arrows, **Fig.7E**). At E15.5, the cords had almost fully

disappeared (broken dotted line, **Fig.7F**). Such irregularities were never observed in

the control testes (**Fig.7A,B**).

Importantly, functional analysis of the genes differentially expressed between

control and NR5A1-deficient SC highlighted extracellular matrix (ECM; 74 genes, *p*

value$<2.6^{e-16}$), cell adhesion (118 genes, *p* value$<5.9^{e-9}$), basement membrane (26

genes, *p* value$<2.1^{e-8}$), as well as integrin binding (25 genes, *p* value$<2.5^{e-6}$)

(**Suppl.Fig.4; Table.S2**). Accordingly, the expression level of some genes involved in

cell junction/adhesion (e.g., *Gja1*, *Jam2*, *Ptk2b*) and integrin signalling (e.g., *Itga6*)

were down-regulated. As to genes for ECM components, some were down (e.g.,

Col4a1, *Col4a2*), while many others (e.g., *Col3a1*, *Col5a2*, *Col6a6*, *Fn1*, *Fbln1*, *Mdk*,

Mgp) were up-regulated (**Fig.7I**). To verify this point, we analysed collagen type IV

(COL-IV) expression by IHC. At E14.5, COL-IV was surrounding the whole periphery

of all cords in control testes (yellow edging, **Fig.7C**). In *Nr5a1*$^{SC–/–}$testes, COL-IV was

greatly reduced and even absent at the periphery of many cords (arrows, **Fig.7G**), in

areas corresponding to regions where the expression of SOX9 was also lost

(arrowheads, **Fig.7H**). As *Col4a1* and *Col4a2* are regulated by SOX9 [28], it is

reasonable to propose that loss of SOX9 in NR5A1-deficient SC yielded loss of COL-

IV and altered basement membrane, which, in turn, disorganized the cord structure.

Searching further for ligand-receptor interactions in our data set, we identified a

deregulated network related to integrin-dependent cell adhesion (**Suppl.Fig.8**).

Together our findings indicate that NR5A1-deficient SC lost their appropriate cell-cell

and cell-ECM interactions.

DISCUSSION

Alteration of gene expression in *Nr5a1*$^{SC–/–}$ testes could result from either loss of

NR5A1 or death of SC. The decreased expression of *Sox9*, *Amh*, *Gja1*, and possibly

Cyp26b1 and *Dhh*, is likely directly related to the loss of NR5A1 in SC. Actually,

NR5A1 initiates SC differentiation by directing *Sox9* expression [9]. Then, it maintains

Sox9 transcription and activates downstream genes by binding to *cis*-regulatory

elements (e.g., *Amh*, *Gja1*) or synergizing through unknown mechanisms (e.g.,

Cyp26a1, *Dhh*) [16,29-32]. As to *Dmrt1*, *Sox8* and *Ptgds*, their decreased expression

was also expected because SOX9 regulates their expression [17,31,33,34].

However, our finding that *Ptgds* is undetectable as early as E13.5, while SOX9 is still

present, suggests that NR5A1 may regulate *Ptgds* expression directly (**Fig.8**).

NR5A1-deficient SC transdifferentiate into pre-granulosa-like cells

Based on genetic studies, SOX9 has emerged as the master regulator

instructing SC differentiation during testis development, notably by repressing the

WNT4/RSPO1-dependent pathways involved in pGrC differentiation [1-3]. However,

human fibroblasts can be reprogramed into SC by forcing expression of NR5A1 and

GATA4, but not SOX9 [35]. Here we show that SC lacking NR5A1 from E13.5 loose

expression of genes defining or maintaining SC identity (e.g., *Sox9*, *Sox8*, *Ptgds*,

Dmrt1), further illustrating the primary role of NR5A1 in instructing SC differentiation*.*

In parallel, they gain expression of genes specifying pGrC differentiation (e.g., *Wnt4*,

Rspo1) and their transcriptome projected to a gonad atlas clearly shows that they

display an identity close to that of pGrC. Nonetheless, these "pGrC-like" cannot reach

a functional state of differentiation since they keep FOXL2-negative. Another study

has shown that SC lacking *Nr5a1* from E12.5 reach a functional, FOXL2-positive,

granulosa cell identity, resulting in complete male-to-female sex reversal [36]. This

indicates that SC devoid of NR5A1 from E12.5 can transdifferentiate into functional

pGrC, while those loosing NR5A1 from E13.5 are no longer licensed to do so. We

further show that these "pGrC-like" subsequently die through a *Trp53*-independent

mechanism since concomitantly removing *Trp53* and *Nr5a1* does not prevent death.

Quite interestingly, SC loosing *Nr5a1* later than E14.5 either die by MDM2/TRP53-

dependent apoptosis or survive, since the resulting testis at adulthood still contains

clearly defined seminiferous tubules [37]. Thus, the sooner NR5A1 is lost in SC, the

highest their plasticity remains, suggesting that NR5A1 progressively locks the SC

identity over time. In addition, SC die by distinct mechanisms, depending on the

stage at which NR5A1 is lost.

304   Sertoli cells lacking NR5A1 from E13.5 die by anoikis

It is accepted that cell adhesion/junction molecules and ECM components play

important roles in maintaining the integrity of seminiferous cords [38-40]. Because

many genes deregulated in NR5A1-deficient SC are coding for such molecules,

including collagen-coding genes, testis cord disruption is likely linked to breakdown of

SC adhesion to their environment. Interestingly enough, some of the upregulated

genes coding for ECM components such as *Col18a1*, *Fn1*, *Nid2* and *Dcn* (**Table.S2**)

are characteristics of pGrC [41].

Cells sense their location and maintain their adhesion through specific

interactions with the ECM. This plays a major role in regulating various processes,

including cell-survival. This function is mediated by integrins, which are the cell-

membrane receptors interacting with components of the ECM. They recruit

nonreceptor tyrosine focal adhesion kinases (FAK), that are activated in response to

adhesion [42]. Therefore, disruption or loss of integrin-ECM adhesion impairs cell-

survival, and often causes detachment-induced cell death, called anoikis [43,44]. We

found that expression of *Itga6*, an integrin subunit crucial for fetal testis organization

[45], and *Ptk2b*, a member of the FAK family expressed in SC in a SOX9-dependent

manner [46] is reduced in NR5A1-deficient SC. Thus, it is conceivable that these

changes may cause SC to lose their anchorage to the modified ECM (see above)

and initiate death of detaching SC by anoikis, altering thereby integrity of the testis

cord epithelium (**Fig.8**). It is worth noting that anoikis still occurs in the absence of

TRP53 [47], which is fully consistent with our finding that NR5A1-deficient SC die

independently of TRP53.

NR5A1-deficient SC and differentiation of the other testicular cell-types

As to the GC fate, the data show that most of them become meiotic and die in

Nr5a1[SC−/−] testes. The fact that expression of *Cyp26b1* is lost in NR5A1-deficient SC

is sufficient to explain why GC entered meiosis (**Fig.8**). Actually, CYP26B1 is

required in SC to prevent meiotic initiation and to maintain GC in an undifferentiated

state [48].

As to FLC, their specification and/or development rely on SC, notably thanks to

DHH- and PDGF-signalling pathways [49,50]. Our finding that FLC are present in

Nr5a1[SC−/−] testes, despite decreased expression of *Dhh* and *Pdgfa* in SC, reveals

that these pathways are no longer required beyond E13.5 to allow proper emergence

or survival of FLC, as observed when SC are ablated after E15.5 [51,52].

Furthermore, these FLC appear functional since testicular descent is occurring in

Nr5a1[SC−/−] mutants, suggesting normal INSL3 production [53].

As to peritubular myoid cell (PTMC), it is admitted that SC are required during

the fetal period to support their differentiation, notably by producing DHH and ECM.

Genetic ablation of SC from E14.5 or *Dhh*-null mutations actually induce seminiferous

cords defects closely resembling those observed in *Nr5a1*[SC–/–] testes [54-56]. Thus,

the decreased expression of *Dhh* in NR5A1-deficient SC and their death questions

the fate of PTMC. As a matter of fact, we could not distinguish PTMC from the

interstitial or coelomic epithelium cell clusters. This is not really surprising as PTMC,

sparse at E14.5, become obvious only from E16.5 [57], i.e., 2 days later than the

stage at which we performed analysis. It is nonetheless worth noting that expression

of *Ptch1*, *Tagln*, *Acta2* and *Tpm1*, which are generally considered as hallmarks of

PTMC [58,59], were significantly reduced in interstitial or coelomic epithelium cluster

cells (**Table.S2**). It is therefore possible that PTMC, included in these clusters, are

altered, or even lost, following ablation of *Nr5a1* in SC. Interestingly, both PTMC and

"dormant" steroidogenic progenitors (SP) at the origin of adult LC stem from a single

population of *Wnt5a*+ SP [60]. In this context, it is conceivable that the absence of

adult LC in PND60 testis is related to an alteration of PTMC during fetal

development. Further investigations are required to trace the fate of these *Wnt5a*+

SP in *Nr5a1*[SC–/–] testis.

Death of the NR5A1-deficient SC affects genital tract development

Müllerian duct regression in males proceeds rostro-caudally, and is dependent

on the action of AMH, which begins as early as E13.5 [61,62]. As *Amh* expression in

Nr5a1[SC–/–] mutants is lost only from E14.5, AMH production starts normally and then

gradually decreases to zero after E14.5. It is therefore not surprising that the rostral

part of the Müllerian ducts regresses normally, making their derivatives (oviducts,

anterior portions of the uterine horns absent in the mutants. On the other hand, it is

reasonable to propose that the caudal part of the Mullerian ducts persists beyond

E14.5, and is at the origin of the posterior portions of the uterine horns, body of the

uterus and the vagina of adult *Nr5a1*[SC–/–] mutants (**Fig.8**).

In addition, *Nr5a1*[SC−/−] mutants display normal Wolffian duct-derived genitalia

(epididymis, vas deferens, seminal vesicles) [63]. These develop under the influence

of testosterone, the synthesis of which starts as early as E13.5 in mice [64]. Their

presence indicates therefore that *Nr5a1*[SC−/−] mutants were exposed to testosterone

during fetal development. This is surprising since SC, converting FLC-produced

androstenedione into testosterone thanks to expression of *Hsd17b1* and *Hsd17b3*

[65], are progressively lost in *Nr5a1*[SC−/−] mutants. Thus, another enzyme necessarily

compensates for the loss of *Hsd17b1* and *Hsd17b3*, as considered elsewhere [66]. In

this regard, the HSD17B12 is a good candidate since it can convert androstenedione

to testosterone [67]. However, the fact that FLC are not able to produce testosterone

from androstenedione [65] means one has to consider HSD17B12 allows

testosterone synthesis in a cell-type other than FLC. This production is nevertheless

insufficient since AGD and penile bone length, both of which are highly sensitive to

testosterone levels between E14.5 and E17.5 in mice [68], are reduced in *Nr5a1*[SC−/−]

mutants (**Fig.8**).

Our study shows that SC lacking NR5A1 from E13.5 transiently

transdifferentiate to acquire a pre-granulosa cell-like identity, profoundly modify the

landscape of the adhesion molecules and ECM they express and die by a TRP53-

independent mechanism related to anoikis. This yields a disorganization of the testis

cords, making GC to prematurely enter meiosis instead of becoming quiescent.

Interestingly, the maintenance of fetal LC is not altered in the *Nr5a1*[SC−/−] mutant, but

the emergence of adult LC is clearly compromised. Further studies are now required

to understand the reason for the absence of adult LC in the postnatal testis.

MATERIALS AND METHODS

Mice

They were on a mixed C57BL/6 (50%)/129/SvPass (50%) genetic background.

The *Nr5a1* conditional mutant mouse line was established at the Institut Clinique de

la Souris (*i*CS, Illkirch, France), in the context of the French National Infrastructure for

Mouse Phenogenomics PHENOMIN (http://www.phenomin.fr). To construct the

targeting vector a 1.9 kb-long DNA fragment encompassing exon 7

(ENSMUSE00000 693512) was amplified by PCR using 129/SvPass genomic DNA

and cloned into an *i*CS proprietary vector containing a *lox*P site, as well as a *lox*P-

and *FRT*-flanked neomycin resistance cassette (step1 plasmid). Then, 3 kb- and

3.7 kb-long fragments corresponding to 5' and 3' homology arms were amplified by

PCR and introduced into step1 plasmid to generate the targeting construct. This

linearized construct was electroporated into 129/SvPass mouse embryonic stem (ES)

cells. After selection, targeted clones were identified by PCR using external primers

and confirmed by Southern blots (5' and 3' digests) hybridized with neomycin, 5' and

3' external probes. One positive ES clone was injected into C57BL/6J blastocysts. To

remove the selection cassette from the *Nr5a1* locus, chimeric males were crossed

with *Gt(ROSA)26Sor*$^{tm1(FLP1)Dym}$ females [69]. Germline transmission was obtained,

and a further breeding step was needed to segregate animals bearing the *Nr5a1* L2

allele from animals bearing the transgene. To inactivate *Nr5a1* in SC, female mice

bearing *Plekha5*$^{Tg(AMH-cre)1Flor}$ [14] and *Gt(ROSA)26Sor*$^{tm1(EYFP)Cos}$ [15] transgenes,

and heterozygous for the L2 allele of *Nr5a1* were mated with males heterozygous or

homozygous for L2 alleles of *Nr5a1*. The resulting *Plekha5*$^{Tg(AMH-cre)1Flor}$;*Nr5a1*$^{+/+}$;

Gt(ROSA)26Sor$^{tm1(EYFP)Cos}$ and *Plekha5*$^{Tg(AMH-cre)1Flor}$;*Nr5a1*$^{L2/L2}$; *Gt(ROSA)26Sor*$^{tm1(EYFP)Cos}$

males are referred to as control and *Nr5a1*$^{SC-/-}$ mutant fetuses, respectively. To test

for the role of TRP53, the L2 allele of *Trp53* gene [21] was further introduced in the

mice described above.

Noon of the day of a vaginal plug was taken as 0.5 day embryonic development

(E0.5). All fetuses were collected by caesarean section (no randomization, no blind

experiments). Yolk sacs or tail biopsies were taken for DNA extraction. Primers 5'-

GTCAAGCGCCCCATGAATGC-3' and 5'-TTAGCCCTCCGATGAGGCTG-3' were

first used to amplify *Sry* gene (230 bp-long fragment) for male sex determination.

Then,   primers   5'-TGAGCCCTGGCACATCCCTCC-3'   and   5'-

CCTCTGCCCTGCAGGCTTC TG-3' were used to detect *Plekha5*[Tg(AMH-cre)1Flor]

transgene (273 bp-long amplicon), and primers 5'-AAGGGAGCTGCAGTGGAGTA-3'

and   5'-GCCAGAGGCCACTTGTGTAG-3'   to   detect   *Gt(ROSA)26Sor*[tm1(EYFP)Cos]

reporter (520 bp-long amplicon). Primers 5'-CTGTCTCCTGTCTTCTACTACCCTG-3'

and 5'-AGCCATTTCAACAGTGCCCC TTCC-3' were used to amplify wild-type (+,

290 bp-long)   and   L2   (400 bp-long)   alleles   of   *Nr5a1*,   while   primers   5'-

GTGGCACATGCATTAGTCCACTTGG-3'   and   5'-

AGCCATTTCAACAGTGCCCCTTCC-3' were used to amplify the excised, null, L–

(243 bp-long)   allele.   Primers   5'-CACAAAAACAGGTTAAACCCAG-3'   and   5'-

AGCACATAGGAGGCAGAGAC-3' were used to amplify the wild-type (288 bp-long)

and L2 (370 bp-long) alleles of *Trp53*. Primers 5'-CACAAAAACAGGTTAAACCCAG-

3' and 5'-GAAGACAGAAAAGGGGAG GG-3' were used to amplify the excised, null,

L– (612 bp-long) allele of *Trp53*. The PCR conditions were 30 cycles with

denaturation at 95°C for 30 seconds, annealing at 61°C for 30 seconds and

elongation at 72°C for 30 seconds. The amplicons were resolved on 1.5% (w/v)

agarose gels, stained by ethidium bromide and visualized under UV light, using

standard protocols.

Blood sample collection and testosterone measurement

After anesthesia with a lethal dose of Xylasin and Ketamine as described

above, the blood of 9-11-week-old adult mice was collected into heparinized

Microvette tubes (Sarstedt, Nümbrecht, Germany) by intra-cardiac sampling. The

tubes were then centrifuged at 5000 g for 5 minutes and the resulting plasma

samples were frozen until further use. Testosterone concentrations were determined

by ELISA using a commercially available kit AR E-8000 (LDN, Labor Diagnostika

Nord, Nordhorn, Germany). Statistical significance was further assessed by using

two-tail Student's *t*-tests.

Morphology, histology and immunohistochemistry

Adult mice (PND60) were anesthetized by intraperitoneal injection of a lethal

anesthetic mixture made of Xylasin (3 mg/ml) and Ketamine (20 mg/ml), and tissues

were immediately fixed by intracardiac perfusion of 4% (w/v) paraformaldehyde

(PFA) dissolved in phosphate buffered saline (PBS). Following collection, E12.5-

E15.5 fetuses and tissues were fixed for 16 hours in 4% (w/v) PFA at 4°C or in

Bouin's fluid at 20°C. After removal of the fixative, samples were rinsed in PBS and

placed in 70% (v/v) ethanol for long-term storage, external morphology evaluation

and organ weight measurement. They were next embedded in paraffin and 5 μm-

thick sections were made. For histology, sections were stained with hematoxylin and

eosin (H&E).

For IHC, antigens were retrieved for 1 hour at 95°C either in 10 mM sodium

citrate buffer at pH 6.0 or in Tris-EDTA at pH 9.0 [10 mM Tris Base, 1 mM EDTA,

0.05% (v/v) Tween 20]. Sections were rinsed in PBS, then incubated with appropriate

dilutions of the primary antibodies (**Table.1**) in PBS containing 0.1% (v/v) Tween 20

(PBST) for 16 hours at 4°C in a humidified chamber. After rinsing in PBST (3 times

for 3 minutes each), detection of the bound primary antibodies was achieved for

45 minutes at 20°C in a humidified chamber using Cy3-conjugated or Alexa Fluor

488-conjugated antibodies. Nuclei were counterstained with 4',6-diamidino-2-phenyl-

indole (DAPI) diluted at 10 µg/ml in the mounting medium (Vectashield; Vector

Laboratories, Newark, CA, USA). ImmPRESS® Polymer Detection Kits MP-7500 and

MP-7405 were used according to the manufacturer's protocol (Vector Laboratories).

The surface area occupied by YFP-positive cells was measured using a macro

command designed for Fiji software (Imaging Center of IGBMC). Data were

expressed as percentage of YFP-positive surface areas relative to the entire testis

section surface areas. At least four samples were analyzed per genotype. Statistical

analysis was done by a two-tail Student *t*-test, assuming equal variances after

arcsine transformation of the percentages.

BrdU incorporation and TUNEL assays

BrdU (Sigma-Aldrich, Saint-Quentin-Fallavier, France) dissolved at 5 mg/ml in

PBS was injected intraperitoneally to pregnant females at 50 mg/kg of body weight.

Two hours later, fetuses were collected (E13.5-E15.5), fixed and embedded as

described above. BrdU incorporation was detected on 5 µm-thick sections by using

an anti-BrdU mouse monoclonal antibody (diluted 1:100) and indirect IHC as

described above. At least three samples per stage and per genotype were analyzed.

Data were expressed as percentages of BrdU-positive cells related to the number of

DDX4-positive cells. Statistical analysis was done by a two-tail Student *t*-test,

assuming equal variances after arcsine transformation of the percentages.

TUNEL-positive cells were detected on sections from PFA-fixed samples using

the In Situ Cell Death Detection Kit, Fluorescein, according to the manufacturer's

instructions (Roche, Mannheim, Germany). At least three samples were analyzed per

genotype. Data were expressed as the ratio between the number of TUNEL-positive

cells quantified on entire sections and the surface areas of the testis sections (μm^2).

Statistical significance was assessed by using two tail Student's *t*-tests.

Real-time RT-qPCR analyses of RNA extracted from whole gonad

Fetal testes were dissected, isolated from mesonephros, snap frozen in liquid

nitrogen and stored at –80°C until use. Whole testis total RNA was extracted using

RNeasy Mini Kit (Qiagen, Les Ulis, France). RT-qPCR was performed on 5 ng RNA

aliquots using Luna® Universal One-Step RT-qPCR Kit, according to the

manufacturer's instructions (New England Biolabs, Evry, France). The primers are

listed in **Table.2**. Triplicates of at least four samples were used for each genotype, at

each stage. The relative transcript levels were determined using the ΔΔCt method,

and normalized to *Hprt* whose expression is not affected by ablation of *Nr5a1*.

Purification of YFP-positive Sertoli cells

To dissociate cells, the testes of controls and mutants additionally bearing the

Gt(ROSA)26Sor[tm1(EYFP)Cos] transgene were incubated for 10 minutes at 37°C in

350 µl of trypsin/EDTA 0.05% (w/v), phenol red solution (Gibco Invitrogen, Auckland,

New-Zealand), filtered through a 40 µm cell strainer to generate single cell

suspensions, centrifuged at 3000g and suspended in 300 µl PBS as described [70].

The YFP-positive and -negative cells were sorted separately by FACS using an

Aria® II flow cytometer (BD Biosciences, Le Pont de Claix, France). Sorted cell

suspensions were then lysed for overnight at 55°C in a proteinase K-containing

buffer, DNA was extracted and genotyped by PCR using standard protocols and

primers as indicated above.

Single cell RNA sequencing and data processing

Gonads from E13.5 and E14.5 control and mutant fetuses were dissected out in

PBS, sexed by their appearance under the microscope and cell suspensions were

prepared as described above. Cell number and viability were determined by a Trypan

Blue exclusion assay on a Neubauer Chamber. Samples consisting of > 90% viable

cells were processed on the Chromium Controller from 10X Genomics (Leiden, The

Netherlands). Ten thousand cells were loaded per well to yield approximately 5000 to

6000 captured cells into nanoliter-scale Gel Beads-in-Emulsion (GEMs). Single cell 3'

mRNA sequencing libraries were generated according to Chromium Single Cell 3'

Reagent Kits User Guide (v3 Chemistry) from 10X Genomics. Briefly, GEMs were

generated by combining barcoded gel beads, a reverse transcription master mix

containing cells, and partitioning oil onto Chromium Chip B. Following full length

cDNA synthesis and barcoding from poly-adenylated mRNA, GEM were broken and

pooled before cDNA amplification by PCR using 11 cycles. After enzymatic

fragmentation and size selection, sequencing libraries were constructed by adding

Illumina P5 and P7 primers (Illumina, Evry, France), as well as sample index via end

repair, A tailing, adaptor ligation and PCR with 10 cycles. Library quantifications and

quality controls were determined using Bioanalyzer 2100 (Agilent Technologies,

Santa Clara, CA, USA). Libraries were then sequenced on Illumina HiSeq 4000^{TM} as

100 bases paired-end reads, using standard protocols.

Sequencing   data   were   processed   using   10X   Genomics   software

(https://support.10xgenomics.com/single-cell-gene-expression/software/overview/welcome).

Fastq files were processed with CellRanger (v6.0) on a chimeric genome composed

of mm10 *Mus musculus* assembly and 753 bp-long YFP sequence (European

Nucleotide Archive accession number AGM20711) [71]. The resulting count matrices

were aggregated with the Read10X function implemented in Seurat (v4.0.1) [72].

Cells with less than 200 detected genes and genes detected in less than 10 cells

were removed. Doublets were filtered out independently in each individual matrix by

using the DoubletFinder R package (v.2.0.2) [73]. Subsequently, the Single-Cell

Analysis Toolkit for Gene Expression Data in R (scater v1.10.1) was used to remove

outlier cells by using several cell features including the proportions of reads mapping

mitochondrial and ribosomal genes, the number of genes and UMIs per cell [74].

Data were normalized using the NormalizeData and the SCT function implemented

into Seurat by regressing out the unwanted variation due to cell cycle and to the

proportion of mitochondrial genes. The top-3000 most varying genes were used to

perform a principal component analysis with the RunPCA function implemented in

Seurat. Cells were then clustered by using Seurat graph-based clustering

(FindNeighbors and FindClusters functions) on the top-30 principal components, with

default parameters. Finally, we used the Uniform Manifold Approximation and

Projection (UMAP) method (RunUMAP function) to project cells in a 2D space. Cell

clusters were annotated using a set of known marker genes. The FindAllMarkers

function implemented in Seurat was used to identify significantly differentially

expressed genes (DEGs) between cell clusters. Gene ontology (GO) and pathway

enrichment analysis was conducted for each gene expression cluster using the

AMEN suite of tools [75] with an BH-adjusted p value of ≤ 0.05. The current dataset

was mapped on top of a reference atlas of gonadal development [25] using

FindTransferAnchors and MapQuery functions with default settings. To address this

issue, the reference atlas was first processed by using the same pipeline.

ACKNOWLEDGEMENTS

The *Nr5a1* mouse mutant line was established at the Institut Clinique de la Souris (PHENOMIN *i*CS) in the Genetic Engineering and Model Validation Department with funds from the ANR (see below). We thank Pr Anne-Marie LEFRANCOIS-MARTINEZ for discussions and advices. We also thank Dr Christelle Thibault-CARPENTIER from the Genomeast platform for her input (http://genomeast.igbmc.fr/). This research was funded by Agence Nationale pour la Recherche Grants (see below). A CC-BY-4.0 public copyright license has been applied by the authors to the present document and will be applied to all subsequent versions up to the Author Accepted Manuscript arising from this submission, in accordance with the grant's open access conditions.

FUNDING

This work was supported by grants from Fondation pour la Recherche Médicale (FRM FDT201904008001), CNRS, INSERM, UNISTRA and Agence Nationale pour la Recherche project ARESSERC (ANR-16-CE14-0017). It was also supported in part by the grant ANR-10-LABX-0030-INRT, a French State fund managed by the ANR under the frame Programme Investissements d'Avenir labelled ANR-10-IDEX-0002-02.

AUTHOR INFORMATION

These authors contributed equally: Frédéric Chalmel, Norbert B. Ghyselinck.

Contributions

SSC, MM and NBG conceptualized and designed the study. SSC and DC performed all the experiments. BF and MK provided technical and material support. EG contributed to image analysis. SSC, VA, MC, MJ and FC performed the scRNA-

seq experiment and analysed data. CM and SN provided datasets for the mouse

gonad atlas. SSC, NV, MM, FC and NBG analysed all experiments, managed the

overall study, wrote the manuscript, and supervised its preparation and submission.

NV, MM, FC and NBG acquired funding for this research.

Corresponding authors

Correspondence to Norbert B. Ghyselinck.

ETHICS DECLARATIONS

Competing interests

The authors declare no competing interests.

Ethics statement

Mice were housed in a licensed animal facility (agreement #C6721837). All

experiments were approved by the local ethical committee (Com'Eth, accreditation

APAFIS#18323- 2018113015272439_v3), and were supervised by N.B.G., M.M. and

N.V., who are qualified in compliance with the European Community guidelines for

laboratory animal care and use (2010/63/UE).

DATA AVAILABILITY

The data sets described in this paper are available at Gene Expression

Omnibus (GEO) database under the accession number GSE219271.

SUPPLEMENTARY INFORMATION

Supplementary information is available at *Cell Death and Differentiation*'s

website.

REFERENCES

1.   Yao HH, Ungewitter E, Franco H & Capel B. Establishment of fetal Sertoli cells

and their role in testis morphogenesis. In: Griswold MD (ed). Sertoli Cell Biology.

2nd edn.   (Academic   Press,   Cambridge,   2015)   57-79.

https://doi.org/10.1016/B978-0-12-417047-6.00002-8

2.   Rotgers E, Jørgensen A & Yao HH. At the Crossroads of Fate-Somatic Cell

Lineage Specification in the Fetal Gonad. *Endocr Rev* **39**, 739-759 (2018)

PMID:29771299

3.   Nef S, Stévant I & Greenfield A. Characterizing the bipotential mammalian

gonad. *Curr Top Dev Biol* **134**, 167-194 (2019) PMID:30999975

4.   Eggers S, Ohnesorg T & Sinclair A. Genetic regulation of mammalian gonad

development. *Nat Rev Endocrinol* **10**, 673-683 (2014) PMID:25246082

5.   Fabbri-Scallet H, de Sousa LM, Maciel-Guerra AT, Guerra-Júnior G & de Mello

MP. Mutation update for the NR5A1 gene involved in DSD and infertility. *Hum*

Mutat **41**, 58-68 (2020) PMID:31513305

6.   Morohashi K, Baba T & Tanaka M. Steroid hormones and the development of

reproductive organs. *Sex Dev* **7**, 61-79 (2013) PMID:22986257

7.   Morohashi KI, Inoue M & Baba T. Coordination of Multiple Cellular Processes by

NR5A1/Nr5a1. *Endocrinol Metab (Seoul)* **35**, 756-764 (2020) PMID:33397036

8.   Luo X, Ikeda Y & Parker KL. A cell-specific nuclear receptor is essential for

adrenal and gonadal development and sexual differentiation. *Cell* **77**, 481-490

(1994) PMID:8187173

9.   Sekido R & Lovell-Badge R. Sex determination involves synergistic action of

SRY and SF1 on a specific Sox9 enhancer. *Nature* **453**, 930-934 (2008)

PMID:18454134

10. Cool J, DeFalco T & Capel B. Testis formation in the fetal mouse: dynamic and

complex de novo tubulogenesis. *Wiley Interdiscip. Rev Dev Biol* **1**, 847-859

(2012) PMID:23799626

11. Griswold MD. 50 years of spermatogenesis: Sertoli cells and their interactions

with germ cells. *Biol Reprod* **99**, 87-100 (2018) PMID:29462262

12. Bitgood MJ, Shen L & McMahon AP. Sertoli cell signaling by Desert hedgehog

regulates the male germline. *Curr Biol* **6**, 298-304 (1996) PMID:8805249

13. Josso N & Picard JY. Genetics of anti-Müllerian hormone and its signaling

pathway. *Best Pract Res Clin Endocrinol Metab* **36**, 101634 (2022)

PMID:35249805

14. Lécureuil C, Fontaine I, Crepieux P & Guillou F. Sertoli and granulosa cell-

specific Cre recombinase activity in transgenic mice. *Genesis* **33**, 114-118 (2002)

PMID:12124943

15. Srinivas S, Watanabe T, Lin CS, William CM, Tanabe Y, Jessell TM *et al.* Cre

reporter strains produced by targeted insertion of EYFP and ECFP into the

ROSA26 locus. *BMC Dev Biol* **1**, 4 (2001) PMID:11299042

16. De Santa Barbara P, Bonneaud N, Boizet B, Desclozeaux M, Moniot B, Sudbeck

P *et al.* Direct interaction of SRY-related protein SOX9 and steroidogenic factor

1 regulates transcription of the human anti-Müllerian hormone gene. *Mol Cell Biol*

18, 6653-6665 (1998) PMID:9774680

17. Wilhelm D, Hiramatsu R, Mizusaki H, Widjaja L, Combes AN, Kanai Y *et al.*

SOX9 regulates prostaglandin D synthase gene transcription in vivo to ensure

testis development. *J Biol Chem* **282**, 10553-10560 (2007) PMID:17277314

18. Malki S, Nef S, Notarnicola C, Thevenet L, Gasca S, Méjean C *et al*.

Prostaglandin D2 induces nuclear import of the sex-determining factor SOX9 via

its cAMP-PKA phosphorylation. *EMBO J* **24**, 1798-1809 (2005) PMID:15889150

19. Tremblay JJ & Viger RS. Transcription factor GATA-4 enhances Müllerian

inhibiting substance gene transcription through a direct interaction with the

nuclear receptor SF-1. *Mol Endocrinol* **13**, 1388-1401 (1999) PMID:10446911

20. Buganim Y, Itskovich E, Hu YC, Cheng AW, Ganz K, Sarkar S *et al*. Direct

reprogramming of fibroblasts into embryonic Sertoli-like cells by defined factors.

Cell Stem Cell **11**, 373-386 (2012) PMID:22958931

21. Jonkers J, Meuwissen R, van der Gulden H, Peterse H, van der Valk M & Berns

A. Synergistic tumor suppressor activity of BRCA2 and p53 in a conditional

mouse model for breast cancer. *Nat Genet* **29**, 418-425 (2001) PMID:11694875

22. Sararols P, Stévant I, Neirijnck Y, Rebourcet D, Darbey A, Curley MK *et al*.

Specific Transcriptomic Signatures and Dual Regulation of Steroidogenesis

Between Fetal and Adult Mouse Leydig Cells. *Front Cell Dev Biol* **9**, 695546

(2021) PMID:34262907

23. Welsh M, Suzuki H & Yamada G. The masculinization programming window.

Endocr Dev **27**, 17-27 (2014) PMID:25247641

24. Kulibin AY & Malolina EA. Formation of the rete testis during mouse embryonic

development. *Dev Dyn* **249**, 1486-1499 (2020) PMID:32852840

25. Mayère C, Regard V, Perea-Gomez A, Bunce C, Neirijnck Y, Djari C *et al*.

Origin, specification and differentiation of a rare supporting-like lineage in the

developing mouse gonad. *Sci Adv* **8**, eabm0972 (2022) PMID:35613264

26. Teletin M, Vernet N, Yu J, Klopfenstein M, Jones JW, Féret B *et al*. Two
functionally redundant sources of retinoic acid secure spermatogonia
differentiation in the seminiferous epithelium. *Development* **146**, dev170225
(2019) PMID:30487180

27. Chassot AA, Gillot I & Chaboissier MC. R-spondin1, WNT4, and the CTNNB1
signaling pathway: strict control over ovarian differentiation. *Reproduction* **148**,
R97-110 (2014) PMID:25187620

28. Sumi E, Iehara N, Akiyama H, Matsubara T, Mima A, Kanamori H *et al*. SRY-
related HMG box 9 regulates the expression of Col4a2 through transactivating its
enhancer element in mesangial cells. *Am J Pathol* **170**, 1854-1864 (2007)
PMID:17525254

29. Arango NA, Lovell-Badge R & Behringer RR. Targeted mutagenesis of the
endogenous mouse Mis gene promoter: in vivo definition of genetic pathways of
vertebrate sexual development. *Cell* **99**, 409-419 (1999) PMID:10571183

30. Kashimada K, Svingen T, Feng CW, Pelosi E, Bagheri-Fam S, Harley VR *et al*.
Antagonistic regulation of Cyp26b1 by transcription factors SOX9/SF1 and
FOXL2 during gonadal development in mice. *FASEB J* **25**, 3561-3569 (2011)
PMID:21757499

31. Li Y, Zheng M & Lau YF. The sex-determining factors SRY and SOX9 regulate
similar target genes and promote testis cord formation during testicular
differentiation. *Cell Rep* **8**, 723-733 (2014) PMID:25088423

32. Couture R & Martin LJ. The transcription factors SF-1 and SOX8 cooperate to
upregulate Cx43 expression in mouse TM4 sertoli cells. *Biochem Biophys Rep*
24, 100828 (2020) PMID:33088929

33. Chaboissier MC, Kobayashi A, Vidal VI, Lützkendorf S, van de Kant HJ, Wegner

M *et al.* Functional analysis of Sox8 and Sox9 during sex determination in the

mouse. *Development* **131**, 1891-1901 (2004) PMID:15056615

34. Rahmoun M, Lavery R, Laurent-Chaballier S, Bellora N, Philip GK, Rossitto M *et*

al. In mammalian foetal testes, SOX9 regulates expression of its target genes by

binding to genomic regions with conserved signatures. *Nucleic Acids Res* **45**,

7191-7211 (2017) PMID:28472341

35. Liang J, Wang N, He J, Du J, Guo Y, Li L *et al.* Induction of Sertoli-like cells from

human   fibroblasts   by   NR5A1   and   GATA4.   *Elife*   **8**,   e48767   (2019)

PMID:31710289

36. Ikeda Y, Tagami A, Maekawa M & Nagai A. The conditional deletion of

steroidogenic   factor   1   (Nr5a1)   in   Sox9-Cre   mice   compromises   testis

differentiation. Sci Rep **11**, 4486 (2021) PMID:33627800

37. Anamthathmakula P, Miryala CSJ, Moreci RS, Kyathanahalli C, Hassan SS,

Condon JC *et al.* Steroidogenic Factor 1 (Nr5a1) is Required for Sertoli Cell

Survival Post Sex Determination. *Sci Rep* **9**, 4452 (2019) PMID:30872705

38. Barrionuevo F, Georg I, Scherthan H, Lécureuil C, Guillou F, Wegner M *et al.*

Testis cord differentiation after the sex determination stage is independent of

Sox9 but fails in the combined absence of Sox9 and Sox8. *Dev Biol* **327**, 301-312

(2009) PMID:19124014

39. Georg I, Barrionuevo F, Wiech T & Scherer G. Sox9 and Sox8 are required for

basal lamina integrity of testis cords and for suppression of FOXL2 during

embryonic   testis   development   in   mice.   *Biol   Reprod*   **87**,   99   (2012)

PMID:22837482

40. Chen SR, Chen M, Wang XN, Zhang J, Wen Q, Ji SY *et al*. The Wilms tumor

gene, Wt1, maintains testicular cord integrity by regulating the expression of

Col4a1 and Col4a2. *Biol Reprod* **88**, 56 (2013) PMID:23325811

41. Piprek RP, Kolasa M, Podkowa D, Kloc M & Kubiak JZ. Transcriptional profiling

validates involvement of extracellular matrix and proteinases genes in mouse

gonad development. *Mech Dev* **149**, 9-19 (2018) PMID:29129619

42. Lu Q & Rounds S. Focal adhesion kinase and endothelial cell apoptosis.

Microvasc Res **83**, 56-63 (2012) PMID:21624380

43. Gilmore AP. Anoikis. *Cell Death Differ* **12 Suppl 2**, 1473-1477 (2005)

PMID:16247493

44. Vachon PH. Integrin signaling, cell survival, and anoikis: distinctions, differences,

and differentiation. *J Signal Transduct* **2011**, 738137 (2011) PMID:21785723

45. Fröjdman K & Pelliniemi LJ. Differential distribution of the alpha 6 subunit of

integrins in the development and sexual differentiation of the mouse testis.

Differentiation **57**, 21-29 (1994) PMID:8070619

46. Beverdam A, Svingen T, Bagheri-Fam S, McClive P, Sinclair AH, Harley VR *et al*.

Protein tyrosine kinase 2 beta (PTK2B), but not focal adhesion kinase (FAK), is

expressed in a sexually dimorphic pattern in developing mouse gonads. *Dev Dyn*

239, 2735-2741 (2010) PMID:20737507

47. McGill G, Shimamura A, Bates RC, Savage RE & Fisher DE. Loss of matrix

adhesion triggers rapid transformation-selective apoptosis in fibroblasts. *J Cell*

Biol **138**, 901-911 (1997) PMID:9265655

48. Li H, MacLean G, Cameron D, Clagett-Dame M & Petkovich M. Cyp26b1

expression in murine Sertoli cells is required to maintain male germ cells in an

undifferentiated state during embryogenesis. *PLoS One* **4**, e7501 (2009)

PMID:19838304

49. Yao HH, Whoriskey W & Capel B. Desert Hedgehog/Patched 1 signaling

specifies fetal Leydig cell fate in testis organogenesis. *Genes Dev* **16**, 1433-1440

(2002) PMID:12050120

50. Brennan J, Tilmann C & Capel B. Pdgfr-alpha mediates testis cord organization

and fetal Leydig cell development in the XY gonad. *Genes Dev* **17**, 800-810

(2003) PMID:1265189

51. Rebourcet D, O'Shaughnessy PJ, Pitetti JL, Monteiro A, O'Hara L, Milne L *et al.*

Sertoli cells control peritubular myoid cell fate and support adult Leydig cell

development in the prepubertal testis. *Development* **141**, 2139-2149 (2014)

PMID:24803659

52. Fouchécourt S, Livera G, Messiaen S, Fumel B, Parent AS, Marine JC *et al.*

Apoptosis of Sertoli cells after conditional ablation of murine double minute 2

(Mdm2) gene is p53-dependent and results in male sterility. *Cell Death Differ* **23**,

521-530 (2016) PMID:26470726

53. Verma-Kurvari S, Nef S & Parada LF. Hormonal regulation of male reproductive

tract development. *Ann N Y Acad Sci* **1061**, 1-8 (2005) PMID:16467252

54. Clark AM, Garland KK & Russell LD. Desert hedgehog (Dhh) gene is required in

the mouse testis for formation of adult-type Leydig cells and normal development

of peritubular cells and seminiferous tubules. *Biol Reprod* **63**, 1825-1838 (2000)

PMID:11090455

55. Pierucci-Alves F, Clark AM & Russell LD. A developmental study of the Desert

hedgehog-null mouse testis. *Biol Reprod* **65**, 1392-1402 (2001) PMID:1167325

56. Wang YQ, Cheng JM, Wen Q, Tang JX, Li J, Chen SR *et al.* An exploration of the

role of Sertoli cells on fetal testis development using cell ablation strategy. *Mol*

Reprod Dev **87**, 223-230 (2020) PMID:32011766

57. Aksel S, Cao M, Derpinghaus A, Baskin LS & Cunha GR. Ontogeny of mouse

Sertoli, Leydig and peritubular myoid cells from embryonic day 10 to adulthood.

Differentiation **Mar7**, S0301-4681(22)00015-9 (2022) PMID:35317954

58. Jeanes A, Wilhelm D, Wilson MJ, Bowles J, McClive PJ, Sinclair AH *et al.*

Evaluation of candidate markers for the peritubular myoid cell lineage in the

developing mouse testis. *Reproduction* **130**, 509-516 (2005) PMID:16183868

59. Sohni A, Tan K, Song HW, Burow D, de Rooij DG, Laurent L *et al.* The Neonatal

and Adult Human Testis Defined at the Single-Cell Level. *Cell Rep* **26**, 1501-

1517.e4 (2019). PMID:30726734

60. Ademi H, Djari C, Mayère C, Neirijnck Y, Sararols P, Rands CM *et al.*

Deciphering the origins and fates of steroidogenic lineages in the mouse testis.

Cell Rep **39**, 110935 (2022). PMID:35705036

61.Staack A, Donjacour AA, Brody J, Cunha GR & Carroll P. Mouse urogenital

development: a practical approach. *Differentiation* **71**, 402-413 (2003)

PMID:12969333

62. Mullen RD & Behringer RR. Molecular genetics of Müllerian duct formation,

regression and differentiation. *Sex Dev* **8**, 281-296 (2014) PMID:25033758

63. Zhao F & Yao HH. A tale of two tracts: history, current advances, and future

directions of research on sexual differentiation of reproductive tracts. *Biol Reprod*

101, 602-616 (2019) PMID:31058957

64. Livera G, Delbes G, Pairault C, Rouiller-Fabre V & Habert R. Organotypic culture,

a powerful model for studying rat and mouse fetal testis development. *Cell Tissue*

Res **324**, 507-521 (2006). PMID:16520975

65. Shima Y, Miyabayashi K, Haraguchi S, Arakawa T, Otake H, Baba T *et al*.

Contribution of Leydig and Sertoli cells to testosterone production in mouse fetal

testes. *Mol Endocrinol* **27**, 63-73 (2013) PMID:23125070

66. Rebourcet D, Mackay R, Darbey A, Curley MK, Jørgensen A, Frederiksen H *et*

al. Ablation of the canonical testosterone production pathway via knockout of the

steroidogenic enzyme HSD17B3, reveals a novel mechanism of testicular

testosterone production. *FASEB J* **34**, 10373-10386 (2020) PMID:32557858

67. Blanchard PG & Luu-The V. Differential androgen and estrogen substrates

specificity in the mouse and primates type 12 17beta-hydroxysteroid

dehydrogenase. *J Endocrinol* **194**, 449-455 (2007) PMID:17641292

68. Welsh M, Suzuki H & Yamada G. The masculinization programming window.

Endocr Dev **27**, 17-27 (2014) PMID:25247641

69. Farley FW, Soriano P, Steffen LS, Dymecki SM.   Widespread recombinase

expression using FLPeR (flipper) mice. *Genesis* **28**, 106-110 (2000) PMID:

11105051

70. Stévant I, Neirijnck Y, Borel C, Escoffier J, Smith LB, Antonarakis SE *et al*.

Deciphering Cell Lineage Specification during Male Sex Determination with

Single-Cell RNA Sequencing. *Cell Rep* **22**, 1589-1599 (2018) PMID:29425512

71. Aliye N, Fabbretti A, Lupidi G, Tsekoa T & Spurio R. Engineering color variants

of green fluorescent protein (GFP) for thermostability, pH-sensitivity, and

improved folding kinetics. *Appl Microbiol Biotechnol* **99**, 1205-1216 (2015)

PMID:25112226.

72. Hao Y, Hao S, Andersen-Nissen E, Mauck WM 3rd, Zheng S, Butler A *et al.*

Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 (2021)

PMID: 34062119

73. McGinnis CS, Murrow LM & Gartner ZJ. DoubletFinder: Doublet Detection in

Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**,

329-337 (2019) PMID:30954475

74. McCarthy DJ, Campbell KR, Lun AT & Wills QF. Scater: pre-processing, quality

control, normalization and visualization of single-cell RNA-seq data in R.

Bioinformatics **33**, 1179-1186 (2017). PMID:28088763

75. Chalmel F & Primig M. The Annotation, Mapping, Expression and Network

(AMEN) suite of tools for molecular systems biology. *BMC Bioinformatics* **9**, 86

(2008) PMID:18254954

76. Best D, Sahlender DA, Walther N, Peden AA & Adams IR. Sdmg1 is a

conserved transmembrane protein associated with germ cell sex determination

and germline-soma interactions in mice. *Development* **135**, 1415-1425 (2008)

PMID:18321981

FIGURE LEGENDS

Figure 1. Ablation of *Nr5a1* is efficient from E13.5 onwards. (A-F) Detection of

NR5A1 (red nuclear signals), YFP (green cytoplasmic signal), and HSD3B1 (red

cytoplasmic signal) by IHC on transverse histological sections of the testis of a

control (A,C,E) and a *Nr5a1*$^{SC-/-}$ mutant fetus (B,D,F) at E12.5 (A,B) and E13.5 (C-

F). Efficient excision of the reporter transgene by cre is assessed by YFP expression

in virtually all SC (arrows) as early as E12.5 (A,B). However, loss of NR5A1 in SC is

only achieved at E13.5 (compare C with D). At this stage, expression of NR5A1 is

maintained in Leydig cells (arrowheads in C,D), as identified on consecutive sections

by their expression of HSD3B1 (E,F). Nuclei are counterstained with DAPI (blue

signal). Scale bar (in F): 50 µm.

Figure 2. Ablation of *Nr5a1* in Sertoli cells impairs AMH and SOX9 expression.

(A-R) Detection of AMH, SOX9 and GATA4 (red signals) on transverse histological

sections of testes of control (A,C,E,G,I,K,M,O,Q) and *Nr5a1*$^{SC-/-}$ mutant fetuses

(B,D,F,H,J,L,N,P,R) at E12.5 (A,B,G,H,M,N), E13.5 (C,D,I,J,O,P) and E14.5

(E,F,K,L,Q,R). In (G-R) nuclei are counterstained with DAPI (blue signal). Insets (in

K,L) are high magnifications showing nuclear localisation of SOX9 in control SC

versus cytoplasmic localisation in mutant SC. Note that at each developmental stage,

AMH, SOX9 and GATA4 IHC are performed on consecutive sections. Scale bar (in

R): 50 µm. (**S**) RT-qPCR analyses comparing the expression levels of *Amh*, *Sox9*,

Sox8 and *Ptgds* mRNAs in whole testis RNA from control (n=4) and *Nr5a1*$^{SC-/-}$

mutant (n=4) fetuses at E13.5 and E14.5. Each point represents the mean value of a

technical triplicate, and the bars indicate the mean value.

Figure 3. NR5A1-deficient Sertoli cells die even when they lack TRP53. (A,D)

Detection of YFP (green signal) and TUNEL-positive cells (red signal) on histological

sections from testis of control (A) and *Nr5a1*$^{SC–/–}$ mutant fetuses (B) at E14.5 (A,B)

and E15.5 (C,D). **(E,F)** Detection of GATA4 (red signal) and TUNEL-positive cells

(green signal) on histological sections of control (E) and *Nr5a1*$^{SC–/–}$ mutant testes (F)

fetuses at E14.5. Nuclei are counterstained with DAPI (blue signal). Arrowheads

point to TUNEL-positive SC in the mutant. **(G,H)** Detection of TRP53 (brown signal)

on histological sections of control (G) and *Nr5a1*$^{SC–/–}$ mutant (H) testes at E14.5. The

insets show larger magnifications. Arrowheads and arrows point to SC and GC,

respectively. **(I,J)** Detection of SOX9 (red signal) and YFP (green signal) by IHC on

histological sections of *Nr5a1*$^{SC–/–}$ (I) and *Nr5a1*$^{SC–/–}$;*Trp53*$^{SC–/–}$ (J) mutants at E14.5.

Arrowheads point to SC. **(K-M)** Haematoxylin and eosin staining (H&E) on

histological sections of testes from control (K), *Nr5a1*$^{SC–/–}$ (L) and *Nr5a1*$^{SC–/–}$;*Trp53*$^{SC–}$

$^{/–}$ (M) mutants at PND15. **(N-P)** Detection of HSB3B1 (red signal) on transverse

histological sections of testes from control (N), *Nr5a1*$^{SC–/–}$ (O) and *Nr5a1*$^{SC–/–}$

;*Trp53*$^{SC–/–}$ mutants (P) at PND15. **(Q-S)** Detection of HSB3B1 (red signal) on

transverse histological sections of testes from control (Q), *Nr5a1*$^{SC–/–}$ (R) and

Nr5a1$^{SC–/–}$;*Trp53*$^{SC–/–}$ mutants (s) at PND60. The dotted white lines (in K,N,Q)

delineate seminiferous tubules. Nuclei are counterstained with DAPI (blue signal).

LC, Leydig cells. Scale bar (in S): 25 µm (A-D,I,J), 15 µm (E,F), 100 µm (G,H), 80 µm

(K-P), 20 µm (Q-S).

Figure 4. Abnormal external genitalia and Müllerian duct retention in *Nr5a1*$^{SC–/–}$

males. (A,C) Anogenital distance in control and *Nr5a1*$^{SC–/–}$ males at PND60. **(B,D)**

Penis bones dissected from control and *Nr5a1*$^{SC–/–}$ males at PND60 and stained with

alizarin red and alcian blue. **(E-H)** Reproductive tracts of control (E) and *Nr5a1*$^{SC–/–}$

(F-H) males at PN60. **(I-J)** Histological sections from the *Nr5a1*$^{SC–/–}$ mutant sample

shown in (H) and stained by H&E. The uterine surface consists of a simple columnar

epithelium and the vaginal epithelium is typically stratified squamous. Legend: A,

ampullary gland; CP, cranial prostate; E, epididymis; F, fat pad; SV, seminal vesicle;

T, testis; UB, uterus body; UH, uterine horn; V, vagina; VD, vas deferens; VP, ventral

prostate. Scale bar (in J): 10 μm (I,J).

Figure 5. Transcriptomes of control and *Nr5a1*[SC−/−] testicular cells at a single-

cell resolution. (A) UMAP plot of single cell transcriptomes from control (blue) and

Nr5a1[SC−/−] (pink) testes at E14.5, partitioned into 23 cell clusters (named C1-C23) by

using Seurat graph-based clustering. The proportion of control and mutant cells in

each cell cluster is indicated below as coloured bars. Legend: Legend: coel. ep.,

coelomic epithelium. **(B,C)** Volcano plots of differential gene expression in GC (panel

B, clusters C22 and C23) and SC (panel C, clusters C18, C19 and C21) from control

and *Nr5a1*[SC−/−] testes. Red dots correspond to genes deregulated more than 1.2-fold.

(D-K) Detection of ALDH1A1, TMEM184A (red signals), TRA98, AMH, DDX4 (green

signals) or TUBB3 and PCP4 (brown signals) by IHC on transverse histological

sections of control (D-G) and *Nr5a1*[SC−/−] mutant (H-K) testes at E14.5. In (D,E,H,I)

nuclei are counterstained with DAPI (blue signal). The dotted lines (in D,I) delimit

seminiferous cords. The arrowheads (in G,L) point to SC nuclei. Scale bar (in K):

25 μm (D,E,H,I), 100 μm (F,J), 15 μm (G,K). (**L**) RT-qPCR analyses comparing the

expression levels of selected mRNAs as indicated in whole testis RNA from control

(n=4) and *Nr5a1*[SC−/−] mutant (n=4) fetuses at E14.5. Each point represents the mean

value of a technical triplicate, and the bars indicate the mean value.

Figure 6. NR5A1-deficient Sertoli change their identity. (A) Predicted projection of

control (blue) and *Nr5a1*[SC−/−] (pink) cells on a reference single-cell transcriptomic

atlas of gonad development. **(B,C)** Dot plots representing the predicted association of

cell clusters from the control and *Nr5a1*[SC−/−] single-cell dataset (y axis) to cell-types

(panel B) and to sex and developmental stages (panel C) according to the atlas

previously published [29] (x axis). The dot size represents the prediction score

(ranging from 0.0 to 1.0) of a given cell cluster to be associated with a given cell-type

based on the TransferData function implemented in Seurat. The color intensity (from

white to dark red) indicates the percentage of cells of a given cell cluster that have

been associated with a given cell-type. Legend: coel., coelomic; ctrl, control; ep.,

epithelium; mes., mesenchyme; meso., mesonephros; mut, mutant; prog., progenitor.

(D,E) Detection of FOXL2 (red signal) and DDX4 (green signal) by IHC on transverse

histological sections of control (D) and *Nr5a1*$^{SC-/-}$ mutant (G) testes at E15.5. Nuclei

are counterstained with DAPI (blue signal). The inset (in D) shows detection of

FOXL2 in a E15.5 fetal ovary, used as a positive control for IHC. Scale bar (in E):

15 μm (D,E).

Figure 7. Loss of NR5A1 in Sertoli cells induces disorganisation of the testis

cords. (A,B,E,F) Histological sections through E14.5 (A,E) and E15.5 (B,F) control

(A,B) and *Nr5a1*$^{SC-/-}$ mutant (E,F) fetuses stained by hematoxylin and eosin (H&E).

In controls, the seminiferous cords are well defined (dotted lines). In mutants, they

are poorly defined from E14.5 onwards (broken dotted lines with arrows). **(C,D,G,H)**

Detection of COL-IV, SOX9 (red signals), and YFP (green signal) on transverse

histological sections of the testis from control (C,D) and *Nr5a1*$^{SC-/-}$ mutant fetuses

(G,H) at E14.5. Note that (C,D) and (G,H) are consecutive sections. Arrows (in G)

point to the loss of COL-IV at the periphery of seminiferous cords, where SOX9 is lost

in SC (arrowheads in H). Scale bar (in D): 10 μm (A,B,E,F), 15 μm (C,D,G,H). **(I)**

Tukey box plots illustrating medians, ranges and variabilities of log normalized

expression of the indicated genes involved in either cell junction/adhesion (CJA) or

ECM composition.

Figure 8. Summary of the alterations induced by the loss of NR5A1 in Sertoli

cells after the sex determination period. (A) Control situation. **(B)** *Nr5a1*[SC−/−]

mutant situation. Blue and yellow ovals are up- and down-regulated genes,

respectively. Legend: AGD, anogenital distance; PGD2, prostaglandin D2.

Table 1. Primary antibodies used for immunohistochemistry experiments.

Antibodies	Species	Dilution	Source	Reference
ALDH1A1	Rabbit polyclonal	1/50	Abcam	ab52492
ALDH1A3	Rabbit polyclonal	1/20	Sigma	HPA046271C106318
AMH	Goat polyclonal	1/100	Santa-Cruz	sc-6886
BHMT	Rabbit polyclonal	1/200	Anticorps-enligne.fr	ABIN310161
BrdU	Rat monoclonal	1/100	Bio Rad	OBT0030S
COL-IV	Rabbit polyclonal	1/50	Abcam	ab19808
DDX4	Rabbit polyclonal	1/2000	Abcam	ab13840
FOXL2	Goat polyclonal	1/250	Abcam	ab5096
GATA 4	Goat polyclonal	1/50	Santa-Cruz	sc-1237
H2AFX	Mouse monoclonal	1/500	Millipore	05-636 (JBW301)
HSD3B1	Rabbit polyclonal	1/1000	Trans Genic's Inc	KO607
NR5A1	Rabbit polyclonal	1/100	Cosmo Bio	KAL-KO611
PAX8	Mouse monoclonal	1/10	Abcam	ab53490
PCP4	Rabbit polyclonal	1/500	Novus Biologicals	NBP1-80929
REC8	Rabbit monoclonal	1/200	Abcam	ab192241
SOX8	Mouse monoclonal	1/100	Novus Biologicals	H00030812-M01
SOX9	Rabbit polyclonal	1/1000	Sigma Aldrich	ab5535
SOX10	Goat polyclonal	1/1000	R&D Systems	AF2864
STRA8	Rabbit polyclonal	1/1000	Abcam	ab49602
TMEM184A	Rabbit polyclonal	1/200		[76]
TRA98	Rat monoclonal	1/500	Abcam	ab82527
TRP53	Rabbit polyclonal	1/200	Santa-Cruz	sc-6243
TUBB3	Mouse monoclonal	1/1000	Covance	MMS-435P-0100 (TUJ1 clone)
WT1	Mouse monoclonal	1/500	Cell Marque	348M-94
YFP	Chicken polyclonal	1/500	Aves	GFP-1020

Table 2. Primers used for quantification of mRNA levels by real-time RT-qPCR.

Gene	Accession number	Forward primer sequence	Reverse primer sequence	Amplicon size (bp)
Amh	NM_007445.3	5'-TCCTACATCTGGCTGAAGTGATATG-3'	5'-CAGGTGGAGGCTCTTGGAACT-3'	66
Col6a6	NM_001102607.1	5'-TGATTGCGTTCAGCAACGTG-3'	5'-TCCGGCTTGCATAGATCATAG-3'	157
Cyp26b1	NM_001177713.1	5'-CGACATCCACCGCAACAAG-3'	5'-GGCCTCCTGATATACATTGATGG-3'	151
Dhh	NM_007857.5	5'-GCACAGGATTCACTCCACTACGA-3'	5'-CCAGTGAGTTATCAGCTTTGACC-3'	172
Fst	NM_001301373.1	5'-CTGCTGCTACTCTGCCAGTT-3'	5'-ACATCCTCCTCGGTCCATGA-3'	167
Hsd17b3	NM_008291.3	5'-ATGGAGTCAAGGAGGAAAGGC-3'	5'-ATGGAGTCAAGGAGGAAAGGC-3'	76
Inhbb	NM_008381.4	5'-GCGTCTCCGAGATCATCAGC-3'	5'-CACCTTGACCCGTACCTTCC-3'	188
Itga6	NM_001277970.1	5'-GCGGCTACTTTCACTAAGGACT-3'	5'-TTCTTTTGTTCTACACGGACGA-3'	92
Pcp4	NM_008791.3	5'-GCGACCAACGGAAAAGACAA-3'	5'-TTCAGGTGGACCAGGAAGCA-3'	194
Ptgds	NM_008963.3	5'-GCTCTTCGCATGCTGTGGAT-3'	5'-GCCCCAGGAACTTGTCTTGTT-3'	118
Sox8	NM_011447.3	5'-ACCCGCATCTCCATAACGCA-3'	5'-TGGTGGCCCAGTTCAGTACC-3'	214
Sox9	NM_011448.4	5'-CAGCAAGACTCTGGGCAAG-3'	5'-TCCACGAAGGGTCTCTTCTC-3'	63
Tubb3	NM_023279.3	5'-CGTGAAGTCAGCATGAGGGA-3'	5'-TCCAAGTCCACCAGAATGGC-3'	218

Control

Nr5a1^{SC-/-}

A B C D

SV
CP
VD
VP
T

SV
CP
VD
VD
UH
VP
T

Nr5a1^{SC-/-}

E
F
U
U
T
VD

UH
UB
V
UH
A
Ur
VD

G H

I J