^{UFM}Track: Under-Flow Migration Tracker enabling analysis of the entire multi-step immune cell

extravasation cascade across the blood-brain barrier in microfluidic devices

Mykhailo Vladymyrov[1, 2], Luca Marchetti[1], Sidar Aydin[1, +], Sasha Soldati[1], Adrien Mossu[1], Arindam

Pal[1, Δ], Laurent Gueissaz[1, ◊], Akitaka Ariga[3, □], and Britta Engelhardt[1]

[1] Theodor Kocher Institute, [2] Data Science Lab, and [3] Laboratory for High Energy Physics, University

of Bern, Bern, Switzerland

present address: [+] Department of Pharmacology, University of California, San Diego; [Δ] Department of

Biomedical Engineering, University of Texas at Dallas; [◊] Department of Neuroscience, Karolinska

Institutet, Stockholm, Sweden; [□] Laboratory for Elementary Particle Physics, Department of Physics,

Chiba University, Chiba, Japan

Correspondence:

Dr. Mykhailo Vladymyrov

Theodor Kocher Institute

University of Bern, Bern, Switzerland

mykhailo.vladymyrov@tki.unibe.ch

Authors contribution

M.V. implemented the ^{UFM}Track software framework, conducted automated analysis and

performance evaluation, and contributed to experimental design and imaging.

M.V. and B.E. wrote the manuscript with contributions from S.S., S.A., and L.M.

M.V., with contributions from S.A., L.M., and S.S., conceptualized the T-cell migration analysis

pipeline.

L.M., S.A., and S.S. conducted wet lab experiments and imaging.

S.A., L.M., A.M., and M.V. performed annotation of the datasets for the segmentation model

training.

S.A., L.G., A.P., and M.V. performed manual data analysis.

A.A. and B.E. designed the research together with M.V., acquired funding, and coordinated the

collaboration.

Abstract

The endothelial blood-brain barrier (BBB) strictly controls immune cell trafficking into the central

nervous system (CNS). In neuroinflammatory diseases such as multiple sclerosis, this tight control is,

however, disturbed, leading to immune cell infiltration into the CNS. The development of in vitro

models of the BBB combined with microfluidic devices has advanced our understanding of the

cellular and molecular mechanisms mediating the multi-step T-cell extravasation across the BBB. A

major bottleneck of these in vitro studies is the absence of a robust and automated pipeline suitable

for analyzing and quantifying the sequential interaction steps of different immune cell subsets with

the BBB under physiological flow in vitro.

Here we present the Under-Flow Migration Tracker ([UFM]Track) framework and a pipeline built

based on it to study the entire multi-step extravasation cascade of immune cells across brain

microvascular endothelial cells under physiological flow in vitro. [UFM]Track achieves 90% track

reconstruction efficiency and allows for scaling due to the reduction of the analysis cost and by

eliminating experimenter bias. This allowed for an in-depth analysis of all behavioral regimes

involved in the multi-step immune cell extravasation cascade. The study summarizes how [UFM]Track

can be employed to delineate the interactions of CD4[+] and CD8[+] T cells with the BBB under

physiological flow. We also demonstrate its applicability to the other BBB models, showcasing

broader applicability of the developed framework to a range of immune cell-endothelial monolayer

interaction studies. The [UFM]Track framework along with the generated datasets is publicly available

in the corresponding repositories.

Author summary

Immune cells continuously travel through our body to perform immune surveillance. They travel

within blood vessels at a very high speed and slow down upon reaching their target organ by the

sequential interaction with different adhesion and signaling molecules on the vascular endothelial

cells.

The study of molecular mechanisms mediating this multi-step extravasation of immune cells has

been significantly advanced by in vitro cultures of microvascular endothelial cell monolayers. The

dynamic interaction of the immune cells with endothelial monolayers can be imaged over time in

vitro in microfluidic devices under physiological flow. The 2-dimensional structure of the endothelial

monolayer allows for reliable visualization of the extravasation process required for the study of the

molecular mechanisms involved. The manual analysis of the acquired imaging data is time-

consuming and prone to experimenter error. Analysis automation is, however, hampered by the

similar appearance of the unlabeled immune and endothelial cells and by the flow causing rapid

immune cell displacement.

Here we introduce ^{UFM}Track, the under-flow migration tracker framework allowing for automated

analysis of immune cell interactions with microvascular endothelial cells under flow in vitro. ^{UFM}Track

performs comparably to the manual analysis of an experienced researcher, eliminates

experimenter's bias, and improves the accuracy of the immune cell tracking. Taken together,

^{UFM}Track sets the stage for scalability of in vitro live cell imaging studies of immune cell

extravasation.

Keywords

Automated analysis, blood-brain barrier, cell tracking under flow, deep learning, flow chamber,

leukocyte trafficking, live cell imaging, machine learning, microfluidics, multiple sclerosis, T cell

Introduction

Immune cells continuously travel throughout our body as a means of immune surveillance.

Moving within the bloodstream allows for their fast transport to even distant sites but requires

extravasation once they have reached their target organ. Immune cell extravasation across the

vascular wall is a multi-step process regulated by the sequential interaction of different signaling and

adhesion molecules on the endothelium and the immune cells (1,2). These molecular interactions

mediate distinct sequential steps, namely tethering and rolling to reduce travel speed, shear-

resistant arrest, polarization and crawling of the immune cell on the luminal surface of the

endothelium, and finally, immune cell diapedesis across the endothelial layer (2–4).

The precise molecular mechanisms mediating the multi-step immune cell extravasation in each

organ depend on the immune cell subset but also the specific characteristics of the vascular bed. For

example, in the central nervous system, the endothelial blood-brain barrier (BBB) establishes a tight

barrier that strictly controls the transport of molecules across the BBB, ensuring tissue homeostasis

required for neuronal function (2,5). The BBB similarly controls immune cell trafficking into the CNS.

Thus, accounting for these special barrier properties, unique characteristics of the multi-step T-cell

migration across the BBB have been described. For instance, T cells crawl for very long distances

against the direction of blood flow on the surface of the BBB endothelium in search of permissive

locations for diapedesis (6–8). Research on T-cell interaction with the BBB has already been

successfully translated into therapies in the clinic (9,10).

Exploring the entire multi-step extravasation of immune cells across the BBB has been

significantly advanced by making use of in vitro BBB models maintaining their barrier properties, and

placing them into microfluidic devices (11). Combined with microscopic setups that allow for in vitro

live-cell imaging (Figure 1A-B) of the immune cell interaction with the brain endothelial monolayer

under physiological flow over time, the molecular mechanisms mediating the sequential interaction

of T cells during extravasation across the BBB have been delineated (2). The live cell imaging is

largely performed in the phase-contrast imaging modality (Figure 1C), which does not require

establishing fluorescent labels of the imaged cells and avoids potential photo-toxicity of the

fluorescent imaging modality. These studies have shown that upon their arrest, T cells polarize and

either crawl at speeds between 3 to 10 µm/minute over the brain endothelial monolayer or probe

the endothelial monolayer by remaining rather stationary and sending cellular protrusions into the

endothelial monolayer (Figure 1D) (12,13). Both behaviors can lead to diapedesis of the T cells across

the brain endothelial monolayers. This process lasts at least 3 to 5 minutes, with some immune cells

observed to protrude and retract several times prior to finalizing a prolonged diapedesis process to

the abluminal side of the brain endothelial monolayer (8,14). Finally, T cells that have successfully

migrated across the brain endothelial monolayer usually continue to migrate underneath the

endothelial monolayer (15,16).

The data analysis of these "in vitro flow assays" requires time-consuming offline frame-by-frame

analysis of the imaging data by individual experimenters, in which the dynamic interactions of each

individual immune cell has to be followed over the entire time of the assay manually and assigned to

specific categories. Such analysis is tedious, and accurately assigning the different T-cell behaviors

requires experience. Thus, this manual analysis is prone to inevitable errors and subjective

judgments of the various experiments. Furthermore, the time-consuming manual T-cell tracking

limits the number of events that can be studied and, thus, the statistical power of the analysis.

Automation of the analysis of the recorded multi-step T-cell extravasation across the BBB in the

microfluidic device would thus be highly desirable. It is, however, hampered as these assays are

usually performed with unlabeled cells and imaged by phase contrast (Figure 1C), which poses a

challenge due to the similar grayscales and morphology of the immune cells interacting with the

brain endothelial cells. Further challenges include perfused on top T cells that, in the presence of

shear flow, instantly appear within the field of view (FoV) and are either suddenly displaced over a

certain distance or completely detached and washed away (Figure 1D). Proper analysis of these

events is mandatory for reliable T-cell tracking but also with respect to the study of the overall

avidity of the dynamic T-cell interaction steps with the underlying brain endothelium. Thus, it was

compulsory to establish a tracking solution that accounts for the effect of the flow on the migrating

cells and the distinct migration regimes.

Segmenting of cells imaged in phase contrast migrating on top of cellular monolayer of similar

appearance is a challenging task. Furthermore, the analysis of T cell interactions requires detection

and quantification of the transmigration. While recent advances in computer vision established

several frameworks for detection (17–20) or segmentation (21,22), no tool is readily available to

complete both tasks simultaneously.

There is a large number of powerful cell tracking solutions developed that focus on different

aspects of cell migration, and are embedded in different frameworks or solutions. The general-

purpose Trackpy (23) allows for simple particle tracking and is not suitable to track cells migrating

under flow. It is used for the tracking of cells imaged in phase contrast in the Usiigaci framework

(24). The widely used TrackMate (25) plugin for ImageJ (26) allows for tracking of particles

performing either the Brownian motion or a linear motion, but not the combination of two. Instant

appearance or disappearance as well as the resolving tracks of under-segmented cells cannot be

easily established in this framework. Similarly, CellTraxx (27) performs matching of cells in the

adjacent frames, prohibiting it's use for tracking of under-segmented, intersecting cells or under

flow. The Bayesian Tracker (btrack) framework (28) is using spatial information as well as

appearance information for track linking focusing on the cell divisions required for cell lineage

tracing.

Here we introduce the developed under-flow migration tracker ([UFM]Track) framework that

systematically addresses the hurdles mentioned above, allowing it to perform automated tracking

and analysis of cell-cell interactions imaged in phase contrast. We also show a successful

implementation of [UFM]Track to build an analysis pipeline for T-cell interactions with brain

microvascular endothelial cells in vitro under physiological flow. [UFM]Track reaches 90% T-cell tracking

efficiency, performing comparably to manual analysis while eliminating the experimenter's bias and

improving the accuracy of T-cell tracking. Therefore, it enables significant savings in the labor force

and time for data analysis.

Figure 1. In vitro analysis of the multistep cascade of T-cell migration across the BBB model under physiological flow. A. In vitro under-flow assay setup. T cells were perfused on top of the pMBMEC monolayer, and their migration under flow was observed using phase-contrast imaging modality.

Imaging was performed with a time step of 10 seconds/timeframe. B. In vitro flow assay timeline. During the accumulation phase under flow with the shear stress of 0.1 dynes/cm^2, T cells adhered to the pMBMEC monolayer. After 5 minutes (timeframe 30), the shear stress was increased to 1.5 dynes/cm^2, leading to rapid detachment of not firmly adhering T cells. Analysis of the post-arrest T-cell behavior was thus starting at 5.5 minutes (timeframe 33). tf = timeframe. C. Example of phase-contrast imaging data. Red arrow – crawling T cell; yellow arrow – fully transmigrated T cell; yellow arrow-head – transmigrated part of a partially transmigrated T cell; red V arrowheads – pMBMECs. D. Schematic representation of distinct T-cell behavior regimes detected and analyzed using the developed ^{UFM}Track framework. Crawling cells migrate continuously while probing cells interact with the pMBMECs and move around the interaction point within two cell-size ($20\ \mu m$) as indicated by the arrows. Side and top views are shown.

Results

To design and develop the automated T-cell under-flow migration analysis framework ^{UFM}Track

framework presented here, we made use of in vitro imaging datasets following T-cell migration

across primary mouse brain microvascular endothelial cells (pMBMECs) under physiological flow in

vitro. The framework combines three components: T-cell segmentation and transmigration

detection, T-cell tracking under flow, and analysis of each of the multistep T-cell migration cascade

steps. Segmentation and transmigration detection of the T cells, migrating on the pMBMECs, is

performed with a 2D+T U-Net-like convolutional neural network (21). The T-cell under-flow tracking

algorithm was formulated as a constrained optimization problem.

Next, we describe the methods and algorithms employed to develop ^{UFM}Track. Links to the code

and the datasets used for model training can be found in the Data Availability section.

I. T-cell segmentation

Reliable cell segmentation is crucial for reliable cell tracking. In the phase-contrast imaging

modality, it was impossible to achieve reliable differentiation between T cells and endothelial cells of

the pMBMEC monolayer based on pixel intensity. For detection of T-cell transmigration across

pMBMEC monolayers (diapedesis), sufficiently reliable cell segmentation of the transmigrated T cells

was also required. To achieve this, we have designed 2D and 2D+T U-Net-like (21) fully convolutional

neural network-based models for multitask learning. The models were trained to predict three

maps: cell probability, the probability that the T cell is below the pMBMEC monolayer, and cell

centroids. The models performed predictions based on the grayscale of the respective phase-

contrast images in the case of the 2D model or sequences of 5 timeframes in the case of the 2D+T

model. The models were implemented in TensorFlow (29). The training was performed on a dataset

corresponding to 154 Megapixels of raw image data using the annotation mask for T cells ("T cell

mask") and the mask of the transmigrated part of the T cells ("transmigration mask"), the centroids

map, and the weight map (Figure 2). For validation 37 Megapixels of raw imaging data were used.

The models have approximately 3M and 8M parameters correspondingly, and the model

architectures are summarized in Supplementary Tables 1, 2. Details on models' implementation,

training, and image processing can be found in the "Segmentation models" section of the

Supplementary Material.

Phase contrast	T cell mask	Transm. mask	T cell centroids	Weight map

Figure 2. Example of data used for training of the T-cell segmentation model. From left to right: Phase-contrast microscopy image of T cells migrating on top of the pMBMEC monolayer; annotated T cell mask; annotated mask of transmigrated part of T cells; T cell centroids; w_{cell} weight map. The bottom row shows zoom-in on the highlighted area.

In the T cell mask prediction task, the 2D+T model outperformed the 2D model by a notable 9%

according to the average precision (AP) metric (Table 1). At the same time, for the transmigration

mask, which is much more difficult to infer, the 2D model performance reached only 54%, which is

insufficient for reliable detection of T cells migrating across the pMBMEC monolayer. In this task, the

2D+T model outperformed the 2D model by 32% AP. We observed that the 2D+T model was

sensitive to frame misalignment, leading to false positive detection of transmigrated T cells. Thus, to

generate the preliminary T-cell masks used for frame alignment and histogram normalization, we

employed the 2D model. Afterward, for the T-cell segmentation and transmigration detection, we

employed the 2D+T model, followed by a watershed algorithm based on the inferred T-cell mask and

the cell centroids as seed points. A stitched phase-contrast image sequence can be seen in

Supplementary Video 1 and overlayed with the segmented cells and highlighted transmigration mask

in Supplementary Video 2.

Table 1. Comparison of the performance of the 2D and 2D+T-cell segmentation models.

	Performance, AP	
	2D	2D+T
T Cell mask	86.12%	95.26%
Transmigration mask	50.56%	82.74%

After segmentation, we suppressed the noise by discarding small objects with an area below 30

pixels (the mean area of a T cell is about 400 pixels). For each T cell, we evaluated its coordinates as

geometric mean coordinate, angle of the longer axis, and elongation $\epsilon = \sqrt{\frac{\sum_i (x_i' - x_0')^2}{\sum_i (y_i - y_0')^2}}$,where x_i', y_i'

are projections of the i-th T cell mask pixel coordinates on the orthogonal long and short axes of the

T cell, and x_0', y_0' are the corresponding projections of the geometrical center of the T cell. The

summation is performed over the area covered by the mask of the corresponding T cell. Additionally,

we evaluated the mean T cell probability $\overline{p_{cell}} = \frac{1}{n_{pix}} \sum_i p_{cell,i}$, mean transmigration probability

$\overline{p_{tr}} = \frac{1}{n_{pix}} \sum_i p_{tr,i}$, and cell transmigration coefficient as $t_c = \frac{1}{n_{pix}} \sum_i \frac{\min(p_{tr,i}, \; p_{cell,i})}{p_{cell,i}}$, where n_{pix} is

number if cell pixels and $p_{cell,i}, p_{tr,i}$ are the T cell and transmigration probabilities for i-th pixel in

the predicted T cell mask.

II. T-cell tracking

The presence of flow causes several types of discontinuous events that were managed with a

specialized tracking algorithm. These are: 1) the sudden appearance of T cells in the field of view

(FoV), 2) the sudden detachment of a T cell followed by its disappearance from FoV, and 3) the

displacement over significant distances (up to hundreds of micrometers) of T cells that do not

adhere firmly to the pMBMECs. The primary inspiration for our approach was the Conservation

Tracking algorithm (30). We could consistently reconstruct the T-cell tracks by performing global

optimization constrained to controlled probabilities of T-cell appearance, disappearance, and

displacement due to the flow at every time point. The standard tracking algorithms like tracking by

association is either unable to detect track links with high displacement and thus reconstructs

several disconnected segments of a cell track, or leads to a high number of false track links. Our

procedure favors the reconstruction of long T-cell tracks while at the same time allowing for tracking

of the T cells which detach or are displaced by the flow ("accelerated T-cell movement") over a

longer distance ($> 8\ \mu m$) with speed significantly higher than their crawling speed.

The tracking consists of four main steps: linking, searching for track candidates, global track

consistency resolving, and resolving the track intersections. The inputs for the tracking are the

segmented cells and the cell centroids (Figure 3A). The centroids and the cell proximity are used to

quantify under-segmentation by introducing the node multiplicity (Figure 3B, see section T-cell

tracking of the supplementary information for details). At the linking step, we identified all possible

connections of T cells between the timeframes (Figure 3C).

Next, during the search for track candidates, we aimed to find continuous track segments of T

cells or under-segmented groups of T cells crawling on top of or below the pMBMEC monolayer

without accelerated movement segments on the T cell track characterized by rapid T cell

displacements. The whole dataset of T cells across all time points was represented as a graph. Each

vertex corresponds to a T cell (multiplicity m=1) or a group of potentially under-segmented T cells

(m>1). The vertices are connected according to links obtained at the linking step. Track segments

were found by performing global optimization to find consistent connectivity of vertices across the

time points (Figure 3D) by employing an approach inspired by the conservation tracking algorithm

summarized in Schiegg et al. (30). Optimization was performed using the CP-SAT constrained

optimization procedure using the open source OR-Tools library (31).

Next, resolving global track consistency was performed. In this step, the scope of T cell tracking is

shifted from individual nodes (representing T cells at particular timeframes as well as groups of

under-segmented T cells) to the track segments (unambiguous sequences of nodes) and vertices at

the endpoints of the segments. These are track start and end points, points of merging and

separation of track segments in case of under-segmentation, as well as ambiguous points on a track.

The latter was identified by sudden T cell displacement, a hallmark of detaching and reattaching T

cells and T cells transitioning from properly segmented to under-segmented T cells or vice versa

(Figure 3E). This was followed by the search for potential missing track segments due to accelerated

T cell movement under flow and missing links in the track crossing points. Both are characterized by

significant displacement lengths such that they were not detected during the linking step. Thus we

will refer to both as "jumps" in this section. We have also eliminated short and therefore unreliable

segments, as well as segments whose multiplicity was found to be $M = 0$ (Figure 3F). Afterward, we

separated the segments into two categories, namely segments with multiplicity $M = 1$, i.e., tracks of

isolated T cells, and segments with multiplicity $M > 1$, i.e., tracks of under-segmented groups of T

cells, where tracks of several T cells intersected. Finally, we extended the tracks of isolated T cells

into the intersection if the branching vertex had a multiplicity of n and was splitting in exactly n

segments with multiplicity $M = 1$ (Figure 3G).

Lastly, we resolved the track intersections to obtain reliable T cell tracks, i.e., identifying track

segments corresponding to the same T cell before and after the under-segmented track region

(Figure 3H).

Detailed information on the developed under-flow T cell tracking algorithm is given in the "T-cell

tracking" section of the Supplementary Material.

III. T-cell migration analysis

Next, we performed the T-cell migration analysis based on the reconstructed T cell tracks. We

selected tracks inside the fiducial area of the FoV, namely coordinates of the T cell at all timepoints

along the track were located at least 25 μm away from the bounding box enclosing all segmented T

cells. Next, tracks of T cells touching another T cell at the end of the assay acquisition were excluded

since T cells directly adjacent to each other can hide the start of T cell transmigration across the

pMBMEC monolayer and thus compromise correct detection and quantification of this step.

Additionally, only tracks with T cells were assigned in at least 6 timeframes during the physiological

flow phase. We also require T cells to be assigned for at least 75% of timeframes along the track.

Under-segmented parts of T cell tracks were not considered. Examples of selected tracks can be

seen in Supplementary Video 3.

Figure 3. The T-cell tracking pipeline. A. Segmentation and centroids of two migrating T cells under flow

over time. B. Nodes corresponding to segmented T cells. Node sizes and colors denote the multiplicity

estimate of a node. C. Nodes, together with links between adjacent in space and time nodes, form a graph. D. Selected links are obtained with global optimization on the graph. E. Graph of segments and vertices obtained according to node multiplicity. F. Resolving global segment multiplicity consistency with global optimization on the graph. Additionally, a search for rapid displacement of segmented T cells due to under-segmentation or flow is performed. G. Extension of tracks of individual T cells into the track intersection. H. Resolving track intersections.

After exclusion of the tracks of the cellular debris, which were misclassified as T cells during the

segmentation step, using a dedicated "not-a-T-cell" classifier, we performed identification of

transmigration (Supplementary Figure 5), probing and crawling migration regimes, as well as

accelerated movement along the T cell track. Detailed information on the analysis of the T cell tracks

is given in the "T-cell migration analysis" section of the Supplementary Material.

Finally, for each track, we evaluated motility parameters for each of the following migration

regimes: probing before the transmigration, crawling before the transmigration, all crawling above

pMBMECs monolayer including T cell crawling segments after the first transmigration attempts, all

crawling below pMBMECs monolayer, whole T-cell track excluding accelerated movement and

tracking inefficiency regions, as well as whole T-cell track. Specifically, we evaluated the following T-

cell motility parameters: duration of each migration regime, the total vector and absolute

displacements, the migration path length, the average migration speed (displacement over time),

average crawling speed (path length over time), and finally the mean and standard deviation of the

instantaneous speed. We evaluated migration time, displacement, and average speed for the

accelerated movement regime.

IV. Analysis of trafficking datasets

Having developed a full pipeline based on the [UFM]Track framework for automated analysis of the

multi-step T-cell migration cascade across the BBB model, we next aimed to compare the results of

the automated analysis with previous studies, assess the capacity of the framework to gain novel

insight into T-cell migration under flow, and evaluate its performance as compared to manual

analysis.

1. CD4[+] T cell analysis

To this end, we first analyzed a total of 18 imaging datasets each corresponding to 32 minutes

recording dedicated to understanding the multi-step migration of CD4[+] T cells across non-stimulated

(n= 6), TNF stimulated (n=5), and IL1-β-stimulated (n=7) pMBMEC monolayers under physiological

flow with the developed pipeline.

In these in vitro live cell imaging datasets, phase-contrast imaging was performed as described in

the "Data acquisition" subsection in the Materials and Methods section. The T-cell accumulation

phase corresponding to the shear stress of 0.1 dynes/cm^2 lasted for 32 timeframes (5 minutes),

followed by conditions of physiological flow (shear stress 1.5 dynes/cm^2) for the subsequent 160

timeframes (27 minutes). The shear stress is controlled by an automated syringe pump. After

increasing to physiological flow rates (12), a significant number of T cells detached from the

pMBMEC monolayers, with higher numbers of T cells detaching from non-stimulated pMBMEC

monolayers when compared to those stimulated with pro-inflammatory cytokines as detected by

the distribution of the track ending times between 5 and 30 minutes. (Figure 4A).

Analyzing these datasets with the established [UFM]Track framework, the T-cell behavior type for

each detected T-cell track and the aggregated T-cell behavior statistic were obtained for the

stimulated and non-stimulated pMBMEC monolayers (Figure 4B). The total number of analyzed T cell

tracks is n=971, n=780, and n=1218 for the non-stimulated, TNF stimulated, and IL1-β stimulated

pMBMEC monolayers correspondingly. The data obtained with [UFM]Track were in accordance with our

previous observations obtained by manual frame-by-frame analysis (12,14,32). We further obtained

the data for T-cell migration speed, T-cell displacement, and T-cell path lengths for the crawling T

cells, as these are the primary cell motility parameters. As shown in Figure 4C-H, we observed

statistically significant differences in the mean T-cell motility parameters depending on the

pMBMECs stimulation condition. Furthermore, we observed differences in the shape of the

distribution of T-cell crawling speeds, consistent with our previous reports of underlying differences

in the mechanisms mediating T-cell crawling on non-stimulated versus cytokine-stimulated

pMBMECs (32). While not statistically significant for the IL1-b and with a low significance for the TNF

stimulations, an increased variance of the T cell instantaneous speed on stimulated compared to

non-stimulated pMBMECs is observed(Figure 4G). This suggests that T-cell crawling is often

interrupted by T-cell recognizing specific cues on the stimulated endothelium. The T cell meandering

index (MI) distribution was high on non-stimulated and lower on stimulated pMBMECs (Figure 4H),

underscoring that cytokine stimulation enhances directed T-cell movement on the pMBMEC

monolayer.

Since the analysis was performed automatically with [UFM]Track, it allowed for deeper insights into

the T-cell migration behavior on the pMBMEC monolayers than obtained by manual frame-by-frame

analysis. Specifically, the detection of accelerated movement by [UFM]Track enables the researcher to

quantify the kinetics of individual T-cell detachment from the endothelium rather than simply

quantifying a bulk T-cell detachment rate. This detachment kinetics is reflected in the average speed

experienced during accelerated movement, which is significantly lower for the IL1-b condition

(Figure 4I), suggesting the need to break more bonds with the endothelium when pro-inflammatory

cytokines are present.

As our [UFM]Track workflow achieves sufficient segmentation efficiency to detect T cells below the

pMBMEC monolayer and T-cell transmigration across the pMBMEC monolayer, it also enables

investigation of T-cell movement after the transmigration step. In Figure 4J, we show the distribution

of the migration speed for the transmigrated CD4$^+$ T cells. Interestingly, cytokine stimulation of

pMBMECs, although applied from the luminal side, also affected T-cell movement at the abluminal

side of the pMBMEC monolayer. While in the microfluidic device used in the present study, the

migration of T cells below the pMBMEC monolayer may not be biologically relevant, this analysis

option will be valuable for future studies involving multilayer in vitro BBB models that include the

vascular basement membrane in addition to pericytes and astrocytes to mimic the entire

neurovascular unit (8,33).

Importantly, our ^{UFM}Track automatically detects and characterizes unusual events, such as reverse

T-cell transmigration, that are easily overlooked with manual counting. We do not present the

statistics here, as only a few such events were observed. However, when applied to the multilayer

BBBs in forthcoming in vitro models, systematic detection and analysis of these rare events will be

important to understand T-cell migration in immune surveillance.

Figure 4. Analysis of CD4+ T-cell tracks. A. Start and end time distribution of CD4+ T-cell tracks on

non-stimulated or TNF or IL-1β stimulated pMBMECs. UFMTrack correctly captures increased T-cell

detachment from the non-stimulated pMBMECs after 5 minutes when the flow is increased to

physiological levels. B. Quantification of CD4+ T-cell behavior in the respective categories obtained

on non-stimulated and TNF and IL-1β stimulated pMBMECs. Error bars show the statistical error

of the mean. (see text for details). C-H. Motility parameters of the crawling CD4+ T cells were

obtained for the three endothelial stimulatory conditions. Distributions of T-cell path length (C),

displacement (D), crawling speed (path/time) (E), migration speed (displacement/time) (F),

variability of instantaneous T-cell crawling speed along the track (standard deviation, G), and

meandering index (H). I. Distribution of CD4+ T-cell accelerated movement (AM) speed is a proxy

metric for the T-cell adhesion to the healthy or inflamed endothelium. J. Migration speed distribution of the transmigrated CD4[+] T cells. Stimulation applied to the luminal side of pMBMECs affects T-cell migration at the abluminal side of pMBMECs after their transmigration.

FT - T cells performed full transmigration; UT – T cells performed uncompleted transmigration.

2.   CD8[+] T cells analysis

As we have previously shown that the multi-step T-cell extravasation across pMBMEC

monolayers differs between CD4[+] and CD8[+] T cells (13), we next analyzed two datasets studying the

multi-step migration of CD8[+] T cells across non-stimulated (NS) and TNF/ interferon-gamma

(TNF/IFN-γ) stimulated pMBMEC monolayers under physiological flow over 161 timeframes

(27 minutes). The CD8[+] T cells were slightly different in size and appearance when compared to the

CD4[+] T cells used for training the segmentation model. To benchmark our established [UFM]Track

pipeline in this more difficult configuration, the datasets were first manually analyzed by four

experimenters: one advanced experimenter with four years of experience (AdEx) and three

inexperienced experimenters who received comparable 2-hour introduction and training (Ex1-Ex3).

The analyses were performed on the subset of the acquisition, 161 timeframes long starting from

timeframe 31.

Manual cell analysis and tracking were performed as described in the "Materials and Methods"

section separately for each of the 8 tiles of the tiled acquisition. Next, all crawling CD8[+] T cells that

did not perform transmigration were manually tracked for the timespan after the flow was increased

to the physiological level using the manual tracking in ImageJ.

Figure 5. Comparison of automated analysis with ^{UFM}Track and manual analysis of the CD8$^+$ T-cell tracks. A-C: CD8$^+$ T-cell behavior statistic obtained for non-stimulated (NS) and cytokine-stimulated pMBMECs as obtained manually by four experimenters, as well as automatically with ^{UFM}Track. A. Quantification of CD8$^+$ T-cell behavior in the respective categories obtained on non-stimulated and TNF/IFN-γ stimulated pMBMECs is consistent with results obtained by manual frame-by-frame analysis. Cytokine stimulation of pMBMECs increases T-cell probing behavior (B) and T-cell transmigration rate (C). Error bars show the standard deviation of the manual analysis and the statistical error of the mean for automated analysis. Points correspond to individual experimenters.

D. Counts of CD8[+] T cells were obtained manually by four experimenters and automatically by [UFM]Track. The T-cell detection efficiency is above 90%. Error bars show the standard deviation of the manual analysis and the statistical error of the mean for the automated analysis. Points correspond to individual experimenters. E. Detection efficiency of crawling CD8[+] T cell tracks between the manual and automated analysis. F-I. Comparison of CD8[+] T-cell crawling speed (path/time) (F, G) and migration speed (displacement/time) (H, I) on non-stimulated (F, H) and cytokine-stimulated (G, I) pMBMECs. The T-cell position assignment error in manual tracking leads to biased crawling speed estimation. J. Comparison of the analysis time (per cell) required for behavior analysis and tracking of CD8[+] T cells. K. Total analysis time (per dataset) for behavior analysis and tracking of CD8[+] T cells. Comparison is shown for manual analysis with tracking of crawling cells only (Crawling manual), the time estimate for manual analysis with tracking of all cells (All manual), and the in-depth automated analysis of all cell tracks with [UFM]Track (All auto).

FT - T cells performed full transmigration; UT – T cells performed uncompleted transmigration; NT - T cells did not perform transmigration.

First, we compared the CD8[+] T cell behavior statistics obtained manually and by our automated

analysis pipeline (Figure 5A-C). The results obtained by the automated analysis were in full

agreement with the manual analysis performed by the experienced experimenter. At the same time,

the data highlight the variability of the results from the inexperienced experimenters, confirming the

potential for significant experimenter bias.

We also show the detection efficiency of T cells by the [UFM]Track. To achieve this goal, we

combined the numbers of CD8[+] T cells detected in each tile by manual analysis. Due to the overlap

between the individual tiles, some CD8[+] T cells are seen more than once. To obtain an estimate of

the total CD8[+] T cell count detected manually, we scaled the number of CD8[+] T cells accordingly to

the number of detected unique T-cell tracks (see next paragraph). By this approach, we found T-cell

detection efficiency to be above 90% (Figure 5D). The increased T-cell density can explain lower

efficiency in the NS condition. These findings set the appropriate T-cell density for automated

analysis of their migration behavior on pMBMEC monolayers to 200-300 cells per FoV of the size of

3.8 mm^2 or 50-80 cells/mm^2.

Next, we compared the T-cells tracks as obtained manually by three experimenters as well as

automatically by our ^{UFM}Track. To this end, we first matched the tracks in adjacent tiles to obtain the

tracks on the whole imaging area, avoiding multiple counts of the same track. This was achieved by

pattern matching with initial offsets between tiles obtained by an automatic frame alignment

procedure performed as part of the automatic analysis. We considered T-cell tracks observed in

different tiles to be tracks of the same T cell if $f_{in} > \frac{1}{2} f_{out} - 0.8$, where f_{in} is the fraction of

timepoints along a track at which the distance between the T cells was below 17 μm and f_{out} was

the fraction of timepoints along a track at which distance between cells was above 25 μm. We used

the same approach to match the T-cell tracks obtained by each experimenter and the automated

^{UFM}Track. To compare performance in an objective manner, we excluded manually obtained tracks

that lay outside of the fiducial volume of the automated analysis, as well as T cells that were

touching another T cell at the end of the acquisition, as those were also excluded from the

automated analysis. We then took as 100% the sum of all T-cell tracks detected manually. We

evaluated the fraction of all T-cell tracks detected by each experimenter as well as automatically by

our ^{UFM}Track.

In Figure 5E, we show that our pipeline achieves a comparable T-cell tracking efficiency when

compared to the manual analysis. While its performance was not superior to that of the

experimenter with four years of expertise in the analysis of the under-flow datasets, it does perform

better than less experienced experimenters.

We also compared the T-cell motility parameters obtained manually and automatically for the

non-stimulated and stimulated pMBMECs (Figure 5F-I). Data obtained for the CD8$^+$ T-cell migration

speed on pMBMECs was comparable between all experimenters and the automated analysis. In

contrast, results obtained for the T-cell crawling speed were significantly different between the

automated approach and manual analysis. Taking a closer look at the T-cell tracks (Supplementary

Figure 6), we readily observed that the manual analysis contains significant errors in the assignment

of the T-cell position. This leads to a jittery pattern in the T-cell migration tracks and overestimating

the T-cell path and, thus, T-cell crawling speed. Employing our novel automated analysis pipeline

eliminated this systematic error in the measurements.

To finally investigate the potential benefit in the time required for automated versus manual data

analysis, we examined the time required for a given experimenter to analyze such datasets manually

using the Clockify time tracker (34). In Figure 5J, we show the average time investment needed to

perform a complete T-cell behavior analysis and T-cell tracking in each imaging tile, as well as on

average. Clearly, the experienced experimenter outperforms the inexperienced experimenters by a

factor of 3. On average, a researcher would thus spend 8.1 hours analyzing a dataset with 300 T cells

when only the tracks of crawling T cells (50%, i.e., 150 cells) are analyzed. If all T-cell tracks were to

be analyzed, this time would further increase to 12.9 hours. Given that on average 10 datasets can

be produced per day, manual analysis becomes a bottleneck, leading to delays in exhaustive data

analysis and thus ultimately in research progress. The [UFM]Track framework reduces analysis time by a

factor of 3 when analyzing only crawling cells and 5 if all cell tracks are to be analyzed (Figure 5K).

With an average analysis time of 2.3 hours, the 10 experiments carried out in one day can be

thoroughly analyzed within one day of machine time. This enables the scalability of flow-based

immune cell migration experiments while simultaneously lifting the burden of tedious and time-

consuming manual analysis from the researchers.

V. Applicability of [UFM]Track to different models of immune cell - endothelial cell interactions

To assess the broader applicability of the [UFM]Track framework to different cell types and BBB

models employed for the *in vitro* under-flow cell trafficking assays. To this end, we have performed

the automated analysis of the multi-step T-cell migration cascade in a set of datasets (one to eight)

across different cell types and assays. Specifically, we attempted to verify the feasibility of analysis of

datasets where the cells forming the monolayer or the migrating cells have noticeably different

appearances, cells from other species are used, specifically human-derived cells, and datasets with a

different imaging modality. These include human peripheral blood mononuclear cells (hPBMCs),

human T cells, bone marrow-derived macrophages (BMDM) interacting with a monolayer of mouse-

and human-derived brain microvascular endothelial cells as well as with immobilized recombinant

BBB adhesion molecules (See "Material and Methods" section for details). To assess the

performance of the model in each condition, we have performed a visual inspection of the cell

masks, transmigration masks, and cell tracks for few datasets in each assay type. We then analyzed

cell behavior distribution or motility parameters relevant in each case.

1.   hPBMC on HBMEC monolayer

The interaction of hPBMCs with the BBB is altered under different disease conditions. We have

therefore analyzed a dataset of human PBMCs on human brain microvascular endothelial cells

(HBMEC) to quantify the cell behavior statistic. While the endothelial cells look significantly different

than the pMBMECs, the segmentation and transmigration detection quality was sufficient to

perform the cell migration analysis. This was also confirmed by the visual inspection of the cell and

transmigration masks and cell tracking results (Figure 6 A). We then quantified the PBMC behavior

distribution and the crawling and migration speed of the crawling PBMC (Figure 6 B, C).

2.   BMDM on the pMBMEC monolayer

Invasion of macrophages into the CNS contributes to many neurological disorders. Therefore, we

have explored if [UFM]Track also allows us to analyze the interaction of BMDMs with the BBB under

physiological flow in vitro. To this end, we have analyzed two datasets of mouse BMDMs interacting

with pMBMEC monolayers (Figure 6 D). Despite the different behavior and appearance of the

macrophages as compared to the T cells, the segmentation quality was sufficient to perform the cell

tracking and analyze cell motility parameters (Figure 6 E, F), as confirmed by visual inspection of the

macrophages masks and cell tracking results (Figure 6 D). Nonetheless, due to the different

appearance of the flattening crawling cells and the transmigrated cells, the transmigration detection

trained on the mouse T cells does not perform sufficiently well for the robust detection of

transmigrating macrophages.

3.   Human T cells on EECM-BMEC monolayer

Next, we analyzed eight datasets of human T cells interacting with the human induced pluripotent

stem cell (hiPSC) derived extended endothelial cell culture method brain microvascular endothelial

cell (EECM-BMEC)-like cell monolayers (Figure 6 G). EECM-BMECs establish barrier properties

comparable to primary human BMECs and display a mature immune phenotype allowing to perform

studies for the first time in a fully autologous manner of immune cell interactions with patient-

derived in vitro models of the BBB (35). Despite the cells being of different species, the T cells were

successfully segmented, and their transmigration was detected, as confirmed by visual inspection of

the T-cell and transmigration masks and cell tracking results (Figure 6 G).

We have confirmed the differences in migration statistics (Figure 6 J, K), migration and crawling

speed of the crawling T cells (Figure 6 H), as well as the previously reported difference in relative

(normalized to track pathlength) displacement (Figure 6 I) of the T cells treated with anti-α4-integrin

+ anti-β2-integrin blocking antibodies (bab) and control (ctrl) condition (36).

4.   Human T cells on immobilized recombinant BBB adhesion molecules

Finally, we have analyzed datasets acquired in a different imaging modality. Seven datasets of

human T cells interacting with immobilized recombinant BBB adhesion molecules, namely ICAM-1,

VCAM-1, and their combination (Figure 6 l). In this case, the fluorescently labeled cells were imaged

with epi-fluorescent microscopy. In the absence of a monolayer, transmigration detection is not

necessary. Thus we performed cell segmentation directly on the fluorescent signal (See Materials

and Methods for details) using the same segmentation strategy as for the T-cell masks. The tracking

was then performed the same way as for other experiments. The T cells were successfully

segmented and tracked, as confirmed by visual inspection. We compared T cell behavior and motility

parameters for three conditions of the recombinant adhesion molecules: ICAM-1, VCAM-1, and

ICAM1 + VCAM1 (Figure 6 M, O). Consistent with previous observations (12,36) where the analysis

was performed using the ImageJ (26) either manually or with the TrackMate (25) plugin, we

observed that in the absence of ICAM-1, T cells were moving in the direction of flow (Figure 6 N). At

the same time, ^{UFM}Track allows for capturing the rapid accelerated motions of the cells and thus

reconstructing longer continuous T cell tracks. This enables capturing the migrations properties of

the consecutive steps of the T cell migration cascade on the call-by-cell basis.

Figure 6. Analysis of different BBB models with ᵁᶠᴹTrack. A-C. Human PBMCs interacting with HBMEC. A. Raw image frame, segmentation, and tracking results. B. PBMCs behavior distribution. C. Crawling and migration speed distribution. D-F. BMDM interacting with the pMBMEC monolayer. D. Raw image frame, segmentation, and tracking results. E. BMDM crawling vs probing behavior distribution. F. Crawling and migration speed distribution. G-K. Human T cells interacting with the EECM BMEC monolayer. G. Raw image frame, segmentation, and tracking results. H. Crawling and migration speed distributions for the control condition (ctrl) and with the blocking anti-bodies (bab). I. X and Y relative cell displacement. J. Crawling vs probing behavior comparison. K. Transmigrations behavior comparison. L-O. Human T cells interacting with immobilized recombinant BBB adhesion molecules. L. Raw fluorescent image frame, thresholding-

based segmentation, and tracking results. M. Crawling vs probing behavior distribution. N. X and Y

relative displacement O. Crawling vs probing speed comparison.

Scale bar 50 µm. Statistical tests are performed with the Mann-Whitney U test.

Discussion

In this study, we have developed the under-flow migration tracker ([UFM]Track) framework. It

consists of independent modules and allows for segmentation (Figure 2), tracking (Figure 3), and

motility analysis of T cells, migrating on, across, and below (Figure 1D) the monolayer of brain

microvascular endothelial cells under physiological flow in vitro. The developed method relies

exclusively on phase-contrast imaging data (Figure 1B). Therefore, it does not require establishing

fluorescent labels of the migrating immune cell population to be studied, avoids potential photo-

toxicity to be considered for fluorescent imaging modalities (37–39), and is thus the preferred choice

for analyzing the trafficking of sensitive cell types. T-cell segmentation and prediction of the

transmigrated T-cell areas are performed using a custom 2D+T U-Net-like convolutional neural

network. It enables reliable segmentation of T cells both above and below the pMBMEC monolayer.

The existing particle and cell tracking toolkits consider migration of one cell type and thus are not

suitable for the detection of distinct migration regimes and cell interactions (23–25,28).

Furthermore, to our knowledge, none of the existing algorithms consider migration under the flow

causing rapid cell displacement. The tracking of T-cell interactions with the pMBMECs during all

migration regimes under physiological flow required designing a new tracking algorithm considering

rapid T-cell appearance, disappearance, and displacement in the field of view caused by the flow.

We have also developed approaches that resolve track intersections, i.e., identifying track segments

corresponding to the same T cell before and after under-segmented track regions, which are

inevitable during T-cell migration. By establishing the detection of T-cell crawling, probing,

transmigration, and accelerated movement combined with reliable T-cell tracking, we have enabled

the in-depth analysis of distinct migration regimes on a cell-by-cell basis. By reducing the dataset

analysis time by a factor of 5, ^{UFM}Track allows for performing a thorough analysis of 10 experiments

that a researcher can carry out in one day within one day of machine-time. We have demonstrated

that the automated analysis performs on par with manual analysis while improving accuracy and

eliminating experimenter bias, and enables scalability of flow-based immune cell migration

experiments by reducing the analysis cost and lifting the burden of time-consuming manual analysis.

In this work, we have demonstrated the applicability of the developed framework to the analysis

of the multi-step extravasation of CD4$^+$ and CD8$^+$ T cells across non-stimulated or cytokine-

stimulated pMBMEC monolayers under physiological flow. We have also demonstrated that the

developed framework allows for automated analysis of T-cell behavior statistics and motility

parameters of distinct T-cell migration regimes. Results of the automated analysis of CD4$^+$ T-cell

behavioral statistics performed with ^{UFM}Track are in agreement with previous studies using manual

data analysis (Figure 4B) (14). The automated analysis of datasets of CD8$^+$ T-cell migration has shown

comparable performance with manual analysis performed by one experimenter with four years of

analysis experience and three less-experienced experimenters. At the same time, the variance of the

results obtained manually showed significant experimenter bias that the automated analysis

eliminated (Figure 5F). Additionally, the automated analysis allowed for in-depth analysis of all T-cell

migration categories and precise evaluation of T-cell motility parameters, which was not achieved by

manual T-cell tracking, even by the most experienced user.

We have further demonstrated the applicability of the ^{UFM}Track framework to the analysis of

different cell types (BMDM, PBMC), species (human-derived T cells, HBMEC, and EECM BMEC) as

well as tracking and analysis of cell migration imaged using the fluorescent imaging modality. In all

cases, the cells could be successfully segmented and tracked, and the cell migration analysis was

performed on a cell-by-cell basis. The main challenge for the system is reliable segmentation of the T

cells and the transmigrated T cells. In the case of HBMEC the monolayer has a dotted pattern unlike

the rather smooth pMBMEC. This leads to more false positives in the T-cell segmentation at the level

of individual frames, which nonetheless could be efficiently filtered at the level of cell tracking. In the

case of the BMDMs, since they have significantly different appearances during transmigration from

the T cells, reliable transmigration detection was not obtained. This prevents the system from

quantifying the transmigration statistics of the BMDMs, yet the cell detection and thus motility

parameter evaluation of BMDMs was successful (Figure 6 D-F). For all the other analyzed cell types,

both the cell segmentation and transmigration detection model trained on the annotated CD4[+] T

cells also produced transmigration maps of sufficient quality for reliable transmigration detection.

The framework was able to successfully detect the cells of sizes that differ by almost a factor of two

(Supplementary Figure 7). This demonstrates, that simply by discriminating cells by size, it is possible

to extend [UFM]Track to study interaction of several types of immune cells migrating on top of a cellular

monolayer under flow and imaged by phase contrast.

Quantifying the fraction of transmigrated T cells is crucial for studying the molecular mechanisms

governing the infiltration of autoaggressive T cells across the BBB into the CNS parenchyma and the

immune surveillance. The role of specific endothelial or T-cell adhesion molecules in influencing the

dynamic interactions of T cells with pMBMECs can, e.g., be probed by quantifying the ratio of T-cell

crawling behavior to T-cell probing (Figure 6M) (14,40,41). The analysis procedure established in the

[UFM]Track framework is flexible and can be easily extended to evaluate further properties of the

migrating cells. For example, the number of times a T cell interrupts its crawling regime by switching

to short probing behavior on the endothelium can be evaluated, thus providing information on the

distribution of "hot spots" on the endothelium. Additionally, experiment scalability and the analysis

on a cell-by-cell basis enable the search for distinct populations of T-cells, e.g., according to the

distribution of probing to crawling behavioral ratios. The strength of T-cell adhesion to the

endothelium can be probed with the developed framework using the measure of T-cell detachment

rate and the distribution of the previously overlooked T-cell accelerated movement occurrences and

accelerated movement speed. Finally, the possibility of quantifying the motility parameters of the

transmigrated T cells enables future studies involving multilayer in vitro BBB models that include the

vascular basement membrane in addition to mural cells such as pericytes and astrocytes mimicking

the entire neurovascular unit.

While in this work we have focused primarily on the analysis of mouse T cells interacting with

pMBMEC under flow or other immune cells interacting with the endothelial monolayer, the methods

we present establish a foundation for a broad range of studies involving in vitro under-flow studies

of immune cell trafficking. The tracking algorithm and motility analysis developed in the ^{UFM}Track

framework are also directly applicable to the analysis of immune cell interactions with recombinant

BBB adhesion molecules under flow. In this case, either phase-contrast or epi-fluorescent imaging

can be employed when studying fluorescently labeled immune cells, as demonstrated. The T-cell

segmentation based on deep neural networks can be applied to studies of the trafficking of other

immune cell subsets on and across the pMBMEC monolayer. We demonstrated the successful

application of the ^{UFM}Track to cells with significantly different appearances, such as the HBMEC

cellular monolayers. The application to immune cell trafficking across other endothelial monolayers,

including those from different species or vascular beds as well as lymphatic endothelial cells where

the cell appearance is drastically different from the ones demonstrated here, is possible but requires

fine-tuning of the trained segmentation model. While training the models presented here required a

large dataset of annotated T-cell masks, and transmigration masks, this can be largely avoided for

future development. Future models can be developed more efficiently by leveraging our existing

annotated dataset of T-cell migration on the pMBMECs while employing transfer learning

approaches in a multitask framework with weak supervision and fluorescent labels as auxiliary

learning targets to adapt the model for the segmentation of cells with different appearances. This

can be further improved by adopting self-supervised contrastive or masked learning methods, which

have demonstrated significant advancements in model pretraining in recent years (42–45). This is

the subject of our future studies. The framework could be employed in combination of other

methods, e.g. by incorporating fluorescently labeled endothelial adherens junctions one could

differentiate the trans- from the paracellular transmigration in the migrating cells. In this case the

transmigration locations detected by the [UFM]Track framework could be automatically extracted for

analysis. One could employ the tracking under flow implemented in the [UFM]Track framework for

tracking the immune cells migrating on the vascular wall *in vivo* to capture the accelerated

movement of cells. By sharing the data and open-source code of the [UFM]Track framework, i.e., the

training data used for the segmentation model training, trained models, as well as the model

architecture, and full under-flow T-cell tracking and migration analysis pipeline, we hope to

encourage the community to pursue these developments to advance the field.

One current limitation of the method is the performance reduction of the T-cell tracking when

the density of migrating T cells significantly increases (>250 cells/dataset). Thus, the recommended T

cell concentration is about 100-150k/ml. However, to avoid non-physiological interactions between

migrating T cells and T cell clumping, the T-cell density should be kept at moderate levels anyway.

Additionally, analyzing data from 8 stitched FoVs allows for obtaining comparable statistics. The

current analysis of the behavioral statistics quantifies the fraction of different T-cell migration

regimes with respect to the number of adhered cells instead of the total T-cell count passing through

the microfluidic device. While the number of fast-moving T cells during the accumulation phase

cannot be directly counted using the imaging modality employed here, it can be estimated indirectly

from the imaging data with additional calibration experiments and a dedicated machine learning

model. Another limitation is that currently, we do not consider dividing immune cells. While cell

division events happen rarely (<1% of T cells) and are not primary events for the study of immune

cell interaction with the BBB model, the modular architecture of our framework will facilitate future

extension to detect cell division.

By enabling experiment scalability, unbiased analysis with advanced accuracy, and an in-depth

analysis of large datasets of T-cell dynamics under flow, the computational and analytical framework

presented here contributes to the 3R principle when studying the interaction of cells derived from

animal models by reducing the number of animals to be sacrificed. [UFM]Track can be employed for

fundamental research of the molecular mechanisms governing immune cell trafficking across a range

of vascular beds, screening of pharmaceutical treatments, as well as for personalized medicine

based on the evaluation of treatment efficacy on patient-derived T-cell migration behavior on

patient-derived endothelial monolayers. One can furthermore envisage that [UFM]Track can be suitable

to study cancer cell metastasis across different endothelial cell monolayers. Eventually, the

developed framework can be extended to real-time operation during image acquisition. Combined

with transgenic photo-convertible immune cells allowing for the photoconversion of immune cells

according to their behavior will allow for subsequent fluorescent cell sorting and scRNA-Seq analysis.

Such advanced studies are needed to reveal the role of genetic differences governing T-cell

migration regimes.

Materials and Methods

1. pMBMEC cell culture

Primary mouse brain microvascular endothelial cells (pMBMECs) were isolated from 8-12 weeks old C57BL/6J WT mice and cultured exactly as described before (12,46). Intact monolayers were stimulated or not with 10 ng/mL of recombinant mouse TNF, 20 ng/mL of recombinant mouse IL1-β for, or 5ng/mL recombinant mouse TNF + 100 U/mL recombinant mouse IFN-γ 16-24 hours prior to the assays as previously described (32).

2. Human brain microvascular endothelial cells

Human brain microvascular endothelial cells (HBMEC) were kindly provided by Prof. Nicholas Schwab (University of Münster, Germany). They were cultured to confluency on speed coating solution coated µ-Dishes (35 mm, low, iBidi) for 2 days as previously described (32).

3. Extended endothelial culture method—brain microvascular endothelial cell-like cells

The Extended Endothelial Cell Culture Method (EECM) was used to differentiate human induced pluripotent stem cells (hiPSCs) to brain microvascular endothelial cell (BMEC)-like cells were employed as a human in vitro model of the BBB. In brief, hiPSCs from one healthy control (HC) (cell line ID: LNISi002-B) were established from erythroblasts in the laboratory of Renaud DuPasquier (University of Lausanne, Switzerland). EECM-BMEC-like cells were cultured to confluency on collagen IV and fibronectin-coated µ-Dishes (35 mm, low, iBidi) for 2 days (36).

4. T cell and hPBMC preparation

Naïve CD4$^+$ and CD8$^+$ T-cell isolation: Peripheral lymph nodes and spleens from 2D2 and OT-I

C57BL/6J mice were harvested, and single-cell suspensions were obtained by homogenization and

filtration through a sterile 100 μm nylon mesh. A second filtration was applied after erythrocyte lysis

(0.83% NH4Cl, Tris-HCl). 2D2 and OT-I cells were isolated respectively with magnetic CD4$^+$ and CD8$^+$ T

cell selection beads (EasySep, STEMCELL Technologies).

In vitro activation of naïve CD8$^+$ T cells: OT-I CD8$^+$ T cells were activated as described before

(13,41). Activated CD8$^+$ T cells were cultured in IL-2-containing media for 3 days post-activation.

In vitro activation of naïve CD4$^+$ T cells: 2D2 CD4$^+$ T cells were activated as described before (32).

Activated CD4$^+$ T cells were cultured in IL-2-containing media for 24 additional hours.

hPBMC were isolated from buffy coats of healthy donors by Ficoll-Paque™ Plus (Cytiva) density

gradient and were frozen and stored in a liquid nitrogen tank until use as previously described in

(36).

Human CD4$^+$ T cells were isolated by employing a CD4$^+$ T cells isolation kit (Miltenyi Biotec kit)

following the provider's instructions. Subsequently, effector/memory CD4$^+$ T cells were sorted by

Fluorescence-Activated Cell Sorting (FACS) into different Th subsets according to their specific

surface expression of chemokine receptors (CCR6$^-$CXCR3$^+$CCR4$^-$ for Th1; CCR6$^+$CXCR3$^+$CCR4$^-$ for

Th1*; CCR6$^-$CXCR3$^-$CCR4$^+$ for Th2; CCR6$^+$CXCR3$^-$CCR4$^+$ for Th17) and expanded as previously

described (1,33,40,47–49).

5. BMDM preparation

BMDMs were prepared as described in (Berve et al., manuscript in preparation). In brief, the

mouse bone marrow was harvested and washed through a 100 μm nylon mesh. After 7 days of cell

culture were plated in a density of 17-20 × 10^6 cells/mL in BMDM medium supplemented with

5ng/mL of recombinant mouse macrophage colony stimulating factor (mCSF, R&D Biosystems,

Minneapolis, USA, 416-ML-500) onto non-treated 100mm tissue culture Petri dishes (Greiner Bio-

One, St. Gallen, Switzerland) for 7 days at 37°C and 5% CO_2 the cells were harvested by incubation

with 0.05% Trypsin.

6.   T cell binding assays under physiological flow conditions

For live-cell imaging of T-cell interaction with recombinant BBB cell adhesion molecules under the

physiological flow condition, μ-Dishes (35 mm, low, iBidi) were coated with 1.54 μg/mL human

recombinant VCAM-1 (Biolegend) and/or 1.14 μg/mL human recombinant ICAM-1 (R&D system) in

DPBS for 1 h at 37°C and blocked with 1.5% (v/v) bovine serum albumin (Sigma-Aldrich) in DPBS

overnight at 4°C.

CMFDA prelabelled Th cells were incubated with the isotype control antibody (30 μg/mL) for 30

minutes at 37°C (8% CO_2) prior to imaging (exact antibody concentrations are illustrated in the figure

legend). The cells were subsequently perfused on top of the pre-coated dishes at a concentration of

10^6/mL in MAM as previously described (12,36).

7.   In vitro under-flow T-cell migration assay

We studied the multi-step T-cell migration across monolayers of primary mouse brain

microvascular endothelial cells (pMBMECs) in a custom made microfluidic device under physiological

flow by in vitro live-cell imaging according to the previously established procedure (14,32). The

custom-made flow chamber was described in depth before (11,46). The flow chamber was placed on

top of the pMBMECs, previously cultured in Ibidi μ-Dish to confluency, and connected to the flow

system filled with migration assay medium (MAM) (12). During the accumulation phase, T cells at a

concentration between 55k/mL and 166k/mL were perfused on top for 5 minutes under low shear

stress of 0.1 dynes/cm^2, allowing them to settle on top of the pMBMEC monolayer. We used lower

T-cell concentration compared to previous studies to enable automated T-cell interaction analysis.

Afterward, the flow was increased to physiological levels with a flow shear stress of 1.5 dynes/cm^2 to

study post-arrest T-cell behavior on pMBMECs for 27 minutes (Figure 1A, B).

8.   Data acquisition

The timelapse imaging was performed during both accumulation and physiological flow phases

using phase-contrast imaging at a framerate of 6 frames/minute or 12 frames/minute, subsampled

to 6 frames/minute and resolution of 0.629 µm/pixel with a 10x objective. In this modality the

acquired images are gray scale, and the T cells, especially after the migration across the pMBMEC

monolayer, have a similar appearance to the pMBMECs, making the T-cell segmentation task very

challenging (Figure 1C). The data was acquired in tiles of 870 × 650 µm^2 (1389 × 1041 pixels) with an

overlap of 100 µm, leading to a total acquired image area of 3170 × 1220 µm^2. For each experiment,

the dataset consists of 30 timeframes of T-cell accumulation and 162 timeframes of dynamic T-cell

interactions with the pMBMEC monolayer under physiological flow. The timestep between

sequential timeframes for each tile is 10 s. The total acquired area is thus limited by the acquisition

speed or by the flow chamber size.

The HBMEC dataset was acquired as previously described (36). In brief, 2 tiles were acquired of

total size 3891 × 1504 pixels with pixel size 0.69 µm and framerate 6 frames/minute. The

accumulation phase was 4 minutes, and the physiological flow phase was 5 minutes.

The BMDM dataset was acquired similarly to the T cell interacting with the pMBMEC datasets. In

brief, one tile was acquired of size of 2048 × 1504 or 1388 × 1040 pixels with pixel size 0.69 or

0.629 µm and framerate 6 frames/minute. The accumulation phase lasted 6 minutes, and the

physiological flow phase was 25 minutes.

The EECM-BMEC dataset was acquired as previously described (36). In brief, 2 tiles were acquired

of total size 3892 × 1504 or 2638 × 1048 pixels with pixel size 0.69 or 0.629 µm and framerate

6 frames/minute.

The dataset of human T cells on immobilized recombinant BBB adhesion molecules was acquired

as previously described (36). In brief, 2 tiles were acquired of total size 3891 × 1504 pixels with pixel

size 0.69 µm and framerate 6 frames/minute. The accumulation phase was 4 minutes, and the

physiological flow phase was 5 minutes. In this dataset, there is only one compartment, thus

excluding any transmigration of the T cells. We, therefore, have employed the preprocessed

fluorescent images instead of the cell probability map and transmigration map produced by the

segmentation neural network. In this case, the centroids are defined as local maxima localized on

the preprocessed fluorescent images. The fluorescent images were preprocessed first by performing

Gaussian blur filtering in 2D with a standard deviation of 0.75. Next, thresholding was applied to

eliminate the background noise, clipping high brightness values (color values with abundance <0.3%

of threshold abundance), scaling to 0-255 8 bit, and saving in a lossless format. Finally, the 2 tiles

were stitched together by matching the integrated color distribution in the X direction in the

overlapping area.

9.   Manual analysis of T-cell migration

Manual cell analysis and tracking were performed according to the previously established

procedure using ImageJ software (ImageJ software, National Institute of Health, Bethesda, MD, USA)

(13). The number of arrested T cells was thus counted at timeframe 33 of the subset. The behavior

of arrested T cells was defined and expressed as fractions of arrested T cells set to 100% as follows:

• T cells that detached during the observation time ("detached")

• T cells that migrated out of the FoV detached during the observation time ("out of FoV")

• T cells that continuously crawled on the pMBMEC monolayer ("crawling")

• T cells that remained in the same location (displacement less than twice the cell size) while

actively interacting with the pMBMEC monolayer ("probing")

• T cells that crossed the pMBMEC monolayer with or without prior crawling ("crawling full

transmigration" and "probing full transmigration"). The event of T-cell transmigration across the

pMBMECs monolayer became obvious due to the change of appearance of the transmigrated part of

the T cells from phase bright (on top of the pMBMECs monolayer) to phase dark.

• T cells that partially crossed the pMBMEC monolayer, then retracted the protrusions and

continued to migrate above the monolayer ("crawling uncompleted transmigration" and "probing

uncompleted transmigration").

10. ^{UFM}Track performance evaluation

The inference of T cell masks and transmigration masks was performed on a dual-CPU Intel(R)

Xeon(R) CPU E5-2670 v3 @ 2.30GHz, 256GB RAM node equipped with 8 Graphical Processing Units

(GPU) NVIDIA GeForce GTX TITAN X / GeForce GTX 1080. Frame alignment and segmentation were

performed on an Intel(R) Core(TM) CPU i7-4771 @ 3.50GHz, 32GB RAM workstation with NVIDIA

GeForce GTX TITAN GPU. Cell tracking was performed on a dual-CPU Intel(R) Xeon(R) CPU E5-2643

v2 @ 3.50GHz, 256GB RAM workstation.

11. Performance analysis.

The statistical tests are performed with the Mann-Whitney U test. Performance of the

segmentation models was evaluated based on the pixel classification metrics evaluated using the

scikit-learn library (50). Specifically, the Average Precision is evaluated on the test set as the area

under the Precision-Recall curve, as the weighted mean of precisions achieved at each threshold,

with the change in recall from the previous threshold used as the weight. The segmentation

thresholds for the cell mask and the transmigrated cell mask predictions were selected to maximize

the F1 score.

Acknowledgments

We thank Kristina Berve for providing BMDM datasets for analysis with the [UFM]Track. We owe

sincere thanks to Dr. James McGrath and Danial Ahmad (Rochester University, NY, USA) for their

detailed and insightful feedback on our manuscript. This work was supported by an UniBe ID Grant

to MV, BE, and AA and the Microscopy Imaging Center (MIC) of the University of Bern.

Data Availability

The source code for the T-cell segmentation models, data preprocessing, training, performance

evaluation, and inference scripts are available on GitHub at

https://github.com/neworldemancer/UFMSegm. This repository also contains win64 binaries used

for the watershed-based segmentation of the predicted T cell probability maps and transmigration

probability maps.

The source code for the under-flow T-cell tracking and migration analysis, along with the scripts

used for performance evaluation of the framework, is available on GitHub at

https://github.com/neworldemancer/UFMTrack.

The training phase-contrast data with manual annotations, the reference datasets for histogram

normalization, and trained models are available on Zenodo:

[http://doi.org/10.5281/zenodo.7489557].

The datasets used for evaluation of our framework are available on Zenodo:

[http://doi.org/10.5281/zenodo.7489972, http://doi.org/10.5281/zenodo.7489984,

http://doi.org/10.5281/zenodo.7489996, http://doi.org/10.5281/zenodo.7490012,

http://doi.org/10.5281/zenodo.7490020, http://doi.org/10.5281/zenodo.7490029].

References

1. Nishihara H, Soldati S, Mossu A, Rosito M, Rudolph H, Muller WA, et al. Human CD4+ T cell
subsets differ in their abilities to cross endothelial and epithelial brain barriers in vitro. Fluids
Barriers CNS. 2020 Feb 3;17(1):1–18.

2. Marchetti L, Engelhardt B. Immune cell trafficking across the blood-brain barrier in the
absence and presence of neuroinflammation. Vascular Biology [Internet]. 2020 Apr 10 [cited
2022 Oct 28];2(1):H1–18. Available from:
https://vb.bioscientifica.com/view/journals/vb/2/1/VB-19-0033.xml

3. Nourshargh S, Alon R. Leukocyte Migration into Inflamed Tissues. Immunity. 2014 Nov
20;41(5):694–707.

4. Nourshargh S, Hordijk PL, Sixt M. Breaching multiple barriers: leukocyte motility through
venular walls and the interstitium. Nature Reviews Molecular Cell Biology 2010 11:5. 2010
May;11(5):366–78.

5. Engelhardt B, Vajkoczy P, Weller RO. The movers and shapers in immune privilege of the CNS
[Internet]. Vol. 18, Nature Immunology. Nature Publishing Group; 2017 [cited 2019 May 3]. p.
123–31. Available from: http://www.nature.com/articles/ni.3666

6. Bartholomäus I, Kawakami N, Odoardi F, Schläger C, Miljkovic D, Ellwart JW, et al. Effector T
cell interactions with meningeal vascular structures in nascent autoimmune CNS lesions.
Nature 2009 462:7269. 2009 Oct 14;462(7269):94–8.

7. Lyck R, Engelhardt B. Going Against the Tide – How Encephalitogenic T Cells Breach the
Blood-Brain Barrier. J Vasc Res. 2012 Oct;49(6):497–509.

8. Dias MC, Quesada AO, Soldati S, Bösch F, Gruber I, Hildbrand T, et al. Brain endothelial
tricellular junctions as novel sites for T cell diapedesis across the blood-brain barrier. J Cell
Sci. 2021 Apr 1;134(8).

9. Steinman L, Zamvil SS. Virtues and pitfalls of EAE for the development of therapies for
multiple sclerosis. Trends Immunol. 2005;26(11):565–71.

10. Singer BA. The role of natalizumab in the treatment of multiple sclerosis: benefits and risks.
Ther Adv Neurol Disord. 2017 Sep 1;10(9):327–36.

11. Lyck R, Nishihara H, Aydin S, Soldati S, Engelhardt B. Modeling Brain Vasculature Immune
Interactions In Vitro. Cold Spring Harb Perspect Med [Internet]. 2023 Sep 12 [cited 2023 Jul
19];13(9):a041185. Available from:
http://perspectivesinmedicine.cshlp.org/lookup/doi/10.1101/cshperspect.a041185

12. Steiner O, Coisne C, Cecchelli R, Boscacci R, Deutsch U, Engelhardt B, et al. Differential Roles
for Endothelial ICAM-1, ICAM-2, and VCAM-1 in Shear-Resistant T Cell Arrest, Polarization,
and Directed Crawling on Blood–Brain Barrier Endothelium. The Journal of Immunology. 2010
Oct 15;185(8):4846–55.

13.   Rudolph H, Klopstein A, Gruber I, Blatti C, Lyck R, Engelhardt B. Postarrest stalling rather than
crawling favors CD8+ over CD4+ T-cell migration across the blood–brain barrier under flow in
vitro. Eur J Immunol [Internet]. 2016 Sep 1 [cited 2020 Mar 9];46(9):2187–203. Available
from: https://onlinelibrary.wiley.com/doi/full/10.1002/eji.201546251

14.   Abadier M, Jahromi NH, Alves LC, Boscacci R, Vestweber D, Barnum S, et al. Cell surface levels
of endothelial ICAM-1 influence the transcellular or paracellular T-cell diapedesis across the
blood–brain barrier. 2015 Apr 1 [cited 2019 Apr 19];45(4):1043–58. Available from:
http://doi.wiley.com/10.1002/eji.201445125

15.   Shulman Z, Shinder V, Klein E, Grabovsky V, Yeger O, Geron E, et al. Lymphocyte Crawling and
Transendothelial Migration Require Chemokine Triggering of High-Affinity LFA-1 Integrin.
Immunity. 2009 Mar 20;30(3):384–96.

16.   Arts JJG, Mahlandt EK, Grönloh MLB, Schimmel L, Noordstra I, Gordon E, et al. Endothelial
junctional membrane protrusions serve as hotspots for neutrophil transmigration. Elife. 2021
Aug 1;10.

17.   He K, Gkioxari G, Dollár P, Girshick R. Mask R-CNN. IEEE Trans Pattern Anal Mach Intell
[Internet]. 2017 Mar 20 [cited 2024 Jul 23];42(2):386–97. Available from:
https://arxiv.org/abs/1703.06870v3

18.   Wang CY, Bochkovskiy A, Liao HYM. YOLOv7: Trainable bag-of-freebies sets new state-of-the-
art for real-time object detectors. 2022 Jul 6 [cited 2024 Jul 23];7464–75. Available from:
https://arxiv.org/abs/2207.02696v1

19.   Carion N, Massa F, Synnaeve G, Usunier N, Kirillov A, Zagoruyko S. End-to-End Object
Detection with Transformers. Lecture Notes in Computer Science (including subseries Lecture
Notes in Artificial Intelligence and Lecture Notes in Bioinformatics) [Internet]. 2020 May 26
[cited 2024 Jul 23];12346 LNCS:213–29. Available from: https://arxiv.org/abs/2005.12872v3

20.   Schmidt U, Weigert M, Broaddus C, Myers G. Cell Detection with Star-convex Polygons.
Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence
and Lecture Notes in Bioinformatics) [Internet]. 2018 Jun 9 [cited 2024 Jul 23];11071
LNCS:265–73. Available from: https://arxiv.org/abs/1806.03535v2

21.   Ronneberger O, Fischer P, Brox T. U-Net: Convolutional Networks for Biomedical Image
Segmentation. 2015 May 18 [cited 2017 Oct 25]; Available from:
http://arxiv.org/abs/1505.04597

22.   Stringer C, Wang T, Michaelos M, Pachitariu M. Cellpose: a generalist algorithm for cellular
segmentation. Nature Methods 2020 18:1 [Internet]. 2020 Dec 14 [cited 2024 Jul
23];18(1):100–6. Available from: https://www.nature.com/articles/s41592-020-01018-x

23.   Allan DB, Caswell T, Keim NC, van der Wel CM, Verweij RW. Trackpy. 2021 Apr 13 [cited 2022
Oct 29]; Available from: https://zenodo.org/record/4682814

24.   Tsai HF, Gajda J, Sloan TFW, Rares A, Shen AQ. Usiigaci: Instance-aware cell tracking in stain-
free phase contrast microscopy enabled by machine learning. SoftwareX. 2019 Jan 1;9:230–7.

25.   Ershov D, Phan MS, Pylvänäinen JW, Rigaud SU, Le Blanc L, Charles-Orszag A, et al. TrackMate
7: integrating state-of-the-art segmentation algorithms into tracking pipelines. Nature

Methods 2022 19:7 [Internet]. 2022 Jun 2 [cited 2022 Oct 29];19(7):829–32. Available from:
https://www.nature.com/articles/s41592-022-01507-1

26. Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, et al. ImageJ2: ImageJ
for the next generation of scientific image data. BMC Bioinformatics [Internet]. 2017 Nov 29
[cited 2024 Jul 4];18(1):1–26. Available from:
https://link.springer.com/articles/10.1186/s12859-017-1934-z

27. Holme B, Bjørnerud B, Pedersen NM, de la Ballina LR, Wesche J, Haugsten EM. Automated
tracking of cell migration in phase contrast images with CellTraxx. Scientific Reports 2023
13:1 [Internet]. 2023 Dec 27 [cited 2024 Jul 23];13(1):1–20. Available from:
https://www.nature.com/articles/s41598-023-50227-9

28. Ulicna K, Vallardi G, Charras G, Lowe AR. Automated Deep Lineage Tree Analysis Using a
Bayesian Single Cell Tracking Approach. Front Comput Sci. 2021 Oct 20;3:92.

29. Abadi M, Barham P, Chen J, Chen Z, Davis A, Dean J, et al. TensorFlow: A system for large-
scale machine learning.

30. Schiegg M, Hanslovsky P, Kausler BX, Hufnagel L, Hamprecht FA. Conservation Tracking. 2013;

31. Perron L, Furnon V. OR-Tools [Internet]. Available from:
https://developers.google.com/optimization/

32. Haghayegh Jahromi N, Marchetti L, Moalli F, Duc D, Basso C, Tardent H, et al. Intercellular
Adhesion Molecule-1 (ICAM-1) and ICAM-2 Differentially Contribute to Peripheral Activation
and CNS Entry of Autoaggressive Th1 and Th17 Cells in Experimental Autoimmune
Encephalomyelitis. Front Immunol. 2020 Jan 14;10:3056.

33. Mossu A, Rosito M, Khire T, Li Chung H, Nishihara H, Gruber I, et al. A silicon nanomembrane
platform for the visualization of immune cell trafficking across the human blood–brain barrier
under flow. Journal of Cerebral Blood Flow and Metabolism. 2019 Mar 1;39(3):395–410.

34. Clockify. Clockify™ - 100% FREE Time Tracking Software [Internet]. 2022 [cited 2022 Oct 23].
Available from: https://clockify.me/

35. Nishihara H, Gastfriend BD, Kasap P, Palecek SP, Shusta E V., Engelhardt B. Differentiation of
human pluripotent stem cells to brain microvascular endothelial cell-like cells suitable to
study immune cell interactions. STAR Protoc. 2021 Jun 18;2(2).

36. Soldati S, Bär A, Vladymyrov M, Glavin D, McGrath JL, Gosselet F, et al. High levels of
endothelial ICAM-1 prohibit natalizumab mediated abrogation of CD4+ T cell arrest on the
inflamed BBB under flow in vitro. J Neuroinflammation. 2023 May 23;20(1):123.

37. Jae Oh D, Min Lee G, Francis K, Palsson BO. Phototoxicity of the Fluorescent Membrane Dyes
PKH2 and PKH26 on the Human Hematopoietic KG1a Progenitor Cell Line.

38. Saetzler RK, Jallo J, Lehr HA, Philips CM, Vasthare U, Arfors KE, et al. Intravital Fluorescence
Microscopy: Impact of Light-induced Phototoxicity on Adhesion of Fluorescently Labeled
Leukocytes. The Journal of Histochemistry & Cytochemistry. 1997;45(4):505–13.

39. Purschke M, Rubio N, Held KD, Redmond RW. Phototoxicity of Hoechst 33342 in time-lapse
fluorescence microscopy. Photochem Photobiol Sci. 2010;9(12):1634–9.

40. Wimmer I, Tietz S, Nishihara H, Deutsch U, Sallusto F, Gosselet F, et al. PECAM-1 stabilizes
blood-brain barrier integrity and favors paracellular T-cell diapedesis across the blood-brain
barrier during neuroinflammation. Front Immunol. 2019 Apr 5;10(APR):447986.

41. Aydin S, Pareja J, Schallenberg VM, Klopstein A, Gruber T, Page N, et al. Antigen recognition
detains CD8+ T cells at the blood-brain barrier and contributes to its breakdown. Nature
Communications 2023 14:1 [Internet]. 2023 May 30 [cited 2023 Jul 31];14(1):1–20. Available
from: https://www.nature.com/articles/s41467-023-38703-2

42. Caron M, Touvron H, Misra I, Jegou H, Mairal J, Bojanowski P, et al. Emerging Properties in
Self-Supervised Vision Transformers. Proceedings of the IEEE International Conference on
Computer Vision [Internet]. 2021 Apr 29 [cited 2024 Jun 24];9630–40. Available from:
https://arxiv.org/abs/2104.14294v2

43. Archit A, Nair S, Khalid N, Hilt P, Rajashekar V, Freitag M, et al. Segment Anything for
Microscopy. bioRxiv [Internet]. 2023 Aug 22 [cited 2024 Jun 24];2023.08.21.554208. Available
from: https://www.biorxiv.org/content/10.1101/2023.08.21.554208v1

44. Kraus O, Kenyon-Dean K, Saberian S, Fallah M, McLean P, Leung J, et al. Masked
Autoencoders for Microscopy are Scalable Learners of Cellular Biology. 2024 Apr 16 [cited
2024 Jun 24]; Available from: https://arxiv.org/abs/2404.10242v1

45. Xie R, Pang K, Bader GD, Wang B. MAESTER: Masked Autoencoder Guided Segmentation at
Pixel Resolution for Accurate, Self-Supervised Subcellular Structure Recognition. In 2023. p.
3292–301.

46. Coisne C, Lyck R, Engelhardt B. Live cell imaging techniques to study T cell trafficking across
the blood-brain barrier in vitro and in vivo. Fluids Barriers CNS [Internet]. 2013 Jan 21 [cited
2020 Mar 9];10(1):7. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23336847

47. Engen SA, Rukke HV, Becattini S, Jarrossay D, Blix IJ, Petersen FC, et al. The Oral Commensal
Streptococcus mitis Shows a Mixed Memory Th Cell Signature That Is Similar to and Cross-
Reactive with Streptococcus pneumoniae. PLoS One. 2014 Aug 13;9(8):e104306.

48. Sallusto F, Lenig D, Mackay CR, Lanzavecchia A. Flexible Programs of Chemokine Receptor
Expression on Human Polarized T Helper 1 and 2 Lymphocytes. Journal of Experimental
Medicine. 1998 Mar 16;187(6):875–83.

49. Zielinski CE, Mele F, Aschenbrenner D, Jarrossay D, Ronchi F, Gattorno M, et al. Pathogen-
induced human TH17 cells produce IFN-γ or IL-10 and are regulated by IL-1β. Nature 2012
484:7395. 2012 Apr 1;484(7395):514–8.

50. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn:
Machine Learning in Python. Journal of Machine Learning Research. 2011;12:2825–30.