# Microbial strong organic ligand production is tightly coupled to iron in hydrothermal plumes

Colleen L. Hoffman[1,2,3*] and Patrick J. Monreal[3,4*], Justine B. Albers[5], Alastair J.M. Lough[6], Alyson E. Santoro[5], Travis Mellett[3,7], Kristen N. Buck[7,8], Alessandro Tagliabue[9], Maeve C. Lohan[6], Joseph A. Resing[1,2,3], Randelle M. Bundy[3]

[1]Joint Institute for the Study of Atmosphere and Ocean, University of Washington, 3737 Brooklyn
       Avenue NE, Seattle, WA 98195, USA

[2]Cooperative Institute for Climate, Ocean, and Ecosystem Studies, University of Washington,
       3737 Brooklyn Avenue NE, Seattle, WA 98195, USA

[3]School of Oceanography, University of Washington, 1501 NE Boat Street, Seattle, WA 98195,
       USA

[4]Earth Systems Program, Stanford University, 473 Via Ortega, Stanford, CA 94305,
       USA

[5]Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara,
       CA 93106, USA

[6]Department of Ocean and Earth Sciences, National Oceanography Centre, University of
       Southampton, European Way, Southampton SO14 3ZH, United Kingdom

[7]College of Marine Science, University of South Florida, 140 7th Avenue South, St. Petersburg,
       FL, 33701, USA

[8]College of Earth, Oceans, and Atmospheric Sciences, Oregon State University, 2651 SW Orchard Ave,
       Corvallis, OR, 97331, USA

[9]Department of Earth, Ocean, and Ecological Sciences, University of Liverpool, 4 Brownlow
       Street, Liverpool l69 3GP, United Kingdom

*Colleen L. Hoffman and Patrick J. Monreal

**Email:** clhoffma@gmail.com, pmonreal@uw.edu

**Abstract.** Hydrothermal vents have emerged as an important source of iron to seawater, yet only a subset of iron is soluble and persists long enough to be available for surface biological uptake. The longevity and solubility of iron in seawater is governed by strong organic ligands, like siderophores, that are produced by marine microorganisms and are a part of the ocean's dissolved iron-binding ligand pool. These ligands have been hypothesized to aid in the persistence of dissolved iron in hydrothermal environments. To explore this hypothesis, we measured iron, iron-binding ligands, and siderophores from 11 geochemically distinct sites along a 1,700 km section of the Mid-Atlantic Ridge. Siderophores were found in hydrothermal plumes at all sites, with proximity to the vent playing an important role in dictating siderophore types and diversity. The notable presence of amphiphilic siderophores may point to microbial utilization of siderophores to access particulate hydrothermal iron, and the exchange of dissolved and particulate iron. The tight coupling between strong ligands and dissolved iron within neutrally buoyant plumes across six distinct hydrothermal environments, and the presence of dissolved siderophores with siderophore-producing microbial genera, suggests that biological production of siderophores exerts a key control on hydrothermal dissolved iron concentrations.

## 1 Introduction

Over the last few decades, observations and modelling efforts have increased our understanding about the critical role organic ligands play in the cycling, transport, and utilization of trace metals (Tagliabue et al., 2017; Buck et al., 2018; Bundy et al., 2018; Moore et al., 2021). Strong iron-binding organic ligands ($L_1$) are a heterogeneous mixture of microbially produced compounds that are operationally classified based on their binding strength with iron (Fe) (log $K_{Fe',FeL}^{cond} > 12$), and are thermodynamically favored to complex and stabilize external sources of Fe to prevent its scavenging and removal. For example, in high dissolved and particulate Fe estuarine systems, only the dissolved Fe (dFe) bound to the strongest Fe-binding ligands remains in solution (Bundy et al., 2015; Buck et al., 2007) and is accessible for downstream biological uptake.

Siderophores are the strongest known Fe-binding organic ligands. They are produced by bacteria and fungi to facilitate Fe uptake and solubilization of otherwise inaccessible phases in the marine environment (Butler, 2005; Manck et al., 2022). They have primarily been considered as an important microbial strategy for Fe acquisition in the low-Fe surface ocean (Vraspir and Butler, 2009; Butler, 2005). However, siderophore uptake and biosynthesis genes were observed in >70% of Fe-related bacterial transcripts at Guaymas Basin (Li et al., 2014), have been identified in oxygen-deficient zones (Moore et al., 2021), and is a common Fe acquisition strategy within terrestrial and pathogenic ecosystems (Sandy and Butler, 2009) where Fe is orders of magnitude higher than seawater.

Although other unknown strong Fe-binding ligands have been observed in hydrothermal plumes (Buck et al., 2018) and throughout the deep ocean as well as siderophores observed below the euphotic zone (Bundy et

al., 2018), no previous studies have ever directly characterized siderophores in hydrothermal systems. Some

form of 'stabilizing agent' (i.e. ligands) has been proposed for the long-range transport of hydrothermal Fe

into the ocean interior. The role of strong Fe-binding ligands in hydrothermal dFe transport represents an

important gap knowledge gap with how hydrothermal vents may impact the ocean dFe inventory (Resing et

al., 2015). Here, for the first time, we identified siderophores and siderophore-producing microbes in 11

geochemically distinct hydrothermal environments along the slow-spreading (20-50 mm/yr) Mid-Atlantic

Ridge (MAR), including four black smokers (high temperature, high Fe), four off-axis sites, one diffuse vent

(low temperature, low Fe), one alkaline vent (pH 9-11, very low Fe), and one non-vent fracture zone, to

understand whether microbial ligand production impacts the supply of hydrothermal dFe to the ocean.

Overall, our results show microbially-produced siderophores were present in all sites, and that strong $L_1$

ligands are tightly coupled to hydrothermal dFe in this system. Strong organic ligands produced by bacteria

thus play a key role in deep ocean Fe delivery from hydrothermal systems.

**2 Results and Discussion**

**2.1 The role of iron-binding ligands and siderophores in hydrothermal plumes**

Strong organic Fe-binding ligands, or $L_1$ ligands, are important for stabilizing Fe in hydrothermal plumes

(Tagliabue et al., 2017; Resing et al., 2015; Buck et al., 2018). The average binding strength and

concentration of organic Fe-binding ligands were quantified in multiple vent systems that spanned a wide

range in dFe concentrations (0.41-90.3 nM) and underlying vent geology. Over 99% of dFe in the neutrally

buoyant plumes were complexed by $L_1$ ligands and the ligands were almost always completely saturated with

dFe, meaning Fe-free 'excess' $L_1$ ligands capable of binding additional Fe were present in low concentrations

(< 1 nM; **Fig. S1**). As a result, dFe concentrations were tightly coupled to $L_1$ ligands in a nearly 1:1 ratio

(**Fig. 1d**), similar to previous studies in other neutrally buoyant plumes (**Fig. 1e**) (Lough et al., 2022; Buck

et al., 2018, 2015). The strong coupling between dFe and ligands was only observed at sites where $L_1$ ligands

were detected. Some sampling locations, such as in the buoyant plume, contained high concentrations of

weaker ligands (log $K_{Fe',FeL}^{cond} < 12$, **Table S2**) with no correlation to dFe.

Our results indicate that $L_1$ ligands uniquely set the dFe concentration in neutrally buoyant plumes. A similar

control of dFe concentrations by $L_1$ ligands has been previously observed in rivers (Buck et al., 2007) and

aerosol solubility experiments (Fishwick et al., 2014). One explanation is that both the dFe and $L_1$ ligands

originate from the vent fluids themselves, yielding a tightly coupled hydrothermal endmember. However, the

concentration of $L_1$ ligands did not correlate with excess mantle Helium-3 ($^3He_{xs}$, **Fig S2**, **Table S2**) (Lough

et al., 2022), a nearly conservative tracer of the mixing of hydrothermal fluids with seawater (Buck et al.,

2018). These results suggest the $L_1$ ligands were not sourced from the vent fluids along with dFe. All known

sources of $L_1$ ligands are biologically produced. Therefore, the $L_1$ ligands observed here could be sourced

either from bacteria that produced them in the surrounding deep ocean seawater that was then entrained, local

production from vent-biota and/or microbial mats, diffusion from microbial production in sediments, or *in-*
*situ* production by bacteria within the neutrally buoyant plume (Mellett et al., *submitted*).

Microbial organic ligand production is thought to tightly control global Fe inventories in both hydrothermal
plumes (Cowen and Bruland, 1985; Cowen et al., 1990) and the open ocean (Lauderdale et al., 2020).
Siderophores are operationally defined as $L_1$ ligands by their binding strength (Moore et al., 2021; Manck et
al., 2022) and have been proposed as important $L_1$ ligands in hydrothermal plumes (Li et al., 2014), though
they have never been directly measured. We used state-of-the-art liquid chromatography coupled to
electrospray ionization mass spectroscopy (Boiteau et al., 2016) in a targeted approach to identify discrete
components of the $L_1$ ligands and to search for known siderophores. We observed a large diversity of
siderophores with high confidence in every vent site using mass-to-charge ratio (*m/z*), MS/MS spectra, and
specific chromatographic characteristics (**Fig. 2a**). Relative peak areas as a proxy for concentrations of
putative siderophores also significantly correlated with dFe, as observed with dFe and $L_1$ ligands (**Fig. 2b**).

Siderophores were present in concentrations similar to the surface ocean (Boiteau et al., 2016; Moore et al.,
2021; Park et al., 2022; Bundy et al., 2018), and comprised 0.01-0.4% of the total $L_1$ ligands (**Table 1**). This
is likely a significant underestimate of siderophore contributions to the $L_1$ ligand pool due to analytical
constraints in identifying unknown siderophores. That is, known siderophores represent a small fraction of
what is expected to be produced in nature (Hider and Kong, 2010). In addition, most siderophores are not
commercially available to use as standards, and individual siderophores have different ionization or
extraction efficiencies. We also restricted our reporting to compounds only identified with very high
confidence (**Fig 2a**, **S3**). The extraction efficiency for the solid phase extraction technique is around 10% for
bulk Fe-binding organics (Bundy et al., 2018) and 40% for a siderophore standard (Waska et al., 2015).
Correcting for extraction efficiency of the identified siderophores yields contributions of 1-4% of the $L_1$ pool.
We are inevitably missing other naturally occurring unknown compounds. Regardless of the small percentage
contribution to total $L_1$ ligands, it is evident that microbially produced siderophores are ubiquitous across all
vent sites.

The high diversity of siderophores across a huge range of hydrothermal vent systems reveals several
surprising aspects of Fe cycling. The biosynthesis of a siderophore is energy-intensive and is regulated by Fe
concentration in the surrounding environment. Siderophore presence suggests that bacteria are producing
these compounds despite the overall higher Fe concentrations in the deep ocean and in hydrothermal plumes.
Consistent with siderophore utilization in terrestrial ecosystems, one hypothesis is that siderophore
production is beneficial to bacteria in the plumes for transforming Fe from otherwise inaccessible forms,
such as particulate nanopyrites or Fe oxyhydroxides. The identification of siderophores — and their
relationship with dFe — provides compelling evidence that microbial production of ligands is responsible
for some portion of the tight coupling between $L_1$ and dFe in hydrothermal systems along the MAR.

**2.2 Identity of siderophores in hydrothermal systems**

Marine ligand composition changes with environmental gradients (Boiteau et al., 2016), making the structure and functional groups of siderophores identified in and around hydrothermal samples of particular interest. Samples collected from on-axis spreading centers contained the highest dFe concentrations (> 20 nM) and wider variety of siderophores than samples from fracture zones, diffuse, and off-axis sites (dFe ≤ 1 nM). The greatest number of distinct siderophores were identified at Lucky Strike, Broken Spur, Rainbow, and TAG (**Fig. 2**). On average, 13 compounds were identified with high confidence per on-axis spreading center sample, compared with 5 per diffuse/fracture zone sample, and 2.5 per off-axis sample (**Fig. 2b, Fig. S4**). Mixed-type siderophores — containing different moieties that bind to Fe(III) — were common at all sites. Hydroxamates were identified at and around spreading centers, yet none of these were detected with high confidence in samples from diffuse/fracture zones (**Fig. S4**). Vent type and proximity played a role in the diversity and abundance of siderophore types produced, likely related to the diversity of the microbial community and/or unique Fe acquisition strategies across sites.

Low Fe surface waters in the ocean have higher concentrations of amphiphilic siderophores compared to high Fe waters and terrestrial systems (Boiteau et al., 2016). Amphiphilic siderophores have long hydrocarbon tails that can be embedded into the lipid bilayer of the bacterial cell membrane providing a mechanism to shuttle Fe into the cell and prevent diffusive loss (Martinez et al., 2003). Amphiphilic siderophores accounted for 57% of the siderophores in our samples (**Fig. S5**), supporting the ubiquity of amphiphilic siderophores in the marine environment (Butler and Theisen, 2010). It was surprising they were so common, due to the elevated Fe concentrations observed relative to the Fe-poor surface ocean. Amphiphilic siderophores were found in concentrations between 0.3-4.7 pM, with highest found at Rainbow (**Fig. 2d**, **Table S5**). These concentrations were similar to those observed in the upper ocean (Boiteau et al., 2016; Bundy et al., 2018; Boiteau et al., 2019). Marine bacteria produce suites of amphiphilic siderophores as a way to adapt to the change in hydrophilicity in the surrounding environment (Sandy and Butler, 2009; Homann et al., 2009). Unlike in the surface ocean where amphiphilic siderophores are observed in Fe-limited regions (Boiteau et al., 2016), amphiphilic siderophores in plumes could be a way for bacteria to access Fe as they are physically transported and cope with strong chemical gradients, similar to the production of multiple siderophores in terrestrial and pathogenetic systems as a means to access inorganic particulate Fe for cellular uptake and storage (Hider and Kong, 2010).

**2.3 Microbial sources of siderophores in hydrothermal plumes**

Microbial production of siderophores is a strategy for organisms to adapt or compete with others for Fe (Sandy and Butler, 2009). We examined microbial community composition around Rainbow (St. 11, 17) and

Lucky Strike (St. 7; **Table 1, Table S1**) using 16S rRNA gene-based amplicon sequencing to detect bacteria

with the metabolic potential to synthesize siderophores (**Fig. 3, S11**), where the presence of taxa encoding

siderophore biosynthetic gene clusters indicates whether the microbial community is genetically capable of

producing the compounds we observed. Bacterial genera containing known siderophore-producers were

found at all three MAR sites examined, and putative siderophore-producers represented 3-20% of the relative

abundance of each community (**Fig. 3**). Putative siderophore-producers were more abundant in the 3 µm

(particle-attached) size fraction than in the 0.2 µm (free-living) fraction, suggesting siderophore production

is more common in particle-associated bacteria in hydrothermal environments.

We found microbial genera in our samples that can produce a subset of the siderophores identified here,

including ferrioxamines, vibrioferrin, and acinetoferrin (Butler, 2005; Vraspir and Butler, 2009; Moore et al.,

2021; Bundy et al., 2018; Boiteau et al., 2016). Genera with the genetic potential to produce ferrioxamines

were present at all three sites, while those known to produce vibrioferrin were present at Lucky Strike and

Rainbow, and those producing acinetoferrin were also present at Rainbow (**Table S1, S6**). Mycobactins were

detected with high confidence in every sample of this study, and genes encoding mycobactin have been

detected in a cultured organism from a hydrothermal system (Gu et al., 2019), but no mycobactin producers

were identified in this study. We detected woodybactin D with high confidence in 5 out of 11 sites. Although

these biosynthetic genes were not identified in any of the genera observed, woodybactin D is a carboxylate

siderophore isolated from *Shewanella* (Carmichael et al., 2019), and groups of deep-sea *Shewanella* (Kato

and Nogi, 2001) were found in the dataset (**Fig. S11**). The biosynthesis genes for many of the siderophores

identified are unknown. Thus, finding genera capable of producing only a subset of the siderophores

characterized is not surprising. The observation that a significant portion of the *in-situ* microbial community

is capable of synthesizing siderophores (**Fig 3**) suggests that siderophore production is more widespread in

the marine environment and deep ocean than previously believed.

Although siderophores are often assumed to be associated primarily with low Fe conditions, evidence that

siderophores are ubiquitous in the marine environment — including higher Fe ones — has been increasing

(Park et al., 2023). The higher dFe associated with hydrothermal plumes may still not be high enough to

suppress siderophore production due to the Fe requirements of bacteria (Tortell et al., 1996). It is also likely

that in hydrothermal plumes not all of the Fe is bio-accessible. For example, soil microbes secrete

siderophores to solubilize particulate Fe (Crowley et al., 1991). Similar processes could be occurring in

hydrothermal plumes, where Fe mineral phases are common (Hoffman et al., 2020). It is telling that most

siderophore-producing genera were found to be particle-associated (**Fig. 3**), providing additional evidence

that siderophores might be produced to solubilize particulate Fe. Further work that assesses why bacteria are

producing siderophores in neutrally buoyant plumes will be important for understanding the role of these

complexes in facilitating the exchange of Fe between dissolved and particulate phases (Fitzsimmons et al.,

2017), and how Fe-siderophore complexes may be stabilizing dFe in hydrothermal plumes.

Organic Fe-binding ligands have been implicated in playing a critical role in the preservation and transport

of hydrothermal Fe into the ocean interior (Hoffman et al., 2018; Resing et al., 2015; Fitzsimmons et al.,

2017; Toner et al., 2009; Bennett et al., 2011, 2008; Buck et al., 2018; Sander and Koschinsky, 2011). In this

work, $L_1$ ligands were tightly coupled to dFe in neutrally buoyant hydrothermal plumes along the MAR and

found to be microbially produced. For the first time, specific siderophores were identified and demonstrated

a bacterial source of $L_1$ ligands being produced in response to external hydrothermal Fe inputs. Most of the

bacteria putatively capable of synthesizing siderophores were particle-associated. This adds to a growing

body of evidence that bacteria are using $L_1$ ligands or siderophores to access particulate Fe pools, which is a

key mechanism for controlling dFe in neutrally buoyant plumes. Exploring whether the Fe-organic ligand

production is tightly coupled across additional hydrothermal vent systems will aid in constraining the

biogeochemical importance of microbial feedbacks in impacting the hydrothermal dFe supply to the deep

ocean.

**Appendix: Materials and Methods**

**Sampling and cruise transect**

Samples were collected as part of the 2017-2018 U.K. GEOTRACES GA13 section cruise along the Mid-

Atlantic Ridge. Water samples from 11 venting and near venting locations were collected using a Seabird

911 conductivity, temperature, and depth (CTD) titanium rosette using conducting Kevlar wire with an

oxidation-reduction potential (ORP) sensor to detect plumes. Teflon coated OTE (Ocean Test Equipment)

bottles were pressurized to approximately 7 psi with 0.2 μm filtered air using an oil free compressor. A

Sartobran 300 (Sartorius) filter capsule (0.2 μm) was used to collect filtered seawater samples into clean 250

mL LDPE sample bottles. Bottles and caps were rinsed 3 times with the filtered sample before being filled.

Samples were stored frozen at -20˚C for Fe-organic ligand characterization by voltammetry and mass

spectrometry.

**Fe-binding ligand concentration and binding strengths Competitive Ligand Exchange-Adsorptive**
**Cathodic Stripping Voltammetry**

Fe-binding ligand concentrations and binding strengths were determined by competitive ligand exchange-

adsorptive cathodic stripping voltammetry (CLE-ACSV) with a BAS*i* controlled growth mercury electrode

(CGME) with an Ag/AgCl$^-$ reference electrode and platinum auxiliary electrode (Bioanalytical Systems

Incorporated). Using previously established methods (Buck et al., 2015, 2018; Bundy et al., 2018; Abualhaija

and van den Berg, 2014; Hawkes et al., 2013b), 40 frozen filtrate (<0.2 μm) samples with dFe concentrations

between 0.41-11.67 nM (**Table S1-S2**) were thawed in a 4˚C fridge prior to analysis. A 15-point titration

curve was analyzed for each sample. Briefly, within each titration, every point sequentially received 10 mL

of sample, 7.5 μM of borate-ammonium buffer, 10 μM salicylaldoxime (SA) added ligand, and a dFe

addition. Data was collected using the *Epsilon Eclipse Electrochemical Analyzer* (v.213) with a deposition

time of 120 seconds and analyzed using *ElectroChemical Data Software* (v2001-2014) and *ProMCC* (v2008-

2018) to determine peak areas and Fe-binding ligand parameters, respectively.

**Reverse Titration-CLE-ACSV**

Reverse titration-CLE-ACSV (RT-CLE-ACSV) (Hawkes et al., 2013a) was completed on 10 samples from

Broken Spur, and TAG hydrothermal vent fields with dFe concentrations between 19.01-90.25 nM (**Table**

**S1-S2**). Briefly, a 10-point titration curve was analyzed for each sample with each titration point consisting

of 10 mL of sample buffered with 7.5 µM boric acid and the competitive ligand 1-nitroso-2-napthol (NN)

additions. All samples were analyzed on a BAS*i* Controlled Growth Mercury Electrode (CGME) with the

*Epsilon Eclipse Electrochemical Analyzer* (v.213) and deposition time of 120 seconds. For each sample,

competitive ligand NN additions were 0.5, 1, 2, 3, 4, 6, 9, 15, 20, and 40 µM. Samples were equilibrated

overnight and purged with $N_2$ (99.99%) for 5 minutes before analysis. At the end of each titration, three Fe

additions (3-15 nM) were added to the final titration point to get the total concentration of Fe in equilibrium

with ligands. Data was analyzed using *ElectroChemical Data Software* (v2001-2014) to acquire peak areas

and a package in R using the model parameters of $\beta_{FeNN3} = 5.12 \times 10^{16}$, $\chi_{min} = 0.8$, $\chi_{max} = 0.9$, and *c1high* =

0.75 to determine the Fe-binding ligand parameters (Hawkes et al., 2013a). These parameters were chosen

based on the recommendations for undersaturated samples and titrations curves where $ip_{max}$ was not reached

(Hawkes et al., 2013a). All other parameters within the model we kept at the default values.

**Fe-binding organic ligand quantification and characterization**

In addition to determining the total concentrations of strong Fe-binding ligands ($L_1$), we also identified and

quantified siderophores that contributed to the $L_1$ ligand pool. Between 0.65-1.5 L of 0.2 µm filtered seawater

pooled from ligand samples at each site (described above) was pumped slowly (15-20 mL min$^{-1}$) onto a

polystyrene-divinylbenzene (Bond Elut ENV) solid phase extraction (SPE) column (Bundy et al., 2018;

Boiteau et al., 2016). SPE columns were rinsed with MilliQ and stored at -20ºC until analysis. For the

analytical measurements, samples were thawed in the dark, eluted in 12 mL of distilled methanol, and dried

down to between 0.2-0.5 mL of sample eluent (**Table S1**). Aliquots were analyzed by reverse-phase liquid

chromatography (LC) on a trace metal clean bio-inert LC (Thermo Dionex 3000 NCS). The LC was

interfaced with an electrospray ionization-mass spectrometer (ESI-MS; Thermo Q-Exactive HF) to identify

and quantify the compounds based on accurate mass (MS$^1$) and the fragmentation (MS$^2$) data (Bundy et al.,

2018; Boiteau et al., 2016). MSconvert (Proteowizard) was used to convert MS data to an open source

mzxML format, and two stages of data processing were conducted using modified versions of previously

reported R scripts (Bundy et al., 2018; Boiteau et al., 2016). In the first stage, mzxML files were read into R

using new package "RaMS" (Kulmer and Ingalls, 2022), and extracted ion chromatograms (EICs) were

generated for each targeted *m/z* of interest from an in-house database of siderophores. The *m/z* targets were

the ionized apo, $^{54}$Fe-bound, and $^{56}$Fe-bound version of each siderophore, with a tolerance of 7.5 ppm.

Putative siderophore candidates were filtered through a series of hard thresholds, such that $MS^1$ spectra were

quality controlled to contain a minimum of 25 datapoints and the maximum intensity of each EIC was greater

than 1e4 counts. Spectra meeting these criteria and containing either $^{54}$Fe-bound and $^{56}$Fe-bound *m/z* peaks

within 30 seconds of each other or an apo peak were displayed for the user to further inspect peak quality

and make the final decision of whether to move on to stage two of processing with a given siderophore

candidate.

Stage two of processing extracted $MS^2$ spectra of the apo and Fe-bound forms of candidate siderophores to

compare with the predicted $MS^2$ generated by *in silico* fragmenter MetFrag (Ruttkies et al., 2016). The *in*

*silico* fragmenter feature was run with a tolerance of 10 ppm on "[M+H]$^+$" and "[M+Na]$^+$" modes. A

confidence level of 1-4, from highest to lowest confidence, was then assigned to putative siderophores based

on the following criteria: (1) peaks were present in $MS^1$ and $MS^2$ spectra, and at least one of the three most-

intense $MS^2$ fragments matched *in silico* fragmentation, (2) peaks were present in $MS^1$ and $MS^2$ spectra, and

smaller-intensity fragments matched *in silico* fragmentation, (3) peaks were present in $MS^1$ and $MS^2$ spectra,

but little to no fragments matched *in silico* fragmentation, and (4) nicely shaped peaks were identified in $MS^1$

spectra but no $MS^2$ spectra was collected (outlined in **Table S4**; example spectra in **Fig. S6-S9**). The

confidence levels were modelled after reporting standards for metabolite identification (Sumner et al., 2007).

MetFrag pulls chemical structures from publicly-available databases like PubChem or COCONUT(Sorokina

et al., 2021), which contain most, but not all variations of siderophores. As such, Fe-bound candidates were

usually run against the apo form available in the database, and for siderophores with similar structures but

variations in fatty chain length or double bond placement, sometimes only one parent structure was available.

A 5-point standard curve with known concentrations of siderophore ferrioxamine E was used for

quantification of putative siderophores, with a limit of detection of 0.257 nM in the eluent (**Fig. S10**), or

0.07-0.21 pM in the sample depending on sample-to-eluent volume ratio at each site (**Table S1**). $MS^1$ peaks

were integrated for all putatively identified siderophores and peak areas were converted to concentration

using the standard curve and the concentration factor of sample volume to eluent volume (**Fig. S10**).

Commercial standards are not available for most siderophores, and different compounds have distinct

ionization efficiencies in ESI-MS. Thus, the siderophore concentrations reported here are estimates of

siderophore concentrations in these environments based on ferrioxamine E. Additionally, 1 mM of

cyanocobalamin was added as an internal standard to each sample aliquot to address any changes in

sensitivity during LC-ESI-MS runs. All putative siderophores that were identified with peak areas less than

the detection limit were discarded, and all remaining putative compounds with at least confidence levels 1

and 2 at one site were included in the manuscript and are referred to as siderophores throughout. Siderophore

identifications remain putative due to inherent uncertainty with assignments by mass, but the confidence

levels were designed such that high confidence candidates contain siderophore-like moieties in their

fragments. Limited sample volumes prevented analysis via LC-ICP-MS like previous studies, which, in

addition to greater availability of commercial standard, would allow definitive identification in future work.

Confidence level 3 and 4 putative siderophores are only included in the Supplementary Information (**Table**

**S5**). In a final step of quality control, EICs for $^{13}$C isotopologues of candidates were inspected to verify

matching peak structure.

**Microbial community analysis**

Microbial community composition was assessed in neutrally buoyant plumes and near venting sites at three

sites: Lucky Strike (Station 7; 1670 m), 10 km S of Rainbow (Station 17; 2000 m), and 200 km E of Rainbow

(Station 11; 600 m, 1600 m and 2250 m). A range of 1- 2 L of seawater were filtered by pressure filtration

through sequential 25 mm membrane filters housed in polypropylene filter holders (Whatman SwinLok, GE

Healthcare, Pittsburgh, Pennsylvania) using a peristaltic pump and silicone tubing. Samples first passed

through a 3 µm pore-size polyester membrane filter (Sterlitech, Auburn, Washington) then onto a 0.2 µm

pore-size polyethersulfone membrane filter (Supor-200, Pall Corporation, Port Washington, New York).

Pump tubing was acid washed with 10% hydrochloric acid and flushed with ultrapure water between each

sample. The filters were flash frozen in liquid nitrogen in 2 mL gasketed bead beating tubes (Fisher Scientific)

at sea.

Nucleic acids (DNA) were extracted as described previously(Santoro et al., 2010), with slight modifications.

Briefly, cells on the filters were lysed directly in the bead beating tubes with sucrose-ethylene diamine

tetraacetic acid (EDTA) lysis buffer (0.75 M sucrose, 20 mM EDTA, 400 mM NaCl, 50 mM Tris) and 1%

sodium dodecyl sulfate. Tubes were then agitated in a bead beating machine (Biospec Products) for 1 min,

and subsequently heated for 2 min. at 99°C in a heat block. Proteinase K (New England Biolabs) was added

to a final concentration of 0.5 mg/mL. Filters were incubated at 55°C for approximately 4 h and the resulting

lysates were purified with the DNeasy kit (Qiagen) using a slightly modified protocol (Santoro et al., 2010).

The purified nucleic acids were eluted in 200 µL of DNase, RNase-free water, and quantified using a

fluorometer (Qubit and Quanti-T HS reagent, Invitrogen Molecular Probes).

The 16S rRNA gene was amplified in all samples using V4 primers (Apprill et al., 2015; Parada et al., 2016)

(515F-Y and 806RB) following a previously established protocol (Stephens et al., 2020). Amplicons were

sequenced using a paired-end 250bp run on an Illumina MiSeq 500 and demultiplexed by the UC Davis

Genome Center. The resulting 16S rRNA amplicon sequences were filtered and trimmed using the DADA2

pipeline in R(Callahan et al., 2016). Taxonomic assignments were made with version 138.1 of the SILVA

SSU    database    (Quast    et    al.,    2013)    (silva_nr99_v138.1_wSpecies_train_set.fa.gz    ;

doi:10.5281/zenodo.4587955; accessed March 2022). Chloroplast and mitochondrial sequences were filtered

out of the dataset using the 'phyloseq' R package (v 1.38.0), after which samples had read depths ranging

from 9375 – 65486 reads (average 28425 $\pm$ 20014 reads) and represented 1010 unique amplicon sequence

variants (ASVs). Read counts were transformed from absolute to relative abundance and taxa were

aggregated to the Family level. The ten most abundant families present in each sample were visualized using
the 'ggplot2' package (v. 3.3.5).

In order to assess the potential of the observed prokaryotic taxa to produce siderophores, we downloaded all
siderophore biosynthetic gene clusters (BGCs) in the antismash secondary metabolite database ($n = 7909$)
and used text-string matching to compare genera containing these BGCs to the genera found in our 16S rRNA
gene dataset(Blin et al., 2021). We cross-referenced the nomenclature of antismash-predicted siderophores
with that of the siderophores identified by LC-ESI-MS in this study, accounting for minor differences in
naming convention between the two databases, to determine if microbial community members present at
each site were predicted to make any of the siderophores that were measured at that site. Station 38 and
Station 12 were the closest sites with siderophore measurements for comparison against the taxonomic
samples taken at 200 km E of Rainbow and 10 km S of Rainbow, respectively. Samples for microbial
taxonomy and siderophore identity were taken from the same location at Lucky Strike and thus directly
compared.

**Data Availability**
The CSV data reported in this study has been deposited at Zenodo under the DOI:
http://doi.org/10.5281/zenodo.7325154. The LC-ES-MS data has been deposited on Massive under the DOI:
http://doi.org/doi.10.25345/C5V97ZW7N. Microbial 16S rRNA data have been deposited on GenBank under
the accession number BioProject #PRJNA865382. All data is freely available on each of these data
repositories.

**Acknowledgments**
We acknowledge the captain and crew of the R/V *James Cook*, Chief Scientist Alessandro Tagliabue, and
Noah Gluschankoff for supporting this work. This study was a part of the FeRidge project (GEOTRACES
section GA13) which was supported by the Natural Environment Research Council funding (NERC United
Kingdom Grants NE/N010396/1 to MCL and NE/N009525/1 to AT). The International GEOTRACES
Programme is possible in part thanks to the support from the U.S. National Science Foundation (Grant OCE-
1840868) to the Scientific Committee on Oceanic Research (SCOR). CLH was funded by JISAO/CICOES
postdoctoral fellowship. PJM was funded through the NOAA Hollings Scholar summer program. JR was
funded by NOAA Ocean Exploration and Research, NOAA Earth-Ocean Interactions programs at NOAA-
Pacific Marine Environmental Labs, and JISAO/CICOES. Part of this work was carried out in the University
of Washington TraceLab, which receives support from the M.J. Murdock Charitable Trust in conjunction
with the University of Washington College of Environment, and the Pacific Marine Environmental Labs at
the National Oceanic and Atmospheric Administration. Parts of this work was also carried out in Dr. Anitra

Ingalls laboratory with the help of Laura Truxal and Dr. Jiwoon Park at the University of Washington-School
of Oceanography.

**Author Contributions:** Manuscript preparation, sample/data processing, CSV analysis, and interpretation
(C.L.H.), manuscript preparation, LC-ESI-MS data analysis and interpretation (P.J.M.), microbial analysis
and interpretation (J.B.A. and A.E.S.), dissolved iron and derived excess $^3\text{He}_{xs}$ measurements, sample
collection (A.J.M. L. and M.C.L.), microbial data collection and ligand data interpretation (T.M. and K.N.B.),
and project design and planning, data interpretation, and mentoring (A.T., M.C.L., J.A.R., and R.M.B.). All
authors were involved in editing manuscript.

**Competing Interest Statement:** The authors declare no competing interests.

## References

Abualhaija, M. M. and van den Berg, C. M. G. G.: Chemical speciation of iron in seawater using catalytic
cathodic stripping voltammetry with ligand competition against salicylaldoxime, Mar. Chem., 164, 60–74,
https://doi.org/10.1016/j.marchem.2014.06.005, 2014.

Apprill, A., Mcnally, S., Parsons, R., and Weber, L.: Minor revision to V4 region SSU rRNA 806R gene
primer greatly increases detection of SAR11 bacterioplankton, Aquat. Microb. Ecol., 75, 129–137,
https://doi.org/10.3354/ame01753, 2015.

Bazylev, B. A.: Allochemical Metamorphism of Mantle Peridoties in the Hayes Fracture Zone of the North
Atlantic, Petrology, 5, 362–379, 1997.

Beaulieu, S. E. and Szafrański, K. M.: InterRidge Global Database of Active Submarine Hydrothermal Vent
Fields Version 3.4, https://doi.org/10.1594/PANGAEA.917894, 2020.

Bennett, S. a., Achterberg, E. P., Connelly, D. P., Statham, P. J., Fones, G. R., and German, C. R.: The
distribution and stabilisation of dissolved Fe in deep-sea hydrothermal plumes, Earth Planet. Sci. Lett., 270,
157–167, https://doi.org/10.1016/j.epsl.2008.01.048, 2008.

Bennett, S. a., Hansman, R. L., Sessions, A. L., Nakamura, K. ichi, and Edwards, K. J.: Tracing iron-fueled
microbial carbon production within the hydrothermal plume at the Loihi seamount, Geochim. Cosmochim.
Acta, 75, 5526–5539, https://doi.org/10.1016/j.gca.2011.06.039, 2011.

Blin, K., Shaw, S., Kautsar, S. A., Medema, M. H., and Weber, T.: The antiSMASH database version 3:
Increased taxonomic coverage and new query features for modular enzymes, Nucleic Acids Res., 49, D639–
D643, https://doi.org/10.1093/nar/gkaa978, 2021.

Boiteau, R. M., Mende, D. R., Hawco, N. J., McIlvin, M. R., Fitzsimmons, J. N., Saito, M. A., Sedwick, P.
N., DeLong, E. F., and Repeta, D. J.: Siderophore-based microbial adaptations to iron scarcity across the
eastern Pacific Ocean, Proc. Natl. Acad. Sci., 113, 14237–14242, https://doi.org/10.1073/pnas.1608594113,
2016.

Boiteau, R. M., Till, C. P., Coale, T. H., Fitzsimmons, J. N., Bruland, K. W., and Repeta, D. J.: Patterns of
iron and siderophore distributions across the California Current System, Limnol. Oceanogr., 64, 376–389,
https://doi.org/10.1002/lno.11046, 2019.

Buck, K. N., Lohan, M. C., Berger, C. J. M., and Bruland, K. W.: Dissolved iron speciation in two distinct
river plumes and an estuary: Implications for riverine iron supply, Limnol. Oceanogr., 52, 843–855,
https://doi.org/10.4319/lo.2007.52.2.0843, 2007.

Buck, K. N., Sohst, B., and Sedwick, P. N.: The organic complexation of dissolved iron along the U.S.
GEOTRACES (GA03) North Atlantic Section, Deep. Res. Part II Top. Stud. Oceanogr., 116, 152–165,
https://doi.org/10.1016/j.dsr2.2014.11.016, 2015.

Buck, K. N., Sedwick, P. N., Sohst, B., and Carlson, C. A.: Organic complexation of iron in the eastern
tropical South Pacific: Results from US GEOTRACES Eastern Pacific Zonal Transect (GEOTRACES cruise
GP16), Mar. Chem., 201, 229–241, https://doi.org/10.1016/j.marchem.2017.11.007, 2018.

Bundy, R. M., Abdulla, H. A. N., Hatcher, P. G., Biller, D. V., Buck, K. N., and Barbeau, K. A.: Iron-binding
ligands and humic substances in the San Francisco Bay estuary and estuarine-influenced shelf regions of
coastal California, Mar. Chem., 173, 183–194, https://doi.org/10.1016/j.marchem.2014.11.005, 2015.

Bundy, R. M., Boiteau, R. M., McLean, C., Turk-Kubo, K. A., McIlvin, M. R., Saito, M. A., Mooy, B. A.
Van, and Repeta, D. J.: Distinct Siderophores Contribute to Iron Cycling in the Mesopelagic at Station
ALOHA, Front. Mar. Sci., 1–15, https://doi.org/10.3389/fmars.2018.00061, 2018.

Butler, A.: Marine siderophores and microbial iron mobilization., Biometals, 18, 369–374,
https://doi.org/10.1007/s10534-005-3711-0, 2005.

Butler, A. and Theisen, R. M.: Iron(III)-siderophore coordination chemistry: Reactivity of marine
siderophores., Coord. Chem. Rev., 254, 288–296, https://doi.org/10.1016/j.ccr.2009.09.010, 2010.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P.: DADA2:
High-resolution sample inference from Illumina amplicon data, Nat. Methods, 13, 581–583,
https://doi.org/10.1038/nmeth.3869, 2016.

Carmichael, J. R., Zhou, H., and Butler, A.: A suite of asymmetric citrate siderophores isolated from a marine
Shewanella species, J. Inorg. Biochem., 198, 1–6, https://doi.org/10.1016/j.jinorgbio.2019.110736, 2019.

Cowen, J. P. and Bruland, K. W.: Metal deposits associated with bacteria: implications for Fe and Mn marine
biogeochemistry, Deep Sea Res. Part A. Oceanogr. Res. Pap., 32, 253–272, https://doi.org/10.1016/0198-
0149(85)90078-0, 1985.

Cowen, J. P., Massoth, G. J., and Feely, R. A.: Scavenging rates of dissolved manganese in a hydrothermal
vent plume, Deep Sea Res. Part A. Oceanogr. Res. Pap., 37, 1619–1637, https://doi.org/10.1016/0198-
0149(90)90065-4, 1990.

Crowley, D. E., Wang, Y. C., Reid, C. P. P., and Szaniszlo, P. J.: Mechanisms of iron acquisition from
siderophores by microorganisms and plants, Plant Soil, 130, 179–198, 1991.

Fishwick, M. P., Sedwick, P. N., Lohan, M. C., Worsfold, P. J., Buck, K. N., Church, T. M., and Ussher, S.
J.: The impact of changing surface ocean conditions on the dissolution of aerosol iron, Global Biogeochem.
Cycles, 28, 1235–1250, https://doi.org/10.1002/2014GB004921, 2014.

Fitzsimmons, J. N., John, S. G., Marsay, C. M., Hoffman, C. L., Nicholas, S. L., Toner, B. M., German, C.
R., and Sherrell, R. M.: Iron persistence in the distal hydrothermal plume supported by dissolved – particulate
exchange, Nat. Geosci., 10, 1–8, https://doi.org/10.1038/ngeo2900, 2017.

Gu, H., Sun, Q., Luo, J., Zhang, J., and Sun, L.: A First Study of the Virulence Potential of a Bacillus subtilis
Isolate From Deep-Sea Hydrothermal Vent, Front. Cell. Infect. Microbiol., 9, 1–14,
https://doi.org/10.3389/fcimb.2019.00183, 2019.

Hawkes, J. A., Gledhill, M., Connelly, D. P., and Achterberg, E. P.: Characterisation of iron binding ligands
in seawater by reverse titration, Anal. Chim. Acta, 766, 53–60, https://doi.org/10.1016/j.aca.2012.12.048,

2013a.

Hawkes, J. A., Connelly, D. P., Gledhill, M., and Achterberg, E. P.: The stabilisation and transportation of
dissolved iron from high temperature hydrothermal vent systems, Earth Planet. Sci. Lett., 375, 280–290,
https://doi.org/10.1016/j.epsl.2013.05.047, 2013b.

Hider, R. C. and Kong, X.: Chemistry and biology of siderophores, Nat. Prod. Rep., 27, 637–657,
https://doi.org/10.1039/b906679a, 2010.

Hoffman, C. L., Nicholas, S. L., Ohnemus, D. C., Fitzsimmons, J. N., Sherrell, R. M., German, C. R., Heller,
M. I., Lee, J. mi, Lam, P. J., and Toner, B. M.: Near-field iron and carbon chemistry of non-buoyant
hydrothermal plume particles, Southern East Pacific Rise 15°S, Mar. Chem., 201, 183–197,
https://doi.org/10.1016/j.marchem.2018.01.011, 2018.

Hoffman, C. L., Schladweiler, C., Seaton, N. C. A., Nicholas, S. L., Fitzsimmons, J., Sherrell, R. M., German,
C. R., Lam, P., and Toner, B. M.: Diagnostic morphology and solid-state chemical speciation of
hydrothermally derived particulate Fe in a long-range dispersing plume, ACS Earth Sp. Chem., 4, 1831–
1842, https://doi.org/10.1021/acsearthspacechem.0c00067, 2020.

Homann, V. V, Sandy, M., Tincu, J. A., Templeton, A. S., Tebo, B. M., and Butler, A.: Loihichelins A - F ,
a Suite of Amphiphilic Siderophores Produced by the Marine Bacterium Halomonas LOB-5, J. Nat. Prod.,
72, 884–888, 2009.

Kato, C. and Nogi, Y.: Correlation between phylogenetic structure and function : examples from deep-sea
Shewanella, 35, 223–230, 2001.

Kelley, D. S. and Shank, T. M.: Hydrothermal systems: A decade of discovery in slow spreading
environments, Geophys. Monogr. Ser., 188, 369–407, https://doi.org/10.1029/2010GM000945, 2010.

Kulmer, W. and Ingalls, A. E.: The R Journal: Tidy Data Neatly Resolves Mass-Spectrometry's Ragged
Arrays, R J., 2022.

Lauderdale, J. M., Braakman, R., Forget, G., Dutkiewicz, S., and Follows, M. J.: Microbial feedbacks
optimize ocean iron availability, Proc. Natl. Acad. Sci. U. S. A., 117, 4842–4849,
https://doi.org/10.1073/pnas.1917277117, 2020.

Li, M., Toner, B. M., Baker, B. J., Breier, J. a, Sheik, C. S., and Dick, G. J.: Microbial iron uptake as a
mechanism for dispersing iron from deep-sea hydrothermal vents., Nat. Commun., 5, 3192,
https://doi.org/10.1038/ncomms4192, 2014.

Lough, A. J. M., Tagliabue, A., Demasy, C., Resing, J. A., Mellett, T., Wyatt, N. J., and Lohan, M. C.: The
impact of hydrothermal vent geochemistry on the addition of iron to the deep ocean, Biogeosciences Discuss.,
[preprint], 1–23, https://doi.org/10.5194/bg-2022-73, 2022.

Manck, L. E., Park, J., Tully, B. J., Poire, A. M., Bundy, R. M., Dupont, C. L., and Barbeau, K. A.:
Petrobactin, a siderophore produced by Alteromonas, mediates community iron acquisition in the global
ocean, ISME J., 16, 358–369, https://doi.org/10.1038/s41396-021-01065-y, 2022.

Martinez, J. S., Carter-Franklin, J. N., Mann, E. L., Martin, J. D., Haygood, M. G., and Butler, A.: Structure
and membrane affinity of a suite of amphiphilic siderophores produced by a marine bacterium, Proc. Natl.
Acad. Sci. U. S. A., 100, 3754–3759, https://doi.org/10.1073/pnas.0637444100, 2003.

Mellett, T., Albers, J. B., Santoro, A., Wang, W., Salaun, P., Resing, J., Lough, A. J. ., Tagliabue, A., Lohan,
M., Bundy, R. M., and Buck, K. N.: Particle exchange mediated by organic ligands in incubation experiments
of hydrothermal vent plumes along the mid-Atlantic Ridge, n.d.

Moore, L. E., Heller, M. I., Barbeau, K. A., Moffett, J. W., and Bundy, R. M.: Organic complexation of iron
by strong ligands and siderophores in the eastern tropical North Pacific oxygen deficient zone, Mar. Chem.,

236, 104021, https://doi.org/10.1016/j.marchem.2021.104021, 2021.

Omanović, D., Garnier, C., and Pižeta, I.: ProMCC: An all-in-one tool for trace metal complexation studies, Mar. Chem., 173, 25–39, https://doi.org/10.1016/j.marchem.2014.10.011, 2015.

Parada, A. E., Needham, D. M., and Fuhrman, J. A.: Every base matters: Assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples, Environ. Microbiol., 18, 1403–1414, https://doi.org/10.1111/1462-2920.13023, 2016.

Park, J., Durham, B. P., Key, R. S., Groussman, R. D., Pinedo-Gonzalez, P., Hawco, N. J., John, S. G., Carlson, M. C. G., Lindell, D., Juranek, L., Ferrón, S., Ribalet, F., Armbrust, E. V., Ingalls, A. E., and Bundy, R. M.: Siderophore production and utilization by microbes in the North Pacific Ocean, bioRxiv, 2022.02.26.482025, https://doi.org/10.1101/2022.02.26.482025, 2022.

Park, J., Durham, B. P., Key, R. S., Groussman, R. D., Pinedo-Gonzalez, P., Hawco, N. J., John, S. G., Carlson, M. C. G., Lindell, D., Juranek, L., Ferrón, S., Ribalet, F., Armbrust, E. V., Ingalls, A. E., and Bundy, R. M.: Siderophore production and utilization by microbes in the North Pacific Ocean, Limnol. Oceanogr., 2022.02.26.482025, https://doi.org/10.1002/lno.12373, 2023.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.: The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools, Nucleic Acids Res., 41, 590–596, https://doi.org/10.1093/nar/gks1219, 2013.

Resing, J. a., Sedwick, P. N., German, C. R., Jenkins, W. J., Moffett, J. W., Sohst, B. M., and Tagliabue, A.: Basin-scale transport of hydrothermal dissolved metals across the South Pacific Ocean, Nature, 523, 200–203, https://doi.org/10.1038/nature14577, 2015.

Ruttkies, C., Schymanski, E. L., Wolf, S., Hollender, J., and Neumann, S.: MetFrag relaunched: incorporating strategies beyond in silico fragmentation, J. Cheminform., 8, 1–16, https://doi.org/10.1186/s13321-016-0115-9, 2016.

Sander, S. G. and Koschinsky, A.: Metal flux from hydrothermal vents increased by organic complexation, Nat. Geosci., 4, 145–150, https://doi.org/10.1038/ngeo1088, 2011.

Sandy, M. and Butler, A.: Microbial iron acquisition: marine and terrestrial siderophores., Chem. Rev., 109, 4580–95, https://doi.org/10.1021/cr9002787, 2009.

Santoro, A. E., Casciotti, K. L., and Francis, C. A.: Activity, abundance and diversity of nitrifying archaea and bacteria in the central California Current, Environ. Microbiol., 12, 1989–2006, https://doi.org/10.1111/j.1462-2920.2010.02205.x, 2010.

Sorokina, M., Merseburger, P., Rajan, K., Yirik, M. A., and Steinbeck, C.: COCONUT online: Collection of Open Natural Products database, J. Cheminform., 13, 1–13, https://doi.org/10.1186/s13321-020-00478-9, 2021.

Stephens, B. M., Opalk, K., Petras, D., Liu, S., Comstock, J., Aluwihare, L. I., Hansell, D. A., and Carlson, C. A.: Organic Matter Composition at Ocean Station Papa Affects Its Bioavailability, Bacterioplankton Growth Efficiency and the Responding Taxa, Front. Mar. Sci., 7, https://doi.org/10.3389/fmars.2020.590273, 2020.

Sumner, L. W., Amberg, A., Barrett, D., Beale, M. H., Beger, R., Daykin, C. A., Fan, T. W.-M., Fiehn, O., Goodacre, R., Griffin, J. L., Hankemeier, T., Hardy, N., Harnly, J., Higashi, R., Kopka, J., Lane, A. N., Lindon, J. C., Marriott, P., Nicholls, A. W., Reily, M. D., Thaden, J. J., and Viant, M. R.: Proposed minimum reporting standards for chemical analysis, Metabolomics, 3, 211–221, https://doi.org/10.1007/s11306-007-0082-2, 2007.

Tagliabue, A., Bowie, A. R., Boyd, P. W., Buck, K. N., Johnson, K. S., and Saito, M. A.: The integral role of iron in ocean biogeochemistry, Nature, 543, 51–59, https://doi.org/10.1038/nature21058, 2017.

Toner, B. M., Fakra, S. C., Manganini, S. J., Santelli, C. M., Marcus, M. a., Moffett, J. W., Rouxel, O.,
German, C. R., and Edwards, K. J.: Preservation of iron(II) by carbon-rich matrices in a hydrothermal plume,
Nat. Geosci., 2, 197–201, https://doi.org/10.1038/ngeo433, 2009.

Tortell, P. D., Maldonado, M. T., and Price, N. M.: The role of heterotrophic bacteria in iron-limited ocean
ecosystems, Nature, 383, 330–332, https://doi.org/10.1038/383330a0, 1996.

Vraspir, J. M. and Butler, A.: Chemistry of marine ligands and siderophores., Ann. Rev. Mar. Sci., 1, 43–63,
https://doi.org/10.1146/annurev.marine.010908.163712, 2009.

Waska, H., Koschinsky, A., Ruiz Chancho, M. J., and Dittmar, T.: Investigating the potential of solid-phase
extraction and Fourier-transform ion cyclotron resonance mass spectrometry (FT-ICR-MS) for the isolation
and identification of dissolved metal-organic complexes from natural waters, Mar. Chem., 173, 78–92,
https://doi.org/10.1016/j.marchem.2014.10.001, 2015.

    **Figures and Tables**

    **Figure 1. Dissolved iron is strongly correlated with L$_1$ iron-binding ligands in diverse hydrothermal**

    **systems.** (a) Station map showing the 11 sites investigated along the MAR. Known hydrothermal vents are

    marked as red triangles(Beaulieu and Szafrański, 2020). Two expanded inset maps for (b) Rainbow and (c)

    TAG hydrothermal vent fields. For additional information about vent site characteristics refer to **Table 1**. (d)

    dFe versus L$_1$ iron-binding ligands at each vent site in this study showing a ~1:1 correlation (m= 0.88, R$^2$=

    0.96) with dFe in neutrally-buoyant plumes along the MAR. (e) dFe versus L$_1$ ligands from previous studies

    over the ridge axis and ~80 km from ridge axis in the Southern East Pacific Rise hydrothermal plume(Buck

    et al., 2018), and over TAG hydrothermal vent field(Buck et al., 2015). The solid black lines in (d) and (e)

    are the 1:1 ratio line between dFe and ligand concentrations, and dashed lines show the linear regression for

    the corresponding data. Square symbols refer to spreading centers, triangles refer to fracture zones, and

591    circles refer to alkaline vents. Error bars represent the 95% confidence interval of the data fit as calculated

592    by ProMCC(Omanović et al., 2015). The map was created using GeoMapApp version 3.6.14.

**Figure 2**. **Siderophore presence in hydrothermal plumes along the MAR.** (a) Heat map of confidence
levels 1-2 (blue gradient, 1 = highest confidence). Gray boxes indicate a detection with lower confidence (see
Methods), and white boxes indicate no detection at those sites. The y-axis is ordered from top to bottom in
terms of descending mass of the apo (without Fe) form of the siderophore. (b) Model II ordinary least squares
regression on dFe versus summed siderophore concentrations (of detections in Fig. 2b), calculated from peak
areas, at each site. Since the siderophore analysis was performed on pooled samples, the dFe values in the
regression are weighted values based on measured dFe and volume of each constituent of the pooled sample.
The vertical error bars represent the standard deviation of dFe of the constituents. TAG (St. 35) was not
included in the regression due to its large range of dFe values and outlier behavior. (c-d) Fe bound versus
total summed concentration of (c) all types of siderophores and (d) amphiphilic siderophores at each station.
The vertical green lines separate fracture/diffuse sites from off-axis sites and vertical red lines separate off-
axis from on-axis sites as defined in Table 1. Symbols follow Fig. 1.

**Figure 3. Relative abundance of putative siderophore-producing taxa.** Bar height indicates the proportion

of 16S rRNA genes recovered in each sample, separated by depth, filter size fraction, and site location. Colors

correspond to taxonomy. Genera found in MAR vent microbial communities with members in the antismash

database predicted to produce siderophores are depicted at the family level.

**Table 1. Characteristics of sample locations along the Mid Atlantic Ridge.**

| Vent Names | Abbr. | Station | Geology | Host rock | Vent type | Spreading rate (mm/yr) | Summed putative siderophore concentration (pM) | Summed Siderophore concentration/ $L_1$ ligand (%)* |
|---|---|---|---|---|---|---|---|---|
| Lucky Strike | LS | 7/8 | Spreading Center | gabbro | Black smoker | 20.2 | 4.38 | 0.034-0.19 |
| 33 km E of Rainbow | CER | 12 | Spreading Center | - | - | - | 0.537 | 0.013-0.017 |
| Rainbow | R | 38 | Spreading Center | ultramafic | Black smoker | 20.6 | 12.1 | *n.a.* |
| Hayes Fracture Zone | HFZ | 21 | Fracture Zone | peridotites/gabbro | - | 21.2 | 2.74 | 0.20-0.39 |
| Lost City | LC | 23 | Fracture Zone | ultramafic/gabbro | Alkaline | 22.6 | 3.47 | 0.27-0.35 |
| Broken Spur | BS | 24 | Spreading Center | gabbro | Black smoker/diffuse | 22.9 | 6.30 | 0.07-0.29 |
| 29 km N of TAG | CNT | 26 | Spreading Center | - | - | - | 0.968 | 0.014-0.079 |
| 30 km W of TAG | CWT | 30 | Spreading Center | - | - | - | 2.91 | 0.15 |
| 30 km E of TAG | CET | 31 | Spreading Center | - | - | - | 2.19 | 0.31 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Trans-Atlantic Geotraverse | TAG | 35 | Spreading Center | gabbro | Black smoker | 23.6 | 6.61 | 0.18 |
| Low Temp Slope | LTS | 37 | - | - | Diffuse fluids | - | 1.13 | 0.079-0.087 |

Spreading rates along the Mid-Atlantic Ridge were gathered from the Interridge Database v3.4. Host rock groups were determined from previously discussed classifications(Bazylev, 1997; Kelley and Shank, 2010). Off-axis sites –33 km E of Rainbow, 29 km N of TAG, 30 km E of TAG, and 30 km W of TAG– were far-field locations of their respective vent field. Low Temp Slope was a diffuse-dominated site that was sampled for the first time as a part of this study. Summed putative siderophore concentrations and the percent of $L_1$ ligand are reported for compounds detected with at least confidence level 1 and 2 at one site. These values do not take into account typical extraction efficiencies of ENV columns for Fe-binding organics. Average $L_1$ ligand and siderophore concentrations can be viewed in **Table S3** and concentrations for individual siderophores can be observed in **Table S5**.

*The siderophore sample at each site was pooled from ligand samples, so the percentage of siderophores in the $L_1$ pool is presented as a range based on the range of $L_1$ concentrations at each site.

*n.a.*= unable to be determined

- = unknown