Microbial strong organic ligand production is tightly coupled to iron in hydrothermal plumes

Colleen L. Hoffman[1,2,3*] and Patrick J. Monreal[3,4*], Justine B. Albers[5], Alastair J.M. Lough[6], Alyson E. Santoro[5], Travis Mellett[3,7], Kristen N. Buck[7,8], Alessandro Tagliabue[9], Maeve C. Lohan[6], Joseph A. Resing[1,2,3], Randelle M. Bundy[3]

[1]Joint Institute for the Study of Atmosphere and Ocean, University of Washington, 3737 Brooklyn
       Avenue NE, Seattle, WA 98195, USA

[2]Cooperative Institute for Climate, Ocean, and Ecosystem Studies, University of Washington,
       3737 Brooklyn Avenue NE, Seattle, WA 98195, USA

[3]School of Oceanography, University of Washington, 1501 NE Boat Street, Seattle, WA 98195,
       USA

[4]Earth Systems Program, Stanford University, 473 Via Ortega, Stanford, CA 94305,
       USA

[5]Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara,
       CA 93106, USA

[6]Department of Ocean and Earth Sciences, National Oceanography Centre, University of
       Southampton, European Way, Southampton SO14 3ZH, United Kingdom

[7]College of Marine Science, University of South Florida, 140 7th Avenue South, St. Petersburg,
       FL, 33701, USA

[8]College of Earth, Oceans, and Atmospheric Sciences, Oregon State University, 2651 SW Orchard Ave,
       Corvallis, OR, 97331, USA

[9]Department of Earth, Ocean, and Ecological Sciences, University of Liverpool, 4 Brownlow
       Street, Liverpool l69 3GP, United Kingdom

*Colleen L. Hoffman and Patrick J. Monreal

Email: clhoffma@gmail.com, pmonreal@uw.edu

Abstract. The oceanic iron inventory exerts a key control on upper ocean primary productivity and thus is intricately linked to the carbon cycle. Hydrothermal vents have emerged as an important source of iron to seawater, yet only a subset of this iron is soluble and persists long enough to be available for surface biological uptake. The longevity and solubility of iron in seawater is governed by organic iron-binding ligands, and strong organic ligands called siderophores are produced by marine microorganisms to access and retain iron in soluble forms. Siderophores and other microbially-produced ligands comprise part of the ocean's dissolved iron-binding ligand pool, which is hypothesized to aid in the persistence of soluble iron in hydrothermal environments. To explore this hypothesis, we measured iron, iron-binding ligands, and siderophores from 11 geochemically distinct sites along a 1,700 km section of the Mid-Atlantic Ridge. For the first time, we identified siderophores in hydrothermal plumes at all sites. Proximity to the vent played an important role in dictating siderophore types and diversity. The notable presence of amphiphilic siderophores may point to microbial utilization of siderophores to access particulate iron in hydrothermal plumes, and the exchange of iron between dissolved and particulate phases in these systems. A tight coupling between strong ligands and dissolved iron was observed in the neutrally buoyant plume across six distinct hydrothermal environments, and the presence of dissolved siderophores with siderophore-producing microbial genera suggests that biological production of siderophores exerts a key control on hydrothermal dissolved iron concentrations.

1 Introduction

Over the last few decades, observations and modelling efforts have increased our understanding about the critical role organic ligands play in the cycling, transport, and utilization of trace metals (Tagliabue et al., 2017; Buck et al., 2018; Bundy et al., 2018; Moore et al., 2021). Strong iron-binding organic ligands (L_1) are a heterogeneous mixture of microbially produced compounds that are operationally classified based on their binding strength ($\log K_{Fe',FeL}^{cond} > 12$) with iron (Fe), and are thermodynamically favored to complex and stabilize external sources of Fe to prevent its scavenging and removal. For example, in high dissolved and particulate Fe estuarine systems, only the dissolved Fe (dFe) bound to the strongest Fe-binding ligands escapes estuarine flocculation (Bundy et al., 2015; Buck et al., 2007) and remains accessible for downstream biological uptake.

Siderophores represent the strongest Fe-binding organic ligands. They are produced by bacteria and fungi to facilitate Fe uptake and to draw Fe out of otherwise inaccessible phases in the marine environment (Butler, 2005; Manck et al., 2022). Historically, due to the energy required to produce a siderophore and the potential for the diffusive loss of the siderophore after excretion, they have primarily been considered as an important microbial strategy for Fe acquisition in the low-Fe surface ocean (Vraspir and Butler, 2009; Butler, 2005). However, siderophores have been identified in oxygen-deficient zones ([dFe] = >2nM) (Moore et al., 2021), and genetic evidence in the Guaymas Basin hydrothermal system showed that >70% of Fe-related bacterial

transcripts were involved in siderophore uptake, regulation, and biosynthesis (Li et al., 2014). Additionally, siderophore production and uptake is also a common Fe acquisition strategy within terrestrial soil and pathogenic ecosystems (Sandy and Butler, 2009) where Fe is orders of magnitude higher than seawater. This suggests siderophores are important for Fe acquisition and offshore transport of Fe away from sediment sources, and provides evidence that siderophore production and uptake may be important even in the higher Fe environments of the deep ocean, similar to terrestrial and pathogenic ecosystems. A recent modeling study explored this idea and showed that organic ligand production could be a widespread microbial Fe acquisition strategy, which then tightly controls the utilization and availability of Fe over longer timescales (Lauderdale et al., 2020).

Although strong ligands have been observed throughout the deep ocean and siderophores have been found below the euphotic zone (Bundy et al., 2018), no previous studies have ever directly characterized siderophores in hydrothermal systems. Strong Fe-binding ligands have been measured in hydrothermal vent systems, but the mechanisms for their production or the identity of these strong ligands in vents has never been explored. Some type of 'stabilizing agent' or ligand is critical for the long-range transport of hydrothermal Fe that has been observed in the ocean interior, and the role of strong Fe-binding ligands and siderophores in hydrothermal dFe transport represents an important gap in our understanding of how hydrothermal sources may regulate the deep ocean dFe inventory (13). Here, for the first time, we identified the presence of siderophores and siderophore-producing microbes in 11 geochemically distinct hydrothermal environments along the slow-spreading (20-50 mm/yr) Mid-Atlantic Ridge (MAR), including four black smokers (high temperature, high Fe), four off-axis sites (within 30 km of black smokers), one diffuse-flow vent (low temperature, low Fe), one alkaline vent (pH 9-11, very low Fe), and one non-vent fracture zone, to understand whether microbial ligand production impacts the supply of hydrothermal dFe to the ocean. Overall, our results show microbially-produced siderophores were present in all sites, and that strong L_1 ligands, which include siderophores, exert a tight control on hydrothermal dFe. This suggests that discrete strong organic ligands produced by bacteria play a key role in setting deep ocean Fe concentrations globally across hydrothermal systems.

2 Results and Discussion

2.1 The role of L_1 iron-binding ligands and siderophores in hydrothermal plumes

Strong organic Fe-binding ligands, or L_1 ligands, have been implicated as important for stabilizing and transporting Fe in hydrothermal plumes (Tagliabue et al., 2017; Resing et al., 2015; Buck et al., 2018). The average binding strength (log $K_{Fe',FeL}^{cond}$) and concentration of organic Fe-binding ligands were quantified in multiple vent systems that spanned a wide range in dFe concentrations (0.41-90.3 nM) and underlying vent geology. Over 99% of the dFe in the neutrally buoyant plumes at these sites was complexed by L_1 ligands and the strong ligands were almost always completely saturated with dFe, meaning Fe-free 'excess' L_1

ligands capable of binding additional Fe were generally present in low concentrations (< 1 nM; **Fig. S1**). As a result, dFe concentrations were tightly coupled to L_1 ligands in a nearly 1:1 ratio (dFe: L_1, **Fig. 1d**) similar to previous studies in other neutrally buoyant plumes (**Fig. 1e**) (Lough et al., 2022; Buck et al., 2018, 2015). The strong coupling between dFe and ligands was only observed at sites where L_1 ligands were detected. Some sampling locations contained high concentrations of weaker ligands (log $K_{Fe',FeL}^{cond} < 12$, **Table S2**) with no correlation to dFe. The weak ligands were either not completely saturated with dFe, or in some cases weak ligands were found in samples with extremely high concentrations of dFe (> 20 nM) in the rising buoyant plume. Although other studies have observed a correlation with dFe and ligands in individual plumes, this is the first study to systematically observe the tight coupling between dFe and L_1 in multiple distinct vent sites in the same study.

Altogether, our results suggest that L_1 ligands uniquely set the dFe concentration in neutrally buoyant plumes. A similar direct control of dFe concentrations by L_1 ligands has been previously observed in rivers (Buck et al., 2007) and aerosol solubility experiments (Fishwick et al., 2014). A possible explanation is that both the dFe and L_1 ligands originate from the vents themselves, yielding a tightly coupled hydrothermal endmember. However, the concentration of L_1 ligands did not correlate with excess mantle Helium-3 ($^3He_{xs}$, **Fig S2**, **Table S2**) (Lough et al., 2022), a nearly conservative tracer of the mixing of hydrothermal fluids with seawater (Buck et al., 2018). These results suggest that L_1 ligands along the MAR are likely not sourced from the vent fluids along with dFe. All known sources of L_1 ligands are biologically produced, so the L_1 ligands observed here could be sourced from either microbes that produced them in the surrounding deep ocean seawater that was then entrained, local production from vent-biota and/or microbial mats, diffusion from microbial production in sediments, or *in-situ* production by microbes within the neutrally buoyant plume (Mellett et al., n.d.).

Microbial organic ligand production has been suggested to tightly control the global oceanic trace metal inventories in both hydrothermal plumes (Cowen and Bruland, 1985; Cowen et al., 1990) and the open ocean (Lauderdale et al., 2020). Specifically, siderophores have Fe-binding constants that would operationally define them as L_1 ligands (Moore et al., 2021; Manck et al., 2022), and have been proposed to be important in hydrothermal plumes (Li et al., 2014), though they have never been directly measured. To further explore this tight coupling between Fe and microbially-mediated ligand production within hydrothermal plumes, we used a state-of-the-art liquid chromatography coupled to electrospray ionization mass spectroscopy (Boiteau et al., 2016) method to identify discrete components of the L_1 ligands we observed. We screened our samples against a library of over 500 known siderophores that have been characterized from bacteria and fungi across marine and terrestrial environments. In a targeted approach, we used mass-to-charge ratio (*m/z*), MS/MS spectra, and accompanying chromatographic characteristics to putatively identify microbially produced siderophores with high confidence in every vent site and observed a high diversity of compounds (**Fig. 2a**). Relative peak areas as a proxy for concentrations of putative siderophores also significantly correlated with

dFe as observed with dFe and L_1 ligands (**Fig. 2b**). The siderophores were present in picomolar concentrations, similar to other upper ocean marine environments (Boiteau et al., 2016; Moore et al., 2021; Park et al., 2022; Bundy et al., 2018).

The siderophores we identified with high confidence comprised from 0.01-0.4% of the L_1 ligands that were measured at the same sites (**Table 1**). Our estimate is likely an underestimate of siderophore contributions to the L_1 ligand pool due to several possibilities. First, the majority of siderophores are not commercially available to use as standards. Secondly, each individual siderophore may have a different ionization or extraction efficiency during our analytical process, and, thirdly, we are only reporting the concentration of siderophores we have identified with high confidence (i.e. accurate mass and fragments matching siderophore-like moieties; 68% of the total identifications, **Fig 2a**, **S3**). Finally, known siderophores only represent a small fraction of the total siderophores expected to be produced in nature (Hider and Kong, 2010). While we quantified our siderophore concentrations based on a standard curve with a commercially available siderophore, previous work has also shown the extraction efficiency in our solid phase extraction technique is around 10% for bulk Fe-binding organics (Bundy et al., 2018) and 40% for a siderophore standard (Waska et al., 2015). With this in mind, correcting for our extraction efficiencies put the concentrations of identified siderophores closer to 1-4% of the L_1 pool. However, we are inevitably missing other naturally occurring unknown compounds. Regardless of the small percentage contribution to total L_1 ligands quantified here, it is evident that microbially produced siderophores are present and widespread across all vent sites.

The presence of a high diversity of siderophores across a huge range of hydrothermal vent systems reveals several surprising aspects of Fe cycling in hydrothermal plumes. The biosynthesis of a siderophore is energy-intensive and is regulated by Fe concentration in the surrounding environment. The presence of siderophores suggests that bacteria are producing these compounds despite the overall higher Fe concentrations in the deep ocean and in hydrothermal plumes. Consistent with siderophore utilization in terrestrial ecosystems, one hypothesis is that siderophore production is beneficial to bacteria in the plumes in transforming Fe from otherwise inaccessible forms, such as particulate nano pyrites or Fe oxyhydroxides. The detection and identification of siderophores in these samples — and their relationship with dFe — provides compelling evidence that microbial production of ligands is responsible for some portion of the tight coupling between L_1 and dFe in hydrothermal systems along the MAR.

2.2 Chemical composition of siderophores in hydrothermal systems

Marine ligand composition has been shown to change with environmental gradients (Boiteau et al., 2016), making the structure and functional groups of siderophores identified in and around hydrothermal samples of particular interest. Samples collected from on-axis spreading centers contained the highest dFe concentrations (> 20 nM) and samples from fracture zones, diffuse, and off-axis sites contained the lowest

dFe ($\leq$ 1 nM). Spreading centers contained a wider variety of siderophore types than diffuse/fracture zones, with the greatest number of distinct siderophore compounds identified at the Lucky Strike, Broken Spur, Rainbow, and TAG vent sites (**Fig. 2**). On average, 13 compounds were identified with high confidence per on-axis spreading center sample, compared with 5 identifications per diffuse/fracture zone sample, and 2.5 identifications per off-axis sample (**Fig. 2b, Fig. S4**). Mixed-type siderophores — siderophores containing different moieties that bind to Fe(III) — were most common at all sites. Hydroxamates were identified at and around spreading centers, yet none of these types were detected with high confidence in samples from diffuse/fracture zones (**Fig. S4**). At spreading centers, neutrally buoyant plumes at on-axis vents averaged more than twice the number of distinct siderophore types than off-axis sites (**Fig. S4**), suggesting the proximity to the vent field plays an important role in the diversity and abundance of siderophore types produced, likely related to the diversity of the microbial community and/or unique Fe acquisition strategies across vent sites.

Low Fe surface waters have been found to contain higher concentrations of amphiphilic siderophores compared to high Fe waters and amphiphilic siderophores appear to be more common in the ocean than terrestrial systems (Boiteau et al., 2016). Amphiphilic siderophores have long hydrocarbon tails that can be embedded into the lipid bilayer of the bacterial cell membrane providing a mechanism to shuttle Fe into the cell and prevent diffusive loss of Fe (Martinez et al., 2003). As a result, these microbes have a competitive advantage over other bacteria that produce non-amphiphilic siderophores which can more easily disperse. Amphiphilic siderophores accounted for over half (57%) of the high confidence identifications in our samples (**Fig. S5**), in line with the ubiquity of amphiphilic siderophores in the marine environment (Butler and Theisen, 2010). It is perhaps surprising that they were so common in our dataset, due to the elevated Fe concentrations observed here relative to the more Fe-poor surface ocean. Amphiphilic siderophores were found in total concentrations between 0.3-4.7 pM, with Rainbow containing the highest concentration (**Fig. 2d**, **Table S5**). Interestingly, these concentrations were on a similar scale to amphiphilic siderophores observed in other upper ocean environments (Boiteau et al., 2016; Bundy et al., 2018; Boiteau et al., 2019). Marine bacteria have been shown to produce suites of amphiphilic siderophores as a way to adapt to the change in hydrophilicity in the surrounding environment (Sandy and Butler, 2009; Homann et al., 2009), which could be an important coping strategy for bacteria living in environments with strong chemical gradients, enabling them to access rapidly changing physical and chemical forms of Fe. Unlike in the surface ocean where high concentrations of amphiphilic siderophores are observed in regions where Fe is a growth-limiting micronutrient for primary production (Boiteau et al., 2016), amphiphilic siderophore production in hydrothermal plumes could be a way for bacteria to store Fe as they are physically transported in and away from the plume. This could be similar to how the production of multiple siderophores in terrestrial and pathogenetic systems are used to access inorganic particulate Fe for cellular uptake, Fe storage, and absorption of other metals in association with Fe (Hider and Kong, 2010).

2.3 Microbial sources of siderophores in hydrothermal plumes

Microbial production of siderophores is a strategy for organisms to adapt or compete with other bacteria for Fe within a specific environment (Sandy and Butler, 2009). This would result in a coupling of organic ligand production and dFe concentrations. We examined the microbial community composition around Rainbow (St. 11- 200 km E of Rainbow and St. 17- 10 km S of Rainbow) and Lucky Strike (St. 7; **Table 1, Table S1**) using 16S rRNA gene-based amplicon sequencing to determine the presence of microbes that may have the metabolic potential to synthesize the siderophores (**Fig. 3, S11**). Diagnosing metabolic function from 16S rRNA data alone is limited, but the presence of taxa in these vent sites encoding siderophore biosynthetic gene clusters indicates whether the microbial community is genetically capable of producing the compounds we observed in the environment. Bacterial genera containing known siderophore-producers were found at all three MAR sites examined, and putative siderophore-producers represented 3-20% of the relative abundance of each community (**Fig. 3**). Putative siderophore-producers were more abundant in the 3 µm (particle-attached) size fraction than in the 0.2 µm (free-living) fraction, suggesting siderophore production is more common in particle-associated bacteria in hydrothermal environments.

We found microbial genera in our samples that have previously been shown to produce a subset of the siderophores identified here, including ferrioxamines, vibrioferrin, and acinetoferrin (Butler, 2005; Vraspir and Butler, 2009; Moore et al., 2021; Bundy et al., 2018; Boiteau et al., 2016). Genera with the genetic potential to produce ferrioxamines were present at all three sites, while those known to produce vibrioferrin were present at Lucky Strike and 10 km S of Rainbow, and those producing acinetoferrin present at 10 km S of Rainbow (**Table S1, S6**). Mycobactins were detected with high confidence in every sample of this study, and genes encoding mycobactin have previously been detected in a cultured organism from a hydrothermal system (Gu et al., 2019), but no putative mycobactin producers were identified in the 16S rRNA data in this study. We also detected woodybactin D with high confidence in five out of our eleven sites. Although these biosynthetic genes were not identified in any of the genera from our 16S rRNA gene amplicon sequences, woodybactin D is a carboxylate siderophore isolated from *Shewanella* (Carmichael et al., 2019), and groups of deep-sea *Shewanella* (Kato and Nogi, 2001) were found in the 16S rRNA data (**Fig. S11**). The biosynthesis genes for many of the siderophores we identified are unknown, and thus the fact that we only found genera capable of producing a subset of the siderophores we characterized in the water column is not surprising. However, the observation that a significant portion of the *in-situ* microbial community is capable of synthesizing siderophores (**Fig 3**) suggests that siderophore production is more widespread in both the marine environment and deep ocean than previously believed.

Although the production of siderophores is often assumed to be associated primarily with low Fe conditions, evidence that siderophores are ubiquitous in marine environments — including those considered to be Fe rich — has been increasing (Park et al., 2023). The higher dFe associated with hydrothermal plumes may still not be high enough to suppress siderophore production due to the requirements for bacteria growth

(Tortell et al., 1996). It is also likely that in higher Fe environments, such as hydrothermal plumes, not all of the Fe is bio-accessible. For example, soil microbes actively secrete siderophores in the high Fe soil environment to solubilize otherwise inaccessible particulate Fe (Crowley et al., 1991). Similar processes could be occurring in hydrothermal plumes, where Fe mineral phases are common (Hoffman et al., 2020). It is telling that most siderophore-producing genera were found in the particulate size fraction (**Fig. 3**), providing additional evidence that siderophores might be produced to solubilize different forms of particulate Fe. Further work that assesses why bacteria are producing siderophores in high Fe neutrally buoyant plumes will be important for understanding the role of these complexes in facilitating the exchange of Fe between dissolved and particulate phases (Fitzsimmons et al., 2017), and how Fe-siderophore complexes may be stabilizing dFe in hydrothermal plumes.

3 Conclusion

Organic Fe-binding ligands have been implicated in playing a critical role in the preservation and transport of hydrothermal Fe into the ocean interior (Hoffman et al., 2018; Resing et al., 2015; Fitzsimmons et al., 2017; Toner et al., 2009; Bennett et al., 2011, 2008; Buck et al., 2018; Sander and Koschinsky, 2011). In this work, we have identified L_1 ligands as being tightly coupled to dFe in neutrally buoyant hydrothermal plumes along the MAR and have provided evidence that these ligands are microbially produced. We identified, for the first time, specific siderophores that demonstrate a conservative lower bound on the concentrations of L_1 ligands being produced by the microbial community in response to external hydrothermal Fe inputs. Bacteria putatively capable of synthesizing the siderophores were largely present in the particulate size fraction along the MAR, which adds to a growing body of evidence that microbes are likely using L_1 ligands like siderophores to access particulate or otherwise inaccessible Fe pools and are a key mechanism for controlling dFe in neutrally buoyant plumes. Exploring additional types of ligands within the L_1 pool and whether the Fe-organic ligand production is tightly coupled across additional hydrothermal vent systems will aid in constraining the biogeochemical importance of microbial feedbacks in impacting the hydrothermal dFe supply to the deep ocean.

Appendix: Materials and Methods

Sampling and cruise transect

Samples were collected as part of the 2017-2018 U.K. GEOTRACES GA13 section cruise along the Mid-Atlantic Ridge. Water samples from 11 venting and near venting locations were collected using a Seabird 911 conductivity, temperature, and depth (CTD) titanium rosette using conducting Kevlar wire with an oxidation-reduction potential (ORP) sensor to detect plumes. Teflon coated OTE (Ocean Test Equipment) bottles were pressurized to approximately 7 psi with 0.2 μm filtered air using an oil free compressor. A Sartobran 300 (Sartorius) filter capsule (0.2 μm) was used to collect filtered seawater samples into clean 250 mL LDPE sample bottles. Bottles and caps were rinsed 3 times with the filtered sample before being filled.

Samples were stored frozen at -20°C for Fe-organic ligand characterization by voltammetry and mass spectrometry.

Fe-binding ligand concentration and binding strengths Competitive Ligand Exchange-Adsorptive Cathodic Stripping Voltammetry

Fe-binding ligand concentrations and binding strengths were determined by competitive ligand exchange-adsorptive cathodic stripping voltammetry (CLE-ACSV) with a BAS*i* controlled growth mercury electrode (CGME) with an Ag/AgCl⁻ reference electrode and platinum auxiliary electrode (Bioanalytical Systems Incorporated). Using previously established methods (Buck et al., 2015, 2018; Bundy et al., 2018; Abualhaija and van den Berg, 2014; Hawkes et al., 2013b), 40 frozen filtrate (<0.2 µm) samples with dFe concentrations between 0.41-11.67 nM (**Table S1-S2**) were thawed in a 4°C fridge prior to analysis. A 15-point titration curve was analyzed for each sample. Briefly, within each titration, every point sequentially received 10 mL of sample, 7.5 µM of borate-ammonium buffer, 10 µM salicylaldoxime (SA) added ligand, and a dFe addition. Data was collected using the *Epsilon Eclipse Electrochemical Analyzer* (v.213) with a deposition time of 120 seconds and analyzed using *ElectroChemical Data Software* (v2001-2014) and *ProMCC* (v2008-2018) to determine peak areas and Fe-binding ligand parameters, respectively.

Reverse Titration-CLE-ACSV

Reverse titration-CLE-ACSV (RT-CLE-ACSV) (Hawkes et al., 2013a) was completed on 10 samples from Broken Spur, and TAG hydrothermal vent fields with dFe concentrations between 19.01-90.25 nM (**Table S1-S2**). Briefly, a 10-point titration curve was analyzed for each sample with each titration point consisting of 10 mL of sample buffered with 7.5 µM boric acid and the competitive ligand 1-nitroso-2-napthol (NN) additions. All samples were analyzed on a BAS*i* Controlled Growth Mercury Electrode (CGME) with the *Epsilon Eclipse Electrochemical Analyzer* (v.213) and deposition time of 120 seconds. For each sample, competitive ligand NN additions were 0.5, 1, 2, 3, 4, 6, 9, 15, 20, and 40 µM. Samples were equilibrated overnight and purged with N_2 (99.99%) for 5 minutes before analysis. At the end of each titration, three Fe additions (3-15 nM) were added to the final titration point to get the total concentration of Fe in equilibrium with ligands. Data was analyzed using *ElectroChemical Data Software* (v2001-2014) to acquire peak areas and a package in R using the model parameters of $\beta_{FeNN3} = 5.12 \times 10^{16}$, $\chi_{min} = 0.8$, $\chi_{max} = 0.9$, and *c1high* = 0.75 to determine the Fe-binding ligand parameters (Hawkes et al., 2013a). These parameters were chosen based on the recommendations for undersaturated samples and titrations curves where *ip_max* was not reached (Hawkes et al., 2013a). All other parameters within the model we kept at the default values.

Fe-binding organic ligand quantification and characterization

In addition to determining the total concentrations of strong Fe-binding ligands (L_1), we also identified and quantified siderophores that contributed to the L_1 ligand pool. Between 0.65-1.5 L of 0.2 μm filtered seawater pooled from ligand samples at each site (described above) was pumped slowly (15-20 mL min⁻¹) onto a polystyrene-divinylbenzene (Bond Elut ENV) solid phase extraction (SPE) column (Bundy et al., 2018;

Boiteau et al., 2016). SPE columns were rinsed with MilliQ and stored at -20ºC until analysis. For the analytical measurements, samples were thawed in the dark, eluted in 12 mL of distilled methanol, and dried down to between 0.2-0.5 mL of sample eluent (**Table S1**). Aliquots were analyzed by reverse-phase liquid chromatography (LC) on a trace metal clean bio-inert LC (Thermo Dionex 3000 NCS). The LC was interfaced with an electrospray ionization-mass spectrometer (ESI-MS; Thermo Q-Exactive HF) to identify and quantify the compounds based on accurate mass (MS^1) and the fragmentation (MS^2) data (Bundy et al., 2018; Boiteau et al., 2016). MSconvert (Proteowizard) was used to convert MS data to an open source mzxML format, and two stages of data processing were conducted using modified versions of previously reported R scripts (Bundy et al., 2018; Boiteau et al., 2016). In the first stage, mzxML files were read into R using new package "RaMS" (Kulmer and Ingalls, 2022), and extracted ion chromatograms (EICs) were generated for each targeted m/z of interest from an in-house database of siderophores. The m/z targets were the ionized apo, ^{54}Fe-bound, and ^{56}Fe-bound version of each siderophore, with a tolerance of 7.5 ppm. Putative siderophore candidates were filtered through a series of hard thresholds, such that MS^1 spectra were quality controlled to contain a minimum of 25 datapoints and the maximum intensity of each EIC was greater than 1e4 counts. Spectra meeting these criteria and containing either ^{54}Fe-bound and ^{56}Fe-bound m/z peaks within 30 seconds of each other or an apo peak were displayed for the user to further inspect peak quality and make the final decision of whether to move on to stage two of processing with a given siderophore candidate.

Stage two of processing extracted MS^2 spectra of the apo and Fe-bound forms of candidate siderophores to compare with the predicted MS^2 generated by *in silico* fragmenter MetFrag (Ruttkies et al., 2016). The *in silico* fragmenter feature was run with a tolerance of 10 ppm on "$[M+H]^+$" and "$[M+Na]^+$" modes. A confidence level of 1-4, from highest to lowest confidence, was then assigned to putative siderophores based on the following criteria: (1) peaks were present in MS^1 and MS^2 spectra, and at least one of the three most-intense MS^2 fragments matched *in silico* fragmentation, (2) peaks were present in MS^1 and MS^2 spectra, and smaller-intensity fragments matched *in silico* fragmentation, (3) peaks were present in MS^1 and MS^2 spectra, but little to no fragments matched *in silico* fragmentation, and (4) nicely shaped peaks were identified in MS^1 spectra but no MS^2 spectra was collected (outlined in **Table S4**; example spectra in **Fig. S6-S9**). The confidence levels were modelled after reporting standards for metabolite identification (Sumner et al., 2007). MetFrag pulls chemical structures from publicly-available databases like PubChem or COCONUT(Sorokina et al., 2021), which contain most, but not all variations of siderophores. As such, Fe-bound candidates were usually run against the apo form available in the database, and for siderophores with similar structures but variations in fatty chain length or double bond placement, sometimes only one parent structure was available.

A 5-point standard curve with known concentrations of siderophore ferrioxamine E was used for quantification of putative siderophores, with a limit of detection of 0.257 nM in the eluent (**Fig. S10**), or 0.07-0.21 pM in the sample depending on sample-to-eluent volume ratio at each site (**Table S1**). MS^1 peaks

were integrated for all putatively identified siderophores and peak areas were converted to concentration using the standard curve and the concentration factor of sample volume to eluent volume (**Fig. S10**). Commercial standards are not available for most siderophores, and different compounds have distinct ionization efficiencies in ESI-MS. Thus, the siderophore concentrations reported here are estimates of siderophore concentrations in these environments based on ferrioxamine E. Additionally, 1 mM of cyanocobalamin was added as an internal standard to each sample aliquot to address any changes in sensitivity during LC-ESI-MS runs. All putative siderophores that were identified with peak areas less than the detection limit were discarded, and all remaining putative compounds with at least confidence levels 1 and 2 at one site were included in the manuscript and are referred to as siderophores throughout. Siderophore identifications remain putative due to inherent uncertainty with assignments by mass, but the confidence levels were designed such that high confidence candidates contain siderophore-like moieties in their fragments. Limited sample volumes prevented analysis via LC-ICP-MS like previous studies, which, in addition to greater availability of commercial standard, would allow definitive identification in future work. Confidence level 3 and 4 putative siderophores are only included in the Supplementary Information (**Table S5**). In a final step of quality control, EICs for ^{13}C isotopologues of candidates were inspected to verify matching peak structure.

Microbial community analysis

Microbial community composition was assessed in neutrally buoyant plumes and near venting sites at three sites: Lucky Strike (Station 7; 1670 m), 10 km S of Rainbow (Station 17; 2000 m), and 200 km E of Rainbow (Station 11; 600 m, 1600 m and 2250 m). A range of 1- 2 L of seawater were filtered by pressure filtration through sequential 25 mm membrane filters housed in polypropylene filter holders (Whatman SwinLok, GE Healthcare, Pittsburgh, Pennsylvania) using a peristaltic pump and silicone tubing. Samples first passed through a 3 µm pore-size polyester membrane filter (Sterlitech, Auburn, Washington) then onto a 0.2 µm pore-size polyethersulfone membrane filter (Supor-200, Pall Corporation, Port Washington, New York). Pump tubing was acid washed with 10% hydrochloric acid and flushed with ultrapure water between each sample. The filters were flash frozen in liquid nitrogen in 2 mL gasketed bead beating tubes (Fisher Scientific) at sea.

Nucleic acids (DNA) were extracted as described previously(Santoro et al., 2010), with slight modifications. Briefly, cells on the filters were lysed directly in the bead beating tubes with sucrose-ethylene diamine tetraacetic acid (EDTA) lysis buffer (0.75 M sucrose, 20 mM EDTA, 400 mM NaCl, 50 mM Tris) and 1% sodium dodecyl sulfate. Tubes were then agitated in a bead beating machine (Biospec Products) for 1 min, and subsequently heated for 2 min. at 99°C in a heat block. Proteinase K (New England Biolabs) was added to a final concentration of 0.5 mg/mL. Filters were incubated at 55°C for approximately 4 h and the resulting lysates were purified with the DNeasy kit (Qiagen) using a slightly modified protocol (Santoro et al., 2010).

The purified nucleic acids were eluted in 200 μL of DNase, RNase-free water, and quantified using a fluorometer (Qubit and Quanti-T HS reagent, Invitrogen Molecular Probes).

The 16S rRNA gene was amplified in all samples using V4 primers (Apprill et al., 2015; Parada et al., 2016) (515F-Y and 806RB) following a previously established protocol (Stephens et al., 2020). Amplicons were sequenced using a paired-end 250bp run on an Illumina MiSeq 500 and demultiplexed by the UC Davis Genome Center. The resulting 16S rRNA amplicon sequences were filtered and trimmed using the DADA2 pipeline in R(Callahan et al., 2016). Taxonomic assignments were made with version 138.1 of the SILVA SSU database (Quast et al., 2013) (silva_nr99_v138.1_wSpecies_train_set.fa.gz ; doi:10.5281/zenodo.4587955; accessed March 2022). Chloroplast and mitochondrial sequences were filtered out of the dataset using the 'phyloseq' R package (v 1.38.0), after which samples had read depths ranging from 9375 – 65486 reads (average 28425 ± 20014 reads) and represented 1010 unique amplicon sequence variants (ASVs). Read counts were transformed from absolute to relative abundance and taxa were aggregated to the Family level. The ten most abundant families present in each sample were visualized using the 'ggplot2' package (v. 3.3.5).

In order to assess the potential of the observed prokaryotic taxa to produce siderophores, we downloaded all siderophore biosynthetic gene clusters (BGCs) in the antismash secondary metabolite database ($n = 7909$) and used text-string matching to compare genera containing these BGCs to the genera found in our 16S rRNA gene dataset(Blin et al., 2021). We cross-referenced the nomenclature of antismash-predicted siderophores with that of the siderophores identified by LC-ESI-MS in this study, accounting for minor differences in naming convention between the two databases, to determine if microbial community members present at each site were predicted to make any of the siderophores that were measured at that site. Station 38 and Station 12 were the closest sites with siderophore measurements for comparison against the taxonomic samples taken at 200 km E of Rainbow and 10 km S of Rainbow, respectively. Samples for microbial taxonomy and siderophore identity were taken from the same location at Lucky Strike and thus directly compared.

Data Availability

The CSV data reported in this study has been deposited at Zenodo under the DOI: http://doi.org/10.5281/zenodo.7325154. The LC-ES-MS data has been deposited on Massive under the DOI: http://doi.org/doi.10.25345/C5V97ZW7N. Microbial 16S rRNA data have been deposited on GenBank under the accession number BioProject #PRJNA865382. All data is freely available on each of these data repositories.

Acknowledgments

We acknowledge the captain and crew of the R/V *James Cook*, Chief Scientist Alessandro Tagliabue, and Noah Gluschankoff for supporting this work. This study was a part of the FeRidge project (GEOTRACES section GA13) which was supported by the Natural Environment Research Council funding (NERC United Kingdom Grants NE/N010396/1 to MCL and NE/N009525/1 to AT). The International GEOTRACES Programme is possible in part thanks to the support from the U.S. National Science Foundation (Grant OCE-1840868) to the Scientific Committee on Oceanic Research (SCOR). CLH was funded by JISAO/CICOES postdoctoral fellowship. PJM was funded through the NOAA Hollings Scholar summer program. JR was funded by NOAA Ocean Exploration and Research, NOAA Earth-Ocean Interactions programs at NOAA-Pacific Marine Environmental Labs, and JISAO/CICOES. Part of this work was carried out in the University of Washington TraceLab, which receives support from the M.J. Murdock Charitable Trust in conjunction with the University of Washington College of Environment, and the Pacific Marine Environmental Labs at the National Oceanic and Atmospheric Administration. Parts of this work was also carried out in Dr. Anitra Ingalls laboratory with the help of Laura Truxal and Dr. Jiwoon Park at the University of Washington-School of Oceanography.

Author Contributions: Manuscript preparation, sample/data processing, CSV analysis, and interpretation (C.L.H.), manuscript preparation, LC-ESI-MS data analysis and interpretation (P.J.M.), microbial analysis and interpretation (J.B.A. and A.E.S.), dissolved iron and derived excess $^3He_{xs}$ measurements, sample collection (A.J.M. L. and M.C.L.), microbial data collection and ligand data interpretation (T.M. and K.N.B.), and project design and planning, data interpretation, and mentoring (A.T., M.C.L., J.A.R., and R.M.B.). All authors were involved in editing manuscript.

Competing Interest Statement: The authors declare no competing interests.

References

Abualhaija, M. M. and van den Berg, C. M. G. G.: Chemical speciation of iron in seawater using catalytic cathodic stripping voltammetry with ligand competition against salicylaldoxime, Mar. Chem., 164, 60–74, https://doi.org/10.1016/j.marchem.2014.06.005, 2014.

Apprill, A., Mcnally, S., Parsons, R., and Weber, L.: Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton, Aquat. Microb. Ecol., 75, 129–137, https://doi.org/10.3354/ame01753, 2015.

Bazylev, B. A.: Allochemical Metamorphism of Mantle Peridoties in the Hayes Fracture Zone of the North Atlantic, Petrology, 5, 362–379, 1997.

Beaulieu, S. E. and Szafrański, K. M.: InterRidge Global Database of Active Submarine Hydrothermal Vent Fields Version 3.4, https://doi.org/10.1594/PANGAEA.917894, 2020.

Bennett, S. a., Achterberg, E. P., Connelly, D. P., Statham, P. J., Fones, G. R., and German, C. R.: The distribution and stabilisation of dissolved Fe in deep-sea hydrothermal plumes, Earth Planet. Sci. Lett., 270,

157–167, https://doi.org/10.1016/j.epsl.2008.01.048, 2008.

Bennett, S. a., Hansman, R. L., Sessions, A. L., Nakamura, K. ichi, and Edwards, K. J.: Tracing iron-fueled microbial carbon production within the hydrothermal plume at the Loihi seamount, Geochim. Cosmochim. Acta, 75, 5526–5539, https://doi.org/10.1016/j.gca.2011.06.039, 2011.

Blin, K., Shaw, S., Kautsar, S. A., Medema, M. H., and Weber, T.: The antiSMASH database version 3: Increased taxonomic coverage and new query features for modular enzymes, Nucleic Acids Res., 49, D639–D643, https://doi.org/10.1093/nar/gkaa978, 2021.

Boiteau, R. M., Mende, D. R., Hawco, N. J., McIlvin, M. R., Fitzsimmons, J. N., Saito, M. A., Sedwick, P. N., DeLong, E. F., and Repeta, D. J.: Siderophore-based microbial adaptations to iron scarcity across the eastern Pacific Ocean, Proc. Natl. Acad. Sci., 113, 14237–14242, https://doi.org/10.1073/pnas.1608594113, 2016.

Boiteau, R. M., Till, C. P., Coale, T. H., Fitzsimmons, J. N., Bruland, K. W., and Repeta, D. J.: Patterns of iron and siderophore distributions across the California Current System, Limnol. Oceanogr., 64, 376–389, https://doi.org/10.1002/lno.11046, 2019.

Buck, K. N., Lohan, M. C., Berger, C. J. M., and Bruland, K. W.: Dissolved iron speciation in two distinct river plumes and an estuary: Implications for riverine iron supply, Limnol. Oceanogr., 52, 843–855, https://doi.org/10.4319/lo.2007.52.2.0843, 2007.

Buck, K. N., Sohst, B., and Sedwick, P. N.: The organic complexation of dissolved iron along the U.S. GEOTRACES (GA03) North Atlantic Section, Deep. Res. Part II Top. Stud. Oceanogr., 116, 152–165, https://doi.org/10.1016/j.dsr2.2014.11.016, 2015.

Buck, K. N., Sedwick, P. N., Sohst, B., and Carlson, C. A.: Organic complexation of iron in the eastern tropical South Pacific: Results from US GEOTRACES Eastern Pacific Zonal Transect (GEOTRACES cruise GP16), Mar. Chem., 201, 229–241, https://doi.org/10.1016/j.marchem.2017.11.007, 2018.

Bundy, R. M., Abdulla, H. A. N., Hatcher, P. G., Biller, D. V., Buck, K. N., and Barbeau, K. A.: Iron-binding ligands and humic substances in the San Francisco Bay estuary and estuarine-influenced shelf regions of coastal California, Mar. Chem., 173, 183–194, https://doi.org/10.1016/j.marchem.2014.11.005, 2015.

Bundy, R. M., Boiteau, R. M., McLean, C., Turk-Kubo, K. A., McIlvin, M. R., Saito, M. A., Mooy, B. A. Van, and Repeta, D. J.: Distinct Siderophores Contribute to Iron Cycling in the Mesopelagic at Station ALOHA, Front. Mar. Sci., 1–15, https://doi.org/10.3389/fmars.2018.00061, 2018.

Butler, A.: Marine siderophores and microbial iron mobilization., Biometals, 18, 369–374, https://doi.org/10.1007/s10534-005-3711-0, 2005.

Butler, A. and Theisen, R. M.: Iron(III)-siderophore coordination chemistry: Reactivity of marine siderophores., Coord. Chem. Rev., 254, 288–296, https://doi.org/10.1016/j.ccr.2009.09.010, 2010.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P.: DADA2: High-resolution sample inference from Illumina amplicon data, Nat. Methods, 13, 581–583, https://doi.org/10.1038/nmeth.3869, 2016.

Carmichael, J. R., Zhou, H., and Butler, A.: A suite of asymmetric citrate siderophores isolated from a marine Shewanella species, J. Inorg. Biochem., 198, 1–6, https://doi.org/10.1016/j.jinorgbio.2019.110736, 2019.

Cowen, J. P. and Bruland, K. W.: Metal deposits associated with bacteria: implications for Fe and Mn marine biogeochemistry, Deep Sea Res. Part A. Oceanogr. Res. Pap., 32, 253–272, https://doi.org/10.1016/0198-0149(85)90078-0, 1985.

Cowen, J. P., Massoth, G. J., and Feely, R. A.: Scavenging rates of dissolved manganese in a hydrothermal vent plume, Deep Sea Res. Part A. Oceanogr. Res. Pap., 37, 1619–1637, https://doi.org/10.1016/0198-

0149(90)90065-4, 1990.

Crowley, D. E., Wang, Y. C., Reid, C. P. P., and Szaniszlo, P. J.: Mechanisms of iron acquisition from siderophores by microorganisms and plants, Plant Soil, 130, 179–198, 1991.

Fishwick, M. P., Sedwick, P. N., Lohan, M. C., Worsfold, P. J., Buck, K. N., Church, T. M., and Ussher, S. J.: The impact of changing surface ocean conditions on the dissolution of aerosol iron, Global Biogeochem. Cycles, 28, 1235–1250, https://doi.org/10.1002/2014GB004921, 2014.

Fitzsimmons, J. N., John, S. G., Marsay, C. M., Hoffman, C. L., Nicholas, S. L., Toner, B. M., German, C. R., and Sherrell, R. M.: Iron persistence in the distal hydrothermal plume supported by dissolved – particulate exchange, Nat. Geosci., 10, 1–8, https://doi.org/10.1038/ngeo2900, 2017.

Gu, H., Sun, Q., Luo, J., Zhang, J., and Sun, L.: A First Study of the Virulence Potential of a Bacillus subtilis Isolate From Deep-Sea Hydrothermal Vent, Front. Cell. Infect. Microbiol., 9, 1–14, https://doi.org/10.3389/fcimb.2019.00183, 2019.

Hawkes, J. A., Gledhill, M., Connelly, D. P., and Achterberg, E. P.: Characterisation of iron binding ligands in seawater by reverse titration, Anal. Chim. Acta, 766, 53–60, https://doi.org/10.1016/j.aca.2012.12.048, 2013a.

Hawkes, J. A., Connelly, D. P., Gledhill, M., and Achterberg, E. P.: The stabilisation and transportation of dissolved iron from high temperature hydrothermal vent systems, Earth Planet. Sci. Lett., 375, 280–290, https://doi.org/10.1016/j.epsl.2013.05.047, 2013b.

Hider, R. C. and Kong, X.: Chemistry and biology of siderophores, Nat. Prod. Rep., 27, 637–657, https://doi.org/10.1039/b906679a, 2010.

Hoffman, C. L., Nicholas, S. L., Ohnemus, D. C., Fitzsimmons, J. N., Sherrell, R. M., German, C. R., Heller, M. I., Lee, J. mi, Lam, P. J., and Toner, B. M.: Near-field iron and carbon chemistry of non-buoyant hydrothermal plume particles, Southern East Pacific Rise 15°S, Mar. Chem., 201, 183–197, https://doi.org/10.1016/j.marchem.2018.01.011, 2018.

Hoffman, C. L., Schladweiler, C., Seaton, N. C. A., Nicholas, S. L., Fitzsimmons, J., Sherrell, R. M., German, C. R., Lam, P., and Toner, B. M.: Diagnostic morphology and solid-state chemical speciation of hydrothermally derived particulate Fe in a long-range dispersing plume, ACS Earth Sp. Chem., 4, 1831–1842, https://doi.org/10.1021/acsearthspacechem.0c00067, 2020.

Homann, V. V, Sandy, M., Tincu, J. A., Templeton, A. S., Tebo, B. M., and Butler, A.: Loihichelins A - F , a Suite of Amphiphilic Siderophores Produced by the Marine Bacterium Halomonas LOB-5, J. Nat. Prod., 72, 884–888, 2009.

Kato, C. and Nogi, Y.: Correlation between phylogenetic structure and function : examples from deep-sea Shewanella, 35, 223–230, 2001.

Kelley, D. S. and Shank, T. M.: Hydrothermal systems: A decade of discovery in slow spreading environments, Geophys. Monogr. Ser., 188, 369–407, https://doi.org/10.1029/2010GM000945, 2010.

Kulmer, W. and Ingalls, A. E.: The R Journal: Tidy Data Neatly Resolves Mass-Spectrometry's Ragged Arrays, R J., 2022.

Lauderdale, J. M., Braakman, R., Forget, G., Dutkiewicz, S., and Follows, M. J.: Microbial feedbacks optimize ocean iron availability, Proc. Natl. Acad. Sci. U. S. A., 117, 4842–4849, https://doi.org/10.1073/pnas.1917277117, 2020.

Li, M., Toner, B. M., Baker, B. J., Breier, J. a, Sheik, C. S., and Dick, G. J.: Microbial iron uptake as a mechanism for dispersing iron from deep-sea hydrothermal vents., Nat. Commun., 5, 3192, https://doi.org/10.1038/ncomms4192, 2014.

Lough, A. J. M., Tagliabue, A., Demasy, C., Resing, J. A., Mellett, T., Wyatt, N. J., and Lohan, M. C.: The impact of hydrothermal vent geochemistry on the addition of iron to the deep ocean, Biogeosciences Discuss., [preprint], 1–23, https://doi.org/10.5194/bg-2022-73, 2022.

Manck, L. E., Park, J., Tully, B. J., Poire, A. M., Bundy, R. M., Dupont, C. L., and Barbeau, K. A.: Petrobactin, a siderophore produced by Alteromonas, mediates community iron acquisition in the global ocean, ISME J., 16, 358–369, https://doi.org/10.1038/s41396-021-01065-y, 2022.

Martinez, J. S., Carter-Franklin, J. N., Mann, E. L., Martin, J. D., Haygood, M. G., and Butler, A.: Structure and membrane affinity of a suite of amphiphilic siderophores produced by a marine bacterium, Proc. Natl. Acad. Sci. U. S. A., 100, 3754–3759, https://doi.org/10.1073/pnas.0637444100, 2003.

Mellett, T., Albers, J. B., Santoro, A., Wang, W., Salaun, P., Resing, J., Lough, A. J. ., Tagliabue, A., Lohan, M., Bundy, R. M., and Buck, K. N.: Particle exchange mediated by organic ligands in incubation experiments of hydrothermal vent plumes along the mid-Atlantic Ridge, n.d.

Moore, L. E., Heller, M. I., Barbeau, K. A., Moffett, J. W., and Bundy, R. M.: Organic complexation of iron by strong ligands and siderophores in the eastern tropical North Pacific oxygen deficient zone, Mar. Chem., 236, 104021, https://doi.org/10.1016/j.marchem.2021.104021, 2021.

Omanović, D., Garnier, C., and Pižeta, I.: ProMCC: An all-in-one tool for trace metal complexation studies, Mar. Chem., 173, 25–39, https://doi.org/10.1016/j.marchem.2014.10.011, 2015.

Parada, A. E., Needham, D. M., and Fuhrman, J. A.: Every base matters: Assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples, Environ. Microbiol., 18, 1403–1414, https://doi.org/10.1111/1462-2920.13023, 2016.

Park, J., Durham, B. P., Key, R. S., Groussman, R. D., Pinedo-Gonzalez, P., Hawco, N. J., John, S. G., Carlson, M. C. G., Lindell, D., Juranek, L., Ferrón, S., Ribalet, F., Armbrust, E. V., Ingalls, A. E., and Bundy, R. M.: Siderophore production and utilization by microbes in the North Pacific Ocean, bioRxiv, 2022.02.26.482025, https://doi.org/10.1101/2022.02.26.482025, 2022.

Park, J., Durham, B. P., Key, R. S., Groussman, R. D., Pinedo-Gonzalez, P., Hawco, N. J., John, S. G., Carlson, M. C. G., Lindell, D., Juranek, L., Ferrón, S., Ribalet, F., Armbrust, E. V., Ingalls, A. E., and Bundy, R. M.: Siderophore production and utilization by microbes in the North Pacific Ocean, Limnol. Oceanogr., 2022.02.26.482025, https://doi.org/10.1002/lno.12373, 2023.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.: The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools, Nucleic Acids Res., 41, 590–596, https://doi.org/10.1093/nar/gks1219, 2013.

Resing, J. a., Sedwick, P. N., German, C. R., Jenkins, W. J., Moffett, J. W., Sohst, B. M., and Tagliabue, A.: Basin-scale transport of hydrothermal dissolved metals across the South Pacific Ocean, Nature, 523, 200–203, https://doi.org/10.1038/nature14577, 2015.

Ruttkies, C., Schymanski, E. L., Wolf, S., Hollender, J., and Neumann, S.: MetFrag relaunched: incorporating strategies beyond in silico fragmentation, J. Cheminform., 8, 1–16, https://doi.org/10.1186/s13321-016-0115-9, 2016.

Sander, S. G. and Koschinsky, A.: Metal flux from hydrothermal vents increased by organic complexation, Nat. Geosci., 4, 145–150, https://doi.org/10.1038/ngeo1088, 2011.

Sandy, M. and Butler, A.: Microbial iron acquisition: marine and terrestrial siderophores., Chem. Rev., 109, 4580–95, https://doi.org/10.1021/cr9002787, 2009.

Santoro, A. E., Casciotti, K. L., and Francis, C. A.: Activity, abundance and diversity of nitrifying archaea and bacteria in the central California Current, Environ. Microbiol., 12, 1989–2006, https://doi.org/10.1111/j.1462-2920.2010.02205.x, 2010.

Sorokina, M., Merseburger, P., Rajan, K., Yirik, M. A., and Steinbeck, C.: COCONUT online: Collection of Open Natural Products database, J. Cheminform., 13, 1–13, https://doi.org/10.1186/s13321-020-00478-9, 2021.

Stephens, B. M., Opalk, K., Petras, D., Liu, S., Comstock, J., Aluwihare, L. I., Hansell, D. A., and Carlson, C. A.: Organic Matter Composition at Ocean Station Papa Affects Its Bioavailability, Bacterioplankton Growth Efficiency and the Responding Taxa, Front. Mar. Sci., 7, https://doi.org/10.3389/fmars.2020.590273, 2020.

Sumner, L. W., Amberg, A., Barrett, D., Beale, M. H., Beger, R., Daykin, C. A., Fan, T. W.-M., Fiehn, O., Goodacre, R., Griffin, J. L., Hankemeier, T., Hardy, N., Harnly, J., Higashi, R., Kopka, J., Lane, A. N., Lindon, J. C., Marriott, P., Nicholls, A. W., Reily, M. D., Thaden, J. J., and Viant, M. R.: Proposed minimum reporting standards for chemical analysis, Metabolomics, 3, 211–221, https://doi.org/10.1007/s11306-007-0082-2, 2007.

Tagliabue, A., Bowie, A. R., Boyd, P. W., Buck, K. N., Johnson, K. S., and Saito, M. A.: The integral role of iron in ocean biogeochemistry, Nature, 543, 51–59, https://doi.org/10.1038/nature21058, 2017.

Toner, B. M., Fakra, S. C., Manganini, S. J., Santelli, C. M., Marcus, M. a., Moffett, J. W., Rouxel, O., German, C. R., and Edwards, K. J.: Preservation of iron(II) by carbon-rich matrices in a hydrothermal plume, Nat. Geosci., 2, 197–201, https://doi.org/10.1038/ngeo433, 2009.

Tortell, P. D., Maldonado, M. T., and Price, N. M.: The role of heterotrophic bacteria in iron-limited ocean ecosystems, Nature, 383, 330–332, https://doi.org/10.1038/383330a0, 1996.

Vraspir, J. M. and Butler, A.: Chemistry of marine ligands and siderophores., Ann. Rev. Mar. Sci., 1, 43–63, https://doi.org/10.1146/annurev.marine.010908.163712, 2009.

Waska, H., Koschinsky, A., Ruiz Chancho, M. J., and Dittmar, T.: Investigating the potential of solid-phase extraction and Fourier-transform ion cyclotron resonance mass spectrometry (FT-ICR-MS) for the isolation and identification of dissolved metal-organic complexes from natural waters, Mar. Chem., 173, 78–92, https://doi.org/10.1016/j.marchem.2014.10.001, 2015.

Figures and Tables

Figure 1. Dissolved iron is strongly correlated with L_1 iron-binding ligands in diverse hydrothermal systems. (a) Station map showing the 11 sites investigated along the MAR. Known hydrothermal vents are marked as red triangles(Beaulieu and Szafrański, 2020). Two expanded inset maps for (b) Rainbow and (c) TAG hydrothermal vent fields. For additional information about vent site characteristics refer to **Table 1**. (d) dFe versus L_1 iron-binding ligands at each vent site in this study showing a ~1:1 correlation (m= 0.88, R^2 = 0.96) with dFe in neutrally-buoyant plumes along the MAR. (e) dFe versus L_1 ligands from previous studies over the ridge axis and ~80 km from ridge axis in the Southern East Pacific Rise hydrothermal plume(Buck et al., 2018), and over TAG hydrothermal vent field(Buck et al., 2015). The solid black lines in (d) and (e) are the 1:1 ratio line between dFe and ligand concentrations, and dashed lines show the linear regression for the corresponding data. Square symbols refer to spreading centers, triangles refer to fracture zones, and

circles refer to alkaline vents. Error bars represent the 95% confidence interval of the data fit as calculated by ProMCC(Omanović et al., 2015). The map was created using GeoMapApp version 3.6.14.

Figure 2. **Siderophore presence in hydrothermal plumes along the MAR.** (a) Heat map of confidence levels 1-2 (blue gradient, 1 = highest confidence). Gray boxes indicate a detection with lower confidence (see Methods), and white boxes indicate no detection at those sites. The y-axis is ordered from top to bottom in terms of descending mass of the apo (without Fe) form of the siderophore. (b) Model II ordinary least squares regression on dFe versus summed siderophore concentrations (of detections in Fig. 2b), calculated from peak areas, at each site. Since the siderophore analysis was performed on pooled samples, the dFe values in the regression are weighted values based on measured dFe and volume of each constituent of the pooled sample. The vertical error bars represent the standard deviation of dFe of the constituents. TAG (St. 35) was not included in the regression due to its large range of dFe values and outlier behavior. (c-d) Fe bound versus total summed concentration of (c) all types of siderophores and (d) amphiphilic siderophores at each station. The vertical green lines separate fracture/diffuse sites from off-axis sites and vertical red lines separate off-axis from on-axis sites as defined in Table 1. Symbols follow Fig. 1.

Figure 3. Relative abundance of putative siderophore-producing taxa. Bar height indicates the proportion of 16S rRNA genes recovered in each sample, separated by depth, filter size fraction, and site location. Colors correspond to taxonomy. Genera found in MAR vent microbial communities with members in the antismash database predicted to produce siderophores are depicted at the family level.

Table 1. Characteristics of sample locations along the Mid Atlantic Ridge.

Vent Names	Abbr.	Station	Geology	Host rock	Vent type	Spreading rate (mm/yr)	Summed putative siderophore concentration (pM)	Summed Siderophore concentration/ L_1 ligand (%)*
Lucky Strike	LS	7/8	Spreading Center	gabbro	Black smoker	20.2	4.38	0.034-0.19
33 km E of Rainbow	CER	12	Spreading Center	-	-	-	0.537	0.013-0.017
Rainbow	R	38	Spreading Center	ultramafic	Black smoker	20.6	12.1	*n.a.*
Hayes Fracture Zone	HFZ	21	Fracture Zone	peridotites/gabbro	-	21.2	2.74	0.20-0.39
Lost City	LC	23	Fracture Zone	ultramafic/gabbro	Alkaline	22.6	3.47	0.27-0.35
Broken Spur	BS	24	Spreading Center	gabbro	Black smoker/diffuse	22.9	6.30	0.07-0.29
29 km N of TAG	CNT	26	Spreading Center	-	-	-	0.968	0.014-0.079
30 km W of TAG	CWT	30	Spreading Center	-	-	-	2.91	0.15
30 km E of TAG	CET	31	Spreading Center	-	-	-	2.19	0.31

| Trans-Atlantic Geotraverse | TAG | 35 | Spreading Center | gabbro | Black smoker | 23.6 | 6.61 | 0.18 |
| Low Temp Slope | LTS | 37 | - | - | Diffuse fluids | - | 1.13 | 0.079-0.087 |

Spreading rates along the Mid-Atlantic Ridge were gathered from the Interridge Database v3.4. Host rock groups were determined from previously discussed classifications(Bazylev, 1997; Kelley and Shank, 2010). Off-axis sites –33 km E of Rainbow, 29 km N of TAG, 30 km E of TAG, and 30 km W of TAG– were far-field locations of their respective vent field. Low Temp Slope was a diffuse-dominated site that was sampled for the first time as a part of this study. Summed putative siderophore concentrations and the percent of L_1 ligand are reported for compounds detected with at least confidence level 1 and 2 at one site. These values do not take into account typical extraction efficiencies of ENV columns for Fe-binding organics. Average L_1 ligand and siderophore concentrations can be viewed in **Table S3** and concentrations for individual siderophores can be observed in **Table S5**.

*The siderophore sample at each site was pooled from ligand samples, so the percentage of siderophores in the L_1 pool is presented as a range based on the range of L_1 concentrations at each site.

n.a.= unable to be determined

- = unknown