Microbial siderophore production is tightly coupled to iron in hydrothermal plumes

Colleen L. Hoffman[1,2,3*], Patrick J. Monreal[3,4], Justine B. Albers[5], Alastair J.M. Lough[6], Alyson

E. Santoro[5], Travis Mellett[3,7], Kristen N. Buck[7], Alessandro Tagliabue[8], Maeve C. Lohan[6],

Joseph A. Resing[1,2,3], Randelle M. Bundy[3*]

1- Joint Institute for the Study of Atmosphere and Ocean, University of Washington, 3737

Brooklyn Avenue NE, Seattle, WA 98195, USA

2-Cooperative Institute for Climate, Ocean, and Ecosystem Studies, University of Washington,

3737 Brooklyn Avenue NE, Seattle, WA 98195, USA

3-School of Oceanography, University of Washington, 1501 NE Boat Street, Seattle, WA 98195,

USA

4- Department of Earth System Science, Stanford University, 473 Via Ortega, Stanford, CA

94305, USA

5- Department of Ecology, Evolution, and Marine Biology, University of California, Santa

Barbara, CA 93106, USA

6-Department of Ocean and Earth Sciences, National Oceanography Centre, University of

Southampton, European Way, Southampton SO14 3ZH, United Kingdom

7- College of Marine Science, University of South Florida, 140 7th Avenue South, St. Petersburg,

FL, 33701, USA

8-Department of Earth, Ocean, and Ecological Sciences, University of Liverpool, 4 Brownlow

Street, Liverpool l69 3GP, United Kingdom

*Corresponding authors clhoffma@uw.edu, rbundy@uw.edu

Abstract

Specialized molecules, such as siderophores, are used to access and retain iron in soluble forms by marine microorganisms. These siderophores form part of the ocean dissolved iron-binding ligand pool and are hypothesized to exert a key control on the persistence of iron in hydrothermal environments. To explore this hypothesis, we measured iron, iron-binding ligands, and siderophores from 11 geochemically distinct sites along a 1,700 km section of the Mid-Atlantic Ridge. We found siderophores at all sites and proximity to the vent played an important role in dictating siderophore types and diversity. The notable presence of amphiphilic siderophores may enable microbes to access particulate iron in hydrothermal plumes. The tight coupling between strong ligands and dissolved iron across six distinct hydrothermal environments, combined with the local presence of siderophore producing microbial genera suggests that biological production of siderophores exerts a key control on hydrothermal dissolved iron concentrations.

Introduction

Over the last few decades, observations and modelling efforts have increased our understanding about the critical role organic ligands play in the cycling, transport, and utilization of trace metals[1–4]. Strong iron-binding organic ligands (L_1), such as siderophores, are thermodynamically favored to complex and stabilize external sources of iron (Fe) and prevent its scavenging and removal. This is particularly important when ligands are present in excess of dissolved Fe (dFe)[5]. For example, in high dissolved and particulate Fe estuarine systems, only the dFe that escapes estuarine flocculation is bound to the strongest Fe-binding ligands[6,7].

Siderophores are a type of strong ligand produced by bacteria and fungi to facilitate uptake of Fe by complexation of Fe in the high affinity siderophore, drawing Fe out of otherwise inaccessible phases in the marine environment[8,9]. Historically, due to the energy required to produce siderophores, they have only primarily been considered as an important microbial strategy for Fe acquisition in the low-Fe surface ocean[8,10]. Compared to terrestrial soil ecosystems – where siderophores were first identified – Fe is several orders of magnitude lower in the marine environment. Their production could be a widespread microbial Fe acquisition strategy, which then tightly controls the utilization and availability of Fe[11]. A recent study identified siderophores in oxygen-deficient zones with relatively high Fe concentrations (>2 nM),

suggesting they are important for both Fe acquisition and offshore transport of Fe away from

sediment sources[2,4,6,12,13]. Genetic evidence in the Guaymas Basin hydrothermal system also

showed that >70% of Fe-related bacterial transcripts were involved in siderophore uptake,

regulation, and biosynthesis, providing additional evidence that siderophore production and

uptake may be important even in the higher Fe environments of the deep ocean[14].

While siderophore production could be common in the deep ocean[3], none of the previous studies

have directly measured siderophores in hydrothermal systems, explored the mechanisms for their

production, or determined their impact on Fe cycling in the ocean interior. This represents an

important gap in our understanding of how hydrothermal sources may regulate the deep ocean

dFe inventory. Here, for the first time, we identified the presence of siderophores and

siderophore producing microbes in 11 geochemically distinct hydrothermal environments,

including four black smokers (high temperature, high Fe), four sites off-axis sites (within 30 km

of black smokers), one diffuse-flow vent (low temperature, low Fe), one alkaline vent (pH 9-11,

very low Fe), and one non-vent fracture zone along the slow-spreading (20-50 mm/yr) Mid-

Atlantic Ridge (MAR) to understand whether microbial ligand production impacts the supply of

hydrothermal dFe to the ocean. Overall, our results show microbially-produced siderophores

were present in all sites and that strong L_1 ligands including siderophores exert a tight control on

hydrothermal dFe. This suggests that strong organic ligands produced by bacteria play a key role

in setting deep ocean Fe concentrations across global hydrothermal systems.

Results and Discussion

L_1 iron-binding ligands and the role of siderophores in hydrothermal plumes

Figure 1. Dissolved iron is strongly correlated with L_1 iron-binding ligands in diverse hydrothermal systems.
(a) Station map showing the 11 sites investigated along the MAR. Known hydrothermal vents are marked as red triangles[15]. Two expanded inset maps for (b) Rainbow and (c) TAG hydrothermal vent fields. For additional information about vent site characteristics refer to **Table 1**. (d) dFe versus L_1 iron-binding ligands at each vent site in this study showing a ~1:1 correlation (m= 0.88, R^2 = 0.96) with dFe in neutrally-buoyant plumes along the MAR. (e) dFe versus L_1 ligands from previous studies over the ridge axis and ~80 km from ridge axis in the Southern East Pacific Rise hydrothermal plume[2], and over TAG hydrothermal vent field[12]. The solid black lines in (d) and (e) are the 1:1 ratio line between dFe and ligand concentrations, and dashed lines show the linear regression for the corresponding data. Square symbols refer to spreading centers, triangles refer to fracture zones, and circles refer to alkaline vents. Error bars represent the 95% confidence interval of the data fit as calculated by ProMCC[16]. The map was created using GeoMapApp version 3.6.14.

The L_1 Fe-binding organic ligand pool is a heterogeneous mixture of compounds that are

operationally classified based on their binding strength with Fe ($\log K^{cond}_{Fe',FeL} > 12$). These L_1

ligands have been proposed to be important for stabilizing and transporting Fe in hydrothermal

plumes[1,2,17]. Here, the average binding strength (log $K^{cond}_{Fe',FeL}$) and concentration of organic Fe-

binding ligands were quantified in multiple vent systems that spanned a wide range in dFe

concentrations (0.41-90.3 nM). Over >99% of the dFe was complexed by L_1 ligands, which

meant Fe-free L_1 ligands capable of binding additional Fe were generally present in low

concentrations (< 1 nM; **Fig. S1**). As a result, dFe concentrations were tightly coupled to L_1

ligands in an ~1:1 ratio (dFe: L_1, **Fig. 1d**) similar to previous studies in other neutrally buoyant

plumes (**Fig. 1e**)[2,12,18]. However, the strong coupling between dFe and ligands was only observed

at sites where L_1 ligands were detected. Some sampling locations contained high concentrations

of weaker ligands (log $K^{cond}_{Fe',FeL}$ < 12, **Supplementary Table 2**) with no correlation to dFe. The

weak ligands were either not saturated with Fe, or contained extremely high concentrations of

dFe (> 20 nM) indicating potential sampling of a rising buoyant plume.

Altogether, our results suggest that L_1 ligands uniquely set the dFe concentration in these

neutrally-buoyant plumes. A similar direct control of dFe concentrations by L_1 ligands has been

previously observed in rivers[7] and aerosol solubility experiments[19]. Both the dFe and L_1 ligands

could originate from the high temperature vents themselves, yielding a tightly coupled

hydrothermal endmember. However, the concentration of L_1 ligands did not correlate with

excess mantle Helium-3 ($^3He_{xs}$, **Fig S2**, **Supplementary Table 2**)[18], a nearly conservative tracer

of the mixing of hydrothermal fluids with seawater[2]. These results suggest that L_1 ligands along

the MAR are likely not sourced from the vent fluids themselves. All known sources of L_1 ligands

ultimately have a biological origin, and the L_1 ligands observed here could be sourced from

either entrainment of surrounding seawater, production from vent-biota and/or microbial mats,

diffusion from sediments, or *in-situ* production by microbes within the neutrally buoyant

plume[20].

Microbial organic ligand production has been suggested to tightly control the global oceanic

trace metal inventories in both hydrothermal plumes[21,22] and the open ocean[11]. To explore this

tight coupling between Fe and microbially-mediated ligand production within hydrothermal

plumes, we used a novel state-of-the-art liquid chromatography coupled to electrospray

ionization mass spectroscopy[23] method to screen our samples against a custom library of 500

known siderophores. Siderophores have Fe-binding constants that would operationally define

them as L_1 ligands[4,9]. While potentially important in hydrothermal plumes, they have never been

directly measured[14]. From our targeted approach, we identified that at least 0.01-0.66% of the L_1

ligands were microbially produced siderophores with high confidence (**Table 1**).

The siderophores we measured were found in picomolar concentrations, similar to other upper

ocean marine environments[3,4,23,24]. Our estimate is a very conservative lower bound on

siderophore concentrations due to (1) reporting only siderophores that we have identified with

high confidence (40% of the total identifications across all samples, **Fig 2a**, **S3**), (2) detection

limits based on small sample volumes, and (3) the 60-80% extraction efficiency for total

dissolved organic matter[25] of our solid phase extraction technique. It is likely that known

siderophores only represent a small fraction of the total siderophores expected to be produced in

nature[26] and we are likely missing many unknown compounds. Yet, based on these results, the

presence of microbially produced siderophores is evident and suggests several surprising aspects

of Fe cycling in hydrothermal plumes. The biosynthesis of a siderophore is energy-intensive and

is regulated by Fe concentration in the surrounding environment. Thus, the presence of

siderophores suggests that bacteria are producing these compounds despite the overall higher Fe

concentrations in the deep sea, and siderophore production is needed to transform Fe from these

otherwise inaccessible forms. Finally, the direct detection of siderophores in these samples

provides evidence that microbial ligand production is responsible for some portion of the tight

coupling between strong ligands and dFe in hydrothermal systems along the MAR.

Chemical composition of siderophores in hydrothermal systems

Figure 2. Siderophore presence in hydrothermal plumes along the MAR. (a) Heat map of confidence levels 1-2
(blue gradient, 1 = highest confidence). White boxes indicate siderophores that were either not detected or detected
with lower confidence at those sites. The y-axis is ordered from top to bottom in terms of descending mass of the
apo-form (without Fe) of the siderophore. (b) Bar graph of the types of siderophores identified with confidence
levels of 1 and 2. Fe bound versus total summed concentration of (c) all types of siderophores and (d) amphiphilic
siderophores at each station. The vertical red lines separate samples from fracture/diffuse sites on the left and
samples from spreading centers on the right as defined in **Table 1**.

Marine ligand composition has been shown to change with environmental gradients[23], making

the structure and functional groups of siderophores identified in and around hydrothermal

samples of particular interest. Samples collected from on-axis spreading centers contained the

highest dFe concentrations (> 20 nM) and samples from fracture zones, diffuse and off-axis sites

contained the lowest dFe (≤ 1 nM). Spreading centers contained a wider variety of siderophore

types than diffuse/fracture zones, with the greatest number of distinct siderophore compounds

identified at the Broken Spur, Rainbow, and TAG vent sites (**Fig. 2**). On average, 10 compounds

were identified with high confidence per spreading center sample, compared with only 6

identifications per diffuse/fracture zone sample (**Fig. 2b, Fig. S4-5**). Mixed-type siderophores —

siderophores containing multiple moieties that bind to Fe(III) — were most common at or around

spreading centers, while carboxylate-type siderophores — siderophores with both carboxyl and

hydroxyl groups — were the most common at diffuse/fracture zones. Hydroxamates, phenolates,

and catecholates were all identified at and around spreading centers, yet none of these types were

detected in samples from diffuse/fracture zones. At spreading centers, neutrally buoyant plumes

at on-axis vents averaged twice the number of distinct siderophore types than off-axis sites (**Fig.**

S6), suggesting the proximity to the vent field plays an important role in the diversity and

abundance of siderophore types produced by the microbial community.

Low Fe surface waters have been found to contain higher concentrations of amphiphilic

siderophores compared to high Fe waters[23]. Amphiphilic siderophores have long hydrocarbon

tails that can be embedded into the lipid bilayer of the bacterial cell membrane providing a

mechanism to shuttle Fe into the cell and preventing diffusive loss of Fe[27]. As a result, these

microbes have a competitive advantage over other bacteria that produce non-amphiphilic

siderophores which more easily disperse away from the cell in solution. Amphiphilic

siderophores accounted for over a third (38%) of the high confidence siderophores observed in

our samples (**Fig. S7**). These siderophores also tended to have higher molecular weights than

those detected with lower confidence (**Fig. S8**). This either suggests our current methods for

identifying siderophores in environmental samples may select for higher molecular weight

compounds, or that amphiphilic siderophores are very common in marine environments[28]. It is

surprising that they were so common in our dataset, due to the elevated Fe concentrations

observed here relative to more Fe-poor surface ocean. Amphiphilic siderophores were found in

total concentrations between 1.0-18.9 pM, with Rainbow and TAG containing the highest

concentrations (8.9±0.76 and 18.9±1.4 pM respectively; **Fig. 2d**, **Supplementary Table 4**).

Interestingly, these concentrations were similar or higher than previously observed in other upper

ocean marine environments[3,23,29]. Marine bacteria have been shown to produce suites of

amphiphilic siderophores as a way to adapt to the change in hydrophilicity in the surrounding

environment[30,31], which could be an important coping strategy for bacteria living in

environments with strong chemical gradients enabling them to access the varying forms of Fe.

Unlike in the surface ocean where high concentrations of amphiphilic siderophores are observed

in regions where Fe is a growth-limiting micronutrient[23], amphiphilic siderophore production in

hydrothermal plumes could be a way for bacteria store Fe as they are physically transported in

and away from the plume. This could be similar to how the production of multiple siderophores

in terrestrial and pathogenetic systems are used to uptake inorganic Fe for cellular uptake, Fe

storage, and absorption of other metals in association with Fe[26].

Microbial sources of siderophores in hydrothermal plumes

Figure 3. Relative abundance of putative siderophore-producing taxa. Bar height indicates the proportion of
16S rRNA genes recovered in each sample, separated by depth, filter size fraction, and site location. Colors
correspond to taxonomy. Genera found in MAR vent microbial communities with members in the antismash
database predicted to produce siderophores are depicted at the family level.

Microbial production of siderophores is a strategy for organisms to adapt or compete with other
bacteria for Fe within a specific environment[30] and points to a coupling of organic ligand
production and dFe concentrations. We examined the microbial community composition around
Rainbow (St. 11- 200 km E of Rainbow and St. 17- 10 km S of Rainbow) and Lucky Strike (St.
7; **Table 1, Supplementary Table 1**) using 16S rRNA gene-based amplicon sequencing to
determine the presence of microbes that may have the metabolic potential to synthesize the
siderophores we observed at Rainbow, 30 km E of Rainbow and Lucky Strike (**Fig. 3, S9**).
Diagnosing metabolic function from 16S rRNA data alone is limited, however exploring whether
taxa present in these vent sites are encoding siderophore biosynthetic gene clusters indicates
whether the microbial community is genetically capable of producing the compounds we
observed in the environment. Bacterial genera containing known siderophore-producers were
found at all three MAR sites examined, and putative siderophore-producers represented 3-20%
of the relative abundance of each community (**Fig. 3**). Putative siderophore-producers were more

abundant in the 3 µm (particle-attached) size fraction than in the 0.2 µm (free-living) fraction,

suggesting siderophore production is more common in particle-associated bacteria in

hydrothermal environments.

We found microbial genera in our samples that have previously been shown to produce a subset

of the siderophores identified here, including ferrioxamines, vibrioferrin, and

acinetoferrin[3,4,8,10,23]. Genera with the genetic potential to produce ferrioxamines were present at

all three sites, while those known to produce vibrioferrin were present at Lucky Strike and 10 km

S of Rainbow and those producing acinetoferrin present at 10 km S of Rainbow (**Supplemental

Table 1, 5**). Mycobactins were detected with high confidence in every sample of this study and

were isolated from a cultured organism from a hydrothermal system[32], but no putative

mycobactin producers were identified in the 16S rRNA data in this study. We also detected

woodybactin D with high confidence in six out of our eleven sites. Although these biosynthetic

genes were not identified in any of the genera from our 16S rRNA gene amplicon sequences,

woodybactin D is a carboxylate siderophore isolated from *Shewanella*[33], and groups of deep-sea

Shewanella[34] were found in the 16S rRNA data (**Fig. S9**). The biosynthesis genes for many of

the siderophores we identified are unknown, and thus the fact that we only found genera capable

of producing a subset of the siderophores we characterized in the water column is not surprising.

However, the observation that a significant portion of the *in-situ* microbial community is capable

of synthesizing siderophores (**Fig 3**) suggests that siderophore production is more widespread in

the marine environment and in the deep ocean than previously believed.

Although the production of siderophores is often assumed to be associated primarily with low Fe

conditions, evidence that siderophores are ubiquitous in marine environments, including those

relatively rich in Fe, has been increasing. These regions are relatively high in Fe in comparison

to the open ocean, but are depleted relative to terrestrial environments and requirements for

bacteria growth[35]. Thus, the higher dFe associated with hydrothermal plumes may still not be

high enough to suppress siderophore production. It is also likely that in higher Fe environments,

such as hydrothermal plumes, not all of the Fe is bio-accessible. For example, soil microbes

actively secrete siderophores in the high Fe soil environment to solubilize otherwise inaccessible

particulate Fe[36]. Similar processes could be occurring in hydrothermal plumes, where Fe mineral

phases are common[37]. It is telling that most siderophore-producing genera were found in the

particulate size fraction (**Fig. 3**), providing additional evidence that siderophores might be

produced to solubilize different forms of particulate Fe. Further work that assesses why bacteria

are producing siderophores in high Fe neutrally buoyant plumes will be important for

understanding the role of these complexes in facilitating the exchange of Fe between dissolved

and particulate phases[38], and how Fe-siderophore complexes may be stabilizing dFe in

hydrothermal plumes.

Organic Fe-binding ligands have been implicated in playing a critical role in the preservation and

transport of hydrothermal Fe into the ocean interior[2,17,38–43]. In this work, we have identified L_1

ligands as being tightly coupled to dFe in neutrally buoyant hydrothermal plumes along the

MAR, and have provided evidence that these ligands are biologically produced. We identified

specific siderophores that demonstrate a lower bound on the concentrations of L_1 ligands being

produced by the microbial community in response to external Fe inputs. Bacteria putatively

capable of synthesizing the siderophores were largely present in the particulate size fraction

along the MAR, which adds to a growing body of evidence that microbes are likely using L_1

ligands to access particulate or otherwise inaccessible Fe pools and siderophores are a key

mechanism for controlling dFe in neutrally buoyant plumes. Understanding whether tightly

coupled Fe inputs and organic ligand production is present across all hydrothermal vent systems

will aid in constraining the biogeochemical importance of microbial feedbacks in impacting the

hydrothermal dFe supply to the deep ocean.

Methods

Sampling and cruise transect

Samples were collected as part of the 2017-2018 U.K. GEOTRACES GA13 section cruise along

the Mid-Atlantic Ridge. Water samples from 11 venting and near venting locations were

collected using a Seabird 911 conductivity, temperature, and depth (CTD) titanium rosette using

conducting Kevlar wire with an oxidation-reduction potential (ORP) sensor to detect plumes.

Teflon coated OTE (Ocean Test Equipment) bottles were pressurized to approximately 7 psi

with 0.2 µm filtered air using an oil free compressor. A Sartobran 300 (Sartorius) filter capsule

(0.2 µm) was used to collect filtered seawater samples into clean 250 mL LDPE sample

bottles. Bottles and caps were rinsed 3 times with the filtered sample before being filled.

Samples were stored frozen at -20˚C for Fe-organic ligand characterization by voltammetry and

mass spectrometry.

299   Fe-binding ligand concentrations and binding strengths

300   Competitive Ligand Exchange-Adsorptive Cathodic Stripping Voltammetry

Fe-binding ligand concentrations and binding strengths were determined by competitive ligand

exchange-adsorptive cathodic stripping voltammetry (CLE-ACSV) with a BAS*i* controlled

growth mercury electrode (CGME) with an Ag/AgCl$^-$ reference electrode and platinum auxiliary

electrode (Bioanalytical Systems Incorporated). Using previously established methods[2,3,12,44,45],

40 frozen filtrate (<0.2 µm) samples with dFe concentrations between 0.41- 11.67 nM

(**Supplementary Table 1-2**) were thawed in a 4˚C fridge prior to analysis. A 15-point titration

curve was analyzed for each sample. Briefly, within each titration, every point sequentially

received 10 mL of sample, 7.5 µM of borate-ammonium buffer, 10 µM salicylaldoxime (SA)

added ligand, and a dFe addition. Data was collected using the *Epsilon Eclipse Electrochemical*

Analyzer (v.213) with a deposition time of 120 seconds and analyzed using *ElectroChemical*

Data Software (v2001-2014) and *ProMCC* (v2008-2018) to determine peak areas and Fe-binding

ligand parameters, respectively.

314   Reverse Titration-CLE-ACSV

Reverse titration-CLE-ACSV (RT-CLE-ACSV)[46] was completed on 10 samples from Broken

Spur, and TAG hydrothermal vent fields with dFe concentrations between 19.01-90.25 nM

(**Supplementary Table 1-2**). Briefly, a 10-point titration curve was analyzed for each sample

with each titration point consisting of 10 mL of sample buffered with 7.5 µM boric acid and the

competitive ligand 1-nitroso-2-napthol (NN) additions. All samples were analyzed on a BAS*i*

Controlled Growth Mercury Electrode (CGME) with the *Epsilon Eclipse Electrochemical*

Analyzer (v.213) and deposition time of 120 seconds. For each sample, competitive ligand NN

additions were 0.5, 1, 2, 3, 4, 6, 9, 15, 20, and 40 µM. Samples were equilibrated overnight and

purged with N_2 (99.99%) for 5 minutes before analysis. At the end of each titration, three Fe

additions (3-15 nM) were added to the final titration point to get the total concentration of Fe in

equilibrium with ligands. Data was analyzed using *ElectroChemical Data Software* (v2001-

2014) to acquire peak areas and a package in R using the model parameters of $\beta_{FeNN3} = 5.12$ x

10^{16}, $\chi_{min} = 0.8$, $\chi_{max} = 0.9$, and *c1high* = 0.75 to determine the Fe-binding ligand parameters[46].

These parameters were chosen based on the recommendations for undersaturated samples and

titrations curves where ip_{max} was not reached[46]. All other parameters within the model we kept at

the default values.

Fe-binding organic ligand quantification and characterization

In addition to determining the total concentrations of strong Fe-binding ligands (L_1), we also

identified and quantified siderophores that contributed to the L_1 ligand pool. Between 0.65-1.5 L

of 0.2 μm filtered seawater sample (described above) was pumped slowly (15-20 mL min^{-1}) onto

a polystyrene-divinylbenzene (Bond Elut ENV) solid phase extraction (SPE) column[3,23]. SPE

columns were rinsed with MilliQ and stored at -20ºC until analysis. For the analytical

measurements, samples were thawed in the dark, eluted in 12 mL of distilled methanol, and dried

down to between 0.2-0.5 mL (**Supplementary Table 1**). Sample aliquots were analyzed by

reverse-phase liquid chromatography (LC) on a trace metal clean bio-inert LC (Thermo Dionex

3000 NCS). The LC was interfaced with an electrospray ionization-mass spectrometer (ESI-MS;

Thermo Q-Exactive HF) to identify and quantify the compounds based on exact mass (MS^1) and

the fragmentation (MS^2) data[3,23]. A 4-point standard curve with known concentrations of

ferrioxamine E was used to quantify the identified siderophores. MS^1 peak areas were

determined for all identified siderophores and peak areas were converted to concentration using

the peak areas from the desferrioxamine E standard curve (**Fig. S14**). Commercial standards are

not available for all of the siderophores identified, and different compounds have distinct

ionization efficiencies in ESI-MS. Thus, the siderophore concentrations reported here are

estimates of siderophore concentrations in these environments. Additionally, 1 mM of

cyanocobalamin was added as an internal standard to each sample aliquot to address any changes

in sensitivity during LC-MS runs.

MSconvert (Proteowizard) was used to convert ESI-MS data to an open source mzxML format,

and two stages of data processing were conducted using modified versions of previously reported

R scripts[3,23]. In the first stage, mzxML files were read into R using new package "RaMS"

(https://github.com/wkumler/RaMS) which eased analysis, searched for masses of interest from

an in-house database of siderophores in MS^1 data, and identified candidates based on a series of

filters. The masses of interest were the ionized apo, ^{54}Fe-bound, and ^{56}Fe-bound version of each

siderophore, with a tolerance of 7.5 ppm. MS^1 spectra were quality controlled to contain a

minimum of 25 datapoints or maximum intensity of each mass of interest greater than 1000

counts. Spectra meeting these criteria and containing either ^{54}Fe-bound and ^{56}Fe-bound peaks

within 30 seconds of each other or peak for the apo form of interest were displayed for the user

to make the final decision of whether to move on to stage two of processing with a given

siderophore.

Stage two of processing compared MS^2 spectra of the apo and Fe-bound forms of candidate

siderophores with the predicted MS^2 generated by *in silico* fragmenter MetFrag[47]. The *in silico*

fragmenter feature was run with a tolerance of 10 ppm on "$[M+H]^+$" and "$[M+Na]^+$" modes. A

confidence level of 1-4, from highest to lowest confidence, was then assigned to candidates

based on the following criteria: (1) peaks were present in MS^1 and MS^2 spectra, and, and at least

one of the three most-intense peaks match *in silico* fragmentation, (2) peaks were present in MS^1

and MS^2 spectra, and small-intensity peaks in the MS^2 match *in silico* fragmentation, (3) peaks

were present in MS^1 and MS^2 spectra, but little to no fragments in the MS^2 match *in silico*

fragmentation, and (4) good peaks were identified in MS^1 spectra but not in MS^2 spectra (**Fig.**

S10-13). MetFrag pulls chemical structures from publicly-available databases like PubChem or

COCONUT[48], which contain most, but not all variations of siderophores. As such, Fe-bound

candidates were usually run against the apo form available in the database, and for siderophores

with similar structures but variations in fatty chain length or double bond placement, sometimes

only one parent structure was available.

Microbial community analysis

Microbial community composition was assessed in neutrally buoyant plumes and near venting

sites at three sites: Lucky Strike (Station 7; 1670 m), 10 km S of Rainbow (Station 17; 2000 m),

and 200 km E of Rainbow (Station 11; 600 m, 1600 m and 2250 m). One to 2 L of seawater were

filtered by pressure filtration through sequential 25 mm membrane filters housed in

polypropylene filter holders (Whatman SwinLok, GE Healthcare, Pittsburgh, Pennsylvania)

using a peristaltic pump and silicone tubing. Samples first passed through a 3 µm pore-size

polyester membrane filter (Sterlitech, Auburn, Washington) then onto a 0.2 µm pore-size

polyethersulfone membrane filter (Supor-200, Pall Corporation, Port Washington, New York).

Pump tubing was acid washed with 10% hydrochloric acid and flushed with ultrapure water

between each sample. The filters were flash frozen in liquid nitrogen in 2 mL gasketed bead

beating tubes (Fisher Scientific) at sea.

Nucleic acids (DNA) were extracted as described previously[49], with slight modifications. Briefly,

cells on the filters were lysed directly in the bead beating tubes with sucrose-ethylene diamine

tetraacetic acid (EDTA) lysis buffer (0.75 M sucrose, 20 mM EDTA, 400 mM NaCl, 50 mM

Tris) and 1% sodium dodecyl sulfate. Tubes were then agitated in a bead beating machine

(Biospec Products) for 1 min, and subsequently heated for 2 min. at 99°C in a heat block.

Proteinase K (New England Biolabs) was added to a final concentration of 0.5 mg/mL. Filters

were incubated at 55°C for approximately 4 h and the resulting lysates were purified with the

DNeasy kit (Qiagen) using a slightly modified protocol[49]. The purified nucleic acids were eluted

in 200 µL of DNase, RNase-free water, and quantified using a fluorometer (Qubit and Quanti-T

HS reagent, Invitrogen Molecular Probes).

The 16S rRNA gene was amplified in all samples using V4 primers[50,51] (515F-Y and 806RB)

following a previously established protocol[52]. Amplicons were sequenced using a paired-end

250bp run on an Illumina MiSeq 500 and demultiplexed by the UC Davis Genome Center. The

resulting 16S rRNA amplicon sequences were filtered and trimmed using the DADA2 pipeline in

R[53]. Taxonomic assignments were made with version 138.1 of the SILVA SSU database[54]

(silva_nr99_v138.1_wSpecies_train_set.fa.gz ; doi:10.5281/zenodo.4587955; accessed March

2022). Chloroplast and mitochondrial sequences were filtered out of the dataset using the

'phyloseq' R package (v 1.38.0), after which samples had read depths ranging from 9375 –

65486 reads (average 28425 $\pm$ 20014 reads) and represented 1010 unique amplicon sequence

variants (ASVs). Read counts were transformed from absolute to relative abundance and taxa

were aggregated to the Family level. The ten most abundant families present in each sample were

visualized using the 'ggplot2' package (v. 3.3.5).

In order to assess the potential of the observed prokaryotic taxa to produce siderophores, we

downloaded all siderophore biosynthetic gene clusters (BGCs) in the antismash secondary

metabolite database ($n = 7909$) and used text-string matching to compare genera containing these

BGCs to the genera found in our 16S rRNA gene dataset[55]. We cross-referenced the

nomenclature of antismash-predicted siderophores with that of the siderophores identified by

LC-ESI-MS in this study, accounting for minor differences in naming convention between the

two databases, to determine if microbial community members present at each site were predicted

to make any of the siderophores that were measured at that site. Station 38 and Station 12 were

the closest sites with siderophore measurements for comparison against the taxonomic samples

taken at 200 km E of Rainbow and 10 km S of Rainbow, respectively. Samples for microbial

taxonomy and siderophore identity were taken from the same location at Lucky Strike and thus

directly compared.

Data Availability

The CSV data reported in this study has been deposited at Zenodo under the DOI:

http://doi.org/10.5281/zenodo.7325154. The LC-ES-MS data has been deposited on Massive

under the DOI: http://doi.org/doi.10.25345/C5V97ZW7N. Microbial 16S rRNA data have been

deposited on GenBank under the accession number BioProject #PRJNA865382. All data is

freely available on each of these data repositories.

Acknowledgments

We acknowledge the captain and crew of the R/V *James Cook*, Chief Scientist Alessandro

Tagliabue, and Noah Gluschankoff for supporting this work. This study was a part of the

FeRidge project (GEOTRACES section GA13) which was supported by the Natural

Environment Research Council funding (NERC United Kingdom Grants NE/N010396/1 to MCL

and NE/N009525/1 to AT). The International GEOTRACES Programme is possible in part

thanks to the support from the U.S. National Science Foundation (Grant OCE-1840868) to the

Scientific Committee on Oceanic Research (SCOR). CLH was funded by JISAO/CICOES

postdoctoral fellowship. PJM was funded through the NOAA Hollings Scholar summer program.

JR was funded by NOAA Ocean Exploration and Research, NOAA Earth-Ocean Interactions

programs at NOAA-Pacific Marine Environmental Labs, and JISAO/CICOES. Part of this work

was carried out in the University of Washington TraceLab, which receives support from the M.J.

Murdock Charitable Trust in conjunction with the University of Washington College of

Environment, and the Pacific Marine Environmental Labs at the National Oceanic and

Atmospheric Administration. Parts of this work was also carried out in Dr. Anitra Ingalls

laboratory with the help of Laura Truxal and Dr. Jiwoon Park at the University of Washington-

School of Oceanography.

Author Contributions

Manuscript preparation, sample/data processing, CSV analysis, and interpretation (C.L.H.),

manuscript preparation, LC-ESI-MS data analysis and interpretation (P.J.M.), microbial analysis

and interpretation (J.B.A. and A.E.S.), dissolved iron and derived excess $^3He_{xs}$ measurements,

sample collection (A.J.M. L. and M.C.L.), microbial data collection and ligand data

interpretation (T.M. and K.N.B.), and project design and planning, data interpretation, and

mentoring (A.T., M.C.L., J.A.R., and R.M.B.). All authors were involved in editing manuscript.

Competing Interests

The authors declare no competing interests.

References

1. Tagliabue, A. *et al.* The integral role of iron in ocean biogeochemistry. *Nature* **543**, 51–59
(2017).

2. Buck, K. N., Sedwick, P. N., Sohst, B. & Carlson, C. A. Organic complexation of iron in
the eastern tropical South Pacific: Results from US GEOTRACES Eastern Pacific Zonal
Transect (GEOTRACES cruise GP16). *Mar. Chem.* **201**, 229–241 (2018).

3. Bundy, R. M. *et al.* Distinct Siderophores Contribute to Iron Cycling in the Mesopelagic
at Station ALOHA. *Front. Mar. Sci.* 1–15 (2018). doi:10.3389/fmars.2018.00061

4. Moore, L. E., Heller, M. I., Barbeau, K. A., Moffett, J. W. & Bundy, R. M. Organic
complexation of iron by strong ligands and siderophores in the eastern tropical North
Pacific oxygen deficient zone. *Mar. Chem.* **236**, 104021 (2021).

5. Fishwick, M. P. *et al.* Global Biogeochemical Cycles on the dissolution of aerosol iron.

Global Biogeochem. Cycles **28**, 1235–1250 (2014).

6.   Bundy, R. M. *et al.* Iron-binding ligands and humic substances in the San Francisco Bay
estuary and estuarine-influenced shelf regions of coastal California. *Mar. Chem.* **173**,
183–194 (2015).

7.   Buck, K. N., Lohan, M. C., Berger, C. J. M. & Bruland, K. W. Dissolved iron speciation
in two distinct river plumes and an estuary: Implications for riverine iron supply. *Limnol.*
Oceanogr. **52**, 843–855 (2007).

8.   Butler, A. Marine siderophores and microbial iron mobilization. *Biometals* **18**, 369–374
(2005).

9.   Manck, L. E. *et al.* Petrobactin, a siderophore produced by Alteromonas, mediates
community iron acquisition in the global ocean. *ISME J.* **16**, 358–369 (2022).

10.   Vraspir, J. M. & Butler, A. Chemistry of marine ligands and siderophores. *Ann. Rev. Mar.*
Sci. **1**, 43–63 (2009).

11.   Lauderdale, J. M., Braakman, R., Forget, G., Dutkiewicz, S. & Follows, M. J. Microbial
feedbacks optimize ocean iron availability. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 4842–4849
(2020).

12.   Buck, K. N., Sohst, B. & Sedwick, P. N. The organic complexation of dissolved iron
along the U.S. GEOTRACES (GA03) North Atlantic Section. *Deep. Res. Part II Top.*
Stud. Oceanogr. **116**, 152–165 (2015).

13.   Kleint, C., Hawkes, J. A., Sander, S. G. & Koschinsky, A. Voltammetric Investigation of
Hydrothermal Iron Speciation. *Front. Mar. Sci.* **3**, 1–11 (2016).

14.   Li, M. *et al.* Microbial iron uptake as a mechanism for dispersing iron from deep-sea
hydrothermal vents. *Nat. Commun.* **5**, 3192 (2014).

15.   Beaulieu, S. E. & Szafrański, K. M. InterRidge Global Database of Active Submarine
Hydrothermal Vent Fields Version 3.4. (2020). doi:10.1594/PANGAEA.917894

16.   Omanović, D., Garnier, C. & Pižeta, I. ProMCC: An all-in-one tool for trace metal
complexation studies. *Mar. Chem.* **173**, 25–39 (2015).

17.   Resing, J. a. *et al.* Basin-scale transport of hydrothermal dissolved metals across the South
Pacific Ocean. *Nature* **523**, 200–203 (2015).

18.   Lough, A. J. M. *et al.* The impact of hydrothermal vent geochemistry on the addition of
iron to the deep ocean. *Biogeosciences Discuss.* **[preprint]**, 1–23 (2022).

19.   Fishwick, M. P. *et al.* The impact of changing surface ocean conditions on the dissolution
of aerosol iron. *Global Biogeochem. Cycles* **28**, 1235–1250 (2014).

20.   Mellett, T. *et al.* Particle exchange mediated by organic ligands in incubation experiments
of hydrothermal vent plumes along the mid-Atlantic Ridge.

21.   Cowen, J. P. & Bruland, K. W. Metal deposits associated with bacteria: implications for
Fe and Mn marine biogeochemistry. *Deep Sea Res. Part A. Oceanogr. Res. Pap.* **32**, 253–

272 (1985).

22. Cowen, J. P., Massoth, G. J. & Feely, R. A. Scavenging rates of dissolved manganese in a
hydrothermal vent plume. *Deep Sea Res. Part A. Oceanogr. Res. Pap.* **37**, 1619–1637
(1990).

23. Boiteau, R. M. *et al.* Siderophore-based microbial adaptations to iron scarcity across the
eastern Pacific Ocean. *Proc. Natl. Acad. Sci.* **113**, 14237–14242 (2016).

24. Park, J. *et al.* Siderophore production and utilization by microbes in the North Pacific
Ocean. *bioRxiv* 2022.02.26.482025 (2022). doi:10.1101/2022.02.26.482025

25. Waska, H., Koschinsky, A., Ruiz Chancho, M. J. & Dittmar, T. Investigating the potential
of solid-phase extraction and Fourier-transform ion cyclotron resonance mass
spectrometry (FT-ICR-MS) for the isolation and identification of dissolved metal-organic
complexes from natural waters. *Mar. Chem.* **173**, 78–92 (2015).

26. Hider, R. C. & Kong, X. Chemistry and biology of siderophores. *Nat. Prod. Rep.* **27**, 637–
657 (2010).

27. Martinez, J. S. *et al.* Structure and membrane affinity of a suite of amphiphilic
siderophores produced by a marine bacterium. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 3754–
3759 (2003).

28. Butler, A. & Theisen, R. M. Iron(III)-siderophore coordination chemistry: Reactivity of
marine siderophores. *Coord. Chem. Rev.* **254**, 288–296 (2010).

29. Boiteau, R. M. *et al.* Patterns of iron and siderophore distributions across the California
Current System. *Limnol. Oceanogr.* **64**, 376–389 (2019).

30. Sandy, M. & Butler, A. Microbial iron acquisition: marine and terrestrial siderophores.
Chem. Rev. **109**, 4580–95 (2009).

31. Homann, V. V *et al.* Loihichelins A - F , a Suite of Amphiphilic Siderophores Produced
by the Marine Bacterium Halomonas LOB-5. *J. Nat. Prod.* **72**, 884–888 (2009).

32. Gu, H., Sun, Q., Luo, J., Zhang, J. & Sun, L. A First Study of the Virulence Potential of a
Bacillus subtilis Isolate From Deep-Sea Hydrothermal Vent. *Front. Cell. Infect.*
Microbiol. **9**, 1–14 (2019).

33. Carmichael, J. R., Zhou, H. & Butler, A. A suite of asymmetric citrate siderophores
isolated from a marine Shewanella species. *J. Inorg. Biochem.* **198**, 1–6 (2019).

34. Kato, C. & Nogi, Y. Correlation between phylogenetic structure and function : examples
from deep-sea Shewanella. **35**, 223–230 (2001).

35. Tortell, P. D., Maldonado, M. T. & Price, N. M. The role of heterotrophic bacteria in iron-
limited ocean ecosystems. *Nature* **383**, 330–332 (1996).

36. Crowley, D. E., Wang, Y. C., Reid, C. P. P. & Szaniszlo, P. J. Mechanisms of iron
acquisition from siderophores by microorganisms and plants. *Plant Soil* **130**, 179–198
(1991).

37. Hoffman, C. L. *et al.* Diagnostic morphology and solid-state chemical speciation of
hydrothermally derived particulate Fe in a long-range dispersing plume. *ACS Earth Sp.*
Chem. **4**, 1831–1842 (2020).

38. Fitzsimmons, J. N. *et al.* Iron persistence in the distal hydrothermal plume supported by
dissolved – particulate exchange. *Nat. Geosci.* **10**, 1–8 (2017).

39. Hoffman, C. L. *et al.* Near-field iron and carbon chemistry of non-buoyant hydrothermal
plume particles, Southern East Pacific Rise 15°S. *Mar. Chem.* **201**, 183–197 (2018).

40. Toner, B. M. *et al.* Preservation of iron(II) by carbon-rich matrices in a hydrothermal
plume. *Nat. Geosci.* **2**, 197–201 (2009).

41. Bennett, S. a., Hansman, R. L., Sessions, A. L., Nakamura, K. ichi & Edwards, K. J.
Tracing iron-fueled microbial carbon production within the hydrothermal plume at the
Loihi seamount. *Geochim. Cosmochim. Acta* **75**, 5526–5539 (2011).

42. Bennett, S. a. *et al.* The distribution and stabilisation of dissolved Fe in deep-sea
hydrothermal plumes. *Earth Planet. Sci. Lett.* **270**, 157–167 (2008).

43. Sander, S. G. & Koschinsky, A. Metal flux from hydrothermal vents increased by organic
complexation. *Nat. Geosci.* **4**, 145–150 (2011).

44. Abualhaija, M. M. & van den Berg, C. M. G. G. Chemical speciation of iron in seawater
using catalytic cathodic stripping voltammetry with ligand competition against
salicylaldoxime. *Mar. Chem.* **164**, 60–74 (2014).

45. Hawkes, J. A., Connelly, D. P., Gledhill, M. & Achterberg, E. P. The stabilisation and
transportation of dissolved iron from high temperature hydrothermal vent systems. *Earth*
Planet. Sci. Lett. **375**, 280–290 (2013).

46. Hawkes, J. A., Gledhill, M., Connelly, D. P. & Achterberg, E. P. Characterisation of iron
binding ligands in seawater by reverse titration. *Anal. Chim. Acta* **766**, 53–60 (2013).

47. Ruttkies, C., Schymanski, E. L., Wolf, S., Hollender, J. & Neumann, S. MetFrag
relaunched: incorporating strategies beyond in silico fragmentation. *J. Cheminform.* **8**, 1–
16 (2016).

48. Sorokina, M., Merseburger, P., Rajan, K., Yirik, M. A. & Steinbeck, C. COCONUT
online: Collection of Open Natural Products database. *J. Cheminform.* **13**, 1–13 (2021).

49. Santoro, A. E., Casciotti, K. L. & Francis, C. A. Activity, abundance and diversity of
nitrifying archaea and bacteria in the central California Current. *Environ. Microbiol.* **12**,
1989–2006 (2010).

50. Apprill, A., Mcnally, S., Parsons, R. & Weber, L. Minor revision to V4 region SSU rRNA
806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquat. Microb.*
Ecol. **75**, 129–137 (2015).

51. Parada, A. E., Needham, D. M. & Fuhrman, J. A. Every base matters: Assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series and
global field samples. *Environ. Microbiol.* **18**, 1403–1414 (2016).

52.   Stephens, B. M. *et al.* Organic Matter Composition at Ocean Station Papa Affects Its
592   Bioavailability, Bacterioplankton Growth Efficiency and the Responding Taxa. *Front.*
593   *Mar. Sci.* **7**, (2020).

594   53.   Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon
595   data. *Nat. Methods* **13**, 581–583 (2016).

596   54.   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: Improved data
597   processing and web-based tools. *Nucleic Acids Res.* **41**, 590–596 (2013).

598   55.   Blin, K., Shaw, S., Kautsar, S. A., Medema, M. H. & Weber, T. The antiSMASH database
599   version 3: Increased taxonomic coverage and new query features for modular enzymes.
600   *Nucleic Acids Res.* **49**, D639–D643 (2021).

601   56.   Bazylev, B. A. Allochemical Metamorphism of Mantle Peridoties in the Hayes Fracture
602   Zone of the North Atlantic. *Petrology* **5**, 362–379 (1997).

603   57.   Kelley, D. S. & Shank, T. M. Hydrothermal systems: A decade of discovery in slow
604   spreading environments. *Geophys. Monogr. Ser.* **188**, 369–407 (2010).

Table 1. Characteristics of sample locations along the Mid Atlantic Ridge

Vent Names	Abbr.	Station	Geology	Host rock	Vent type	Spreading rate (mm/yr)	Siderophore concentration (pM)	Siderophore/L_1 ligand (%)*
Lucky Strike	LS	7/8	Spreading Center	gabbro	Black smoker	20.2	0.4-1.3	0.001±5.23
33 km E of Rainbow	CER	12	Spreading Center	-	-	-	0.5-2.4	0.02±0.27
Rainbow	R	38	Spreading Center	ultramafic	Black smoker	20.6	0.5-17.5	*n.a.*
Hayes Fracture Zone	HFZ	21	Fracture Zone	peridotites/gabbro	-	21.2	0.7-33.7	0.66±0.35
Lost City	LC	23	Fracture Zone	ultramafic/gabbro	Alkaline	22.6	0.4-3.7	0.12±0.08
Broken Spur	BS	24	Spreading Center	gabbro	Black smoker/diffuse	22.9	0.1-10.4	0.03±2.30
29 km N of TAG	CNT	26	Spreading Center	-	-	-	0.6-4.0	0.06±1.75
30 km W of TAG	CWT	30	Spreading Center	-	-	-	0.4-2.8	0.09±0.001
30 km E of TAG	CET	31	Spreading Center	-	-	-	0.3-1.2	0.11±0.0002
Trans-Atlantic Geotraverse	TAG	35	Spreading Center	gabbro	Black smoker	23.6	0.2-8.5	0.05±0.0004
Low Temp Slope	LTS	37	-	-	Diffuse fluids	-	0.3-1.6	0.06±0.03

Spreading rates along the Mid-Atlantic Ridge were gathered from the Interridge Database v3.4. Host rock groups were determined from previously discussed classifications[56,57]. Off-axis sites –33 km E of Rainbow, 29 km N of TAG, 30 km E of TAG, and 30 km W of TAG– were far-field locations of their respective vent field. Low Temp Slope was a diffuse-dominated site that was sampled for the first time as a part of this study.

Siderophore concentrations and the percent of L_1 ligand are reported for both confidence level 1 and 2. Average L_1 ligand and siderophore concentrations can be viewed in **Supplementary Table 3** and concentrations for individual siderophores can be observed in **Supplementary Table 4**.

*Standard deviations are reported to understand the spread of both the average siderophore and L_1 ligand data.
n.a.= unable to be determined
- = unknown

613