A high-throughput platform for single-molecule tracking identifies drug interaction and

cellular mechanisms

David T. McSwiggen, Helen Liu, Ruensern Tan, Sebastia Agramunt Puig, Lakshmi B. Akella,

Russell Berman, Mason Bretan, Hanzhe Chen, Xavier Darzacq, Kelsey Ford, Ruth Godbey, Eric

Gonzalez, Adi Hanuka, Alec Heckert, Jaclyn J. Ho, Stephanie L. Johnson, Reed Kelso, Aaron

Klammer, Ruchira Krishnamurthy, Jifu Li, Kevin Lin, Brian Margolin, Patrick McNamara,

Laurence Meyer, Sarah E. Pierce, Akshay Sule, Connor Stashko, Yangzhong Tang, Daniel J.

Anderson, Hilary P. Beck

Abstract

The regulation of cell physiology depends largely upon interactions of functionally distinct

proteins and cellular components. These interactions may be transient or long-lived, but often

affect protein motion. Measurement of protein dynamics within a cellular environment,

particularly while perturbing protein function with small molecules, may enable dissection of

key interactions and facilitate drug discovery; however, current approaches are limited by

throughput with respect to data acquisition and analysis. As a result, studies using super-

resolution imaging are typically drawing conclusions from tens of cells and a few experimental

conditions tested. We addressed these limitations by developing a high-throughput single-

molecule tracking (htSMT) platform for pharmacologic dissection of protein dynamics in living

cells at an unprecedented scale (capable of imaging $> 10^6$ cells/day and screening $> 10^4$

compounds). We applied htSMT to measure the cellular dynamics of fluorescently tagged

estrogen receptor (ER) and screened a diverse library to identify small molecules that perturbed

ER function in real time. With this one experimental modality, we determined the potency,

pathway selectivity, target engagement, and mechanism of action for identified hits. Kinetic

htSMT experiments were capable of distinguishing between on-target and on-pathway

modulators of ER signaling. Integrated pathway analysis recapitulated the network of known ER

interaction partners and suggested potentially novel, kinase-mediated regulatory mechanisms.

The sensitivity of htSMT revealed a new correlation between ER dynamics and the ability of ER

antagonists to suppress cancer cell growth. Therefore, measuring protein motion at scale is a

powerful method to investigate dynamic interactions among proteins and may facilitate the

identification and characterization of novel therapeutics.

Introduction

Both the activity and mobility of proteins within the crowded cellular environment are

profoundly influenced by interactions with their surroundings[1–3]. Under these conditions, where

diffusion is no longer well described by the Stokes radius of the protein monomer[4], changes in

protein motion might be expected to correlate closely with changes in the activity of these

proteins. The development of increasingly sophisticated live-cell microscopy techniques,

including early ensemble methods like fluorescence recovery after photobleaching (FRAP) and

fluctuation correlation spectroscopy (FCS), have informed our understanding of protein

dynamics in cellular biology[5]. A myriad of technical improvements, such as enhanced labeling

methods[6,7], better live-cell compatible fluorophores[8], new forms of light microscopy[9], dramatic

increases in computational power[10], and the addition of machine learning approaches to data

analysis[10,11] have together enabled a new era of imaging-based studies across biological

contexts[3,5,12].

Applying any microscopy technique at scale presents challenges, but recent advances have

shown the power of high content imaging techniques to address both mechanistic biological

questions as well as to generate leads for new chemical matter in drug discovery[13]. Even these

conceptually simple experiments involving fixed cells stained with well-characterized

commercial reagents take careful experiment design and sophisticated computational approaches

to execute[14]. It is no wonder, then, that attempts to combine high content imaging workflows

with more advanced super-resolution microscopy methods have thus far been limited. Such

advances have enabled the development of systems for fixed-cell STORM imaging at an

impressive 10,000 cells per day[15], though the appropriate application of this increase in scale

remains an open question.

In single molecule tracking (SMT), a fluorescently labeled protein of interest is imaged at

high spatiotemporal resolution to track its motion in a live cell[10]. The information embedded in

these trajectories has been used to investigate diverse cellular phenomena including protein

oligomerization state and function[15–17], inter-organelle communication[18], nuclear organization[19],

and transcription regulation [20–23]. Of particular utility are "fast-SMT" approaches which use high

frame rates and stroboscopic illumination to minimize motion-induced blurring, and hence can

measure diffusive states over a large dynamic range[24,25]. Specifically, proteins that diffuse rapidly

throughout the cell are often missed in alternative tracking approaches, biasing the resulting data.

In spite of the potential biological discoveries that depend on the application of SMT on a large

scale, SMT in general (and fast-SMT in particular) has not been adapted to a high-throughput

setting that would enable the analysis of complex, multi-component systems, or the identification

of compounds that affect protein motion.

Steroid hormone receptors (SHRs) are a class of transcription factors that play crucial roles in

normal human development and in disease pathogenesis. SHRs like the estrogen receptor (genes

ESR1 and ESR2), androgen receptor (AR), and progesterone receptor (PR), as examples,

contribute decisively to the acquisition of secondary sex characteristics, while the glucocorticoid

receptor (GR) helps to orchestrate both metabolism and inflammation[26]. In their ligand-free state,

SHRs are kept sequestered in multiprotein complexes by the chaperone HSP90[27]. Canonically, in

the presence of hormone they dimerize and bind their cognate genomic response elements,

recruiting epigenetic modifiers and transcription machinery[26–28]. At the same time, steroid

hormone receptor-derived signals impose a large disease burden by promoting the growth of

breast cancers (ER)[26,29] or prostate cancers (AR)[26]; or by imposing immune and metabolic

dysfunction (GR)[26,30]. SHRs therefore provide an excellent proof-of-concept system to study the

relationship between protein dynamics and protein function due to the wealth of information and

reagents[29] already available for these systems as well as previous reports characterizing some

aspects of their cellular dynamics [20,22,31–33].

Here, we present the first industrial-scale, high-throughput, fast-SMT (htSMT) platform

capable of measuring protein motion from more than 13,000 individual assay wells (>1,000,000

individual cells) per day. Using ER as a test system, we demonstrate that chemical screening

using htSMT is specific, robust, and reproducible. The increase in throughput enables classical

drug discovery activities, including compound library screening and the elucidation of structure-

activity relationships (SAR), yielding accurate and reproducible results that are inaccessible or

unmeasurable with other techniques or using SMT on a smaller scale. Importantly, we

demonstrate that htSMT can be used to characterize both known and novel pathway contributions

to the ER protein interaction network. More than a proof-of-concept for the htSMT platform,

these data confirm that analysis of protein motion itself on a large scale reveals detailed

information about pathway interactions and signaling.

Results

Creation and validation of a high throughput SMT platform

We developed a robotic system capable of handling reagents, collecting high-quality fast

SMT image series, processing time-ordered raw images to yield molecular trajectories, and

extracting features of biological interest within defined cellular compartments (Figure 1 – figure

supplement 1A, B). Samples start as cells seeded into 384-well plates in a hotel incubator. A

central robotic arm retrieves the plates, and delivers them to an Echo 650 acoustic dispenser to

add dye. After incubating, excess dye is washed away and Echo 650 is again used to administer

compound treatment. Stained and compound-treated plates are then delivered to any of up to four

identical SMT microscopes for imaging. Both SMT and accompanying Hoechst images are

collected and automatically processed to identify individual molecule positions, reconnect the

spot coordinates into trajectories, and then associate each trajectory with a nuclear mask. Finally

the processed SMT data are subjected to quality control to omit aberrant fields of view using a

convolutional neural network trained to identify technical errors in the images (Figure 1 – figure

supplement 1C), and finally stored for downstream analysis.

To examine htSMT system performance across a broad spectrum of diffusion coefficients, we

generated three U2OS cell lines ectopically expressing HaloTag[6] fused proteins with well-

established behaviors in the cell. These HaloTag fusions allow the subsequent addition of bright

and photostable organic fluorophores like JF_{549}[8] which produce high signal spots to detect and

track. Histone H2B-Halo, which is predominantly incorporated into chromatin and therefore

effectively immobile over short timescales[25], was employed to estimate localization error. A

prenylation motif (Halo-CaaX) embedded in the plasma membrane exhibits moderate diffusion[34].

Unfused HaloTag was chosen to represent the upper limit of cellular "free" diffusion. Single-

molecule trajectories measured in these cell lines yielded diffusion coefficients for CaaX similar

to published results[34], and the diffusion for H2B-HaloTag was consistent with the theoretical

lower bounds that can be approximated from the localization error and 10 msec frame interval

$(D_{apparent} = D_{true} + \frac{localization\ error^2}{\Delta\tau})$ (Figure 1A). Localization error can be measured

directly from the single molecule trajectories using the jump covariance of slow or immobile

particles[35]. Using the immobile H2B-Halo trajectories, we found the localization error of the

htSMT system to be 39 nm (Figure 1 – figure supplement 2A), comparable to other benchmark

stroboscopic illumination datasets[25,35]. The diffusion coefficient for free HaloTag is consistent

with previous SMT reports[25], but is also within the theoretical upper bounds of an MSD

estimator of diffusion coefficient for a 10 msec frame interval and 1.25 µm search radius

$(D_{apparent} \leq \frac{R^2_{search}}{8\Delta\tau})$, thus we consider the distribution recovered from the free HaloTag to

represent the upper limit of trackable particles with this assay configuration.

We then tested whether our htSMT platform can extract accurate molecular trajectories at

scale. We employed 384-well plates where free Halo, Halo-CaaX, and H2B-Halo cell lines were

mixed in equal proportions in each well. Imaging with a 94 µm by 94 µm field-of-view (FOV),

we achieved an average of 10 nuclei simultaneously (Figure 1B, Figure 1 – figure supplement

2B, Supplemental video 1), enough that most FOVs contained cells from each cell line. To limit

ambiguity in cell assignment, we considered only the trajectories that fell within nuclear

segmented regions. The probability distribution of diffusion states cleanly distinguishes between

the three cell types (Figure 1C). More importantly, by looking at the single-cell state distribution

profiles of 103,757 cells from five separate 384-well plates, grouped by their distribution profile,

we recovered highly consistent estimates of protein dynamics at the single cell level, comparable

to the pure populations (Figure 1D, Figure 1 – figure supplement 2C).

While single-cell measurements are powerful, the number of trajectories in one cell is

limited, and so estimates of diffusive states can be broad. Combining trajectories from multiple

cells, however, provides the expected distribution of diffusive states (Figure 1E). Moreover,

combining trajectories derived from many cells makes model-based[25] or model-agnostic[35] state

analysis possible, where as few as 10^3 trajectories permit satisfactory inference of the underlying

diffusion states. We determined that imaging six fields of view (FOV), for 1.5 seconds each,

yielded enough trajectories ($> 10,000$) to accurately estimate protein dynamics, bringing the

overall throughput of the platform to more than 13,000 individual wells ($>90,000$ FOVs; $>$

1,000,000 cells/day), a rate of data acquisition that enables compound screening on a feasible

timescale (Figure 1 – figure supplement 2D).

Using htSMT to measure protein dynamics of SHRs

Equipped with an htSMT system capable of measuring protein dynamics broadly, we sought

to understand how measuring protein motion can be used to characterize protein activity. SHRs

transition between inactive and active states via ligand binding (Figure 2 – figure supplement

2A), a phenomenon that has been previously observed at the single molecule level[20,22], and we

hypothesized that the large dynamic range and orders-of-magnitude increase in throughput of our

platform could capture these differences in the context of compound screening. We generated

HaloTag fusions to ER, AR, and PR through ectopic expression, and GR through endogenous

knock-in. Similar to previous approaches[20,36], we used clonal cell lines in a U2OS cell

background to minimize effects of comparing dynamics in different cell types. Clones were

carefully selected such that the HaloTag fusion SHRs were comparable to each other in transcript

abundance, and not higher than transcript levels in tissue-specific cell lines like MCF7 and T47d,

which are both ER and PR positive (Figure 2 – figure supplement 2B).

In the absence of hormone, all four SHR proteins exhibit similar dynamic profiles: a small

immobile fraction and a large freely diffusing fraction with a $3.4 - 4.3$ µm^2/sec average diffusion

coefficient (Figure 2A, Figure 2 – figure supplement 2C). No correlation between diffusion and

protein molecular weight (138 kDa for Halo-AR, 102 kDa for ER-Halo, 122 kDa for Halo-GR,

and 135 kDa for Halo-PR) was observed, highlighting the differences between cellular protein

dynamics versus purified systems. Upon addition of agonist, a dramatic increase in immobile

trajectories is observed, which we attribute to chromatin binding. Using a conservative upper-

bound of chromatin mobility in the nucleus and chromatin-associated transcription factors[35], we

define the bound fraction (f_{bound}) for each SHR as the fraction of trajectories diffusing less than

0.1 µm^2/sec (Figure 2A). Using this threshold, f_{bound} of histone H2B is 0.92 on average,

consistent with previous reports[25]. D_{free} we defined as the occupation-weighted average diffusion

coefficient of the non-bound states (Figure 2 – figure supplement 2C). Some SHRs had a higher

proportion of bound molecules than others; The ligand-induced effect is most pronounced for

ER, with 34% bound in basal conditions and 87% bound after estradiol treatment (Figure 2A,

Supplemental video 2 and 3).

SHRs are highly selective for their cognate agonists in biochemical binding assays, which we

confirmed by measuring the dose-dependent change in dynamics as a function of agonist

concentration. The maximal change in f_{bound} (Figure 2B) and decrease in D_{free} (Figure 2 – figure

supplement 2D) differed between SHRs. The dose titration curves also showed variable potencies

(EC_{50}) for each SHR/hormone pair, with ER-estradiol being both the most potent and most

selective pair. RNA-seq after estradiol stimulation showed a marked induction of hallmark ER-

dependent gene sets[37], confirming that the increase in chromatin binding we observed by SMT

has a functional effect in promoting ER-responsive gene programs, even in the ectopic

expression setting (Figure 2 – figure supplement 2E, F). Thus, SMT can detect functionally-

relevant changes in transcription factor dynamics and accurately differentiate the ligand/target

specificity directly within the cellular environment.

Screening a diverse bioactive chemical set identifies known and novel modulators of ER

dynamics

Our characterization efforts of ligand selectivity for AR, ER, GR, and PR collectively

suggested that we could use SMT to interrogate the effects of compounds on protein dynamics at

a throughput conducive to high throughput screening. We first identified a structurally diverse set

of 5,067 molecules with heterogeneous biological activities as a useful screening set. Having

determined that the same data acquisition parameters (6 FOVs imaged for 1.5 sec each) were

sufficient to recover more than 10,000 trajectories per well, we could achieve a throughput of

15,000 wells per day which would permit us to interrogate the whole library in a single day

(Figure 3 – figure supplement 1A). For cells at steady state, such a brief acquisition time also

means that the dynamical state of the cell remains largely constant within an FOV (Figure 3 –

figure supplement 1B), allowing all of the trajectories from an FOV to be considered largely

representative of the same cellular state. We chose to screen this bioactive compound set against

ER, assessing change in f_{bound} at 1 µM compound versus the vehicle DMSO (Figure 3, Figure 3 –

figure supplement 1C, D, Figure 3 – supplemental table 1). The screen was run twice to assess

reproducibility, showing a high degree of agreement between replicates for ER-active molecules

(Figure 3 – figure supplement 2A). This screen illustrates some important advantages of our

htSMT platform over more manual lower-throughput approaches.

From plate to plate, the assay window for the screen was robust[38] (average Z'-factor = 0.79

over 72 plates), the measured potency of the control estradiol in each instance remained within

three-fold of the mean (Figure 3 – figure supplement 2B-D), and the distribution of negative

control wells centered tightly on zero (Figure 3 – figure supplement 2E, F). Each compound

measurement was averaged from SMT trajectories of between 94 and 161 cells (25^{th} to 75^{th}

percentiles; Figure 3 – figure supplement 2G). Of the 30 compounds we identified from the

bioactive set that we expected to modulate ER, either as agonists or antagonists[29,39], all

significantly increased f_{bound} measured by SMT. This includes agonistic molecules like estradiol,

but also notable examples such as 4-hydroxytamoxifen, fulvestrant, and bazedoxifene (Figure 3,

Extended Data 3B, H).

With a large dataset to work from, collected across multiple assay plates on multiple

independent microscopes, we examined the sources of variability within the screening assay.

Using random sampling of individual jumps within the screening dataset while holding constant

the source of the jumps (for example, sampling jumps within a specific assay well), we could

estimate the relative contributions of microscope-to-microscope, plate-to-plate, well-to-well,

FOV-to-FOV and cell-to-cell variability. Cell-to-cell variability was the single largest contributor

to overall assay variability, especially when considering only vehicle treated controls (Figure 3 –

figure supplement 3B), and stabilized after ~1000 jumps (Figure 3 – figure supplement 3C).

These results support the use of a short acquisition time per FOV and multiple FOVs to stabilize

the dynamical state estimate.

The somewhat counter-intuitive finding that either strong agonism or antagonism can lead to

an increase in chromatin binding has been reported for ER[31], but this appears not to a general

feature of SHRs. While the PR antagonist mifepristone[40] behaves similarly to ER (Figure 3 –

figure supplement 4A), antagonists of AR like Enzalutamide and Darolutamide[41], and antagonists

of GR like AL082D06[42] cause a decrease in chromatin binding. This decrease occurs when

administered singly or when co-administered in competition with the cognate agonist (Figure 3 –

figure supplement 4B, C). These results show how the cellular context and interaction partners

are critical to understand the effect of a compound on its intended target. To underscore this

point, in addition to binders of the ER ligand-binding domain, we also identified a number of

active compounds targeting diverse nodes in the ER interaction network, including modulators of

the proteasome, chaperones, kinases, and others (Figure 3).

Cellular ER dynamics elucidate structure-activity relationships (SAR) of ER

modulators

Our screen revealed that, surprisingly, all the known ER modulators—both agonists like

estradiol and potent antagonists like fulvestrant—caused an increase in f_{bound} (Figure 4A). We

therefore characterized a subset of selective ER modulators (SERMs) and selective ER degraders

(SERDs) in more detail. These molecules all bind competitively to the ER ligand binding

domain[29]. As in the bioactive screen, both SERDs and SERMS increased f_{bound} (Figure 4A) and

slightly decreased measured D_{free} (Figure 4 – figure supplement 1A), with potencies ranging

from 9 pM for GDC-0927 to 4.8 nM for GDC-0810 (Figure 4B). To understand how quickly

these compound effects take place, we measured the change in f_{bound} as a function of time,

collecting timepoints roughly every 2 minutes beginning immediately after compound addition.

Despite different physical-chemical properties, all five increased f_{bound} within minutes of

compound addition (Figure 4C, Figure 4 – figure supplement 1B), with no evidence of transient

states distinct from the free diffusion and chromatin bound peaks (Figure 4 – figure supplement

1C). Presumably, ER dissociation from the chaperone complex, dimerization, and chromatin

binding occur on rapid and seemingly comparable timescales. Since we cannot distinguish

individual steps in these transitions, we consider the on rate of the entire process to have the

effective rate constant k^*_{on}. Importantly, selective antagonists of AR and GR did not induce

significant modulation of ER dynamics, further highlighting the utility of htSMT in

characterizing the specificity of interactions between small molecule modulators of protein

function and their cognate targets (Figure 4 – figure supplement 1D).

Interestingly, SERMs 4-hydroxytamoxifen (4OHT) and GDC-0810 show lower maximal

increases in f_{bound} compared with the SERDs fulvestrant and GDC-0927 (Figure 4D). Similar

effects have been described previously using fluorescence recovery after photobleaching[31]

(FRAP), which we confirmed using our Halo-ER cell line (Figure 4E). The delay in ER signal

recovery after two minutes in FRAP was consistent with the changes in f_{bound} measured by SMT

(Figure 4F). Although FRAP was used to measure these f_{bound} differences, the technique suffers

from challenges in scalability and depends heavily on prior assumption of the underlying

dynamical model in the sample. SMT, alternatively, enables detailed characterization of the

potency of 4OHT and GDC-0810 relative to other ER ligands in their ability to increase ER

chromatin binding on a tractable timescale.

Neither FRAP nor htSMT can discriminate between recovery driven by an increase in

residence time (decreasing k^*_{off}) or increasing the rate of chromatin binding (increasing k^*_{on}),

either of which would result in increasing f_{bound}. By changing SMT acquisition conditions to

reduce the illumination intensity and collect long, 250 msec continuous frame exposures, only

immobile proteins form spots. Under these imaging conditions, the distribution of track lengths

provides a measure of relative residence times[20,21,43,44]. Both agonist and antagonist treatment led

to longer binding times compared to DMSO, suggesting that ligand binding decreases k^*_{off}

(Figure 4G, Figure 4 – figure supplement 1E). Consistent with FRAP, estradiol, GDC-0927, and

fulvestrant show longer binding times compared with other ER modulators. Using f_{bound} and k^*_{off}

measurements, one can infer the k^*_{on}. In all cases the changes in dissociation rate are not

proportional to the increase in f_{bound}, and so ligand-imposed increases in k^*_{on} likely contribute to

the observed change in the chromatin associated ER fraction (Figure 4 – figure supplement 1F).

These data are consistent with a model wherein ER rapidly binds to chromatin irrespective of

which molecule occupies the ligand binding domain, but some ligands induce a conformation

that can be further stabilized on chromatin by cofactors. Consequently, these data support the

hypothesis that ER may engage chromatin in mechanistically different ways[31]. An efficacious ER

inhibitor may promote rapid and transient chromatin binding that fails to effectively recruit

necessary cofactors to drive transcription[33].

htSMT reveals a relationship between ER dynamics and efficacy in ER-dependent cell

toxicity

As the name implies, next-generation ER degraders like GDC-0927, AZD9833, and

GDC9545 were optimized to enhance degradation of ER[31,45]. We indeed observed compound-

induced ER degradation via immunofluorescence, both in established breast cancer model lines

and our U2OS ectopic expression system (Figure 5 – figure supplement 1A, B). Structural

analogs of GDC0927 have been reported and optimized for ER degradation, however the

correlation between ER degradation and cell proliferation is poor[29,46] (Figure 5A, Figure 5 –

figure supplement 1C-E). We hypothesized that by measuring protein dynamics we might obtain

more precise measurements of inhibitory activity than can be achieved by assessing protein

degradation. We therefore determined the potency and maximal effect of structural analogues of

GDC-0927 using htSMT.

Overall, these analogues exhibited a potency range of 15 pM to 12 nM and increased ER f_{bound} by

0.4 to 0.56 (Figure 5B). Small changes in the chemical structure produced measurable changes in

both compound potency and maximal efficacy as determined using SMT, a critical feature if an

assay is to be used to iteratively optimize a compound for potency or efficacy.

We compared the potencies of GDC-0927 and analogues determined either via ER

degradation or SMT, with the ability of each of these compounds to block estrogen-induced

breast cancer cell proliferation. Potency assessed by ER degradation was not a good predictor of

potency in the cell proliferation assay (Figure 5A). By contrast, SMT measurements of f_{bound}

strongly correlate with cell viability (Figure 5C; R^2 of 0.83 for T47d and 0.84 for MCF7).

Intriguingly, SMT EC_{50} values were on average 10-fold lower than those observed in the cell

growth assay, suggesting that SMT may be sensitive enough to identify chemical series that

would not show effects in other cellular assays, enabling the identification of starting points for

medicinal chemistry that could not be obtained by other methods. This correlation between

effects on protein dynamics (f_{bound}) and protein function (suppression of cell proliferation)

coupled with the throughput of the SMT system make this an attractive approach for the

identification of protein modulators with novel properties.

Screening of a diverse chemical library using htSMT enables unbiased pathway

interaction analysis by monitoring protein dynamics

In addition to known ER active modulators, many other compounds in our bioactive library

provoked easily measurable changes in f_{bound}. To define a threshold for calling a molecule from

the screen "active," we selected 92 compounds with different magnitudes of change in f_{bound} to

retest in a dose titration (Figure 6 – figure supplement 1A, B). We found that a 5% change in

f_{bound} was sufficient to reproducibly distinguish active compounds. Using this approach, we

identified 239 compounds in the bioactive library that affected the ER mobility (Figure 6 – figure

supplement 1B). Among these compounds, the correlation between the two screen replicates was

high ($R^2 = 0.92$) and the level of activity was reproducible (the slope for active molecules was

0.94). Some active compounds could be clustered based on scaffold homology, but most clusters

consisted of one or only a few members (Figure 6 – figure supplement 1C, D). Structural

clustering was employed to identify known ER modulators where the vendor-provided

annotation was poorly defined (Figure 6 – figure supplement 1C). Our results demonstrate that

htSMT is reproducible and robust when screening large collections of molecules.

Most active molecules from the screen were not structurally related to steroids (Figure 6 –

figure supplement 1C, D). On the other hand, many compounds could be grouped based on their

reported biological targets or pathways (Figure 3, Figure 6 – figure supplement 2A, B). For

example, heat shock protein (HSP) and proteasome inhibitors consistently increased f_{bound},

whereas cyclin dependent kinase (CDK) and mTOR inhibitors decreased f_{bound}. Though many

CDK inhibitors lack within-family specificity[47] (Figure 6 – figure supplement 2A, pan-CDK), we

found that CDK9-specific inhibitors more strongly affected ER dynamics than did CDK4/6-

specific inhibitors. Furthermore, as with selective AR and GR antagonists, inhibitors targeting

ALK, BTK, and FLT3 kinases that have not been shown to interact with ER have no impact on

ER dynamics when assessed using SMT (Figure 3).

For the inhibitors of cellular pathways we identified, we used a dose titration to better

characterize the effect of each on ER dynamics. Potencies ranged from the sub-nanomolar to low

micromolar (Figure 6B), similar to the reported potencies of these compounds against their

cellular targets[48–59]. Additionally, we tested these molecules against AR (Figure 6 – figure

supplement 2C) and PR (Figure 6 – figure supplement 2D). Each SHR differed meaningfully

from the others in terms of the response to compounds identified through an ER-focused

screening effort. Again, the magnitude of ER SMT effect was largely consistent within a target

class (Figure 6A, Figure 6 – figure supplement 2A-D). The finding that structurally distinct

compounds exhibited similar effects based on their biological targets favors the view that these

biological targets must themselves interact with ER, and that the compounds therefore affect ER

dynamics indirectly. HSP90 is a chaperone for many proteins, including SHRs. In the canonical

model, hormone binding releases the SHR-HSP90 complex. Indeed, HSP90 inhibitors increased

f_{bound} for ER, AR, and PR, consistent with the hypothesis that one function of the chaperone may

be to adjust the equilibrium of SHR binding to chromatin (Figure 6 – figure supplement 2B-D).

Proteasome inhibition also leads to ER immobilization on chromatin[60], which aligns with the

results that we obtained in our htSMT screen of bioactive compounds. ER has been shown to be

phosphorylated by CDK[61], Src[62], or GSK-3 through MAPK and PI3K/AKT signaling

pathways[63], and therefore inhibition of these pathways could reasonably be expected to affect ER

dynamics measured using SMT. While CDK inhibition led to an increase in ER mobility,

inhibition of PI3K, AKT, or other upstream kinases showed no effect (Figure 3).

Interestingly, SMT dynamics of an ER triple point mutant engineered to lack previously

defined phosphorylation sites important for transactivation[63] (S104A/S106A/S118A) were

affected by CDK and mTOR pathway inhibitors (Figure 6 – figure supplement 3), suggesting that

additional phosphorylation sites can mediate the effects of CDK9 and PI3K/AKT signaling, or

that other molecular targets of CDK and PI3K/AKT can act indirectly to alter the motion of ER.

The change in ER protein dynamics for characterized pathway inhibitors such as those targeting

CDK and mTOR is subtle but consistent across compounds, suggesting biological meaning in

these observations and highlight the need for accurate and precise SMT measurements. Hence

htSMT screening offers the promise of providing comprehensive pathway interaction

information or revealing novel interaction mechanisms.

Kinetic htSMT Facilitates Evaluation of Small Molecule Mechanism of Action

Since SMT can identify compounds that act either directly on a fluorescent target, or through

some intermediary process, we sought to distinguish between these alternative modes of action.

We hypothesized that by investigating the rate at which changes in protein motion emerge, SMT

could be used to distinguish direct versus indirect effects on ER activity. Given the live-cell

setting of SMT, we configured a data collection mode that allows for measurement of protein

dynamics in set intervals after compound addition (kinetic SMT or kSMT). Both ER agonists and

antagonists rapidly induce ER immobilization on chromatin when measured in kSMT ($t_{1/2} = 1.6$

minutes for estradiol; Figure 4C). On the other hand, HSP90 inhibitors like ganetespib and

HSP990 exhibit a delay of 5 to 7 minutes before alterations in ER dynamics appear, after which

we observed an increase in f_{bound} with a $t_{1/2}$ of 19.3 and 17.5 minutes, respectively. The overall

effect of these compounds reached a plateau after an hour (Figure 6B). Proteasome inhibitors,

e.g. bortezomib and carfilzomib, acted even more slowly, with changes in ER dynamics

emerging only after 40 minutes, and slowly increasing over the four-hour measurement window

(Figure 6C). Hence this exploration of SMT kinetics represents an important tool that can

facilitate differentiation between on-target and on-pathway modulators. We believe that this

approach will permit, for example, rapid mechanistic characterization of active compounds in a

drug discovery setting.

To further differentiate the effect of pathway inhibitors on ER protein dynamics, we sought to

characterize relative ER residence times for each such molecule. In contrast to the SERDs and

SERNs, although HSP90 inhibition by HSP990 and ganetespib resulted in an increase in f_{bound},

we observed a decrease in the total number long binding events by two- and four-fold,

respectively, while the binding times were similar to that observed with DMSO alone (Figure

6D-E). These results suggest HSP90 inhibition primarily increases k^*_{on} while leaving k^*_{off}

largely unaffected. On the other hand, inhibition of the proteasome led to an increase in both the

number and duration of long binding events. These results demonstrate that ER-chromatin

binding can be modulated by changing the rate of association or disassociation, and that the

inhibition of specific cellular partners can affect these rates differentially. Taken together with the

different kinetics for direct ER, HSP90, and proteasome modulators, our data suggest that each

class of molecule alters ER dynamics through distinct mechanisms.

Discussion

Many pathways that regulate the fundamental biochemistry of cells depend upon the

interaction of protein "sensors" with distinct protein "effectors" that engage transiently to trigger

a change in cell physiology. Although the fundamentals of this process have long been

appreciated, biochemical investigation of these protein interactions has typically required *in vitro*

reconstitution or has been interrogated through pull-down assays after cell permeabilization.

Here we report the combination of SMT, a type of super-resolution microscopy, with high

content microscopy as a means of visualizing individual protein motion in millions of live cells,

and under circumstances where the effect of small molecule inhibitors can be assessed

quantitatively. We demonstrate the capabilities of the htSMT platform by analyzing the behavior

of SHRs, a class of sensors that mediate hormone-induced modulation of gene expression, and in

particular the dynamics of the ER.

To validate our htSMT platform, our analysis initially focused on the very rapid

immobilization of SHRs on chromatin observed in cells exposed to their established, cognate

steroid ligands. The technique proved to be highly quantitative, effectively evaluating ligands

whose potency differ by more than four orders of magnitude, and readily characterizing

differences in both the sensitivity and the selectivity of the steroid receptor family. These

observations prompted us to apply htSMT to screen thousands of bioactive compounds, which

we hypothesized would reveal new chemical matter as well as enable comprehensive

pharmacologic dissection of ER pathway interactions. As further validation of the technique,

automated screening of ER dynamics using htSMT identified all 30 known steroid ligands from a

library of 5,067 bioactive compounds. The potency of these steroid ligands with respect to

alterations in ER dynamics varied across a thousand-fold range, demonstrating the dynamic

range of this single experimental setup. Among molecules known to behave as ER signaling

inhibitors, the change in ER dynamics correlated closely with the ability of these compounds to

block estrogen-induced proliferation of estrogen dependent breast cancer cells, demonstrating the

ability of htSMT to document structure-activity relationships in a chemical series across

disparate and biologically relevant readouts. Since our analysis relies only on detection of

changes in protein dynamics, this unbiased readout will prove broadly useful in screening

libraries of compounds to identify starting points for the development of new therapeutics. In

fact, our analysis identified 209 non-steroidal molecules that affect ER dynamics, which are

likely to act elsewhere in the network of ER-interacting proteins.

Our characterization of the htSMT platform and subsequent screen highlighted some

important considerations for future screening efforts using single molecule tracking. Notably, the

observation that cell-to-cell variability is the dominant driver of assay variance, when compared

to other sources like the microscope or the assay plate, suggests that an even larger field of view

would sample cells more effectively and result in a more stable dynamical estimate (Figure 3 –

figure supplement 3). This could be particularly important for detecting subtle dynamical

changes such as those seen with the mTOR inhibitors where the maximal change in f_{bound} was

only around 5% (Figure 6). Similarly, for our screening assay we chose a frame interval of 10

msec, which proved very sensitive to detecting a wide range of compound effects but is

necessarily limited in the types of perturbations it can detect. The diffusion of the most mobile

ER population was well below the upper limit of detection for 10 msec, suggesting that faster

frame rates were not necessary, but this may not be the case for other protein targets. On the

other hand, ER has been reported to have multiple low-mobility chromatin binding states (Wagh

2023), but these low mobility states are below the assay lower bound set by our localization error

$(D_{loc} = \frac{localization\ error^2}{\Delta\tau})$ and would require a slower frame interval to differentiate.

The observation that by screening large libraries of bioactive compounds for an effect on

protein motion, htSMT can define networks of biochemical signaling pathways is a critical

outcome of this high-throughput platform. Protein dynamics in the cell are not governed by

singular interactions between any two proteins but by biochemical networks within the cell that

intersect with the protein under observation. Our work enables construction of a putative

interaction network connecting nodes involving different proteins, the identities of which were

deduced based on the impact of known inhibitors on the dynamics of steroid receptors (Figure

6F). The interaction map derived from unbiased htSMT screening recapitulates many known

biological interaction partners of the ER, in a single experimental setup using protein motion as a

sole readout. More experiments are necessary to determine which nodes represent direct physical

interactions and which occur through intermediaries. To our knowledge, the impact of HSP inhibition

in increasing ER-chromatin association has never been described, neither has the link between inhibition

post-translation modifying enzymes like the CDKs or mTOR and ER dynamics ever been described. We

believe that identifying additional cellular interaction networks through htSMT will provide an

important foundation for the broader understanding of biochemical regulatory mechanisms. For

example, CDK4/6 inhibitors are co-administered with SERDs and SERMs to improve

therapeutic outcomes[64,65]. CDK4/6 inhibition only minimally affects ER dynamics in SMT

(Figure 3), supporting the view that the combination of CDK4/6 inhibition with ER antagonists

functions through a non-redundant ER-independent mechanism[66].

Most cell-based assays cannot easily be configured for kinetic analysis of treatment effects,

as these typically involve fixed-endpoint, aggregate readouts. With such endpoint assays,

biochemical feedback and compensatory mechanisms often confound interpretation of the direct

effect of a change in treatment conditions. SMT, however, when implemented in a high-

throughput system that exhibits consistency over long time intervals, permits kinetic analysis of

treatment effects on protein dynamics in real time with high reliability. Such kinetic analyses

help to define pathway cascades; in a first instance, they can be used to rapidly identify

compounds that likely engage directly with a therapeutic target.

Mechanistically, an increase in f_{bound} provoked by a change in cell conditions suggests that

molecular interactions with the protein being analyzed have been stabilized (either made more

probable or longer-lasting), while a decrease in f_{bound} suggests the opposite: molecular

interactions that have been made less robust. Unexpectedly, we observed both SERDs and

SERMs, which antagonize ER, cause an increase in nuclear bound fraction, likely via chromatin

binding, in a way that mimics what is seen with traditional ER agonists. We showed that this

mechanism is not a commonality among all SHRs, and indeed inhibitors of AR and GR behaved

more like the prototypical competitive antagonist. Inhibition of HSP90 also produces a marked

increase in ER f_{bound}, though these binding events are more transient (Figure 6E). Transcription

factors are thought to find binding sites through free 3D diffusion, 1D sliding, and transient, non-

specific DNA binding[5,12,20,21,24]. ER antagonists may function by promoting ER binding to non-

specific decoy chromatin sites, thus reducing the amount of ER able to activate transcription at

ER-responsive genes[31–33]. Previous work has also shown that SERDs, and to a lesser extent

SERMs, induce an alternate conformation of ER, thereby inhibiting co-factor recruitment[31]. If

co-factors like CBP and p300 stabilize ER-chromatin binding, then efficacious inhibitors might

exhibit shorter binding times compared with agonist stimulation[31]. The SERMs 4OHT and GDC-

0810 dramatically increase ER-chromatin binding frequency and only modestly increase binding

times; fulvestrant and GDC-0927 strongly increase both ER-chromatin binding frequency and

residence time. Therefore, htSMT suggests that agonists and antagonists of SHR signaling

operate through a previously unappreciated, unified mechanism of chromatin immobilization,

meriting further investigation.

Lastly, we note that fast-SMT, as implemented here, can define relationships between the

structures of chemical inhibitors and their effects on a fundamental property of protein regulatory

elements: their dynamics in living cells. In our hands, these measurements proved far more

reliable in predicting the ability of an ER antagonist to block cell proliferation than an ER

degradation assay, a conclusion that was only able to be drawn due to the scale of htSMT

screening capabilities. Notably, saturable dose responses were observed for ER antagonists at

much lower concentrations using htSMT than with ER degradation assays, suggesting that this

method will be more sensitive for identifying novel therapeutic agents than with corresponding

traditional assay formats. Thus, SMT can identify promising compounds the activity of which

might not otherwise be measurable, provide insight into their mechanism(s) of action, and in the

native environment of the cell.While ER is a transcription factor, the same principles may apply

broadly to regulatory mechanisms of proteins with diverse function. In this context, we conclude

that the application of technologies for measuring protein dynamics at scale will prove broadly

applicable to the elucidation of biological mechanisms.

Author Contributions

D. A., H. P. B., D. T. M., & Y. T. conceived the research project; D. A., H. P. B., X. D., E. G.,

A. He., J. H., H. L., D. T. M., S. E. P., R. T., & Y. T. contributed to study design; H. C., K. F., E.

G., R.Kr., H. L., D. T. M., S. E. P., A. S., & R. T. performed experiments; H. C., E. G., H. L., D.

T. M., A. S., & R. T. generated reagents; L. A., D. A., A. Ha, A. He., H. L., D. T. M., C. S. & R.

T. analyzed data; S. A., R. B., M. B., R. G., A. Ha, A. He, S. L. J., R. Ke., A. K., J. L., & L. M.

designed and wrote software; R. B., X. D., R. Ke., K. L., B. M., & D. T. M. designed and built

hardware; D. A., H. P. B., R. B., J. H., A. K ., & R. Ke. Supervised the work; L. A., D. A., A. He.,

H. L., D. T. M., S. E. P., & R. T. wrote the original draft; L. A., D. A., H. P. B, R. B., X. D., A.

He., J. H., H. L., D. T. M., R. T., & Y. T. reviewed and revised of the original draft.

Acknowledgements

The authors extend their deepest gratitude to all the employees and consultants of Eikon, past

and present, especially Caitlyn Bonilla, Tiffany Cheng, Michael Hirte, David Hoffman, Fedor

Ilkov, Yan Li, Anuja Lohia, Edith Martinez-Soto, Eugene Masterov, Mai Nguyen, and Gregory

Snyder. Their tireless work enabled the experiments described here. We thank Rand Miller,

Roger Perlmutter, Yan Li, Robert Tjian for helpful discussions and critical feedback on the

direction of our investigation and on the resulting manuscript. Eikon Therapeutics provided all

funding.

Figure captions

Main Figures

Figure 1. Benchmarking a high throughput single molecule tracking platform

a) Diffusion state probability distributions from three cell lines expressing Histone H2B-

540   Halo, Halo-CaaX, or Halo alone. Shaded bins represent the diffusive states characteristic

541   of each cell line.

542   b) Example field of view. Equal mixture of H2B-Halo, Halo-CaaX and free Halo cell line

543   single molecule images (top) and reference Hoechst image (bottom). Insets show zoom-

544   ins to individual cells and to sequential frames of individual molecules. Image intensities

545   are equivalently scaled across panels.

546   c) Single-cell diffusive states extracted from©), colored based on similarity to the H2B-,

547   CaaX-, or free-Halo dynamics reported in (b).

548   d) Heatmap representation of 103757 cell nuclei measured from a mixture of Halo-H2B,

549   Halo-CaaX, or Free Halo mixed within each well over 1540 unique wells in five 384-well

550   plates. Each horizontal line represents a nucleus. Cells were clustered using k-means

551   clustering and labels assigned based on the diffusive profiles determined in (b).

552   e) Ensemble state occupation of all trajectories recovered from a mixture of Halo-H2B,

553   Halo-CaaX, free Halo cells. Mean state occupation from 308 assay wells.

554   **Figure 2**. Chromatin binding by steroid hormone receptors is affected by compound treatment.

555   a) Distribution of diffusive states for Halo-AR, Halo-ER, Halo-GR, and Halo-PR in U2OS

556   cells before and after stimulation with an activating ligand. The area in the shaded region

557   is f_{bound}. Shaded error bands represent S.D.

558   b) Selectivity of individual SHRs to their cognate ligand compared with other steroids, as

559   determined by f_{bound}. Error bars represent SEM across three biological replicates.

Figure 3. Bioactive screen results of change in estrogen receptor f_{bound} for two biological

replicates, each with 2 well replicates per compound of 5067 compounds. Select inhibitors are

grouped and uniquely colored by pathway or target. Error bars are SEM of all replicates.

Figure 4. Selective ER modulators and degraders induce DNA binding measurable through

htSMT

a) Diffusion state probability distribution for ER treated with 100 nM of exemplified

566   SERMs and SERDs. Each state distribution is generated from 10,000 randomly sampled

567   nuclear trajectories per assay well. Shaded regions represent the S.D.

568   b) Change in f_{bound} as a function of a 12-pt dose titration of fulvestrant (5), 4-OHT (6),

569   GDC-0810 (8), AZD9496 (9), or GDC-0927 (10) with fitted curve. Compounds colored

570   as in (a). Error bars represent the SEM across three biological replicates.

571   c) Change in f_{bound} as a function of time after agonist or antagonist addition, fitted with a

572   single exponential. Compounds colored as in (a). Estradiol (1, green) and DMSO added

573   for comparison. Error bars represent SEM across three biological replicates.

574   d) Maximum effect of SERMs and SERDs on f_{bound}. Each box represents quartiles while

575   whiskers denote the 5-95th percentiles of single well measurements, measured over a

576   minimum of four days with at least 8 wells per compound per day.

577   e) Fluorescence Recovery After Photobleaching of ER-Halo cells, treated either with

578   DMSO alone or with 100 nM SERM/D. Curves are the mean ± SEM for 18-24 cells,

579   colored as in (a).

580   f) Quantification of FRAP recovery curves to measure recovery 2 minutes after

581   photobleaching. Whiskers denote the 5-95[th] percentiles of single cell measurements.

582   g) Quantification long-lived tracks, each point representing the fraction of trajectories

583   greater than 10 seconds for a single biological replicate consisting of 3-10 wells per

584   condition. Dashed line represents the median fraction of trajectories lasting longer than

585   10 sec for Histone H2B-Halo, which is the upper limit of measurement sensitivity. *

586   indicates sample with $p < 0.05$ as measured by t-test.

587   **Figure 5.** htSMT can be used to determine chemical structure-activity relationships.

588   a) Correlation of potency measured by ER degradation and cell proliferation in MCF7 cells

589   (black) and T47d cells (magenta) for compounds in the GDC-0927 structural series. Cell

590   proliferation and ER degradation-derived potencies are the mean of 4 and 3 biological

591   replicates, respectively.

592   b) Change in f_{bound} across a 12-pt dose titration of compounds 11 through 16, colored by

593   structure. Points are the mean ± SEM across three biological replicates.

594   c) Correlation of potency measured by change in f_{bound} and cell proliferation in MCF7 cells

595   (black) and T47d cells (magenta) for compounds in the GDC-0927 structural series. Cell

596   proliferation and SMT-derived potencies are the mean of 4 and 3 biological replicates,

597   respectively.

Figure 6. Bioactive molecules targeting pathways associated with ER affect its dynamics

a) Bioactive screen results with select inhibitors grouped and uniquely colored by pathway.

Change in f_{bound} across a 12-pt dose titration of three representative compounds targeting

each of HSP90, mTOR, CDK9 and the proteasome. Individual molecules are denoted by

specific shapes. Error bars represent SEM across three biological replicates.

b) Change in f_{bound} as a function of time after compound addition. Estradiol treatment

(black) is compared to ganetespib (blue circles) and HSP990 (blue squares). Points are

bins of 4 minutes. Error bars represent SEM across three biological replicates.

c) Change in f_{bound} as a function of time after compound addition. Estradiol treatment

(black) is compared to HSP90 inhibitors ganetespib (blue circles) and HSP990 (blue

squares); proteasome inhibitors bortezomib (red circles) and carfilzomib (red squares).

Points are bins of 7.5 minutes of htSMT data, marking the mean ± SEM. The shaded

region denotes the window of time used during htSMT screening.

d) Track length survival curve of ER-Halo cells treated either with DMSO alone, with

estradiol stimulation, or with 100 nM HSP90 or proteasome inhibition. Track survival is

plotted as the 1-CDF of the track length distribution; faster decay means shorter binding

times.

e) Quantification of the number of long-lived trajectories as a function of treatment

condition. All conditions were normalized to the median number of trajectories in

DMSO.

f) Diagram summarizing pathway interactions based on htSMT results for ER, AR, and PR.

Supplementary Figures

Figure 1 – figure supplement 1. Overview of the htSMT workflow

a) Schematic representation of the end-to-end automated htSMT process, beginning with the physical preparation of the sample; image collection on a custom-assembled super-resolution microscope; detection of fluorescent spots and subsequent trajectory reconstruction; image processing and QC; extraction of relevant features by combining trajectories with cellular image data; and finally interpretation by a biologist.

b) Schematic view of the automated htSMT platform.

c) Example Hoechst images to be detected and removed during the data QC phase. Debris, incorrect focus, uneven illumination and empty FOVs which can contaminate screening data are are removed using a deep-learning based classifier of the Hoechst image.

Figure 1 – figure supplement 2. Characterization of htSMT system performance.

a) Distribution of localization error measured across multiple independent wells using Histone H2B-Halo cells. Median localization error of 39 nm.

b) Number of measurable nuclei in a 94 by 94 µm FOV. Each box plot is the distribution over wells in one 384-well plate.

c) Single-cell state distributions for Histone H2B, CaaX and free HaloTag cell lines.

d) Cumulative number of trajectories as a function of imaging time for each individual cell type for cells labeled with 10 pM JF_{549}. Shaded error bars are 1 S.D.

Figure 2 – figure supplement 1. Activation of steroid hormone receptors changes free diffusion and impacts downstream gene expression.

a) Cartoon illustration of SHR function. Under basal conditions, SHRs are sequestered in complex with HSP90 and other cofactors. Upon ligand binding, the receptor dissociates from the inactive complex, dimerizes, and binds to DNA.

b) mRNA transcript levels in log(FPKM) of AR, ESR1, PR, and NR3C1 for the engineered

U2OS cell lines compared to the parental U2OS line as well as three reference breast

cancer cell lines.

c) Distribution of diffusive states for Halo-AR, Halo-ER, Halo-GR, and Halo-PR in U2OS

cells before and after stimulation with an activating ligand. Arrows indicate the mean

D_{free}. Shaded error bands represent 1 S.D.

d) Selectivity of individual SHRs to their cognate ligand compared with other steroids, as

determined by D_{free}. Error bars represent SEM.

e) Reference gene sets induced after 24 hrs of stimulation with 25 nM estradiol. The top five

induced gene sets for Halo-ER ectopic expression increases were not significantly

induced in the parental U2OS line.

f) Bar plot showing the -log(q value) from the top 50 most significantly induced gene sets

after estradiol stimulation. Gene sets characteristic of ESR1 or the estrogen response are

colored green.

Figure 3 – figure supplement 1. Experimental design for bioactive molecule screen.

a) Cumulative number of trajectories as a function of imaging time for DMSO-treated cells

labeled with 20 pM JF_{549}. Shaded error bars are 1 S.D.

b) Median of the jump length distribution, reported as a function of frame number and

calculated in 10-frame increments comparing DMSO- and estradiol-treated wells. Curves

are the mean, shaded error bands are 1 S.D.

c) Filtering criteria for bioactive compounds in the screen: compounds without SMILES

data; compounds with a molecular weight less than 226 Da or more than 800 Da; heavy

metals and compounds with more than 60% non-carbon atoms; and compounds with

known fluorescent structures (e.g. fluorescein) were removed before screening. During

and after screening additional compounds were removed if they were observed to be

fluorescent or if they failed to meet data quality criteria.

d) Composition of the bioactive screening set based on the biological pathway targeted as

provided by the manufacturer.

Figure 3 – figure supplement 2. Screen of bioactive molecules produces robust data with good

assay performance.

a) Change in f_{bound} repeatability of screening results assessed over two biological replicates.

Compounds in magenta were identified as active if their magnitude of change in f_{bound}

when averaged across both replicates was greater than 0.05. Of the expected 30 positive

control compounds, 26 significantly increased f_{bound} in both replicates (gray outline).

Three compounds increased f_{bound} but were only present in one replicate after filtering.

Exemplified positive controls including the agonist estradiol (1) and the antagonists

fulvestrant (5), 4-OHT (6) and bazedoxifene (7) are exemplified. A linear regression was

fit to the magenta compounds to determine slope and correlation.

b) Dose titration of estradiol for each plate in the bioactive molecules screen fit with a

logistic regression to determine potency.

c) Z'-factor analysis for the bioactive screen. Each point represents the Z'-factor of a single

384-well plate, measuring the difference between DMSO and 25 nM estradiol treatment.

Plates with very low Z'-factors were removed from further analysis.

d) EC_{50} values extracted from each curve fit in (c). Shaded region is a three-fold range in

potency.

e) Distribution of change in f_{bound} for control DMSO wells.

f) Change in f_{bound} for unfiltered screening results plotted per assay plate with the negative

control (DMSO, blue) and positive control (Estradiol, red) highlighted. Each tick mark on

the x-axis is a separate assay plate. Each point is the mean per compound per assay plate,

error bars are the SEM.

g) Boxplot showing the number of cell nuclei measured for each compound tested.

Whiskers are the 1st and 99th percentiles.

Figure 3 – figure supplement 3. Contributions of different sources of variability of jump length

in the htSMT bioactive compound screen against ER.

a) Change in f_{bound} for 30 known ER-interaction molecules circled in (b), ordered by the

magnitude of effect. Error bars are the SEM.

b) Variance in mean jump length after subsampling increasing numbers of jumps from the

bioactive screen, resampling at different levels of aggregation (microscope, plate, well,

etc.). Law of large numbers scaling would result in a line with slope -1, and deviations

from linearity are the result of intrinsic sources of variance over microscopes, plates, etc.

c) Sources of variance attributed to differences between microscopes, plates, sample wells,

fields of view (FOV) and cells. Solid bars are analyzed from all wells in the bioactive

screen, half-tone bars are assessing DMSO wells only. The asterisk highlights the

expected well-to-well variability introduced when assaying compounds.

Figure 3 – figure supplement 4. Antagonists of ER (a), PR (b), AR (c) and GR(d) show distinct

effects on target protein dynamics. Change in f_{bound} after addition of 1 µM antagonist either in the

absence or presence of the cognate agonist.

Figure 3 – supplemental table 1. A list of the compounds composing the bioactive screening

library.

Figure 4 – figure supplement 1. SERMs and SERDs decrease free diffusion and increase f_{bound}

rapidly after addition.

a) Normalized occupation of diffusive states (diffusion coefficient 0.2 – 100 µm^2/sec) for

100 nM SERD or SERM-treated samples compared with DMSO. Histograms are

normalized to integrate to 1. Shaded regions represent bin-wise standard deviations.

Curves are the average of 3-4 biological replicates, with 8 well replicates per condition

(>1200 cells total)

b) Results from a single-exponential association fit to the data in Figure 4C.

c) Change in diffusive state distribution as a function of time after compound addition for

estradiol and fulvestrant. Each curve represents the mean of three biological replicates.

Shaded regions are the bin-wise standard deviation.

d) Change in f_{bound} of selective AR or GR antagonists from the bioactive screening set.

e) Track length survival curve of ER-Halo cells treated either with DMSO alone or with

SERM/D. Track survival is plotted as the 1-CDF of the trajectory length distribution;

faster decay means shorter binding times.

f) Table of results of the slow decay rate constant (k_{slow}) from residence time SMT curve

fits. Fits were performed on the aggregate of three biological replicates. Taken together

with the f_{bound} determined in 4D, an upper limit of the inferred k_{on} can be calculated from

the equation assuming all bound molecules will have $k^{*}_{off} = k_{slow}$. Cells marked with an

asterisk are ones in which k_{slow} could not reliably be distinguished from photobleaching.

Figure 5 – figure supplement 1. GDC-0927 structural variants characterized by ER degradation

or cell proliferation assays.

a) Western blot of the ER-expressing breast cancer cell lines MCF7 and T47d compared to

ER expression in ER-null lines SK-BR-3 or U2OS. Samples were treated with fulvestrant

for 24 hours prior to lysis. Fulvestrant treatment leads to the degradation of ER, even

when fused to HaloTag.

b) Example of GDC-0927 induced degradation of ER in different cell lines measured by

immunofluorescence against ER. Cells were exposed to compound for 24 hours prior to

fixation. Image pixel intensities are equivalently scaled.

c) Example compound dose titrations showing change in mean nuclear intensity as a

function of compound concentration.

d) Example compound dose titrations measuring cell proliferation, normalized to DMSO-

treated cells, of cells treated with analogs of GDC-0927.

Figure 6 – figure supplement 1. Deeper investigation of bioactive screening data identifies

cutoff for active molecules.

a) Examples of compound effects on ER f_{bound} from a dose titration experiment. 92

compounds from the primary screen are ranked based on the magnitude of their effect.

Compounds colored in black repeatably showed dose-dependent changes; magenta

compounds were inactive in a dose titration.

b) Example dose titration of compounds of increasing overall change in ER f_{bound}.

c) Two replicates of the bioactive screen as in Figure 3. Active compounds are colored

based on structural scaffold.

d) A quantification of the number of compounds associated with any given cluster.

Singletons (cluster 21) represent the majority of active compounds.

Figure 6 – figure supplement 2. Some pathway inhibitors modulating ER dynamics are specific

to ER.

a) Change in f_{bound} of ER following treatment with inhibitors of HSP90, the proteasome,

mTOR, and CDK from the initial bioactive screen. CDK inhibitors were further stratified

into CDK4/6 inhibitors, CDK9 inhibitors, or inhibitors without strong selectivity for a

particular family (pan-CDK). Line represents the median for each target. * indicates

sample with $p < 0.05$ as measured by t-test.

b) Change in f_{bound} of ER for 97 bioactive molecules, colored by their pathway annotation.

Compounds are plotted in their rank order based on their effect in ER. Error bars are the

SEM of three biological replicates.

c) Change in f_{bound} of AR for the same molecules from (b). Compounds are plotted in their

rank order based on their effect in ER. Error bars are the SEM of three biological

replicates.

d) Change in f_{bound} of PR for the same molecules from (b). Compounds are plotted in their

rank order based on their effect in ER. Error bars are the SEM of three biological

replicates.

Figure 6 – figure supplement 3. Dose titration plots of ER(S104A/S106A/S118A) with mTOR

and CDK9 compounds. Each point represents the mean and SEM of three biological

replicates.

Supplemental movie 1. Example SMT field of view as shown in Figure 1. A representative

mixture of H2B-Halo, Halo-CaaX, and free Halo expressing cells. Video is played back

at 10 frames per second, 10x slower than real time.

Supplemental movie 2. Example field of view of SMT from Halo-ER cells. Video is played

back at 10 frames per second, 10x slower than real time.

Supplemental movie 3. Example field of view of SMT from Halo-ER cells treated with 25 nM

estradiol for 1 hr prior to imaging. Video is played back at 10 frames per second, 10x

slower than real time.

785
Methods

Cell Lines

U2OS (ATCC Cat. No. HTB-96), MCF7 (ATCC Cat. No. HTB-22), T47d (ATCC Cat. No.

HTB-133) and SK-BR-3 (ATCC Cat. No. HTB-30) were grown in DMEM (Cat. No. 1056601,

Gibco DMEM, high glucose, GlutaMAX Supplement, Thermofisher) supplemented with 10%

Fetal Bovine Serum (Cat. No. 16000044, Thermofisher) and 1% pen-strep (Cat. No 15140122,

Thermo Fisher) and maintained in a humidified 37 °C incubator at 5% CO_2 and subcultivated

approximately every two to three days.

HaloTag-expressing cell lines

For H2B, CaaX, ER, AR, and PR-HaloTag fusions, mammalian expression vectors

containing the fusion gene under the control of a weak L30 promoter and containing a Neomycin

resistance marker were transfected into U2OS cells at 70% confluence using FuGENE 6 (Cat.

No. E2691, Promega). Transfected cells were selected with G418 (Cat. No. 10131027, Thermo

Fisher) at 500 µg/mL, then clonally isolated. Clones expressing the desired fusion gene were

determined first by staining with 100 nM JF_{549}-HTL (Cat. No. GA1110, Promega) and 50 nM

Hoechst 33342 and identifying clones with the expected distribution of JF_{549} signal. Between

three and six clones were subsequently tested using SMT conditions for response to a control

compound, and the most homogenous clones were subsequently expanded for further testing.

Unless otherwise specified, all experiments are with a single, clonally isolated cell line. Because

U2OS cells express GR endogenously, HaloTag was inserted right before the stop codon of

endogenous NR3C1 via homology-directed repair using CRISPR/Cas9. The HaloTag knock-in

was validated by imaging using HTL-JF$_{646}$ staining and through DNA sequencing[67].

Western Blot

Cells were grown in the same conditions as described previously. 1.5×10^6 cells were seeded

per well in a 6-well plate in DMEM overnight, followed by compound treatment (DMSO or

100nM fulvestrant) the following day for 24 hours. Cells are lysed in 200 μL 1X Cell Lysis

Buffer (catalogue number 9803, Cell Signaling). Protein lysate concentration is then determined

using BCA protein assay kit (Catalog number 23225, Pierce™ BCA Protein Assay Kit) following

manufacturer instructions. Capillary Western Immunoassay were performed using Jess Protein

Simple following manufacturer's instruction (protein simple, USA). Levels of ⍰ER

(1:100, RM-9101) were normalized to loading control β-tubulin (1:100, NC0244815 LI-COR

92642213, Thermo Fisher). The peaks were analyzed with the Compass software (Protein

Simple, USA).

RNA-seq

Cells were seeded into 12-well tissue-culture treated plates at densities of 250,000 cells

(U2OS-WT), 200,000 cells (U2OS-ER), or 300,000 cells (MCF7, SKBR3, T47d) per well. 24

hours later, cells were treated with estradiol at a final concentration of 25nM for 24 hours. To

process cells for total RNA, cells were washed twice with ice-cold PBS, lysed with 350uL Buffer

RLT (Qiagen 79216), scraped off the plate (Fisher 08100241), frozen on dry ice and stored at –20

degrees C. Cell lysates were then thawed, homogenized using QIAshredder columns (Qiagen

79656), and processed through the Qiagen RNeasy Micro kit (Qiagen 74004) using the standard

protocol and including the optional on-column DNase digestion step (Qiagen 79254). All

samples had a RIN score of 10 by

TapeStation (Agilent 5067-5576). RNA sequencing libraries were prepared from total RNA by

Novogene (CA). In brief, mRNA was purified from total RNA using poly-T oligo-attached

magnetic beads and fragmented. First-strand synthesis was performed using random hexamer

primers, second-strand synthesis was performed using dTTP, and libraries were prepared after

end repair, A-tailing, adapter ligation, amplification, and purification. Libraries were sequenced

on an Illumina NovaSeq with paired 150 cycle reads. For data analysis, paired-end reads were

aligned to the hg38 reference genome using Hisat2 v2.0.5, featureCounts v1.5.0-p3 was used to

count the number of reads mapped to each gene, and differential expression analysis was

performed using DESeq2 (1.20.0).

Single Molecule Tracking Sample preparation

Cells were seed on tissue culture treated 384-well glass-bottom plates at 6000 cells per well.

Seeded cells were then incubated at 37 °C and 5% CO_2 to allow adhesion overnight. For all SMT

experiments, cells were incubated with 5-100 pM of JF_{549}-HTL (Cat. No. GA1110, Promega) and

50 nM Hoechst 33342 for an hour in complete medium. Cells were then washed three times in

DPBS and twice in imaging media using an EL406 plate washer, which is fluoroBrite DMEM

media (Cat. No. A1896701, Thermo Fisher) supplemented with GlutaMAX (Cat. No. 35050079,

Thermo Fisher) and the same serum and antibiotics as growth media. Where appropriate,

compounds were serially diluted in an Echo Qualified 384-Well Low Dead Volume Source

Microplate (0018544, Beckman Coulter) to generate dose-titration source material. Compounds

were administered at a final 1:1000 dilution in cell culture medium. Each dose of a compound

has at least 2 replicates per plate and 3 plate replicates, 20 DMSO control wells and 2 no dye

control wells were randomized across each plate. Unless otherwise specified, compounds were

allowed to incubate for an hour at 37 °C prior to image acquisition.

Image Acquisition

Unless otherwise stated, all image acquisition using SMT was performed on a custom-built

HILO microscope based on a Nikon Ti2, motorized stage, stage top environmental chamber

(OKO labs), quad-band filter cube (Chroma), custom laser launch with 405 nm, and 561 nm

wavelengths, coupled to a Nikon TIRF illumination module by fiber optic element (KineFlex

HPV-P-3-S-405..640-0.7-APC-P2) delivering >10 mW and >600 mW of power in a gaussian

beam with a FWHM of approximately 250 µm to the back focal plane of the objective. Angle of

inclination and beam direction were adjusted by micrometer on the TIRF illuminator and

empirically set to maximize and flatten the signal across the camera field of view. Fluorescence

emission was passed through a high-speed filter wheel (Finger Lakes Instruments) and collected

with a backlit CMOS camera (Prime 95b, Teledyne). Images were acquired with a 60X 1.27 NA

water immersion objective (Nikon). Environmental chamber was set to 37^o Celsius, 95%

humidity, and 5% CO_2. For each field of view, 200 SMT frames were collected at a frame rate of

100 Hz, with a 2 msec stroboscopic laser pulse. 10 frames of the Hoechst channel were collected

at the same frame rate for downstream registration of trajectories to nuclei. Sample focus was

maintained using the reflection-based Perfect Focus System to determine the position of the

coverglass and apply an empirically-determined offset to focus into the sample.

Experiment design and sample size

All assays were designed with high throughput screening in mind. Unless otherwise stated,

experiments were performed with three biological replicates and within each assay plate having

at least three well replicates. In instances where a specific plate or a portion of a plate failed to

meet assay quality criteria (e.g. no respons from a positive control), those data point were

omitted and, where possible, the assay was repeated. For htSMT experiments, samples were

prepared and acquired such that a minimum 20 cells were sampled per assay well, resulting in an

minimum of 21,000 detections per well; 60 cells and 63,000 detections minimum per condition

per assay plate.

htSMT Image Analysis

Image acquisition produced one JF_{549} movie and one Hoechst per field of view. The JF_{549}

movie was used to track the motion of individual JF_{549} molecules, while the Hoechst movie was

used for nuclear segmentation. Tracking was accomplished in three sequential steps – detection,

subpixel localization, and linking – using a combination of existing methods. Briefly, spots were

detected using a generalized log likelihood ratio detector to test every 11x11 pixel window using

a gaussian kernel with a radius of 1.25 pixels and with a log likelihood detection threshold 14[68].

After detection, the estimated position of each emitter was refined to subpixel resolution using

Levenberg-Marquardt fitting[69–71] with an integrated 2D Gaussian spot model[72] starting from an

initial guess afforded by the radial symmetry method[73]. Detected spots were linked into

trajectories using a custom modification of a hill-climbing algorithm with a maximum linking

radius of 1.25 µm and allowing a maximum of 2 gap frames where a spot may go undetected but

still be linked within the same trajectory[10,74]. The same detection, subpixel localization, and

linking settings were used for all movies used in this manuscript.

For nuclear segmentation, all frames of the Hoechst movie were averaged to generate a mean

projection. This mean projection was then segmented with a neural network trained on human-

labeled nuclei, the output of which is a mask assigning a semantic category to each pixel in the

image[75]. Each spot was assigned to at most one nucleus using its subpixel coordinates.

To recover dynamical information from trajectories, we used state arrays[35], a Bayesian

inference approach, with the "RBME" likelihood function and a grid of 100 diffusion

coefficients from 0.01 to 100.0 $\mu m^2 s^{-1}$ and 31 localization error magnitudes from 0.02 to 0.08

μm. For each assay well, 10,000 trajectories were randomly sampled from the aggregated pool of

nuclear trajectories, and this set of trajectories was used for state inference. After inference,

localization error was marginalized out to yield a one-dimensional distribution over the diffusion

coefficient for each field of view. For single-cell analysis, we performed SMT and nuclear

segmentation on a mixture of U2OS cells bearing H2B-HaloTag,

HaloTag-CaaX, or free HaloTag. We then evaluated the marginal likelihood of each of a set of

100 diffusion coefficients on the set of trajectories within each segmented nucleus[30]. These

marginal likelihood functions were clustered with k-means (3 clusters), and the marginal

likelihood functions for each cell were ordered by their cluster index to produce the heat map. To

estimate the fraction bound (f_{bound}), we integrated the state array posterior distribution below 0.1

$\mu m^2 s^{-1}$. To estimate the free diffusion coefficient (D_{free}), we computed the mean of the posterior

distribution above 0.1 $\mu m^2 s^{-1}$ using the following equation:

$$D_{free} = \frac{\sum_{i=25}^{100} D_i * P(D_i)}{\sum_{i=25}^{100} P(D_i)}$$

htSMT data quality control

High content imaging approaches require image quanity control to systematically remove

aberrant measurements from the set. Primarily these are fields of view that are empty, that are out

of focus, or contain large piece of debris in the well or occlusion of the objective. To detect and

remove these fields of view, we trained a convolutional neural network classifier to score

Hoechst image quality on a scale of -1 to 1 (low quality to high quality). Annotations from five

independent annotators on a training set of ~1000 Hoechst images were used as input for the

model. After training, a threshold for filtering was empirically set so as to remove problem FOVs

such as those exemplified in Figure 1 – figure supplement 1c). Screening wells were only

considered if more than two fields of view passed QC, and screening plates were only considered

if more than 8 control wells could be included for normalization. During screening, compounds

with a standard deviation in f_{bound} greater than 0.15 were omitted from analysis and rescreened

where possible.

924
htSMT data analysis

Tracking results from the automated processing pipeline were analyzed using KNIME or

Spotfire (TIBCO). Individual f_{bound} or D_{free} measurements were associated with experimental

metadata and aggregated by condition. Change in f_{bound} was calculated as the difference between

the f_{bound} of each well and the median f_{bound} of DMSO in the same plate. Wells that had no cells in

the field of view or in which the field of view was out of focus were omitted from further

analysis. Compounds were assessed for assay interference using the median fluorescence

intensity of the tracking channel and omitted if it was more than 3 standard deviations higher

than the median intensity of the DMSO wells. Similarly, plates where the active and negative

controls could not be clearly resolved or where the significantly deviated from the performance

of the rest of the screen were removed from further analysis. Finally, compound with a variance

more than three standard deviations higher than the average compound variance (41 compounds;

0.08%) were removed from downstream analysis. Z'-factor between the active controls on a

plate and DMSO was calculated as previously described[38]. EC_{50} values were calculated in Prism

(GraphPad) by first log-transforming the molecule concentrations and then fitting to a four

parameter logistic curve.

Sources of variability analysis

The contribution of microscope-to-microscope, plate-to-plate, well-to-well, FOV-to-FOV and

cell-to-cell variability on the 2D jump distribution was estimated using a subsampling approach.

We consider a simplified model where the observed jump length distribution (Y) is a function of

the intrinsic stochasticity in jump length due to diffusion (X) with additional biases introduced at

the cell, FOV, well, plate or microscope level:

$$Y = B_{microscope} + B_{plate} + B_{well} + B_{FOV} + B_{cell} + X$$

For simplicity the biases are assumed independent, although these biases may indeed have

some dependence.

To estimate the variance of each bias term, we computed the variance in sample means for

different resampling schemes. Firstly we sample N jumps from all data from one replicate of the

bioactive screen and compute the average. The resulting sample mean is the average over all

sources of variability ($B_{microscope}$, etc.). In a second step, we sample a microscope and randomly

sample all jumps from all plates on that microscope. This averages over all other sources of

variability except $B_{microscope}$ can be expected to approach Var($B_{microscope}$) for large N. We

continune this same sampling scheme at the Plate-, Well-, FOV- and cell-level, such that for large

N results in estimated Var(B_{plate}), Var(B_{well}), Var(B_{FOV}), and Var(B_{cell}) respectively. 1000

rounds of sampling were performed for each resampling scheme, either including all wells or

only those containing the vehicle DMSO.

Clustering active molecules

Chemical structure-based clustering was performed on molecules identified as active (239 in

total). Molecular frameworks were computed as described by Murcko et al and as implemented

in Pipeline Pilot[76]. Molecular frameworks were clustered using functional class fingerprints

(FCFP_4) with a similarity threshold cut-off of 0.3 Tanimoto distance. A total of 21 clusters were

obtained with singletons being the major class (124 molecules). The next largest group was the

flavone class represented by 27 members, followed by a couple of diverse classes within the

steroidal class with 14 and 20 members respectively. The other category is the stilbene class with

7 members representing tamoxifen as one of the members. The remaining actives (47 molecules)

were grouped into one 3-membered cluster and all the others with 2 members per cluster.

Kinetic Experiments

Cells were seeded into a 384-well plate the day before, dyed, and washed as described above.

1 well with 25 FOVs per well were taken as a baseline reading. Then, while imaging, compound

was manually added to each well to a final concentration of 100 nM. Data was then collected for

20 wells. A pause was included between each FOV such that the entire imaging regime covers

the assay window. Change in f_{bound} was determined per-well relative to t=0.

For assays extending to 4 hours, the plate was imaged twice with 8 FOVs per well with

different FOV locations per readthrough to prevent photobleaching from impacting data. All data

presented represents was performed in three different biological replicates.

Residence Time Imaging

Sample preparation and execution of residence time imaging experiments were conducted in

a similar manner to the single molecule tracking assay described above with a few exceptions.

Samples were dyed with 1 – 10 pM JF$_{549}$ (Promega) and 50 nM Hoechst 33342 for an hour. 400

frames per field of view were collected with a camera integration time was set to 250 msec, and

laser sources reduced to 5 mW at the objective. During image acquisition, lasers were on

continuously. Compound incubation ranged from 1 to 4 hours. At least 8 well replicates were

collected per condition.

Residence Time analysis

Quantifying transcription factor binding times on DNA is an open problem with multiple

proposed solutions[21,77,78]. Here we adopted an approach similar to Hansen et. al. [19]. Image

processing, including spot detection, localization, and track reconnection were performed using

the same methods described above. Because residence time imaging selectively tracks slow-

diffusing molecules, individual localizations were limited to a 300 nm maximum displacement

for individual jump reconnections. Sets of trajectories for each field of view were binned into 1-

CDF distributions as previously described and fit to a two exponent decay model:

$CDF(t) = A(Fe^{-k_{fast}t} + (1 - F)e^{-k_{slow}t}).$

The k_{slow} term comprises both the rate of molecule unbinding (k_{off}) as well as photobleaching

(k_{bleach}) and diffusion of chromatin out of the focal volume. Often approaches attempt to derive

the unbinding rate by applying a correction such as subtracting a bleaching rate measured either

directly in the sample or by using a separate control sample[19,21], where $T_{corrected} = (k_{off}-k_{bleach})^{-1}$.

The inverse relationship between $T_{corrected}$ and k_{bleach} makes it highly sensitive and nonlinear to

noise in k_{bleach}. Because the 1-CDF distributions of compound-treated ER samples are so close in

decay rate to the control Histone H2B, background subtraction yields unfeasible k_{off} values after

correction. Instead we report only the uncorrected k_{slow} values with the understanding that these

represent a lower-bound of the actual k_{off}, but that within-experiment comparisons can be made

between conditions.

Fluorescence Recovery After Photobleaching

Images were acquired on a custom-built HiLo microscope as described above with a Spectra

Light Engine RS-232. Stimulation was directed using a miniscanner coupled with a Coherent

OBIS 561nm 100 mW laser. All imaging was performed using a 60X 1.27 NA water immersion

objective (Nikon). All experiments were performed at 37^o Celsius. For FRAP experiments, Cells

were seeded into a 384-well plate the day before, labeled with 50 nM HTL-JF$_{549}$, and washed as

described above. Compound was added to 100 nM final an hour before imaging. Then, a

prebleach image was acquired by averaging 10 consecutive images. Then 8-10 regions were

bleached (2 background, 6-8 cells) and 2 regions in cells were unbleached. Regions that were

bleached were bleached at 10% power without scanning. For the next 30 seconds, an image was

acquired every 200 ms, then every 1 second for 2 minutes. The background-subtracted average

intensity was measured in the region of interest over time and normalized to the average of the

fluorescence in the baseline images, then normalized to the unbleached regions to account for

readout-induced photobleaching of fluorophores. Data from 18-24 cells were pooled per

experiment for 3 biological experiments.

Immunofluorescence

Cells were grown in conditions as described previously. Cells were seeded in glass bottom

384-well plates coated with 0.05mg/ml PDL (Cat. No. A3890401, Thermofisher) at 6000 cells

per well for Halo- ER U2OS cells and 8000 for MCF7 and T47d cells. Cells were grown

overnight followed by compound treatment on the second day for 24 hours at 37 °C and 5% CO_2.

Compounds were serially diluted in an Echo® Qualified 384-Well Low Dead Volume Source

Microplate (0018544, Beckman Coulter) to generate a 21-point dose response at 1:3 dilution

starting from a concentration of 10mM. Compounds were administered at a final 1:1000 dilution

in cell culture medium. An 8 to 12-point dose response was selected based on the potency of each

compound. Each concentration was replicated at least once per plate and has at least 2 plate

replicates. Cells were fixed by addition of paraformaldehyde (Cat. No. 15710-S; Electron

Microscopy Sciences), with a final concentration of 4% for 20 minutes. Cells were then

permeabilized using blocking buffer containing 1% bovine serum albumin and 0.3% TritonX100

in 1x PBS for an hour at room temperature. Immunofluorescent staining of ER was carried out

using ⬚ER antibody (1:500, RM-9101) diluted in the same blocking buffer for 1 hour at room

temperature. Extensive washing with PBS was performed prior to secondary antibody staining.

Secondary antibody staining was carried out using Alexa fluor 488 conjugate anti-rabbit IgG

(1:1000, Cat. No. A32731, thermos Fisher) for an hour. Nuclear staining was carried out using

Hoechst 33342 solution at 1mg/ml. Imaging of immunofluorescence was done using the

ImageXpress Micro (Molecular Devices) at 10x magnification and 4 field-of-view per well.

Fluorescence intensity within the nucleus were quantified using CellProfiler[79]. All analysis and

curve fitting were carried out using Prism with DMSO as a baseline. ER degradation

experiments were performed in three biological replicates with the same source compounds.

Cell proliferation

Cells were grown and seeded in conditions as described above. Cells were seeded in 384well

plates (Cat. No. 353963, Corning) at 1000 cells per well for Halo-ER U2OS, 1200 cells for

SKBR3 and 1800 cells for MCF7 and T47d. Cells were grown overnight, then treated with

compounds the following day. Compound concentration and administration are the same as

described previously for the immunofluorescence assay. Plates are scanned in the IncuCyte live

cell analysis system (Sartorius) at 24-hour intervals for a total of 5 days using phase contrast.

Cell proliferation quantification was carried out by the built-in analysis function using whole

well confluency mask. All analysis and curve fitting were carried out using Prism with DMSO as

a baseline. MCF7 and T47d cell proliferation experiments were performed in four biological

replicates with the same source compounds.

1054

1055   # References

(1)   Heo; Sugita; Feig. Protein Assembly and Crowding Simulations. *Curr Opin Struct Biol* **2022**, *73*. https://doi.org/10.1016/j.sbi.2022.102340.

(2)   Guin, D.; Gruebele, M. Weak Chemical Interactions That Drive Protein Evolution: Crowding, Sticking, and Quinary Structure in Folding and Function. *Chem Rev* **2019**, *119* (18), 10691–10717. https://doi.org/10.1021/acs.chemrev.8b00753.

(3)   Shen, H.; Tauzin, L. J.; Baiyasi, R.; Wang, W.; Moringo, N.; Shuang, B.; Landes, C. F. Single Particle Tracking: From Theory to Biophysical Applications. *Chem Rev* **2017**, *117* (11), 7331–7376. https://doi.org/10.1021/acs.chemrev.6b00815.

(4)   Skóra, T.; Vaghefikia, F.; Fitter, J.; Kondrat, S. Macromolecular Crowding: How Shape and Interactions Affect Diffusion. *J Phys Chem B* **2020**, *124* (35), 7537–7543. https://doi.org/10.1021/acs.jpcb.0c04846.

(5)   Liu; Lavis; Betzig. Imaging Live-Cell Dynamics and Structure at the Single-Molecule Level. *Mol Cell* **2015**, *58* (4), 644–659. https://doi.org/10.1016/j.molcel.2015.02.033.

(6)   Los, G. V.; Encell, L. P.; McDougall, M. G.; Hartzell, D. D.; Karassina, N.; Zimprich, C.; Wood, M. G.; Learish, R.; Ohana, R. F.; Urh, M.; et al. HaloTag: A Novel Protein Labeling Technology for Cell Imaging and Protein Analysis. *ACS Chem Biol* **2008**, *3* (6), 373–382. https://doi.org/10.1021/cb800025k.

(7)   Mollwitz, B.; Brunk, E.; Schmitt, S.; Pojer, F.; Bannwarth, M.; Schiltz, M.; Rothlisberger, U.; Johnsson, K. Directed Evolution of the Suicide Protein O6-Alkylguanine-DNA Alkyltransferase for Increased Reactivity Results in an Alkylated Protein with Exceptional Stability. *Biochemistry* **2012**, *51* (5), 986–994. https://doi.org/10.1021/bi2016537.

(8)   Grimm Jonathan B; English Brian P; Chen Jiji; Slaughter Joel P; Zhang Zhengjian; Revyakin Andrey; Patel Ronak; Macklin John J; Normanno Davide; Singer Robert H; et al. A General Method to Improve Fluorophores for Live-Cell and Single-Molecule Microscopy. *Nature Methods* **2015**, *12* (3), 244–250. https://doi.org/10.1038/nmeth.3256.

(9)   Tokunaga Makio; Imamoto Naoko; Sakata-Sogawa Kumiko. Highly Inclined Thin Illumination Enables Clear Single-Molecule Imaging in Cells. *Nature Methods* **2008**, *5* (2), 159–161. https://doi.org/10.1038/nmeth1171.

(10)   Chenouard Nicolas; Smal Ihor; de Chaumont Fabrice; Maška Martin; Sbalzarini Ivo F; Gong Yuanhao; Cardinale Janick; Carthel Craig; Coraluppi Stefano; Winter Mark; et al. Objective Comparison of Particle Tracking Methods. *Nature Methods* **2014**, *11* (3), 281–289. https://doi.org/10.1038/nmeth.2808.

(11)   Speiser, A.; Müller, L.-R.; Hoess, P.; Matti, U.; Obara, C. J.; Legant, W. R.; Kreshuk, A.; Macke, J. H.; Ries, J.; Turaga, S. C. Deep Learning Enables Fast and Dense Single-Molecule Localization with High Accuracy. *Nature Methods* **2021**, *18* (9), 1082–1090. https://doi.org/10.1038/s41592-021-01236-x.

(12)   Boka, A. P.; Mukherjee, A.; Mir, M. Single-Molecule Tracking Technologies for Quantifying the Dynamics of Gene Regulation in Cells, Tissue and Embryos. *Development* **2021**, *148* (18). https://doi.org/10.1242/dev.199744.

(13)   Chandrasekaran, S. N.; Ceulemans, H.; Boyd, J. D.; Carpenter, A. E. Image-Based Profiling for Drug Discovery: Due for a Machine-Learning Upgrade? *Nature Reviews Drug Discovery* **2021**, *20* (2), 145–159. https://doi.org/10.1038/s41573-020-00117-w.

(14)   Caicedo Juan C; Cooper Sam; Heigwer Florian; Warchal Scott; Qiu Peng; Molnar Csaba; Vasilevich Aliaksei S; Barry Joseph D; Bansal Harmanjit Singh; Kraus Oren; et al. Data-Analysis Strategies for Image-Based Cell Profiling. *Nature Methods* **2017**, *14* (9), 849–

1102   863. https://doi.org/10.1038/nmeth.4397.

1103   (15)   Barentine, A. E. S.; Lin, Y.; Courvan, E. M.; Kidd, P.; Liu, M.; Balduf, L.; Phan, T.;
1104   Rivera-Molina, F.; Grace, M. R.; Marin, Z.; et al. An Integrated Platform for High-
1105   Throughput Nanoscopy. *Nature Biotechnology* **2023**, 1–8.
1106   https://doi.org/10.1038/s41587-023-01702-1.

1107   (16)   Needham Sarah R.; Roberts Selene K.; Arkhipov Anton; Mysore Venkatesh P.; Tynan
1108   Christopher J.; Zanetti-Domingues Laura C.; Kim Eric T.; Losasso Valeria; Korovesis
1109   Dimitrios; Hirsch Michael; et al. EGFR Oligomerization Organizes Kinase-Active
1110   Dimers into Competent Signalling Platforms. *Nature Communications* **2016**, *7*, 13307.
1111   https://doi.org/10.1038/ncomms13307.

1112   (17)   Yasui Masato; Hiroshima Michio; Kozuka Jun; Sako Yasushi; Ueda Masahiro.
1113   Automated Single-Molecule Imaging in Living Cells. *Nature Communications* **2018**, *9*
1114   (1), 3061. https://doi.org/10.1038/s41467-018-05524-7.

1115   (18)   Nixon-Abell, J.; Obara, C. J.; Weigel, A. V.; Li, D.; Legant, W. R.; Xu, C. S.; Pasolli, H.
1116   A.; Harvey, K.; Hess, H. F.; Betzig, E.; et al. Increased Spatiotemporal Resolution
1117   Reveals Highly Dynamic Dense Tubular Matrices in the Peripheral ER. *Science* **2016**,
1118   *354* (6311), AAF3928. https://doi.org/10.1126/science.aaf3928.

1119   (19)   Hansen, A. S.; Pustova, I.; Cattoglio, C.; Tjian, R.; Darzacq, X. CTCF and Cohesin
1120   Regulate Chromatin Loop Stability with Distinct Dynamics. *Elife* **2017**, *6*.
1121   https://doi.org/10.7554/eLife.25776.

1122   (20)   Paakinaho Ville; Presman Diego M.; Ball David A.; Johnson Thomas A.; Schiltz R.
1123   Louis; Levitt Peter; Mazza Davide; Morisaki Tatsuya; Karpova Tatiana S.; Hager Gordon
1124   L. Single-Molecule Analysis of Steroid Receptor and Cofactor Action in Living Cells.
1125   *Nature Communications* **2017**, *8*, 15896. https://doi.org/10.1038/ncomms15896.

1126   (21)   Mazza, D.; Abernathy, A.; Golob, N.; Morisaki, T.; McNally, J. G. A Benchmark for
1127   Chromatin Binding Measurements in Live Cells. *Nucleic Acids Res* **2012**, *40* (15), E119.
1128   https://doi.org/10.1093/nar/gks701.

1129   (22)   Swinstead; Miranda; Paakinaho; Baek; Goldstein; Hawkins; Karpova; Ball; Mazza;
1130   Lavis; et al. Steroid Receptors Reprogram FoxA1 Occupancy through Dynamic
1131   Chromatin Transitions. *Cell* **2016**, *165* (3), 593–605.
1132   https://doi.org/10.1016/j.cell.2016.02.067.

1133   (23)   Presman; Ball; Paakinaho; Grimm; Lavis; Karpova; Hager. Quantifying Transcription
1134   Factor Binding Dynamics at the Single-Molecule Level in Live Cells. *Methods* **2017**,
1135   *123*, 76–88. https://doi.org/10.1016/j.ymeth.2017.03.014.

1136   (24)   Izeddin, I. Single-Molecule Tracking in Live Cells Reveals Distinct Target-Search
1137   Strategies of Transcription Factors in the Nucleus. *Elife* **2014**, *3*, 26–87.
1138   https://doi.org/10.7554/ELIFE.02230.

1139   (25)   Hansen, A. S.; Woringer, M.; Grimm, J. B.; Lavis, L. D.; Tjian, R.; Darzacq, X. Robust
1140   Model-Based Analysis of Single-Particle Tracking Experiments with Spot-On. *Elife*
1141   **2018**, *7*. https://doi.org/10.7554/eLife.33125.

1142   (26)   Ahmad; Kumar. Steroid Hormone Receptors in Cancer Development: A Target for
1143   Cancer Therapeutics. *Cancer Lett* **2011**, *300* (1), 1–9.
1144   https://doi.org/10.1016/j.canlet.2010.09.008.

1145   (27)   Saha, S.; Dey, S.; Nath, S. Steroid Hormone Receptors: Links With Cell Cycle
1146   Machinery and Breast Cancer Progression. *Front Oncol* **2021**, *11*.
1147   https://doi.org/10.3389/fonc.2021.620214.

(28)   Papageorgiou, L.; Shalzi, L.; Efthimiadou, A.; Bacopoulou, F.; Chrousos, G.; Eliopoulos,
E.; Vlachakis, D. Conserved Functional Motifs of the Nuclear Receptor Superfamily as
Potential Pharmacological Targets. *International Journal of Epigenetics* **2021**, *1* (2).
https://doi.org/10.3892/ije.2021.3.
(29)   Lu, Y.; Liu, W. Selective Estrogen Receptor Degraders (SERDs): A Promising Strategy
for Estrogen Receptor Positive Endocrine-Resistant Breast Cancer. *J Med Chem* **2020**, *63*
(24), 15094–15114. https://doi.org/10.1021/acs.jmedchem.0c00913.
(30)   Quatrini, L.; Ugolini, S. New Insights into the Cell- and Tissue-Specificity of
Glucocorticoid Actions. *Cell Mol Immunol* **2021**, *18* (2), 269–278.
https://doi.org/10.1038/s41423-020-00526-2.
(31)   Guan; Zhou; Hafner; Blake; Chalouni; Chen; De Bruyn; Giltnane; Hartman;
Heidersbach; et al. Therapeutic Ligands Antagonize Estrogen Receptor Function by
Impairing Its Mobility. *Cell* **2019**, *178* (4), 949–963.
https://doi.org/10.1016/j.cell.2019.06.026.
(32)   Van Royen, M. E.; van Cappellen, W. A.; Geverts, B.; Schmidt, T.; Houtsmuller, A. B.;
Schaaf, M. J. M. Androgen Receptor Complexes Probe DNA for Recognition Sequences
by Short Random Interactions. *J Cell Sci* **2014**, *127* (7), 1406–1416.
https://doi.org/10.1242/jcs.135228.
(33)   van Royen, M. E.; van Cappellen, W. A.; de Vos, C.; Houtsmuller, A. B.; Trapman, J.
Stepwise Androgen Receptor Dimerization. *J Cell Sci* **2012**, *125* (8), 1970–1979.
https://doi.org/10.1242/jcs.096792.
(34)   Natwick, D. E.; Collins, S. R. Optimized ILID Membrane Anchors for Local Optogenetic
Protein Recruitment. *ACS Synth Biol* **2021**, *10* (5), 1009–1023.
https://doi.org/10.1021/acssynbio.0c00511.
(35)   Heckert, A.; Dahal, L.; Tjian, R.; Darzacq, X. Recovering Mixtures of Fast-Diffusing
States from Short Single-Particle Trajectories. *Elife* **2022**, *11*.
https://doi.org/10.7554/eLife.70169.
(36)   Wagh, K.; Stavreva, D. A.; Jensen, R. A. M.; Paakinaho, V.; Fettweis, G.; Schiltz, R. L.;
Wüstner, D.; Mandrup, S.; Presman, D. M.; Upadhyaya, A.; et al. Dynamic Switching of
Transcriptional Regulators between Two Distinct Low-Mobility Chromatin States. *Sci*
Adv **2023**, *9* (24). https://doi.org/10.1126/sciadv.ade1122.
(37)   Liberzon, A.; Birger, C.; Thorvaldsdóttir, H.; Ghandi, M.; Mesirov, J. P.; Tamayo, P. The
Molecular Signatures Database (MSigDB) Hallmark Gene Set Collection. *Cell Syst* **2015**,
1 (6), 417–425. https://doi.org/10.1016/j.cels.2015.12.004.
(38)   Zhang, J.-H.; Chung, T. D. Y.; Oldenburg, K. R. A Simple Statistical Parameter for Use
in Evaluation and Validation of High Throughput Screening Assays. *J Biomol Screen*
1999, *4* (2), 67–73. https://doi.org/10.1177/108705719900400206.
(39)   Kuiper, G. G. J. M.; Carlsson, B.; Grandien, K.; Enmark, E.; Häggblad, J.; Nilsson, S.;
Gustafsson, J.-Å. Comparison of the Ligand Binding Specificity and Transcript Tissue
Distribution of Estrogen Receptors α and β. *Endocrinology* **1997**, *138* (3), 863–870.
https://doi.org/10.1210/endo.138.3.4979.
(40)   Goyeneche, A. A.; Car n, R. n W.; Telleria, C. M. Mifepristone Inhibits Ovarian Cancer
Cell Growth In Vitro and In Vivo. *Clin Cancer Res* **2007**, *13* (11), 3370–3379.
https://doi.org/10.1158/1078-0432.CCR-07-0164.
(41)   Rajaram, P.; Rivera, A.; Muthima, K.; Olveda, N.; Muchalski, H.; Chen, Q.-H. Second-
Generation Androgen Receptor Antagonists as Hormonal Therapeutics for Three Forms

of Prostate Cancer. *Molecules* **2020**, *25* (10). https://doi.org/10.3390/molecules25102448.

(42) Miner, J. N.; Tyree, C.; Hu, J.; Berger, E.; Marschke, K.; Nakane, M.; Coghlan, M. J.;
Clemm, D.; Lane, B.; Rosen, J. A Nonsteroidal Glucocorticoid Receptor Antagonist. *Mol*
Endocrinol **2003**, *17* (1), 117–127. https://doi.org/10.1210/me.2002-0010.

(43) Liu, Z.; Legant, W. R.; Chen, B.-C.; Li, L.; Grimm, J. B.; Lavis, L. D.; Betzig, E.; Tjian,
R. 3D Imaging of Sox2 Enhancer Clusters in Embryonic Stem Cells. *Elife* **2014**, *3*,
E4236. https://doi.org/10.7554/eLife.04236.

(44) Elf, J.; Li, G.-W.; Xie, X. S. Probing Transcription Factor Dynamics at the Single-
Molecule Level in a Living Cell. *Science* **2007**, *316* (5828), 1191–1194.
https://doi.org/10.1126/science.1141967.

(45) Chen, Y.-C.; Yu, J.; Metcalfe, C.; De Bruyn, T.; Gelzleichter, T.; Malhi, V.; Perez-
Moreno, P. D.; Wang, X. Latest Generation Estrogen Receptor Degraders for the
Treatment of Hormone Receptor-Positive Breast Cancer. *Expert Opin Investig Drugs*
2022, *31* (6), 515–529. https://doi.org/10.1080/13543784.2021.1983542.

(46) Kahraman, M.; Govek, S. P.; Nagasawa, J. Y.; Lai, A.; Bonnefous, C.; Douglas, K.;
Sensintaffar, J.; Liu, N.; Lee, K.; Aparicio, A.; et al. Maximizing ER-α Degradation
Maximizes Activity in a Tamoxifen-Resistant Breast Cancer Model: Identification of
GDC-0927. *ACS Med Chem Lett* **2019**, *10* (1), 50–55.
https://doi.org/10.1021/acsmedchemlett.8b00414.

(47) Wells, C. I.; Vasta, J. D.; Corona, C. R.; Wilkinson, J.; Zimprich, C. A.; Ingold, M. R.;
Pickett, J. E.; Drewry, D. H.; Pugh, K. M.; Schwinn, M. K.; et al. Quantifying CDK
Inhibitor Selectivity in Live Cells. *Nature Communications* **2020**, *11* (1).
https://doi.org/10.1038/s41467-020-16559-0.

(48) Wang; Wu; Wu; Chen; Zou; Wang; Wu; Hu; Jiang; Liu; et al. Discovery of 4-(((4-(5-
Chloro-2-(((1s,4s)-4-((2-Methoxyethyl)Amino)Cyclohexyl)Amino)Pyridin-4-Yl)Thiazol-
2-Yl)Amino)Methyl)Tetrahydro-2H-Pyran-4-Carbonitrile (JSH-150) as a Novel Highly
Selective and Potent CDK9 Kinase Inhibitor. *Eur J Med Chem* **2018**, *158*, 896–916.
https://doi.org/10.1016/j.ejmech.2018.09.025.

(49) Olson Calla M; Jiang Baishan; Erb Michael A; Liang Yanke; Doctor Zainab M; Zhang
Zinan; Zhang Tinghu; Kwiatkowski Nicholas; Boukhali Myriam; Green Jennifer L; et al.
Pharmacological Perturbation of CDK9 Using Selective CDK9 Inhibition or Degradation.
Nature Chemical Biology **2018**, *14* (2), 163–170. https://doi.org/10.1038/nchembio.2538.

(50) Parry, D.; Guzi, T.; Shanahan, F.; Davis, N.; Prabhavalkar, D.; Wiswell, D.; Seghezzi,
W.; Paruch, K.; Dwyer, M. P.; Doll, R.; et al. Dinaciclib (SCH 727965), a Novel and
Potent Cyclin-Dependent Kinase Inhibitor. *Mol Cancer Ther* **2010**, *9* (8), 2344–2353.
https://doi.org/10.1158/1535-7163.MCT-10-0324.

(51) Menezes, D. L.; Taverna, P.; Jensen, M. R.; Abrams, T.; Stuart, D.; Yu, G. K.; Duhl, D.;
Machajewski, T.; Sellers, W. R.; Pryer, N. K.; et al. The Novel Oral Hsp90 Inhibitor
NVP-HSP990 Exhibits Potent and Broad-Spectrum Antitumor Activities In Vitro and In
Vivo. *Mol Cancer Ther* **2012**, *11* (3), 730–739. https://doi.org/10.1158/1535-7163.MCT-
11-0667.

(52) McCleese, J. K.; Bear, M. D.; Fossey, S. L.; Mihalek, R. M.; Foley, K. P.; Ying, W.;
Barsoum, J.; London, C. A. The Novel HSP90 Inhibitor STA□1474 Exhibits Biologic
Activity against Osteosarcoma Cell Lines. *Int J Cancer* **2009**, *125* (12), 2792–2801.
https://doi.org/10.1002/ijc.24660.

(53) Bussenius, J.; Blazey, C. M.; Aay, N.; Anand, N. K.; Arcalas, A.; Baik, T.; Bowles, O. J.;

Buhr, C. A.; Costanzo, S.; Curtis, J. K.; et al. Discovery of XL888: A Novel Tropane-
Derived Small Molecule Inhibitor of HSP90. *Bioorg Med Chem Lett* **2012**, *22* (17),
5396–5404. https://doi.org/10.1016/j.bmcl.2012.07.052.

(54) Apsel Beth; Blair Jimmy A; Gonzalez Beatriz; Nazif Tamim M; Feldman Morri E;
Aizenstein Brian; Hoffman Randy; Williams Roger L; Shokat Kevan M; Knight Zachary
A. Targeted Polypharmacology: Discovery of Dual Inhibitors of Tyrosine and
Phosphoinositide Kinases. *Nature Chemical Biology* **2008**, *4* (11), 691–699.
https://doi.org/10.1038/nchembio.117.

(55) Thoreen, C. An ATP-Competitive Mammalian Target of Rapamycin Inhibitor Reveals
Rapamycin-Resistant Functions of MTORC1. *J Biol Chem* **2009**, *284* (12), 8020–8029.
https://doi.org/10.1074/JBC.M900301200.

(56) Yu, K.; Toral-Barza, L.; Shi, C.; Zhang, W.-G.; Lucas, J.; Shor, B.; Kim, J.; Verheijen, J.;
Curran, K.; Malwitz, D. J.; et al. Biochemical, Cellular, and In Vivo Activity of Novel
ATP-Competitive and Selective Inhibitors of the Mammalian Target of Rapamycin.
Cancer Res **2009**, *69* (15), 6232–6240. https://doi.org/10.1158/0008-5472.CAN-09-0299.

(57) Adams, J.; Palombella, V. J.; Sausville, E. A.; Johnson, J.; Destree, A.; Lazarus, D. D.;
Maas, J.; Pien, C. S.; Prakash, S.; Elliott, P. J. Proteasome Inhibitors: A Novel Class of
Potent and Effective Antitumor Agents. *Cancer Res* **1999**, *59* (11), 2615–2622.

(58) Kuhn; Chen; Voorhees; Strader; Shenk; Sun; Demo; Bennett; van Leeuwen; Chanan-
Khan; et al. Potent Activity of Carfilzomib, a Novel, Irreversible Inhibitor of the
Ubiquitin-Proteasome Pathway, against Preclinical Models of Multiple Myeloma. *Blood*
2007, *110* (9), 3281–3290. https://doi.org/10.1182/blood-2007-01-065888.

(59) Mroczkiewicz, M.; Winkler, K.; Nowis, D.; Placha, G.; Golab, J.; Ostaszewski, R.
Studies of the Synthesis of All Stereoisomers of MG-132 Proteasome Inhibitors in the
Tumor Targeting Approach. *J Med Chem* **2010**, *53* (4), 1509–1518.
https://doi.org/10.1021/jm901619n.

(60) Stenoien David L.; Patel Kavita; Mancini Maureen G.; Dutertre Martin; Smith Carolyn
L.; O'Malley Bert W.; Mancini Michael A. FRAP Reveals That Mobility of Oestrogen
Receptor-a Is Ligand- and Proteasome-Dependent. *Nature cell biology* **2001**, *3* (1), 15–
23. https://doi.org/10.1038/35050515.

(61) Franco, L. C.; Morales, F.; Boffo, S.; Giordano, A. CDK9: A Key Player in Cancer and
Other Diseases. *J Cell Biochem* **2017**, *119* (2), 1273–1284.
https://doi.org/10.1002/jcb.26293.

(62) Shah, Y. M.; Rowan, B. G. The Src Kinase Pathway Promotes Tamoxifen Agonist
Action in Ishikawa Endometrial Cells through Phosphorylation-Dependent Stabilization
of Estrogen Receptor Promoter Interaction and Elevated Steroid Receptor Coactivator 1
Activity. *Mol Endocrinol* **2005**, *19* (3), 732–748. https://doi.org/10.1210/ME.2004-0298.

(63) Anbalagan; Rowan. Estrogen Receptor Alpha Phosphorylation and Its Functional Impact
in Human Breast Cancer. *Mol Cell Endocrinol* **2015**, *418*, 264–272.
https://doi.org/10.1016/j.mce.2015.01.016.

(64) Liang, J.; Zbieg, J. R.; Blake, R. A.; Chang, J. H.; Daly, S.; DiPasquale, A. G.; Friedman,
L. S.; Gelzleichter, T.; Gill, M.; Giltnane, J. M.; et al. GDC-9545 (Giredestrant): A Potent
and Orally Bioavailable Selective Estrogen Receptor Antagonist and Degrader with an
Exceptional Preclinical Profile for ER+ Breast Cancer. *J Med Chem* **2021**, *64* (16),
11841–11856. https://doi.org/10.1021/acs.jmedchem.1c00847.

(65) Wardell, S. E.; Ellis, M. J.; Alley, H. M.; Eisele, K.; VanArsdale, T.; Dann, S. G.; Arndt,

K. T.; Primeau, T.; Griffin, E.; Shao, J.; et al. Efficacy of SERD/SERM Hybrid-CDK4/6
Inhibitor Combinations in Models of Endocrine Therapy–Resistant Breast Cancer. *Clin*
Cancer Res **2015**, *21* (22), 5121–5130. https://doi.org/10.1158/1078-0432.CCR-15-0360.

(66)   Finn, R. S.; Aleshin, A.; Slamon, D. J. Targeting the Cyclin-Dependent Kinases (CDK)
4/6 in Estrogen Receptor-Positive Breast Cancers. *Breast Cancer Research* **2016**, *18* (1).
https://doi.org/10.1186/s13058-015-0661-5.

(67)   Chu, Q.; Wang, J.; Du, Y.; Zhou, T.; Shi, A.; Xiong, J.; Ji, W.-K.; Deng, L. Oligomeric
CHMP7 Mediates Three-Way ER Junctions and ER-Mitochondria Interactions. *Cell*
Death and Differentiation **2023**, *30* (1), 94–110. https://doi.org/10.1038/s41418-022-
01048-2.

(68)   Sergé Arnauld; Bertaux Nicolas; Rigneault Hervé; Marguet Didier. Dynamic Multiple-
Target Tracing to Probe Spatiotemporal Cartography of Cell Membranes. *Nature*
Methods **2008**, *5* (8), 687–694. https://doi.org/10.1038/nmeth.1233.

(69)   Levenberg, K. A Method for the Solution of Certain Non-Linear Problems in Least
Squares. *Quarterly of applied mathematics* **1944**, *2* (2), 164–168.
https://doi.org/10.1090/qam/10666.

(70)   Donald W. Marquardt. An Algorithm for Least-Squares Estimation of Nonlinear
Parameters. *Journal of the Society for Industrial and Applied Mathematics* **1963**, *11* (2),
431–441. https://doi.org/10.1137/0111030.

(71)   Laurence Ted A; Chromy Brett A. Efficient Maximum Likelihood Estimator Fitting of
Histograms. *Nature Methods* **2010**, *7* (5), 338–339. https://doi.org/10.1038/nmeth0510-
338.

(72)   Smith Carlas S; Joseph Nikolai; Rieger Bernd; Lidke Keith A. Fast, Single-Molecule
Localization That Achieves Theoretically Minimum Uncertainty. *Nature Methods* **2010**,
No. 5, 373–375. https://doi.org/10.1038/nmeth.1449.

(73)   Parthasarathy Raghuveer. Rapid, Accurate Particle Tracking by Calculation of Radial
Symmetry Centers. *Nature Methods* **2012**, *9* (7), 724–726.
https://doi.org/10.1038/nmeth.2071.

(74)   Sbalzarini, I. F.; Koumoutsakos, P. Feature Point Tracking and Trajectory Analysis for
Video Imaging in Cell Biology. *J Struct Biol* **2005**, *151* (2), 182–195.
https://doi.org/10.1016/j.jsb.2005.06.002.

(75)   Olaf Ronneberger; Philipp Fischer; Thomas Brox. U-Net: Convolutional Networks for
Biomedical Image Segmentation. In *Medical Image Computing and Computer-Assisted*
Intervention – MICCAI 2015; Springer International Publishing, 2015; pp 234–241.
https://doi.org/10.1007/978-3-319-24574-4_28.

(76)   Bemis, G. W.; Murcko, M. A. The Properties of Known Drugs. 1. Molecular
Frameworks. *J Med Chem* **1996**, *39* (15), 2887–2893. https://doi.org/10.1021/jm9602928.

(77)   Garcia, D. A.; Fettweis, G.; Presman, D. M.; Paakinaho, V.; Jarzynski, C.; Upadhyaya,
A.; Hager, G. L. Power-Law Behavior of Transcription Factor Dynamics at the Single-
Molecule Level Implies a Continuum Affinity Model. *Nucleic acids research* **2021**, *49*
(12), 6605–6620. https://doi.org/10.1093/nar/gkab072.

(78)   Reisser, M.; Hettich, J.; Kuhn, T.; Popp, A. P.; Große-Berkenbusch, A.; Gebhardt, J. C.
M. Inferring Quantity and Qualities of Superimposed Reaction Rates from Single
Molecule Survival Time Distributions. *Sci Rep* **2020**, *10* (1).
https://doi.org/10.1038/s41598-020-58634-y.

(79)   Stirling, D. R.; Swain-Bowden, M. J.; Lucas, A. M.; Carpenter, A. E.; Cimini, B. A.;

Goodman, A. CellProfiler 4: Improvements in Speed, Utility and Usability. *BMC*
Bioinformatics **2021**, *22* (1). https://doi.org/10.1186/s12859-021-04344-9.
1334

A State occupations for each cell line

Histone H2B
>0.1 µm²/sec

f_{bound} = 0.9

CaaX
0.5-5 µm²/sec

free Halo
7-30 µm²/sec

State Occupation

Diffusion Coefficient (µm²/sec)

B Mixture of Histone H2B, CaaX, and Free Halo

JF$_{549}$

Hoechst

1: Free Halo

2: CaaX

3: Histone H2B

D

Cell number

State Occupation

Diffusion Coefficient (µm²/sec)

C Single-cell state distribution

Histone H2B   CaaX   Free Halo

State Occupation

Diffusion Coefficient (µm²/sec)

E Ensemble of cell mixture

State Occupation

Diffusion Coefficient (µm²/sec)

Figure 1 - figure supplement 1

A

Reagent handling | Super-resolution imaging | Particle detection and tracking | Image processing and QC | Feature extraction and storage | Biological interpretation

B

Cell Incubator
Plate Carousel
Microscopes 1 & 2
Microscopes 3 & 4
Robotic Arm
Computer
Plate Sealer
Plate Washer
Echo disepnser

C

High quality images | Debris | Out of focus | Uneven illumination | Not enough cells

Figure 1 - figure supplement 2

Figure 2

A

B

estradiol 2-hydroxytestosterone progesterone dexamethasone

Figure 3

5067 Compound screened against Estrogen Receptor

Figure 3 - figure supplement 1

Figure 3 - figure supplement 2

Figure 3 - figure supplement 3

Figure 3 - figure supplement 4

Figure 4

Figure 4 - figure supplement 1

A Diffusing states only (> 0.1 μm²/sec)

fulvestrant | 4-hydroxytamoxifen | GDC-0810 | AZD9496 | GDC-0927

(Norm. State Occupation vs Diffusion Coefficient (μm²/sec))

B

Compound	X_o (min)	$T_{1/2}$ (min)	E_{Max}
Estradiol	0	1.6	0.53
Fulvestrant	0.3	3.9	0.55
Tamoxifen	0	1.7	0.35
GDC-0810	0.6	4.3	0.43
GDC-0927	0.2	3.1	0.51
AZD9496	0.6	2.5	0.41

C

Estradiol | Fulvestrant

Untreated, 1 min, 3 min, 5 min, 7 min, 9 min

(Norm. State Occupation vs Diffusion Coefficient (μm²/sec))

D

Δf_{bound}

AR Antagonists | GR Antagonists

Estradiol, Bicalutamide, Enzalutamide, Cyproterone Acetate, Apalutamide, RAD140, RU58841, Darolutamide, Loteprednol etabonate, Hydrocortisone, Methylprednisolone, Mometasone furoate, Diflorasone, Hydrocortisone 21-acetate, Methylprednisolone hemisuccinate, Hydrocortisone butyrate

* p < 0.01

E

Fraction of surviving tracks vs Track length (sec)

- fulvestrant
- 4-hydroxytamoxifen
- GDC-0810
- AZD9496
- GDC-0927
- DMSO

F

$$k^*_{on,inferred} = \frac{f_{bound} \cdot k^*_{off}}{1 - f_{bound}}$$

Compound	k^*_{off} (s⁻¹) *lower bound*	Rel. to DMSO	f_{bound}	$k_{on, inferred}$ *lower bound*
DMSO	0.17	1	0.34	0.08
Estradiol	0.11*	1.60	0.87	*0.73*
Fulvestrant	0.11*	1.55	0.85	*0.62*
Tamoxifen	0.12	1.38	0.77	0.40
GDC-0810	0.14	1.27	0.78	0.50
GDC-0927	0.09*	1.82	0.87	*0.60*
AZD9496	0.12	1.39	0.88	0.88

Figure 5

Figure 5 - figure supplement 1

Figure 6

Figure 6 - figure supplement 3