Loss of secondary motor cortex neurons in chronic

neuropathic pain

Amrita Das Gupta[a,#], Hongwei Zheng[a,#], Livia Asan[a,b], Jennifer John[a], Sanjeev Kaushalya[c,d], Carlo

A. Beretta[a,c], Rohini Kuner[c], Johannes Knabbe[a,e], Thomas Kuner[a]

a Department of Functional Neuroanatomy, Institute for Anatomy and Cell Biology, Heidelberg

University, Germany.

b Current address: Department of Neurology and Center for Translational Neuro- and Behavioral

Sciences (C-TNBS), University Hospital Essen, Germany.

c Department of Molecular Pharmacology, Pharmacology Institute, Heidelberg University,

Germany.

d Current address: Cold-Spring-Harbour Laboratory, Neuroscience Center

e Department of General Psychiatry, Centre for Psychosocial Medicine, Heidelberg University,

Germany.

# These authors contributed equally.

Correspondence: thomas.kuner@uni-heidelberg.de

Abstract

Chronic neuropathic pain is associated with structural plasticity of the brain on different spatial

scales, yet, on the mesoscopic scale of tissue composition, little is known. Here, we determined

the cellular composition of cortical areas and structural variability of entire neurons during

chronic neuropathic pain using longitudinal in vivo two-photon microscopy and behavioral

assessment. When monitoring cell type composition in response to spared-nerve injury in 84

cortical volumes containing ~25000 cells each, we found neuronal loss in the secondary motor

cortex region M2 immediately adjacent to the cingulate cortex already one week after surgery.

Loss of mostly interneurons was also evident when monitoring individual M2 neurons over

time. This neuronal loss was preceded by decreased spine density and loss of distal dendritic

branches. In conclusion, our work delineates M2 as a novel site and neuronal loss as a so far

underappreciated mechanism underlying chronic neuropathic pain states.

Introduction

Chronic neuropathic pain has been shown to be associated with structural alterations of the

brain, ranging from macroscopic grey-matter-volume changes (Apkarian et al. 2004; Seminowicz

et al. 2009; Bilbao et al. 2018; Silva and Seminowicz 2019) to microscopic alterations on the

cellular level (Metz et al. 2009; Kim et al. 2011; Kim and Nabekura 2011; Kassem et al. 2013). In

principle, structural neocortical changes associated with chronic pain could be based on a

multitude of mechanisms. The most commonly discussed mechanism concerns structural

neuronal plasticity, i.e. changes in dendritic architecture and spine density (Holtmaat and

Svoboda 2009). Such changes have been shown in many cortical areas to underly functional

changes in multifarious physiological and pathophysiological functions such as learning and

memory, fear processing, addiction and others (Forrest et al. 2018), and can be viewed as a

general disease-relevant mechanism. Changes in the microenvironment of the neuropil

including extracellular space regulation (Mascio et al. 2022), ion homeostasis and others could

also undergo long-lasting adaptive changes and affect neuronal communication. Such changes

would imply astroglial, microglial, oligodendroglial functions and likely also endothelial cells and

pericytes. Plasticity could arise from changes in structure and function of these cells, but also by

changes in their distribution pattern and number. The latter is well known in neuroinflammation

and neurodegeneration, where astrogliosis and proliferation of microglia is particularly well

established (Salter and Stevens 2017; Ji et al. 2019). Hence, to address structural mechanisms

underlying chronic neuropathic pain states, monitoring the cellular composition of cortical pain

areas after induction of chronic pain is a promising avenue to follow. We recently developed an

approach called "NuClear" that allows longitudinal, comprehensive, monitoring of cell type-

composition in living animals (Das-Gupta et al. 2022). The resulting 3D matrix of identified cells

and their location can be repetitively acquired from the identical cortical area and changes in

number and location of any cell type can be identified, for example after establishing chronic

neuropathic pain using the spared-nerve injury (SNI) model (Decosterd and Woolf 2000).

Corresponding changes in pain behavior can be interspersed and allow correlations with

structural changes.

Cortical pain processing occurs in multiple brain regions also referred to as the pain

matrix (Tracey and Mantyh 2007), with each area contributing specialized functions. The

cingulate cortex (CC) and its subdivisions is consistently reported to represent emotional and

motor aspects of pain perception (Bliss et al. 2016; Kuner and Kuner 2021) and confers pain

hypersensitivity via a direct pathway to the posterior insula (Tan et al. 2017). Studies of

dendritic structural plasticity have so far addressed the somatosensory (S1) and medial

prefrontal cortices (Metz et al. 2009; Kim et al. 2011; Kim and Nabekura 2011), yet has not been

studied much in the CC.

Here, we attempted to address structural changes of the CC in response to SNI using

NuClear and imaging of individual neurons in an in vivo longitudinal study design. In response to

SNI, we surprisingly found a loss of neurons in the medial M2 cortex, immediately lateral to the

CC. Most of the lost neurons were interneurons, suggesting that loss of inhibition leads to

stronger activation of local pyramidal neurons. Neurons prone to die showed a loss of distal

dendritic branches and a loss of dendritic spines, suggesting that changes in synaptic inputs

may, together with other factors, induce neuronal cell death. In conclusion, we identified medial

M2 as a novel player in chronic pain and a loss of M2 neurons as a mechanism previously not

considered for chronic pain.

Results

We assessed structural changes associated with nerve injury in the cingulate cortex (CC) and

adjoining secondary motor cortex (M2) on two levels: (1) changes in tissue composition

represented by the number of major brain cell types within large two-photon (2P) imaging

volumes (~0.25 mm^3) and (2) cellular and subcellular changes on the level of entire neurons and

their dendritic spines (Fig. 1A). Both levels were investigated in a longitudinal paradigm

combining SNI, two-photon in vivo imaging and pain behavior (Fig. 1B). Tissue composition was

determined by imaging Histon2B-GFP-labelled nuclei (Asan et al. 2021) and subsequent

classification of the cell type derived from specific features of the nucleus (Das-Gupta et al.

2022)(Das-Gupta et al. 2022). Subcellular changes were investigated by repetitively imaging

individually labeled or small groups of labeled neurons in their entirety.

SNI reliably induced chronic neuropathic pain states

As a foundation for the structural analysis aimed at by this study, we first report the pain

behavioral readouts for both cohorts of mice studied. All mice underwent sensory withdrawal

testing using von Frey filaments at the timepoints relative to the SNI or sham surgery as

indicated in Figure 1B. Allodynia and hypersensitivity were detected when testing the left hind

foot on which surgery was performed (Fig. 1C, Supp. Fig. 1A), but not when testing the right side

(Supp. Fig. 1B). Furthermore, cold plate withdrawal was found to be significantly different in

SNI-treated mice (Fig. 1D). To test the emotional aspect of pain, we assessed behavior of mice in

the open field. SNI-treated mice showed longer lasting immobility, consistent with the presence

of a chronic pain state (Fig. 1 E).

Tissue composition in volumes encompassing cingulate cortex and adjacent M2 regions

After establishing the development of a chronic pain state in all mice tested, we now focus on

cohort 1 mice and tissue composition analysis. We defined four 700 µm x 700 µm x 500 µm

large imaging volumes in both hemispheres that were placed as close to the midline as

permitted by the sagittal sinus, a structure that obstructs direct 2P imaging of brain areas

residing below it (Fig. 2A, Supp. Fig. 2A). Owing to a variable position of large bridging veins

towards the sagittal sinus, imaging volumes could not be placed at identical positions in each

mouse (indicated by light grey squares in Fig. 2A). We took advantage of this situation by

dividing the imaging volumes into cingulate/M2 and M2/M1 volumes, the latter serving as

controls for the expected cingulate-specific effects. Nucleus imaging was done before, 1 week

and 12 weeks after SNI surgery and yielded 20000 to 30000 cell nuclei per imaging volume (Fig.

2B). Using the NuClear approach (Das-Gupta et al. 2022), a method allowing for non-biased

neuronal network-based identification of all major cell types residing within the imaging volume

solely relying on a set of radiomic features of the cell nucleus (Das-Gupta et al. 2022)we

determined 3D matrices of the localization of all neurons, astroglia, oligodendroglia, microglia

and endothelial cells present in the volume (Fig. 2C). When applying this analysis to all 84

imaging volumes taken from 21 mice, encompassing >2 million cells, no significant changes in

cell numbers were found (Fig. 2D). As expected, the most frequent cell type are neurons which

were about as abundant as all other cells taken together. When re-slicing the large volumes into

coronal sections, the cortical layering becomes evident and the different spatial distributions of

the cell types can be appreciated. For example, neurons are organized in layers, microglia and

astroglia are evenly distributed at rather constant nearest-neighbor distances (Fig. 2E).

Chronic neuropathic pain associated with changes in cell number contralateral to the lesion

We next compared the number of cells in each category at different timepoints post-SNI or

sham surgery. Because the medial imaging volumes contain both cingulate cortex and M2 cortex

(Fig. 3, center), we digitally separated the volumes into a medial wedge representing the

cingulate cortex (black) and a lateral wedge representing the adjoining M2 cortex (light blue).

We identified several significant changes in cell number on the contralateral side to the lesion,

where such changes are expected to occur (Fig. 3B,D). In contrast, no differences between SNI

and sham were detected in ipsilateral volumes (Fig. 3A,C). Hence, we now focus on changes in

cell number found for the different cell types on the contralateral side specifically affected by

the lesion.

Loss of neurons in the medial part of M2 associated with chronic neuropathic pain

A decreased number of neurons was evident already one week after SNI in the medial M2 and

the CC (Fig. 3B,D). This decrease persisted throughout week 12 in M2, but not in CC. The weakly

significant difference in CC may arise from an imperfect separation of CC versus medial M2

regions, with imaging volumes analysed for CC containing a certain fraction of M2 territory. The

much stronger and highly significant difference in the number of neurons found in medial M2

may thereby 'contaminate' the CC imaging volume. Hence, our data suggests that SNI causes a

persistent loss of neurons in the contralateral medial M2 cortex, and to some extent possibly

also in the CC. This decrease may contribute to the chronic pain state.

Changes in the number of non-neuronal cells

We further found a decrease in the number of astroglia in medial M2 after one week that was,

however, no longer evident 12 weeks after the lesion (Fig. 3B). Also microglia show a trend to be

decreased in M2. We further detected a significant increase in oligodendroglia in the CC after 12

weeks, while the large variability in M2 prevented a similar readout (Fig. 3B). This increase in

oligodendrocyte number may suggest an activity-dependent differentiation of oligodendrocyte

precursor cells (Faria et al. 2021). Lastly, we found a decrease in endothelial cell number already

at week one that persisted until week 12 (Fig. 3B). This may correspond to a reduced capillary

density in medial M2 and could go along with a reduced supply of oxygen and nutrients, both of

which could be related to neuronal loss.

Taken together, our data introduce the medial part of M2 as a new cortical region

involved in pain processing and establishes neocortical neuronal cell death as a cellular

mechanism underlying chronic neuropathic pain.

Longitudinal imaging of individually labeled neurons confirms neuronal loss in medial M2

To validate the findings obtained using the NuClear approach and to learn more about structural

changes of individual neurons after SNI surgery, we sparsely labeled neurons within the medial

M2 region (Supp. Fig. 3A) and repetitively imaged the same neurons over a time course of 16

weeks (see Methods). Figure 4A shows eight labeled and fully reconstructed as well as spatially

registered neurons at basal state prior to SNI and at 1, 9 and 16 weeks thereafter. Evidently, only

three neurons persist for 16 weeks (N6-8), but already after one week neuron 1 disappeared,

followed by neurons 2, 3 and 4 at week 9 (further examples in Supp. Fig 3). On average,

approximately 40% of labeled neurons were lost within 16 weeks after SNI (Fig. 4C, red curve).

In sharp contrast, nearly all neurons survived over the entire imaging period in sham-treated

mice (Fig. 4B, D), indicating that the loss of mGFP expression with time or toxicity triggered by

high level mGFP expression over extended time periods are unlikely explanations for neuronal

loss. Therefore, we conclude that neurons in the medial M2 region are lost by some mechanism

of cell death triggered by SNI.

Interneurons are lost more frequently than pyramidal neurons

A differential loss of interneurons versus pyramidal neurons could have a strong impact on

network function, with a loss of interneurons reducing local inhibition and thus making

pyramidal cells more active, and a loss of pyramidal neurons attenuating the output of the

neuronal ensemble in the medial M2. To address this, we sorted the imaged neurons according

to morphological parameters into interneurons and pyramidal neurons (see Methods). Our

analysis revealed that interneurons are more prone for cell death than pyramidal neurons (Fig.

4D). Hence, a more pronounced loss of interneurons predicts a reduction in inhibitory drive

within the local M2 network.

Structural changes of the dendritic tree precede neuronal loss after SNI

To assess if neurons prone to die undergo some prodromal state of structural changes, we

analysed their entire dendritic tree including their postsynaptic spines (see Supp. Fig. 4 for

illustration and Methods Supplement for detailed description). For example, neuron 4 (already

displayed in Figure 4) underwent a pronounced loss in spine density already within the first

week after SNI and barely retained any spines one week later (Figure 5A), to finally disappear

before week four. In other neurons we found rearrangements of the dendritic tree. Figure 5B

shows neurons 2 and 5 (as in Fig. 4). Each panel shows both neurons at the earlier time point in

red and at the later time point in green, hence, structures appearing in yellow exist at both time

points while red alone indicates loss of a structure. The pink rectangle marks a dynamic region

while the blue rectangle delineates a rather stable region. While dendritic structures are mostly

overlapping between basal and day 3 (Fig. 5B1), a loss of distal dendritic branches is evident at

week 1 (Fig. 5B2) and progresses further in the delineated area (Fig. 5B3,4). At week 7, neuron 2

has disappeared (Fig. 5B5,6). Similar observations were made in all neurons analyzed (Supp. Fig.

5). In conclusion, before disappearing, neurons undergo structural changes ranging from a loss

of spines to a loss of dendritic branches. These structural changes may reflect alterations in

synaptic inputs, possibly triggered by SNI.

Discussion

Using NuClear, an unbiased and comprehensive tissue analysis approach, we monitored the

cellular composition of the cingulate and M2 cortices after SNI. While monitoring a large tissue

volume (in total approximately 21 mm^3) distributed over the medial part of both hemispheres,

we found significant changes in cellular composition solely within a small subvolume

encompassing the medial part of M2. The most prominent change was a reduction of the

number of neurons in medial M2, but also changes in non-neuronal cells, in particular

endothelial cells, were observed. The loss of neurons could be validated by longitudinally

imaging individually labeled neurons in medial M2. Up to 40% of labelled neurons disappeared,

mostly affecting interneurons. Neurons prone to die lost dendritic branches and spines

beforehand. In conclusion, our work adds the medial M2 cortex as a novel player in chronic pain

processing and suggests selective loss of cortical neurons as a mechanism of chronic pain.

What is particular about the medial M2?

The medial M2 in rodents is considered to belong to the mPFC and is thought to mediate

context-dependent and choice-related decisions, to map sensory cues to motor actions, and to

potentially also contribute to depressive-like behavior (Barthas and Kwan 2017). Consistent with

these diverse functions, this area receives abundant multimodal sensory inputs and projects to

sensory, parietal and retrosplenial cortices (Barthas and Kwan 2017). This constellation of

functionalities and in particular connectivity makes a role of the medial M2 in pain processing

highly plausible, consistent with reports on patients suffering from complex regional pain

syndrome (Maihöfner et al. 2007; Jung et al. 2019) However, the detailed connectivity of cells in

medial M2 lost after SNI, both in terms of afferent and efferent connections, changes in activity

in these projections and their impact on pain behavior await further elucidation.

A neurodegenerative mechanisms contributing to chronic pain?

Our finding of neuronal loss starting as early as one week after SNI suggests that a mechanism

of neuronal cell death may get triggered by SNI-induced changes in activity of neurons upstream

of M2. The distinct reduction in spine density and loss of distal dendritic branches observed

within one week after SNI further supports the idea that the loss of certain synaptic inputs or

their altered activity patterns are involved in early steps, thereby eventually resulting in a

denervation of distinct M2 neurons. Such loss of afferent input has been reported to result in

cell loss (Mostafapour et al. 2000; Inquimbert et al. 2018). Persistent changes in activity, either

silence or overactivity, are known to trigger different mechanisms of cell death (Chi et al. 2018;

Fricker et al. 2018; Moujalled et al. 2021). Which of these are active in SNI-induced loss of M2

neurons remains to be investigated. Alternatively, or synergistically, the neuronal

microenvironment defined by glia and endothelial cells might respond to the SNI impact and

affect neuron survival. A decreased number of endothelial cells may reflect changes in

microvasculature that in turn may negatively affect neuronal fidelity. In conclusion, our work

implies neurodegeneration as a mechanism underlying chronic pain states, thereby extending a

concept so far limited to the peripheral nervous system (Reichling and Levine 2011) to

neocortical circuits.

How can neuronal loss contribute to the development of a chronic pain state?

If neurons get lost, they can no longer contribute to function, yet chronic pain is an active

percept generated by as yet unknown interactions of brain areas. As a simple approximation, an

increased firing of excitatory M2 output neurons could elicit activity in other areas of the pain

matrix and thereby contribute to the pain percept. Likewise, disinhibition via long-range

inhibitory connections could achieve the same. Given that we found a more pronounced loss of

interneurons within M2, the surviving pyramidal neurons may become more active and thereby

contribute to eliciting a pain percept. Interestingly, such local disinhibition in the anterior

cingulate network has been described after nerve injury (Blom et al. 2014). Likewise,

hyperactive pyramidal neurons in the anterior cingulate have been linked to neuropathic pain

(Zhao et al. 2018). In conclusion, while published work supports the mechanisms discussed

here, additional work will be needed for a deeper understanding.

Role of non-neuronal cells

In addition to changes in neuronal number we found a consistent change in endothelial cell

number. Limited to the medial M2 region, the number of endothelial cells was reduced in SNI-

treated mice compared with sham. This effect was already evident at week one and persisted

until week 12. At the moment, we can only speculate about the pathophysiological contribution

of this change. A possible explanation could be that the microvascular network becomes more

sparse after SNI. This could in turn affect the metabolic support of M2 neurons and may be a

factor synergistically acting with the structural changes in dendrites and concomitant

connectivity. Astroglia were significantly reduced after SNI at week one, but not at week 12. This

could be an apparent difference related to the higher variability of the data at week 12.

However, the lack of astrogliosis would rather argue against a neurodegenerative mechanism of

neuronal cell death in M2 after SNI.

Technical aspects of our imaging and analysis approach

The approach used for comprehensive tissue analysis, NuClear, delivers an unbiased and

comprehensive analysis of cell type composition. The number of cell types detected depends on

the training data used for the neuronal network and is not limited to the cell types chosen here.

The detection accuracy differs between cell types and ranges from 95% for glial cells to 99% for

neurons and endothelial cells. By precisely re-identifying imaging volumes acquired at different

time points, artificial alterations of cell numbers by misregistration is unlikely to happen.

Variations in imaging quality between individual session may in some cases affect the outcome.

However, given the large number of volumes analyzed here, such an effect would cancel out

given the study design used here (SNI versus control, volumes ipsi- and contralateral to the

lesion, medial versus more lateral volumes).

Materials and Methods

<u>Ethical approval</u>

The entire study was conducted in line with the European Communities Council Directive

(86/609/ECC) to reduce animal pain and/or discomfort. All experiments followed the German

animal welfare guidelines specified in the TierSchG. The study has been approved by the local

animal care and use council (Regierungspräsidium Karlsruhe, Baden-Wuerttemberg) under the

reference numbers G206/11, G294/15 and G27/20. All experiments followed the 3R principle

and complied with the ARRIVE guidelines (Percie du Sert, Hurst et al. 2020).

<u>Animals</u>

Transgenic mice expressing the human histone 2B protein (HIST1H2B) fused with enhanced

green fluorescence protein (eGFP) under the control of the chicken beta actin promoter (CAG-

H2B-eGFP) (Hadjantonakis and Papaioannou 2004) (B6.Cg-Tg(HIST1H2BB/EGFP)1Pa/J, Jackson

Laboratory; # 006069) were used in animal experiments of cohort 1 (21 mice in total). C57BL6-J

mice were used for cohort 2 (16 mice in total).

<u>mGFP expression in single neurons (cohort 2)</u>

Two adeno-associated virus constructs were used: (1) A DIO cassette containing the synapsin

promotor and membrane-bound GFP to achieve long-term neuron-specific expression, (2) a

cassette expressing Cre-recombinase under the control of the CBA promotor. The in house-

produced AAV1/2 viral particles were stereotaxically injected into the medial M2 region before

the cranial window surgery using a high titer of construct (1) and a 10000-20000x dilution of

construct (2), both virus solutions were mixed at equal amounts. Stereotaxic coordinates were

AP: 1.8 or -0.1 to -0.5, ML: -0.25 to -0.3, DV: − 0.35. This strategy resulted in sufficiently strong

expression of mGFP in a few neurons within the target volume.

<u>Chronic cranial window implantation</u>

A craniectomy procedure was performed on each mouse to enable in vivo two-photon imaging

following a predefined protocol as described before (Holtmaat, Bonhoeffer et al. 2009). Briefly,

the mice were anesthetized with an intraperitoneal injection (i.p.) of a combination of 60 µl

Medetomidine (1mg/ml), 160 µl Midazolam (5 mg/ml) and 40 µl Fentanyl (0.05 mg/ml) at a

dosage of 3 µl/g body weight. The head was shaved, and the mice were fixed with ear bars in

the stereotactic apparatus and eye ointment was applied. Xylocain® 1 % (100 µl, Lidocaine

hydrochloride) was applied under the cranial skin and 250 µl Carprofen (0.5 mg/ml) was

injected subcutaneously (s.c.). The skin was removed to expose the skull and the surface of the

skull was made rough to allow the cement to adhere better. A skull island (approx. 6mm Ø) was

drilled centered at bregma using a dental drill and removed with a forceps (#2 FST by Dumont)

making sure not to damage the cortical surface. For improved imaging condition, the dura was

carefully removed using a fine forceps (#5 FST by Dumont). Normal rat ringer solution (NRR) was

applied on the exposed brain surface to keep it moist, and a curved cover glass was placed on

top to cover it (Asan, Falfan-Melgoza et al. 2021). With dental acrylic cement (powder: Paladur,

Kulzer; activator: Cyano FAST, Hager & Werken GmbH) the cover glass was sealed, and excess

cement was used to cover the exposed skull and edges of the skin. A custom designed holder

was placed on top of the window and any gaps were filled with cement to ensure maximum

adhesion with the skull. After the procedure, the mice were injected (i.p. / s.c.) with a mix of 30

µl Atipamezole (5 mg/ml), 30 µl Flumazenil (0.1 mg/m) and 180 µl Naloxon (0.4 mg/ml) at a

dosage of 6 µl per gram body weight. To ensure proper recovery of the mice, 3 more doses of

Carprofen were given every 8-12 hours and the mice were placed onto a heating plate and

monitored.

<u>In vivo two photon imaging</u>

Two photon imaging was carried out with a two-photon microscope (TriM Scope II, LaVision

BioTec GmbH) equipped with a tunable pulsed Titanium-Sapphire (Ti:Sa) laser (Chameleon Ultra

2; Coherent). Prior to each imaging session, the laser power was adjusted to achieve the best

signal to noise ratio. Adaptations were made to minimize the effect of laser attenuation due to

tissue depth by creating a z-profile and setting the laser power for different imaging depths

while making sure to minimize oversaturation of image pixels.

<u>In vivo two photon imaging of the H2B-GFP mice (cohort 1)</u>

Imaging of H2B-GFP was performed at a laser wavelength of 960 nm. A water immersion

objective (16x; NA 0.8, Nikon) was used to obtain volumetric 3D stacks of the H2B-GFP mice.

Individual frames consisted of 694 µm x 694 µm in XY with a resolution of 0.29 µm/pixel. Stacks

were obtained at varying depths up to 702 µm from the cortical surface with a step size of 2 µm

in Z. Mice were initially anesthetized using a vaporizer with 6% isoflurane and eye ointment was

applied. For imaging, mice were fixed in a custom-built holder on the microscope stage.

Isoflurane (Baxter) was adjusted between 0.5 – 1.5% depending on the breathing rate of each

mouse to achieve a stable breathing rate of 55 – 65 breaths per minute with an oxygen flow rate

of 0.9 – 1.1 l/min. A heating pad was placed underneath the mouse to regulate body

temperature. An infrared camera was used for monitoring of the mice during the imaging.

<u>In vivo two photon imaging of single cell labelled mice (cohort 2)</u>

Imaging of membrane-bound GFP (mGFP) was performed at a laser wavelength of 960 nm. A

water immersion objective (25x, NA 1.1, Nikon) was used to obtain overlapping volumetric 3D

stacks of whole neurons labelled with mGFP. Individual frames covered an XY area of 292 µm x

292 µm, corresponding to a resolution of 143 nm/pixel when sampling at 2048 x 2048 pixels at

16 bit gray scale resolution. Using a Z-step size of 1 µm, image stacks consisted typically of 500

sections. Four to six adjacent image stacks were acquired with a 15-20% overlap for stitching. A

cropped superstack typically consisted of 3600 x 4500 x 500 pixels. Further details are provided

in Supplementary Methods.

Mice were anesthetized using an intraperitoneal injection (i.p.) of a combination of 60 µl

Medetomidine (1mg/ml), 160 µl Midazolam (5 mg/ml) and 40 µl Fentanyl (0.05 mg/ml) at a

dosage of 3 µl/g body weight. For imaging, mice were fixed in a custom-built holder on the

microscope stage. After the imaging session, mice were injected (i.p. / s.c.) with a mix of 30 µl

Atipamezole (5 mg/ml), 30 µl Flumazenil (0.1 mg/m) and 180 µl Naloxon (0.4 mg/ml) at a

dosage of 6 µl per gram body weight. For recovery, mice were kept on a heating plate.

<u>SNI surgery</u>

For the induction of neuropathic pain, a spared nerve injury (SNI) of the sciatic nerve was

performed after baseline imaging as described elsewhere (Gangadharan et al. 2022). Briefly,

mice were anesthetized using isoflurane as described above. After applying a local anesthetic,

an incision on the left hindpaw was made above the sciatic nerve. The peroneal and tibial nerve

were ligated and cut and the sural nerve (innervating the lateral part of the ipsilateral paw) was

left intact.

<u>Behavior analysis</u>

For habituation, mice were placed in the dark behavior room for 3 days, one hour each prior to

the day of the experiment as well as the day of the experiment. All behavior experiments were

performed in the dark cycle of the circadian rhythm of the mice under red light conditions.

<u>von Frey test</u>

To test hypersensitivity after chronic pain induction, the von Frey test was used as described

elsewhere (Gangadharan et al. 2022). Briefly, all mice were habituated on the von Frey grid

prior to the experiment. Each mouse was stimulated on the lateral part of both the ipsilateral

and contralateral paw with von Frey filaments ranging from 0.07 g to 1.4 g. Each filament test

consisted of 5 trials and trials were either marked 0 (negative response) or 1 (positive response)

depending on whether the mouse showed any response like paw flicking, licking or flinching

behavior. For analysis, a mouse was classified as being responsive to a certain filament when

more than 50% of the trials were positive.

<u>Cold plate test</u>

To test allodynia, the cold plate test way used (Tan et al. 2017). The cold plate was set at 4°C

and mice were lowered onto the plate and the latency for a reaction was tested for 30s. The

time for a first aversive response (paw withdrawal, jumping, licking) was noted and any mouse

not responding during the first 30 s was assigned a latency of 30 s.

<u>Open field test</u>

The open field test was used to test for the emotional aspect of pain. The setup consists of a

50cm x 50cm box with 15 cm high walls and a separate distinction for the center of the box.

Mice were placed in the center of the box and were subsequently recorded with a video

camera for 5 minutes. A custom script was designed in Bonsai for mouse tracking and analysis

of time spent in the center or outside the center of the box. Immobility time excluding

grooming was manually scored by an investigator blinded for the mice.

<u>Reidentification of stack positions over time in cohort 1</u>

To be able to compare cell type changes in the same volume over time, the imaging positions of

each imaging volume had to be identical. To achieve this, three or more large vessels on the

cortical surface were used as fiducials. The center of the first vessel was set as zero on both X

and Y coordinates of the microscope stage. The coordinates of the subsequent vessels and

stacks were noted with respect to the first vessel. For imaging of the subsequent timepoint, the

coordinates were used to identify the approximate location of the stack. After roughly

reidentifying the imaging position, the microscopi field of view in different imaging planes along

the Z-axis was aligned to fit to the previous position. A registration system based on the

distance and angle of the first vessel from bregma was created using a self-written script in

Python, which was used to assign the position of all imaging volumes with respect to bregma.

Using a mouse brain atlas (Paxinos & Franklin, 2012), imaging volumes were selected which

were located in the secondary motor area (M2) and the cingulate cortex (CC). Each imaging

volume was divided into diagonal wedges (Fig. 3) corresponding the M2 and CC regions for both

ipsilateral and contralateral hemispheres for further analysis. Due to a "pseudo"-dura consisting

of probably fibroblasts growing on top of the brain, imaging was started from ~76 μm below

occurrence of the first fluorescent signal. Imaging was performed until 550 μm below the first

fluorescent signal due signal-to-noise ratio increasing with depth.

<u>Image analysis of individual neurons used in cohort 2</u>

The image analysis pipeline is illustrated in Supp. Figure 4 and described in detail in the

Methods supplement.

Morphometric sorting of neurons

Interneurons were identified based on their multipoloar shape, lower spine density and small

size. Pyramidal neurons were identified by the typical pyramidal shape of the soma and a strong

apical dendrite, high spine density, and large size.

<u>Antibody staining</u>

Antibody staining was performed on 50 μm thick brain slices with following antibodies: Anti-

NeuN (monoclonal mouse) antibody (Synaptic Systems; #266 011; dilution 1:500), Anti-IBA1

(polyclonal rabbit) antibody (Fujifilm – Cellular dynamics; #019-19741; dilution 1:250), Anti-

S100ß (polyclonal chicken) antibody (Synaptic Systems; #287 006; dilution 1:500) for astroglia.

Secondary antibodies were used with 1:500 dilution.

<u>NuClear analysis for identification of cell types belonging to individual H2B nuclei</u>

Details of this method are published elsewhere (Das-Gupta et al. 2022). Briefly, automated

segmentation of all nuclei was performed with the StarDist neuronal network using a high

performance cluster environment (bwForCluster MLS&WISO). Afterwards, radiomics features

for every segmented nucleus were extracted using the PyRadiomics library and self-written

python scripts. Every segmented nucleus was classified with five pre-trained neuronal network

classifiers and assigned a unique class (neurons, astroglia, microglia, oligodendroglia,

endothelial cells). Nuclei that were assigned to two or more classes were assigned to the

"undecided" class; nuclei that were assigned to no class were labelled as "unclassified".

<u>Statistical analysis</u>

In general, statistical analysis was performed in R (R. Core Team 2021). Depending on the

distribution of the data either t-tests or non-parametric statistical test (Wilcoxon signed-rank

test) were used for statistical testing, p-values were corrected for multiple comparisons using

the "Bonferroni"-method. Plots were created using the R package "ggplot2" (Wickham 2016).

<u>Statistical analysis of 3D cell type distribution</u>

All data were stored in a local MySQL database (MySQL Workbench 8.0). Data were imported

into R. Cell count and first nearest neighbor distance were calculated for each cell type for

different timepoints. The nearest neighbor distance was calculated using the "spatstat"

package for R (Baddeley and Turner 2005).

Contributions

ADG: Cohort 1. Cranial window and SNI surgeries, imaging data acquisition, implementation

and development of analysis pipeline, data analysis, conceptual inputs, generation of figures.

HZ: Cohort 2. Analysis of 2P in vivo imaging data, development of automated tracing algorithm

compatible with large data volumes, generation of figures.

JJ: Cohort 1. Cranial window and SNI surgeries, behavior and imaging data acquisition, tissue

volume reidentification.

LA: Cohort 1. Development of the initial analysis pipeline for nucleus-based cell type

identification, tissue volume reidentification script.

CAB: Cohort 1. Development of segmentation, application of StarDist.

SK: Cohort 2: Cranial window implantations, SNI surgeries and 2P imaging data acquisition.

RK: Initial conceptual input.

JK: Conceptualization, implementation, experimentation, analysis, writing of the MS, project

management and supervision.

TK: Project idea and concept, writing of the MS, supervision.

Acknowledgements

We want to thank Michaela Kaiser, Dunja Baumgartl-Ahlert and Nadine Gehrig for their

excellent technical support during preparation and execution of experiments. We thank

Deepitha Männich for help with posthoc analysis. We gratefully acknowledge the support by

the German Research Foundation (DFG) (SFB1158, project B08 awarded to TK), the data storage

service SDS@hd, supported by the Ministry of Science, Research and the Arts Baden-

Württemberg (MWK) and the DFG through grant INST 35/1314-1 FUGG, as well as the high

performance cluster bwForCluster MLS&WISO and HELIX, supported by the MWK and the DFG

through Grant INST 35/1134-1 FUGG.

Figure legends

Figure 1: Experimental design (a) A spared nerve injury (SNI) surgery was carried to induce

chronic pain in mice. Effects of SNI were studied in the secondary motor area (M2) and anterior

cingulate cortex (ACC) using in vivo 2-photon (2P) microscopy of cell nuclei and single neurons.

Nuclei were classified into cell types using a neural network classifier. **(b)** Longitudinal study

design for Cohort 1 (nuclei imaging) and cohort 2 (single neuron imaging) **(c)** von Frey test

shows significant differences of the threshold filament between sham and SNI animals at 50%

paw withdrawal threshold and **(d)** cold plate latency is significantly lower in SNI mice. **(e)**

Immobility time showing significant difference between SNI and Sham. All behavior data N=10

for SNI and sham.

Figure 2: Imaging analysis of cohort 1 (a) Schematic representation of imaging areas for 2P

imaging in cohort 1 **(b)** 3D reconstruction of a volumetric image. Zoom in shows a sub volume

and a 2D maximum intensity projection (50 μm). **(c)** Imaging positions were reidentified over

time using large vessels as fiducials. Each imaging volume was further classified into different

cell types using a pre-trained neuronal network. **(d)** Absolute cell count per mm^3 for different

imaging timepoints. N=10 for SNI and sham. **(e)** Coronal projections (700 μm and 100 μm)

showing distribution of classified cells in distinct pattens.

Figure 3: Changes in cell type composition in CC and M2 (a) Cell type changes in ipsilateral and

contralateral hemispheres. Inlay depicts a schematic representation of the imaging volumes

with regard to brain areas (dotted line), ACC nuclei are shown in blue, M2 nuclei in black. N left

panel: SNI =7 / Sham = 10; right panel: SNI = 9 / Sham = 7. Wilcox test. M2: week 1 contralateral

neurons: *** (p = 0.000175); Aglia: ** (p = 0.005240); EC: * (p = 0.022900); week 12:

contralateral neurons : * (p = 0.03110); EC = ** (p = 0.00122). CC: Week 1 contralateral neurons

: * (p = 0.0311); Week 12 contralateral Olig: * (p = 0.0418).

Figure 4: Loss of cells in medial M2 after SNI. (a) A group of eight M2 neurons imaged before

SNI (basal) and 1, 9 and 12 weeks after SNI. Images show cells after image processing as

summarized in Supp. Fig. 4 and the Methods supplement. Shown are fully reconstructed

neurons with the raw fluorescence excised based on the traced skeleton (hence no background

visible). Scale bar 100 μm. (b) Neuron imaged in M2 of a sham mouse at the same time points

reported in (a). Scale bar 100 μm. (c) Quantitative comparison of the number of cells counted at

different time points in sham and SNI treated mice. N=20 cells for sham and N=28 for SNI, taken

from a total of 16 mice. Mean ± SEM. ANOVA with Bonferoni multiple comparison test, * depict

P <0.05. (d) Quantitative comparison of the number of lost interneurons versus pyramidal

neurons after SNI. N=14 pyramidal cells and N=18 interneurons, taken from a total of 16 mice.

Mean ± SEM. ANOVA with Bonferoni multiple comparison test, * depict P <0.05.

Figure 5: Spine loss and changes in dendritic morphology. (a) Neuron 4 (from Fig. 4) imaged at

basal state, 3 days, 1 week and 2 weeks after SNI. Neuron disappeared before week 4. Scale bar

20 μm. (b) Neurons 2 and 5 (from Fig. 4) imaged at multiple time points as indicated. The

respectively earlier time point is shown in red, the later time point in green, so that preserved

cellular morphology will display yellow, but lost structures appear in red. The two colour-coded

insets (scale bars 20 μm) show details of a dynamic region (magenta) and a static region (blue).

Scale bar in upper panels = 50 μm.

Supplementary Figure 1: Sensory testing of cohort 2 mice. Von Frey withdrawal thresholds for

SNI (N=7) and sham mice (N=6). Mean ± SEM.

Supplementary Figure 2: Scheme illustrating the imaging situation at the cingulate cortex. The

red square represents the imaging volume of 700 μm width. The triangle demarks the sagittal

sinus. Drawing approximately to scale. AP position of coronal section at approximately -0.2. M1

primary motor cortex, M2 secondary motor cortex, Cgd dorsal cingulate cortex, Cgv ventral

cingulate cortex.

Supplementary Figure 3: Post-hoc analysis of neurons located in the medial M2. Two examples

(A1, A2) of neurons stained with anti-GFP antibody in fixed coronal sections. Parvalbumin

staining (red). Triangles represent the medial M2 region. Scale bar 200 μm.

Supplementary Figure 4: Illustration of image analysis pipeline. (a) Panels showing different

steps in image processing. (a1) raw data. (a2) data excised from (a1) using the traced skeleton

(a3). (b) Workflow of Bayesian learning-based iterative image analysis strategy. A detailed

description can be found in the Supplemental Methods.

Supplementary Figure 5: Changes in spine density in neurons prone to die. Comparison of

spine density determined for entire neurons over time. Neurons imaged in sham mice (black),

neurons imaged in SNI mice surviving 16 weeks, and neurons surviving four weeks or less (blue).

Mean ± SEM, N=4 mice in each group, ANOVA with Bonferroni multiple comparison (* = $P <$

0.05).

References

Apkarian AV, Sosa Y, Sonty S, et al (2004) Chronic back pain is associated with decreased
574   prefrontal and thalamic gray matter density. The Journal of neuroscience : the official
575   journal of the Society for Neuroscience 24:10410–10415.
576   https://doi.org/10.1523/jneurosci.2541-04.2004

577   Asan L, Falfán-Melgoza C, Beretta CA, et al (2021) Cellular correlates of gray matter volume
578   changes in magnetic resonance morphometry identified by two-photon microscopy.
579   Scientific Reports 11:4234. https://doi.org/10.1038/s41598-021-83491-8

580   Barthas F, Kwan AC (2017) Secondary Motor Cortex: Where 'Sensory' Meets 'Motor' in the
581   Rodent Frontal Cortex. Trends Neurosci 40:181–193.
582   https://doi.org/10.1016/j.tins.2016.11.006

583   Bilbao A, Falfán-Melgoza C, Leixner S, et al (2018) Longitudinal Structural and Functional Brain
584   Network Alterations in a Mouse Model of Neuropathic Pain. Neuroscience 387:104–115.
585   https://doi.org/10.1016/j.neuroscience.2018.04.020

586   Bliss TVP, Collingridge GL, Kaang B-K, Zhuo M (2016) Synaptic plasticity in the anterior cingulate
587   cortex in acute and chronic pain. Nat Rev Neurosci 17:485–96.
588   https://doi.org/10.1038/nrn.2016.68

589   Blom SM, Pfister JP, Santello M, et al (2014) Nerve Injury-Induced Neuropathic Pain Causes
590   Disinhibition of the Anterior Cingulate Cortex. Journal of Neuroscience 34:5754–5764.
591   https://doi.org/10.1523/jneurosci.3667-13.2014

592   Chi H, Chang H-Y, Sang T-K (2018) Neuronal Cell Death Mechanisms in Major
593   Neurodegenerative Diseases. Int J Mol Sci 19:3082. https://doi.org/10.3390/ijms19103082

594   Das-Gupta A, Asan L, John J, et al (2022) Comprehensive monitoring of tissue composition using
595   in vivo imaging of cell nuclei and deep learning. Biorxiv 2022.10.03.510670.
596   https://doi.org/10.1101/2022.10.03.510670

597   Decosterd I, Woolf CJ (2000) Spared nerve injury: an animal model of persistent peripheral
598   neuropathic pain. Pain 87:149–158. https://doi.org/10.1016/s0304-3959(00)00276-1

599   Faria O de, Pivonkova H, Varga B, et al (2021) Periods of synchronized myelin changes shape
600   brain function and plasticity. Nat Neurosci 24:1508–1521. https://doi.org/10.1038/s41593-
601   021-00917-2

Forrest MP, Parnell E, Penzes P (2018) Dendritic structural plasticity and neuropsychiatric
disease. Nat Rev Neurosci 19:215–234. https://doi.org/10.1038/nrn.2018.16

Fricker M, Tolkovsky AM, Borutaite V, et al (2018) Neuronal Cell Death. Physiol Rev 98:813–880.
https://doi.org/10.1152/physrev.00011.2017

Gangadharan V, Zheng H, Taberner FJ, et al (2022) Neuropathic pain caused by miswiring and
abnormal end organ targeting. Nature 1–9. https://doi.org/10.1038/s41586-022-04777-z

Holtmaat A, Svoboda K (2009) Experience-dependent structural synaptic plasticity in the
mammalian brain. Nature Reviews Neuroscience 10:647–658.
https://doi.org/10.1038/nrn2699

Inquimbert P, Moll M, Latremoliere A, et al (2018) NMDA Receptor Activation Underlies the
Loss of Spinal Dorsal Horn Neurons and the Transition to Persistent Pain after Peripheral
Nerve Injury. Cell Reports 23:2678–2689. https://doi.org/10.1016/j.celrep.2018.04.107

Ji R-R, Donnelly CR, Nedergaard M (2019) Astrocytes in chronic pain and itch. Nature Reviews
Neuroscience 20:667–685. https://doi.org/10.1038/s41583-019-0218-1

Jung H-Y, Kim J-H, Lee WJ, et al (2019) Microstructural Abnormalities of the Secondary Motor
Area Coordinating Sensory and Motor Functions in Patients with Complex Regional Pain
Syndrome. Neuropsychiatry 9:2419–2427

Kassem MS, Lagopoulos J, Stait-Gardner T, et al (2013) Stress-induced grey matter loss
determined by MRI is primarily due to loss of dendrites and their synapses. Molecular
neurobiology 47:645–661. https://doi.org/10.1007/s12035-012-8365-7

Kim S, Kato G, Ishikawa T, Nabekura J (2011) Phase-specific plasticity of synaptic structures in
the somatosensory cortex of living mice during neuropathic pain. Molecular Pain 7:87.
https://doi.org/10.1186/1744-8069-7-87

Kim SK, Nabekura J (2011) Rapid synaptic remodeling in the adult somatosensory cortex
following peripheral nerve injury and its association with neuropathic pain. 31:5477–5482.
https://doi.org/10.1523/jneurosci.0328-11.2011

Kuner R, Kuner T (2021) Cellular Circuits in the Brain and Their Modulation in Acute and Chronic
Pain. Physiol Rev 101:213–258. https://doi.org/10.1152/physrev.00040.2019

Maihöfner C, Baron R, DeCol R, et al (2007) The motor system shows adaptive changes in
complex regional pain syndrome. Brain 130:2671–2687.
https://doi.org/10.1093/brain/awm131

Mascio G, Notartomaso S, Martinello K, et al (2022) A Progressive Build-up of Perineuronal Nets
in the Somatosensory Cortex Is Associated with the Development of Chronic Pain in Mice. J
Neurosci 42:3037–3048. https://doi.org/10.1523/jneurosci.1714-21.2022

Metz AE, Yau H-J, Centeno MV, et al (2009) Morphological and functional reorganization of rat
medial prefrontal cortex in neuropathic pain. Proceedings of the Royal Society of London
Series B, Containing papers of a Biological character Royal Society (Great Britain) 106:2423–
2428. https://doi.org/10.1073/pnas.0809897106

Mostafapour SP, Cochran SL, Puerto NMD, Rubel EW (2000) Patterns of cell death in mouse
anteroventral cochlear nucleus neurons after unilateral cochlea removal. J Comp Neurol
426:561–571. https://doi.org/10.1002/1096-9861(20001030)426:4<561::aid-cne5>3.0.co;2-
g

Moujalled D, Strasser A, Liddell JR (2021) Molecular mechanisms of cell death in neurological
diseases. Cell Death Differ 28:2029–2044. https://doi.org/10.1038/s41418-021-00814-y

Reichling DB, Levine JD (2011) Pain and death: Neurodegenerative disease mechanisms in the
nociceptor. Ann Neurol 69:13–21. https://doi.org/10.1002/ana.22351

Salter MW, Stevens B (2017) Microglia emerge as central players in brain disease. Nat Med
23:1018–1027. https://doi.org/10.1038/nm.4397

Seminowicz DA, Laferriere AL, Millecamps M, et al (2009) MRI structural brain changes
associated with sensory and emotional function in a rat model of long-term neuropathic
pain. NeuroImage 47:1007–1014. https://doi.org/10.1016/j.neuroimage.2009.05.068

Silva JTD, Seminowicz DA (2019) Neuroimaging of pain in animal models: a review of recent
literature. Pain Reports 4:e732. https://doi.org/10.1097/pr9.0000000000000732

Tan LL, Pelzer P, Heinl C, et al (2017) A pathway from midcingulate cortex to posterior insula
gates nociceptive hypersensitivity. Nat Neurosci 20:1591–1601.
https://doi.org/10.1038/nn.4645

Tracey I, Mantyh PW (2007) The Cerebral Signature for Pain Perception and Its Modulation.
Neuron 55:377–391. https://doi.org/10.1016/j.neuron.2007.07.012

Zhao R, Zhou H, Huang L, et al (2018) Neuropathic Pain Causes Pyramidal Neuronal
Hyperactivity in the Anterior Cingulate Cortex. Front Cell Neurosci 12:107.
https://doi.org/10.3389/fncel.2018.00107

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

a1   a2   10 µm   a3

b

3D image stack

Accuracy evaluation of posterior

Incremental Bayesian graph structure learning

Structure voting

Direct detected broken fibers, no connectivity

Parametric optimization, structure prediction

Refine

Estimate

self-learned prior

Structure detection, connect broken fibes

Iterative adaptive local and global optimization

Vectorization of detected dendrites and spines

Supplementary Figure 5