Discovery of lipid binding sites in a ligand-gated ion channel by integrating simulations and cryo-EM

Cathrine Bergh[1], **Urška Rovšnik**[2], **Rebecca J. Howard**[1,2], **Erik Lindahl**[1,2]*

*For correspondence:
erik.lindahl@scilifelab.se

[1]Science for Life Laboratory & Swedish e-Science Research Center, Department of Applied Physics, KTH Royal Institute of Technology, PO Box 1031, SE-171 21 Solna, Sweden; [2]Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, PO Box 1031, SE-171 21 Solna, Sweden

---

Abstract   Ligand-gated ion channels transduce electrochemical signals in neurons and other excitable cells. Aside from canonical ligands, phospholipids are thought to bind specifically to the transmembrane domain of several ion channels. However, structural details of such lipid contacts remain elusive, partly due to limited resolution of these regions in experimental structures. Here, we discovered multiple lipid interactions in the channel GLIC by integrating cryo-electron microscopy and large-scale molecular simulations. We identified 25 bound lipids in the GLIC closed state, a conformation where none, to our knowledge, were previously known. Three lipids were associated with each subunit in the inner leaflet, including a buried interaction disrupted in mutant simulations. In the outer leaflet, two intrasubunit sites were evident in both closed and open states, while a putative intersubunit site was preferred in open-state simulations. This work offers molecular details of GLIC-lipid contacts particularly in the ill-characterized closed state, testable hypotheses for state-dependent binding, and a multidisciplinary strategy for modeling protein-lipid interactions.

Introduction

Pentameric ligand-gated ion channels (pLGICs) can be found in numerous cell types, particularly in the postsynaptic membrane of the vertebrate nervous system, where they transduce electrochem­ical signals upon neurotransmitter binding (*Tasneem et al., 2005*). Gating in pLGICs is initiated by ligand binding in the extracellular domain (ECD), which is believed to lead to large twisting and blooming motions in the ECD, followed by rearrangements in the transmembrane domain (TMD), which eventually lead to opening of the central transmembrane pore. In addition to soluble ligands and numerous allosteric modulators, lipid components of the plasma membrane have been shown to bind to a variety of membrane proteins, potentially modifying their stability or gating (*Duncan et al., 2020*). For pLGICs, lipid composition was shown to influence reconstitution of functional activity in the *Torpedo* nicotinic acetylcholine receptor (nAChR) as early as the 1980s (*Popot et al., 1981*; *Criado et al., 1982*; *Fong and McNamee, 1986*; *daCosta et al., 2002*). Multiple members of the pLGIC family, particularly nAChRs and prokaryotic channels cloned from *Gloeobacter violaceus* (GLIC) and *Dickeya dadantii* (formerly *Erwinia chrysanthemi*, reflected in the channel name ELIC), have since demonstrated functional sensitivity or co-purification with membrane lipids (*Thomp­son and Baenziger, 2020*). It has been suggested that lipids interact with membrane-facing loops from the ECD, or with the four membrane-spanning helices (M1–M4) of the TMD. However, many

structural details of these lipid interactions remain poorly understood (*Poveda et al., 2017*).

A critical challenge to elucidating lipid-protein interactions is their typically poor definition in experimental densities. Among more than 100 deposited structures in the protein data bank for GLIC crystallized under activating conditions, several lipid poses have been reported (*Zhu et al., 2018*; *Masiulis et al., 2019*; *Laverty et al., 2019*; *Walsh et al., 2018*; *Gharpure et al., 2019*; *Rahman et al., 2020*; *Zhao et al., 2021*). These include an outer-leaflet site on the complementary face of a single GLIC subunit, an inner-leaflet site on the principal face of a subunit, and an interfacial inner-leaflet site making contacts with two neighboring subunits (*Thompson and Baenziger, 2020*). In one case, an additional outer-leaflet density was built as a detergent molecule (*Hu et al., 2018*). However, the positions and orientations of these membrane components or mimetics vary between structures and typically include only a subset of their constituent atoms.

Another issue is the potential dependence of lipid interactions on the functional state of the membrane protein. In most members of the pLGIC family, binding of a chemical stimulus to the ECD favors a transition from an unliganded, nonconducting state to a liganded, conducting state, and in some cases to a subsequent liganded, desensitized state. Although terminology in the field may vary, conditions predicted to favor the first versus second of these states are referred to here as *resting* versus *activating*; the corresponding conformations of the TMD pore are designated *closed* versus *open*. No lipids were reported with the lone X-ray structure of GLIC under resting conditions, raising the possibility that interactions could be state-dependent, or obscured by the relatively low overall resolution of this dataset (4.0 Å) (*Sauguet et al., 2014*). Independent validation of phospholipid interactions in the closed as well as open states of GLIC would provide a valuable baseline for investigating such contacts in the larger family of pLGICs, including pharmacologically important effects of components such as cholesterol and neurosteroids that modulate LGICs (*Thompson and Baenziger, 2020*; *Kim and Hibbs, 2021*).

Molecular dynamics simulations offer an alternative approach to visualizing lipid-protein interactions. All-atom simulations of membrane proteins typically involve inserting an experimental structure in a simplified lipid bilayer surrounded by water and ions, all of which move freely according to physical principles encoded in the force field. Simulations on both atomistic and coarse-grained levels have recently enabled the identification of specific lipid interactions in pLGICs such as ELIC (*Sridhar et al., 2021*; *Dietzen et al., 2022*), nAChRs (*Sharp and Brannigan, 2021*), glycine (*Dämgen and Biggin, 2021*) and serotonin-3 receptors (*Crnjar and Molteni, 2020*), but were typically limited to shorter timescales or specific protein conformations. We recently reported free-energy landscapes for GLIC gating using Markov state modeling of 120 μs unrestrained molecular dynamics simulations (*Bergh et al., 2021*). Because GLIC is activated by external acidification (pH < 6), resting versus activating conditions could be approximated throughout these simulations by setting titratable amino-acid residues as deprotonated versus protonated, respectively. Under these two conditions, different distributions of functional states were captured, consistent with the relative stabilization of open versus closed channels upon activation. Although our previous analyses focused on conformational changes of the protein alone, the same simulation data encodes a wealth of additional features, including membrane lipids. We also recently reported the first structural data for GLIC using cryogenic electron microscopy (cryo-EM), which enabled the reconstruction of multiple new closed structures (*Rovšnik et al., 2021*).

Here, we present the first cryo-EM structure of GLIC in a closed conformation with multiple bound lipids resolved in each subunit. These were built into non-protein densities evident in the cryo-EM data, further informed by closed-state lipid occupancies independently identified and characterized from extensive molecular dynamics simulations. Five distinct closed-state lipid-interaction sites per subunit covered intra- and intersubunit cavities in the inner and outer leaflets. We also compared open-state lipid occupancies from our simulations with previous open X-ray structures, identifying six possible poses. These analyses allowed prediction of novel lipid interaction features in GLIC, including a potentially state-dependent binding site at the outer subunit interface, and a potential role for lipid-tail saturation in binding at the inner leaflet, further validated through mutant

92 simulations. In addition to casting light on selective lipid interactions with LGICs, this work demon-
93 strates the combined power of experimental densities and molecular simulations to support the
94 characterization of lipid binding sites in systems challenged by heterogeneous or otherwise low-
95 resolution data.

Results

Markov state modeling and cryo-EM data resolve lipid-protein interactions

We previously reported cryo-EM structures of GLIC in multiple closed states (*Rovšnik et al., 2021*)
and demonstrated how Markov state modeling (MSM) can predict shifts in the GLIC gating land-
scape under resting versus activating conditions (*Bergh et al., 2021*). In this work, we have com-
bined new cryo-EM reconstructions with simulations newly extended to achieve better sampling
of lipids, which made it possible to identify five previously uncharacterized lipid poses interacting
with each subunit in the closed state of GLIC (*Figure 1*A).

For computational quantification of lipid interactions and binding sites, we used molecular sim-
ulations of GLIC conducted under either resting or activating conditions (*Bergh et al., 2021*). As
described in Methods, resting conditions corresponded to neutral pH with most acidic residues
deprotonated; activating conditions corresponded to acidic pH with several acidic residues proto-
nated. Both open and closed conformations were present in both conditions, albeit with different
probabilities. Simulation frames previously identified as either closed or open (*Figure 1*B) were
clustered to their corresponding state, and used to generate densities occupied by lipids that con-
tacted the protein in at least 40% of simulation frames (*Figure 1*D,F). Although the stochastic nature
of simulations resulted in nonidentical lipid densities associated with the five GLIC subunits, pat-
terns of lipid association were notably symmetric (*Figure 1*—**??**). In order to distinguish presumed
functional endpoints, snapshots classified into macrostates other than closed or open (*Figure 1*B,
colored boundaries) were excluded from occupancy calculations, except as indicated below. In
parallel, mean duration times of lipid contacts were quantified for each amino acid residue in each
1.7 μs trajectory (50 seeds ×2 conditions ×5 subunits = 500 trajectories; *Figure 1*C). In some cases,
we validated notable contacts by additional simulations in the presence of targeted mutations (*Figure 1*E).

In our cryo-EM work, a new GLIC reconstruction was generated by merging previously reported
datasets collected at pH 7, 5, and 3 (*Rovšnik et al., 2021*). The predominant class from the merged
data corresponded to an apparently closed channel at an overall resolution of 2.9 Å, the highest
resolution yet reported for GLIC in this state (*Figure 1*—**??**, *Table 1*). After building all residues,
additional non-protein densities were evident in both the outer and inner leaflets of the predicted
membrane region. These were further sharpened by post-processing with ResolveCryoEM (*Terwilliger et al., 2020*) and LocScale (*Jakobi et al., 2017*). From the combined observations of these
cryo-EM densities (*Figure 1*G) and occupied densities in closed-state simulations (*Figure 1*F), five
lipids were built in association with each of the five GLIC subunits, for a total of 25 lipid molecules
(*Figure 1*H, *Figure 1*—**??**C,D).

Lipids built in this structure formed close contacts with GLIC residues in several membrane-
proximal regions. These included an ECD motif with a characteristic proline/cysteine-rich turn,
known as the Pro-loop (or Cys-loop in eukaryotes); N-terminal residues of the M1 helix; regions in
and surrounding the M2–M3 loop; interfacial residues at the C-terminal end of M3; and inward- and
outward-facing regions of M4 (*Figure 2*A). In several cases, these regions corresponded to longer-
duration lipid interactions in molecular simulations (*Figure 2*B,C). In the inner leaflet, relatively
long-lasting interactions were found at the subunit interface, involving buried residues in both the
M1 and M3 helices. In the outer leaflet, longer interactions were focused around the pre-M1 and
M2-M3 loops. Lateral lipid diffusion coefficients were estimated to 1.47 nm^2/μs for bulk lipids and
0.68 nm^2/μs for lipids of the first lipid shell (*Figure 2*—**??**A), which is relatively slow compared to the
timescales of each trajectory (1.7 μs). However, multiple residues throughout the M1, M3, and M4

Figure 1. Lipid-protein interactions mapped by Markov state modeling and cryo-electron microscopy.
(A) Overview of the MD simulation system with GLIC (gray surface) embedded in a lipid bilayer (light blue carbon atoms) with interacting lipids highlighted (yellow carbon atoms). Lipid oxygens are shown in red; phosphorus, orange; nitrogen, dark blue. Water molecules and ions have been omitted for clarity. (B) Markov state models were used to cluster simulations conducted under resting (R) or activating (A) conditions into five states, including closed (left of the light or dark orange lines) and open (right of the light or dark blue lines). Black lines mark edges of other state clusters derived from MSM eigenvectors. Experimental structures are highlighted as white circles. (C) To characterize the stability of lipid interactions, duration times were measured in individual, unclustered, trajectories. (D) To identify lipid binding sites, lipid-occupied densities were obtained from state-clustered frames for closed (orange) and open (blue) states. (E) Further simulations were performed to test the role of residues involved in extensive lipid interactions through single residue mutations around lipid binding sites. (F) View of a manually built cryo-EM model of closed GLIC (gray) with computationally derived lipid occupancies in orange (semi-transparent). (G) Cryo-EM reconstruction of closed GLIC (gray), with partly defined non-protein densities (orange). (H) Equivalent view of a manually built cryo-EM model as in F, with newly built lipids shown as sticks (yellow, heteroatom; phosphorus, orange; oxygen, red; nitrogen, blue).

Figure 2. Cryo-EM and simulations reveal lipid interaction sites. (A) Cryo-EM reconstruction of closed GLIC with distance to the interacting lipids colored according to the bar below. (B) Mean duration time of lipid-residue interactions, scaled (white–red) according to the color bar below, projected onto the closed-state GLIC structure (PDB ID: 4NPQ). For both panels, a single subunit is displayed in cartoon. (C) Secondary structure schematic for GLIC (top), mean duration time of each lipid-residue interaction (middle), and number of lipids interacting with each residue during simulations (bottom). Light gray points represent numbers from individual trajectories and black lines ensemble averages (500 trajectories).

helices exchanged contacts with 2-4 different lipid molecules in individual simulations (*Figure 2*C). Furthermore, 1.7-μs root mean square displacement of lipids originally in the first lipid shell was 2.15 nm, and 3.16 nm in the bulk bilayer, indicating such exchanges are not limited to nearby lipids (*Figure 2*—**??**B). Thus, exchange events and diffusion estimates indicate that the duration of lipid contacts observed in this work can be at least partly attributed to interaction stabilities and not solely to sampling limitations.

Protein and lipid determinants of state-independent inner-leaflet binding

In the inner leaflet, occupied densities from closed-state simulation snapshots were largely super-imposable with those from open snapshots (*Figure 3*A, *Figure 1*—**??**), indicating lipid interactions in this region were not notably state-dependent. In both states, three types of inner-leaflet compu-tational densities could be observed; accordingly, our cryo-EM reconstruction included three types of forked densities, which could be built as three independent lipids per subunit (*Figure 3*B). One density spanned the principal M3 and complementary M1 helices of neighboring subunits; one was located between M3 and M4 on the principal face of each subunit; and another spanned M1 and M4 on the complementary face of each subunit. Interfacial and principal-face inner-leaflet lipids have been resolved in previous open-state X-ray structures of GLIC (*Bocquet et al., 2009*) as well as other pLGIC structures (*Thompson and Baenziger, 2020*), while the complementary inner-leaflet GLIC site, to our knowledge, has not been reported before.

The lipid occupying the subunit interface accounted for some of the longest-lived interactions in the simulations, both with M1 and M3 (*Figure 3*C,F). A particularly frequent contact at this site in our simulations was the unsaturated C = C bond of 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC) (*Figure 3*D,E). Interestingly, lipids adopted two types of poses in this site, depending on the placement of the unsaturated tail. The double bond was frequently wedged between M3-T274 and M1-W217 at the intersubunit cleft, with the remainder of the lipid projecting out towards the membrane (*Figure 3*D, left). When the double bond exited the pocket, the more flexible saturated tail was able to enter the intersubunit cleft and sample a range of poses characterized by contacts distributed more broadly across the lipid tail (*Figure 3*E), resulting in some deeply buried poses (*Figure 3*D, right).

To further explore the influence of specific amino acid contacts on this interfacial lipid pose, we

Figure 3. Protein and lipid determinants of state-independent inner-leaflet binding. (A) Computationally derived densities from the open (blue) and closed (orange) states generally agree. (B) Modeled lower leaflet lipids (yellow, stick, heteroatom coloring) from cryo-EM densities (brown mesh). (C) Zoom-in view of the lower leaflet buried lipid interaction site at the M1-M3 subunit interface. Lipids particularly interact with residues T274 (purple) and W217 (magenta). (D) The 100 simulation frames that best fit the computational densities clustered into two distinct binding poses at the T274 binding site. When the tail with a double bond (yellow sphere) is situated in the pocket lipid heads are directed out from the channel (left panel), while a larger variety of poses are sampled when the lipid tail without a double bond occupies the site, allowing the head to enter into a crevice on the bottom of the channel, approaching the pore (right panel). The M4 helix is not shown for clarity in the top panels. (E) Number of contacts made by specific POPC lipid atoms with residue T274 in simulations at both resting and activating conditions (colors span 0 contacts, white, to 137015 contacts, dark red). For the tail with an unsaturation, interactions are concentrated around the double-bond region while the saturated tail displays interactions interspersed along the tail. (F) Mutation of residue W217, lining this pocket, reveals shortened interactions at the T274 binding site (magenta).

performed additional MD simulations with mutations in targeted sites. Interestingly, the buried M3 position 274 remained the contact of longest duration, even when mutated from threonine to a smaller (alanine) or larger (tryptophan) residue (*Figure 3*—**??**). Conversely, removing the bulky tryptophan sidechain at position 217 (W217A) dramatically shortened lipid contacts at the more deeply buried T274 (*Figure 3*F). These results confirm the importance of specific sidechain identities on the protein surface (W217) in influencing lipid binding pose, whereas even long-lived contacts (T274) may be sustained more by their buried position than side chain size.

Intrasubunit lipids and state dependence in the outer leaflet

In the outer leaflet, closed-state cryo-EM densities enabled confident building of two lipid poses, associated with either the principal or complementary face of each subunit (*Figure 4*A, left). Lipids have previously been resolved in open and inhibited X-ray structures of GLIC at a complementary-face outer-leaflet site between the M1 and M4 helices (*Bocquet et al., 2009*; *Nury et al., 2011*; *Basak et al., 2017*; *Hu et al., 2018*) (*Figure 4*A, right), comparable to that observed here in the cryo-EM closed state. To our knowledge, no lipid has been reported at any other outer-leaflet site in GLIC, though a principal-face density was built as a detergent molecule in at least one open-state X-ray structure (*Hu et al., 2018*) (*Figure 4*A, right).

In closed-state simulation snapshots, outer-leaflet densities largely corresponded to our closed-state cryo-EM data, including distinct interaction sites on the complementary and principal faces (*Figure 4*B, left, *Figure 1*—**??**C). In these and other sites, occupancies and distributions were largely superimposable when calculated from Markov state models constructed independently under resting versus activating conditions (*Figure 4*B,C, *Figure 1*—**??**), indicating reproducibility of the approach and relative insensitivity of the lipid-interaction landscape to pH. However, comparing open- versus closed-state occupancies suggested state-dependent differences in lipid interactions, as described below (*Figure 4*B).

At the complementary face, lipid tails were more likely to enter a buried region of the intrasubunit pocket in the open state (*Figure 4*—**??**). Although the difference was subtle, the distribution of outer-leaflet lipids shifted in snapshots of open compared to closed conformations, increasing (5% to 10%) the probability of lipids penetrating 18 Å from the pore axis (*Figure 4*C). Interestingly, lipid tails entered even deeper (15 Å from the pore) in an alternative state within the GLIC gating landscape (*Figure 4*—**??**A), previously associated with channel desensitization due to relative contraction at the intracellular gate (*Figure 4*—**??**B) (*Bergh et al., 2021*). Although the lipid tails were too poorly resolved in our structural data to definitively capture buried interactions at the complementary site, it is interesting that the simulations implicated them in channel opening (and possibly in transitions further along the activation pathway) given that the deeper parts of this site have been shown to bind allosteric modulators such as propofol (*Heusser et al., 2018*).

At the principal face, interactions with lipid heads were more diverse than those with lipid tails. In our closed cryo-EM and a previously reported open X-ray structure (*Hu et al., 2018*), the lipid or detergent head in this site tilted away from the pore axis, towards M4 (*Figure 4*A). On initial inspection, lipid-head densities appeared less tilted in closed- versus open-state simulation frames (*Figure 4*—**??**C). However, lipid heads in this site were relatively flexible and did align with cryo-EM densities in several closed-state snapshots (*Figure 4*D). In fact, lowering the occupancy threshold (from 40% to 30%) revealed an alternative head-group density, tilted towards M4 and covering most of the lipid built in our cryo-EM structure (*Figure 4*—**??**D). Thus, despite an apparent discrepancy between highly populated densities in simulations and cryo-EM data, both methods supported at least partial occupancy in the built pose. Although this lipid head was primarily associated with M3–M4 on the principal face of one subunit, it also made contacts with residues on the complementary M1 helix (Q193, S196) in closed- but not open-state simulation frames (*Figure 4*—**??**A,B,C), likely associated with conformational changes during gating.

In addition to the intrasubunit sites described above, a third outer-leaflet lipid density was observed in open-state simulation frames, wedged between two subunits (*Figure 4*B, right). This

Figure 4. State-dependent protein-lipid interactions in the upper leaflet. (A) Zoom view of the outer transmembrane domain of the closed GLIC cryo-EM structure (left, gray), showing non-protein cryo-EM densities (brown mesh) overlaid with built lipids (yellow). For comparison, the same region is shown of an open state x-ray structure (PDB ID: 6HZW) (*Hu et al., 2018*) with built lipid and detergent in equivalent positions (right). Conformational changes of the M2-M3 loop, including residue P250, in channel opening are highlighted. Lipids are colored by heteroatom (phosphorus, orange; oxygen, red; nitrogen, blue). (B) Same region as in (A) showing densities occupied by lipids (>40%) in simulation frames clustered as closed (left, orange) or open (right, blue) states (top). Open-state differences include an enhanced density on the right-hand side of the intersubunit cleft, and a more deeply penetrating density in the intrasubunit cleft (*Figure 4—??*). Overlaid densities for each state represent simulations conducted under resting (dark shades) or activating (light shades) conditions, which were largely superimposable within each state. (C) Radial distribution of the lipid atoms closest to the upper pore, showing closer association of lipids in the open (light blue, dark blue) versus closed (orange, red) states. (D) Same region and cryo-EM densities (mesh) as in (A) overlaid with lipids from a few simulation frames where the lipids had the highest correlation with the closed-state computational occupancies in (B). (E) Snapshots from simulation frames that had the highest correlation with the occupancies (activating conditions) in (B). The open states are characterized by an additional lipid at the intersubunit site, interacting with the oxygen of P250. (F) The minimum distance between P250 and the closest lipid projected onto the free energy landscapes obtained from Markov state modeling. P250-lipid interactions are possible in the open state, but not in the closed state.

intersubunit pose was not apparent in closed-state simulations (*Figure 4*B, left), nor in closed or open experimental GLIC structures (*Figure 4*A). The lipid head made particularly close contacts with residue P250 on the M2-M3 loop, which undergoes substantial conformational change away from the pore upon channel opening, along with outer-leaflet regions of M1–M3 (*Figure 4*E, *Figure 4*— **??**A,B,C, *Figure 4*—**??**). These conformational changes were accompanied by a flip of M1 residue F195, which blocked the site in the closed state but rotated inward to allow closer lipid interactions in the open state (*Figure 4*—**??**C, *Figure 4*—**??**). Indeed, P250 was predominantly located within 3 Å of the nearest lipid atom in open- but not closed-state frames (*Figure 4*F). Despite being restricted to the open state, interactions with P250 were among the longest duration in all simulations (*Figure 2*C) and as these binding events preceded pore opening, it is plausible to infer a role for this state-dependent lipid interaction in the gating process (*Figure 4*—**??**D).

Discussion

Lipid interactions have increasingly been shown to play important roles in stabilizing and regulating membrane proteins (*Phillips et al., 2009*; *Cordero-Morales and Vásquez, 2018*). Particularly since the emergence of cryo-EM, a growing number of ion channel structures have been reported with detergents or lipids in membrane-facing regions (*Thompson and Baenziger, 2020*; *Levental and Lyman, 2022*). However, the resolution of bound lipids is typically low compared to the protein, making it challenging to define precise poses or interactions. As a result, lipids and detergents are often built with truncated heads, modeled as isolated hydrocarbon chains, or disregarded altogether to avoid overfitting. Here, we present the building of lipids in the closed state of GLIC by comparing non-protein densities in a relatively high-resolution cryo-EM map with lipid interactions sampled in molecular dynamics simulations.

Our 2.9-Å cryo-EM structure represents, to our knowledge, the highest-resolution closed state of wild-type GLIC yet reported, and the first in which lipids could be confidently built. Notably, the GLIC sample was solubilized in detergent, which is typically thought to replace biological lipids. Nonetheless, the forked shape of non-protein densities associated with each channel subunit indicated the presence of phospholipids, evidently bound tightly enough to resist substitution by detergent. Similarly persistent lipids have been reported in at least three sites per subunit of open-state GLIC; here, we find evidence for five lipids per subunit in the closed state (*Figure 5*A). Continued improvements in membrane-protein cryo-EM may reveal the full extent of persistent lipid binding at key transmembrane sites in the larger pLGIC family.

Molecular dynamics simulations have recently been used to identify lipid binding sites corresponding to experimental data for both prokaryotic and eukaryotic ligand-gated ion channels (*Sridhar et al., 2021*; *Dietzen et al., 2022*), including nAChRs (*Sharp and Brannigan, 2021*), glycine (*Dämgen and Biggin, 2021*) and serotonin-3A receptors (*Crnjar and Molteni, 2020*). In most cases, the slow rate of lateral lipid diffusion has resulted in the preferential use of coarse-grained force fields able to sample longer timescales, but that may overestimate the strength of interactions (*Lamprakis et al., 2021*). In addition, such studies typically apply harmonic restraints to the protein, potentially obscuring contributions of protein dynamics to lipid interactions. Indeed, the unrestrained atomistic MD simulations studied here were not expected to capture the maximal duration of stable contacts, as indicated by some interaction times approaching the full 1.7-μs trajectory (*Figure 2*). Nevertheless, simulations were of sufficient length to sample exchange of up to four lipids, particularly around the M4 helix. Calculation of lipid lateral diffusion coefficients resulted in average displacements at the end of simulations of 2.15 nm for lipids initially interacting with the protein surface, roughly corresponding to lipids diffusing out to the 4th lipid shell. Diffusion of bulk lipids was faster, allowing lipids originally 3.16 nm away from the protein surface to ingress the first lipid shell. This observation underscores the potential for lipid exchange events even among lipids initially distant from the protein surface. Of course, duration of exceptionally stable interactions, such as those involving T274 (*Figure 2*C), inevitably remain bounded by the length of our

Figure 5. Lipid sites identified by cryo-EM and simulations. (A) In the closed state, five lipid interaction sites were independently identified through both cryo-EM and simulations. In the outer leaflet, two intrasubunit sites are situated at the principal (P) and complementary (C) subunits, with the lipid at the principal subunit site interacting with S196 of the pre-M1 loop. Particularly, no lipid site could be found at the subunit interface. In the inner leaflet, a buried lipid pose was found at the subunit interface, determined by interactions with T274 and W217 of the M3 and M1 helices, respectively. (B) In the open state, six lipid interaction sites were identified from simulations, out of which three sites were known from previous structures and one presumably occupied by a detergent molecule (*Hu et al., 2018*) (thin blue stripes). In the outer leaflet, an additional lipid site could be found at the subunit interface in the open state, interacting with P250 on the M2-M3 loop. Additionally, the site at the principal intrasubunit site is slightly closer to the M4 helix compared to the closed state, and the lipid site at the principal intrasubunit site has an increased likelihood of tail penetration into an allosteric pocket compared to the closed state. Lipid sites in the inner leaflet are at positions equivalent to those in the closed state.

simulations. Still, diffusion metrics, supported by robust statistical analysis encompassing diverse
starting conditions (500 trajectories), enable confident estimation of relative interaction times.

Sampling was further simplified by performing simulations in a uniform POPC membrane. Prior
experiments have been conducted to assess the sensitivity of GLIC in varying lipid environments
(*Labriola et al., 2013*; *Carswell et al., 2015*; *Menny et al., 2017*)), indicating that GLIC remains
fully functional in pure POPC bilayers. In our cryo-EM experiments, the protein was recombi-
nantly expressed from *E. coli*, which means that the experimental density would likely represent
phosphatidylglycerol or phosphatidylethanolamine lipids. However, as the molecular identities of
bound lipids could not be precisely determined, POPC lipids were built for straightforward compar-
ison with simulation poses. While it appears that GLIC is capable of gating in a pure POPC bilayer,
it remains plausible that its function could be influenced by different lipid species, especially due
to the presence of multiple charged residues around the TMD/ECD interface which might inter-
act differently with different lipid head groups. Further experiments would be needed to confirm
whether the state dependence observed in simulations is also lipid-dependent. It is possible that
certain types of lipids bind in one but not the other state, or that certain states are stabilized by a
particular lipid type.

In simulations, we were able to identify both long-lasting and highly occupied lipid sites by an-
alyzing a large number of parallel trajectories (500 instances of each subunit). Simulation frames
could further be classified into closed or open states by projection onto a previously published
Markov state model of GLIC gating, which had been validated against electrophysiology record-
ings and structural features (*Bergh et al., 2021*). In particular, as simulations were initiated by
placing the GLIC structures in generated POPC bilayers, it is reassuring that lipid binding sites iden-
tified in simulations appear to correspond closely to densities representing lipids bound tightly
enough in the cryo-EM data to be retained during reconstitution (*Figure 1*—**??**C, D). The resulting
occupancy plots enabled confident distinction between functional annotations while preserving
the protein's flexibility and capacity to transition between states throughout the simulations. Inter-
estingly, the occupational densities varied remarkably little between resting and activating condi-
tions (*Figure 1*—**??**), indicating state- rather than pH-dependence in lipid interactions, also further
justifying the approach of merging closed-state GLIC cryo-EM datasets collected at different pH
conditions to resolve lipids.

The three lipids identified in the inner leaflet may play primarily structural roles, as they did not
vary substantially between closed and open states (*Figure 5*). Principal-face and intersubunit sites
in our closed cryo-EM structure were largely superimposable with those in open X-ray structures,
and all three sites contained comparable densities in closed- and open-state simulation frames
(*Figure 3*A, *Figure 1*—**??**D). A particularly stable contact in this leaflet featured deep intercalation of
a lipid tail between channel subunits (*Figure 4*—**??**), as represented by lipid interactions persisting
over 1 μs at residue T274, the longest-lasting in the entire protein (*Figure 2*). A lipid in this site was
also reported in previous open-state X-ray structures of GLIC (*Bocquet et al., 2009*; *Hu et al., 2018*).
Removal of the sidechain at W217 substantially abbreviated interactions with its buried neighbor
T274 (*Figure 3*), suggesting this hydrophobic contact helps to stabilize lipid penetration. Notably,
the same W217A substitution was previously shown to suppress GLIC function and was consid-
ered the most impactful of several aromatic residues on the transmembrane surface (*Therien and
Baenziger, 2017*). Interestingly, a lack of double bonds in the intercalating tail was associated with
the lipid head group interacting more closely with the protein surface, indicating a possible role
for saturation in overall lipid-binding modes (*Figure 3*). The relevance of lipid intercalation and sat-
uration has been difficult to establish on the basis of structural data, given that resolved lipid tails
are often limited to a few hydrocarbons beyond the glycerol group. An important extension of this
work may be to apply similar analyses to simulations of mixed-lipid systems, albeit at substantially
higher computational cost.

Lipid interactions in the outer leaflet were more variable, both between functional states and ex-
perimental methods, possibly reflecting larger conformational changes of the protein in this region.

At the complementary-face site - one of the most consistently documented in previous X-ray structures - simulations indicated that lipid tails could penetrate deeper into the helical bundle upon channel opening (*Figure 5*B), and even further in a state associated with desensitization (*Figure 4*—**??**A). Interestingly, buried portions of this site have been shown to host allosteric modulators such as propofol, substituting for lipid-tail interactions (*Heusser et al., 2018*). State dependence was also suggested at the principal-face site, where the lipid head preferred a pose tilted in towards M3 in the closed state, but tilted out towards M4 in the open state. However, our closed cryo-EM structure was more consistent with open-state simulations, and with a detergent built in an open X-ray structure (*Hu et al., 2018*). Indeed, the outward orientation was also sampled in closed-state simulations, albeit with lower frequency (*Figure 4*—**??**D). The relevance of the inward-facing pose, possibly in the context of an alternative lipid head group, remains to be determined. The most provocative indication of state dependence was at the outer subunit interface, which did not contain strong density in either cryo-EM or simulations data for the closed state (*Figure 5*A). In contrast, lipid occupancy at this site was evident in open simulations (*Figure 5*B), including state-dependent contacts with P250, a conserved M2–M3 residue implicated in channel gating *Kaczor et al.* (*2022*). Although lipids have yet to be resolved at this intersubunit site in GLIC, they have been reported in open-state structures of both ELIC (*Petroff et al., 2022*) and GluCl *Althoff et al.* (*2014*), and in a desensitized neuronal GABA(A) receptor *Laverty et al.* (*2019*). The state-dependent binding event at this site preceded pore opening in MSMs, where lipid binding coincided with crossing a smaller energy barrier between closed and intermediate states, followed by pore opening at the main energy barrier between intermediate and open states (*Figure 4*—**??**D). Further, since the P250-lipid interaction was characterized by relatively long residence times (*Figure 2*), it is possible this lipid interaction has a role to play in GLIC gating.

Finally, this work demonstrates the combined power of cryo-EM and MD simulations to characterize membrane-protein lipid interactions. In leveraging extensive MD simulations to build lipids for the first time into a closed-state structure of GLIC, we were also able to capture features of lipid-tail saturation and potential state dependence that were not available from structural data alone. Knowledge of such interactions may prove useful in the development of lipid-like drugs or lipid-focused treatments of diseases related to malfunction of pLGICs.

Methods and Materials

Molecular dynamics simulations and analysis

We analyzed previously published MSMs of GLIC gating under both resting and activating conditions (*Bergh et al., 2021*). Resting conditions corresponded to pH 7, at which GLIC is nonconductive in functional experiments, with all acidic residues modeled as deprotonated. Activating conditions corresponded to pH 4.6, at which GLIC is conductive and has been crystallized in an open state (*Bocquet et al., 2009*). These conditions were modeled by protonating a group of acidic residues (E26, E35, E67, E75, E82, D86, D88, E177, E243; H277 doubly protonated) as previously described (*Nury et al., 2011*). In order to sample lipid movement more extensively, each of the 100 simulations was extended to 1.7 μs, resulting in an additional 50 μs sampling compared to (*Bergh et al., 2021*). For state classification, the additional simulation frames were projected onto the MSMs trained on the previously published simulations.

Time-based measures of protein-lipid interactions, such as mean duration times and exchange of interactions, were calculated for the 100×1.7 μs-long simulations using prolintpy (*Sejdiu and Tieleman, 2021*) with a 4 Å interaction cutoff. Analysis of lateral lipid diffusion in individual simulations was carried out for two disjoint sets of lipids: the first lipid shell defined as lipids with any part within 4 Å of the protein surface (∼90 lipids), and bulk lipids consisting of all other lipids (∼280 lipids). Mean square displacements of each lipid set were calculated using GROMACS 2021.5 (*Abraham et al., 2015*) with contributions from the protein center of mass removed. Diffusion coefficients for each set, D_A, were calculated using the Einstein relation (*Equation 1*) by estimating the slope of the

369 linear curve fit to the data.

$$\lim_{t\to\infty}\langle \|\boldsymbol{r}_i(t) - \boldsymbol{r}_i(0)\|^2\rangle_{i\in A} = 4D_A t \qquad (1)$$

where $\boldsymbol{r}_i(t)$ is the coordinate of the center of mass of lipid i of set A at time t and D_A is the self-
diffusion coefficient.

To investigate state-dependent protein-lipid interactions, all trajectory frames were classified
according to the five clusters in (*Bergh et al., 2021*), resulting in 19555 and 21941 frames for the
closed and open protonated states, respectively, and 20632 and 13452 frames for the closed and
open deprotonated states, respectively. Occupied densities were calculated from the frames cor-
responding to each cluster using VMD's volmap tool (*Humphrey et al., 1996*) for a selection of all
whole lipids within 3 Å of the protein surface. In order to get representative snapshots of the
protein-lipid interactions, we calculated global correlations between the computational densities
and the selection of lipids used to derive the same densities using GROmaps (*Briones et al., 2019*)
and extracted 100 frames with the highest correlations. To ensure that all frames would not orig-
inate from a single trajectory, a maximum of 40% of the highest-correlating frames were allowed
to originate from a single trajectory.

To explore the effect of mutations, we ran sets of 8×1.7 μs simulations on open wild-type GLIC
(PDB ID: 4HFI) (*Sauguet et al., 2013*) with the modification W217A, T274A, or T274W. Each mu-
tated protein was independently embedded in a 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine
(POPC) lipid bilayer, solvated by water and 0.1 M NaCl. Acidic residues (E26, E35, E67, E75, E82,
D86, D88, E177, E243; H277 doubly protonated) were protonated to approximate the probable
pattern at pH 4.6, as previously described (*Nury et al., 2011*; *Bergh et al., 2021*). Constant pres-
sure and temperature relaxation was carried out in four steps, restraining all heavy atoms in the
first cycle, followed by only backbone atoms, then Cα atoms, and finally the M2 helices, for a to-
tal of 76 ns. Equilibration and production runs were performed using the Amber99SB-ILDN force
field with Berger lipids (*Berger et al., 1997*), together with the TIP3P water model. Temperature
coupling was achieved with velocity rescaling (*Bussi et al., 2007*) and pressure coupling with the
Parrinello-Rahman barostat (*Parrinello and Rahman, 1981*). Open-state mutagenesis simulations
were prepared and run with GROMACS versions 2018.4 and 2020.5 (*Abraham et al., 2015*). Sim-
ulations extended from those described in (*Bergh et al., 2021*) were prepared in a similar way as
described above but with GROMACS versions 2018.4 and 2019.3.

Conformational analysis and visualization of protein and lipids were performed with MDAnal-
ysis (*Michaud-Agrawal et al., 2011*; *Richard J. Gowers et al., 2016*), PyEMMA 2.5.7 (*Scherer et al.,*
2015), and VMD (*Humphrey et al., 1996*).

Cryo-EM sample preparation and data acquisition

Experiments used in this project were previously reported in (*Rovšnik et al., 2021*). Briefly, C43(DE3)
E. coli transformed with GLIC-MBP in vector pET-20b were cultured overnight at 37° C. Cells were
then inoculated 1:100 into the 2xYT media containing 100 μg/mL ampicillin and grown at 37° C until
they reached OD$_{600}$ = 0.7. Next, the cells were induced with 100 μM isopropyl-β-D-1-thiogalactopyranoside
(IPTG), and shaken overnight at 20° C. Membranes were harvested from cell pellets by sonication
and ultracentrifugation in buffer A (300 mM NaCl, 20 mM Tris-HCl pH 7.4) supplemented with 1
mg/mL lysozyme, 20 μg/mL DNase I, 5 mM MgCl$_2$, and protease inhibitors. At this point, the cells
were either frozen or immediately solubilized in 2% n-dodecyl-β-D-maltoside (DDM). Amylose affin-
ity resin (NEB) was used for purification of fusion protein in batch which was eluted in buffer B
(buffer A with 0.02% DDM) with 2–20 mM maltose followed by size exclusion chromatography in
buffer B. After overnight thrombin digestion, GLIC was isolated from its fusion partner by size ex-
clusion in buffer B at pH 7, or in buffer B with citrate at pH 5 or 3 substituted for Tris. The purified
protein was concentrated to 3–5 mg/mL by centrifugation.

Quantifoil 1.2/1.3 Cu 300 mesh grids (Quantifoil Micro Tools) were used for sample preparation.

The grids were glow-discharged in methanol vapor directly before 3 µl of sample was applied to them. Following a 1.5 s blot they were plunge-frozen into liquid ethane using a FEI Vitrobot Mark IV. Movies were collected on an FEI Titan Krios 300 kV microscope with a K2-Summit direct electron detector camera at nominal 165,000x magnification, equivalent to a pixel spacing of 0.82Å. A total dose of 40.8 e$^-$/Å^2 was used to collect 40 frames over 6 sec, with defocus values ranging from -2.0 to -3.8 µm.

Image processing

Processing was performed through the RELION 4.0-beta-2 pipeline (*Kimanius et al., 2021*). Data from three different grids, at pH 7, 5, and 3, were merged and processed together. Motion correction was performed with Relion's own implementation (*Zivanov et al., 2019*), followed by a defocus estimation from the motion-corrected micrographs using CtfFind4 (*Rohou and Grigorieff, 2015*). Following manual picking, initial 2D classification was performed to generate references for autopicking. Particles were extracted after autopicking and binned, followed by a 2D classification of the entire dataset. A smaller subset of particles was used to generate an initial model. All subsequent processing steps were done using 5-fold symmetry. The initial model was used in a consecutive 3D auto-refinement and the acquired alignment parameters were used to identify and remove noisy particles through multiple rounds of pre-aligned 2D- and 3D-classification. The final set of particles was then refined, using the best 3D reconstruction as reference.

Particle polishing and per-particle CTF parameters were estimated from the resulting reconstruction using RELION 4.0-beta-2. Global beam-tilt was estimated from the micrographs and correction applied. Micelle density was subtracted and the final 3D auto-refinement was performed using a soft mask covering the protein, followed by post-processing. Local resolution was estimated using the Relion implementation. Autosharpen map tool in PHENIX 1.19.2-4158 (*Adams et al., 2010*) was used to improve the visibility of peripheral lipid and detergent densities around the protein.

Model building

The model was built from a template cryo-EM structure determined at pH 7 (PDB ID: 6ZGD (*Rovšnik et al., 2021*)). A monomer of that model was fit to the reconstructed density and 5-fold symmetry was applied with PHENIX 1.19.2-4158 through NCS restraints detected from the reconstructed cryo-EM map, to generate a complete channel. The model was incrementally adjusted in COOT 0.9.6 EL (*Emsley and Cowtan, 2004*) and re-refined until conventional quality metrics were optimized in agreement with the reconstruction. Model statistics are summarized in *Table 1*.

Lipids were manually built in COOT by importing a canonical SMILES format of POPC (*Kim et al., 2021*) and adjusting it individually into the cryo-EM density in each of the sites associated with a single subunit, based in part on visual inspection of lipid densities from simulations, as described above. After building, 5-fold symmetry was applied to generate lipids at the same sites in the remaining four subunits.

Molecular visualizations were created with VMD (*Humphrey et al., 1996*) and ChimeraX (*Goddard et al., 2018*).

Acknowledgments

The authors would like to thank the Swedish Cryo-EM National Facility staff, especially Marta Carroni and Stefan Fleischmann from Stockholm and Michael Hall from Umeå, for kind assistance with data collection. This work was supported by the Knut and Alice Wallenberg Foundation, and the Swedish Research Council (2019-04433, 2021-05806), the Swedish e-Science Research Centre (SeRC), the BioExcel Center of Excellence (EU-823830), and the Sven and Lily Lawskis Fond (UR). Cryo-EM data were collected at the Swedish national cryo-EM facility funded by the Knut and Alice Wallenberg Foundation, Erling Persson, and Kempe Foundations. Computational resources were provided by the Swedish National Infrastructure for Computing (SNIC).

Data Availability

Cryo-EM density maps of the pentameric ligand-gated ion channel GLIC in detergent micelles have been deposited in the Electron Microscopy Data Bank under accession number EMD-15649. The deposition includes the cryo-EM sharpened and unsharpened maps, both half-maps and the mask used for final FSC calculation. Coordinates of the model have been deposited in the Protein Data Bank under accession number 8ATG. State-clustered simulation frames, computational lipid densities, parameter files, and full trajectories of mutant simulations as reported in this work can be accessed at https://doi.org/10.5281/zenodo.7058272. Previously published computational models and simulation files can be accessed at https://doi.org/10.5281/zenodo.5500174.

Competing interests

The authors declare no conflicts of interest.

References

Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, Lindahl E. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX. 2015; 1-2:19 – 25. http://www.sciencedirect.com/science/article/pii/S2352711015000059, doi: https://doi.org/10.1016/j.softx.2015.06.001.

Adams PD, Afonine PV, Bunkóczi G, Chen VB, Davis IW, Echols N, Headd JJ, Hung LW, Kapral GJ, Grosse-Kunstleve RW, et al. PHENIX: a comprehensive Python-based system for macromolecular structure solution. Acta Crystallographica Section D: Biological Crystallography. 2010; 66(2):213–221.

Althoff T, Hibbs RE, Banerjee S, Gouaux E. X-ray structures of GluCl in apo states reveal a gating mechanism of Cys-loop receptors. Nature. 2014; 512(7514):333–337.

Basak S, Schmandt N, Gicheru Y, Chakrapani S. Crystal structure and dynamics of a lipid-induced potential desensitized-state of a pentameric ligand-gated channel. eLife. 2017 mar; 6:e23886. https://doi.org/10.7554/eLife.23886, doi: 10.7554/eLife.23886.

Berger O, Edholm O, Jähnig F. Molecular dynamics simulations of a fluid bilayer of dipalmitoylphosphatidylcholine at full hydration, constant pressure, and constant temperature. Biophysical Journal. 1997; 72(5):2002–2013. https://www.sciencedirect.com/science/article/pii/S0006349597788453, doi: https://doi.org/10.1016/S0006-3495(97)78845-3.

Bergh C, Heusser SA, Howard R, Lindahl E. Markov state models of proton- and pore-dependent activation in a pentameric ligand-gated ion channel. eLife. 2021 oct; 10:e68369. https://doi.org/10.7554/eLife.68369, doi: 10.7554/eLife.68369.

Bocquet N, Nury H, Baaden M, Le Poupon C, Changeux JP, Delarue M, Corringer PJ. X-ray structure of a pentameric ligand-gated ion channel in an apparently open conformation. Nature. 2009; 457(7225):111–114.

Briones R, Blau C, Kutzner C, de Groot BL, Aponte-Santamaría C. GROmaps: A GROMACS-Based Toolset to Analyze Density Maps Derived from Molecular Dynamics Simulations. Biophysical Journal. 2019; 116(1):4–11. https://www.sciencedirect.com/science/article/pii/S0006349518344485, doi: https://doi.org/10.1016/j.bpj.2018.11.3126.

Bussi G, Donadio D, Parrinello M. Canonical sampling through velocity rescaling. The Journal of Chemical Physics. 2007; 126(1):014101. doi: 10.1063/1.2408420.

Carswell CL, Sun J, Baenziger JE. Intramembrane Aromatic Interactions Influence the Lipid Sensitivities of Pentameric Ligand-gated Ion Channels *. Journal of Biological Chemistry. 2015 Jan; 290(4):2496–2507. https://doi.org/10.1074/jbc.M114.624395, doi: 10.1074/jbc.M114.624395.

Cordero-Morales JF, Vásquez V. How lipids contribute to ion channel function, a fat perspective on direct and indirect interactions. Current opinion in structural biology. 2018; 51:92–98.

Criado M, Eibl H, Barrantes FJ. Effects of lipids on acetylcholine receptor. Essential need of cholesterol for maintenance of agonist-induced state transitions in lipid vesicles. Biochemistry. 1982 Jul; 21(15):3622–3629. https://doi.org/10.1021/bi00258a015, doi: 10.1021/bi00258a015.

Crnjar A, Molteni C. Cholesterol content in the membrane promotes key lipid-protein interactions in a pentameric serotonin-gated ion channel. Biointerphases. 2020; 15(6):061018. https://doi.org/10.1116/6.0000561, doi: 10.1116/6.0000561.

daCosta CJB, Ogrel AA, McCardy EA, Blanton MP, Baenziger JE. Lipid-Protein Interactions at the Nicotinic Acetylcholine Receptor: A FUNCTIONAL COUPLING BETWEEN NICOTINIC RECEPTORS AND PHOSPHATIDIC ACID-CONTAINING LIPID BILAYERS*. Journal of Biological Chemistry. 2002; 277(1):201–208. https://www.sciencedirect.com/science/article/pii/S002192582087999X, doi: https://doi.org/10.1074/jbc.M108341200.

Dietzen NM, Arcario MJ, Chen LJ, Petroff I John T, Moreland KT, Krishnan K, Brannigan G, Covey DF, Cheng WW. Polyunsaturated fatty acids inhibit a pentameric ligand-gated ion channel through one of two binding sites. eLife. 2022 jan; 11:e74306. https://doi.org/10.7554/eLife.74306, doi: 10.7554/eLife.74306.

Duncan AL, Song W, Sansom MS. Lipid-dependent regulation of ion channels and G protein–coupled receptors: insights from structures and simulations. Annual Review of Pharmacology and Toxicology. 2020; 60:31–50.

Dämgen MA, Biggin PC. State-dependent protein-lipid interactions of a pentameric ligand-gated ion channel in a neuronal membrane. PLOS Computational Biology. 2021 02; 17(2):1–24. https://doi.org/10.1371/journal.pcbi.1007856, doi: 10.1371/journal.pcbi.1007856.

Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. Acta crystallographica section D: biological crystallography. 2004; 60(12):2126–2132.

Fong TM, McNamee MG. Correlation between acetylcholine receptor function and structural properties of membranes. Biochemistry. 1986 Feb; 25(4):830–840. doi: 10.1021/bi00352a015.

Gharpure A, Teng J, Zhuang Y, Noviello CM, Walsh Jr RM, Cabuco R, Howard RJ, Zaveri NT, Lindahl E, Hibbs RE. Agonist selectivity and ion permeation in the $\alpha3\beta4$ ganglionic nicotinic receptor. Neuron. 2019; 104(3):501–511.

Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE. UCSF ChimeraX: Meeting modern challenges in visualization and analysis. Protein Science. 2018; 27(1):14–25.

Heusser SA, Lycksell M, Wang X, McComas SE, Howard RJ, Lindahl E. Allosteric potentiation of a ligand-gated ion channel is mediated by access to a deep membrane-facing cavity. Proceedings of the National Academy of Sciences. 2018; 115(42):10672–10677. https://www.pnas.org/doi/abs/10.1073/pnas.1809650115, doi: 10.1073/pnas.1809650115.

Hu H, Ataka K, Menny A, Fourati Z, Sauguet L, Corringer PJ, Koehl P, Heberle J, Delarue M. Electrostatics, proton sensor, and networks governing the gating transition in GLIC, a proton-gated pentameric ion channel. Proceedings of the National Academy of Sciences. 2018; 115(52):E12172–E12181.

Humphrey W, Dalke A, Schulten K. VMD – Visual Molecular Dynamics. Journal of Molecular Graphics. 1996; 14:33–38.

Richard J Gowers, Max Linke, Jonathan Barnoud, Tyler J E Reddy, Manuel N Melo, Sean L Seyler, Jan Domański, David L Dotson, Sébastien Buchoux, Ian M Kenney, Oliver Beckstein. MDAnalysis: A Python Package for the Rapid Analysis of Molecular Dynamics Simulations. In: Sebastian Benthall, Scott Rostrup, editors. *Proceedings of the 15th Python in Science Conference*; 2016. p. 98 – 105. doi: 10.25080/Majora-629e541a-00e.

Jakobi AJ, Wilmanns M, Sachse C. Model-based local density sharpening of cryo-EM maps. Elife. 2017; 6:e27131.

Kaczor PT, Michałowski MA, Mozrzymas JW. $\alpha1$ Proline 277 Residues Regulate GABAAR Gating through M2-M3 Loop Interaction in the Interface Region. ACS Chemical Neuroscience. 2022; .

Kim JJ, Hibbs RE. Direct Structural Insights into GABAA Receptor Pharmacology. Trends in Biochemical Sciences. 2021; 46(6):502–517. https://www.sciencedirect.com/science/article/pii/S0968000421000256, doi: https://doi.org/10.1016/j.tibs.2021.01.011.

Kim S, Chen J, Cheng T, Gindulyte A, He J, He S, Li Q, Shoemaker BA, Thiessen PA, Yu B, et al. PubChem in 2021: new data content and improved web interfaces. Nucleic acids research. 2021; 49(D1):D1388–D1395.

Kimanius D, Dong L, Sharov G, Nakane T, Scheres SH. New tools for automated cryo-EM single-particle analysis in RELION-4.0. Biochemical Journal. 2021; 478(24):4169–4185.

Labriola JM, Pandhare A, Jansen M, Blanton MP, Corringer PJ, Baenziger JE. Structural sensitivity of a prokaryotic
pentameric ligand-gated ion channel to its membrane environment. Journal of Biological Chemistry. 2013;
288(16):11294–11303.

Lamprakis C, Andreadelis I, Manchester J, Velez-Vega C, Duca JS, Cournia Z. Evaluating the Efficiency of the
Martini Force Field to Study Protein Dimerization in Aqueous and Membrane Environments. Journal of
Chemical Theory and Computation. 2021 May; 17(5):3088–3102. https://doi.org/10.1021/acs.jctc.0c00507,
doi: 10.1021/acs.jctc.0c00507.

Laverty D, Desai R, Uchański T, Masiulis S, Stec WJ, Malinauskas T, Zivanov J, Pardon E, Steyaert J, Miller KW, et al.
Cryo-EM structure of the human $\alpha 1 \beta 3 \gamma 2$ GABAA receptor in a lipid bilayer. Nature. 2019; 565(7740):516–520.

Levental I, Lyman E. Regulation of membrane protein structure and function by their lipid nano-
environment. Nature Reviews Molecular Cell Biology. 2022 Sep; https://doi.org/10.1038/s41580-022-00524-4,
doi: 10.1038/s41580-022-00524-4.

Masiulis S, Desai R, Uchański T, Serna Martin I, Laverty D, Karia D, Malinauskas T, Zivanov J, Pardon E, Kotecha
A, et al. GABAA receptor signalling mechanisms revealed by structural pharmacology. Nature. 2019;
565(7740):454–459.

Menny A, Lefebvre SN, Schmidpeter PA, Drege E, Fourati Z, Delarue M, Edelstein SJ, Nimigean CM, Joseph
D, Corringer PJ. Identification of a pre-active conformation of a pentameric channel receptor. Elife. 2017;
6:e23955.

Michaud-Agrawal N, Denning EJ, Woolf TB, Beckstein O. MDAnalysis: A toolkit for the analysis of molecular
dynamics simulations. Journal of Computational Chemistry. 2011; 32(10):2319–2327. https://onlinelibrary.
wiley.com/doi/abs/10.1002/jcc.21787, doi: https://doi.org/10.1002/jcc.21787.

Nury H, Van Renterghem C, Weng Y, Tran A, Baaden M, Dufresne V, Changeux JP, Sonner JM, Delarue M, Cor-
ringer PJ. X-ray structures of general anaesthetics bound to a pentameric ligand-gated ion channel. Nature.
2011 Jan; 469(7330):428–431. https://doi.org/10.1038/nature09647, doi: 10.1038/nature09647.

Parrinello M, Rahman A. Polymorphic transitions in single crystals: A new molecular dynamics method. Journal
of Applied Physics. 1981; 52(12):7182–7190. doi: 10.1063/1.328693.

Petroff JT, Dietzen NM, Santiago-McRae E, Deng B, Washington MS, Chen LJ, Trent Moreland K, Deng Z, Rau M,
Fitzpatrick JA, et al. Open-channel structure of a pentameric ligand-gated ion channel reveals a mechanism
of leaflet-specific phospholipid modulation. Nature Communications. 2022; 13(1):1–16.

Phillips R, Ursell T, Wiggins P, Sens P. Emerging roles for lipids in shaping membrane-protein function. Nature.
2009; 459(7245):379–385.

Popot JL, Cartaud J, Changeux JP. Reconstitution of a Functional Acetylcholine Receptor. European Journal of
Biochemistry. 1981; 118(2):203–214. https://febs.onlinelibrary.wiley.com/doi/abs/10.1111/j.1432-1033.1981.
tb06388.x, doi: https://doi.org/10.1111/j.1432-1033.1981.tb06388.x.

Poveda JA, Giudici AM, Renart ML, Morales A, González-Ros JM. Towards understanding the molecular basis
of ion channel modulation by lipids: Mechanistic models and current paradigms. Biochimica Et Biophysica
Acta (BBA)-Biomembranes. 2017; 1859(9):1507–1516.

Rahman MM, Teng J, Worrell BT, Noviello CM, Lee M, Karlin A, Stowell MH, Hibbs RE. Structure of the native
muscle-type nicotinic receptor and inhibition by snake venom toxins. Neuron. 2020; 106(6):952–962.

Rohou A, Grigorieff N. CTFFIND4: Fast and accurate defocus estimation from electron micrographs. Journal of
structural biology. 2015; 192(2):216–221.

Rovšnik U, Zhuang Y, Forsberg BO, Carroni M, Yvonnesdotter L, Howard RJ, Lindahl E. Dynamic closed states
of a ligand-gated ion channel captured by cryo-EM and simulations. Life Science Alliance. 2021; 4(8). https:
//www.life-science-alliance.org/content/4/8/e202101011, doi: 10.26508/lsa.202101011.

Sauguet L, Poitevin F, Murail S, Van Renterghem C, Moraga-Cid G, Malherbe L, Thompson AW, Koehl P, Cor-
ringer PJ, Baaden M, Delarue M. Structural basis for ion permeation mechanism in pentameric ligand-gated
ion channels. The EMBO Journal. 2013; 32(5):728–741. https://www.embopress.org/doi/abs/10.1038/emboj.
2013.17, doi: 10.1038/emboj.2013.17.

Sauguet L, Shahsavar A, Poitevin F, Huon C, Menny A, Àkos Nemecz, Haouz A, Changeux JP, Corringer PJ, De-
larue M. Crystal structures of a pentameric ligand-gated ion channel provide a mechanism for activation.
Proceedings of the National Academy of Sciences. 2014; 111(3):966–971. https://www.pnas.org/doi/abs/10.
1073/pnas.1314997111, doi: 10.1073/pnas.1314997111.

Scherer MK, Trendelkamp-Schroer B, Paul F, Pérez-Hernández G, Hoffmann M, Plattner N, Wehmeyer C, Prinz
JH, Noé F. PyEMMA 2: A Software Package for Estimation, Validation, and Analysis of Markov Models. Journal
of Chemical Theory and Computation. 2015 Oct; 11:5525–5542. http://dx.doi.org/10.1021/acs.jctc.5b00743,
doi: 10.1021/acs.jctc.5b00743.

Sejdiu BI, Tieleman DP. ProLint: a web-based framework for the automated data analysis and visualization
of lipid–protein interactions. Nucleic Acids Research. 2021 05; https://doi.org/10.1093/nar/gkab409, doi:
10.1093/nar/gkab409, gkab409.

Sharp L, Brannigan G. Spontaneous lipid binding to the nicotinic acetylcholine receptor in a native mem-
brane. The Journal of Chemical Physics. 2021; 154(18):185102. https://doi.org/10.1063/5.0046333, doi:
10.1063/5.0046333.

Sridhar A, Lummis SCR, Pasini D, Mehregan A, Brams M, Kambara K, Bertrand D, Lindahl E, Howard RJ,
Ulens C. Regulation of a pentameric ligand-gated ion channel by a semiconserved cationic lipid-binding
site. Journal of Biological Chemistry. 2021 Aug; 297(2). https://doi.org/10.1016/j.jbc.2021.100899, doi:
10.1016/j.jbc.2021.100899.

Tasneem A, Iyer LM, Jakobsson E, Aravind L. Identification of the prokaryotic ligand-gated ion channels and
their implications for the mechanisms and origins of animal Cys-loop ion channels. Genome biology. 2005;
6(1):1–12.

Terwilliger TC, Ludtke SJ, Read RJ, Adams PD, Afonine PV. Improvement of cryo-EM maps by density modifica-
tion. Nature methods. 2020; 17(9):923–927.

Therien JPD, Baenziger JE. Pentameric ligand-gated ion channels exhibit distinct transmembrane domain
archetypes for folding/expression and function. Scientific Reports. 2017 Mar; 7(1):450. https://doi.org/10.
1038/s41598-017-00573-2, doi: 10.1038/s41598-017-00573-2.

Thompson MJ, Baenziger JE. Structural basis for the modulation of pentameric ligand-gated
ion channel function by lipids. Biochimica et Biophysica Acta (BBA) - Biomembranes. 2020;
1862(9):183304. https://www.sciencedirect.com/science/article/pii/S0005273620301358, doi:
https://doi.org/10.1016/j.bbamem.2020.183304.

Walsh RM, Roh SH, Gharpure A, Morales-Perez CL, Teng J, Hibbs RE. Structural principles of distinct assemblies
of the human $\alpha 4\beta 2$ nicotinic receptor. Nature. 2018; 557(7704):261–265.

Zhao Y, Liu S, Zhou Y, Zhang M, Chen H, Eric Xu H, Sun D, Liu L, Tian C. Structural basis of human $\alpha 7$ nicotinic
acetylcholine receptor activation. Cell research. 2021; 31(6):713–716.

Zhu S, Noviello CM, Teng J, Walsh RM, Kim JJ, Hibbs RE. Structure of a human synaptic GABAA receptor. Nature.
2018; 559(7712):67–72.

Zivanov J, Nakane T, Scheres SH. A Bayesian approach to beam-induced motion correction in cryo-EM single-
particle analysis. IUCrJ. 2019; 6(1):5–17.

Table 1. Cryo-EM data collection and model refinement statistics

Data Collection	
Microscope	FEI Titan Krios
Magnification	165,000
Voltage (kV)	300
Electron exposure (e$^-$/Å^2)	~ 40
Defocus Range (μm)	-2.6 – -3.8
Pixel Size (Å)	0.82
Symmetry Imposed	C5
Number of Images	~ 18982
Particles Picked	$\sim 2,7$ mil
Particles Refined	16,586
Refinement	
Resolution (Å)	2.9
FSC Threshold	0.143
Map sharpening B-factor	-80
Model Composition	
Non-hydrogen Protein atoms	12,010
Protein Residues	1555
Ligands	0
B-factor (Å^2)	144
RMSD	
Bond Lengths (Å)	0.004
Bond Angles (°)	0.584
Validation	
Molprobity Score	1.84
Clashscore	6.82
Poor Rotamers (%)	0
Ramachandran Plot	
Favored (%)	92.6
Allowed (%)	7.4
Outliers (%)	0