Title

Leaf decomposing fungi influence *Saccharomyces paradoxus* growth across carbon environments

Authors

Samer El-Khatib (0000-0003-1531-3002)[1], Madeleine G. Lambert (0000-0002-5652-2305)[1,2], Meghan N. Reed (0000-0002-6661-553X)[1,3], Melane Brito Goncalves (0000-0003-3707-4583)[1,4], Primrose J. Boynton[1*] (0000-0003-0720-8966)

1) Wheaton College, Norton, MA, USA
2) Georgetown University Medical Center, Washington, DC, USA
3) Lonza, Portsmouth, NH, USA
4) Rhode Island College, Providence, RI, USA
* Submitting and corresponding author: boynton_primrose@wheatoncollege.edu

Abstract

Saccharomyces paradoxus is a model organism in ecology and evolution. However, its metabolism in its native habitat remains mysterious: it is frequently found growing on leaf litter, a habitat with few carbon sources that *S. paradoxus* can metabolize. We hypothesized that leaf-decomposing fungi from the same habitat break down the cellulose in leaf litter extracellularly and release glucose, supporting *S. paradoxus* growth. We found that facilitation by leaf-decomposing fungi was possible on cellulose and inhibition was common on glucose, suggesting diverse interactions between *S. paradoxus* and other fungi that have the potential to support *S. paradoxus* in nature.

Figure

Legend

Figure 1: Leaf litter decomposing fungi and their interactions with *S. paradoxus*. **(a)** Five filamentous fungi isolated from forest oak leaf litter growing on solid glucose medium. **(b)** The five fungi in flasks of liquid cellulose (top) and liquid glucose (bottom) media. **(c)** The total number of *S. paradoxus* colony forming units (CFUs) grown after 26 hours in 5 ml cellulose medium conditioned by each of five litter-decomposing fungi, or in control unaltered media (five replicate *S. paradoxus* strains per treatment; each *S. paradoxus* strain is represented by a single dot per treatment). Conditioned media were prepared after 96 hours of fungal growth in liquid media, followed by filter-sterilization of used media. *Nemania* increased the number of *S. paradoxus* CFUs to 4.7 times the control (one-way ANOVA adjusted R^2 = 0.72, F = 15.6, degrees of freedom = 5,24, p < 10^{-6}). In contrast, *S. paradoxus* produced equivalent numbers of CFUs in the unaltered control medium as in media conditioned by *Trichoderma, Biscogniauxia, Coniochaeta,* and *Chaetomium*. **(d)** The total number of *S. paradoxus* CFUs grown in glucose medium conditioned by each of the five litter-decomposing fungi, plus an untreated control. *S.*

paradoxus produced considerably fewer CFUs after conditioning by *Trichoderma* (0.06 times the number of CFUs produced in untreated glucose medium; one-way ANOVA adjusted $R^2 = 0.81$, $F = 25.69$, degrees of freedom = 5,24, $p < 10^{-8}$). *S. paradoxus* also produced fewer CFUs after conditioning by *Biscogniauxia, Coniochaeta, and Chaetomium* than in unaltered glucose medium (0.61 times the control number of CFUs). *Nemania* did not significantly change the mean number of *S. paradoxus* CFUs compared to the control. The y-axes are presented on a $\log_{10}$ scale in figures 1c and 1d, and different letters in each of figures 1c and 1d indicate significantly different means (Tukey post-hoc HSD $p < 0.05$).

Description

Leaf litter decomposers may solve the mystery of *Saccharomyces paradoxus* metabolism in nature. *S. paradoxus* is a wild yeast and the closest described relative of *Saccharomyces cerevisiae*, the most widely researched yeast and one of the most widely researched eukaryotic model organisms (Goffeau et al. 1996; Boynton and Greig 2014). *S. paradoxus* can be found growing in leaf litter and soil close to oak trees, and less abundantly on oak tree bark (Glushakova et al. 2007; Sampaio and Gonçalves 2008; Kowallik and Greig 2016). It can grow both aerobically and anaerobically, utilizing simple sugars such as glucose as carbon sources. One common source of simple sugars in forests, honeydew (*i.e.,* aphid excretions), has been proposed as a potential carbon source for *S. paradoxus* (Brysch-Herzberg and Seidel 2017), but surveys in New Zealand and Germany have not been able to uncover any association between *Saccharomyces* and honeydew (Serjeant et al. 2008; Kowallik 2015). *S. paradoxus* also cannot metabolize cellulose, one of the most common polymers in leaf litter (Van Rensburg et al. 1998; Dashko et al. 2014). The primary component of the plant leaf cell walls in leaf litter is cellulose, and glucose and other simple monosaccharides are generally scarce on leaf litter (Sampaio and Gonçalves 2008; Keegstra 2010). It is unclear how *S. paradoxus* survives and reproduces in such cellulose-rich and glucose-poor environments: here, we investigate how interactions with nearby fungi might facilitate *S. paradoxus* metabolism.

Diverse leaf fungal communities are the main decomposers of leaf litter, playing an essential role globally in plant polymer decomposition and the carbon cycle (Kjøller and Struwe 1982; Benocci et al. 2017). Fungi that occupy the same habitat as *S. paradoxus* decompose leaf litter through extracellular cellulolytic enzymes, such as cellulases and glucanases (Krishna and Mohan 2017). Previous research has shown that nearby microorganisms can facilitate or inhibit *S. paradoxus* growth on oak bark (Kowallik et al. 2015). However, the mechanisms behind these interactions remain unknown. We hypothesize that cellulose-degrading fungi facilitate the survival of *S. paradoxus* in leaf litter through extracellular breakdown of leaf cellulose and the consequent release of more accessible sugars such as glucose. Conversely, we hypothesize that cellulose-degrading fungi might consume free glucose, a carbon source otherwise conducive to *S. paradoxus* growth, and inhibit *S. paradoxus* when cellulose is not available.

To test our hypotheses, we first isolated five *S. paradoxus* strains and five leaf-degrading fungal isolates with diverse morphologies (**Figure 1a-b**) from partially decomposed oak leaves in a New England forest. The leaf-degrading fungi were identified using ribosomal DNA sequences as members of the genera *Trichoderma*, *Biscogniauxia*, *Coniochaeta, Chaetomium,* and

Nemania. These genera include characterized leaf endophytes, leaf litter decomposers, and wood decomposing fungi (Nugent et al. 2005; Fukasawa et al. 2011; Osono et al. 2011; Helaly et al. 2018; Mäkelä et al. 2021). We confirmed that all five litter fungi, but not *S. paradoxus*, can grow on a growth medium containing cellulose as the sole carbon source, and then further explored the interactions between *S. paradoxus* and the other fungi in different carbon environments.

We carried out spent-media experiments to test our hypothesis that the leaf litter decomposing fungi would facilitate *S. paradoxus* growth on cellulose by breaking down the polymer into simpler sugars. We first allowed each fungus to grow on cellulose medium and to change the environment, including presumably through the release of cellulolytic enzymes or metabolites into the medium. Cellulose medium, used for spent-media experiments, differs from cellulose screening medium, used to confirm growth with cellulose as a sole carbon source, in that it contains peptone and can support some *S. paradoxus* growth. We then filtered out each fungus and grew *S. paradoxus* on the used medium, quantifying *S. paradoxus* growth using colony-forming unit (CFU) counts for each fungal treatment after 26 hours. One of the five fungi, *Nemania*, increased the mean number of *S. paradoxus* CFUs produced to 4.7 times the control untreated medium, but the other four fungi had no significant effect on growth (**Figure 1c,** ANOVA adjusted R^2 = 0.72, F = 15.6, degrees of freedom = 24,5, $p < 10^{-6}$; Tukey post-hoc comparison between the control and *Nemania* treatments $p < 10^{-5}$).

To further investigate the effects that leaf litter decomposing fungi have on *S. paradoxus* growth, we repeated the experiment with glucose medium, an environment permissive to *S. paradoxus* growth. We hypothesized that the litter decomposing fungi would use up free glucose, inhibiting *S. paradoxus* growth. In glucose medium, the previously identified facilitator, *Nemania*, had no significant effect on *S. paradoxus* growth, while *Trichoderma* had an extremely inhibitory effect, decreasing the mean number of *S. paradoxus* CFUs to 0.06 times the control. *Biscogniauxia, Coniochaeta,* and *Chaetomium* all also inhibited growth compared to the control but to a lesser extent, decreasing the mean number of CFUs to 0.61 times the control (**Figure 1d,** ANOVA adjusted R^2 = 0.8098, F = 25.69, degrees of freedom = 24,5, $p < 10^{-8}$; Tukey post-hoc $p = 10^{-7}$ for the comparison between *the Trichoderma* and the control, and $p \leq 0.036$ for comparisons between each of *Biscogniauxia*, *Coniochaeta*, and *Chaetomium* and control). The varying degrees of inhibition by different fungi on glucose indicates that certain leaf litter species are more efficiently antagonistic to *S. paradoxus* than others.

Our results partially support our hypotheses and highlight the diversity of possible interactions between *S. paradoxus* and other litter decomposing fungi. *S. paradoxus* growth on cellulose was facilitated by at least one litter decomposing fungus, while growth on glucose was inhibited to varying degrees by four of the fungi. *Nemania* likely facilitates *S. paradoxus* on cellulose by releasing cellulolytic enzymes and enabling opportunistic *S. paradoxus* glucose consumption after the cellulose is broken down. *Nemania*'s lack of inhibition in glucose could also indicate that *Nemania* doesn't consume glucose quickly enough to prevent *S. paradoxus* from accessing it, lending a rare but well-timed opportunity for *S. paradoxus* to grow in unfavorable carbon environments such as leaf litter.

Other interactions between *S. paradoxus* and leaf decomposing fungi (**Figure 1c-d**) may be due to more complex interactions, possibly involving primary metabolism or secreted antagonistic molecules. We speculate that cellulose-degrading fungi might be especially effective at removing glucose from the environment when cellulolytic enzymes are not needed. High concentrations of glucose repress cellulolytic enzyme production (Strauss et al. 1995; Suto and Tomita, 2001), allowing more resources to be devoted to other cell functions, perhaps including glucose uptake. This increased glucose uptake and removal from the environment may allow litter decomposing fungi to more effectively inhibit *S. paradoxus* in glucose than cellulose (**Figures 1c-d**). Litter decomposing fungi might also release metabolites that could directly inhibit *S. paradoxus*. Qualitative observations of differences in pigmentation among the fungi in cellulose and glucose growth media (**Figure 1b**) support this mechanism. Additionally, fungi in the *Sordariomycetes* class, to which *Chaetomium, Coniochaeta*, and *Trichoderma* belong, and others in the *Xylariales* order, to which *Biscogniauxia* and *Nemania* belong, release active secondary metabolites, including pigments (Stamps et al. 2015; Bills and Gloer, 2016; Helaly et al. 2018; Becker and Stadler, 2021), with some fungi doing so when associated with other organisms (Lamacchia et al. 2016). In this case, fungi growing in glucose might be able to expend more energy towards the production of active secondary antagonistic metabolites because cellulolytic enzyme production is not necessary.

S. paradoxus growth on different carbon sources is contingent on interactions among *S. paradoxus*, leaf litter fungi, and the environment. Litter decomposers can create conditions conducive to *S. paradoxus*, or they can inhibit *S. paradoxus*, depending on the nutrient environment. *S. paradoxus* growth in its natural cellulose environment most likely depends on the presence of facilitator fungi such as *Nemania*, the lack of antagonistic species such as *Trichoderma*, and well-timed opportunistic glucose consumption. Future research will explore the roles of cellulases and secondary metabolites in interactions between litter decomposing fungi and *S. paradoxus* in nature.

Methods
Isolation and identification of *S. paradoxus* and leaf litter decomposing fungi
All fungi were isolated from oak leaf litter in a mixed temperate forest in Norton, MA, USA (coordinates 41.96, -71.18). Five *S. paradoxus* strains were isolated from leaf litter in the summer of 2021 as in Boynton et al. (2019). First, 134 yeast colonies were isolated by directly plating soil directly underneath leaf litter onto the isolation medium PIM1 (Sniegowski et al. 2002; Boynton et al., 2019). All media recipes are provided in the reagents section below. Colonies were then screened for *Saccharomyces*-like spore morphology, and *S. paradoxus* colonies were identified using *S. paradoxus* specific PCR primers, as in Robinson et al. (2016), or by sequencing the ribosomal internal transcribed spacer region using primers ITS1 and ITS4 (White et al. 1990).

Filamentous fungi were isolated from leaf litter next to three oak trees; partially degraded fallen leaves (edges were degraded but not the center) were collected in the fall of 2021, aseptically contained in sterile plastic bags, and stored at approximately 4°C for up to 24 hours before

fungal isolation. Seven-millimeter diameter leaf discs were punched out from each leaf three times with a sterile hole punch. Discs were surface sterilized as in Osono & Takeda (2001) and were arranged on filamentous fungi isolation medium and incubated at 28°C until fungal mycelium could be seen growing from leaf cut edges. Five fungi were subcultured onto glucose medium and pure cultures were maintained on glucose medium for the duration of the study. Growth on cellulose as a sole carbon source was confirmed by observing growth on cellulose screening medium. The fungi were identified by sequencing portions of large ribosomal subunit and ribosomal internal transcribed spacer DNA using primers detailed in the reagents section below. Fungi were identified to genera using NCBI BLAST (Altschul et al. 1990; Sayers et al. 2022) and querying databases associated with 26S RNA and internal transcribed spacer regions from Fungi type and reference material. Internal transcribed spacer and partial 26S RNA sequences were deposited in GenBank (Accession numbers OQ145425-OQ145429, OQ145437-145441).

Media conditioning experiments
To understand the impacts of each leaf litter fungus on *S. paradoxus*, we grew *S. paradoxus* on cellulose and glucose media that had first been conditioned by each leaf litter fungus. Each of the leaf litter fungi was inoculated into a flask containing 75 mL of liquid medium. Flasks were incubated at 28°C for 96 hours with 140 rpm shaking. After growth, the conditioned medium from each flask was filtered through a 0.2 μm filter, dispensed into 5 tubes (5 ml conditioned medium per tube), and each tube was inoculated with one of five *S. paradoxus* strains by transferring colonies into each tube from a petri dish using a sterile wooden stick. Tubes were incubated at 28°C with 140 rpm shaking for 26 hours, after which *S. paradoxus* colony forming units (CFUs) were counted on YEPD solid medium. In untreated cellulose medium, *S. paradoxus* could grow a little bit, most likely by using peptone in the medium as a carbon source. One-way ANOVAs with Tukey's HSD post-hoc tests were performed for each medium with the conditioning fungus as a predictor and number of *S. paradoxus* cells produced as a response variable. Statistical tests were performed using R version 4.1.2, with plots produced using the ggplot2 package (Wickham 2016; R Core Team, 2021).

Reagents
Fungal strains

Type of fungus	Genotype	Available from
Yeast Strain 130	*Saccharomyces paradoxus*	The corresponding author
Yeast Strain 132	*Saccharomyces paradoxus*	The corresponding author
Yeast Strain 154	*Saccharomyces paradoxus*	The corresponding author
Yeast Strain 200	*Saccharomyces paradoxus*	The corresponding author
Yeast Strain 293	*Saccharomyces paradoxus*	The corresponding author
Leaf litter decomposing fungus strain 3a	*Trichoderma sp.*	The corresponding author
Leaf litter decomposing fungus strain 4a	*Biscogniauxia sp.*	The corresponding author
Leaf litter decomposing fungus strain 4b	*Coniochaeta sp.*	The corresponding author
Leaf litter decomposing fungus strain 6b	*Chaetomium sp.*	The corresponding author
Leaf litter decomposing fungus strain 8b	*Nemania sp.*	The corresponding author

Primers

Name	Sequence (5' to 3')	Target	Citation
cepC6f	GAACCTGTCGAAGTTGTTAATGC	*Saccharomyces* rapidly evolving centromere	Robinson et al. 2016
cepC6r	TRGAAGGTTTCTTTGGWGCCAT	*Saccharomyces* rapidly evolving centromere	Robinson et al. 2016
parC15f	TCATATTTATAACAAGCGATCAAAGC	*S. paradoxus* rapidly evolving centromere	Robinson et al. 2016
parC15r	GAAGTACATCCTGATTTTTAGAAGCC	*S. paradoxus* rapidly evolving centromere	Robinson et al. 2016
ITS1	TCCGTAGGTGAACCTGCGG	Ribosomal internal transcribed spacer	White et al. 1990
ITS4	TCCTCCGCTTATTGATATGC	Ribosomal internal transcribed spacer	White et al. 1990
NL1	GCATATCAATAAGCGGAGGAAAAG	Large ribosomal subunit	O'Donnell 1992
NLr	GGTCCGTGTTTCAAGACGG	Large ribosomal subunit	O'Donnell 1992

Media

Medium	Ingredients	Citation
Filamentous Fungi Isolation Medium	peptone 5.0 g/L, glucose 10.0 g/L, monopotassium phosphate 1.0 g/L, magnesium sulfate 0.5 g/L, agar 20.0 g/L, chloramphenicol 54 µL of 20 mg/ml	modified from Sun et al. 2020
Cellulose Medium	sodium carboxymethylcellulose 10.0 g/L, ammonium sulfate 4.0 g/L, monopotassium phosphate 2.0 g/L, magnesium sulfate, heptahydrate 0.5 g/L, peptone 1.0 g/L	modified from Sun et al. 2020
Cellulose Screening Medium	sodium carboxymethylcellulose 10.0 g/L, ammonium sulfate 5.0 g/L, monopotassium phosphate 2.0 g/L, magnesium sulfate, heptahydrate 0.5 g/L, yeast nitrogen base w/ amino acids 1.7 g/L, agar 15 g/L	modified from Cellulose Medium (above)
Glucose medium (liquid)	monopotassium phosphate 1.0 g/L, magnesium sulfate 0.5 g/L, peptone 5.0 g/L, glucose (dextrose) 10.0 g/L	modified from Sun et al. 2020
Glucose medium (solid)	monopotassium phosphate 1.0 g/L, magnesium sulfate 0.5 g/L, peptone 5.0 g/L, glucose (dextrose) 10.0 g/L, agar 20.0 g/L	modified from Sun et al. 2020
YEPD medium	yeast extract 10 g/L, peptone 20 g/L, dextrose (glucose) 20 g/L, agar 25 g/L	modified from Liti et al. 2017
PIM1 medium	yeast extract 3 g/L, peptone 5 g/L, sucrose 10 g/L, malt extract 3 g/L, chloramphenicol 1 mg/L, ethanol 80 ml/L, 1 M HCl 5.2 ml/L, agar 20 g/L	Sneigowski et al. 2002; Boynton et al. 2019

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. J Mol Biol 215(3):403–410. 10.1016/S0022-2836(05)80360-2
PMID: **2231712**

Becker K, Stadler M. 2021. Recent progress in biodiversity research on the Xylariales and their secondary metabolism. J Antibiot 74(1):1–23. 10.1038/s41429-020-00376-0
PMID: **33097836**

Benocci T, Aguilar-Pontes MV, Zhou M, Seiboth B, de Vries RP. 2017. Regulators of plant biomass degradation in ascomycetous fungi. Biotechnol Biofuels 10:152. 10.1186/s13068-017-0841-x
PMID: **28616076**

Bills GF, Gloer JB. 2016. Biologically Active Secondary Metabolites from the Fungi. Microbiol Spectr 4(6). 10.1128/microbiolspec.FUNK-0009-2016
PMID: **27809954**

Brysch-Herzberg M, Seidel M. 2017. Distribution patterns of Saccharomyces species in cultural landscapes of Germany. FEMS Yeast Res 17(5):fox033. 10.1093/femsyr/fox033. PMID: **28520895**

Boynton PJ, Greig D. 2014. The ecology and evolution of non-domesticated Saccharomyces species. Yeast 31(12):449–462. 10.1002/yea.3040
PMID: **25242436**

Boynton PJ, Kowallik V, Landermann D, Stukenbrock EH. 2019. Quantifying the efficiency and biases of forest Saccharomyces sampling strategies. Yeast 36(11):657–668. 10.1002/yea.3435
PMID: **31348543**

Dashko S, Zhou N, Compagno C, Piškur J. 2014. Why, when, and how did yeast evolve alcoholic fermentation? FEMS Yeast Res 14(6):826–832. 10.1111/1567-1364.12161
PMID: **24824836**

Fukasawa Y, Osono T, Takeda H. 2011. Wood decomposing abilities of diverse lignicolous fungi on nondecayed and decayed beech wood. Mycologia 103(3):474–482. 10.3852/10-246
PMID: **21262989**

Glushakova AM, Ivannikova YV, Naumova ES, Chernov IY, Naumov GI. 2007. Massive isolation and identification of Saccharomyces paradoxus yeasts from plant phyllosphere. Mikrobiologiia 76(2):205–210. 10.1134/S0026261707020129
PMID: **17583221**

Goffeau A, Barrell BG, Bussey H, Davis RW, Dujon B, Feldmann H, …, Oliver SG. 1996. Life with 6000 genes. Science 274(5287):546–567. 10.1126/science.274.5287.546
PMID: **8849441**

Helaly SE, Thongbai B, Stadler M. 2018. Diversity of biologically active secondary metabolites from endophytic and saprotrophic fungi of the ascomycete order Xylariales. Nat Prod Rep 35(9):992–1014. 10.1039/c8np00010g
PMID: **29774351**

Keegstra K. 2010. Plant cell walls. Plant Physiol 154(2):483–486. 10.1104/pp.110.161240
PMID: **20921169**

Kjøller A, Struwe S. 1982. Microfungi in Ecosystems: Fungal Occurrence and Activity in Litter and Soil. Oikos 39(3):391–422. 10.2307/3544690
PMID: **n/a**

Kowallik V. 2015. The natural ecology of *Saccharomyces* yeasts [dissertation]. [Christian Albrechts University of Kiel (Germany)]:Christian Albrechts University of Kiel.
PMID: **n/a**

Kowallik V, Greig D. 2016. A systematic forest survey showing an association of Saccharomyces paradoxus with oak leaf litter. Environ Microbiol Rep 8(5):833–841. 10.1111/1758-2229.12446
PMID: **27481438**

Kowallik V, Miller E, Greig D. 2015. The interaction of Saccharomyces paradoxus with its natural competitors on oak bark. Mol Ecol 24(7):1596–1610. 10.1111/mec.13120
PMID: **25706044**

Krishna MP, Mohan M. 2017. Litter decomposition in forest ecosystems: a review. Energy Ecol Environ 2:236–249. 10.1007/s40974-017-0064-9
PMID: **n/a**

Lamacchia M, Dyrka W, Breton A, Saupe SJ, Paoletti M. 2016. Overlapping Podospora anserina Transcriptional Responses to Bacterial and Fungal Non Self Indicate a Multilayered Innate Immune Response. Front Microbiol 7:471. 10.3389/fmicb.2016.00471
PMID: **27148175**

Liti G, Warringer J, Blomberg A. 2017. Isolation and Laboratory Domestication of Natural Yeast Strains. Cold Spring Harb Protoc 2017(8):pdb.prot089052. 10.1101/pdb.prot089052
PMID: **28765292**

Mäkelä M, Hildén K, Kuuskeri J. 2021. Fungal Lignin-Modifying Peroxidases and H2O2-Producing Enzymes. In: Zaragoza Ó, Casadeveall A, Balestrini R, Mäkelä M, Nosanchuk

J, Viegas C, Vizzini A, de Vries RP, editors. Encyclopedia of Mycology Volume 2. Amsterdam: Elsevier. p. 247–259. 10.1016/B978-0-12-809633-8.21127-8
PMID: **n/a**

Nugent LK, Sihanonth P, Thienhirun S, Whalley AJS. 2005. Biscogniauxia: a genus of latent invaders. Mycologist 19(1):40–43. 10.1017/S0269-915X(05)00106-0
PMID: **n/a**

O'Donnell K. 1992. Ribosomal DNA internal transcribed spacers are highly divergent in the phytopathogenic ascomycete Fusarium sambucinum (Gibberella pulicaris). Curr Genet 22(3):213–220. 10.1007/BF00351728
PMID: **1525873**

Osono T, Takeda H. 2001. Effects of organic chemical quality and mineral nitrogen addition on lignin and holocellulose decomposition of beech leaf litter by Xylaria sp. Eur J Soil Biol 37(1):17–23. 10.1016/S1164-5563(01)01066-4
PMID: **n/a**

Osono T, To-Anun C, Hagiwara Y, Hirose D. 2011. Decomposition of wood, petiole and leaf litter by Xylaria species from northern Thailand. Fungal Ecol 4(3):210–218. 10.1016/j.funeco.2010.11.003
PMID: **n/a**

R Core Team 2021. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/
PMID: **n/a**

Robinson HA, Pinharanda A, Bensasson D. 2016. Summer temperature can predict the distribution of wild yeast populations. Ecol Evol 6(4):1236–1250. 10.1002/ece3.1919
PMID: **26941949**

Sampaio JP, Gonçalves P. 2008. Natural populations of Saccharomyces kudriavzevii in Portugal are associated with oak bark and are sympatric with S. cerevisiae and S. paradoxus. Appl Environ Microbiol 74(7):2144–2152. 10.1128/AEM.02396-07
PMID: **18281431**

Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, …, Sherry ST. 2022. Database resources of the national center for biotechnology information. Nucleic Acids Res 50(D1):D20–D26. 10.1093/nar/gkab1112
PMID: **34850941**

Serjeant K, Tang R, Anfang N, Beggs JR, Goddard MR. 2008. Yeasts associated with the New Zealand *Nothofagus* honeydew system. N Z J Ecol 32(2): 209-213.
PMID: **n/a**

Sniegowski PD, Dombrowski PG, Fingerman E. 2002. Saccharomyces cerevisiae and Saccharomyces paradoxus coexist in a natural woodland site in North America and display different levels of reproductive isolation from European conspecifics. FEMS Yeast Res 1(4):299–306. 10.1111/j.1567-1364.2002.tb00048.x
PMID: **12702333**

Stamps BW, Du L, Mitchell CA, Cichecwicz RH, Stevenson BS. 2015. Draft genomes of two sordariomycete fungi that produce novel secondary metabolites. Genome Announc 3(2):e00291-15. 10.1128/genomeA.00291-15
PMID: **25883292**

Strauss J, Mach RL, Zeilinger S, Hartler G, Stöffler G, Wolschek M, Kubicek CP. 1995. Cre1, the carbon catabolite repressor protein from Trichoderma reesei. FEBS Lett 376(1-2):103–107. 10.1016/0014-5793(95)01255-5
PMID: **8521952**

Sun S, Weng Y, Di X, Liu Z, Yang G. 2020. Screening of cellulose-degrading fungi in forest litter and fungal effects on litter decomposition. Bioresources 15(2):2937–2946. 10.15376/biores.15.2.2937-2946
PMID: **n/a**

Suto M, Tomita F. 2001. Induction and catabolite repression mechanisms of cellulase in fungi. J Biosci Bioeng 92(4):305–311. 10.1263/jbb.92.305
PMID: **16233102**

Van Rensburg P, Van Zyl WH, Pretorius IS. 1998. Engineering yeast for efficient cellulose degradation. Yeast 14(1):67–76. 10.1002/(SICI)1097-0061(19980115)14:1%3C67::AID-YEA200%3E3.0.CO;2-T
PMID: **9483796**

White TJ, Bruns T, Lee SB, Taylor JW. 1990. Amplification and Direct Sequencing of Fungal Ribosomal RNA Genes for Phylogenetics. In: Innis MA, Gelfand DH, Sninksy JJ, White TJ, editors. PCR Protocols: A Guide to Methods and Applications. New York (NY): Academic Press. p. 315-322. 10.1016/B978-0-12-372180-8.50042-1
PMID: **n/a**

Wickham H. 2016. ggplot2: Elegant graphics for data analysis. New York: Springer.
PMID: **n/a**

Extended Data
File name: Saccharomyces paradoxus conditioning media experiment.csv
Description: Data table (.csv format) of treatments and CFUs for Figure 1c-d.

Funding

Funding support came from Wheaton College Faculty-Student Summer Internship grants for PJB, MNR, MBG, and SEK and Wheaton College Foundation grants for MGL and SEK.

Author contributions

Samer El-Khatib: Data curation, formal analysis, funding acquisition, investigation, methodology, visualization, writing–original draft, writing–review & editing

Madeleine G. Lambert: Conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, visualization, writing–review & editing

Meghan N. Reed: Funding acquisition, investigation, methodology, writing–review & editing

Melane Bitro Goncalves: Funding acquisition, investigation, writing–review & editing

Primrose J. Boynton: Conceptualization, data curation, formal analysis, funding acquisition, methodology, project administration, supervision, validation, visualization, writing–original draft, writing–review & editing