1

Core Ideas:

1) North American Popcorn is composed of two distinct populations that differ genetically

and phenotypically.

2) North American Yellow Pearl Popcorns contain limited genetic diversity and are highly

inbred.

3) Pointed and Latin American Popcorns exhibit considerable genetic diversity and rapid

linkage disequilibrium decay.

4) Kernel color does not affect nicosulfuron sensitivity and instead reflects differences

between the populations.

5) Popcorn-specific candidate genes for nicosulfuron tolerance are distinct from dent corn.

Keywords:

Popcorn, Genetic Diversity, Nicosulfuron Tolerance, Plant Genetic Resources

Genetic diversity of North American popcorn germplasm and the effect of population structure on nicosulfuron response

Madsen Sullivan[1], Martin M. Williams II[2], Anthony J. Studer[1,†]

Affiliations:

[1] Department of Crop Sciences, University of Illinois, Urbana, IL 61801, USA.

[2] Global Change and Photosynthesis Research Unit, USDA-ARS, Urbana, IL 61801, USA.

[†] Corresponding author (astuder@illinois.edu).

Abbreviations:

ALS, acetolactate synthase; BCAA, branched-chain amino acid; CIMMYT, International Maize and Wheat Improvement Center; DAPC, discriminant analysis of principal coordinates; GBS, genotyping-by-sequencing; GRIN, Germplasm Resources Information Network; GWAS, genome-wide association studies; IBS, identity by state; LD, linkage disequilibrium; MAF, minor allele frequency; MAPQ, mapping quality; PCoA, principal coordinate analysis; SNP, single nucleotide polymorphism

ABSTRACT

Popcorn is an important crop in the United States; however, genetic analyses of popcorn are limited and tend to utilize relatively few markers that cannot capture the total genomic variation. To improve the genomic resources in popcorn, a panel of 362 popcorn accessions was evaluated using 417,218 single nucleotide polymorphisms generated using a genotyping-by-sequencing approach. Using this genomic data, a model-based clustering analysis identified two populations. The first comprised North American Yellow Pearl Popcorns and several accessions of the Chilean Curagua landrace. The second, the Pointed and Latin American Popcorns, included all remaining North American (pointed and early popcorns), Latin American, and global accessions. The two populations exhibited large differences in population structure and genetic diversity. The North American Yellow Pearl Popcorns constitute a highly inbred population with limited genetic diversity compared to the Pointed and Latin American Popcorns. Additionally, phenotypic differences between the two populations were observed in kernel color and nicosulfuron sensitivity. A filtered set of SNPs was curated and used for genome-wide association studies and popcorn-specific candidate genes for nicosulfuron tolerance were identified. The genomic characterization described here can be used by breeding programs to accelerate the rate of genetic gain and incorporate genetic diversity into elite popcorn germplasm.

1 INTRODUCTION

Popcorn (*Zea mays var. everta*) is a popular snack and a valuable commercial crop in the United

States. In a global market worth 10 billion USD, the United States has the highest per capita

popcorn consumption of any country at over 60 quarts per year, representing a total of 17 billion

quarts consumed annually (Serna-Saldivar 2022). Likewise, the United States grows more

popcorn than any other nation, with approximately 90,000 hectares of popcorn planted annually,

most of which are grown in Nebraska, Indiana, Ohio, and Illinois (USDA NASS, 2022).

Popcorn has been cultivated within the territory of the modern United States since pre-

Columbian times, although the range of pre-Columbian popcorn was limited to the American

Southwest. Popcorn landraces first reached this region by 4,200 BP (Merrill et al. 2009), likely

first from a highland origin, including the landraces Palomero Toluqueño and Palomero de

Jalisco, followed by a later mixing with lowland landraces, such as Chapalote and Reventador

(da Fonseca et al. 2015; Wang et al. 2017). Although these landraces do pop, the popping quality

is poor compared to modern lines in terms of expansion, hull fragmentation, and percentage of

unpopped kernels (Vazquez-Carrillo et al. 2019; Bautista-Ramirez et al. 2020).

Despite cultivation in the American Southwest for millennia, popcorn was not widely distributed

in North America until the late 18th and early 19th centuries, and even then, the amount grown

was limited (Erwin 1950). During the first half of the 19th century, popcorn was typically grown

in the United States as a small-scale garden crop for household use (Ziegler 2000). By the late

19th and early 20th century, the popularity of popcorn rose as open-pollinated varieties began

reaching more consumers, following the first successful processors and commercial brands in the

1880s and the introduction of the mobile popcorn machine in 1893 (Smith 1999). The

predominant types of popcorn grown during this time were the North American Pointed Rice

Popcorns or 'hulless' types, characterized as having a pointed upper kernel shape and frequently

colored white. The first hybrid popcorn used in commercial production was generated using two

selections of Japanese Hulless, a white pointed rice variety (Brunson 1937). However, since the

mid-20th century, another group of popcorn has largely replaced the pointed rice type

commercially – the North American Yellow Pearl Popcorns. This group has a rounded kernel

shape and is predominantly yellow, although color variation has been introduced by several

breeding programs (Ziegler 2000). While the North American Yellow Pearl Popcorns are the

most important for commercial production in the United States, North American Pointed Rice

Popcorns are still commercially grown and used in breeding programs.

Characterizing the genetic relationships and identifying variation within these distinct

populations of North American popcorn is an important aspect of breeding improved popcorn.

Early molecular techniques were used to assess the genetic diversity of North American Yellow

Pearl Popcorns and identified three heterotic groups – Supergold, South American, and Amber

Pearl (Kantety et al. 1995). Later work by Santacruz-Varela et al. defined three major groups of

North American popcorn – the aforementioned North American Pointed Rice Popcorn and North

American Yellow Pearl Popcorn, as well as the North American Early Popcorn (2004).

Additionally, each group's phylogenetic relationship and historical origin were investigated, and

possible germplasm sources were identified. The North American Pointed Rice Popcorns

appeared to originate from the traditional pointed rice popcorns of Latin America, based on both

morphological and genotypic similarities. Furthermore, the North American Yellow Pearl

Popcorns were most closely related to, and likely principally derived from, the Chilean Curagua

landrace. Finally, the North American Early Popcorns may have been developed from Northern

Flint corn and subsequently influenced by European popcorn varieties (Santacruz-Varela et al.

2004). Although this information has been useful in understanding breeding history and

relatedness, the limited number of markers and lines has restricted its use in modern breeding

programs to heterotic pool development.

Compared with dent corn germplasm, popcorn exhibits considerably inferior agronomic traits,

including herbicide tolerance, disease resistance, lodging, yield, and overall vigor (Ziegler 2000).

Additionally, differences in herbicide tolerance have been reported within popcorn, with white

popcorn generally exhibiting greater herbicide sensitivity than yellow (Loux et al. 2017).

Historically, the poorer agronomic traits of popcorn are due partly to the amount of time and

resources breeders have invested relative to dent corn and also because of the emphasis placed

on popping quality traits. In addition to agronomic traits, popcorn breeders must consider quality

traits directly related to popping and eating quality, including popping expansion, hull

separation, frequency of unpopped kernels, texture, and appearance (Matz 1984). Furthermore,

dent corn breeding pipelines have integrated molecular characterization and resources, resulting

in improved selection and rates of genetic gain, while publicly available resources within

popcorn are limited.

High-quality genetic and genomic resources in maize have empowered researchers and

facilitated discoveries involving the genetic architecture of traits, candidate gene association and

identification, and population structure (Buckler et al. 2009; Cook et al. 2012; Romay et al.

2013; Peiffer et al. 2014; Wallace et al. 2014; Gage et al. 2020). While the genetic resources for

mapping populations, diversity panels, and seed banks have been used extensively within dent

corn, popcorn germplasm comprises a genetically distinct subpopulation of maize that is

underrepresented in these studies. Recently, genotyping-by-sequencing (GBS) was performed on

popcorn accessions from CIMMYT (Li et al. 2019; Li et al. 2021) and Chinese breeding

programs (Yu et al. 2021) and used to characterize genetic diversity, describe population

structure, and perform genome-wide association studies (GWAS). The genomic characterization

of these diverse popcorn accessions can be used to assist breeders in incorporating genetic

variation into breeding programs and identifying loci controlling important traits in popcorn.

Likewise, the rate of genetic gain in North American popcorn breeding programs would increase

dramatically with the incorporation of high-quality genetic and genomic information, as has been

utilized in dent corn breeding for decades.

In this study, molecular analysis was performed on 362 publicly available popcorn accessions

genotyped with GBS representing the diversity of North American popcorn. The goals of this

study were to i) characterize the genetic diversity of North American popcorn, ii) identify

population structure within North American popcorn and other groups of popcorn, iii) describe

genomic differences between these populations, iv) perform genome-wide association studies,

and v) compare findings with previous attempts to characterize North American popcorn.

2 MATERIALS AND METHODS

2.1 Plant materials and DNA extraction

Of publicly available lines of popcorn, 341 unique accessions were selected from the Germplasm

Resources Information Network (GRIN), and 21 accessions were selected from the International

Maize and Wheat Improvement Center (CIMMYT). Of these, 79 accessions had available

sequence information (Romay et al. 2013). Thus, 283 new popcorn lines were sequenced, with

68 selected for replicate sequencing. Additionally, four dent inbreds with available sequence

information were included as outgroup accessions: B73, Mo17, PH207, and Oh43.

Samples were prepared for DNA extraction by bulking tissue from five seedlings of each

accession in a 2 mL tube, collected from the youngest leaf of each approximately ten days after

germination. DNA was extracted from each sample using a modified CTAB DNA extraction

protocol and resuspended in water. DNA was quantified in 96-well plates using PicoGreen

(Invitrogen, Carlsbad, CA), and DNA concentrations were normalized to 50ng/μL by diluting

with 10 mM Tris, then stored at 4°C.

2.2 Genotyping-by-sequencing library construction

Genotyping-by-sequencing libraries were constructed using the protocol described by Elshire *et

al.* (2011). Briefly, restriction digestion and ligation were performed in 96-well plates using

*Ape*KI by adding 250 ng of DNA from each sample, 1.5μL of 0.1 μM rare barcoded adapter, and

0.5μL of 10 μM common barcoded adapter per well. Libraries were pooled, size-selected and

cleaned with Agencourt AMPure XP beads (Beckman Coulter Inc., Brea, CA), and amplified for

15 cycles using KAPA HiFi HotStart ReadyMix (KAPA BIOSYSTEMS). Following a final

purification with Agencourt AMPure XP beads, libraries were quantified using PicoGreen. A 1

ng/μL dilution of the library was made using 10 mM Tris, and 5 μL was run on a 1% agarose gel

153   to verify size selection and primer-dimer removal. Fragment size was estimated using a

154   Bioanalyzer 2100 (Agilent, Santa Clara, CA) High Sensitivity DNA Analysis. Pooled libraries

155   were diluted to 10 nmol using 10 mM Tris and sequenced with 100-bp, single-end reads on an

156   Illumina NovaSeq 6000 (Illumina, San Diego, CA). The sequencing data has been submitted to

157   NCBI SRA (PRJNA911304).

158   **2.3 Marker calling**

The reads from the 79 popcorn accessions previously sequenced by Romay et al. (2013) were

downloaded from NCBI SRA (SRP021921) and processed with the new raw sequencing reads

using the GBS v2 discovery pipeline from TASSEL 5 (release 5.2.59) (Glaubitz et al. 2014). The

pipeline tools and select parameters were run in the following order: First,

GBSSeqToTagDBPlugin, with ApeKI and a kmer length of 64, followed by

TagExportToFastqPlugin. Next, two alignment indices were built with Bowtie2 (Langmead &

Salzberg 2012) using the B73 v5 and HP301 v1 reference genomes (Hufford et al. 2021). Both

genome indices and output files were processed using the same parameters afterward. The

Bowtie2 alignment command was run using the --very-sensitive preset, resulting in an overall

alignment of 90.6% and 91.2% for the B73 v5 and HP301 v1 indices, respectively. The SAM

files were then sorted using samtools (Li et al. 2009), and mapping quality (MAPQ) scores were

analyzed using Qualimap 2 (Okonechnikov et al. 2016). The SAM files were read using the

SAMToGBSdbPlugin with a minimum MAPQ score of 20, after which the

DiscoverySNPCallPluginV2 was run using a minimum minor allele frequency of 0.05. Finally,

the ProductionSNPCallerPluginV2 was run using the same parameters as the

GBSSeqToTagDBPlugin.

175   **2.4 Marker coverage and characterization**

The genomic distribution of single nucleotide polymorphisms (SNPs) was based on the HP301

v1 reference genome, as the popcorn reads alignments were slightly better than when using the

B73 v5 genome. Markers were reported from TASSEL 5 for the Core Analysis set and each

chromosome. An additional marker set was generated for imputation and GWAS after filtering

out sites with more than 0.75 missing data from the Core Analysis set. Nucleotide substitutions

were calculated from the Core Analysis set using the TASSEL 5 GenotypeSummaryPlugin.

Markers were identified as intragenic by searching their positions relative to the start and stop

positions of the HP301 v1 gene models. The density and distribution of polymorphic SNPs were

determined in *R* (R Core Team, 2021) using the *R* package *rMVP* (Yin et al. 2021).

2.5 Population structure

A kinship matrix of centered identity by state (IBS) values was calculated for all possible

pairwise taxa comparisons in TASSEL. A plot of each pairwise comparison was generated using

the *R* package *ggplot2* (Wickham 2016). To formally test for population structure using

molecular data, an unsupervised, model-based clustering analysis was performed using

ADMIXTURE (Alexander et al. 2009). One through five *K* populations were tested using tenfold

cross-validation, with errors minimized at *K* equals two. A second ADMIXTURE analysis was

performed in validation after combining the 362 popcorn accessions with 2,284 additional

accessions sequenced from the Ames Diversity Panel (Romay et al. 2013). Due to computational

restraints, accessions designated stiff stalk or non-stiff stalk were excluded. Reads from the

combined set of accessions were aligned to the HP301 v1 reference genome using the TASSEL 5

GBS v2 discovery pipeline without changing any of the aforementioned parameters. In

ADMIXTURE, one through twenty *K* populations were tested using fivefold cross-validation,

with errors minimized at *K* equals fifteen.

Principal coordinates were generated in TASSEL from the IBS kinship matrix. Three of the

coordinates from the matrix with high eigenvalues were visualized using *ggplot2*. After

removing 30 accessions with missing data greater than 0.95, phylogenetic analysis was

performed using the *R* package *ape* (Paradis et al. 2004) using the neighbor-joining method from

a modified Euclidean distance matrix and visualized with the *R* package *ggtree* (Yu et al. 2017).

The neighbor-joining algorithm was selected because of the considerable differences in selection

and breeding history of the populations included in this study, making alternate methods such as

UPGMA untenable. To further investigate the relationship between teosinte and the Pointed and

Latin American Popcorn population, an additional neighbor-joining tree was generated. A

distance matrix was produced from the seventeen teosinte accessions and ten randomly selected

individuals from each of the fifteen ADMIXTURE groups identified from the combined popcorn

– Ames Diversity Panel analysis. This distance matrix was used by *ape* to create the second tree

and also visualized with *ggtree*.

2.6 Population genetics and diversity

To evaluate differences in selection and genetic drift between the two populations, the proportion

of monomorphic SNPs within each population was calculated. Both populations were filtered

from the Core Analysis set, after which markers in each population were filtered to contain a

minimum of two observations. Markers without minor alleles were then filtered out, allowing the

number of monomorphic markers to be calculated by subtracting the remaining polymorphic

markers. After filtering monomorphic sites, marker minor allele frequencies were calculated

from the Core Analysis set using the TASSEL 5 GenotypeSummaryPlugin. The distribution of

minor allele frequencies was calculated for each popcorn population.

A marker subset was produced for linkage disequilibrium (LD) analysis by removing SNPs with

more than 50% missing data, a MAF less than 0.05, and heterozygosity greater than 0, resulting

in 13,031 retained markers. Within each population, LD was calculated between the subset of

markers on each chromosome using the TASSEL 5 LinkageDisequilibriumPlugin. The level of

LD decay was measured using the squared Pearson correlation coefficient (r^2) between each pair

of markers. Pairwise comparisons were binned according to distance using logarithmically

increasing bins.

2.7 Herbicide treatment

A subset of popcorn accessions was evaluated for response to nicosulfuron at the University of

Illinois Crop Sciences Research and Education Center, Urbana, Illinois, during the summer of

2022. A total of 294 genotypes were included in the trial, with 111 North American Yellow Pearl

Popcorns and 178 Pointed and Latin American Popcorns. Additionally, five controls were

included in the panel – a tolerant sweet corn hybrid (GSS1477), a nicosulfuron-sensitive sweet

corn hybrid (Merit), a partially sensitive yellow popcorn hybrid, a tolerant yellow popcorn

hybrid, and a tolerant white popcorn hybrid. Genotypes were arranged in a randomized complete

block design with three replications. An experimental unit comprised a single 3-meter row

containing 10-20 plants.

When the majority of plants reached the V3/V4 growth stage, nicosulfuron was applied to one-

half of each row at a rate of 35 g ai ha^{-1} with a 0.25% (v/v) nonionic surfactant and 2.5% (v/v)

urea ammonium nitrate. This treatment was delivered using a pressurized CO_2 backpack sprayer

equipped with AI110025 nozzles (TeeJet Technologies, P.O. Box 7900, Wheaton, IL 60187)

spaced 51 cm apart on a 1.5-meter boom calibrated to apply 187 L ha^{-1} at 276 kPa. Popcorn

plants were evaluated ten days after nicosulfuron application. Injury was assessed visually on a

scale of 0-4, with 0 representing no injury and 4 representing plant death.

2.8 Imputation and genome-wide association studies (GWAS)

To evaluate the imputation accuracy of different methods, 1% of the GWAS Filtered set was

masked and imputed. Imputation was performed with FILLIN (Swarts et al. 2014) and Beagle

v5.1 (Browning et al. 2018). FILLIN was run using a haplotype block size of 4,096. Beagle was

run using an effective population size of 100,000 and an HP301 genetic map. The genetic map

for HP301 was constructed in *R/qtl* (Broman et al. 2003) using GBS data from the HP301 RILs

(Elshire et al. 2011), aligned to the HP301 v1 reference genome using the TASSEL 5 GBS v2

discovery pipeline. To compare the two imputation methods, the unmasked, masked, and

imputed marker sets were used as input for TASSEL's ImputationAccuracyPlugin, and error

rates were generated.

As the popcorn populations differ in kernel color and response to nicosulfuron, GWAS was

performed on both these phenotypes. Both GWAS analyses were performed using a weighted

MLM in TASSEL with the FILLIN-imputed marker set, a kinship matrix of normalized IBS

values, and a weight matrix of principal coordinates with high eigenvalues. A Bonferroni-

corrected p-value of 0.01 resulted in a -log(p) significance threshold of 7.3. Results of both

GWAS analyses were visualized using the *R* package *qqman* (Turner 2018).

261   **3 RESULTS**

262   **3.1 Germplasm**

To improve the genomic resources and investigate the genetic diversity of popcorn, GBS was

performed on 362 popcorn accessions. In GRIN, a total of 640 available accessions were listed

as popcorn. Accessions that were also labeled as other kernel types were removed, reducing the

count to 379. Then all accessions that appeared to be duplicates based on passport information

were removed, bringing the number of unique popcorn accessions down to 304. Of the

accessions with multiple kernel types listed, 37 were added back, as they were known popcorn

lines. Finally, an additional 21 popcorn landraces from CIMMYT were included, bringing the

total number of accessions to 362 (Table S1). Of these, 72 are listed as landraces, 109 as

cultivated material, and the remaining 181 as breeding material. In addition to the 362 popcorn

accessions, four previously sequenced dent inbreds were included to serve as outgroup lines:

B73, Mo17, PH207, and Oh43.

274   **3.2 Marker density and distribution**

From the GBS data, a Core Analysis set of 417,118 markers with an average minor allele

frequency of 0.169 was generated after passing SNP filtering and quality control. A Filtered

GWAS set of 200,101 markers with an average minor allele frequency of 0.216 was generated

by removing all SNPs with greater than 0.75 missing data in preparation for imputation. Of the

nucleotide substitutions from biallelic SNPs, adenine-guanine and thymine-cytosine transitions

were the most common (29.5% each), while adenine-thymine transversions were the least

common (7.7%). Of the Core Analysis set of markers, 327,876 (78.6%) were identified as

intragenic using the HP301 v1 gene models (Hufford et al. 2021), consistent with the observation

that SNP density increased from the centromeric to the telomeric regions (Supplementary Fig.

S1). Across the 10 maize chromosomes, the Core Analysis set of markers had an average marker

density of 197.3 SNPs per Mb, with the highest and lowest marker densities observed on

chromosomes 5 (216 SNPs per Mb) and 6 (184 SNPs per Mb), respectively (Table 1). The

Filtered GWAS set had an average marker density of 94.6 SNPs per MB, with the highest and

lowest densities on chromosomes 1 (108 SNPs per Mb) and 4 (74.7 SNPs per Mb), respectively.

Chromosome	Chromosome Length (b)	Core Analysis Set			Filtered GWAS Set		
		Number of SNPs	SNPs per Mb	MAF	Number of SNPs	SNPs per Mb	MAF
1	305,741,490	64450	210.8	0.16807	32903	107.6	0.21527
2	245,520,863	51470	209.6	0.17452	24514	99.8	0.22231
3	240,329,036	47791	198.9	0.17328	24554	102.2	0.22337
4	251,015,817	41118	163.8	0.16409	18748	74.7	0.20173
5	221,904,911	47873	215.7	0.1712	23121	104.2	0.22609
6	177,672,499	32691	184.0	0.16488	14358	80.8	0.21071
7	180,265,189	35984	199.6	0.16534	15815	87.7	0.21834
8	177,329,642	34910	196.9	0.16316	16579	93.5	0.2199
9	164,337,916	31699	192.9	0.1707	16053	97.7	0.21045
10	150,435,773	29232	194.3	0.16794	13456	89.4	0.21866
Total	2,114,553,136	417218	197.3	0.16872	200101	94.6	0.21576

290 Table 1 SNP characteristics summary for the Core Analysis and Filtered GWAS SNP sets

3.3 Population structure

To identify relationships among accessions included in the panel, the distribution of IBS pairwise

comparisons for the 362 accessions was calculated using the Core Analysis set (Fig. 1). A

bimodal distribution was identified, with a large main peak centered on 0.72 and an additional

smaller peak centered on 0.82. These results indicate that although most accessions have a weak

relationship with other material in this panel, a subset appears to be more closely related.

The North American popcorn populations are comprised of accessions with different breeding

histories, geographic distributions, and selection pressures. An ADMIXTURE analysis was

performed on all taxa to formally test for population structure using one to five K populations.

Figure 1 Identity by state (IBS) distribution across 362 popcorn accessions

Cross-validation error was minimized at $K=2$ (Supplementary Fig. S2), and taxa were clustered

into two groups of similar size, with 147 individuals assigned to the first group and 215 assigned

to the second (Fig. 2, Table S1). Most taxa assigned to the first group are North American

Yellow Pearl Popcorns from breeding programs in the United States with known pedigrees and

generally fall into one of the previously described heterotic pools. However, several notable taxa

from Chile were assigned to this group, including CHZM 05 006, CHZM 06 004, CHZM 06 012,

and CHZM 07 097, all of which are members of the Curagua race. The remaining 23 members of

Curagua were assigned to the other group of popcorn, which will be referred to as the Pointed

and Latin American Popcorns. This group contains many diverse accessions, including all North

American Pointed Rice Popcorns (n=21), North American Early Popcorns (n=32), all Latin

American pointed and pearl popcorns accessions outside the four members of Curagua (n=98),

and nearly all lines developed outside of the Americas (n=64).

Figure 2 Bar plot of assignment values from ADMIXTURE analysis. Each bar corresponds to

one of 362 popcorn accessions and is ordered by the likelihood of assignment. The complete list

319   of assignments can be found in Table S1.

Since more than two popcorn groups have been reported previously in the literature, a second

analysis was performed to ensure that the identification of additional popcorn groups was not

limited by an insufficient number of related accessions. A combined dataset was generated after

merging the 362 popcorn accessions with 2,284 accessions from the Ames Diversity Panel that

were not categorized as dent (stiff stalk or non-stiff stalk). After minimizing the ADMIXTURE

cross-validation error at K=15, the combined 2,646 accessions were assigned to 15 groups, with

no additional popcorn groups identified. Although dent lines were excluded, 885 unclassified

accessions were assigned to known public subgroups, including B73, B14, B52, Hy, Mo17,

W22, Oh43, and WF9, demonstrating that dent material was still well-represented in the

analysis. Notably, the analysis assigned the 17 teosinte inbred lines included in the Ames

Diversity Panel to the Pointed and Latin American Popcorn group. Thus, the teosinte inbred lines

are most genetically similar to the Pointed and Latin American Popcorns, followed by North

American Yellow Pearl Popcorns, then sweet corn, and finally dent and flint tropical and

temperate material (Supplementary Fig. S3). Other notable taxa assigned to the Pointed and

Latin American Popcorn group include accessions of Palomero Toluqueño, Reventador,

Chapalote, Pororo, Avati Pichinga, and Pisankalla.

Figure 3 Principal coordinate analysis (PCoA) of 362 popcorn accessions estimated with the

340   Core Analysis set and associated scree plot.

Principal coordinate analysis (PCoA) was also performed to investigate population structure and

identify individual taxa grouping, and three of the coordinates explained 83.3% of the variation

(Fig. 3). The first coordinate separates North American Yellow Pearl Popcorns from the Pointed

and Latin American Popcorns, supporting the $K=2$ optimum number of clusters of the popcorn

ADMIXTURE analysis. The second coordinate separates the Pointed and Latin American

Popcorns into three general divisions. Most of the North American Pointed Rice Popcorns cluster

together near the bottom, with several accessions of Pisankalla. Next, most of the Latin Pointed

Rice Popcorns group in the center with members of Curagua and Reventador. Finally, near the

top, lowland tropical popcorn landraces such as Avati Pichinga, Pipoca, and Pororo are located

together. The third coordinate separates the North American Yellow Pearls, with accessions from

the South American heterotic group clustered near the bottom, those from Supergold clustered at

the top, and lines developed by Iowa State's Popcorn Breeding Program located in the center.

The neighbor-joining tree based on the distance matrix supported the population structure

findings of the ADMIXTURE analysis and PCoA (Fig. 4). The major division of clades within

the tree corresponded to the separation between North American Yellow Pearl Popcorns and

Pointed and Latin American Popcorns. Several Chilean Curagua accessions fall within and basal

to the North American Yellow Pearl clade, while the remainder are located within the Pointed

and Latin American Popcorns. Two additional Latin American Taxa assigned to the North

American Yellow Pearl Popcorn population, BRAZ 2832 and SONO 147, occur within the

Supergold clade, indicating that they likely represent a recent movement of North American

Yellow Pearl Popcorn germplasm (Fig. 4).

Figure 4 Neighbor-joining tree of 332 popcorn accessions and 4 dent accessions based on a

modified Euclidean distance matrix calculated from the Core Analysis set. The three heterotic

pools of North American Yellow Pearl Popcorns are labeled, along with the dent outgroup.

Accessions SONO 147 and BRAZ 2832 are marked with an asterisk (∗). Accessions belonging

to Curagua are marked with a dot (•).

3.4 Population genetics and allele diversity

To evaluate differences in allele frequencies and genetic diversity within popcorn germplasm,

each population was evaluated for the proportion of monomorphic SNPs, minor allele

frequencies (MAF), and LD decay. After filtering markers in each group to account for structural

variation, monomorphism was evaluated. The proportion of monomorphic SNPs was

considerably different between the two groups, with 5.5 times more monomorphic SNPs in the

North American Yellow Pearl Popcorns (0.328) than in the Pointed and Latin American

Popcorns (0.059). Additionally, there were differences in MAF between the two populations

(Fig. 5).

Figure 5 Minor allele frequency (MAF) distributions of the two popcorn populations. Y-axis

shows the proportion of SNPs, as differing counts of monomorphic SNPs were removed from

each population prior to calculating frequencies.

The relative frequency of SNPs with a MAF of 0.05 or less in North American Yellow Pearl

Popcorns and Pointed and Latin American Popcorns were 28.4% and 7.7%, respectively. For all

MAF greater than 0.05, the Pointed and Latin American Popcorns contained more SNPs than the

North American Yellow Pearl Popcorns. Overall, the North American Yellow Pearl Popcorns

contained more SNPs with low MAF than the Pointed and Latin American Popcorns.

Linkage disequilibrium decay also differed between the two populations (Fig. 6). Decay was low

in the North American Yellow Pearl Popcorns, reaching a median r^2 of 0.3 within 10 – 99 Kb. In

contrast, LD decayed rapidly in the Pointed and Latin American Popcorns, reaching a median r^2

of 0.1 within 10 – 99 bp. When compared with previous evaluations of LD decay in temperate

and tropical populations (Romay et al. 2013, White et al. 2020), North American Yellow Pearl

Popcorns exhibit slower LD decay than even the B73 heterotic subgroup (r^2 of ~0.25 within 10 –

99 Kb), while LD decay in the Pointed and Latin American Popcorns was slightly faster than in

CIMMYT tropical germplasm (r^2 of 0.1 within 100 – 999 bp).

Figure 6 Median linkage disequilibrium (LD) decay of the two popcorn populations. Pairwise

comparisons were binned by distance using logarithmically increasing bins.

3.5 Genome-wide association studies

A significant limitation of GWAS is the *p*-value correction required to account for multiple tests performed (Tam et al. 2019). Reducing the number of markers tested is one way to ease the stringency of the commonly used Bonferroni correction method. By removing all markers with greater than 0.75 missing data, the resulting Filtered GWAS set reduced the number of markers to approximately half of the Core Analysis set. Furthermore, since the imputation strategies tested here rely on generating haplotypes from existing marker data, any reduction in missing sites improves haplotype generation and subsequent imputation. Two commonly used imputation strategies in maize are Beagle and FILLIN. Both were used to impute the Filtered GWAS set, and error metrics were calculated for each. FILLIN and Beagle had overall error rates of 0.02321 and 0.08173, respectively. Because of the difference in error rates, only the FILLIN-imputed marker set was used to perform GWAS.

The North American Yellow Pearl Popcorns and Pointed and Latin American Popcorns represent two distinct populations with significant differences in population structure, genetic diversity, and breeding history. Additionally, phenotypic differences are known between these two populations, such as kernel color. North American Yellow Pearl Popcorns are typically yellow, with only four white accessions in the panel (3.3%), while the Pointed and Latin American Popcorns have a much higher frequency of white kernels or endosperm, with 111 accessions in the panel (54.7%). Since kernel color is a simple trait that differs among the germplasm assessed here, it was selected as a proof of concept to assess the utility of using this panel for GWAS (Fig. 7). Several significant SNPs were identified, some of which have been associated with kernel color previously (Table S2). The SNP with the strongest association with kernel color is located in *Y1*, which is known to affect the concentration of carotenoid pigments in the endosperm. The

SNP with the next strongest association with kernel color is located 200 bases away from

Zm00027ab258670, a gene model predicted to have carotene desaturase activity. Three other

peak SNPs were identified, none of which have been previously associated with kernel color.

These results indicate that despite the known differences in population structure and phenotype

frequencies, the panel can identify associations between loci and traits.

Figure 7 Manhattan plot of genome-wide association study (GWAS) of kernel color from the

Filtered GWAS set. The red line represents the threshold after correcting for multiple testing

430   using a Bonferroni correction.

A trait anecdotally associated with kernel color in popcorn is herbicide injury. Herbicide labels

often exclude white popcorn from approved use, even when yellow popcorn is included (Loux et

al. 2017). While the white color itself should have no bearing on injury, herbicide sensitivity

differences between the two popcorn populations may have been reduced to kernel color for

simplicity. In the herbicide trial, 4 of the 111 North American Yellow Pearl Popcorns (3.6%) and

32 of 178 Pointed and Latin American Popcorns (18%) showed typical nicosulfuron injury

(Table S3). Of the 32 injured Pointed and Latin American Popcorns, several are important

historical lines grown in the American Midwest, including Japanese Hulless, Australian Hulless,

H5505, IA DS 38-W, and IA DS 43-W. Notably, these pointed lines are all white and represent

much of the variation of the pointed rice popcorns traditionally grown in North America.

The Filtered GWAS set was used to map nicosulfuron injury. A total of nine SNPs representing

seven peaks were found to be significant (Fig. 8, Table S4). The SNP with the strongest

association with injury is located in *cl11433_2b*, a transcription factor associated with the

thylakoid. The next strongest association was with a subunit of NADH dehydrogenase. Other

SNPs with statistically significant associations include a predicted protein kinase, the vacuole-

associated phospholipase *pco139881*, a locus 1.3 kb upstream from a predicted aminoacyl-tRNA

synthetase, the cytochrome p450 *CYP72A27*, and a predicted monooxygenase

glycosyltransferase. Although these genes have not previously been implicated in nicosulfuron

injury, many of the classes to which they belong are associated with herbicide tolerance.

Figure 8 Manhattan plot of genome-wide association study (GWAS) of nicosulfuron injury from

the Filtered GWAS set. The red line represents the threshold after correcting for multiple testing

using a Bonferroni correction.

4 DISCUSSION

4.1 Population Structure

The results of IBS, ADMIXTURE, PCoA, and the neighbor-joining tree support two populations

of popcorn in the panel investigated here: the North American Yellow Pearl Popcorns and the

Pointed and Latin American Popcorns. Both the PCoA and neighbor-joining tree indicate a

degree of sub-grouping within each group, but these subgroups were not significantly different

based on the genomic data. Within the North American Yellow Pearl Popcorns, differences in

population structure of the Supergold and South American heterotic groups were apparent, and

there was a minor grouping effect on lines originating from Iowa State's Popcorn Breeding

Program. Divisions within the Pointed and Latin American Popcorns resulted in three general

subgroups – North American Pointed Rice Popcorns, Latin Pointed Rice and highland popcorn

landraces, and lowland tropical popcorn landraces. It is important to note that not all individuals

typically assigned to one of these subgroups clustered with their subgroup, though the majority

do. Although sub-groups were identified in both populations, they are not comparable in

magnitude to the division between North American Yellow Pearl Popcorns and Pointed and

Latin American Popcorns. Therefore, two groups are proposed here that explain most of the

variation in population structure in this popcorn diversity panel.

Foundational work on the phylogenetic relationships of North American popcorn identified more

than two groups of popcorn. After evaluating 56 popcorn populations for morphological traits,

isozymes, and SSRs, six major divisions were previously identified (Santacruz-Varela et al.

2004). Two groups of pointed rice popcorn of North American and Latin American origin were

proposed, followed by the related North American Early Popcorns. A single population of North

American Yellow Pearl Popcorns was identified, with minimal divisions between the heterotic

groups. Finally, two additional groups of Latin America were identified, one corresponding to

lowland races of Argentina, Brazil, and Paraguay, and the other composed of various accessions

from Chile, Uruguay, and Mexico.

Besides the number of lines, a major difference between previous studies and the results

presented here is that morphological traits were not included when determining phylogenetic

relationships and population structure. All traits are composed of genotypic and environmental

effects, and many of the vegetative and phenological traits included in previous analyses of

popcorn are known to be affected by photoperiod sensitivity, including plant height, ear height,

leaf number, and days to flowering (Fei et al. 2022). As few as four genomic regions have been

identified that control photoperiod responses in these traits (Coles et al. 2010). Therefore,

including these traits in measures of phylogenetic divergence likely does not reflect genomic

differences between populations and could be heavily influenced by photoperiod sensitivity. This

is supported by the conclusion that the six previously identified popcorn populations were

primarily assigned based on results of the morphological characterization (Santacruz-Varela et

al. 2004). The inclusion of morphological traits that do not reflect genome-wide differences

explains the discrepancy between the six populations of popcorn identified previously and the

two populations presented here.

Investigation into the genetic diversity and population structure of CIMMYT accessions and

Chinese germplasm has identified distinct popcorn populations in their panels. A panel

composed of teosinte, popcorn landraces, non-popcorn landraces, North American Yellow Pearl

Popcorns, and elite CIMMYT inbreds was used to identify loci associated with temperate and

tropical adaptation (Li et al. 2019). This panel identified two groups of popcorn, one

corresponding to the North American Yellow Pearl Popcorns and the other associated with

landraces from Latin America. The genetic diversity and population structure within a popcorn

collection of Chinese accessions, CIMMYT landraces, and inbreds from the United States were

recently characterized (Yu et al. 2021). In this study, popcorn accessions were assigned to one of

three groups – Latin American landraces, North American Yellow Pearl Popcorns, and recently

developed inbreds from breeding programs, likely of North American Yellow Pearl origin. Based

on these studies and the findings presented here, North American Yellow Pearl Popcorns have

consistently been shown to constitute a genetically distinct pool of germplasm compared to Latin

American landraces.

In contrast with the uniform separation of North American Yellow Pearl Popcorns from other

types of maize, the assignment of the Pointed and Latin American Popcorns is not as consistent.

Single-group assignment of popcorn landraces of Latin America has been previously reported

when compared with Chinese germplasm as discussed above (Yu et al. 2021), Brazilian popcorn

inbreds (Vittorazzi et al. 2018), and highland and lowland Argentine landraces (Bracco et al.

2012). However, an investigation of Latin American and Caribbean landraces identified sixteen

populations, three of which included popcorn (Bedoya et al. 2017). The first of the three

populations contained Palomero Toluqueño, Chapalote, and Reventador, all of which are

considered primitive popcorns of Northern Mexico. The second group was related to the first and

contained Canguil, Confite Puneño, and Pisankalla, all of which are pointed. The third group was

unrelated to the first two and contained lowland Avati popcorn. More recently, an analysis of

575 popcorn landraces from Latin America identified nine genetic clusters using discriminant

analysis of principal components (DAPC) without considering a priori groups (De Almeida Silva

et al. 2020). However, it is important to consider differences in methods when comparing these

findings.

Most popcorn groups have been identified using model-based clustering approaches such as

STRUCTURE (Pritchard et al. 2000) or ADMIXTURE. Furthermore, model-based clustering

methods assign individuals to populations relative to the entire panel – populations identified by

an initial clustering analysis can be further divided into sub-groups by subsequent clustering

analyses, as with the American and Caribbean landraces discussed above. Additionally, methods

based on PCA, like DAPC, tend to provide higher estimates of the total number of populations

than model-based clustering approaches (Intarapanich et al. 2009, Linck & Battey 2019) and are

especially sensitive to the influence of a priori vs. de novo groupings (Miller et al. 2020).

Because of these considerations, a single assignment by the model-based clustering method

ADMIXTURE on the Core Analysis set was used to assess population structure for the entire

panel. The findings reported here largely agree with similar approaches to estimate population

structure in popcorn, and differences with other reports largely stem from assignment methods.

536   **4.2 Genetic diversity**

The Pointed and Latin American Popcorns contain a variety of accessions, including pointed

popcorns from the American Midwest. It is unlikely that the genetic similarities between lines of

pointed popcorn from North America and Latin America indicate a recent introduction or

transfer of germplasm between continents, as the lines constituting this group were

geographically widely distributed. Instead, these similarities may indicate a pre-Columbian

interchange of popcorn between North and South America, as has been previously suggested

(Bedoya et al. 2017). Additionally, the genetic similarity may reflect reduced admixture with

other landraces due to the high frequency of the dent-sterile $Ga1$-S allele in North and South

American popcorn (Goodman et al. 2021). This aligns with the finding that popcorn landraces

tend to group separately from other maize types, as discussed above.

North American Yellow Pearl Popcorns share genetic similarities with several Curagua

accessions from Chile. Genetic similarity between United States varieties and South American

material has been noted before (Vigouroux et al. 2008, van Heerwaarden et al. 2010). However,

this has generally been attributed to the post-Columbian movement of germplasm from the

United States to South America. Many Chilean, Argentine, and Brazilian races are of United

States origin, including Dulce Golden Bantam, Hickory King, and Dente Branco. The presence

of Curagua accessions in the North American Yellow Pearls is evidence that intercontinental

movement from South to North also occurred.

The North American Yellow Pearl Popcorns have a much higher LD and proportion of

monomorphic markers than Pointed and Latin American Popcorns and other populations of

maize (Romay 2013). North American Yellow Pearl Popcorns also exhibit higher IBS values

when compared with other North American Yellow Pearl Popcorns and lower minor allele

frequencies, indicating a high rate of inbreeding and lower genetic variation. Considering the few

Curagua accessions clustered among the North American Yellow Pearls (Fig. 4), this reduced

genome variation is consistent with a small founder population and fewer meiotic events relative

to the considerably older Pointed and Latin American Popcorns. The genetic variation within the

three accessions of Curagua assigned to the North American Yellow Pearls represents only a

portion of the total variation present in all members of the Curagua race. If members of Curagua

were transported to New England by sailors, as previously suggested (Smith 1999), they would

have likely contained a limited gene pool, and subsequent inbreeding beginning in the 1940s

would have exacerbated the limited genetic diversity. However, it appears that after development

in the United States, some of this germplasm returned to Latin America, as evidenced by the

SONO 147 and BRAZ 2832 accessions (Fig. 4). Although little additional information is known

about BRAZ 2832, the passport data for SONO 147 (National Genetic Resources Program,

2022) contains the alternate name "Reventador Gringo", suggesting that growers associated this

population with non-native material.

4.3 Kernel color, popcorn populations, and nicosulfuron injury

The product of the *Y1* gene catalyzes the first step in the maize carotenoid biosynthesis pathway.

A SNP in this gene has been identified as a causal mutation that results in a Thr to Asn

substitution (Fu et al. 2013). The peak SNP identified in the GWAS is tightly linked with the

causal SNP, separated by 346 bases. Another SNP occurs close to a gene model predicted to

have carotene desaturase activity and is located 3 Mb away from a locus previously associated

with popcorn flake color in CIMMYT germplasm (Li et al. 2021). Given the structural variation

between popcorn germplasm and most material used to evaluate kernel color, it is plausible that

additional loci may be identified outside the core genome (Lu et al. 2015, Hufford et al. 2021).

The three remaining SNPs reported here have not been previously associated with kernel color,

although two occur in known genes, *wrky11* and *nc3*, and the other in an uncharacterized gene.

Further investigation is necessary to determine whether these additional associations represent

biologically meaningful relationships or spurious correlations from population structure.

Recent studies have been unable to identify differences in herbicide sensitivity among several

(i.e., up to eight entries) commercial white and yellow popcorn hybrids (Barnes et al. 2020a;

Barnes et al. 2020b). However, given the unknown origin of the hybrid parents used in these

studies and the prevalence of North American Yellow Pearl Popcorns in commercial germplasm,

these hybrids were likely selected using kernel color alone and do not represent the two North

American popcorn populations. The results presented here show significant differences in the

two popcorn populations for susceptibility to nicosulfuron. Kernel color has no bearing on

herbicide injury and demonstrates the importance of population structure in genetic analyses.

Kernel color represents a major phenotypic difference between the generally nicosulfuron-

tolerant North American Yellow Pearl Popcorns and the more susceptible Pointed and Latin

American Popcorns.

Nicosulfuron, along with other sulfonylurea herbicides, inhibits acetolactate synthase (ALS), the

first enzyme in the biosynthesis of branched-chain amino acids (BCAAs). Although encoded by

the nuclear genome, ALS is localized in the chloroplast (Mazur et al. 1987). Along with other

enzymes involved in BCAA biosynthesis, ALS interacts with many substrates and products of

cellular respiration, including pyruvate, NAD, and FAD (Duggleby et al. 2008). While target-site

resistance of sulfonylurea herbicides has been demonstrated in maize (Anderson & Georgeson

1989; Li et al. 2020), non-target site resistance is more common and complex (Délye 2013).

Non-target site resistance involves initial detoxification, often in the form of cytochrome p450s

and other oxidases, followed by conjugation of thiols or glucosyl groups, and subsequent

transport and degradation in the vacuole or extracellular spaces (Yuan et al. 2007).

In sweet and dent corn, detoxification of several post-emergent herbicides, including

nicosulfuron, is driven largely by *CYP81A9*, a cytochrome p450 produced by the *Nsf1* locus

(Nordby et al. 2008, Choe and Williams 2020). Although this does not appear to be the case with

nicosulfuron response in popcorn, other candidate genes were identified. Since the pool of

available NADH in oxidative phosphorylation is driven largely by pyruvate, variation in NADH

dehydrogenase may allow plants to tolerate changes in pyruvate as ALS becomes bound by

nicosulfuron. Additionally, variation in glycosylation and changes in the vacuole is consistent

with non-target site resistance. Furthermore, altered amino acid synthesis and phosphorylation

may be involved in the alternate metabolic flux of some BCAAs (Joshi et al. 2010). Finally, as

many of these processes occur within, or are related to, the chloroplast, changes in the

transcription factors of chloroplast-associated genes would be expected.

Although *CYP72A27* is a cytochrome p450, its identification here is somewhat unexpected, as it

belongs to the CYP clan 72 group of cytochrome p450s. Members of this clan are generally

associated with secondary metabolism, while clan 71 enzymes, such as *CYP81A9*, are

responsible for herbicide detoxification (Prall et al. 2016; Brazier-Hicks et al. 2022). However,

sulfonylurea herbicide metabolism via clan 72 enzymes has been shown in other crops, such as

CYP72A31 in *Oryza indica* (Saika et al. 2014) and *CYP749A16* in *Gossypium hirsutum* (Thyssen

et al. 2018). Furthermore, several members of CYP72 were shown to be differentially expressed

in nicosulfuron tolerant and susceptible lines of maize when treated with the herbicide (Liu et al.

2015). Alternatively, the role of *CYP72A27* in detoxifying nicosulfuron may be unique to the

Pointed and Latin American Popcorns and simply not previously reported. However, it is

possible that rather than directly detoxifying nicosulfuron, *CYP72A27* and other genes identified

here respond more like safener-induced enzymes, conferring protection via a stress signaling

pathway (Riechers et al. 2010). Finally, as with other candidate genes identified using GWAS,

CYP72A27 may have a spurious correlation with nicosulfuron response. Determining the roles

and mechanisms of the genes presented here with nicosulfuron tolerance would require further

investigation, including the phenotyping and genotyping of additional members of the Pointed

and Latin American Popcorns.

5 CONCLUSION

High-quality genetic resources are important tools for maize breeding programs. A collection of

362 popcorn accessions were analyzed here using 417,218 SNPs. Two distinct populations of

popcorn were identified that exhibit genetic and phenotypic differences. North American Yellow

Pearl Popcorns constitute a pool of germplasm with reduced genetic diversity and greater rates of

inbreeding when compared with the Pointed and Latin American Popcorns. Kernel color

frequencies were shown to differ between the two populations, along with sensitivity to

nicosulfuron. Novel candidate genes responsible for nicosulfuron tolerance were identified in

popcorn. The genomic characterization provided here should be incorporated into popcorn

breeding programs to direct improvement and accelerate the rate of genetic gain. Additionally,

this collection of popcorn germplasm and genetic data can be used by the maize research

community to explore the genetic architecture of traits in popcorn and be compared with loci

identified in other populations of maize.

SUPPLEMENTAL MATERIAL

Supplemental Table S1 ADMIXTURE assignment probabilities of 362 popcorn accessions by

population.

Supplemental Table S2 Significant markers identified by GWAS of kernel color.

Supplemental Table S3 Nicosulfuron injury of 294 accessions, including 111 North American

Yellow Pearl Popcorns, 178 Pointed and Latin American Popcorns, and five controls.

Supplemental Table S4 Significant markers identified by GWAS of nicosulfuron injury.

Supplemental Figure S1 SNP density and distribution from the Core Analysis set across

chromosomes at a 1Mb window size.

Supplemental Figure S2 Admixture cross-validation plot for 362 popcorn accessions using the

Core Analysis set.

Supplemental Figure S3 Neighbor-joining tree of 150 maize and 17 teosinte accessions based on

a modified Euclidean distance matrix calculated from the Core Analysis set.

ACKNOWLEDGMENTS

The authors thank Nicholas Hausman for managing the field design, planting, and nicosulfuron

application, Crookham Company for the popcorn hybrids used in the herbicide trial, and Marlee

Labroo, Lucas Roberts, and Lynn Doran for assistance with library preparation.

AUTHOR CONTRIBUTIONS

Madsen Sullivan: Conceptualization, Data curation, Formal analysis, Investigation,

Methodology, Software, Validation, Visualization, Writing – original draft, Writing – review &

editing. Martin M. Williams II: Conceptualization, Methodology, Resources, Writing – review &

editing. Anthony J. Studer: Conceptualization, Funding acquisition, Methodology, Project

administration, Resources, Supervision, Writing – original draft, Writing – review & editing.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

673
Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in
unrelated individuals. *Genome research*, *19*(9), 1655-1664.
Anderson, P. C., & Georgeson, M. (1989). Herbicide-tolerant mutants of corn. *Genome*, *31*(2),
994-999.Li, Y., Zhu, J., Wu, H., Liu, C., Huang, C., Lan, J., ... & Xie, C. (2020). Precise base
editing of non-allelic acetolactate synthase genes confers sulfonylurea herbicide resistance in
maize. *The Crop Journal*, *8*(3), 449-456.
Barnes, E. R., Lawrence, N. C., Knezevic, S. Z., Irmak, S., Rodriguez, O., & Jhala, A. J. (2020).
Dose response of yellow and white popcorn hybrids to glyphosate, a premix of 2, 4-D choline
and glyphosate, or dicamba. *Agronomy Journal*, *112*(4), 2956-2967.
Barnes, E. R., Lawrence, N. C., Knezevic, S. Z., Rodriguez, O., Irmak, S., & Jhala, A. J. (2020).
Weed control and response of yellow and white popcorn hybrids to herbicides. *Agronomy*
Journal, *112*(1), 458-469.
Bautista-Ramírez, E., Santacruz-Varela, A., Córdova-Téllez, L., Muñoz Orozco, A., López-
Sánchez, H., & Esquivel-Esquivel, G. (2020). Yield and expansion capacity of the corn grain in
the Palomero Toluqueño race. *Revista mexicana de ciencias agrícolas*, *11*(7), 1607-1618.
Bedoya, C. A., Dreisigacker, S., Hearne, S., Franco, J., Mir, C., Prasanna, B. M., ... &
Warburton, M. L. (2017). Genetic diversity and population structure of native maize populations
in Latin America and the Caribbean. *PloS one*, *12*(4), e0173488.
Bracco, M., Lia, V. V., Hernández, J. C., Poggio, L., & Gottlieb, A. M. (2012). Genetic diversity
of maize landraces from lowland and highland agro-ecosystems of Southern South America:
implications for the conservation of native resources. *Annals of Applied Biology*, *160*(3), 308-
321.
Brazier-Hicks, M., Franco-Ortega, S., Watson, P., Rougemont, B., Cohn, J., Dale, R., ... &
Edwards, R. (2022). Characterization of Cytochrome P450s with Key Roles in Determining
Herbicide Selectivity in Maize. *ACS omega*.
Broman, K. W., Wu, H., Sen, Ś., & Churchill, G. A. (2003). R/qtl: QTL mapping in experimental
crosses. *bioinformatics*, *19*(7), 889-890.
Browning, B. L. (2018). Beagle 5.0. *University of Washington, Division of Medical Genetics*.
Brunson, A. M. (1937). Popcorn breeding.
Buckler, E. S., Holland, J. B., Bradbury, P. J., Acharya, C. B., Brown, P. J., Browne, C., ... &
McMullen, M. D. (2009). The genetic architecture of maize flowering time. *Science*, *325*(5941),
714-718.
Choe, E., & Williams, M. M. (2020). Expression and comparison of sweet corn CYP81A9s in
relation to nicosulfuron sensitivity. *Pest Management Science*, *76*(9), 3012-3019.
Coles, N. D., McMullen, M. D., Balint-Kurti, P. J., Pratt, R. C., & Holland, J. B. (2010). Genetic
control of photoperiod sensitivity in maize revealed by joint multiple population
analysis. *Genetics*, *184*(3), 799-812.
Cook, J. P., McMullen, M. D., Holland, J. B., Tian, F., Bradbury, P., Ross-Ibarra, J., ... & Flint-
Garcia, S. A. (2012). Genetic architecture of maize kernel composition in the nested association
mapping and inbred association panels. *Plant physiology*, *158*(2), 824-834.
Da Fonseca, R. R., Smith, B. D., Wales, N., Cappellini, E., Skoglund, P., Fumagalli, M., ... &
Gilbert, M. T. P. (2015). The origin and evolution of maize in the Southwestern United
States. *Nature plants*, *1*(1), 1-5.

De Almeida Silva, N. C., Vidal, R., Bernardi Ogliari, J., E Costich, D., & Chen, J. (2020).
Relationships among American popcorn and their links with landraces conserved in a
microcenter of diversity. *Genetic Resources and Crop Evolution*, *67*(7), 1733-1753.
Délye, C. (2013). Unravelling the genetic bases of non-target-site-based resistance (NTSR) to
herbicides: a major challenge for weed science in the forthcoming decade. *Pest management*
science, *69*(2), 176-187.
Duggleby, R. G., McCourt, J. A., & Guddat, L. W. (2008). Structure and mechanism of
inhibition of plant acetohydroxyacid synthase. *Plant Physiology and Biochemistry*, *46*(3), 309-
324.
Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K., Buckler, E. S., & Mitchell,
S. E. (2011). A robust, simple genotyping-by-sequencing (GBS) approach for high diversity
species. *PloS one*, *6*(5), e19379.
Erwin, A. T. (1950). The origin and history of pop corn. *Economic Botany*, *4*(3), 294-299.
Fei, J., Jiang, Q., Guo, M., Lu, J., Wang, P., Liu, S., ... & Guan, S. (2022). Fine Mapping and
Functional Research of Key Genes for Photoperiod Sensitivity in Maize. *Frontiers in Plant*
Science, *13*.
Fu, Z., Chai, Y., Zhou, Y., Yang, X., Warburton, M. L., Xu, S., ... & Yan, J. (2013). Natural
variation in the sequence of PSY1 and frequency of favorable polymorphisms among tropical
and temperate maize germplasm. *Theoretical and Applied Genetics*, *126*(4), 923-935.
Gage, J. L., Monier, B., Giri, A., & Buckler, E. S. (2020). Ten years of the maize nested
association mapping population: impact, limitations, and future directions. *The Plant Cell*, *32*(7),
2083-2093.
Glaubitz, J. C., Casstevens, T. M., Lu, F., Harriman, J., Elshire, R. J., Sun, Q., & Buckler, E. S.
(2014). TASSEL-GBS: a high capacity genotyping by sequencing analysis pipeline. *PloS*
one, *9*(2), e90346.
Goodman, M. M., Jones, Z. G., Sanchez, G. J., & Kermicle, J. L. (2021). Maize Cross
Incompatibility and the Promiscuous Ga1-m Allele. *Plant breeding reviews*, *44*, 31-56.
Hufford, M. B., Seetharam, A. S., Woodhouse, M. R., Chougule, K. M., Ou, S., Liu, J., ... &
Dawe, R. K. (2021). De novo assembly, annotation, and comparative analysis of 26 diverse
maize genomes. *Science*, *373*(6555), 655-662.
Intarapanich, A., Shaw, P. J., Assawamakin, A., Wangkumhang, P., Ngamphiw, C.,
Chaichoompu, K., ... & Tongsima, S. (2009). Iterative pruning PCA improves resolution of
highly structured populations. *BMC bioinformatics*, *10*(1), 1-17.
Joshi, V., Joung, J. G., Fei, Z., & Jander, G. (2010). Interdependence of threonine, methionine
and isoleucine metabolism in plants: accumulation and transcriptional regulation under abiotic
stress. *Amino acids*, *39*(4), 933-947.
Kantety, R. V., Zeng, X., Bennetzen, J. L., & Zehr, B. E. (1995). Assessment of genetic diversity
in dent and popcorn (Zea mays L.) inbred lines using inter-simple sequence repeat (ISSR)
amplification. *Molecular breeding*, *1*(4), 365-373.
Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*
methods, *9*(4), 357-359.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... & Durbin, R. (2009).
The sequence alignment/map format and SAMtools. *Bioinformatics*, *25*(16), 2078-2079.
Li, J., Chen, G. B., Rasheed, A., Li, D., Sonder, K., Zavala Espinosa, C., ... & Li, H. (2019).
Identifying loci with breeding potential across temperate and tropical adaptation via
EigenGWAS and EnvGWAS. *Molecular ecology*, *28*(15), 3544-3560.

Li, J., Li, D., Espinosa, C. Z., Pastor, V. T., Rasheed, A., Rojas, N. P., ... & Li, H. (2021). Genome-wide analyses reveal footprints of divergent selection and popping-related traits in CIMMYT's maize inbred lines. *Journal of experimental botany*, *72*(4), 1307-1320.

Linck, E., & Battey, C. J. (2019). Minor allele frequency thresholds strongly affect population structure inference with genomic data sets. *Molecular Ecology Resources*, *19*(3), 639-647.

Liu, X., Xu, X., Li, B., Wang, X., Wang, G., & Li, M. (2015). RNA-seq transcriptome analysis of maize inbred carrying nicosulfuron-tolerant and nicosulfuron-susceptible alleles. *International Journal of Molecular Sciences*, *16*(3), 5975-5989.

Loux, M. M., Doohan, D., Dobbels, A. F., Johnson, W. G., Young, B. G., Legleiter, T. R., & Hager, A. (2017). Weed control guide for Ohio, Indiana and Illinois.

Lu, F., Romay, M. C., Glaubitz, J. C., Bradbury, P. J., Elshire, R. J., Wang, T., ... & Buckler, E. S. (2015). High-resolution genetic mapping of maize pan-genome sequence anchors. *Nature communications*, *6*(1), 1-8.

Matz, S. A. (1984). Snacks based on popcorn. In *Snack food technology* (pp. 138-149). Springer, Dordrecht.

Matz, Samuel A. "Snacks based on popcorn." *Snack food technology*. Springer, Dordrecht, 1984. 138-149.

Mazur, B. J., Chui, C. F., & Smith, J. K. (1987). Isolation and characterization of plant genes coding for acetolactate synthase, the target enzyme for two classes of herbicides. *Plant Physiology*, *85*(4), 1110-1117.

Merrill, W. L., Hard, R. J., Mabry, J. B., Fritz, G. J., Adams, K. R., Roney, J. R., & MacWilliams, A. C. (2009). The diffusion of maize to the southwestern United States and its impact. *Proceedings of the National Academy of Sciences*, *106*(50), 21019-21026.

Miller, J. M., Cullingham, C. I., & Peery, R. M. (2020). The influence of a priori grouping on inference of genetic clusters: simulation study and literature review of the DAPC method. *Heredity*, *125*(5), 269-280.

Nordby, J. N., Williams, M. M., Pataky, J. K., Riechers, D. E., & Lutz, J. D. (2008). A common genetic basis in sweet corn inbred Cr1 for cross sensitivity to multiple cytochrome P450-metabolized herbicides. *Weed Science*, *56*(3), 376-382.

Okonechnikov, K., Conesa, A., & García-Alcalde, F. (2016). Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics*, *32*(2), 292-294.

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in R language. *Bioinformatics*, *20*(2), 289-290.

Peiffer, J. A., Romay, M. C., Gore, M. A., Flint-Garcia, S. A., Zhang, Z., Millard, M. J., ... & Buckler, E. S. (2014). The genetic architecture of maize height. *Genetics*, *196*(4), 1337-1356.

Prall, W., Hendy, O., & Thornton, L. E. (2016). Utility of a phylogenetic perspective in structural analysis of CYP72A enzymes from flowering plants. *PloS one*, *11*(9), e0163024.

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, *155*(2), 945-959.

Riechers, D. E., Kreuz, K., & Zhang, Q. (2010). Detoxification without intoxication: herbicide safeners activate plant defense gene expression. *Plant Physiology*, *153*(1), 3-13.

Romay, M. C., Millard, M. J., Glaubitz, J. C., Peiffer, J. A., Swarts, K. L., Casstevens, T. M., ... & Gardner, C. A. (2013). Comprehensive genotyping of the USA national maize inbred seed bank. *Genome biology*, *14*(6), 1-18.

Saika, H., Horita, J., Taguchi-Shiobara, F., Nonaka, S., Nishizawa-Yokoi, A., Iwakami, S., ... & Toki, S. (2014). A novel rice cytochrome P450 gene, CYP72A31, confers tolerance to

acetolactate synthase-inhibiting herbicides in rice and Arabidopsis. *Plant physiology*, *166*(3),
1232-1240.
Santacruz-Varela, A., Widrlechner, M. P., Ziegler, K. E., Salvador, R. J., Millard, M. J., &
Bretting, P. K. (2004). Phylogenetic relationships among North American popcorns and their
evolutionary links to Mexican and South American popcorns. *Crop Science*, *44*(4), 1456-1467.
Serna-Saldivar, S. O. (2022). Popcorn and Other Puffed Grains. In *Snack Foods* (pp. 201-220).
CRC Press.
Smith, A. F. (1999). *Popped culture: A social history of popcorn in America*. Univ of South
Carolina Press.
Swarts, K., Li, H., Romero Navarro, J. A., An, D., Romay, M. C., Hearne, S., ... & Bradbury, P.
J. (2014). Novel methods to optimize genotypic imputation for low-coverage, next-generation
sequence data in crop plants. *The Plant Genome*, *7*(3), plantgenome2014-05.
Tam, V., Patel, N., Turcotte, M., Bossé, Y., Paré, G., & Meyre, D. (2019). Benefits and
limitations of genome-wide association studies. *Nature Reviews Genetics*, *20*(8), 467-484.
Team, R. C. (2021). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. 2012.
Thyssen, G. N., Naoumkina, M., McCarty, J. C., Jenkins, J. N., Florane, C., Li, P., & Fang, D. D.
(2018). The P450 gene CYP749A16 is required for tolerance to the sulfonylurea herbicide
trifloxysulfuron sodium in cotton (Gossypium hirsutum L.). *BMC plant biology*, *18*(1), 1-8.
Turner, S. D. (2018). qqman: an R package for visualizing GWAS results using QQ and
manhattan plots. J. Open Source Softw. 3: 731.
USDA NASS. (2022). Quick Stats. US Department of Agriculture. National Agricultural
Statistics Service. https://quickstats.nass.usda.gov/
Van Heerwaarden, J., Doebley, J., Briggs, W. H., Glaubitz, J. C., Goodman, M. M., de Jesus
Sanchez Gonzalez, J., & Ross-Ibarra, J. (2011). Genetic signals of origin, spread, and
introgression in a large sample of maize landraces. *Proceedings of the National Academy of*
Sciences, *108*(3), 1088-1092.
Vázquez-Carrillo, M. G., Santiago-Ramos, D., & de Dios Figueroa-Cárdenas, J. (2019). Kernel
properties and popping potential of Chapalote, a Mexican ancient native maize. *Journal of*
Cereal Science, *86*, 69-76.
Vigouroux, Y., Glaubitz, J. C., Matsuoka, Y., Goodman, M. M., Sánchez G, J., & Doebley, J.
(2008). Population structure and genetic diversity of New World maize races assessed by DNA
microsatellites. *American journal of botany*, *95*(10), 1240-1253.
Vittorazzi, C., Junior, A. A., Guimarães, A. G., Silva, F. H. L., Pena, G. F., Daher, R. F., ... &
Lima, V. J. (2018). Evaluation of genetic variability to form heterotic groups in popcorn. *Genet.*
Mol. Res, *17*, 18083.
Wallace, J. G., Larsson, S. J., & Buckler, E. S. (2014). Entering the second century of maize
quantitative genetics. *Heredity*, *112*(1), 30-38.
Wang, L., Beissinger, T. M., Lorant, A., Ross-Ibarra, C., Ross-Ibarra, J., & Hufford, M. B.
(2017). The interplay of demography and selection during maize domestication and
expansion. *Genome biology*, *18*(1), 1-13.
White, M. R., Mikel, M. A., de Leon, N., & Kaeppler, S. M. (2020). Diversity and heterotic
patterns in North American proprietary dent maize germplasm. *Crop Science*, *60*(1), 100-114.
Wickham, H., Chang, W., & Wickham, M. H. (2016). Package 'ggplot2'. *Create elegant data*
visualisations using the grammar of graphics. Version, *2*(1), 1-189.

Yin, L., Zhang, H., Tang, Z., Xu, J., Yin, D., Zhang, Z., ... & Liu, X. (2021). rMVP: a memory-
efficient, visualization-enhanced, and parallel-accelerated tool for genome-wide association
study. *Genomics, proteomics & bioinformatics*, *19*(4), 619-628.
Yu, D., Wang, H., Gu, W., Qin, T., Sun, P., Lu, Y., ... & Zheng, H. (2021). Genetic diversity and
population structure of popcorn germplasm resources using genome-wide SNPs through
genotyping-by-sequencing. *Genetic Resources and Crop Evolution*, *68*(6), 2379-2389.
Yu, G., Smith, D. K., Zhu, H., Guan, Y., & Lam, T. T. Y. (2017). ggtree: an R package for
visualization and annotation of phylogenetic trees with their covariates and other associated
data. *Methods in Ecology and Evolution*, *8*(1), 28-36.
Yuan, J. S., Tranel, P. J., & Stewart Jr, C. N. (2007). Non-target-site herbicide resistance: a
family business. *Trends in plant science*, *12*(1), 6-13.
Ziegler, Kenneth E. "Popcorn." *Specialty corns*. CRC Press, 2000. 211-246.