M. tuberculosis antigen-responsive IL17[+] CD4 T cells are disproportionately spared in

ART-suppressed HIV

Paul Ogongo[1,2], Anthony Tran[1], Florence Marzan[3], David Gingrich[3], Melissa Krone[4],

Francesca Aweeka[3], Cecilia S Lindestam Arlehamn[5,6], Jeffrey N. Martin[4], Steven G. Deeks[7],

Peter W. Hunt[1#*] and Joel D. Ernst[1#*]

[1]Division of Experimental Medicine, University of California, San Francisco, CA, USA

[2]Department of Tropical and Infectious Diseases, Institute of Primate Research, Nairobi,

Kenya

[3]Drug Research Unit, Department of Clinical Pharmacy, School of Pharmacy, University of

California, San Francisco, CA, USA

[4]Department of Epidemiology and Biostatistics, University of California, San Francisco, CA,

USA

[5]Center for Infectious Disease and Vaccine Research, La Jolla Institute for Immunology, La

Jolla, CA 92037, USA

[6]Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy

Chase, MD, USA

[7]Division of HIV, Infectious Diseases, and Global Medicine, University of California, San

Francisco, CA, USA

[#] Joint senior authors

* Correspondence

Running title: Mtb-responsive Th17 cells during HIV-ART

Abstract

Background. Interleukin 17 producing CD4 T cells contribute to control of *Mycobacterium tuberculosis (Mtb)* infection in humans; whether infection with Human Immunodeficiency Virus (HIV) disproportionately affects distinct Th17 cell subsets that respond to Mtb are incompletely defined.

Methods. We performed high-definition characterization of circulating Mtb-specific Th17 cells by spectral flow cytometry in people with latent TB and treated HIV (HIV-ART). We also measured kynurenine pathway activity in plasma by LC/MS and tested the hypothesis that tryptophan catabolism influences Th17 cell differentiation in this context.

Results. We identified two categories of Th17 cells: T_H17 ($CD4^+V\alpha7.2^-CD161^+CD26^+$) and T17 ($CD4^+V\alpha7.2^-CCR6^+CXCR3^-$) cells that were disproportionately reduced in LTBI with HIV-ART, yet Mtb-responsive IL17-producing CD4 T cells were preserved; we found that IL17-producing CD4 T cells dominate the response to Mtb antigen but not CMV antigen or staphylococcal enterotoxin B (SEB); and tryptophan catabolism negatively correlates with T_H17 and T1T17 but not T17 cell frequencies.

Conclusions. We found differential effects of ART-suppressed HIV on distinct categories of Th17 cells, that IL17-producing CD4 T cells dominate responses to Mtb but not CMV antigen or SEB, and that kynurenine pathway activity is associated with selective decreases of circulating Th17 cells that may contribute to tuberculosis immunity.

Key words: Interleukin 17; CD4 T cells; antigen-responsive; immunity; Tuberculosis; ART-suppressed HIV; kynurenine pathway

Introduction

Introduction

Tuberculosis (TB) remains the leading cause of death among people with HIV (PWH) [1] at

least in part because of CD4 T cell depletion [2]. CD4 T cells are critical for immunity to TB,

although the CD4 T cell phenotypes, functions, and antigen specificity that contribute to TB

immunity are incompletely understood. Importantly, the risk of TB in PWH remains higher

than in those without HIV, even after reconstitution by antiretroviral therapy (ART) [3,4],

indicating that the effects of HIV on TB immunity are not simply because of loss of bulk CD4

T cells. Indeed, HIV may preferentially infect Mtb-specific CD4 T cells [5,6], and one or more

defects in Mtb-specific CD4 T cells might be incompletely restored by ART.

Increasing evidence indicates that CD4 T cells that produce interleukin 17 (IL17), broadly

termed T helper 17 (Th17) cells, contribute to control of TB. In Mtb-infected adolescents, IL17

transcriptional signatures decreased in blood of those who progressed to active TB compared

to non-progressors [7]; another study found a subset of CD4 T cells that produce IL-17 in

response to Mtb antigens that is less abundant in TB progressors than in non-progressors [8].

Mtb-responsive CD4$^+$ T cells producing IL17 are enriched in human lungs compared to

matched blood and inversely correlated with plasma IL-1β, suggesting a role in control of Mtb

[9]. IL17 producing CD4 T cells are depleted in HIV infection and their depletion contributes

to progression to AIDS through breakdown in intestinal mucosal barrier function [10,11].

Several approaches are used to identify T helper 17 cells (herein denoted 'Th17' cells to include all

subsets regardless of criteria used). IL17 production upon stimulation is the canonical signature of

Th17 cells [12,13]. Although this identifies Th17 cells, it does not classify all cells in the Th17

lineage, and other cells can produce IL17. In blood of healthy donors and synovial fluid of

rheumatoid arthritis patients, a CCR6[+] subset contained all IL17-producing T cells expressing

RORC mRNA [14], the transcription factor that supports Th17 differentiation in humans. Use of

CCR6 as a marker of Th17 cells has been replicated in naïve cord blood [15], inflammatory

diseases [16] and infections like TB [17]. The cell surface markers CD26 [18,19] and CD161

[20,21] have also been used to identify Th17 cells. CD26[hi] CD4[+] T cells were demonstrated to co-

express CD161 and CCR6 and enriched for production of IL17 [18]. In transcriptomic studies,

Th17 cells are identified by expression of *RORC, IL23R* (supports expansion of IL17-producing

cells) and *IL17* mRNA.

Host metabolism impacts immune responses [22] by driving cell differentiation. One example

is the kynurenine pathway of tryptophan catabolism whose products, collectively called

kynurenines, inhibit Th17 while promoting Treg development. [23–26]. In TB, circulating

tryptophan concentrations ([Trp]) decline in persons with LTBI progressing to active TB [27]

and [Trp] increases with TB treatment [23]. Similarly, HIV infected persons have low

circulating [Trp] with a link to pathogenesis [28]. Plasma kynurenine/tryptophan ratios (K/T)

can distinguish humans with active TB from those without and are significantly higher in

MDR-TB patients with or without HIV coinfection compared to controls [23]. Plasma K/T is

also elevated in PWH compared to controls, especially in those who have progressed to

AIDS[25,29]. While a link between K/T and Treg/Th17 ratio has been established in HIV,

similar data are lacking in TB.

Here, we characterized subsets of Th17 cells defined by distinct criteria in participants with

ART-suppressed HIV (HIV-ART), to determine whether IL17-producing CD4 T cells that

respond to Mtb antigens are altered in HIV-ART, and to test the hypothesis that indoleamine

2,3-dioxygenase (IDO) activity and generation of kynurenines influences Th17 differentiation

in HIV-ART.

Materials and Methods

Participants and sample collection

PBMC and plasma were obtained from adults with or without HIV enrolled in the UCSF

SCOPE cohort [25]. All participants gave written informed consent using protocols approved

by the UCSF Instititutional Review Board. Blood was collected into EDTA-containing tubes

for cell counts and into ACD-containing tubes for purification of plasma and PBMCs.

Processed plasma was stored at -80°C while PBMC were cryopreserved in liquid nitrogen.

Demographic characteristics of the participants are in Tables 1 and 2.

Plasma Tryptophan and Kynurenine measurement

Tryptophan and Kynurenine concentrations were measured by liquid chromatography-mass

spectrometry as previously described [25,29].

PBMC antigen stimulation and Intracellular cytokine staining

Cryopreserved PBMCs were thawed in a 37°C water bath and transferred into pre-warmed R10

media (RPMI 1640 containing L-glutamine with 10% FBS, 1%PenStrep and 1% Hepes). Cells

were centrifuged at 2000rpm for 5 minutes at room temperature, supernatants discarded, and

cells resuspended before adding 5ml of warm R10 and transfer to a 6-well culture plate and

resting overnight in 37°C/5%CO$_2$ incubator. After resting, $1\times10^{\wedge6}$ live cells resuspended in

200μL R10 were transferred to each well of a 96-well round bottom plate and duplicate wells

were stimulated with Mtb peptide mega pool (Mtb300) [30](2μg/ml), CMV pp65 peptide

(1μg/ml) or Staphyloccal enterotoxin B (SEB) positive control (1μg/ml) in the presence of

costimulating antibodies anti CD28 (1µg/ml) and anti CD49d (1µg/ml) (BD). Negative

controls received no stimulation. After two hours, GolgiStop and GolgiPlug (BD) were added

to one well of each of the stimulation conditions, and the cells were incubated for additional

18 hours.

After 20-hour total stimulation, cells were washed, stained with Live/Dead Fixable Blue Dead

Cell stain kit (Invitrogen), washed again, then surface antibody cocktail (αCD3 BV510 clone

UCHT1, αCD8 BV570 clone RPA-T8, αCCR7 BV785 clone G043H7, αCD95 Alexa Fluor700

clone DX2, αCCR6 FITC clone G034E3, αCXCR3 BV 605 clone G025H7, αCD161 APC-

Fire750 clone HP-3G10, αCD69 BV650 clone FN50, αCD137 BV711 clone 4B4-1, αOX40

PE-Cyanine7 clone (Ber-ACT35) and αTRAV1-2 (Vα7.2) BV421 clone 3C10 (all from

BioLegend), αCD4 BUV496 clone SK3, αCD45RA BUV395 clone 5H9, αCD27 BUV615

clone L128, αCD25 BUV563 clone 2A3, αCD39 BUV737 clone TU66 and αCD26 BUV805

clone M-A261 (all BD)) diluted in Brilliant Violet buffer (BD) was added then kept in the dark

for 20 minutes at room temperature., After washing, cells were fixed and permeabilized using

eBioscience FOXP3/Transcription Factor staining kit (Invitrogen). Cells were washed twice

with 1X eBioscience Perm diluent, then resuspended in intracellular antibody mix (αRORγT

Alexa Fluor 647 Clone Q21-559, αIFNγ BB700 clone 2B7 (both from BD Biosciences), αIL17

PE clone BL168, αT-bet PE-Dazzle 594 clone 4B10 (both from BioLegend) and αFoxP3 PE-

Cyanine5.5 clone PCH101 (Thermofisher)) in eBiosciences perm diluent. Cells were then

washed twice with 1X eBioscience Perm diluent and fixed in 2% PFA. Data acquired on a 5-

Laser Aurora Spectral Flow cytometer (Cytek).

Data and statistical analysis

Spectral flow fcs files were initially analysed using SpectroFlo v3.0 (Cytek) for unmixing and

autofluorescence correction. Identification of T cell subsets from unmixed fcs files was

performed using Flowjo v10 (Flowjo LLC). Antigen specific cytokine levels are reported after

subtraction of values of unstimulated cells for each participant. K/T ratio was determined from

the concentrations of tryptophan and kynurenines after quality control checks. Non-parametric

tests were used for comparison between two groups for paired (Wilcoxon test) or unpaired

(Mann-Whitney test) and $p < 0.05$ considered significant. Adjusted comparisons between

groups in continuous variables were performed with linear regression, transforming outcome

variables to satisfy model assumptions. Spearman correlation was used to test the association

between parameters with a $p < 0.05$ considered significant. All statistical analyses used Prism

9.0 (GraphPad).

Results

Subsets of circulating T helper 17 cells are disproportionately reduced in LTBI with treated

HIV

To study the effects of LTBI and ART-treated HIV (HIV-ART) on Th17 cells defined by

different criteria, we designed a spectral flow cytometry panel encompassing cytokines,

chemokine receptors, transcription factors and cell surface markers (Figure 1A&C). We used

the following nomenclature: (a) T_H17 are $CD26^+CD161^+$ [18,19]; (b) T17 are $CCR6^+CXCR3^-$

[14,17]; and (c) T1T17 are $CCR6^+$ $CXCR3^+$ [17,31]. We first gated out $V\alpha7.2^+$ $CD4^+$ T cells

to exclude MAIT cells that share certain markers with Th17 cells [32], since MAITs can

produce IL17 [33].

We also stained for RORγT, the Th17 lineage defining transcription factor [34] and used

unsupervised clustering and t-distributed stochastic neighbor embedding (tSNE) to determine

whether T_H17 and T17 cells express RORγT and/or produce IL17 upon stimulation with an

Mtb peptide mega pool (Mtb300) (Figure 1C). To identify T1T17 (also termed Th1*) cells

that respond to Mtb antigens [17], we included T-bet, the Th1 lineage transcription factor [35].

We found a positive and significant association (r = 0.4472, p = 0.0002) (Figure 1B) between

T_H17 and T17 cells, and found that cells in both populations express RORγT and produce IL17

(Figure 1C). Therefore, T_H17 and T17 cells express other properties of bona fide Th17 cell

populations. T-bet expressing cells were present within T_H17 and T17 cell populations

suggesting a Th1* population that expresses both CCR6 and CXCR3 and likely produce both

IFNγ and IL17 (Figure 1C).

As expected, circulating CD4 T cells were reduced in HIV-ART compared with HIV-

uninfected participants, (Figure 2A). We first phenotyped $CD4^+$ T cell populations based on

CD45RA and CCR7 [9] to identify naïve, central memory, effector memory and terminally

differentiated T cells (Figure S1 A, left). The predominant populations were naïve and central

memory T cells and there was no difference in the frequency of cells in these memory

populations in participants with or without HIV (Figure 2B). $CD4^+CD45RA^+CCR7^+$ (naïve T

cells) are a heterogenous population, including cells that produce cytokines upon short term

stimulation [36], so we characterised the naïve T cells further based on co-expression of CD95

and CD27 to identify cells with a stem cell-like memory phenotype (Figure S1 A, right). We

found no difference in frequencies of $CD95^+CD27^+$ stem cell-like memory cells or $CD95^-$

$CD27^+$ 'true naïve' cells in those with or without HIV (Figure S1 B). Thus, whereas individuals

with treated HIV have lower overall frequencies of circulating $CD4^+$ T cells than those without

HIV (Tables 1 and 2, Figure 2A), their memory states do not differ.

Next, we quantitated Th17 cells in those with LTBI without HIV or with HIV-ART and found

significantly lower frequencies of T_H17 (CD4$^+$Vα7.2$^-$CD26$^+$CD161$^+$) and T1T17 (CD4$^+$Vα7.2$^-$

CCR6$^+$CXCR3$^+$) cells (expressed as % of CD4$^+$ T cells) in HIV-ART than in those without

HIV (Figure 2C). Although T17 cells (CD4$^+$Vα7.2$^-$CCR6$^+$CXCR3$^-$) and Th1 (CD4$^+$Vα7.2$^-$

CCR6$^-$CXCR3$^+$) cells were also present at lower frequencies in HIV-ART, the difference was

not significant (Figure 2C). These data demonstrate that certain circulating Th17 cell subsets

are disproportionately depleted in HIV-ART, while others are relatively preserved.

Th17 and regulatory T cells (Tregs) develop under related but distinct cytokine environments

[37]. Tregs contribute to immune homeostasis by maintaining unresponsiveness to self-

antigens, suppressing exaggerated immune responses, and promoting epithelial tissue integrity

[38]. Human Tregs are identified by the expression of high levels of CD25 and the transcription

factor FoxP3 [39]. Since the kynurenine pathway has differential effects on Th17 and Treg

differentiation, we characterised Treg cells (Figure S2A, left) and found that the frequency of

circulating Treg was similar in HIV-ART and those without HIV (Figure S2B). We also found

that the frequency of activated (CD39$^+$) Tregs did not differ in the two groups (Figure S2A,

right) although there was considerable variation in CD39$^+$ Treg, especially in HIV-ART

(Figure S2 C). We also compared the ratio of T_H17 and T17 cells to Tregs between the groups

and found a trend toward a lower T_H17/Treg in HIV-ART than in those without HIV (Figure

S2D), but the difference was not significant, and T17/Treg ratios were similar in HIV-ART

and those without HIV (Figure S2E). Taken together, we found that certain circulating Th17

cell subsets are disproportionately reduced in LTBI with treated HIV, suggesting that even

effective treatment of HIV does not reconstitute all Th17 cell subsets equally. We did not find

differences in Treg by HIV status.

Mtb antigen-responsive IL17-producing cells are enriched in T_H17 compared with T17 populations in HIV-ART

Having established that $CD26^+CD161^+$ T_H17 and $CCR6^+$ T17 cells are bona fide Th17 populations, we determined whether T_H17 and T17 cells differ in IL17 production after stimulation. HIV-ART participants with LTBI had significantly more $IL17^+$ cells in T_H17 cells than in T17 cells upon stimulation with the Mtb300 peptide pool (Figure 3A, left). All except one participant had $IL17^+$ cells in T_H17 cells compared with 11 of 21 participants with $IL17^+$ cells in T17 cells (Figure 3A, left). We did not have enough cells from participants without HIV (cohort 1) for cytokine analysis. We confirmed this observation in cohort 2; regardless of the sampling time point, there were significantly more $IL17^+$ cells in T_H17 cells than in T17 cells (Figure 3A, right). We performed a similar analysis after stimulation with SEB and found an opposite pattern: IL17 production was significantly higher in T17 than T_H17 cells (Figure 3B). Additionally, CMV peptide stimulation induced equivalent production of IL17 by T_H17 and T17 cells, although there was a trend towards higher responses in T17 cells (Figure S3A). These data reveal enrichment of IL17 production to Mtb antigens by a subset of Th17 cells marked by expression of CD26 and CD161 (T_H17) which are distinct from IL17 producing cells that are marked by expression of chemokine receptor CCR6 (T17) and those that produce IL17 in response to SEB or CMV.

Mtb antigen-responsive IL17-producing $CD4^+$ T cells are relatively preserved in people with LTBI and HIV-ART

Next, we assessed IFNγ and IL17 production after Mtb300 stimulation. In HIV-ART, only 4 of 21 (19%) had detectable IFNγ responses compared with 11 of 21 (52%) who had detectable IL17 responses (p = 0.0516, Fisher's exact test). Additionally, the median IL17 response was

higher than the median IFNγ response although this did not reach significance (Figure 3C, left).

Similar IL-17 responses to Mtb antigen stimulation were observed in people with HIV before

and after LTBI diagnosis and treatment. Treatment of LTBI did not change either IFNγ or IL17

responses although the response magnitudes varied by participant (Figure 3C, right). Similarly,

more participants had detectable IL17 responses than detectable IFNγ responses at all 3

sampling times: prior to being TST positive, and at INH initiation and treatment completion:

2/11 vs 10/11, 3/11 vs 9/11 and 6/14 vs 12/14, respectively (Figure 3C, right).

To determine whether predominant production of IL17 over IFNγ was specific to Mtb antigens,

we used a CMV peptide pool to stimulate PBMC from participants (Tables 1 and 2) who had

sufficient cells (Figure 3D). This revealed a pattern distinct from that with Mtb300: participants

with LTBI and HIV-ART (cohort 1) exhibited higher magnitude and higher prevalence of IFNγ

than IL17 responses (IFNγ: 11/16 (69%) vs IL17: 5/16 (31%)) (Figure 3D, left) (p = 0.0756,

Fisher's exact test). Similarly, in Cohort 2 there was a higher magnitude of IFNγ responses

than IL17 responses at all time points, reaching statistical significance at baseline (prior to

TST$^+$) timepoint (Figure 3D – right). As with CMV and in contrast to Mtb300, SEB induced

higher magnitude IFNγ than IL17 responses (Figure S3B). Together, these data demonstrate

that in LTBI and HIV-ART, IL17 responses to Mtb are relatively preserved compared with

IFNγ responses, and this is distinct compared to CMV and SEB responses. This is also contrary

to the commonly held impression that IFNγ is the predominant response to Mtb.

IDO activity is increased in people with LTBI and HIV-ART, and negatively correlates with

circulating Th17 cells

We quantitated tryptophan and kynurenine concentrations in plasma of participants with LTBI

with or without HIV-ART and inferred IDO activity based on the K/T ratio. We found

significantly lower plasma tryptophan concentrations in participants with HIV-ART compared

with those without HIV (Figure 4A) and kynurenine concentrations were higher in HIV-ART

but did not achieve statistical significance when compared with the HIV uninfected group

(Figure 4B). Finally, there was a significantly higher K/T ratio in people with HIV-ART

compared to those without HIV (Figure 4C). Although we cannot ascertain the contribution of

LTBI to the observed IDO activity because we did not have participants who were not exposed

to Mtb, these observations agree with prior reports showing that IDO-1 mediated tryptophan

catabolism is elevated in PWH, including those initiated on early treatment and ART-

suppressed viral loads [29].

Since kynurenines favour development of Tregs while suppressing Th17 generation [25], we

determined the correlation of plasma K/T and Th17 cells, Tregs, and Th17/Treg ratios in

Cohort 1 participants. This revealed a significant inverse relationship between K/T and

circulating T_H17 and T17 (Figure 4D-E), while there was no correlation between K/T and Tregs

(Figure 4D, middle). We also found significant inverse correlations between K/T and

T_H17/Treg and T17/Treg (Figure 4D-E). Importantly, we found no correlation between K/T

and T1T17, Th1, or IL17$^+$ cell frequencies (Fig. S4A-B and data not shown). Although we

confirmed high IDO activity (Figure 4) and found reduction in circulating Th17 (especially

T_H17) cells in HIV-ART compared to HIV uninfected people (Figure 2), Tregs and the

Th17/Treg ratio were similar in the two groups (Figure S2). Thus, increased IDO activity is

associated with a reduction in circulating Th17 cell subsets, but the frequency of IL17

producing CD4 T cells does not correlate with IDO activity.

Discussion

Studies that demonstrate Th17 cells contribute to the control of Mtb infection in humans [7–9]

used different approaches to identify the cells. We used a combination of cell markers [14,17–

21] to broadly characterize Th17 cells and then confirmed that they produce IL17, instead of

using IL17 production alone to define Th17 cells [5,9]. Since there are cytokine-independent

mechanisms of T cell responses, and since cytokine production can be influenced by

stimulation conditions, our approach affords a broader characterization of Th17 cells to

determine the impact of HIV coinfection. Indeed, a recent study in macaques showed

granulomas that controlled Mtb were enriched for T1T17 (Th1*) cells that did not express IL17

transcripts [40]. Furthermore, since Mtb-antigen specific cells must traffic to sites of disease,

using chemokine receptor expression alone in blood may underreport the population of these

cells. Our finding of disproportionate reduction of specific circulating Th17 cell categories in

HIV-ART reveals the importance of using multiple criteria to identify CD4 T cells of interest,

and that ART-mediated suppression of HIV does not reconstitute all Th17 cell types equally.

Studies to identify CD4 T cell defects that contribute to TB susceptibility in HIV and HIV-

ART will benefit from higher resolution characterization of CD4 T cell subsets.

Our finding of a higher magnitude of IL17- than IFNγ-producing CD4$^+$ T cells in HIV-ART

upon Mtb300 stimulation is contrary to the impression that IFNγ production is the dominant

response to Mtb infection [41]. This may have implications for TB diagnosis since current tests

for detection of responses to Mtb antigens rely on IFNγ production, whose sensitivity is

variable and is impacted by HIV infection [42]. Recent studies revealed evidence that people

who have been exposed to Mtb but remain negative by IFNγ responses have detectable

responses to Mtb antigens when other readouts are used [43]. Until recently, those individuals

were thought to remain uninfected by Mtb; our results add weight to the evolving concept that

considering IFNγ as the canonical responses to Mtb infection requires reconsideration.

Our findings indicate that IL17 responses to Mtb but not CMV antigens or SEB can be

preserved in HIV-ART. We hypothesize that IL17-producing cells may be less activated than

IFNγ-producing cells and/or express the CCR5 coreceptor at lower levels or at lower

frequencies and are thus preserved from HIV infection. The technical challenge of optimal

measurement of T cell activation and cytokine production in one condition (detection of

cytokines requires protein transport inhibitors which hinder T cell activation marker

expression) precluded us from testing this hypothesis.

The K/T ratio is a strong predictor of mortality risk in ART-suppressed PWH [44], does not

normalize in PWH initiating ART early, and durably suppressed early-ART initiators have high

plasma KT ratios compared with HIV-negative controls [29]. Therefore, the elevated

tryptophan catabolism in HIV-ART reported here and previously [25,29] may explain, in part,

the persistently increased risk of tuberculosis in treated HIV. Inhibition of IDO-1 with 1-

methyl-D-tryptophan (indoximod) in macaques was reported to decrease Mtb burdens and

pathology, and increased proliferation of CD4$^+$ and CD8$^+$ memory T cells [45] Together, these

results suggest that enhancing Th17 cell responses by IDO inhibition may be beneficial in the

context of TB.

Our study has limitations: 1) the participants were from a study investigating the long-term

clinical and immunological consequences of HIV infections and their treatment, therefore, we

had few HIV uninfected participants; 2) 10 of 13 PWH initially with a negative TST prior to

TST conversion had high viral loads that probably reduced the sensitivity of diagnosis; 3) INH

treatment was not given as directly observed therapy (DOT) and we could not ascertain

treatment compliance by the study participants. Nevertheless, we found that Th17 cells are a

heterogenous population and IL17 responses to Mtb are preserved in HIV-TB coinfected

individuals, and we demonstrated the link between IDO activity and circulating Mtb-

responsive Th17 cells in humans. With IDO-1 inhibitors in clinical trials [46], management of

TB disease might benefit from simultaneous therapeutic inhibition of IDO-1 activity to enhance

Th17 cell responses, particularly in PWH where HIV further enhances tryptophan catabolism

via the kynurenine pathway.

NOTES

Funding: This research was supported by a grant from the National Institutes of Health: UCSF-

Gladstone Center for AIDS Research (P30 AI027763).

Acknowledgments: We thank the study participants in the SCOPE study for providing

samples, UCSF Core Immunology Lab and UCSF-CFAR Specimen Processing & Banking

Subcore.

Conflict of interest: The authors have no conflict of interest to declare.

Presentations: This work was presented at the 1st Bay Area Tuberculosis Symposium

(BATS) September 9, 2022, San Francisco, CA.

Correspondence: Joel D. Ernst, M.D.

368   Division of Experimental Medicine

369   University of California, San Francisco

370   1001 Potrero Ave, Bldg. 3, Room 601 | San Francisco, CA 94110

371   joel.ernst@ucsf.edu

372   Peter W. Hunt, MD

373   Division of Experimental Medicine

374   Zuckerberg San Francisco General Hospital

375   1001 Potrero Avenue

376   Bldg 3, Room 605B, San Francisco, CA. 94110

377   peter.hunt@ucsf.edu

References

1. Global tuberculosis report 2021. https://www.who.int/publications-detail-redirect/9789240037021.

2. Kwan, C. K. & Ernst, J. D. HIV and tuberculosis: a deadly human syndemic. *Clin. Microbiol. Rev.* **24**, 351–376 (2011).

3. Lawn, S. D., Myer, L., Edwards, D., Bekker, L.-G. & Wood, R. Short-term and long-term risk of tuberculosis associated with CD4 cell recovery during antiretroviral therapy in South Africa. *AIDS* **23**, 1717–1725 (2009).

4. Gupta, A., Wood, R., Kaplan, R., Bekker, L.-G. & Lawn, S. D. Tuberculosis incidence rates during 8 years of follow-up of an antiretroviral treatment cohort in South Africa: comparison with rates in the community. *PLoS ONE* **7**, e34156 (2012).

5. Scriba, T. J. *et al.* Distinct, specific IL-17- and IL-22-producing CD4+ T cell subsets contribute to the human anti-mycobacterial immune response. *J. Immunol.* **180**, 1962–1970 (2008).

6. Geldmacher, C. *et al.* Preferential infection and depletion of Mycobacterium tuberculosis-specific CD4 T cells after HIV-1 infection. *J. Exp. Med.* **207**, 2869–2881 (2010).

7. Scriba, T. J. *et al.* Sequential inflammatory processes define human progression from M. tuberculosis infection to tuberculosis disease. *PLoS Pathog.* **13**, e1006687 (2017).

8. Nathan, A. *et al.* Multimodally profiling memory T cells from a tuberculosis cohort identifies cell state associations with demographics, environment and disease. *Nat Immunol* **22**, 781–793 (2021).

9. Ogongo, P. *et al.* Tissue-resident-like CD4+ T cells secreting IL-17 control Mycobacterium tuberculosis in the human lung. *J Clin Invest* **131**, 142014 (2021).

10.   Ma, W.-T., Yao, X.-T., Peng, Q. & Chen, D.-K. The protective and pathogenic roles of IL-

17 in viral infections: friend or foe? *Open Biol* **9**, 190109 (2019).

11.   Wacleche, V. S., Landay, A., Routy, J.-P. & Ancuta, P. The Th17 Lineage: From Barrier

Surfaces Homeostasis to Autoimmunity, Cancer, and HIV-1 Pathogenesis. *Viruses* **9**,

E303 (2017).

12.   Iwakura, Y., Ishigame, H., Saijo, S. & Nakae, S. Functional specialization of interleukin-

17 family members. *Immunity* **34**, 149–162 (2011).

13.   Shen, H. & Chen, Z. W. The crucial roles of Th17-related cytokines/signal pathways in

M. tuberculosis infection. *Cell Mol Immunol* **15**, 216–225 (2018).

14.   Acosta-Rodriguez, E. V. *et al.* Surface phenotype and antigenic specificity of human

interleukin 17-producing T helper memory cells. *Nat. Immunol.* **8**, 639–646 (2007).

15.   Singh, S. P., Zhang, H. H., Foley, J. F., Hedrick, M. N. & Farber, J. M. Human T cells that

are able to produce IL-17 express the chemokine receptor CCR6. *J Immunol* **180**, 214–

221 (2008).

16.   Annunziato, F. *et al.* Phenotypic and functional features of human Th17 cells. *J. Exp.*

Med. **204**, 1849–1861 (2007).

17.   Lindestam Arlehamn, C. S. *et al.* Memory T cells in latent Mycobacterium tuberculosis

infection are directed against three antigenic islands and largely contained in a

CXCR3+CCR6+ Th1 subset. *PLoS Pathog.* **9**, e1003130 (2013).

18.   Bengsch, B. *et al.* Human Th17 cells express high levels of enzymatically active

dipeptidylpeptidase IV (CD26). *J. Immunol.* **188**, 5438–5447 (2012).

19.   Zhao, X. *et al.* Involvement of CD26 in Differentiation and Functions of Th1 and Th17

Subpopulations of T Lymphocytes. *J Immunol Res* **2021**, 6671410 (2021).

20. Cosmi, L. *et al.* Human interleukin 17-producing cells originate from a CD161+CD4+ T

cell precursor. *J. Exp. Med.* **205**, 1903–1916 (2008).

21. Maggi, L. *et al.* CD161 is a marker of all human IL-17-producing T-cell subsets and is

induced by RORC. *European Journal of Immunology* **40**, 2174–2181 (2010).

22. Li, S. *et al.* Metabolic Phenotypes of Response to Vaccination in Humans. *Cell* **169**, 862-

877.e17 (2017).

23. Collins, J. M. *et al.* Tryptophan catabolism reflects disease activity in human

tuberculosis. *JCI Insight* (2020) doi:10.1172/jci.insight.137131.

24. Adu-Gyamfi, C. G. *et al.* Plasma Indoleamine 2, 3-Dioxygenase, a Biomarker for

Tuberculosis in Human Immunodeficiency Virus-Infected Patients. *Clin Infect Dis* **65**,

1356–1358 (2017).

25. Favre, D. *et al.* Tryptophan catabolism by indoleamine 2,3-dioxygenase 1 alters the

balance of TH17 to regulatory T cells in HIV disease. *Sci Transl Med* **2**, 32ra36 (2010).

26. Desvignes, L. & Ernst, J. D. Interferon-gamma-responsive nonhematopoietic cells

regulate the immune response to Mycobacterium tuberculosis. *Immunity* **31**, 974–985

(2009).

27. Weiner, J. *et al.* Metabolite changes in blood predict the onset of tuberculosis. *Nat*

Commun **9**, 5208 (2018).

28. Murray, M. F. Tryptophan depletion and HIV infection: a metabolic link to

pathogenesis. *Lancet Infect Dis* **3**, 644–652 (2003).

29. Schnittman, S. R. *et al.* Abnormal Levels of Some Biomarkers of Immune Activation

Despite Very Early Treatment of Human Immunodeficiency Virus. *J Infect Dis* **223**,

1621–1630 (2021).

30.   Lindestam Arlehamn, C. S. *et al.* A Quantitative Analysis of Complexity of Human

Pathogen-Specific CD4 T Cell Responses in Healthy M. tuberculosis Infected South

Africans. *PLoS Pathog* **12**, e1005760 (2016).

31.   Foreman, T. W. *et al.* CD4 T cells are rapidly depleted from tuberculosis granulomas

following acute SIV co-infection. *Cell Rep* **39**, 110896 (2022).

32.   Wong, E. B., Ndung'u, T. & Kasprowicz, V. O. The role of mucosal-associated invariant T

cells in infectious diseases. *Immunology* **150**, 45–54 (2017).

33.   Lu, B. *et al.* IL-17 production by tissue-resident MAIT cells is locally induced in children

with pneumonia. *Mucosal Immunology* **13**, 824–835 (2020).

34.   Ivanov, I. I. *et al.* The Orphan Nuclear Receptor RORγt Directs the Differentiation

Program of Proinflammatory IL-17+ T Helper Cells. *Cell* **126**, 1121–1133 (2006).

35.   Szabo, S. J. *et al.* A Novel Transcription Factor, T-bet, Directs Th1 Lineage Commitment.

Cell **100**, 655–669 (2000).

36.   Mpande, C. A. M. *et al.* Functional, Antigen-Specific Stem Cell Memory (TSCM) CD4+ T

Cells Are Induced by Human Mycobacterium tuberculosis Infection. *Front Immunol* **9**,

324 (2018).

37.   Wang, T. *et al.* Regulatory T cells in rheumatoid arthritis showed increased plasticity

toward Th17 but retained suppressive function in peripheral blood. *Ann Rheum Dis* **74**,

1293–1301 (2015).

38.   Boothby, I. C., Cohen, J. N. & Rosenblum, M. D. Regulatory T cells in skin injury: At the

crossroads of tolerance and tissue repair. *Sci Immunol* **5**, eaaz9631 (2020).

39.   Fontenot, J. D., Gavin, M. A. & Rudensky, A. Y. Foxp3 programs the development and

function of CD4+CD25+ regulatory T cells. *Nat Immunol* **4**, 330–336 (2003).

40.   Gideon, H. P. *et al.* Multimodal profiling of lung granulomas in macaques reveals

472   cellular correlates of tuberculosis control. *Immunity* **55**, 827-846.e10 (2022).

473   41.   Flynn, J. L. & Chan, J. Immune cell interactions in tuberculosis. *Cell* **185**, 4682–4702

474   (2022).

475   42.   Chen, H. *et al.* Diagnostic accuracy of the interferon-gamma release assay in acquired

476   immunodeficiency syndrome patients with suspected tuberculosis infection: a meta-

477   analysis. *Infection* **50**, 597–606 (2022).

478   43.   Lu, L. L. *et al.* IFN-γ-independent immune markers of Mycobacterium tuberculosis

479   exposure. *Nat. Med.* **25**, 977–987 (2019).

480   44.   Byakwaga, H. *et al.* The kynurenine pathway of tryptophan catabolism, CD4+ T-cell

481   recovery, and mortality among HIV-infected Ugandans initiating antiretroviral therapy.

482   *J Infect Dis* **210**, 383–391 (2014).

483   45.   Gautam, U. S. *et al.* In vivo inhibition of tryptophan catabolism reorganizes the

484   tuberculoma and augments immune-mediated control of Mycobacterium tuberculosis.

485   *Proc. Natl. Acad. Sci. U.S.A.* **115**, E62–E71 (2018).

486   46.   Tang, K., Wu, Y.-H., Song, Y. & Yu, B. Indoleamine 2,3-dioxygenase 1 (IDO1) inhibitors in

487   clinical trials for cancer immunotherapy. *J Hematol Oncol* **14**, 68 (2021).

Table 1: Cohort 1 (Characteristics of participants with and without HIV with Latent Tuberculosis Infection)

Characteristic	HIV negative (n =7)	HIV positive (n = 21)	P
Age, median (IQR), y	58 (44 -61)	47 (38 - 58)	0.5395
Sex, Male (%)	7 (100)	21 (100)	
Race/Ethnicity (n, (%))			
Black	6 (86)	8 (38)	
Asian	1 (14)	2 (10)	
White	0	7 (30)	
Middle East	0	1 (5)	
Multiracial	0	2 (10)	
Pacific Islander	0	1 (5)	
Combined antiretroviral therapy (Yes/No)	No	Yes	
CD4 Counts, median (IQR), cells/mm3	1297 (707 -1349)	640 (487 - 906)	0.0082
Plasma viral load (copies/mL)	None	< 40	

Table 2: Cohort 2 (Characteristics of participants with HIV before and after LTBI diagnosis and treatment)

Characteristic	TST neg	TST+ INH initiation	TST+INH completion	P
Age, median (IQR), y	45 (42 - 51)	_	_	
Sex, Male (%)	10 (71)	_	_	
Race/Ethnicity (n, (%))				
Black	5 (36)	_	_	
Asian	1 (7)	_	_	
White	7 (50)	_	_	
Pacific Islander	1 (7)	_	_	
CD4 Counts, median (IQR), cells/mm3	527 (230 - 781)	442 (367 - 588)	464 (361 - 724)	0.9172
HIV infection (n, %)	13 (100)	11 (100)	13 (100)	
Detectable viral load (n; median (IQR), copies/mL)	10; 3455 (614 - 24888)	6; 4895 (497 - 35920)	6; 2980 (170 - 59542)	0.9632
Suppressed viral load (n)	3	5	7	
Combined antiretroviral therapy				
On treatment (n)	7	6	10	
Not on treatment (n)	6	5	3	

Figure 1:

A.

Mtb300: Gated on CD4$^+$Vα7.2$^-$ cells

B.

p = 0.0002
r = 0.4472

■ TST$^+$ HIV$^+$
● TST$^+$ HIV$^-$
● Prior to TST$^+$
● Start of INH
● End of INH

C.

Teal = CD4$^+$
Blue = CD8$^+$
Black = Other CD3$^+$

Orange = T$_H$17
Cyan = T17
Black = Other CD3$^+$

Green = RORγT
Pink = T-bet
Black = Other CD3$^+$

Red = CD4$^+$IL17$^+$
Yellow = CD4$^+$IFNγ^+
Black = Other CD3$^+$

Figure 1. Different definitions of T helper 17 cells reveal high but incomplete concordance. (A) Representative flow cytometry plots and gating schemes for definition of T helper 17 cells by published marker criteria independent of cytokine expression. Left panel, chemokine receptor CCR6 and CXCR3 expression used to distinguish T17, T1T17, and Th1 non-MAIT CD4 T cells. Right panel: Use of surface expression of CD26 and CD161 to identify T$_H$17 non-MAIT CD4 T cells. Cells in both panels were stimulated with the MTB300 antigenic peptide pool. **(B)** High correlation of T$_H$17 (CD26$^+$CD161$^+$) and T17 (CCR6$^+$CXCR3$^-$) non-MAIT CD4 T cells in cell populations from participants with varying TB and HIV status. Spearman correlation p and r values are shown for results pooled from all participants. **(C)** t-stochastic neighbor embedding (t-SNE) analysis reveals incomplete concordance of transcription factor (RORγt and T-bet) and cytokine (IFNγ and IL17) production and surface phenotypes of non-MAIT CD4$^+$ T cells. In this analysis, protein transport inhibitors (golgi stop and golgi plug) were not added to PBMC during antigen stimulation thus weak cytokine signals which are best shown in contour format.

Figure 2:

Figure 2. ART treated HIV infection is associated with disproportionate depletion of distinct subsets of T helper cells. (A) Reduced frequency of bulk CD4 T cells in HIV+ vs HIV- participants in Cohort 1. **(B)** ART-treated HIV is not associated with differential distribution of CD4 T cells in memory T cell subsets. Naive: CD45RA+CCR7+; Central Memory (CM): CD45RA-CCR7+); Effector Memory: CD45RA-CCR7-; TEMRA: CD45RA+CCR7-). **(C)** Disproportionate reduction in frequencies of T_H17 and T1T17 but not T17 or Th1 cells in ART-treated HIV. T_H17=CD4+Vα7.2-CD26+CD161+; T1T17 (also termed Th1*): CD4+Vα7.2-CCR6+CXCR3+; T17: CD4+Vα7.2-CCR6+CXCR3-; and Th1: CD4+Vα7.2-CCR6-CXCR3+. Statistical comparisons were by Mann-Whitney test.

492
493

Figure 3:

Figure 3. **In contrast to responses to other stimuli, *Mtb* antigen responsive IL17 expression is enriched in CD26⁺CD161⁺ T_H17 rather than CCR6⁺CXCR3⁻ T17 cells. (A)** Left panel (Cohort 1): CD4 T cells that produce IL17 in response to MTB300 are predominantly in the T_H17 subset. Right panel (Cohort 2): MTB300-stimulated IL17 production predominates in T_H17 rather than T17 cells regardless of TST status and is not altered by INH preventive therapy. **(B)** Left panel (Cohort 1): In contrast to MTB300-responsive cells, IL17 production predominates in T17 cells in response to SEB activation. Right panel (Cohort 2): INH preventive therapy does not alter the pattern of IL17 production by SEB-stimulated T17 or T_H17 cells. **(C)** Left panel (Cohort 1): In ART-treated HIV⁺ participants, CD4 T cells that produce IFNγ in response to MTB300 stimulation are present at low frequencies. Right panel (Cohort 2): **(D)** Left panel (Cohort 1): CMV pp56 antigen stimulation induces higher frequencies of IFNγ than IL17-producing CD4 T cells. Right panel (Cohort 2): INH treatment does not alter CD4 T cell responses to CMV pp65 antigen stimulation.

Figure 4:

Figure 4. In TST[+] participants, HIV infection is associated with increased indoleamine-2,3-dioxygenase (IDO) activity, which negatively correlates with TH17 and T17 cell frequencies. (A) Plasma tryptophan concentrations; (B) Plasma kynurenine concentrations; (C) Plasma Kynurenine (K)/Tryptophan (T) ratios as a reflection of IDO activity. (D) Left panel: correlation of TH17 cell frequencies with plasma K/T ratios; Middle panel: Correlation of regulatory T cells and plasma K/T ratios; Right panel: Correlation of TH17/Treg ratios and plasma K/T ratios. (E) Left panel: correlation of T17 cell frequencies with plasma K/T ratio; Right panel: Correlation of T17/Treg ratios and plasma K/T ratios. Statistical analyses were by Mann-Whitney (panels A-C) or Spearman correlation (D and E).

496
497

26