Genomes of the human filarial parasites *Mansonella perstans* and

Mansonella ozzardi

Amit Sinha[1], Zhiru Li[1], Catherine B. Poole[1], Richard D. Morgan[1], Laurence Ettwiller[1],
Nathália F. Lima[2], Marcelo U. Ferreira[2,3], Fanny F. Fombad[4], Samuel Wanji[4], Clotilde K. S.
Carlow[1*]

[1]New England Biolabs, Ipswich Massachusetts, 01938 USA;

[2]Department of Parasitology, Institute of Biomedical Sciences, University of São Paulo, São Paulo,
Brazil;

[3]Global Health and Tropical Medicine, Institute of Hygiene and Tropical Medicine, NOVA
University of Lisbon, Lisbon, Portugal;

[4]Department of Microbiology and Parasitology, University of Buea, Buea, Cameroon

* Correspondence: Clotilde Carlow, Division of Genome Biology, New England Biolabs, Ipswich,
MA 01938. Tel: 978 380 7263, Fax: 978 921 1350, e-mail: carlow@neb.com.
14

15 **Keywords:** *Mansonella*, **Filaria, Neglected Tropical Disease, Genomics, Evolution.**

Abstract

The filarial parasites *Mansonella ozzardi* and *Mansonella perstans*, causative agents of

mansonellosis, infect hundreds of millions of people worldwide, yet remain among the most

understudied of the human filarial pathogens. *M. ozzardi* is highly prevalent in Latin American

countries and Caribbean Islands, while *M. perstans* is predominantly found in sub-Saharan Africa as

well as in a few areas in South America. In addition to the differences in their geographical

distribution, the two parasites are transmitted by different insect vectors, as well as exhibit

differences in their responses to commonly used anthelminthic drugs. The lack of genome

information has hindered investigations into the biology and evolution of *Mansonella* parasites and

understanding the molecular basis of the clinical differences between species. In the current study,

high quality genomes of two independent clinical isolates of *M. perstans* from Cameroon and two *M.*

ozzardi isolates one from Brazil and one from Venezuela are reported. The genomes are

approximately 76 Mb in size, encode about 10,000 genes each, and are largely complete based on

BUSCO scores of about 90%, similar to other completed filarial genomes. These sequences represent

the first genomes from *Mansonella* parasites and enabled a comparative genomic analysis of the

similarities and differences between *Mansonella* and other filarial parasites. Horizontal DNA

transfers (HDT) from mitochondria (nuMTs) as well as transfers from genomes of endosymbiotic

Wolbachia bacteria (nuWTs) to the host nuclear genome were identified and analyzed. Sequence

comparisons and phylogenetic analysis of known targets of anti-filarial drugs diethylcarbamazine

(DEC), ivermectin and mebendazole revealed that *M. ozzardi* lacks an ortholog of *gon-2*, a target for

DEC, while *M. perstans* lacks an ortholog of *avr-14*, a target of ivermectin. These new reference

genome sequences will provide a valuable resource for further studies on biology, symbiosis,

evolution and drug discovery.

1   Introduction

A large part of the world's population is infected with one or more filarial nematode parasites

resulting in several distinct diseases often referred to as a group under the general term filariasis. The

filarial parasites *Mansonella ozzardi* and *Mansonella perstans* are the two main causes of

mansonellosis in humans. Infections are transmitted by infected biting midges (mostly of the genus

Culicoides) or blackflies (genus *Simulium*) that release infective stage larvae during a blood meal.

Adult male and female worms have been observed rarely in humans since they are presumed to live

in the serous body cavities and/or subcutaneous tissues. Viviparous females release unsheathed

microfilaria which circulate in the blood, available for ingestion by an appropriate vector during

feeding. Unlike some of the other human filarial diseases, *Mansonella* infections do not present with

a distinct pathology or specific clinical picture. Infected individuals are often asymptomatic or may

have a variety of clinical features including subcutaneous swellings, aches, pain, fever, headache,

pruritis, corneal lesions and eosinophilia (Simonsen et al., 2011; Lima et al., 2016; Mediannikov &

Ranque 2018; Vianna et al., 2012; Ferreira et al 2021). Other manifestations include

immunosuppression (Ritter et al., 2018) with important ramifications such as increased susceptibility

to other infections and reduced efficacy of vaccine programs (Ta-Tang et al., 2018).

Even though mansonellosis ranks first in prevalence among the human filariases, affecting hundreds

of millions of people in Africa and Central and South America, it remains one of the least studied and

most neglected of all filarial diseases (Downes & Jacobsen, 2010; Simonsen et al., 2011; Lima et al.,

2016; Mediannikov & Ranque, 2018; Ta-Tang et al., 2018; Ta-Tang et al., 2021; Sandri et al., 2021;

Bélard & Gehringer 2021). Indeed, much of the somewhat limited information available has been

obtained serendipitously during studies on other filarial parasites or malaria, in which individuals

were co-infected with *Mansonella* parasites. *M. perstans* is considered the most common filariasis in

Central Africa with infection prevalence often very high (up to 96%) in endemic areas, even among

children (Downes and Jacobsen, 2010; Simonsen et al., 2011; TaTang et al., 2018). It is also found in

northern regions of the Amazon rainforest and the Caribbean coast in South America (TaTang et al.,

2018; Bélard and Gehringer, 2021), with molecular evidence supporting introduction of *M. perstans*

to the American continent from Africa because of the slave trade (Tavares da Silva et al., 2017). *M.*

ozzardi infections have been reported from many countries in South America, Central America, and

some Caribbean Islands (Lima et al., 2018; TaTang et al.,2018; Tavares de Silva et al., 2017; Raccurt

et al., 2018; Ferriera et al., 2021). Co-infections of *M. perstans* and *M. ozzardi*, together with other

filarial parasites, have also been reported in South America (Kozek et al., 1983; Calvopina et al.,

2019), making their treatment challenging as the two species respond differently to the commonly

used anti-filarial drugs such as ivermectin, diethylcarbamazine (DEC) or mebendazole (Chadee et al.,

1995; TaTang et al., 2018). The recent development of highly specific, simple molecular tests which

can distinguish *M. perstans* and *M. ozzardi*, and do not cross-react with other filarial parasites (Poole

et al., 2019), will facilitate an effective treatment regimen. The antibiotic doxycycline, targeting the

endosymbiotic *Wolbachia* bacteria of *M. perstans* and *M. ozzardi*, has been shown to clear the

microfilarial stage from the blood (Coulibaly et al., 2009; Debrah et al., 2019; Crainey et al., 2020)

highlighting the importance of *Wolbachia* in these species. The *Wolbachia* genomes, *w*Mpe from *M.*

perstans, and *w*Moz from *M. ozzardi* have recently been published (Sinha et al., 2022), providing a

valuable resource for the discovery of new antibiotics and studies on the symbiotic relationship and

evolution.

Here we provide the first *de novo* assemblies of the nuclear genomes from *M. perstans* and *M.*

ozzardi using combined PacBio long-read and Illumina paired-end sequencing data. We used

customized metagenomic assembly and binning pipelines, and extensive quality checks to ensure

accurate assemblies from a mixture of DNA from the *Mansonella* parasites, their human host, and the

associated microbiome. These methods were applied to generate genomes from two clinical isolates

of each species, providing a unique opportunity for intra-species and inter-species comparisons and

exploring features of sequence evolution of these parasites. We analyzed the genomic regions

containing horizontally transferred DNA from mitochondria (nuMTs), as well horizontally

transferred DNA from their endosymbiont *Wolbachia* (nuWTs). Phylogenetic analysis of orthologs

of genes that encode drug targets of common anti-filarial drugs such as DEC, ivermectin and

mebendazole revealed intriguing differences in the presence and absence of these targets between *M.*

perstans and *M. ozzardi* that might have clinical implications for the treatment of mansonellosis.

2   Methods

2.1   Ethics statement

All research involving human subjects was approved by the appropriate committee and performed in

accordance with all relevant guidelines and regulations. Informed consent was obtained from all

participants or their legal guardians. These samples have also been described and utilized in previous

studies (Poole et al., 2019; Sinha et al., 2022).

Ethical clearance for *M. perstans* samples was obtained from the National Institutional Review

board, Yaoundé (REF: N° 2015/09/639/CE/CNERSH/SP) and administrative clearance from the

Delegation of Public Health, South-West region of Cameroon (Re: R11/MINSANTE/SWR/

RDPH/PS/259/382). Approval for the study was granted by the "National Ethics Committee of

Research for Human Health" in Cameroon. Special consideration was taken to minimize the health

risks to which any participant in this study was exposed. The objectives of the study were explained

to the consenting donor after which they signed an informed consent form. The participant's

documents were given a code to protect the privacy of the study subject. At the end of the study, the

donor received a cure of mebendazole (100 mg twice a day for 30 days).

The study protocols for *M. ozzardi* were approved by the Institutional Review Board of the Institute

of Biomedical Sciences, University of São Paulo, Brazil (1133/CEP, 2013) as part of a previous

study (Lima et al., 2018). Written informed consent was obtained from all patients, or their parents or

guardians if participants were minors younger than 18 years. Diagnosed infections were treated with

a single dose of 0.2 mg/kg of ivermectin after blood sampling (Lima et al., 2018).

2.2   Illumina library construction and sequencing

The parasite materials, DNA extractions, Illumina library preparation and sequencing have been

described in detail previously (Poole et al., 2019; Sinha et al., 2022). Briefly, two clinical isolates of

M. perstans, named Mpe-Cam-1 and Mpe-Cam-2, were obtained as microfilariae in a blood draw

from consenting individuals in Cameroon. One clinical isolate of *M. ozzardi*, named Moz-Brazil-1,

was obtained from a consenting individual as part of a previous study in Brazil, while another isolate

from Venezuela, named Moz-Venz-1, was generously donated by Izaskun Petralanda in 1989. The

DNA samples from Mpe-Cam-1, Mpe-Cam-2, and Moz-Brazil-1 were treated with the NEBNext

Microbiome DNA enrichment to reduce human DNA contamination before being used for library

preparation. Illumina libraries for all 4 isolates were constructed using the NEBNext Ultra II FS

DNA Library Prep Kit (New England Biolabs Inc., USA) as per standard protocol and sequenced

using the Illumina MiSeq and NextSeq500 platforms (paired end, 150 bps). Bioinformatic analysis

programs were run with their default parameters unless otherwise stated.

2.3   PacBio library sequencing and *de novo* assembly

A total of 250 ng of genomic DNA from *M. perstans* isolate Mpe-Cam-1 was used for sequencing on

PacBio RSII platform as per manufacturer's instructions. Consensus reads from raw PacBio reads

were obtained using the protocol RS_PreAssembler.1 (parameters minLen=1000, minQual=0.80,

genomeSize=100000000) on PacBio SMRT Portal (smrtanalysis_2.3.0.140936.p5.167094). Reads

originating from human genomic DNA were removed by first by mapping the consensus reads to the

human genome (grch38) using minimap2 v2.17-r941 (Li, 2018) followed by discarding all mapped

reads (MAPQ score > 20). The remaining reads were assembled using canu v2.2 (Koren et al., 2017)

(parameters genomeSize=90m correctedErrorRate=0.045). The assembly was iteratively polished

over 12 rounds using "Resequencing" protocol on PacBio SMRT Portal (v 6.0.0.47841) (parameters -

-minMatch 12 --bestn 10 --minPctSimilarity 70.0 --refineConcordantAlignments) till no new variants

could be detected by the Resequencing pipeline. The assembly contiguity was further improved by

using finisherSC software v2.1 (Lam et al., 2015), a scaffolder that uses raw PacBio reads while

accounting for any genomic repeats in the assembly. The assembly was polished using the Illumina

data for the same isolate via the polca software (Zimin and Salzberg, 2020). Finally, gap-filling and

heterozygosity removal was performed on this assembly using Illumina reads as an input to the

Redundans pipeline v0.14a (Pryszcz and Gabaldón, 2016). The assembly quality was evaluated using

quast (Gurevich et al., 2013) at each step.

2.4   Metagenomic assembly using Illumina reads

Illumina raw reads were processed using the BBTools package (https://jgi.doe.gov/data-and-

tools/bbtools/). Duplicate raw reads and bad tiles were removed using the clumpify.sh and

filterbytile.sh utilities. Adapter trimming, removal of phiX reads, and reads with 13-nt or longer

stretches of the same nucleotide (poly-G, poly-C, poly-A, poly-T) was performed using bbduk.sh.

Human host reads were removed by mapping against the human genome (grch38) using bbmap.sh

utility. The quality metrics of the processed reads at each step were assessed using FastqC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Reads from different runs of the

libraries prepared from the same genomic DNA sample were combined and used as an input for the

assembly of the metagenome using metaSpades (Nurk et al., 2017).

2.5   Metagenomic binning to identify *Mansonella* contigs

Binning of metagenomic contigs to identify sequences originating from *Mansonella* genomes was

carried out based on the BlobTools software (Laetsch and Blaxter, 2017) and additional curation, as

described previously (Sinha et al., 2022). Briefly, for each of the isolates Mpe-Cam-2, Moz-Brazil-1

and Moz-Venz-1, the Illumina reads were mapped back to the metagenomic contigs produced by

metaSpades using bowtie2 (Langmead and Salzberg 2012). For the PacBio assembly obtained from

the Mpe-Cam-1 sample, the corresponding Illumina reads were mapped to this assembly using

bowtie2. Sequence similarities of assembled contigs to the NCBI nt database and Uniport database

was carried out using blastn and diamond in BlobTools package. The bins annotated as "Nematoda"

and "Proteobacteria" by BlobTools were analyzed further to retrieve sequences originating from

Mansonella. Contigs that were marked as Proteobacteria but co-localized with the Nematoda blobs in

the blobplots, were included in the *Mansonella* bins as they represented potential examples of

horizontal gene transfer from *Wolbachia* to *Mansonella* host. These HGT candidates were validated

by inspection of blastn and blastx hits of these contigs to the NCBI nt and nr databases respectively.

Contigs originating from mitochondrial DNA were identified by analysis of nucmer alignments

between assembled sequences and the previously reported mitochondrial genomes, GenBank

accessions MT361687.1 and KX822021.1 for *M. perstans* (Chung et al., 2020) and *M. ozzardi*

(Crainey et al., 2018) respectively. The contigs with more than 98% sequence identity to published

mitochondrial genomes were marked as mtDNA and removed from the nuclear assembly.

2.6   Assembly refinements and heterozygosity removal for Illumina assemblies

For each Illumina-only assembly, scaffolding, gap-filling and heterozygosity removal was performed

using the corresponding Illumina reads in the Redundans pipeline v0.14a (Pryszcz and Gabaldón,

2016).

2.7   Genome annotations and comparative analysis

Identification of repetitive elements in the assembled genomes were performed using RepeatModeler

version 2.0.3 (Flynn et al., 2020) and classified against the Dfam database (Storer et al., 2021). First

round of repeat annotations was performed on a fasta file combining all assembled genomes. The

repeat sequence families identified from this run were used as a reference library in RepeatMasker

4.1.2-p1 (http://repeatmasker.org/) to mask repeats in each of the 4 genome assemblies

independently. The outputs from RepeatMasker were processed with parseRM.pl utility (Kapusta et

al. 2017) to generate summary statistics on repats classified as DNA transposons and retroelements.

Gene predictions were carried out for each assembly using the braker2 pipeline v2.2.6 (Bruna et al.,

2020), with a reference proteome comprised of all metazoan proteins in OrthoDB version 10.1

(Kriventseva et al., 2019) and all filarial proteomes downloaded from WormBase ParaSite (Howe et

al., 2017). The completeness of the protein-coding content of genomes was assessed using the

BUSCO pipeline v5.0 beta using the "nematoda_odb10" reference dataset (Simão et al., 2015).

OrthoFinder v2.4 (Emms and Kelly, 2019) was used to infer orthologs of protein coding genes across

multiple nematode genomes. The set of proteomes included all 16 filarial species with publicly

available genomes. The species *Setaria digitata* and *Thelazia callipaeda* from the same infraorder

Spiruromorpha were included as the immediate outgroups, while *Caenorhabditis elegans* and

Pristionchus pacificus were included as relatively more distant outgroups. The protein and transcript

sequences were downloaded from WormBase parasite where available. The recently reported

genome sequences for filarial nematodes *Cruorifilaria tuberocauda*, *Dipetalonema caudispina*,

Dirofilaria immitis, *Litomosoides brasiliensis* and *Madathamugadia hiepei* (Lefoulon et al., 2020)

were available from NCBI. However, the corresponding gene annotations for these assemblies were

not available in NCBI. Hence, repeat-modeling, repeat-masking and gene annotation using braker2

pipeline was performed on each of these species as described for *M. perstans* and *M. ozzardi* above.

Whole-genome alignments for global sequence similarity and synteny between various repeat-

masked genomes were performed using minimap2 v2.17-r941 (Li, 2018) and visualized using the

JupiterPlot tool (https://github.com/JustinChu/JupiterPlot). For global sequence comparisons across

multiple filarial genomes, the average nucleotide identity (ANI) scores were calculated using the

OrthoANIu tool (Yoon et al., 2017).

Comparison and analysis of sequence differences, including structural difference, between different

pairs of genomes, namely (A) Mpe-Cam-1versus Mpe-Cam-2 (B) Moz-Brazil-1 versus Moz-Venz-1

(C) Mpe-Cam-1 versus Moz-Brazil-1 were performed using nucdiff version 2.0.3 (Khelik et al.,

2017). The localization of the sequence variants within gene bodies and exons was calculated through

the intersectBed tool in the bedtools package v2.29.2 (Quinlan and Hall, 2010). The effect of

sequence variants on the coding potential of affected genes was determined using snpEff version 5.1d

(Cingolani et al., 2012).

2.8   Phylogenetic analysis

For genome-wide phylogenetic analysis, single copy orthologs across various nematodes were

identified from the OrthoFinder analysis described above. The corresponding protein sequences were

aligned using mafft v7.149b (Nakamura et al., 2018) and trimmed using Gblocks v0.91b (Talavera

and Castresana, 2007). Phylogenetic analysis of the resulting supermatrix of 823,832 amino acids

was carried out using iqTree v2.1.2 (Nguyen et al., 2015). The best fit substitution model was chosen

automatically using ModelFinder implementation within iqTree (Kalyaanamoorthy et al., 2017).

Bootstrap support values were calculated based on (i) the SH-like approximate likelihood ratio test

(Guindon et al., 2010), and (ii) ultrafast bootstrap (Hoang et al., 2018) with 1000 replicates each

(iqTree command-line options '-B 1000 -alrt 1000 -abayes -lbp 1000').

For phylogenetic analysis of specific genes and gene families, their homologs and the corresponding

protein sequences were obtained from the OrthoFinder output. The protein domains for all the

sequences were annotated using NCBI CD search and the NCBI CDD database (Lu et al., 2020). The

protein sequences which did not have the expected domain structures as compared to the *C. elegans*

homologs or were too short in length were excluded from the final analysis. The multiple sequence

alignments and tree inference was carried out using mafft and iqtree as described above. The

consensus trees were edited and annotated using the iTOL webserver (Letunic and Bork, 2021) and

Adobe Illustrator.

2.9   Identifying nuWTs and nuMTs

The assembled nuclear genomes were aligned to the respective *Wolbachia w*Mpe or *w*Moz (Sinha et

al., 2022) genomes using nucmer v4.0.0beta2 (Marçais et al., 2018). Aligned fragments ("MuM"s)

that were separated by less than 50 bp, were combined into a single nuWT locus. For manual

validation of HGT of *Wolbachia* sequences into filarial contigs, the *Wolbachia* regions were verified

to be flanked by nematode-like sequences and protein coding genes from nematodes.

A similar strategy was employed for nuMT identification, with nucmer alignments against respective

mitochondrial sequences, (Chung et al., 2020, Crainey et al., 2018), and blastx against nematode

nuclear proteomes. The nuMT loci with more than 85% AT-content were removed to avoid spurious

matches to low-complexity regions.

Comparison of nuMT or nuWT loci for conservation of their sequence and position across different

genomes was based on their sequence comparisons using megablast coupled with the synteny

analysis described above. The sequences for nuMT or nuWT loci were extracted for each genome

and compared using megablast, and only hits with more than 90% identity over 90% of the sequences

were analyzed further as candidates for conserved loci. In the next step, if these candidate loci were

present on syntenic contigs, they were marked as conserved loci across strains or species. A blastx

analysis of the nuWT or nuMT sequences against the respective mitochondrial or *Wolbachia*

proteomes was performed to identify any intact protein coding genes.

3   Results

3.1   PacBio sequencing and *de novo* assembly generates a highly contiguous *M. perstans* genome

259
260

PacBio sequencing and *de novo* genome assembly of genomic DNA isolated from *M. perstans* isolate

"Mpe-Cam-1", together with BlobTools analysis for metagenomic binning resulted in a high-quality

M. perstans genome. The genome assembly is comprised of 581 scaffolds with a total size of 76 Mb,

with the largest scaffold greater than 1.2 Mb and a N50 size greater than 285 kb (Table 1). Based on

these quality metrics, the Mpe-Cam-1 assembly ranks highly in comparison to all filarial genomes

available so far (Supplementary Material Table 1). Repetitive elements account for 9.7% of the

genome, of which about 27% of the repeat sequence length was comprised of low complexity and

simple repeats. Long terminal repeat (LTR) retrotransposons comprised 17% of all the repeats, while

50% of the repeat content is comprised of unknown, novel repeat families (Supplementary Material

Table 2). The genome has 9,460 protein coding genes, with a BUSCO score of 93.2 %

(Supplementary Figure 1) when evaluated against a set of 3,131 core proteins conserved across the

nematode phylum, further indicating the high quality of this genome assembly.

Another clinical isolate from Cameroon, "Mpe-Cam-2" was sequenced and assembled *de novo* based

on Illumina sequencing. Although the quality metrics of this Illumina short-reads assembly are lower

than the PacBio long-read Mpe-Cam-1 assembly (Table 1), the repeat content, (Supplementary

Material Table 2), the number of predicted genes and their BUSCO scores are similar to Mpe-Cam-1

(Supplementary Figure 1).

3.2   Genome sequencing and assembly of *M. ozzardi* from South America

280 For *M. ozzardi*, two genome assemblies from Latin American isolates, "Moz-Brazil-1" from Brazil

281 and "Moz-Venz-1" from Venezuela, were generated based on Illumina sequencing reads. These

282 assembled genomes are about 76 Mb (Table 1), similar in size to the *M. perstans* genomes described

283 above. The number of predicted genes, their BUSCO scores (Supplementary Figure 1) and repeat

284 content (Supplementary Material Table 2) are also similar to those present in *M. perstans* genomes.

3.3   Analysis of genome variation across *M. perstans* and *M. ozzardi* and among multiple

     clinical isolates

The availability of genomes from two *Mansonella* species and multiple isolates provided a unique

opportunity for analysis of whole genome sequence variation both between and within the two

species. The genome-wide average nucleotide identity (gANI) scores for isolates within species are

99.5% (Table 2) in comparison to the score of 89.69% between the two *Mansonella* species.

Visualizations of whole genome alignments between isolates as Jupiter plots (Figure 1) reveal high

levels of synteny. The isolates Mpe-Cam-1and Mpe-Cam-2, collected from the same geographical

region within a few years apart are most similar to each other (Figure 1A). Relatively more genomic

re-arrangements (represented as crisscrossing lines in Jupiter plots) can be seen between *M. ozzardi*

isolates Moz-Brazil-1 and Moz-Venz-1 (Figure 1B), which have been collected from different

geographical regions (Brazil, Venezuela) and at about 30 years apart. This indicates that the

evolution of *Mansonella* genomes is dependent on geographical and temporal separation. Whole-

genome alignments comparing *M. perstans* and *M. ozzardi* genomes also demonstrate high levels of

synteny (Figure 1C), although lower than the within-species levels, consistent with the trend in their

gANI scores.

At the nucleotide level, the numbers of simple substitutions, insertions and deletions (Table 2) were

10 to 30-fold higher in across-species comparisons (6.22%), as compared to within-species (0.2%

and 0.29%). In all the pairwise comparisons, most of the sequence variation (>52.8%) was localized

to the intergenic portions of the genomes outside the protein-coding genes CDS regions. The

proportion of variants within protein-coding genes ranged from about 5% to 10% (Table 2). Most of

the nucleotide changes (54% to 60%) that occurred within coding regions, were mutationally silent as

they resulted in synonymous amino-acid changes, 39 to 44% of coding variants did result in missense

or non-synonymous changes in the predicted protein sequences and only a small fraction (0.21 % to

0.77%) of the coding variants resulted in a nonsense mutation. Overall, the intra-species differences

between the two *M. ozzardi* isolates from different geographical areas were higher than the

corresponding intra-species differences of two *M. perstans* isolates which were both collected in

Cameroon. A similar trend was observed for structural differences as well, as the number of

relocations, reshufflings and inversions of contig fragments was observed to be much higher in cross-

species comparisons (Table 2).

3.4   Orthology and phylogenomic analysis of filarial nematodes

An Orthofinder analysis of proteomes from all available filarial genomes, with *S. digitata* and *T.*

callipaeda as immediate outgroups from the sub-order Spiruromorpha, and *C. elegans* and *P.*

pacificus as distant outgroups identified 3,755 orthogroups conserved across all these nematodes

(Figure 2). This analysis also highlighted 91 orthogroups specific to *Mansonella* genus, 317

orthogroups specific to *Onchocerca* genus, and 130 orthogroups specific to *Brugia*. Most of these

proteins had only weak similarities to other nematode proteins (median sequence identity and

coverage both around 33%), making them good candidates for genus-specific biomarkers

(Supplementary Material Table 3).

Among the orthogroups conserved across all nematode genomes analyzed here, 465 orthogroups

comprised of single copy orthogroups (SCOs) and were used for analysis. The phylogenomic tree

(Figure 3) generated using these 465 orthogroups recapitulated the phylogenetic relationships based

on analysis of mitochondrial genomes (McCann et al., 2021) and the monophyletic feature of major

clades ONC2, ONC3, ONC4 and ONC5 based on 7 genes (Lefoulon et al., 2015).

3.5   Analysis of nuMTs, horizontal transfers of mitochondrial DNA into nuclear genomes

Transfer of mitochondrial DNA fragments into the nuclear genome has been observed for multiple

filarial nematodes (Grote et al., 2017) as well other eukaryotes (Hazkani-Covo et al., 2010). Since

mitochondrial markers are often used as barcodes for species identification and phylogenetic analysis

(Crainey et al., 2018), it is important to distinguish between genuine mitochondrial sequences versus

nuMTs (Song et al., 2008, Crainey et al., 2018). In the newly sequenced *Mansonella* genomes, 39 to

94 nuMT loci with median sizes ranging around 250 bp were found across different isolates, with the

largest loci ranging from 2.35 kb to 3.69 kb in size (Table 3). None of the nuMTs was found to carry

an intact protein coding gene. The total number and the combined size of nuMTs in was observed to

be higher in *M. ozzardi* as compared to *M. perstans* (Table 3). The locations and sequences of nuMTs

were poorly conserved across *M. perstans* and *M. ozzardi*. For example, in a comparison of the 48

Mpe-Cam-1 nuMTs with the 82 Moz-Brazil-1 nuMTs, only 2 conserved loci were found

(Supplementary Material Table 4). Even within the isolates of each species, approximately 35% and

60% of nuMT loci were conserved in *M. perstans* and *M. ozzardi* respectively (Supplementary

Material Table 4), while the remainder were isolate-specific, indicating a fast turnover of these

elements. The existence of multiple cox1 derived nuMT pseudogenes has been reported in *M. ozzardi*

(Crainey et al., 2018). Similarly, multiple fragments of the *cox1* gene were also observed in both *M.*

ozzardi isolates (Supplementary Figure 2). However, an exact match for these sequences could not be

found in either of the Moz-Brazil-1 or Moz-Venz-1 assemblies, providing further evidence for poor

conservation of such nuMTs across different isolates.

When the nuMT sequences from each isolate were mapped back to the mitochondrial genomes, they

were found to cover almost the entire mitochondrial genome, without any preference for specific

regions to be transferred into the nuclear genome (Supplementary Figures 2 and 3).

3.6   Analysis of nuWTs, horizontal transfers of *Wolbachia* endosymbiont DNA into host

nuclear genomes

Instances of transfer of DNA sequences from *Wolbachia* endosymbionts into the nuclear genomes of

their hosts, termed "nuWTs" for nuclear-*Wolbachia* transfers, have been observed frequently in

filarial nematodes (McNulty et al., 2010; Cotton et al., 2016; Grote et al., 2017; Hotopp et al., 2017)

as well as arthropod hosts (Klasson et al., 2014; Tvedte et al., 2022). The *Mansonella* isolates

reported here harbor about 300 nuWT loci in each isolate, with median size about 180 bp, adding up

to a total of 76 kb to 88 kb), with the largest nuWT (2.73 kb) observed in Mpe-Cam-1 (Table 4).

None of these nuWTs were found to harbor any intact protein coding gene. Analysis of synteny and

sequence identity of nuWTs across species identified only 34 conserved nuWTs between *M. perstans*

and *M. ozzardi* (Supplementary Material Table 5). Interestingly, about 80% (260 from *M. perstans*

and 280 from *M. ozzardi*) were found to be conserved in within-species comparisons of *M. perstans*

and *M. ozzardi* respectively (Supplementary Material Table 5).

3.7 Analysis and comparison of genes encoding drug targets for the filaricides

diethylcarbamazine, ivermectin and albendazole

The main preventive intervention for filarial infections is based on mass drug administration (MDA)

using diethylcarbamazine (DEC), ivermectin (IVM) and albendazole (ALB) (Simonsen et al 2010).

Although DEC has been successfully used to treat *M. perstans* infections, the drug has been found to

be ineffective against *M. ozzardi* (Chadee et al., 1995). To investigate potential reasons for

differences in drug susceptibility between *M. perstans* and *M. ozzardi*, the genes encoding the drug

targets for DEC, ivermectin and albendazole, were identified and analyzed. A search for the DEC

target *gon-2* (Verma et al., 2020), a gene that encodes a Transient Receptor Protein M (TRPM)

channel subunit, identified two potential homologs in *M. perstans* but one in *M. ozzardi*.

Phylogenetic analysis of the *gon-2* homologs in *Mansonella* and other nematode genomes revealed

two distinct, paralogous clades, denoted as Clade 1 and Clade 2, which represent orthologs of *Cel-*

gon-2 and *Cel-gtl-2* respectively. The nematodes known to be susceptible to DEC, including *C.*

elegans and filarial parasites *B. malayi L. loa*, and *W. bancrofti*, all have a *gon-2* ortholog in Clade 1.

Interestingly, *M. perstans*, which is treatable with DEC, also has a corresponding ortholog in Clade 1

while *M. ozzardi*, which is refractory to DEC, lacks the ortholog for its target gene *gon-2* (Figure 4).

Both *M. perstans* and *M. ozzardi* do possess orthologs of *Cel-gtl-2* in Clade 2, which share only 52 %

sequence identity with the *C. elegans* GON-2 protein in Clade 1 and are unlikely to share their

functions. The absence of a *gon-2* ortholog in *M. ozzardi* is likely to be real and not the consequence

of incomplete assemblies as the gene is absent in both isolates whose genomes were assembled

independently. Similar analysis for the other known DEC target genes, *trp-2*, *ced-11* and *slo*-1

identified the corresponding one-to-one orthologs for all three genes in all the filarial species

including both isolates of *M. perstans* and *M. ozzardi* (Supplementary Figures 4, 5 and 6). Taken

together, this suggests that the absence the *gon-2* ortholog in *M. ozzardi* is most likely responsible for

the ineffectiveness of DEC against this parasite.

The targets of ivermectin as identified through genetic studies in *C. elegans* include various subunits

of the Glutamate☐gated Cl– channels (GluCl), encoded by genes *glc-1*, *glc-2*, *glc-3*, *glc-4*, *avr-14*

and *avr-15* (Dent et al., 2000; Wolstenholme and Rogers, 2005). Phylogenetic analysis of these genes

across filarial parasites shows that the homologs of *glc-1*, *glc-2*, *glc-3*, *avr-14* and *avr-15* form a

family of paralogous genes with complex patterns of homology across filarial parasites (Figure 5),

while *glc-4* has one-to-one orthologs across all genomes analyzed (Supplementary Figure 7).

Orthologs for *Cle-glc-1*, *Cel-glc-2* and *Cel-avr-15* are not present in either of the *Mansonella* species

(Figure 5). Both isolates of *M. perstans* and *M. ozzardi* have one ortholog most closely related to the

Cel-glc-3. Although both isolates of *M. ozzardi* have a homolog most closely related to *Cel-avr-14*,

the corresponding homologs are absent in both *M. perstans* isolates. Mutations in the *avr-14* gene

have been shown to contribute to ivermectin resistance in *C. elegans* (Dent et al., 2000) while *M.*

ozzardi, which has orthologs for ivermectin target *avr-14*, is susceptible to ivermectin (Basano et al.,

2018). The absence of an ortholog for *avr-14* in *M. perstans* (Figure 5) could contribute to the

reported lack of effectiveness of ivermectin against *M. perstans* (Bregani et al., 2006).

The anthelminthic drug mebendazole acts by binding to beta-tubulin protein and inhibiting

microtubule formation (Lacey, 1990). The beta-tubulin gene family in *C. elegans* consists of 6

different members (*ben-1*, *tbb-1*, *tbb-2*, *tbb-4*, *tbb-6*, and *mec-7*) all of which have varying effects on

drug sensitivity (Driscoll et al., 1989; Pallotto et al., 2022). A phylogenetic analysis of the beta-

tubulin gene family in *Mansonella* and other nematodes reveals a complex pattern of expansion and

contraction of this gene family across filarial species, with no direct orthologs for *ben-1* identifiable

in any filarial nematodes (Supplementary Figure 8). Interestingly, there are 5 homologs present in *M.*

ozzardi but only 4 in *M. perstans*, which is lacking an ortholog of the *tbb-4* gene. Mebendazole is

commonly used in combination DEC or ivermectin for treating *M. perstans* infections (Bregani et al.,

2006) but its use for *M. ozzardi* has not been reported. The exact roles of different beta-tubulin

homologs on drug susceptibility in *Mansonella* will require further study.

4   Discussion

To gain some insight into the neglected pathogens responsible for human mansonellosis, we report

the first *de novo* genome assemblies from the filarial parasites *M. perstans* and *M. ozzardi*, providing

reference genomes for each species. The genomes encode approximately 10,000 genes each and are

high quality and largely complete based on universally conserved orthologs (BUSCO scores greater

than 90%) relative to other filarial species. The availability and sequencing of two isolates for *M.*

perstans and *M. ozzardi* also allowed higher confidence in calling presence or absence of genes in

each species as it could be confirmed in two independent isolates. The *Mansonella* genomes average

76 Mb in length and are similar in size to the genomes described from other human and animal

filarial parasites (Ghedin et al., 2007; Godel et al., 2012; Desjardins et al., 2013; Tallon et al., 2014;

Lau et al., 2015; Foster et al., 2020, McCann et al., 2021). The sequencing of two clinical isolates of

M. perstans and *M. ozzardi* also enabled a quantification of the genetic variability both between

species as well in the same parasite from different regions. Most of the genetic differences observed

between isolates and species are in the intergenic regions, and only a small fraction (10%) of variants

affects coding exons. Relatively little is known about natural variation in the nuclear genomes of

filarial parasites (McNulty et al 2013., Qing et al., 2021). In the heartworm *Dirofilaria immitis*, a

closely related filarial parasite of dogs and cats, sequencing of two isolates from Europe and the

United States revealed very little genetic variation in their nuclear or mitochondrial genomes (Godel

et al., 2012). Other studies of genetic variation within isolates of a filarial species have used

mitochondrial *cox1* gene (McNulty et al., 2013) or the mitochondrial genomes (Qing et al., 2021).

Our study presents a more comprehensive analysis of variation in nuclear genomes within isolates. It

will be interesting to investigate how these patterns evolve in other human filarial species.

Transfer of mitochondrial DNA fragments into the nuclear genome has been observed for multiple

filarial nematodes (Grote et al., 2017). These nuMT loci are annotated for all *Mansonella* isolates

using their reported mitochondrial genomes (Crainey et al., 2018; Chung et al., 2020). Most filarial

parasites possess the intracellular bacterial symbiont, *Wolbachia*, capable of transferring genetic

material to the nuclear genome of its host. The genomes of *Wolbachia* endosymbionts *w*Mpe and

*w*Moz from *M. perstans* and *M. ozzardi* have also been reported (Sinha et al., 2022), enabling an

analysis of nuWTs, regions of DNA transfers from *Wolbachia* to its host nuclear genomes. While

reported in other filarial species, the biological function and significance of such nuclear transfer

events are not well understood. Moreover, the levels of conservation of such loci across multiple

isolates of a species, or closely related species has not been previously studied. Our analysis

combining synteny and sequence identity shows that the nuMT loci are poorly conserved in

Mansonella even within the same species, and rarely harbor intact coding genes, suggesting that most

of these transfers are not evolutionarily selected, and non-functional. For the nuWTs, relatively more

loci were found to be conserved across the two species, but isolate specific acquisition of these nuWT

fragments was also observed. However, none of these loci harbored any intact protein-coding genes,

suggesting a lack of functional constraints on their evolution. This observation is consistent with the

detection of mostly non-functional nuWTs in other filarial parasites (Grote et al., 2017) and suggests

that these loci most likely arise randomly and degrade via genetic drift over shorter time scales, as

seen in other organisms (Tvedte et al., 2022; Hazkani-Covo et al., 2010; Bensasson et al. 2003).

Despite their genetic relatedness, *M. perstans* and *M. ozzardi* differ in their geographical distribution,

adaptation to insect vectors, and in their response to drug treatments. We have used these new

genomes to explore the genetic basis for the reported differences in their response to commonly used

anti-filarial drugs. Our analysis of the sequences and phylogeny of gene products targeted by DEC

revealed that *M. ozzardi* does not have an ortholog for *gon-2*, one of the targets of this drug. In

contrast, *M. perstans* as well as other filarial parasites that are susceptible to DEC have a *gon-2*

ortholog. It can therefore be hypothesized that the absence of *gon-2* ortholog in *M. ozzardi* has

rendered it unsusceptible to DEC. Interestingly, a reciprocal pattern was observed for the ivermectin

target encoded by the *avr-14* gene, where it was found to be present in the ivermectin-sensitive *M.*

ozzardi but absent in *M. perstans*, which does not respond to this drug. In contrast, although

mebendazole is used for the treatment *M. perstans*, it was found to lack an ortholog of one of the

targets encoded by the beta-tubulin gene *tbb-4*, while the corresponding ortholog is present in *M.*

ozzardi. Given the complex patterns of homology for this gene family, and the absence of any

reported effects of mebendazole on *M. ozzardi*, the significance of this observation remains to be

determined. In addition to these observations, there could be yet other undiscovered filarial genes that

have also have effects on drug susceptibility and could be responsible for differences between *M.*

perstans and *M. ozzardi*. At least one of the parasites, *M. perstans*, can now be successfully cultured

in vitro enabling functional investigations into its biology (Njouendou et al., 2019). Recent

developments in genome editing of non-model organisms open further possibilities of genetic studies

(Lok et al., 2016; Zamanian and Andersen, 2016) and in the control of parasites (Kwarteng et al.,

2021). The availability of the new *Mansonella* genomes reported here will be a critical genomic

resource for such endeavors and for further studies on drug targets, as well as enabling insights into

the biology and evolution of these parasites.

5   Conflict of Interest

Authors AS, ZL, CBP, LE, RDM, and CKSC are employees at New England Biolabs Inc., Ipswich,

MA 01938 (USA), which manufactures biotechnology reagents including those used in Next

Generation Sequencing. The authors declare that the research was conducted in the absence of any

commercial or financial relationships that could be construed as a potential conflict of interest.

6   Author Contributions

NFL, MUF, FFF, SW collected the clinical samples. AS, ZL, CBP, LE, RDM performed the

sequencing experiments and bioinformatics analysis. AS, ZL, CBP, LE, RDM, CKSC analysed the

data. AS, ZL, CBP, CKSC wrote the initial manuscript. AS, ZL, CBP, LE, RDM, MUF, SW, CKSC

and all authors contributed to the final manuscript.

7   Funding

Research funding was provided by New England Biolabs, Ipswich, MA 01938, USA to AS, ZL, LE,

RDM, CBP, CKSC. FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) supported

the field research in Brazil through a research grant to M.U.F (2013/12723-7) and a doctoral

scholarship to N.F.L. (2013/ 26928-0).

8   Acknowledgments

The authors thank the DNA sequencing core at New England Biolabs. The authors are grateful to the

late Don Comb for initiating and inspiring research into NTDs at NEB.

9   Data Availability Statement

Raw sequencing reads for the *M. per*stans isolates Mpe-Cam-1 and Mpe-Cam-2 have been deposited

in the NCBI SRA database under BioProject accession numbers PRJNA917716 and PRJNA917721

respectively. Illumina sequencing data for *M. ozzardi* isolates Moz-Brazil-1 and Moz-Venz-1 have

been deposited in the NCBI SRA database under BioProject accession numbers PRJNA917722 and

PRJNA917766 respectively.

10   Figure Legends

10.1   Figure 1. Comparisons of syntenic regions across the assembled *Mansonella* genomes.

Whole genome alignments were performed using minimap with default parameters. Syntenic regions

were visualized using Jupiter plots, comparing (A) The two isolates of *M. perstans* (B) The two

isolates of *M. ozzardi*, and (C) the cross-species comparison between Mpe-Cam-1 and Moz-Brazil-1.

10.2   Figure 2. Orthology analysis across filarial parasites.

Homologs conserved across various filarial species, as well as the Onchocercid nematodes *S. digitata*

and *T. callipaeda*, and free-living model nematodes *P. pacificus* and *C. elegans* from Clade V were

identified using OrthoFinder. The set of all 263,019 proteins from these nematode genomes were

grouped into 16,564 orthogroups. The number of orthogroups identified in each species are

represented by horizontal bars in the Upset plot, and the number of orthogroups common across

different species are represented by vertical bars. The presence of an orthogroup in a species is

indicated by a filled, colored dot while the absence is indicated by empty gray dots. Genus-specific

orthogroups are shown as orange dots/bars for *Mansonella*, and blue dots/bars for *Onchocerca* and

Brugia.

10.3   Figure 3. Phylogenomic analysis of filarial parasites.

Phylogenomic tree was generated based on protein sequences of 465 single copy orthologs identified

across all nematode genomes. Onchocercid specie *S. digitata* and *T. callipaeda* were used as

immediate outgroups, while free-living nematodes *C. elegans* and *P. pacificus* were used as more

distant outgroups. The previously described Onchocercid clades ONC2 to ONC5 are highlighted.

10.4   Figure 4. Phylogenetic analysis of genes encoding the targets of the anti-filarial drug

Diethylcarbamazine (DEC).

Phylogenetic analysis of the *gon-2* and *gtl-2* gene family was performed based on their protein

sequences and the tree was rooted at the midpoint. The orthologs of the *Cel-gon-2* gene are placed in

Clade 1, highlighted in blue, and orthologs of *Cel-gtl-2* orthologs are placed in Clade 2, highlighted

green. Genes from *M. perstans* and *M. ozzardi* are in red, and *C. elegans* genes are in blue.

10.5   Figure 5. Phylogenetic analysis of genes encoding Glutamate Chloride channel subunits,

targets of ivermectin.

Phylogenetic analysis of the GluCl gene family was based on their protein sequences and the tree was

rooted at the midpoint. The clade with *Cel-avr-14* orthologs is highlighted in blue, and the clade with

Cel-glc-3 orthologs is highlighted in green. *M. perstans* and *M. ozzardi* genes are in red, and *C.*

elegans genes are in blue.

11 Tables

11.1 Table 1. Summary statistics of *Mansonella* genome assemblies

	M. perstans	*M. perstans*	*M. ozzardi*	*M. ozzardi*
Isolate Name	Mpe1-Cam-1	Mpe-Cam-2	Moz-Brazil-1	Moz-Venz-1
Sampling site	Cameroon	Cameroon	Brazil	Venezuela
Sequencing Platform	PacBio	Illumina	Illumina	Illumina
Assembled genome size	76 Mb	74.39 Mb	75.78 Mb	74.86 Mb
Number of scaffolds	581	6,700	4,789	4,561
Largest scaffold	1.25 Mb	217 kb	317 kb	258 kb
Scaffold N50 size	285.8 kb	28kb	37.75 kb	38.90 kb
%GC	30.31	30.06	29.65	29.67
% Repeats	9.70 %	7.77 %	8.57%	7.91 %
Predicted genes	9,460	9,994	9,899	10,029

549 **11.2 Table 2. Structural and local differences between *Mansonella* genomes**

	Mpe-Cam-2 versus Mpe-Cam-1	Moz-Venz-1 versus Moz-Brazil-1	Moz-Brazil-1 versus Mpe-Cam-1
gANI score	**99.51 %**	**99.50 %**	**89.69**

All local variants categorized by type (% of genome size)	146,854 (0.19 %)	220, 466 (0.29 %)	4,726,827 (6.22 %)
Substitutions, Single Nucleotide Variants	92,837 (0.12 %)	169, 489 (0.22 %)	3,575,532 (4.70 %)
Insertions	34,473 (0.05 %)	19,983 (0.03 %)	401,196 (0.53 %)
Deletions	14,419 (0.02 %)	20,018 (0.03 %)	429,726 (0.57 %)
Substitutions, Multiple Nucleotide variants	5,125 (0.01 %)	10,976 (0.01 %)	320,373 (0.42 %)
All local variants categorized by location in the genome	146,854	220,466	4,726, 827
Variants outside genes (% of all variants)	92,028 (62.7 %)	132, 043 (59.9 %)	2,495, 765 (52.8 %)
Variants within introns (% of all variants)	47,318 (32.2 %)	63,934 (29.0 %)	1,786, 741 (37.8 %)
Variants within exons (% of all variants)	7,508 (5.1 %)	24,471 (8.4 %)	444,321 (9.4 %)
Total number of synonymous and non-synonymous changes caused by variants	8,757	18,502	577,029
Synonymous changes	4,797 (54.8 %)	11,112 (60%)	330,755 (57.32 %)
Non-Synonymous, Missense changes	3,893 (44.5 %)	7,300 (39.5 %)	245,069 (42.47 %)
Non-Synonymous, Nonsense changes	67 (0.77%)	90 (0.48 %)	1,205 (0.21 %)

Total number of Structural differences	1,273	4,114	3,828
Translocations	1,139	4,060	2,975
Relocations	63	1	292
Reshufflings	29	15	176
Inversions	42	38	385

551 **11.3 Table 3. Nuclear mitochondria Transfers (nuMTs) annotated across all genomes**

Species	Isolate	Total number of nuMTs	Total size of nuMTs	Median size of nuMTs	Largest nuMT	Median percent identity to mitochondria
M. perstans	Mpe-Cam1	48	19.240 kb	238 bp	2,591 bp	87.81 %
M. perstans	Mpe-Cam2	39	14.330 kb	249 bp	2,351 bp	86.48 %
M. ozzardi	Moz-Brazil-1	82	35.070 kb	271 bp	3,170 bp	89.57 %
M. ozzardi	Moz-Venz-1	94	35.851 kb	223 bp	3,697 bp	89.50 %

553 **11.4 Table 4. Nuclear Wolbachia Transfers (nuWTs) annotated across all genomes**

Species	Isolate	Total number of nuWTs	Total size of nuWTs	Median size of nuWTS	Largest nuWT	Median percent identity to *Wolbachia*
M. perstans	Mpe-Cam1	328	87.83 kb	173 bp	2,733 bp	91.51 %

M. perstans	Mpe-Cam2	340	76.26 kb	171 bp	1,236 bp	91.84 %
M. ozzardi	Moz-Brazil-1	308	77.33 kb	198 bp	1,515 bp	91.37 %
M. ozzardi	Moz-Venz-1	294	73.54 kb	195 bp	2,151 bp	90.97 %

12   References

Basano, S. de A., Camargo, J. de S. A. A., Fontes, G., Pereira, A. R., Medeiros, J. F., Laudisse, M. C.
de O., et al. (2018). Phase III Clinical Trial to Evaluate Ivermectin in the Reduction of *Mansonella*
ozzardi infection in the Brazilian Amazon. *The American Journal of Tropical Medicine and Hygiene*
98, 786–790. doi: 10.4269/ajtmh.17-0698.

Bélard, S., and Gehringer, C. (2021). High Prevalence of *Mansonella* Species and Parasitic
Coinfections in Gabon Calls for an End to the Neglect of *Mansonella* Research. *The Journal of*
Infectious Diseases 223, 187–188. doi: 10.1093/infdis/jiaa671.

Bensasson, D., Feldman, M. W., and Petrov, D. A. (2003). Rates of DNA Duplication and
Mitochondrial DNA Insertion in the Human Genome. *J Mol Evol* 57, 343–354. doi: 10.1007/s00239-
003-2485-7.

Bregani, E. R., Rovellini, A., Mbaïdoum, N., and Magnini, M. G. (2006). Comparison of different
anthelminthic drug regimens against Mansonella perstans filariasis. *Transactions of The Royal*
Society of Tropical Medicine and Hygiene 100, 458–463. doi: 10.1016/j.trstmh.2005.07.009.

Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M., and Borodovsky, M. (2021). BRAKER2: automatic
eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein
database. *NAR Genomics and Bioinformatics* 3. doi: 10.1093/nargab/lqaa108.

Calvopina, M., Chiluisa-Guacho, C., Toapanta, A., Fonseca, D., and Villacres, I. (2019). High
Prevalence of *Mansonella ozzardi* Infection in the Amazon Region, Ecuador. *Emerging Infectious*
Diseases 25, 2081–2083. doi: 10.3201/eid2511.181964.

Chadee, D. D., Tilluckdharry, C. C., Rawlins, S. C., Doon, R., and Nathan, M. B. (1995). Mass
Chemotherapy with Diethylcarbamazine for the Control of Bancroftian Filariasis: A Twelve-Year
Follow-Up in Northern Trinidad, Including Observations on Mansonella ozzardi. *The American*
Journal of Tropical Medicine and Hygiene 52, 174–176. doi: 10.4269/ajtmh.1995.52.174.

Chung, M., Aluvathingal, J., Bromley, R. E., Nadendla, S., Fombad, F. F., Kien, C. A., et al. (2020).
Complete Mitochondrial Genome Sequence of Mansonella perstans. *Microbiol Resour Announc* 9.
doi: 10.1128/MRA.00490-20.

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., et al. (2012). A program for
annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. *Fly* 6, 80–92. doi:
10.4161/fly.19695.

Cotton, J. A., Bennuru, S., Grote, A., Harsha, B., Tracey, A., Beech, R., et al. (2016). The genome of
Onchocerca volvulus, agent of river blindness. *Nature Microbiology* 2, 16216. doi:
10.1038/nmicrobiol.2016.216.

Coulibaly, Y. I., Dembele, B., Diallo, A. A., Lipner, E. M., Doumbia, S. S., Coulibaly, S. Y., et al.
(2009). A Randomized Trial of Doxycycline for *Mansonella perstans* Infection. *New England*
Journal of Medicine 361, 1448–1458. doi: 10.1056/NEJMoa0900863.

Crainey, J. L., Costa, C. H. A., de Oliveira Leles, L. F., Ribeiro da Silva, T. R., de Aquino Narzetti,
L. H., Serra dos Santos, Y. V., et al. (2020). Deep Sequencing Reveals Occult Mansonellosis
Coinfections in Residents From the Brazilian Amazon Village of São Gabriel da Cachoeira. *Clinical*
Infectious Diseases 71, 1990–1993. doi: 10.1093/cid/ciaa082.

Crainey, J. L., Marín, M. A., Silva, T. R. R. da, Medeiros, J. F. de, Pessoa, F. A. C., Santos, Y. V., et
al. (2018). *Mansonella ozzardi* mitogenome and pseudogene characterisation provides new
perspectives on filarial parasite systematics and CO-1 barcoding. *Scientific Reports* 8, 6158. doi:
10.1038/s41598-018-24382-3.

Debrah, L. B., Phillips, R. O., Pfarr, K., Klarmann-Schulz, U., Opoku, V. S., Nausch, N., et al.
(2019). The Efficacy of Doxycycline Treatment on *Mansonella perstans* Infection: An Open-Label,
Randomized Trial in Ghana. *The American Journal of Tropical Medicine and Hygiene* 101, 84–92.
doi: 10.4269/ajtmh.18-0491.

Dent, J. A., Smith, M. M., Vassilatis, D. K., and Avery, L. (2000). The genetics of ivermectin
resistance in Caenorhabditis elegans. *Proc Natl Acad Sci U S A* 97, 2674–2679.

Desjardins, C. A., Cerqueira, G. C., Goldberg, J. M., Hotopp, J. C. D., Haas, B. J., Zucker, J., et al.
(2013). Genomics of Loa loa, a Wolbachia-free filarial parasite of humans. *Nat. Genet.* doi:
10.1038/ng.2585.

Downes, B., and Jacobsen, K. (2010). A systematic review of the epidemiology of mansonelliasis.
African Journal of Infectious Diseases 4. doi: 10.4314/ajid.v4i1.55085.

Driscoll, M., Dean, E., Reilly, E., Bergholz, E., and Chalfie, M. (1989). Genetic and molecular
analysis of a Caenorhabditis elegans beta-tubulin that conveys benzimidazole sensitivity. *Journal of*
Cell Biology 109, 2993–3003. doi: 10.1083/jcb.109.6.2993.

Emms, D. M., and Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biology* 20, 238. doi: 10.1186/s13059-019-1832-y.

Ferreira, M. U., Crainey, J. L., and Luz, S. L. B. (2021). *Mansonella ozzardi*. *Trends in Parasitology*
37, 90–91. doi: 10.1016/j.pt.2020.03.005.

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., et al. (2020).
RepeatModeler2 for automated genomic discovery of transposable element families. *PNAS* 117,
9451–9457. doi: 10.1073/pnas.1921046117.

Foster, J. M., Grote, A., Mattick, J., Tracey, A., Tsai, Y.-C., Chung, M., et al. (2020). Sex
chromosome evolution in parasitic nematodes of humans. *Nature Communications* 11, 1964. doi:
10.1038/s41467-020-15654-6.

Ghedin, E., Wang, S., Spiro, D., Caler, E., Zhao, Q., Crabtree, J., et al. (2007). Draft Genome of the
Filarial Nematode Parasite Brugia malayi. *Science* 317, 1756–1760. doi: 10.1126/science.1145406.

Godel, C., Kumar, S., Koutsovoulos, G., Ludin, P., Nilsson, D., Comandatore, F., et al. (2012). The
genome of the heartworm, Dirofilaria immitis, reveals drug and vaccine targets. *FASEB J* 26, 4650–
4661. doi: 10.1096/fj.12-205096.

Grote, A., Lustigman, S., and Ghedin, E. (2017). Lessons from the genomes and transcriptomes of
filarial nematodes. *Molecular and Biochemical Parasitology* 215, 23–29. doi:
10.1016/j.molbiopara.2017.01.004.

Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010). New
Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance
of PhyML 3.0. *Systematic Biology* 59, 307–321. doi: 10.1093/sysbio/syq010.

Gurevich, A., Saveliev, V., Vyahhi, N., and Tesler, G. (2013). QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* 29, 1072–1075. doi: 10.1093/bioinformatics/btt086.

Hazkani-Covo, E., Zeller, R. M., and Martin, W. (2010). Molecular Poltergeists: Mitochondrial DNA
Copies (numts) in Sequenced Nuclear Genomes. *PLOS Genetics* 6, e1000834. doi:
10.1371/journal.pgen.1000834.

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., and Vinh, L. S. (2018). UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution* 35, 518–522.
doi: 10.1093/molbev/msx281.

Hotopp, J. C. D., Slatko, B. E., and Foster, J. M. (2017). Targeted Enrichment and Sequencing of
Recent Endosymbiont-Host Lateral Gene Transfers. *Sci Rep* 7, 1–10. doi: 10.1038/s41598-017-
00814-4.

Howe, K. L., Bolt, B. J., Shafie, M., Kersey, P., and Berriman, M. (2017). WormBase ParaSite – a
comprehensive resource for helminth genomics. *Molecular and Biochemical Parasitology* 215, 2–10.
doi: 10.1016/j.molbiopara.2016.11.005.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., and Jermiin, L. S. (2017).
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* 14, 587–589.
doi: 10.1038/nmeth.4285.

Khelik, K., Lagesen, K., Sandve, G. K., Rognes, T., and Nederbragt, A. J. (2017). NucDiff: in-depth
characterization and annotation of differences between two sets of DNA sequences. *BMC*
Bioinformatics 18, 338. doi: 10.1186/s12859-017-1748-z.

Klasson, L., Kumar, N., Bromley, R., Sieber, K., Flowers, M., Ott, S. H., et al. (2014). Extensive
duplication of the Wolbachia DNA in chromosome four of Drosophila ananassae. *BMC Genomics*
15, 1097. doi: 10.1186/1471-2164-15-1097.

Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., and Phillippy, A. M. (2017).
Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Res. 27, 722–736. doi: 10.1101/gr.215087.116.

Kozek, W. J., Palma, G., Henao, A., García, H., and Hoyos, M. (1983). Filariasis in Colombia:
Prevalence and Distribution of *Mansonella Ozzardi* and *Mansonella (=Dipetalonema) Perstans*
Infections in the Comisaría Del Guainía. *The American Journal of Tropical Medicine and Hygiene*
32, 379–384. doi: 10.4269/ajtmh.1983.32.379.

Kriventseva, E. V., Tegenfeldt, F., Petty, T. J., Waterhouse, R. M., Simão, F. A., Pozdnyakov, I. A.,
et al. (2015). OrthoDB v8: update of the hierarchical catalog of orthologs and the underlying free
software. *Nucl. Acids Res.* 43, D250–D256. doi: 10.1093/nar/gku1220.

Kwarteng, A., Sylverken, A., Asiedu, E., and Ahuno, S. T. (2021). Genome editing as control tool for
filarial infections. *Biomedicine & Pharmacotherapy* 137, 111292. doi:
10.1016/j.biopha.2021.111292.

Lacey, E. (1990). Mode of action of benzimidazoles. *Parasitology Today* 6, 112–115. doi:
10.1016/0169-4758(90)90227-U.

Laetsch, D. R., and Blaxter, M. L. (2017). BlobTools: Interrogation of genome assemblies.
F1000Research 6, 1287. doi: 10.12688/f1000research.12232.1.

Lam, K.-K., LaButti, K., Khalak, A., and Tse, D. (2015). FinisherSC: a repeat-aware tool for
upgrading de novo assembly using long reads. *Bioinformatics* 31, 3207–3209. doi:
10.1093/bioinformatics/btv280.

Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*
Methods 9, 357–359. doi: 10.1038/nmeth.1923.

Lau, Y.-L., Lee, W.-C., Xia, J., Zhang, G., Razali, R., Anwar, A., et al. (2015). Draft genome of
Brugia pahangi: high similarity between B. pahangi and B. malayi. *Parasites & Vectors* 8, 451. doi:
10.1186/s13071-015-1064-2.

Lefoulon, E., Bain, O., Makepeace, B. L., d'Haese, C., Uni, S., Martin, C., et al. (2016). Breakdown
of coevolution between symbiotic bacteria *Wolbachia* and their filarial hosts. *PeerJ* 4, e1840. doi:
10.7717/peerj.1840.

Lefoulon, E., Clark, T., Guerrero, R., Cañizales, I., Cardenas-Calligros, J. M., Junker, K., et al.
(2020). Diminutive, degraded but dissimilar: *Wolbachia* genomes from filarial nematodes do not
conform to a single paradigm. *Microbial Genomics* 6, e000487. doi: 10.1099/mgen.0.000487.

Letunic, I., and Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. *Nucleic Acids Research* 49, W293–W296. doi: 10.1093/nar/gkab301.

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34, 3094–
3100. doi: 10.1093/bioinformatics/bty191.

Lima, N. F., Aybar, C. A. V., Juri, M. J. D., and Ferreira, M. U. (2016). *Mansonella ozzardi*: a
neglected New World filarial nematode. *Pathogens and Global Health* 110, 97–107. doi:
10.1080/20477724.2016.1190544.

Lima, N. F., Gonçalves-Lopes, R. M., Kruize, Y. C. M., Yazdanbakhsh, M., and Ferreira, M. U.
(2018). CD39 and immune regulation in a chronic helminth infection: The puzzling case of
Mansonella ozzardi. PLOS Neglected Tropical Diseases 12, e0006327. doi:
10.1371/journal.pntd.0006327.

Lok, J. B., Shao, H., Massey, H. C., and Li, X. (2016). Transgenesis in Strongyloides and related
parasitic nematodes: historical perspectives, current functional genomic applications and progress
towards gene disruption and editing. *Parasitology*, 1–16. doi: 10.1017/S0031182016000391.

Lu, S., Wang, J., Chitsaz, F., Derbyshire, M. K., Geer, R. C., Gonzales, N. R., et al. (2020).
CDD/SPARCLE: the conserved domain database in 2020. *Nucleic Acids Research* 48, D265–D268.
doi: 10.1093/nar/gkz991.

Marçais, G., Delcher, A. L., Phillippy, A. M., Coston, R., Salzberg, S. L., and Zimin, A. (2018).
MUMmer4: A fast and versatile genome alignment system. *PLOS Computational Biology* 14,
e1005944. doi: 10.1371/journal.pcbi.1005944.

McCann, K., Grant, W., and Doyle, S. R. (2021). The genome sequence of the Australian filarial
nematode, Cercopithifilaria johnstoni. *Wellcome Open Res* 6, 259. doi:
10.12688/wellcomeopenres.17258.2.

McNulty, S. N., Foster, J. M., Mitreva, M., Hotopp, J. C. D., Martin, J., Fischer, K., et al. (2010).
Endosymbiont DNA in Endobacteria-Free Filarial Nematodes Indicates Ancient Horizontal Genetic
Transfer. *PLOS ONE* 5, e11029. doi: 10.1371/journal.pone.0011029.

McNulty, S. N., Mitreva, M., Weil, G. J., and Fischer, P. U. (2013). Inter and intra-specific diversity
of parasites that cause lymphatic filariasis. *Infection, Genetics and Evolution* 14, 137–146. doi:
10.1016/j.meegid.2012.11.002.

Mediannikov, O., and Ranque, S. (2018). Mansonellosis, the most neglected human filariasis. *New*
Microbes New Infect 26, S19–S22. doi: 10.1016/j.nmni.2018.08.016.

Nakamura, T., Yamada, K. D., Tomii, K., and Katoh, K. (2018). Parallelization of MAFFT for large-
scale multiple sequence alignments. *Bioinformatics* 34, 2490–2492. doi:
10.1093/bioinformatics/bty121.

Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., and Minh, B. Q. (2015). IQ-TREE: A Fast and
Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular*
Biology and Evolution 32, 268–274. doi: 10.1093/molbev/msu300.

Njouendou, A. J., Ritter, M., Ndongmo, W. P. C., Kien, C. A., Narcisse, G. T. V., Fombad, F. F., et
al. (2017). Successful long-term maintenance of Mansonella perstans in an in vitro culture system.
Parasites & Vectors 10, 563. doi: 10.1186/s13071-017-2515-8.

Nurk, S., Meleshko, D., Korobeynikov, A., and Pevzner, P. A. (2017). metaSPAdes: a new versatile
metagenomic assembler. *Genome Res.* 27, 824–834. doi: 10.1101/gr.213959.116.

Pallotto, L. M., Dilks, C. M., Park, Y.-J., Smit, R. B., Lu, B. T., Gopalakrishnan, C., et al. (2022).
Interactions of Caenorhabditis elegans β-tubulins with the microtubule inhibitor and anthelmintic
drug albendazole. *Genetics* 221, iyac093. doi: 10.1093/genetics/iyac093.

Poole, C. B., Sinha, A., Ettwiller, L., Apone, L., McKay, K., Panchapakesa, V., et al. (2019). In
Silico Identification of Novel Biomarkers and Development of New Rapid Diagnostic Tests for the
Filarial Parasites *Mansonella perstans* and *Mansonella ozzardi*. *Scientific Reports* 9, 10275. doi:
10.1038/s41598-019-46550-9.

Pryszcz, L. P., and Gabaldón, T. (2016). Redundans: an assembly pipeline for highly heterozygous
genomes. *Nucleic Acids Res* 44, e113–e113. doi: 10.1093/nar/gkw294.

Qing, X., Kulkeaw, K., Wongkamchai, S., and Tsui, S. K.-W. (2021). Mitochondrial Genome of
Brugia malayi Microfilariae Isolated From a Clinical Sample. *Frontiers in Ecology and Evolution* 9.
Available at: https://www.frontiersin.org/article/10.3389/fevo.2021.637805 [Accessed May 17,
2022].

Quinlan, A. R., and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* 26, 841–842. doi: 10.1093/bioinformatics/btq033.

Raccurt, C. P. (2018). *Mansonella ozzardi* and its vectors in the New World: an update with
emphasis on the current situation in Haiti. *Journal of Helminthology* 92, 655–661. doi:
10.1017/S0022149X17000955.

Ritter, M., Ndongmo, W. P. C., Njouendou, A. J., Nghochuzie, N. N., Nchang, L. C., Tayong, D. B.,
et al. (2018). *Mansonella perstans* microfilaremic individuals are characterized by enhanced type 2
helper T and regulatory T and B cell subsets and dampened systemic innate and adaptive immune
responses. *PLOS Neglected Tropical Diseases* 12, e0006184. doi: 10.1371/journal.pntd.0006184.

Sandri, T. L., Kreidenweiss, A., Cavallo, S., Weber, D., Juhas, S., Rodi, M., et al. (2021). Molecular
Epidemiology of *Mansonella* Species in Gabon. *The Journal of Infectious Diseases* 223, 287–296.
doi: 10.1093/infdis/jiaa670.

Saunders, G. I., Wasmuth, J. D., Beech, R., Laing, R., Hunt, M., Naghra, H., et al. (2013).
Characterization and comparative analysis of the complete Haemonchus contortus β-tubulin gene
family and implications for benzimidazole resistance in strongylid nematodes. *International Journal*
for Parasitology 43, 465–475. doi: 10.1016/j.ijpara.2012.12.011.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., and Zdobnov, E. M. (2015).
BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics 31, 3210–3212. doi: 10.1093/bioinformatics/btv351.

Simonsen, P. E., Onapa, A. W., and Asio, S. M. (2011). *Mansonella perstans* filariasis in Africa.
Acta Tropica 120, S109–S120. doi: 10.1016/j.actatropica.2010.01.014.

Sinha, A., Li, Z., Poole, C. B., Ettwiller, L., Lima, N. F., Ferreira, M. U., et al. (2021). Genomes of
Wolbachia endosymbionts from the human filarial parasites Mansonella perstans and Mansonella
ozzardi. *bioRxiv*, 2021.03.23.436630. doi: 10.1101/2021.03.23.436630.

Sinha, A., Li, Z., Poole, C. B., Ettwiller, L., Lima, N. F., Ferreira, M. U., et al. (2022). Multiple
origins of nematode-Wolbachia symbiosis in supergroup F and convergent loss of bacterioferritin in
filarial Wolbachia. 2021.03.23.436630. doi: 10.1101/2021.03.23.436630.

Song, H., Buhay, J. E., Whiting, M. F., and Crandall, K. A. (2008). Many species in one: DNA
barcoding overestimates the number of species when nuclear mitochondrial pseudogenes are
coamplified. *Proceedings of the National Academy of Sciences* 105, 13486–13491. doi:
10.1073/pnas.0803076105.

Storer, J., Hubley, R., Rosen, J., Wheeler, T. J., and Smit, A. F. (2021). The Dfam community
resource of transposable element families, sequence models, and genome annotations. *Mobile DNA*
12, 2. doi: 10.1186/s13100-020-00230-y.

Talavera, G., and Castresana, J. (2007). Improvement of Phylogenies after Removing Divergent and
Ambiguously Aligned Blocks from Protein Sequence Alignments. *Systematic Biology* 56, 564–577.
doi: 10.1080/10635150701472164.

Tallon, L. J., Liu, X., Bennuru, S., Chibucos, M. C., Godinez, A., Ott, S., et al. (2014). Single
molecule sequencing and genome assembly of a clinical specimen of Loa loa, the causative agent of
loiasis. *BMC Genomics* 15, 788. doi: 10.1186/1471-2164-15-788.

Ta-Tang, T.-H., Crainey, J. L., Post, R. J., Luz, S. L., and Rubio, J. M. (2018). Mansonellosis:
current perspectives. *Res Rep Trop Med* 9, 9–24. doi: 10.2147/RRTM.S125750.

Ta-Tang, T.-H., Luz, S. L., Crainey, J. L., and Rubio, J. M. (2021). An Overview of the Management
of Mansonellosis. *RRTM* 12, 93–105. doi: 10.2147/RRTM.S274684.

Tavares da Silva, L. B., Crainey, J. L., Ribeiro da Silva, T. R., Suwa, U. F., Vicente, A. C. P.,
Fernandes de Medeiros, J., et al. (2017). Molecular Verification of New World *Mansonella perstans*
Parasitemias. *Emerging Infect. Dis.* 23, 545–547. doi: 10.3201/eid2303.161159.

Tvedte, E. S., Gasser, M., Zhao, X., Tallon, L. J., Sadzewicz, L., Bromley, R. E., et al. (2022).
Accumulation of endosymbiont genomes in an insect autosome followed by endosymbiont
replacement. *Current Biology* 32, 2786-2795.e5. doi: 10.1016/j.cub.2022.05.024.

Verma, S., Kashyap, S. S., Robertson, A. P., and Martin, R. J. (2020). Diethylcarbamazine activates
TRP channels including TRP-2 in filaria, Brugia malayi. *Commun Biol* 3, 1–11. doi: 10.1038/s42003-
020-01128-4.

Vianna, L. M. M., Martins, M., Cohen, M. J., Cohen, J. M., and Belfort, R. (2012). Mansonella
ozzardi corneal lesions in the Amazon: a cross-sectional study. *BMJ Open* 2, e001266. doi:
10.1136/bmjopen-2012-001266.

Wolstenholme, A. J., and Rogers, A. T. (2005). Glutamate-gated chloride channels and the mode of
action of the avermectin/milbemycin anthelmintics. *Parasitology* 131, S85–S95. doi:
10.1017/S0031182005008218.

Yoon, S.-H., Ha, S., Lim, J., Kwon, S., and Chun, J. (2017). A large-scale evaluation of algorithms to
calculate average nucleotide identity. *Antonie van Leeuwenhoek* 110, 1281–1286. doi:
10.1007/s10482-017-0844-4.

Zamanian, M., and Andersen, E. C. (2016). Prospects and challenges of CRISPR/Cas genome editing
for the study and control of neglected vector-borne nematode diseases. *The FEBS Journal* 283, 3204–
3221. doi: 10.1111/febs.13781.

Zimin, A. V., and Salzberg, S. L. (2020). The genome polishing tool POLCA makes fast and accurate
corrections in genome assemblies. *PLOS Computational Biology* 16, e1007981. doi:
10.1371/journal.pcbi.1007981.

Figure 1

(A)

Mpe-Cam-1   Mpe-Cam-2

(B)

Moz-Brazil-1   Moz-Venz-1

(C)

Mpe-Cam-1   Moz-Brazil-1

Figure 2

Figure 3

Figure 4

Figure 5