Long timescale anti-directional rotation in *Drosophila* optomotor behavior

Omer Mano[1], Minseung Choi[2], Ryosuke Tanaka[3], Matthew S. Creamer[3], Natalia C.B. Matos[3],
Joseph Shomar[4], Bara A. Badwan[5], Thomas R. Clandinin[2], Damon A. Clark[1,3,4,6,#]

1 – Department of Molecular, Cellular, and Developmental Biology, Yale University, New
Haven, CT 06511, USA

2 – Department of Neurobiology, Stanford University, Stanford, CA 94305, USA

3 – Interdepartmental Neuroscience Program, Yale University, New Haven, CT 06511, USA

4 – Department of Physics, Yale University, New Haven, CT 06511, USA

5 – Department of Chemical Engineering, Yale University, New Haven, CT 06511, USA

6 – Department of Neuroscience, Yale University, New Haven, CT 06511, USA

– Lead author: damon.clark@yale.edu

Abstract
Locomotor movements cause visual images to be displaced across the eye, a retinal slip that is
counteracted by stabilizing reflexes in many animals. In insects, optomotor turning causes the
animal to turn in the direction of rotating visual stimuli, thereby reducing retinal slip and
stabilizing trajectories through the world. This behavior has formed the basis for extensive
dissections of motion vision. Here, we report that under certain stimulus conditions, two
Drosophila species, including the widely studied *D. melanogaster*, can suppress and even
reverse the optomotor turning response over several seconds. Such 'anti-directional turning' is
most strongly evoked by long-lasting, high-contrast, slow-moving visual stimuli that are distinct
from those that promote syn-directional optomotor turning. Anti-directional turning, like the syn-
directional optomotor response, requires the local motion detecting neurons T4 and T5. A subset
of lobula plate tangential cells, CH cells, show involvement in these responses. Imaging from a
variety of direction-selective cells in the lobula plate shows no evidence of dynamics that match
the behavior, suggesting that the observed inversion in turning direction emerges downstream of
the lobula plate. Further, anti-directional turning declines with age and exposure to light. These
results show that *Drosophila* optomotor turning behaviors contain rich, stimulus-dependent
dynamics that are inconsistent with simple reflexive stabilization responses.

Intro
Visual navigation requires active mechanisms to stabilize trajectories through the world. Insects
exhibit an optomotor turning response, a behavior in which they rotate their bodies in the
direction of visual patterns that rotate about them [1-3]. This behavior is analogous to optomotor

turning responses in fish [4] and the optokinetic response in mammals [5]. In insects, this response is
thought to be a course-stabilization mechanism that minimizes retinal slip, allowing animals to
maintain their trajectory in the face of external or unexpected rotational forces [2,6]. For instance, if
an insect attempts to walk in a straight line, it may slip and turn to the right. From the point of
view of the insect, this turn is observed as optic flow rotating to the left. By responding to this
leftward optic flow with a leftward turn, the insect can recover its original trajectory.

In fruit flies, the optomotor response relies on well-characterized circuitry [7]. Photoreceptor
signals are split into parallel ON and OFF pathways in the lamina and medulla [8-11], which are not
direction-selective. These signals provide input to T4 and T5 cells, which compute direction-
selective responses along four directions at every point in the fly visual field [12-16]. The outputs of
T4 and T5 cells are then summed across visual space by lobula plate tangential cells (LPTCs)
[12,17-20]. Different LPTCs provide distinct signals about the overall pattern of motion surrounding
the fly, and have been linked to head and body movements [21-23].

Interestingly, there have been several reports of flies turning in the direction opposite to what is
predicted by the optomotor turning response. In some cases, these counter-intuitive behaviors
were observed using periodic stimuli with spatial wavelengths smaller than the receptive field of
individual ommatidia, and thus can be accounted for by aliasing [3,24,25]. Work in a tethered flight
simulator showed that when a moving pattern is presented in front of the fly, the animal turned in
the direction of the stimulus motion [26], as expected [27]. However, if the moving pattern was
presented behind the fly, it attempted to turn in the direction opposite to stimulus motion [26]. In a
different experimental preparation, rotational patterns were presented on a dome around freely-
walking flies [28]. Under these conditions, flies generally turned in the direction of motion of the
stimulus, but these rotations were often punctuated by brief, large-magnitude saccades in the
opposite direction. Similarly, experiments using flight simulators have reported spikes in the
torque in the direction opposite the stimulus rotation [29].

Here we show that rotational stimuli can elicit strong, consistent anti-directional turning behavior
in two drosophilid species, *D. melanogaster* and *D. yakuba*. We report that flies respond to high
contrast, high luminance rotational motion stimuli by first turning in the direction of stimulus
motion, and then reversing their trajectory after approximately one second, depending on the
species. In *Drosophila melanogaster*, we characterize the dynamics of this behavior and the
stimuli that drive it, showing that it is distinct from prior observations of anti-directional turning.
The behavior depends critically on adaptation to back-to-front motion. We use the genetic tools
available in *Drosophila melanogaster* to show that this behavior relies on the motion detecting
neurons T4 and T5. Silencing HS and CH, two widefield neurons downstream of T4 and T5,
resulted in small changes in this complex turning behavior. However, the visually evoked
responses of these direction-selective neurons could not account for the anti-directional behavior.
Thus, the observed reversal must be mediated by downstream circuitry. Overall, these results
show that circuits in the fly generate behaviors that oppose the direction of wide-field visual
motion, showing that *Drosophila* turning responses are more complex than a simple stabilizing
reflex.

Results

Anti-directional turning responses to high contrast stimuli

Optomotor turning responses are central to gaze stabilization, so we sought to examine this response across different conditions. Many studies have investigated this behavior using stimuli with low contrast, low light intensity, or both [2,3,30-33], at a variety of different speeds. However, natural scenes can have relatively high contrast and luminance, conditions have been poorly explored in the laboratory. In this experiment, we presented flies with rotational stimuli using high contrast and relatively high luminance.

We tethered individual female *D. melanogaster* above a freely rotating ball to characterize the optomotor response [3,34] (**Fig. 1a**). As expected, low contrast, slow-moving sinusoidal gratings caused flies to turn in the same direction as the moving gratings via the classical optomotor turning response (**Fig. 1b**) [1,3,8,25,26,30,31,35-42]. However, when we changed the stimulus to high contrast sinusoidal gratings (nominal 100% Weber contrast), flies turned in the stimulus direction for approximately 1 second, but then reversed course, and turned in the direction opposite to the stimulus motion for the duration of the stimulus presentation. Because this turning response is in the opposite direction of stimulus and the syn-directional optomotor turning response, we refer to it as anti-directional turning.

We swept a range of contrasts and compared the fly turning in the first 500 milliseconds to the turning after one second (**Fig. 1c**). As contrast increased, the flies turned faster during the first half second of stimulus presentation, reaching a plateau at around 0.5 contrast, consistent with previous results [3,35,36,43-45]. Fly behavior after the first second of stimulation was more complex. As contrast increased from 0 to 0.25, flies turned in the same direction as the stimulus, with faster turning as the contrast increased. When the contrast was greater than 0.25, turning decreased, lowering to no net sustained turning at around 0.8 contrast. Above a contrast of 0.8, flies began to turn in the direction opposite the stimulus.

These initial experiments took place in the lab of author DAC. To confirm that these unexpected responses did not reflect some idiosyncrasy of one specific behavioral apparatus or environment, we repeated these experiments in a second lab, that of author TRC. Under similar conditions, using the same strain of *Drosophila melanogaster*, we reproduced the rapid deceleration after an initial, transient syn-directional response (**Fig. 1d**), with some individual flies exhibiting significant anti-directional turning (**Supp. Fig. S1**). This demonstrates that the key features of this behavioral response are stable across experimental systems and laboratories, though the magnitude of anti-directional turning behavior in *D. melanogaster* is sensitive to some unknown experimental parameter differences between the laboratories.

Individual strains of *D. melanogaster*, and other drosophilid species, display significant variation in their locomotor patterns during walking [46]. Indeed, when we tested a Canton-S *D. melanogaster* strain, we observed milder but significant anti-directional turning at long timescales (**Supp. Fig. S2b**). We reasoned that a strong test of the generality of anti-directional turning would be to examine turning behavior in another species, and selected *D. yakuba*. Strikingly, *D. yakuba* also displayed anti-directional turning behavior under similar conditions (**Fig. 1e**). Thus, this behavior is not an idiosyncratic feature of a single laboratory strain.

Figure 1. Flies turn opposite to the stimulus direction in high contrast conditions

a) We measured fly turning behavior as they walked on an air-suspended ball. Stimuli were presented over 270 degrees around the fly.

b) We presented drifting sinusoidal gratings for 5 seconds (shaded region) with either high contrast (c = 1.0) or low contrast (c = 0.25). When high contrast sinusoidal gratings were presented, flies initially turned in the same direction as the stimulus, then started turning

119
120

121
122
123

124	in the opposite direction after ~1 second of stimulation. Under low contrast conditions,
125	flies turned continuously in the same direction as the stimulus. In these experiments, the
126	sine waves had a wavelength of 60º and a temporal frequency of 1 Hz. Shaded patches
127	represent ±1 SEM. N= 10 flies.
128	**c)** We swept contrast between 0 and 1 and measured the mean turning response during the
129	first 0.5 seconds (purple, purple bar in **b**) and during the last 4 seconds of the stimulus
130	(brown, brown line in **b**). The response in the first 0.5 seconds increased with increasing
131	contrast, while the response in the last four seconds increased from c = 0 to c = 0.25, and
132	then decreased with increasing contrast, until flies turned in the direction opposite the
133	stimulus direction at the highest contrasts. N = 20 flies.
134	**d)** We repeated the presentation of drifting sinusoidal gratings, this time in the lab of author
135	TRC, using a similar behavioral apparatus. Stimulus parameters were as described in (b).
136	In these experiments, the population average shows that flies proceeded to zero net
137	turning at high contrasts, but some individual flies exhibited anti-directional turning
138	responses. N = 20 flies.
139	**e)** We repeated the experiments with *D. yakuba*, also in the lab of TRC, and observed that
140	this species exhibited a robust anti-directional turning response to high contrast gratings
141	and a classical syn-directional turning response to low contrast gratings. N = 11 flies.

Conditions for anti-directional turning behaviors

While anti-directional turning behaviors have been reported before, other groups have presented
similar stimuli without observing anti-directional behavior [2,3,30-33]. We wondered what aspects of
our experimental setup could lead to these behavioral differences. In our experiments, anti-
directional turning was strongly linked to display brightness (**Supp. Fig. S2a**). When the mean
brightness of the screens was reduced from 100 cd/m^2 to 1 cd/m^2, we saw no anti-directional
turning in 5 second trials (though average optomotor behavior did decrease over the course of the
stimulus presentation). When we further reduced the mean brightness to 0.1 cd/m^2, flies persisted
in their optomotor behavior throughout the stimulus presentation. We note that in these low
luminance experiments, low levels of ambient light in the nominally dark experimental rig could
also reduce the effective contrast of the stimulus.

We tested a variety of other factors that might affect anti-directional turning. Anti-directional
turning occurred when experiments were run both at hot temperatures and at room temperature
(**Supp. Fig. S2b**). We also observed anti-directional behavior when flies were reared in the dark
and on different media. We also tested several other experiment conditions (**Supp. Fig. S2c**).
Flies responded with anti-directional turning to high contrast stimuli presented at both blue and
green wavelengths. We glued fly heads to their thorax to ensure stimuli could not be affected by
head movements [21,22], but found no difference between head-fixed and head-free flies. We did
find a few factors that modulated anti-directional turning behavior. In particular, rearing *D.*
melanogaster at 25°C instead of 20°C or testing flies that were two weeks old instead of 12-60
hours old both reduced overall turning behavior and eliminated anti-directional turning. In these
cases, optomotor turning still decreased over the course of the 5 second, high contrast trials, but

did not reverse. As details of rearing temperature and the age at which behavior tests are run
often vary across labs, it is likely that these factors, as well as stimulus brightness, account for
the differences between our observations and the previous literature.

Distinct spatiotemporal tuning of the anti-directional behavioral response

To further characterize the anti-directional response, we swept the spatial and temporal
frequency of the sinusoidal grating stimulus. Using only Weber contrasts of 1, we compared the
early response (first quarter second, **Fig. 2a**) to the late response (after one second, **Fig. 2b**).
Drosophila melanogaster always turned in the optomotor direction during the early stimulus
response. In this early response, flies turned most vigorously to stimuli with short spatial
frequencies (~20º wavelength) and fast temporal frequencies (~8 Hz), in agreement with earlier
studies [26,37,39]. However, during the longer-timescale response to high-contrast stimuli, flies only
turned in the optomotor direction at very high temporal frequencies (> ~16 Hz) and at very low
temporal frequencies (<0.5 Hz). At intermediate temporal frequencies, flies showed a sustained
anti-directional response. The maximal anti-directional response was achieved at 1 Hz and 45º
wavelength, distinct from the conditions for peak classical turning responses. Interestingly, the
stimuli that elicit the strongest anti-directional response appear similar to those that maximally
activate T4 and T5 neurons when those neurons are measured in head-fixed flies [12,37,39,47-49].

Anti-directional turning results from adaptation effects

We were intrigued by the switch from syn-directional to anti-directional turning behavior. To
investigate the dynamics of these changes, we presented a rotating sinusoidal stimulus at contrast
1 for five seconds, and then changed the contrast to 0.25 (**Fig. 2c**). After the switch to low
contrast, the flies quickly reverted classical, syn-directional optomotor behavior, demonstrating
that no long-term switch in directional turning occurs during high contrast stimulus presentation.
This effect did not depend on the periodic nature of these stimuli: a rotating stimulus consisting
of 5º-wide vertical bars with randomly-chosen, binary contrasts [38] yielded similar behavioral
responses (**Fig. 2d**).

To further isolate the causes of this switch in behavior, we developed a stimulus to adapt the fly
to different stimuli before presenting high-contrast rotational sinusoidal gratings to elicit the anti-
directional turning response. This adapting stimulus consisted of five seconds of high contrast
'translational' stimuli, which was then followed by a rotational stimulus (**Fig. 2e**). The
translational stimuli consisted of both left and right hemifields moving either front-to-back or
back-to-front across the fly's two eyes [39]. These stimuli resulted in no net turning by the flies
[39,42]. Adapting the fly with front-to-back stimuli did not have a strong effect on the subsequent
response to rotational stimuli. However, adapting with back-to-front stimuli generated responses
that no longer showed an initial syn-directional turning response, but instead exhibited anti-
directional turning immediately after the rotational stimulus began. This result indicates that the
anti-directional turning results from slow-timescale changes that depend on strong back-to-front
motion stimulation.

Figure 2. Anti-directional turning behavior has distinct tuning and is driven by adaptation.

a) Heatmap of fly turning velocity during the first 0.5 seconds of sinusoidal grating stimulation under high contrast conditions and variable temporal and spatial frequencies. The flies turned in the direction of the stimulus across all conditions and responded most to 8 Hz, 22-degree stimuli. N = 16,21,17,21,7, and 22 flies for spatial frequencies 1/120, 1/90, 1/60, 1/45, 1/30 and 1/22 degrees respectively.

b) Heatmap as in (**a**), measured during the last four seconds of stimulation. Flies turned in the same direction as the stimulus at high and low temporal frequencies, but in the opposite direction of the stimulus at intermediate temporal frequencies, with a maximal anti-directional response at wavelengths between 30º and 60º.

c) Switching stimulus contrast from high to low after 5 seconds caused flies to revert to syn-directional behavior after the anti-directional response. N = 7 flies.

d) Presenting rotating random binary patterns (5-degree vertical strips rotating at 150 degrees/second) induced anti-directional turning similar to that elicited by rotating sine wave gratings. N = 7 flies.

e) We presented flies with five seconds of "translational" stimuli (dark shaded region), with high contrast sinusoidal gratings moving either front-to-back or back-to-front, bilaterally, for five seconds. After that, we presented high contrast rotational sinusoidal grating stimuli (60º wavelength, 1 Hz). Front-to-back stimulation did not affect the subsequent response to rotational stimuli, but back-to-front stimuli caused flies to turn immediately in the opposite direction of the stimulus. N = 18 flies.

Anti-directional turning is elicited when stimuli are presented in front of the fly

A previous report of anti-directional turning behavior in flying tethered flies showed that flies turn in the opposite direction to stimuli that are presented behind their midline [26]. To test whether our results were caused by this effect, we split our stimulus into three regions: 90 degrees in front of the fly, 45 degrees in front of the midline on either side of the fly, and 45 degrees behind the midline on either side of the fly (**Fig. 3a**). We found that flies displayed anti-directional turning when presented with stimuli only in the front region or only just in front of the midline (**Fig. 3bc**). They did not display anti-directional turning when moving stimuli were presented behind the midline (**Fig. 3bc**). This suggests a different mechanism from the behaviors that depend on posterior spatial location to elicit reverse-turning [26].

Anti-directional responses do not depend on saccades

Anti-directional saccades have been reported in walking and flying flies [28,29]. In walking flies [28], flies largely turned in syn-directionally, but these turns were sometimes interrupted by brief, high-amplitude saccades in the opposite direction, against the stimulus direction. If such saccades were frequent or high amplitude, the net effect could shift the average turning we measured, creating apparent anti-directional turning. To investigate this possibility, we plotted the turning response on a per-trial basis (**Fig. 3d**). We then discarded information about the magnitude of the turns and considered only the direction of the turning at each point in time (**Fig. 3e**). Strikingly, in many trials, flies continued to turn opposite to the stimulus for several seconds, a behavior unlike brief saccades. We then calculated a turning index for each response timepoint (sampled at 60 Hz). This turning index represented the fraction of trials where the fly turned in the direction of the stimulus at each timepoint minus the fraction of trials where the fly turned in the opposite direction (**Fig. 3f**). Since this turning index does not include the magnitude of turning, it is strongly affected by sustained low-amplitude turns and discounts any brief high-amplitude saccades. When presented with high contrast stimuli, flies maintained a negative turning index, indicating that sustained turns, and not high velocity saccades, underlie this anti-directional turning behavior. As such, it appears distinct from the reports of anti-directional saccades.

Figure 3. Anti-directional turning is driven by stimuli in the forward-facing visual field and is not driven by saccades.

a) We divided our panoramic display into three sections — the front 90º, the 45º behind the fly on either side, and a middle 45º.

b) High contrast sinusoidal gratings were presented on each of these three display sections, with the remaining sections blank. Flies turned syn-directionally when stimuli were presented behind the fly, and turned anti-directionally when stimuli were presented in front of the fly. Shaded patches represent ±1 SEM. N = 55 flies.

c) Average turning in the last 4 seconds of the stimulus (black bar in **b**), in low contrast and high contrast conditions. Shaded patches in the time trace plots represent ±1 SEM. N = 55 flies.

d) A single fly responds to many trials of sinusoidal grating stimuli at high contrast (blue bar) and low contrast (orange bar). We show a heatmap of the fly's responses over time (horizontal axis) and across trials (vertical axis).

e) We can ignore the magnitude of the turning and instead only quantify whether the fly was turning in the same direction as the stimulus (white area) or in the opposite direction (dark gray area). This shows sustained anti-directional turning, not brief saccades.

f) Averaging the direction (but not magnitude) of turning across trials and across flies yields a turning index for each point in time. Shaded patches in the time trace plots represent ±1 SEM. N = 7 flies.

Anti-directional turning requires elementary motion detectors

What neurons are involved in this anti-directional turning behavior? Previous work demonstrated that T4 and T5 are required for directional neural responses [18], as well as for optomotor turning [12,40,50], for walking speed regulation [39], and for responses to visual looming stimuli [51]. We

silenced the neurons T4 and T5 using shibire[ts] [52] and measured responses to sinusoidal stimuli
that switched from high to low contrast (**Fig. 4a**). Flies in which T4 and T5 had been silenced
displayed only minimal responses to motion stimuli, with anti-directional turning suppressed
along with classical syn-directional turning. Thus, we conclude that, like optomotor turning
behaviors, this anti-directional behavior depends critically on signals from T4 and T5.

Anti-directional turning requires the CH lobula plate tangential cell
Since the switch from optomotor to anti-directional behavior seems to be dependent on the
direction of motion adaptation (**Fig. 2e**), we reasoned that neurons involved in this behavior were
likely to be downstream from T4 and T5. Relatively little is known about circuitry that connects
the neurons T4 and T5 to optomotor turning behavior. However, Horizontal System (HS) cells
are well-studied postsynaptic partners of T4 and T5 [9,20]. These lobula plate tangential cells
integrate information from front-to-back and back-to-front selective T4 and T5 cells across the
fly's visual field [17]. HS cells have been implicated in visually-evoked head turns [21] and body
rotations in flight [22] and in maintenance of direction during walking [53]. When we silenced HS
neurons, we found small deficits in syn-directional turning behavior, consistent with prior
results, but no deficits in anti-directional turning (**Fig. 4b**), indicating that HS cells synaptic
output is not required specifically for anti-directional turning behavior.

Next, we turned to the CH lobula plate tangential cells. These cells are GABAergic and are both
pre-synaptic and post-synaptic in the lobula plate [54]. In blowflies, these neurons play an
inhibitory role in an interconnected LPTC circuit that shapes behavior [55]. When we silenced CH
neurons, we found a small increase in syn-directional turning and a decrease in anti-directional
turning (**Fig. 4c**). Overall, silencing this neuron type caused the flies to turn more in the direction
of motion. This result suggests that CH activity contributes to the anti-directional turning
response. However, since adapting to back-to-front translational stimuli significantly affected the
dynamics of anti-directional turning, it seems likely that other neurons beyond HS and CH are
involved, since these two neurons both respond selectively to front-to-back motion [20,56].

308

Figure 4. Syn-directional and anti-directional turning share common circuitry

310   **a)** We silenced T4 and T5 neurons by expressing shibire[ts] selectively in those neurons. We
311   measured turning behavior during a contrast-switching stimulus (as in **Fig. 2c**). Results
312   from flies with T4 and T5 silenced shown in dark red, while controls are in light red and
313   gray. Average fly behavior during the last four seconds of the first contrast (black bar on
314   left) shown as bars on the right, with individual fly behavior shown as dots. Note that the
315   data labeled "low contrast" are from experiments in which the low-contrast stimulus was
316   shown before the high contrast stimulus. Shaded patches in the time trace plots represent
317   ±1 SEM, as do vertical lines on bar plots. *** indicates experimental results are
318   significantly different from results, $P < 0.001$ via a two-sample Student t-test. * indicates
319 $P < 0.05$. N = 17, 24, 19 flies with genotypes T4T5/Shibire[ts], T4T5/+, +/Shibire[ts].
320   **b)** Results from HS silencing as in **a**. Silencing HS reduced syn-directional turning behavior
321   ($P < 0.001$) but did not have a strong effect on anti-directional turning. N = 34, 21, 19
322   flies with genotypes HS/Shibire[ts], HS/+, +/Shibire[ts].
323   **c)** Results from CH silencing as in **a**. CH silencing reduced the degree of anti-directional
324   turning ($P < 0.001$). N = 63, 57, 70 flies with genotypes CH/Shibire[ts], CH/+, +/Shibire[ts].

325

326   Early direction-selective cells do not adapt to the stimulus
327   The anti-directional turning response is preceded by an initial syn-directional response. This
328   change in behavior must be the result of changes in neural activity, but this change could happen

at any point along the neural pathway between photoreceptors and motor neurons. In order to constrain possible mechanisms for generating the anti-directional turning behaviors, we used calcium imaging to interrogate the activity of direction selective neurons during high and low contrast stimulation (**Fig. 5a**). However, as calcium imaging experiments using two photon microscopy require additional spectral filtering of the projector, we first confirmed that these spectral differences did not alter anti-directional turning responses. To do this, we re-measuring the anti-directional turning behavior using optical filtering matched to the conditions needed for imaging. Using this spectrally distinct illuminant, we observed both syn-directional and anti-directional turning behaviors, following the previously observed dynamics (**Supp. Fig. S3**).

As T4 and T5 neurons play a critical role in both the syn- and anti-directional turning responses, we first measured the calcium activity of these neurons as they responded to sine wave gratings at a range of contrasts in their preferred and null directions. The T4 and T5 neurons responded to sine wave gratings in their preferred direction by increasing their calcium activity for the full duration of the stimulus presentation, reaching a plateau after approximately 1 second (**Fig. 5bc**, middle). As we increased the contrast of the preferred direction stimuli, we found that both T4 and T5 cells had increased calcium activity throughout the contrast range (**Fig. 5bc**, right), consistent with prior measurements [12]. Thus, the responses of T4 and T5 cells do not capture the transition from syn-directional to anti-directional turning behavior.

Next we examined two LPTCs downstream of T4 and T5 cells. Calcium activity in HS cells followed similar trends to T4 and T5. Calcium signals increased at the start of preferred direction stimuli presentation and stayed high until the end of the presentation (**Fig. 5d**, middle). Increasing contrast caused stronger calcium responses with a mild saturation effect at high contrast (**Fig. 5d**, right), consistent with prior voltage measurements [20]. These results indicate that the changes in the time course of optomotor behavior at high contrast are not related to changes in HS activity. Finally, we measured calcium activity in CH cells. CH cells responded to visual stimuli more quickly than HS cells (**Fig. 5e**, middle), and showed decreased calcium signals in response to null direction stimuli (**Fig. 5e**, right). However, they also showed sustained responses to high contrast stimuli, as in T4, T5, and HS. These measurements suggest that the switch from syn- to anti-directional turning behavior is driven by cells downstream of or parallel to T4, T5, HS, and CH.

Figure 5. Responses in early direction-selective cells do not show a reduction or reversal of response on the timescale of the behavior.

    a) We used two-photon microscopy to measure calcium activity in lobula plate neurons while presenting sinusoidal gratings at a range of contrasts.

    b) T4 cells, marked in orange (*left*), responded to drifting sinusoidal gratings with increased calcium activity (*middle*). Darker colors indicate higher contrast, preferred direction in blue, null direction in red. When integrated across the stimulus presentation (*right*), calcium activity increased with stimulus contrast. N = 8 flies.

    c-e) As in **b)** measuring calcium activity in T5, HS, and CH cells. N = 8, 10, 15 flies.

Adult plasticity in anti-directional turning behavior

In behaving flies, the strength of anti-directional turning was dependent both on rearing temperature, which alters the rate of growth, and on age (**Supp. Fig. S2**). This raises the possibility that syn- and anti-directional turning responses might be plastic during the early adult stages of development. To probe this possibility, we presented 1 Hz, high-contrast, rotating

sinusoidal grating at various stages during early adulthood (**Fig. 6**). Strikingly, as flies aged from 0.5 to 4 days post eclosion (dpe), the initial syn-directional turning became less transient and more sustained, indicative of a weaker anti-directional turning drive. We then wondered whether this plasticity was intrinsically programmed, or dependent on visual input. To disambiguate these possibilities, we reared flies in darkness to 2 or 4 dpe and measured their turning responses (**Fig. 6**, *gray*). Dark-reared flies exhibited a stronger deceleration away from syn-directional turning, similar to that found in more juvenile flies, arguing that visual input may sculpt the balance of syn- and anti-directional turning. Finally, we examined whether optomotor response plasticity could be detected in *D. yakuba*. However, in this species, anti-directional responses were stable across the first four days of adulthood, arguing that the role of visual experience in shaping these responses is itself evolutionarily tuned in drosophilids (**Supp. Fig. 4**).

Figure 6. Maturation of optomotor response in early adulthood

a) Adult flies at various ages post eclosion were presented with 5-second, high-contrast, rotating sinusoidal gratings as in **Fig 2b**. As the flies aged from 1 day post eclosion (dpe) to 2, 4, and 8 dpe, the initial anti-directional turning response transitioned into syn-directional turning. Dark-rearing flies at 2 dpe reduced this maturation effect. Shaded patches represent ±1 SEM. N = 5-14 flies.

b) The last 1.5 seconds of the mean turning velocity of each fly was averaged, and the population response was plotted.

c) As in (a) but in the TRC lab, using 0.5, 1, 2, and 4 dpe, with dark rearing for 4 dpe. With maturation, the syn-directional turning became less transient. N = 9-15 flies.

d) As in (b) but for data in (c).

Discussion

In this study, we found we could elicit robust turning in the opposite direction of high contrast motion stimuli (**Fig. 1**). This behavior is qualitatively different from other turning behaviors reported in the literature (**Figs. 2** and **3**), but shares elements with the circuitry necessary for optomotor behavior (**Fig. 4**). However, the switch from syn-directional turning behavior to anti-directional turning behavior is not a reflection of changes in the activity of known direction-selective neuron types in the early visual system (**Fig. 5**). Moreover, this anti-directional turning behavior exhibits a degree of experience-dependent plasticity (**Fig. 6**).

Anti-directional turning is distinct from other against-stimuli behaviors

The anti-directional turning behavior we have characterized is distinct from previous reports of flies turning in the direction opposite to the stimulus motion. First, some opposite-direction turning behaviors can be explained by stimulus aliasing [3]. Aliasing cannot explain our results because the stimulus that maximally activates anti-directional behavior has a spatial frequency of 1/60 cycles per degree, well below the Nyquist frequency of the fly eye (~1/10 cycles per degree) [3,24] and below reports of higher acuity vision in flies [57]. Aliasing would also not explain the dependence on stimulus contrast.

Second, our observations also cannot be explained by stimuli to the rear of the fly driving it in the opposite direction [26], since we observe anti-directional turning even when stimuli are only presented in only the 90 degrees in front of the fly (**Fig. 3**).

Third, it is also distinct from previous reports of reverse body saccades [28] since it manifests in persistent turns in the opposite direction of the stimulus and can be measured even when the magnitude of the turns is discarded (**Fig. 3**).

Fourth, the behavior observed here also appears to be distinct from previously-observed stimulus-density dependent behavioral reversals [58]. Those previously reported behaviors showed immediate reversals, but it took ~1 second for flies in our paradigm to switch between optomotor and anti-directional behaviors.

Anti-directional turning is unlikely to be due to adaptation to contrast alone

In mammalian retina, the direction preference of cells can switch because of upstream circuit adaptation [59,60]. However, we do not believe the anti-directional turning we observe has similar causes. In the mammalian retina, direction switching occurs when non-direction-selective neurons adapt to high contrast stimuli, which distorts the downstream direction-selective computation. Since the adaptation in those experiments occurs in non-direction-selective neurons, it cannot be affected by the direction of the adapter stimulus. However, we see

differences in turning behavior depending on whether we adapt with front-to-back or back-to-
front stimuli (**Fig. 2e**). This observation rules out a mechanism based solely on contrast, since the
contrast content of front-to-back and back-to-front stimuli are identical.

The fly's visual system, however, adapts its gain to stimulus contrast [61,62]. Importantly, the
phenomenology of the anti-directional turning also argues that the contrast adaptation is
incomplete or heterogeneous among neurons, since contrast 1 and contrast 0.25 stimuli result in
such different behaviors. Contrast adaptation reported in the fly is also faster than the 1-2
seconds preceding the shift to anti-directional turning in these experiments.

Anti-directional turning behavior may require specific experimental and rearing conditions
Despite these previous reports of anti-directional turning under certain conditions, other labs
have measured sustained optomotor turning in response to high contrast stimuli [2,31,32,36]. Two
major causes of this difference are likely display brightness and rearing conditions. Some
experiments employ displays with mean luminances less than 5 cd/m^2 [31,33,36]. Our screens, with a
mean luminance of 100 cd/m^2, are substantially brighter, but not especially bright when
compared to natural scenes. In daytime natural scenes, foliage and the ground have average
luminances of 200-500 cd/m^2 and the sky has an average luminance of around 4000 cd/m^2 [63]. We
suspect that as researchers move to using displays that can more accurately depict natural scene
luminances, anti-directional turning behaviors will be encountered more frequently.

Rearing conditions also significant influenced anti-directional turning behavior. Flies reared at
25°C showed less anti-directional behavior than those reared at 20°C. Temperature has known
developmental effects on neural connectivity [64]. We also found differences based on fly age and
fly strain. Notably, all three of these parameters vary significantly across the field, with prior
studies varying rearing temperatures from 18 to 20 to 25°C (see for instance [36,39,57,65,66]), ages
from 1 day to 10 days (see for instance [26,30,42]), and strain between CantonS or OregonR (see for
instance [8,33]). Thus, these factors likely account for the fact that this phenomenon has not
previously been reported, even as rotating sinusoids have been widely used in behavioral
experiments.

Tuning of anti-directional turning matches tuning of direction selective neurons
The study of anti-directional turning behavior may yield clues about the temporal tuning of fly
motion detectors. Optomotor behavior is tuned to visual stimuli in the range of 8-22 Hz [26,37,39,67],
while anti-directional behavior is tuned to stimuli in the range of 0.5-4 Hz (**Fig. 2**). Intriguingly,
this slower tuning matches the tuning of T4, T5, and HS neurons, as measured via calcium
imaging or electrophysiology [12,20,39,68]. Previous studies have suggested that the difference in
tuning between behavior and imaging are due to octopamine that is released during behavior but
not necessarily released during imaging [37,47,68]. In this work, we demonstrate a motion-related
behavior tuned to low frequencies, comparable to those in neural measurements, during behavior
that requires T4 and T5 neurons. Overall, this suggests that T4 and T5 are required for behaviors
with very different temporal tuning, which in turn suggests that the temporal tuning of behavior
is not determined solely by T4 and T5 tuning, but by other, parallel pathways as well.

Anti-directional turning reveals circuits that turn the fly counter to visual motion

Experiments that show a decrease of turning over time to high contrast stimuli (e.g., **Fig. 1d**) could plausibly be explained by some kind of gain reduction or adaptation over time. However, the existence of turning in the direction opposite the stimulus motion in *D. melanogaster* and in *D. yakuba* requires a different explanation. These experiments reveal that over long timescales, a circuit that opposes the syn-directional optomotor turning response can dominate the behavioral response. Thus, this circuit is not simply scaling the magnitude of turning responses, but rather must be implementing an antagonistic, subtractive operation. Measurements of free walking behavior have shown that the time constant of the autocorrelation of fly turning is around 100 ms [69,70]. Opposing syn- and anti-directional turning circuits could be used to balance and tune the strength of turning responses on short timescales, while the anti-directional turning is revealed on longer timescales. This sort of subtractive processing predominates in computing motion signals in the visual systems of insects [17] and mammals [71], and could also explain the existence of syn- and anti-directional turning behaviors.

In summary, we have presented evidence of a transition from syn-directional turning to no turning or to anti-directional turning when high contrast stimuli are presented to the fly. This persists across laboratory environments and across *Drosophila* species and shows plasticity with age. This behavior suggests that turning in response to rotational stimuli is not a simple reflex. Instead, the turning is likely driven by circuits with opposing influences on turning direction. These circuits appear to differentially adapt to the direction and contrast of the stimulus. This complexity makes the optomotor response a model for studying the interactions of circuits as they control the low-dimensional behaviors that change an animal's orientation.

Contributions

OM, MC, RT, MSC, NCBM, JS, and BAB collected data. OM, MC, TRC, and DAC wrote the paper.

Acknowledgements

This work was supported by NIH R01EY026555 (DAC), R01EY022638 (TRC), and by a Chan-Zuckerberg Investigator Award (TRC). MC was supported by an NDSEG Fellowship; RT was supported by the Takenaka Foundation; MSC was supported by an NSF GRFP; NCBM was supported by a CAPES fellowship; JS was supported by a Ford Foundation Fellowship. We also gratefully acknowledge Irving E. Wang, who performed initial experiments examining optomotor responses in many drosophilid species in the lab of TRC.

Methods

Data availability

Raw behavioral and imaging data, along with code to run the analyses and create the plots in this paper, are available on Dryad: https://doi.org/10.5061/dryad.stqjq2c77.

Fly strains

Strains used in these experiments are listed in the tables below:

Table 1: Parental stock genotypes

Name	Genotype	Source	Stock #
Wildtype	+; +; + (IsoD1)	[72]	N/A
T4T5-Gal4	+; +; R42F06-Gal4 (IsoD1 background)	BDSC	BDSC 41253
HS-Gal4	+; +; R27B03-Gal4 (IsoD1 bg)	[31]	BDSC 49211
CH-Gal4	w; +; R35A10-Gal4 (Janelia bg)	BDSC	BDSC 49897
UAS-Shibire^{ts}	+; +; UAS-Shibire^{ts} (IsoD1 bg)	[42]	N/A
Empty Gal4	w; +; pBDPGAL4.1Uw (Janelia bg)	BDSC	BDSC 68384
GCaMP6f	w; UAS-GCaMP6f; +	BDSC	BDSC 42747
jGCaMP7b	w; +; UAS-jGCaMP7b	BDSC	BDSC 79029
mtdTomato	w; +; UAS-mtdTomato	BDSC	BDSC 30124

Table 2: Genotypes of flies used in behavior experiments

Experimental	Gal4 Control	UAS Control	Background Control
T4T5-Gal4 x UAS-Shibire^{ts}: +; +; R42F06-Gal4/UAS-Shibire^{ts}	T4T5-Gal4 x IsoD1: +;+;R42F06-Gal4/+	IsoD1 x UAS-Shibire^{ts}: +; +; +/UAS-Shibire^{ts}	IsoD1: +; +; +
HS-Gal4 x UAS-Shibire^{ts}: +; +; R27B03-Gal4/UAS-Shibire^{ts}	HS-Gal4 x IsoD1: +; +; R27B03-Gal4/+	IsoD1 x UAS-Shibire^{ts}: +; +; +/UAS-Shibire^{ts}	IsoD1: +; +; +
CH-Gal4 x UAS-Shibire^{ts}: w/+; +; R35A10-Gal4/UAS-Shibire^{ts}	CH-Gal4 x IsoD1: w/+; +; R35A10-Gal4/+	Empty Gal4 x UAS-Shibire^{ts}: +/w; +; pBDPGAL4.1Uw /UAS-Shibire^{ts}	Empty Gal4 X IsoD1: +/w; +; +/ pBDPGAL4.1Uw

Genotypes of files used in imaging experiments: +; +; HS-Gal4/UAS-jGCaMP7b, +; UAS-GC6f/+; T4T5-Gal4/UAS-mtdTomato, w/+; +; CH-Gal4/UAS-jGCaMP7b.

Fly rearing (DAC lab)

Unless otherwise noted, flies were reared at 20 degrees Celsius in Panasonic MIR-154-PA incubators (Panasonic/PHC, Tokyo, Japan). The flies were circadian entrained on 12-hour light-dark cycles. Flies were raised on Archon Scientific glucose food (recipe D20102, Archon Scientific, Durham, NC). We used CO_2 to anesthetize flies more than 12 hours before the behavioral experiments.

Flies were tested for behavior in rigs built in the labs of DAC and TRC. Behavior shown in **Figs.**
1d, **1e**, **6c**, **6d**, **S1**, and **S4** was acquired in the lab of TRC, while the rest was obtained in the lab
of DAC.

Fly rearing (TRC lab)
Flies were reared at 25°C, on molasses-based food, and circadian entrained on 12-hour light-dark
cycles. Flies were collected within three hours of eclosion using brief CO_2 anesthetization. *D.*
melanogaster and *D. yakuba* were raised under identical conditions. Dark-reared flies were put
in a dark chamber within 3 hours of eclosion. Flies tested at 0.5 days post eclosion were
collected during the first two hours of the light cycle and were exposed to light until they were
tested.

Stimulus generation and behavioral turning assays (DAC lab)
Stimuli were presented using DLP Lightcrafter (Texas Instruments, Dallas, TX) projectors [34].
Mirrors were used to bounce the projected light onto three screens made of back-projection
material, surrounding the fly. The screens covered the front 270 degrees around the fly, and ~45
degrees in elevation above and below the fly. The projectors were set to monochrome mode
(green unless otherwise noted), updating at 180 Hz. Stimulus video was generated through a
custom MATLAB (Mathworks, Natick, MA) application using PsychToolbox [73]. Stimuli were
mapped onto a virtual cylinder around the fly and the MATLAB application generated a
viewpoint-corrected video signal.

Behavioral experiments were performed 12-60 hours after staging. For behavioral experiments,
we selected female flies, and co-housed them with males after staging. Flies were cold-
anesthetized and fixed to needles using UV-cured epoxy (Norland optical adhesive #63, Norland
Products, Cranbury, NJ). Flies were then placed above air-suspended polypropylene balls. These
balls were 6 mm in diameter and weighed ~120 mg. The balls were painted with two layers of
marker coatings- a base silver layer and a red top layer. The motion of balls was detected by
either a Parallax mouse sensor board (Parallax, Rocklin, CA) with an MCS-12086 sensor (Unity
Opto Technology, Taipei, Taiwan), or a custom board with an ADNS 2080 sensor (Avago
Technologies / Broadcom Inc, San Jose, TX). The data from these sensors were transferred to a
custom MATLAB application via an Arduino Uno board.

Stimulus generation and behavioral turning assays (TRC lab)
Stimuli were presented using a DLP Lightcrafter (Texas Instruments, Dallas, TX) projector.
Three coherent optic fibers were used to direct the projected light onto three screens made of
back-projection material, surrounding the fly [8,38]. The screens covered the front 270 degrees
around the fly, and ~45 degrees in elevation above and below the fly. The projectors were set to
monochrome mode, updating at 120 Hz. Stimulus video was generated through Flystim
(https://github.com/ClandininLab/flystim), a custom Python application developed in the
Clandinin Lab [74]. Stimuli were mapped onto a virtual cylinder around the fly and Flystim
generated a viewpoint-corrected video signal.

Behavioral experiments were performed 12-48 hours after eclosion, as described in the figures.
Flies were cold-anesthetized and fixed to needles using UV-cured adhesive (Bondic, Niagara
Falls, NY). Flies were then placed above air-suspended balls made with LAST-A-FOAM FR-
4615 polyurethane foam (General Plastics, Tacoma, WA). These balls were 9 mm in diameter
and weighed ~91.7 mg. The motion of balls was detected by a Flea3 FL3-U3-13Y3M camera
(Teledyne Flir, Wilsonville, OR) and Fictrac software [75].

Imaging procedures
Two photon imaging (**Fig. 5**) was performed as previously described [76]. Briefly, two-photon
images were acquired with a Scientifica microscope at between 6 and 13 Hz using a 930 nm
femtosecond laser (SpectraPhysics, Santa Clara, USA) using ScanImage [77]. Visual stimuli were
presented on three screens occupying 270º of azimuthal angle about the fly using projectors [34].
Optical filters on the projector and emission filters prevented the visual stimulus light from
leaking into the two-photon images.

Regions of interest (ROIs) were extracted from image timeseries using a watershed algorithm.
Responsive ROIs were included in the analyses. For T4 and T5 neurons, each ROI was identified
as a T4-dominant or T5-dominant ROI by its response to light vs. dark edges, following prior
procedures [78]. For all neuron types, responses were averaged over ROIs and over trials of each
stimulus type to obtain a measurement for each fly; these fly measurements acted as the
independent measurements to compute means and standard error bars for the figure.

Statistical tests
Throughout the paper, each fly was considered an independent sample for statistical purposes.
Means and standard errors were computed over flies. For imaging experiments, regions of
interest from a specific neuron type were first averaged within each fly, creating a value for each
fly's response. These values were used to calculate means and standard errors over the tested
flies. In the silencing experiments, a 2-sample Student t-test was used to test for significant
differences between the experimental genotype and parental controls.

Supplementary Figure S1. Individual *D. melanogaster* flies in TRC lab experiments show anti-directional turning.

a) Mean time traces of individual fly responses to the high contrast stimulus, averaged over trials. The flies are those in **Fig. 1d**.
b) Long-timescale responses of individual flies, averaged over the last 1.5 s of the 5-second stimulus in panel (a) (indicated by thick black line). Mean and SEM shown are over the trials presented to that fly.

Supplementary Figure S2. Flies perform anti-directional turning under a wide range of stimulus and growing conditions.

a) Fly turning behavior at different mean screen brightness. We swept brightness from 100 cd/m² to 0.1 cd/m² and measured turning responses to high and low contrast stimuli. Flies performed the most anti-directional behavior in response to high brightness stimuli. At 1 cd/m², flies never turned in the opposite direction of the stimulus, and at 0.1 cd/m², flies turned continuously in the same direction as the stimulus, even in high contrast conditions. We also measured average turning during the last four seconds of stimulation (black bar above time traces). Average fly behavior shown as bars on the right, with individual fly behavior shown as dots. Shaded patches in the time trace plots represent ±1 SEM, as do vertical lines on bar plots. N = 19, 10, 9, 8 flies, top to bottom.

b) Our wildtype flies were Oregon-R strain [72] raised at 20 degrees. They were grown on glucose-based food media with 12-hour light-dark cycles. Experiments were run at high temperature, 12-60 hours after eclosion. We used uniform, red balls to avoid visual feedback from walking. The response of these wildtype flies to a contrast-switching stimulus (as in **Fig. 2c**) is shown in the upper left corner. We also tested different variations of all these parameters. Canton-S flies turned less overall, and showed less anti-directional turning, but still turned in the opposite direction after 5 seconds of high

contrast stimuli. We tested flies walking on highly-visible silver balls with black dots and
saw behavior similar to wildtype. Two-week-old flies showed reduced turning and much
reduced anti-directional behavior. Flies raised at 25 degrees Celsius had behavior similar
to two-week-old flies. When we performed experiments at 25 degrees, we saw much less
optomotor turning, but anti-directional turning persisted. Rearing on molasses-based
media or in the dark did not have strong effects on behavior. N = 22, 8, 12, 12, 24, 19, 19,
13 flies top to bottom, left to right.

c) Other changes to the experimental setup did not cause large differences in behavior. We
compared responses to high contrast stimuli presented with green light (peak wavelength:
525nm) and blue light (peak wavelength: 450), and did not see large differences in
behavior. Head-fixed flies (middle) showed similar behavior to head-free flies (**a**, *top*). N
= 5 and 11 flies, top to bottom.

Supplementary Figure S3. Anti-directional turning behavior occurs when using the optical
filters also employed in the two-photon imaging experiments. High and low contrast
sinusoidal stimuli were presented as in Figure 2c, but using the bandpass filters also used in our
two-photon microscope stimulus presentation. N = 30 flies.

Supplementary Figure S4. *D. yakuba* lacks plasticity of anti-directional responses in adulthood that is observed *D. melanogaster*.

    a) Adult *yakuba* flies at various ages post eclosion were presented with 5-second, high-contrast, rotating sinusoidal gratings as in Fig. 6. Data was acquired in the TRC lab. Anti-directional responses stayed consistent from 0.5 days post eclosion (dpe) to 1, 2, and 4 dpe, although the initial optomotor response became smaller as the flies aged. Shaded patches represent ±1 SEM. N = 7-11 flies.

    b) The last 1.5 seconds of the mean turning velocity of each fly was averaged, and the population response was plotted.

Citations

1. Hassenstein, B., and Reichardt, W. (1956). Systemtheoretische Analyse der Zeit-, Reihenfolgen-und Vorzeichenauswertung bei der Bewegungsperzeption des Rüsselkäfers Chlorophanus. Zeits. Naturforsch. *11*, 513–524.

2. Götz, K., and Wenking, H. (1973). Visual control of locomotion in the walking fruitfly Drosophila. J. Comp. Physiol. A *85*, 235-266.

3. Buchner, E. (1976). Elementary movement detectors in an insect visual system. Biol. Cybern. *24*, 85-101.

4. Clark, D. (1981). Visual responses in developing zebrafish. Brachydanio rerio.

5. Koerner, F., and Schiller, P.H. (1972). The optokinetic response under open and closed loop conditions in the monkey. Exp. Brain Res. *14*, 318-330.

6. Götz, K.G. (1975). The optomotor equilibrium of theDrosophila navigation system. Journal of comparative physiology *99*, 187-210. 10.1007/BF00613835.

7. Yang, H.H., and Clandinin, T.R. (2018). Elementary motion detection in Drosophila: algorithms and mechanisms. Ann. Rev. Vis. Sci. *4*, 143-163.

8. Clark, D.A., Bursztyn, L., Horowitz, M.A., Schnitzer, M.J., and Clandinin, T.R. (2011). Defining the computational structure of the motion detector in Drosophila. Neuron *70*, 1165-1177.

9. Joesch, M., Schnell, B., Raghu, S., Reiff, D., and Borst, A. (2010). ON and OFF pathways in Drosophila motion vision. Nature *468*, 300-304.

10. Strother, J.A., Nern, A., and Reiser, M.B. (2014). Direct observation of ON and OFF pathways in the Drosophila visual system. Curr. Biol. *24*, 976-983.

11. Behnia, R., Clark, D.A., Carter, A.G., Clandinin, T.R., and Desplan, C. (2014). Processing properties of ON and OFF pathways for Drosophila motion detection. Nature *512*, 427-430.

12. Maisak, M.S., Haag, J., Ammer, G., Serbe, E., Meier, M., Leonhardt, A., Schilling, T., Bahl, A., Rubin, G.M., Nern, A., et al. (2013). A directional tuning map of Drosophila elementary motion detectors. Nature *500*, 212-216.

13. Takemura, S.-y., Bharioke, A., Lu, Z., Nern, A., Vitaladevuni, S., Rivlin, P.K., Katz, W.T., Olbris, D.J., Plaza, S.M., Winston, P., et al. (2013). A visual motion detection circuit suggested by Drosophila connectomics. Nature *500*, 175-181.

14. Shinomiya, K., Huang, G., Lu, Z., Parag, T., Xu, C.S., Aniceto, R., Ansari, N., Cheatham, N., Lauchie, S., Neace, E., et al. (2019). Comparisons between the ON-and OFF-edge motion pathways in the Drosophila brain. eLife *8*, e40025.

15. Bausenwein, B., Dittrich, A., and Fischbach, K.-F. (1992). The optic lobe of Drosophila
melanogaster. Cell Tissue Res. *267*, 17-28.

16. Henning, M., Ramos-Traslosheros, G., Gür, B., and Silies, M. (2022). Populations of
local direction–selective cells encode global motion patterns generated by self-motion.
Science advances *8*, eabi7112.

17. Mauss, A.S., Pankova, K., Arenz, A., Nern, A., Rubin, G.M., and Borst, A. (2015).
Neural circuit to integrate opposing motions in the visual field. Cell *162*, 351-362.

18. Schnell, B., Raghu, S.V., Nern, A., and Borst, A. (2012). Columnar cells necessary for
motion responses of wide-field visual interneurons in Drosophila. J. Comp. Physiol. A
198, 389-395.

19. Barnhart, E.L., Wang, I.E., Wei, H., Desplan, C., and Clandinin, T.R. (2018). Sequential
nonlinear filtering of local motion cues by global motion circuits. Neuron *100*, 229-243.
e223.

20. Joesch, M., Plett, J., Borst, A., and Reiff, D. (2008). Response properties of motion-
sensitive visual interneurons in the lobula plate of Drosophila melanogaster. Curr. Biol.
18, 368-374.

21. Kim, A.J., Fenk, L.M., Lyu, C., and Maimon, G. (2017). Quantitative predictions
orchestrate visual signaling in Drosophila. Cell *168*, 280-294. e212.

22. Haikala, V., Joesch, M., Borst, A., and Mauss, A.S. (2013). Optogenetic control of fly
optomotor responses. J. Neurosci. *33*, 13927-13934.

23. Krapp, H.G., and Hengstenberg, R. (1996). Estimation of self-motion by optic flow
processing in single visual interneurons. Nature *384*, 463-466.

24. Götz, K.G. (1970). Fractionation of *Drosophila* Populations According to Optomotor
Traits. Journal of Experimental Biology *52*, 419-436.

25. Götz, K. (1964). Optomotorische untersuchung des visuellen systems einiger
augenmutanten der fruchtfliege Drosophila. Biol. Cybern. *2*, 77-92.

26. Tammero, L., Frye, M., and Dickinson, M. (2004). Spatial organization of visuomotor
reflexes in Drosophila. J. Exp. Biol. *207*, 113-122.

27. Goetz, K.G. (1968). Flight control in Drosophila by visual perception of motion. Biol.
Cybern. *4*, 199-208.

28. Williamson, W.R., Peek, M.Y., Breads, P., Coop, B., and Card, G.M. (2018). Tools for
Rapid High-Resolution Behavioral Phenotyping of Automatically Isolated Drosophila.
Cell Rep. *25*, 1636-1649. e1635.

29.   Wolf, R., and Heisenberg, M. (1990). Visual control of straight flight in Drosophila
719   melanogaster. Journal of Comparative Physiology A: Neuroethology, Sensory, Neural,
720   and Behavioral Physiology *167*, 269-283.

721   30.   Bahl, A., Ammer, G., Schilling, T., and Borst, A. (2013). Object tracking in motion-blind
722   flies. Nat. Neurosci. *16*, 730-738.

723   31.   Seelig, J., Chiappe, M., Lott, G., Dutta, A., Osborne, J., Reiser, M., and Jayaraman, V.
724   (2010). Two-photon calcium imaging from head-fixed Drosophila during optomotor
725   walking behavior. Nat. Methods.

726   32.   Bosch, D.S., van Swinderen, B., and Millard, S.S. (2015). Dscam2 affects visual
727   perception in Drosophila melanogaster. Frontiers in behavioral neuroscience *9*, 149.

728   33.   Rister, J., Pauls, D., Schnell, B., Ting, C., Lee, C., Sinakevitch, I., Morante, J., Strausfeld,
729   N., Ito, K., and Heisenberg, M. (2007). Dissection of the peripheral motion channel in the
730   visual system of Drosophila melanogaster. Neuron *56*, 155-170.

731   34.   Creamer, M.S., Mano, O., Tanaka, R., and Clark, D.A. (2019). A flexible geometry for
732   panoramic visual and optogenetic stimulation during behavior and physiology. J.
733   Neurosci. Methods *323*, 48-55.

734   35.   Bahl, A., Serbe, E., Meier, M., Ammer, G., and Borst, A. (2015). Neural mechanisms for
735   Drosophila contrast vision. Neuron *88*, 1240-1252.

736   36.   Strother, J.A., Wu, S.-T., Wong, A.M., Nern, A., Rogers, E.M., Le, J.Q., Rubin, G.M.,
737   and Reiser, M.B. (2017). The emergence of directional selectivity in the visual motion
738   pathway of Drosophila. Neuron *94*, 168-182. e110.

739   37.   Strother, J.A., Wu, S.-T., Rogers, E.M., Eliason, J.L., Wong, A.M., Nern, A., and Reiser,
740   M.B. (2018). Behavioral state modulates the ON visual motion pathway of Drosophila.
741   Proc. Natl. Acad. Sci. USA *115*, E102-E111.

742   38.   Clark, D.A., Fitzgerald, J.E., Ales, J.M., Gohl, D.M., Silies, M., Norcia, A.M., and
743   Clandinin, T.R. (2014). Flies and humans share a motion estimation strategy that exploits
744   natural scene statistics. Nat. Neurosci. *17*, 296-303.

745   39.   Creamer, M.S., Mano, O., and Clark, D.A. (2018). Visual Control of Walking Speed in
746   Drosophila. Neuron *100*, 1460-1473.

747   40.   Salazar-Gatzimas, E., Chen, J., Creamer, M.S., Mano, O., Mandel, H.B., Matulis, C.A.,
748   Pottackal, J., and Clark, D.A. (2016). Direct measurement of correlation responses in
749   Drosophila elementary motion detectors reveals fast timescale tuning. Neuron *92*, 227-
750   239.

751   41.   Leonhardt, A., Ammer, G., Meier, M., Serbe, E., Bahl, A., and Borst, A. (2016).
752   Asymmetry of Drosophila ON and OFF motion detectors enhances real-world velocity
753   estimation. Nat. Neurosci. *19*, 706–715.

42. Silies, M., Gohl, D.M., Fisher, Y.E., Freifeld, L., Clark, D.A., and Clandinin, T.R.
(2013). Modular use of peripheral input channels tunes motion-detecting circuitry.
Neuron *79*, 111-127.

43. Heisenberg, M., and Buchner, E. (1977). The role of retinula cell types in visual behavior
ofDrosophila melanogaster. Journal of Comparative Physiology A: Neuroethology,
Sensory, Neural, and Behavioral Physiology *117*, 127-162.

44. McCann, G.D., and MacGinitie, G. (1965). Optomotor response studies of insect vision.
Proceedings of the Royal Society of London. Series B. Biological Sciences *163*, 369-401.

45. Duistermars, B., Chow, D., Condro, M., and Frye, M. (2007). The spatial, temporal and
contrast properties of expansion and rotation flight optomotor responses in Drosophila. J.
Exp. Biol. *210*, 3218.

46. York, R.A., Brezovec, L.E., Coughlan, J., Herbst, S., Krieger, A., Lee, S.-Y., Pratt, B.,
Smart, A.D., Song, E., and Suvorov, A. (2022). The evolutionary trajectory of
drosophilid walking. Curr. Biol.

47. Arenz, A., Drews, M.S., Richter, F.G., Ammer, G., and Borst, A. (2017). The temporal
tuning of the Drosophila motion detectors is determined by the dynamics of their input
elements. Curr. Biol. *27*, 929-944.

48. Leong, J.C.S., Esch, J.J., Poole, B., Ganguli, S., and Clandinin, T.R. (2016). Direction
selectivity in Drosophila emerges from preferred-direction enhancement and null-
direction suppression. J. Neurosci. *36*, 8078-8092.

49. Wienecke, C.F., Leong, J.C., and Clandinin, T.R. (2018). Linear Summation Underlies
Direction Selectivity in Drosophila. Neuron.

50. Salazar-Gatzimas, E., Agrochao, M., Fitzgerald, J.E., and Clark, D.A. (2018). The
Neuronal Basis of an Illusory Motion Percept Is Explained by Decorrelation of Parallel
Motion Pathways. Curr. Biol. *28*, 3748-3762.

51. Schilling, T., and Borst, A. (2015). Local motion detectors are required for the
computation of expansion flow-fields. Biology open, bio. 012690.

52. Kitamoto, T. (2001). Conditional modification of behavior in Drosophila by targeted
expression of a temperature-sensitive shibire allele in defined neurons. J. Neurobiol. *47*,
81-92.

53. Fujiwara, T., Brotas, M., and Chiappe, M.E. (2022). Walking strides direct rapid and
flexible recruitment of visual circuits for course control in Drosophila. Neuron.

54. Wei, H., Kyung, H.Y., Kim, P.J., and Desplan, C. (2020). The diversity of lobula plate
tangential cells (LPTCs) in the Drosophila motion vision system. J. Comp. Physiol. A
206, 139-148.

55. Borst, A., and Weber, F. (2011). Neural Action Fields for Optic Flow Based Navigation:
A Simulation Study of the Fly Lobula Plate Network. PLOS ONE *6*, e16303.
10.1371/journal.pone.0016303.

56. Eckert, H., and Dvorak, D.R. (1983). The centrifugal horizontal cells in the lobula plate
of the blowfly, Phaenicia sericata. J. Insect Physiol. *29*, 547-560.

57. Juusola, M., Dau, A., Song, Z., Solanki, N., Rien, D., Jaciuch, D., Dongre, S.A.,
Blanchard, F., de Polavieja, G.G., and Hardie, R.C. (2017). Microsaccadic sampling of
moving image information provides Drosophila hyperacute vision. Elife *6*, e26117.

58. Katsov, A., and Clandinin, T. (2008). Motion processing streams in Drosophila are
behaviorally specialized. Neuron *59*, 322-335.

59. Rivlin-Etzion, M., Wei, W., and Feller, M.B. (2012). Visual stimulation reverses the
directional preference of direction-selective retinal ganglion cells. Neuron *76*, 518-525.

60. Vlasits, A.L., Bos, R., Morrie, R.D., Fortuny, C., Flannery, J.G., Feller, M.B., and Rivlin-
Etzion, M. (2014). Visual stimulation switches the polarity of excitatory input to starburst
amacrine cells. Neuron *83*, 1172-1184.

61. Matulis, C.A., Chen, J., Gonzalez-Suarez, A., Behnia, R., and Clark, D.A. (2020).
Heterogeneous temporal contrast adaptation in Drosophila direction-selective circuits.
Curr. Biol. *30*, 222-236.

62. Drews, M.S., Leonhardt, A., Pirogova, N., Richter, F.G., Schuetzenberger, A., Braun, L.,
Serbe, E., and Borst, A. (2020). Dynamic Signal Compression for Robust Motion Vision
in Flies. Curr. Biol.

63. Frazor, R.A., and Geisler, W.S. (2006). Local luminance and contrast in natural images.
Vision Res. *46*, 1585-1598.

64. Kiral, F.R., Dutta, S.B., Linneweber, G.A., Hilgert, S., Poppa, C., Duch, C., von Kleist,
M., Hassan, B.A., and Hiesinger, P.R. (2021). Brain connectivity inversely scales with
developmental temperature in Drosophila. Cell Rep. *37*, 110145.

65. Mongeau, J.-M., and Frye, M.A. (2017). Drosophila spatiotemporally integrates visual
signals to control saccades. Curr. Biol. *27*, 2901-2914. e2902.

66. Ketkar, M.D., Sporar, K., Gür, B., Ramos-Traslosheros, G., Seifert, M., and Silies, M.
(2020). Luminance information is required for the accurate estimation of contrast in
rapidly changing visual contexts. Curr. Biol. *30*, 657-669. e654.

67. Tuthill, J.C., Nern, A., Holtz, S.L., Rubin, G.M., and Reiser, M.B. (2013). Contributions
of the 12 neuron classes in the fly lamina to motion vision. Neuron *79*, 128-140.

68. Chiappe, M.E., Seelig, J.D., Reiser, M.B., and Jayaraman, V. (2010). Walking modulates
speed sensitivity in Drosophila motion vision. Curr. Biol. *20*, 1470-1475.

824 69. DeAngelis, B.D., Zavatone-Veth, J.A., and Clark, D.A. (2019). The manifold structure of
825   limb coordination in walking Drosophila. eLife *8*, e46409.

826 70. Katsov, A.Y., Freifeld, L., Horowitz, M.A., Kuehn, S., and Clandinin, T.R. (2017).
827   Dynamic structure of locomotor behavior in walking fruit flies. eLife *6*, e26410.

828 71. Rust, N.C., Mante, V., Simoncelli, E.P., and Movshon, J.A. (2006). How MT cells
829   analyze the motion of visual patterns. Nat. Neurosci. *9*, 1421-1431.

830 72. Gohl, D.M., Silies, M.A., Gao, X.J., Bhalerao, S., Luongo, F.J., Lin, C.C., Potter, C.J.,
831   and Clandinin, T.R. (2011). A versatile in vivo system for directed dissection of gene
832   expression patterns. Nat. Methods *8*, 231-237.

833 73. Kleiner, M., Brainard, D., Pelli, D., Ingling, A., Murray, R., and Broussard, C. (2007).
834   What's new in Psychtoolbox-3. Perception *36*, 1.

835 74. Turner, M.H., Krieger, A., Pang, M.M., and Clandinin, T.R. (2022). Visual and motor
836   signatures of locomotion dynamically shape a population code for visual features in
837   Drosophila. bioRxiv.

838 75. Moore, R.J., Taylor, G.J., Paulk, A.C., Pearson, T., van Swinderen, B., and Srinivasan,
839   M.V. (2014). FicTrac: a visual method for tracking spherical motion and generating
840   fictive animal paths. J. Neurosci. Methods *225*, 106-119.

841 76. Tanaka, R., and Clark, D.A. (2022). Neural mechanisms to exploit positional geometry
842   for collision avoidance. Curr. Biol. *32*, 2357-2374.

843 77. Pologruto, T.A., Sabatini, B.L., and Svoboda, K. (2003). ScanImage: flexible software
844   for operating laser scanning microscopes. Biomed. Eng. Online *2*, 13.

845 78. Agrochao, M., Tanaka, R., Salazar-Gatzimas, E., and Clark, D.A. (2020). Mechanism for
846   analogous illusory motion perception in flies and humans. Proc. Natl. Acad. Sci. *117*,
847   23044-23053.