Long timescale anti-directional rotation in *Drosophila* optomotor behavior

Omer Mano[1], Minseung Choi[2], Ryosuke Tanaka[3], Matthew S. Creamer[3], Natalia C.B. Matos[3],
Joseph Shomar[4], Bara A. Badwan[5], Thomas R. Clandinin[2], Damon A. Clark[1,3,4,6,#]

1 – Department of Molecular, Cellular, and Developmental Biology, Yale University, New
Haven, CT 06511, USA

2 – Department of Neurobiology, Stanford University, Stanford, CA 94305, USA

3 – Interdepartmental Neuroscience Program, Yale University, New Haven, CT 06511, USA

4 – Department of Physics, Yale University, New Haven, CT 06511, USA

5 – Department of Chemical Engineering, Yale University, New Haven, CT 06511, USA

6 – Department of Neuroscience, Yale University, New Haven, CT 06511, USA

– Lead author: damon.clark@yale.edu

Abstract
Locomotor movements cause visual images to be displaced across the eye, a retinal slip that is
counteracted by stabilizing reflexes in many animals. In insects, optomotor turning causes the
animal to turn in the direction of rotating visual stimuli, thereby reducing retinal slip and
stabilizing trajectories through the world. This behavior has formed the basis for extensive
dissections of motion vision. Here, we report that under certain stimulus conditions, two
Drosophila species, including the widely studied *D. melanogaster*, can suppress and even
reverse the optomotor turning response over several seconds. Such 'anti-directional turning' is
most strongly evoked by long-lasting, high-contrast, slow-moving visual stimuli that are distinct
from those that promote syn-directional optomotor turning. Anti-directional turning, like the syn-
directional optomotor response, requires the local motion detecting neurons T4 and T5; a subset
of lobula plate tangential cells, CH cells, show involvement in these responses. Imaging from a
variety of direction-selective cells in the lobula plate shows no evidence of dynamics that match
the behavior, suggesting that the observed inversion in turning direction emerges downstream of
the lobula plate. Further, anti-directional turning declines with age and exposure to light. These
results show that *Drosophila* optomotor turning behaviors contain rich, stimulus-dependent
dynamics that are inconsistent with simple reflexive stabilization responses.

Intro
Visual navigation requires active mechanisms to stabilize trajectories through the world. Insects
exhibit an optomotor turning response, a behavior in which they rotate their bodies in the
direction of visual patterns that rotate about them (Buchner, 1976; Götz and Wenking, 1973;

Hassenstein and Reichardt, 1956). This behavior is analogous to optomotor turning responses in
fish (Clark, 1981) and the optokinetic response in mammals (Koerner and Schiller, 1972). In
insects, this response is thought to be a course-stabilization mechanism that minimizes retinal
slip, allowing animals to maintain their trajectory in the face of external or unexpected rotational
forces (Götz and Wenking, 1973; Götz, 1975). For instance, if an insect attempts to walk in a
straight line, it may slip and turn to the right. From the point of view of the insect, this turn is
observed as optic flow rotating to the left. By responding to this leftward optic flow with a
leftward turn, the insect can recover its original trajectory. The optokinetic response, similarly,
acts to stabilize eye position relative to the visual scene, even as the head rotates (Schweigart et
al., 1997).

In fruit flies, the optomotor response relies on well-characterized circuitry (Yang and Clandinin,
2018). Photoreceptor signals are split into parallel ON and OFF pathways in the lamina and
medulla (Behnia et al., 2014; Clark et al., 2011; Joesch et al., 2010; Strother et al., 2014), that are
not direction-selective. These signals provide input to T4 and T5 cells, which compute direction-
selective responses along four directions for every point in the fly visual field (Bausenwein et al.,
1992; Henning et al., 2022; Maisak et al., 2013; Shinomiya et al., 2019; Takemura et al., 2013).
The outputs of T4 and T5 cells are then summed across visual space by lobula plate tangential
cells (LPTCs) (Barnhart et al., 2018; Joesch et al., 2008; Maisak *et al.*, 2013; Mauss et al., 2015;
Schnell et al., 2012). Different LPTCs provide distinct signals about the overall pattern of motion
surrounding the fly, and have been linked to head and body movements (Haikala et al., 2013;
Kim et al., 2017; Krapp and Hengstenberg, 1996).

Interestingly, there have been several reports of flies turning in the direction opposite to what
would be predicted from the optomotor turning response. In some cases, these counter-intuitive
behaviors were observed using periodic stimuli with spatial wavelengths smaller than the
receptive field of individual ommatidia, and thus can be accounted for by aliasing (Buchner,
1976; Götz, 1964; Götz, 1970). Work in a tethered flight simulator showed that when a moving
pattern is presented in front of the fly, the animal turned in the direction of the stimulus motion
(Tammero et al., 2004), as expected (Goetz, 1968). However, if the moving pattern was
presented behind the fly, it attempted to turn in the direction opposite to stimulus motion
(Tammero *et al.*, 2004). In a different experimental preparation, rotational patterns were
presented on a dome around freely-walking flies (Williamson et al., 2018). Under these
conditions, flies generally turned in the direction of motion of the stimulus, but these rotations
were often punctuated by brief, large-magnitude saccades in the opposite direction. Similarly,
experiments using flight simulators have reported spikes in the torque in the direction opposite
the stimulus rotation (Wolf and Heisenberg, 1990).

Here we show that rotational stimuli can elicit strong, consistent anti-directional behavior in two
drosophilid species, *D. melanogaster* and *D. yakuba*. We report that flies respond to high
contrast, high luminance rotational motion stimuli by first turning in the direction of stimulus
motion, and then reversing their trajectory after approximately one second, depending on the
species. In *Drosophila melanogaster*, we characterize the dynamics of this behavior and the
stimuli that drive it. The behavior depends critically on adaptation to back-to-front motion. We

use the genetic tools available in *Drosophila melanogaster* to show that this behavior relies on the motion detecting neurons T4 and T5. Silencing HS and CH, two widefield neurons downstream of T4 and T5, resulted in small changes in this complex turning behavior. However, the visually evoked responses of these direction-selective neurons could not account for these behaviors. Thus, behavioral reversal must be mediated by more downstream circuitry. Overall, these results show that flies generate behavioral signals that oppose the direction of visual motion, showing that *Drosophila* turning responses to wide-field visual motion stimuli are more complex than a simple stabilizing reflex.

Results

Anti-directional turning responses to high contrast stimuli

Optomotor turning responses are central for gaze stabilization, so we sought to examine the stability of this response across different conditions. Many studies have investigated this behavior using low contrast and low light intensity stimuli or both (Bahl et al., 2013; Bosch et al., 2015; Buchner, 1976; Götz and Wenking, 1973; Rister et al., 2007; Seelig et al., 2010), at a variety of different speeds. However, natural scenes can have relatively high contrast and luminance, conditions have been poorly explored in the laboratory. In this experiment, we presented flies with rotational stimuli using high contrast and relatively high luminance.

We tethered individual female *D. melanogaster* above a freely rotating ball to characterize the optomotor response (Buchner, 1976; Creamer et al., 2019) (**Fig. 1a**). As expected, low contrast, slow-moving sinusoidal gratings caused flies to turn in the same direction as the moving gratings via the classical optomotor turning response (**Fig. 1b**) (Bahl *et al.*, 2013; Bahl et al., 2015; Buchner, 1976; Clark *et al.*, 2011; Clark et al., 2014; Creamer et al., 2018; Götz, 1964; Hassenstein and Reichardt, 1956; Leonhardt et al., 2016; Salazar-Gatzimas et al., 2016; Seelig *et al.*, 2010; Silies et al., 2013; Strother et al., 2018; Strother et al., 2017; Tammero *et al.*, 2004). However, when we changed the stimulus to high contrast sinusoidal gratings (nominal 100% Weber contrast), flies turned in the stimulus direction for approximately 1 second, but then reversed course, and turned in the direction opposite to the stimulus motion for the duration of the stimulus presentation. Because this turning response is in the opposite direction of stimulus and the syn-directional optomotor turning response, we refer to it as anti-directional turning.

We swept a range of contrasts and compared the fly turning in the first 500 milliseconds to the turning after one second (**Fig. 1c**). As contrast increased, the flies turned faster during the first half second of stimulus presentation, reaching a plateau at around 0.5 contrast, consistent with previous results (Bahl *et al.*, 2015; Buchner, 1976; Duistermars et al., 2007; Heisenberg and Buchner, 1977; McCann and MacGinitie, 1965; Strother *et al.*, 2017). Fly behavior after the first second of stimulation was more complex. As contrast increased from 0 to 0.25, flies turned in the same direction as the stimulus, with faster turning as the contrast increased. When the contrast was greater than 0.25, turning decreased, lowering to no net sustained turning at around 0.8 contrast. Above a contrast of 0.8, flies began to turn in the direction opposite the stimulus.

These initial experiments took place in the lab of author DAC. To confirm that these unexpected responses did not reflect some idiosyncrasy of one specific behavioral apparatus or environment, we repeated these experiments in a second lab, that of author TRC. Under similar conditions,

using the same strain of *Drosophila melanogaster*, we reproduced the rapid deceleration after an
initial, transient syn-directional response (**Fig. 1d**), with some individual flies exhibiting
significant anti-directional turning, as in the experiments in the first lab (**Supp. Fig. S1**). This
demonstrates that the key features of this behavioral response are stable across experimental
systems and laboratories, though the magnitude of reverse-turning behavior in *D. melanogaster*
is sensitive to some unknown experimental parameter differences between the laboratories.

Individual strains of *D. melanogaster*, and other drosophilid species, display significant variation
in their locomotor patterns during walking (York et al., 2022). Indeed, when we tested a Canton-
S *D. melanogaster* strain, we observed mild but significant anti-directional turning at long
timescales (**Supp. Fig. S2b**). We reasoned that a strong test of the generality of anti-directional
turning would be to examine turning behavior in another species, and selected *D. yakuba*.
Strikingly, *D. yakuba* also displayed anti-directional turning behavior under similar conditions
(**Fig. 1e**). Thus, this behavior is not an idiosyncratic feature of a single laboratory strain.

Figure 1. Flies turn opposite to the stimulus direction in high contrast conditions

a) We measured fly turning behavior as they walked on an air-suspended ball. Stimuli were presented over 270 degrees around the fly.

b) We presented drifting sinusoidal gratings for 5 seconds (shaded region) with either high contrast (c = 1.0) or low contrast (c = 0.25). When high contrast sinusoidal gratings were presented, flies initially turned in the same direction as the stimulus, then started turning

in the opposite direction after ~1 second of stimulation. Under low contrast conditions,
flies turned continuously in the same direction as the stimulus. In these experiments, the
sine waves had a wavelength of 60º and a temporal frequency of 1 Hz. Shaded patches
represent ±1 SEM. N= 10 flies.

c) We swept contrast between 0 and 1 and measured the mean turning response during the
first 0.5 seconds (purple, purple bar in **b**) and during the last 4 seconds of the stimulus
(brown, brown line in **b**). The response in the first 0.5 seconds increased with increasing
contrast, while the response in the last four seconds increased from c = 0 to c = 0.25, and
then decreased with increasing contrast, until flies turned in the direction opposite the
stimulus direction at the highest contrasts. N = 20 flies.

d) We repeated the presentation of drifting sinusoidal gratings, this time in the lab of author
TRC, using a similar behavioral apparatus. Stimulus parameters were as described in (b).
In these experiments, the population average shows that flies proceeded to zero net
turning at high contrasts, but some individual flies exhibited anti-directional turning
responses. N = 20 flies.

e) We repeated the experiments with *D. yakuba*, also in the lab of TRC, and observed that
this species exhibited a robust anti-directional turning response to high contrast gratings
and a classical syn-directional turning response to low contrast gratings. N = 11 flies.

Conditions for anti-directional turning behaviors

While anti-directional turning behaviors have been reported before, other groups have presented
similar stimuli without observing anti-directional behavior (Bahl *et al.*, 2013; Bosch *et al.*, 2015;
Buchner, 1976; Götz and Wenking, 1973; Seelig *et al.*, 2010). We wondered what aspects of our
experimental setup could lead to these behavioral differences. In our experiments, anti-
directional turning was strongly linked to display brightness (**Supp. Fig. S2a**). When the mean
brightness of the screens was reduced from 100 cd/m^2 to 1 cd/m^2, we saw no anti-directional
turning in 5 second trials (though average optomotor behavior did decrease over the course of the
stimulus presentation). When we further reduced the mean brightness to 0.1 cd/m^2, flies persisted
in their optomotor behavior throughout the stimulus presentation. We note, however, that at low
luminance, low levels of ambient light in the nominally dark experimental rig could also reduce
the effective contrast of the stimulus.

We tested a variety of other factors that might affect anti-directional turning. Anti-directional
turning occurred when experiments were run both at hot temperatures and at room temperature
(**Supp. Fig. S2b**). We also observed anti-directional behavior when flies were reared in the dark
and on different media. We also tested a number of other experiment conditions (**Supp. Fig.
S2c**). Flies responded with anti-directional turning to high contrast stimuli presented at both blue
and green wavelengths. We glued fly heads to their thorax to ensure stimuli could not be affected
by head movements (Haikala *et al.*, 2013; Kim *et al.*, 2017), but found no difference between
head-fixed and head-free flies. There were, however, a few factors that did modulate anti-
directional turning behavior. In particular, rearing *D. melanogaster* at 25°C instead of 20°C or
testing flies that were two weeks old instead of 12-60 hours old both reduced overall turning

behavior and eliminated anti-directional turning. In these cases, optomotor turning still decreased
over the course of the 5 second, high contrast trials, but did not reverse. As details of rearing
temperature and the age at which behavior tests are run often vary across labs, it is likely that
these factors account for the differences between our observations and the previous literature.

Distinct spatiotemporal tuning of the anti-directional behavioral response
To further characterize the anti-directional response, we swept the spatial and temporal
frequency of the sinusoidal grating stimulus. Using only Weber contrasts of 1, we compared the
early response (first quarter second, **Fig. 2a**) to the late response (after one second, **Fig. 2b**).
Drosophila melanogaster always turned in the optomotor direction during the early stimulus
response. In this early response, flies turned most vigorously to stimuli with short spatial
frequencies (~20º wavelength) and fast temporal frequencies (~8 Hz), in agreement with earlier
studies (Creamer *et al.*, 2018; Strother *et al.*, 2018; Tammero *et al.*, 2004). However, during the
long-term response to high-contrast stimuli, flies only turned in the optomotor direction at very
high temporal frequencies (> ~16 Hz) and at very low temporal frequencies (<0.5 Hz). At
intermediate temporal frequencies, flies showed a sustained anti-directional response. The
maximal anti-directional response was achieved at 1 Hz and 45º wavelength, distinct from the
conditions for peak classical turning responses. Interestingly, the stimuli that elicit the strongest
anti-directional response appear similar to those that maximally activate T4 and T5 neurons
when those neurons are measured in head-fixed flies (Arenz et al., 2017; Creamer *et al.*, 2018;
Leong et al., 2016; Maisak *et al.*, 2013; Strother *et al.*, 2018; Wienecke et al., 2018).

Anti-directional turning results from adaptation effects
We were intrigued by the switch from syn-directional to anti-directional turning behavior. To
investigate the dynamics of these changes, we presented a rotating sinusoidal stimulus at contrast
1 for five seconds, and then changed the contrast to 0.25 (**Fig. 2c**). After the switch to low
contrast, the flies quickly reverted classical, syn-directional optomotor behavior, demonstrating
that no long-term switch in directional turning occurs during high contrast stimulus presentation.
This effect did not depend on the periodic nature of these stimuli; a rotating stimulus consisting
of 5º-wide vertical bars with randomly-chosen, binary contrasts (Clark *et al.*, 2014) yielded
similar behavioral responses (**Fig. 2d**).

To further isolate the causes of this switch in behavior, we developed a stimulus to adapt the fly
to different stimuli before presenting high-contrast rotational sinusoidal gratings to elicit the anti-
directional turning response. This adapting stimulus consisted of five seconds of high contrast
'translational' stimuli, which was then followed by a rotational stimulus (**Fig. 2e**). The
translational stimuli consisted of both left and right hemifields moving either front-to-back or
back-to-front across the fly's two eyes (Creamer *et al.*, 2018). These stimuli resulted in no net
turning by the flies (Creamer *et al.*, 2018; Silies *et al.*, 2013). Adapting the fly with front-to-back
stimuli did not have a strong effect on the subsequent response to rotational stimuli. However,
adapting with back-to-front stimuli generated responses that no longer showed an initial syn-

directional turning response, but instead exhibited anti-directional turning immediately after the rotational stimulus began. This result indicates that the anti-directional turning results from slow-timescale changes that depend on strong back-to-front motion stimulation.

Figure 2. Anti-directional turning behavior has distinct tuning and is driven by adaptation.

a) Heatmap of fly turning velocity during the first 0.5 seconds of sinusoidal grating stimulation under high contrast conditions and variable temporal and spatial frequencies. The flies turned in the direction of the stimulus across all conditions and responded most to 8 Hz, 22-degree stimuli. N = 16,21,17,21,7, and 22 flies for spatial frequencies 1/120, 1/90, 1/60, 1/45, 1/30 and 1/22 degrees respectively.

b) Heatmap as in (**a**), measured during the last four seconds of stimulation. Flies turned in the same direction as the stimulus at high and low temporal frequencies, but in the opposite direction of the stimulus at intermediate temporal frequencies, with a maximal anti-directional response at wavelengths between 30º and 60º.

c) Switching stimulus contrast from high to low after 5 seconds caused flies to revert to syn-directional behavior after the anti-directional response. N = 7 flies.

d) Presenting rotating random binary patterns (5-degree vertical strips rotating at 150 degrees/second) induced anti-directional turning similar to that elicited by rotating sine wave gratings. N = 7 flies.

e) We presented flies with five seconds of "translational" stimuli (dark shaded region), with high contrast sinusoidal gratings moving either front-to-back or back-to-front, bilaterally, for five seconds. After that, we presented high contrast rotational sinusoidal grating stimuli (60º wavelength, 1 Hz). Front-to-back stimulation did not affect the subsequent response to rotational stimuli, but back-to-front stimuli caused flies to turn immediately in the opposite direction of the stimulus. N = 18 flies.

Anti-directional turning is elicited when stimuli are presented in front of the fly

A previous report of anti-directional turning behavior in flying tethered flies showed that flies turn in the opposite direction to stimuli that are presented behind their midline (Tammero *et al.*, 2004). To test whether our results were caused by this effect, we split our stimulus into three regions: 90 degrees in front of the fly, 45 degrees in front of the midline on either side of the fly, and 45 degrees behind the midline on either side of the fly (**Fig. 3a**). We found that flies displayed anti-directional turning when presented with stimuli only in the front region or only just in front of the midline (**Fig. 3bc**). They did not display anti-directional turning when moving stimuli were presented behind the midline (**Fig. 3bc**). This suggests a different mechanism from the behaviors that depend on posterior spatial location to elicit reverse-turning (Tammero *et al.*, 2004).

Anti-directional responses do not depend on saccades

Anti-directional saccades have been reported in walking and flying flies (Williamson *et al.*, 2018; Wolf and Heisenberg, 1990). In walking flies (Williamson *et al.*, 2018), flies largely turned in syn-directionally, but these turns were sometimes interrupted by brief, high-amplitude saccades in the opposite direction, against the stimulus direction. If such saccades were frequent or high amplitude, the net effect could shift the average turning we measured, creating apparent anti-directional turning. To investigate this possibility, we plotted the turning response on a per-trial basis (**Fig. 3d**). We then discarded information about the magnitude of the turns and considered only the direction of the turning at each point in time (**Fig. 3e**). Strikingly, in many trials, flies continued to turn opposite to the stimulus for several seconds, a behavior unlike brief saccades. We then calculated a turning index for each response timepoint (sampled at 60 Hz). This turning index represented the fraction of trials where the fly turned in the direction of the stimulus at each timepoint minus the fraction of trials where the fly turned in the opposite direction (**Fig. 3f**). Since this turning index does not include the magnitude of turning, it is strongly affected by sustained low-amplitude turns and discounts any brief high-amplitude saccades. When presented with high contrast stimuli, flies maintained a negative turning index, indicating that sustained turns, and not high velocity saccades, underlie this anti-directional turning behavior.

Figure 3. Anti-directional turning is driven by stimuli in the forward-facing visual field and is not driven by saccades.

a) We divided our panoramic display into three sections — the front 90º, the 45º behind the fly on either side, and a middle 45º.

b) High contrast sinusoidal gratings were presented on each of these three display sections, with the remaining sections blank. Flies turned syn-directionally when stimuli were presented behind the fly, and turned anti-directionally when stimuli were presented in front of the fly. Shaded patches represent ±1 SEM. N = 55 flies.

c) Average turning in the last 4 seconds of the stimulus (black bar in **b**), in low contrast and high contrast conditions. Shaded patches in the time trace plots represent ±1 SEM. N = 55 flies.

d) A single fly responds to many trials of sinusoidal grating stimuli at high contrast (blue bar) and low contrast (orange bar). We show a heatmap of the fly's responses over time (horizontal axis) and across trials (vertical axis).

e) We can ignore the magnitude of the turning and instead only quantify whether the fly was turning in the same direction as the stimulus (white area) or in the opposite direction (dark gray area). This shows sustained anti-directional turning, not brief saccades.

f) Averaging the direction (but not magnitude) of turning across trials and across flies yields a turning index for each point in time. Shaded patches in the time trace plots represent ±1 SEM. N = 7 flies.

Anti-directional turning requires elementary motion detectors

What neurons are involved in this anti-directional turning behavior? Previous work demonstrated that T4 and T5 are required for directional neural responses (Schnell *et al.*, 2012), as well as for optomotor turning (Maisak *et al.*, 2013; Salazar-Gatzimas et al., 2018; Salazar-Gatzimas *et al.*,

2016), for walking speed regulation (Creamer *et al.*, 2018), and for responses to visual looming stimuli (Schilling and Borst, 2015). We silenced the neurons T4 and T5 using shibire^ts (Kitamoto, 2001) and measured responses to sinusoidal stimuli that switched from high to low contrast (**Fig. 4a**). Flies in which T4 and T5 had been silenced displayed only minimal responses to motion stimuli, with anti-directional turning suppressed along with classical syn-directional turning. Thus, we conclude that, like optomotor turning behaviors, this anti-directional behavior depends critically on signals from T4 and T5.

Anti-directional turning requires the CH lobula plate tangential cell

Since the switch from optomotor to anti-directional behavior seems to be dependent on the direction of motion adaptation (**Fig. 2e**), we reasoned that neurons involved in this behavior were likely to be downstream from T4 and T5. Horizontal System (HS) cells are well-studied postsynaptic partners of T4 and T5 (Joesch *et al.*, 2008; Joesch *et al.*, 2010). These lobula plate tangential cells integrate information from front-to-back and back-to-front selective T4 and T5 cells across the fly's visual field (Mauss *et al.*, 2015). HS cells have been implicated in visually-evoked head turns (Kim *et al.*, 2017) and body rotations in flight (Haikala *et al.*, 2013), as well as in maintenance of walking direction (Fujiwara et al., 2022). When we silenced HS neurons, we found small deficits in syn-directional turning behavior, but not in anti-directional turning behavior (**Fig. 4b**), indicating that HS cells synaptic output is not required specifically for anti-directional turning behavior.

Next, we turned to the CH lobula plate tangential cells. These cells are GABAergic and are both pre-synaptic and post-synaptic in the lobula plate (Wei et al., 2020). In blowflies, these neurons play an inhibitory role in an interconnected LPTC circuit that shapes behavior (Borst and Weber, 2011). When we silenced these neurons, we found a small increase in syn-directional turning and a decrease in anti-directional turning (**Fig. 4c**). Overall, silencing this neuron type caused the flies to turn more in the direction of motion. This result suggests that CH activity contributes to the anti-directional turning response. However, since adapting to back-to-front translational stimuli significantly affected the dynamics of anti-directional turning, it seems likely that other neurons beyond HS and CH are involved, since these two neurons both respond selectively to front-to-back motion (Eckert and Dvorak, 1983; Joesch *et al.*, 2008).

Figure 4. Syn-directional and anti-directional turning share common circuitry

a) We silenced T4 and T5 neurons by expressing shibire^{ts} selectively in those neurons. We measured turning behavior during a contrast-switching stimulus (as in **Fig. 2c**). Results from flies with T4 and T5 silenced shown in dark red, while controls are in light red and gray. Average fly behavior during the last four seconds of the first contrast (black bar on left) shown as bars on the right, with individual fly behavior shown as dots. Note that the data labeled "low contrast" are from experiments in which the low-contrast stimulus was shown before the high contrast stimulus. Shaded patches in the time trace plots represent ±1 SEM, as do vertical lines on bar plots. *** indicates experimental results are significantly different from results, P < 0.001 via a two-sample Student t-test. * indicates P < 0.05. N = 17, 24, 19 flies with genotypes T4T5/Shibire^{ts}, T4T5/+, +/Shibire^{ts}.

b) Results from HS silencing as in **a**. Silencing HS reduced syn-directional turning behavior (P < 0.001) but did not have a strong effect on anti-directional turning. N = 34, 21, 19 flies with genotypes HS/Shibire^{ts}, HS/+, +/Shibire^{ts}.

c) Results from CH silencing as in **a**. CH silencing reduced the degree of anti-directional turning (P < 0.001). N = 63, 57, 70 flies with genotypes CH/Shibire^{ts}, CH/+, +/Shibire^{ts}.

Early direction-selective cells do not adapt to the stimulus

The anti-directional turning response is preceded by an initial syn-directional response. This change in behavior must be the result of changes in neural activity, but this change could happen

at any point along the neural pathway between photoreceptors and motor neurons. In order to
constrain possible mechanisms for generating the anti-directional turning behaviors, we used
calcium imaging to interrogate the activity of direction selective neurons during high and low
contrast stimulation (**Fig. 5a**). However, as calcium imaging experiments using two photon
microscopy require additional spectral filtering of the projector, we first confirmed that these
spectral differences did not alter anti-directional turning responses. To do this, we re-measuring
the anti-directional turning behavior using optical filtering matched to the conditions needed for
imaging. Using this spectrally distinct illuminant, we observed both syn-directional and anti-
directional turning behaviors, following the previously observed dynamics (**Supp. Fig. S3**).

As T4 and T5 neurons play a critical role in both the syn- and anti-directional turning responses,
we first measured the calcium activity of these neurons as they responded to sine wave gratings
at a range of contrasts in their preferred and null directions. The T4 and T5 neurons responded to
sine wave gratings in their preferred direction by increasing their calcium activity for the full
duration of the stimulus presentation, reaching a plateau after approximately 1 second (**Fig. 5bc**,
middle). As we increased the contrast of the preferred direction stimuli, we found that both T4
and T5 cells had increased calcium activity throughout the contrast range (**Fig. 5bc**, right),
consistent with prior measurements (Maisak *et al.*, 2013). Thus, the responses of T4 and T5 cells
do not capture the transition from syn-directional to anti-directional turning behavior.

Next we examined two LPTCs downstream of T4 and T5 cells. Calcium activity in HS cells
followed similar trends to T4 and T5. Calcium signals increased at the start of preferred direction
stimuli presentation and stayed high until the end of the presentation (**Fig. 5d**, middle).
Increasing contrast caused stronger calcium responses with a mild saturation effect at high
contrast (**Fig. 5d**, right), consistent with prior voltage measurements (Joesch *et al.*, 2008). These
results indicate that the changes in the time course of optomotor behavior at high contrast are not
related to changes in HS activity. Finally, we measured calcium activity in CH cells. CH cells
responded to visual stimuli more quickly than HS cells (**Fig. 5e**, middle), and showed decreased
calcium signals in response to null direction stimuli (**Fig. 5e**, right). However, they showed
sustained responses to high contrast stimuli, as in T4, T5, and HS. These measurements suggest
that the switch from syn- to anti-directional turning behavior is driven by cells downstream of or
parallel to T4, T5, HS, and CH.

Figure 5. Responses in early direction-selective cells do not show a reduction or reversal of response on the timescale of the behavior.

    a) We used two-photon microscopy to measure calcium activity in lobula plate neurons while presenting sinusoidal gratings at a range of contrasts.

    b) T4 cells, marked in orange (*left*), responded to drifting sinusoidal gratings with increased calcium activity (*middle*). Darker colors indicate higher contrast, preferred direction in blue, null direction in red. When integrated across the stimulus presentation (*right*), calcium activity increased with stimulus contrast. N = 8 flies.

    c-e) As in **b)** measuring calcium activity in T5, HS, and CH cells. N = 8, 10, 15 flies.

Adult plasticity in anti-directional turning behavior

In behaving flies, the strength of anti-directional turning was dependent both on rearing temperature, which alters the rate of growth, and on age (**Supp. Fig. S2**). This raises the possibility that syn- and anti-directional turning responses might be plastic during the early adult stages of development. To probe this possibility, we presented 1 Hz, high-contrast, rotating

sinusoidal grating at various stages during early adulthood (**Fig. 6**). Strikingly, as flies aged from
0.5 to 4 days post eclosion (dpe), the initial syn-directional turning became less transient and
more sustained, indicative of a weaker anti-directional turning drive. We then wondered whether
this plasticity was intrinsically programmed, or dependent on visual input. To explore this
possibility, we reared flies in darkness to 2 or 4 dpe and measured their turning responses (**Fig. 6**,
gray). Dark-reared flies exhibited a stronger deceleration away from syn-directional turning,
similar to that found in more juvenile flies, arguing that visual input may sculpt the balance of
syn- and anti-directional turning. Finally, we examined whether optomotor response plasticity
could be detected in *D. yakuba*. However, in this species, anti-directional responses were stable
across the first four days of adulthood, arguing that the role of visual experience in shaping these
responses is itself evolutionarily tuned in drosophilids (**Supp. Fig. 4**).

Figure 6. Maturation of optomotor response in early adulthood

    a) Adult flies at various ages post eclosion were presented with 5-second, high-contrast,
rotating sinusoidal gratings as in **Fig 2b**. As the flies aged from 1 day post eclosion (dpe)
to 2, 4, and 8 dpe, the initial anti-directional turning response transitioned into syn-
directional turning. Dark-rearing flies at 2 dpe reduced this maturation effect. Shaded
patches represent ±1 SEM. N = 5-14 flies.

416   **b)** The last 1.5 seconds of the mean turning velocity of each fly was averaged, and the
417   population response was plotted.

418   **c)** As in (a) but in the TRC lab, using 0.5, 1, 2, and 4 dpe, with dark rearing for 4 dpe. With
419   maturation, the syn-directional turning became less transient. N = 9-15 flies.

420   **d)** As in (b) but for data in (c).

Discussion

In this study, we found we could elicit robust turning in the opposite direction of high contrast motion stimuli (**Fig. 1**). This behavior is qualitatively different from other turning behaviors reported in the literature (**Figs. 2** and **3**), but shares elements with the circuitry necessary for optomotor behavior (**Fig. 4**). However, the switch from optomotor behavior to anti-directional turning behavior is not a reflection of changes in the activity of known direction-selective neuron types in the early visual system (**Fig. 5**). Moreover, this anti-directional turning behavior exhibits a degree of experience-dependent plasticity (**Fig. 6**).

Anti-directional turning is distinct from other against-stimuli behaviors

The anti-directional turning behavior we have characterized is distinct from previous reports of flies turning in the direction opposite to the stimulus motion. First, some opposite-direction turning behaviors can be explained by stimulus aliasing (Buchner, 1976). Aliasing cannot explain our results because the stimulus that maximally activates anti-directional behavior has a spatial frequency of 1/60 cycles per degree, well below the Nyquist frequency of the fly eye (~1/10 cycles per degree) (Buchner, 1976; Götz, 1970) and below reports of higher acuity vision in flies (Juusola et al., 2017). Aliasing would also not explain the dependence on stimulus contrast.

Second, our observations also cannot be explained by stimuli to the rear of the fly driving it in the opposite direction (Tammero *et al.*, 2004), since we observe anti-directional turning even when stimuli are only presented in only the 90 degrees in front of the fly (**Fig. 3**).

Third, it is also distinct from previous reports of reverse body saccades (Williamson *et al.*, 2018) since it manifests in persistent turns in the opposite direction of the stimulus and can be measured even when the magnitude of the turns is discarded (**Fig. 3**).

Fourth, the behavior observed here also appears to be distinct from previously-observed stimulus-density dependent behavioral reversals (Katsov and Clandinin, 2008). Those previously reported behaviors showed immediate reversals, but it took ~1 second for flies in our paradigm to switch between optomotor and anti-directional behaviors.

Anti-directional turning is unlikely to be due to adaptation to contrast alone

In mammalian retina, the direction preference of cells can switch because of upstream circuit adaptation (Rivlin-Etzion et al., 2012; Vlasits et al., 2014). However, we do not believe the anti-directional turning we observe has similar causes. In the mammalian retina, direction switching occurs when non-direction-selective neurons adapt to high contrast stimuli, which distorts the downstream direction-selective computation. Since the adaptation in those experiments occurs in

non-direction-selective neurons, it cannot be affected by the direction of the adapter stimulus.
However, we see differences in turning behavior depending on whether we adapt with front-to-
back or back-to-front stimuli (**Fig. 2e**). This observation rules out a mechanism based solely on
contrast, since the contrast content of front-to-back and back-to-front stimuli are identical.

The fly's visual system, however, adapts its gain to stimulus contrast (Drews et al., 2020;
Matulis et al., 2020). Importantly, the phenomenology of the anti-directional turning also argues
that the contrast adaptation is incomplete or heterogeneous among neurons, since contrast 1 and
contrast 0.25 stimuli result in such different behaviors. Contrast adaptation reported in the fly is
also faster than the 1-2 seconds preceding the shift to anti-directional turning in these
experiments.

Anti-directional turning behavior may require specific experimental and rearing conditions
Despite these previous reports of anti-directional turning under certain conditions, other labs
have measured sustained optomotor turning in response to high contrast stimuli (Bosch *et al.*,
2015; Götz and Wenking, 1973; Seelig *et al.*, 2010; Strother *et al.*, 2017). We suspect that the
two major causes of this difference are display brightness and rearing temperature. Some
experiments employ displays with mean luminances less than 5 cd/m^2 (Rister *et al.*, 2007; Seelig
et al., 2010; Strother *et al.*, 2017). Our screens, with a mean luminance of 100 cd/m^2, are
substantially brighter, but not especially bright when compared to natural scenes. In daytime
natural scenes, foliage and the ground have average luminances of 200-500 cd/m^2 and the sky
has an average luminance of around 4000 cd/m^2 (Frazor and Geisler, 2006). We therefore suspect
that as researchers move to using displays that can more accurately depict natural scene
luminances, anti-directional turning behaviors will be encountered more frequently.

Rearing conditions also had a significant influence on anti-directional turning behavior. Flies
reared at 25°C showed less anti-directional behavior than those reared at 20°C. We also found
differences based on fly age and fly strain. Our rearing conditions and choice of fly strain have
all been optimized during previous experiments to yield strong optomotor responses. Since anti-
directional behavior at high contrast usually occurs under conditions that yield strong optomotor
turning at low contrast, these optimized conditions may be required to observe anti-directional
behavior. Temperature, in particular, has developmental effects on neural connectivity (Kiral et
al., 2021). Notably, all three of these parameters vary significantly across the field, with prior
studies varying rearing temperatures from 18 to 20 to 25ºC (see for instance (Creamer *et al.*,
2018; Juusola *et al.*, 2017; Ketkar et al., 2020; Mongeau and Frye, 2017; Strother *et al.*, 2017)),
ages from 1 day to 10 days (see for instance (Bahl *et al.*, 2013; Silies *et al.*, 2013; Tammero *et*
al., 2004)), and strain between CantonS or OregonR (see for instance (Clark *et al.*, 2011; Rister
et al., 2007)). Thus, we believe that these factors likely account for the fact that this phenomenon
has not previously been reported.

Tuning of anti-directional turning matches tuning of direction selective neurons
The study of anti-directional turning behavior may yield clues about the temporal tuning of fly
motion detectors. Optomotor behavior is tuned to visual stimuli in the range of 8-22 Hz (Creamer
et al., 2018; Strother *et al.*, 2018; Tammero *et al.*, 2004; Tuthill et al., 2013), while anti-
directional behavior is tuned to stimuli in the 0.5-4 Hz range (**Fig. 1**). Intriguingly, this slower

tuning matches the tuning of T4, T5, and HS neurons, as measured via calcium imaging or electrophysiology (Chiappe et al., 2010; Creamer *et al.*, 2018; Joesch *et al.*, 2008; Maisak *et al.*, 2013). Previous studies have suggested that the difference in tuning between behavior and imaging are due to octopamine released during behavior, and not necessarily released during imaging (Arenz *et al.*, 2017; Chiappe *et al.*, 2010; Strother *et al.*, 2018). In this work, we demonstrate a motion-related behavior tuned to low frequencies, comparable to those in neural measurements, during behavior that requires T4 and T5 neurons. Overall, this suggests that T4 and T5 are required for behaviors with very different temporal tuning, which in turn suggests that the temporal tuning of behavior is not determined solely by T4 and T5 tuning, but by other, parallel pathways as well.

Anti-directional turning is unlikely to occur in nature

In a natural environment, flies are unlikely to encounter a situation where they see continuous motion in the same direction for more than 1 second. Measurements of free walking behavior have shown that the time constant of the autocorrelation of fly turning is around 100 ms (DeAngelis et al., 2019; Katsov et al., 2017). This means that the anti-directional turning studied here has likely not been directly subject to evolutionary pressures. However, the fact that we observed strong anti-directional turning in *D. yakuba* indicates that anti-directional turning is not an idiosyncratic behavior of *D. melanogaster*. It seems likely that the behavior reflects some other requirement of fly behavior, whose circuits are engaged by this stimulus. In this context, the anti-directional turning response represents a promising avenue to further constrain the underlying mechanisms of motion detection and integration. Indeed, just as illusory motion stimuli have placed key constraints on the circuits and algorithms for visual motion detection in flies and vertebrates (Agrochao et al., 2020; Clark *et al.*, 2011; Clark *et al.*, 2014; Eichner et al., 2011; Leonhardt *et al.*, 2016; Salazar-Gatzimas *et al.*, 2018; Salazar-Gatzimas *et al.*, 2016; Theobald et al., 2008; Tuthill et al., 2011) (Adelson and Bergen, 1985; Anstis and Rogers, 1975; Conway et al., 2005; Hassenstein and Reichardt, 1956; Hu and Victor, 2010; Livingstone et al., 2001; Livingstone and Conway, 2003; Mo and Koch, 2003; Orger et al., 2000), we believe that the anti-directional turning we describe here will provide additional insights.

In summary, we have presented evidence of a transition from syn-directional turning to no turning or to anti-directional turning when high contrast stimuli are presented to the fly. This persists across laboratory environments and across *Drosophila* species and shows plasticity with age. This behavior suggests than turning in response to rotational stimuli is not a simple reflex. Instead, the turning likely represents a superposition of behaviors driven by distinct circuits and elicited by different characteristics of the stimulus and different states of the fly. This complexity makes the optomotor response a model for studying the interactions of circuits as they control the low-dimensional behaviors that change an animal's orientation.

Contributions

OM, MC, RT, MSC, NCBM, JS, and BAB collected data. OM, MC, TRC, and DAC wrote the paper.

Acknowledgements

This work was supported by NIH R01EY026555 (DAC), R01EY022638 (TRC), and by a Chan-Zuckerberg Investigator Award (TRC). MC was supported by an NDSEG Fellowship; RT was supported by the Takenaka Foundation; MSC was supported by an NSF GRFP; NCBM was supported by a CAPES fellowship; JS was supported by a Ford Foundation Fellowship.

Methods

Fly strains

Strains used in these experiments are listed in the tables below:

Table 1: Parental stock genotypes

Name	Genotype	Source	Stock #
Wildtype	+; +; + (IsoD1)	(Gohl et al., 2011)	N/A
T4T5-Gal4	+; +; R42F06-Gal4 (IsoD1 background)	BDSC	BDSC 41253
HS-Gal4	+; +; R27B03-Gal4 (IsoD1 bg)	(Seelig et al., 2010)	BDSC 49211
CH-Gal4	w; +; R35A10-Gal4 (Janelia bg)	BDSC	BDSC 49897
UAS-Shibire^{ts}	+; +; UAS-Shibire^{ts} (IsoD1 bg)	(Silies et al., 2013)	N/A
Empty Gal4	w; +; pBDPGAL4.1Uw (Janelia bg)	BDSC	BDSC 68384
GCaMP6f	w; UAS-GCaMP6f; +	BDSC	BDSC 42747
jGCaMP7b	w; +; UAS-jGCaMP7b	BDSC	BDSC 79029
mtdTomato	w; +; UAS-mtdTomato	BDSC	BDSC 30124

Table 2: Genotypes of flies used in behavior experiments

Experimental	Gal4 Control	UAS Control	Background Control
T4T5-Gal4 x UAS-Shibire^{ts}: +; +; R42F06-Gal4/UAS-Shibire^{ts}	T4T5-Gal4 x IsoD1: +;+;R42F06-Gal4/+	IsoD1 x UAS-Shibire^{ts}: +; +; +/UAS-Shibire^{ts}	IsoD1: +; +; +
HS-Gal4 x UAS-Shibire^{ts}: +; +; R27B03-Gal4/UAS-Shibire^{ts}	HS-Gal4 x IsoD1: +; +; R27B03-Gal4/+	IsoD1 x UAS-Shibire^{ts}: +; +; +/UAS-Shibire^{ts}	IsoD1: +; +; +
CH-Gal4 x UAS-Shibire^{ts}: w/+; +; R35A10-Gal4/UAS-Shibire^{ts}	CH-Gal4 x IsoD1: w/+; +; R35A10-Gal4/+	Empty Gal4 x UAS-Shibire^{ts}: +/w; +; pBDPGAL4.1Uw /UAS-Shibire^{ts}	Empty Gal4 X IsoD1: +/w; +; +/ pBDPGAL4.1Uw

Genotypes of files used in imaging experiments: +; +; HS-Gal4/UAS-jGCaMP7b, +; UAS-GC6f/+; T4T5-Gal4/UAS-mtdTomato, w/+; +; CH-Gal4/UAS-jGCaMP7b.

Fly rearing (DAC lab)

Unless otherwise noted, flies were reared at 20 degrees Celsius in Panasonic MIR-154-PA incubators (Panasonic/PHC, Tokyo, Japan). The flies were circadian entrained on 12-hour light-dark cycles. Flies were raised on Archon Scientific glucose food (recipe D20102, Archon Scientific, Durham, NC). We used CO_2 to anesthetize flies more than 12 hours before the behavioral experiments.

Flies were tested for behavior in rigs built in the labs of DAC and TRC. Behavior shown in **Figs. 1d**, **1e**, **6c**, **6d**, **S1**, and **S4** was acquired in the lab of TRC, while the rest was obtained in the lab of DAC.

Fly rearing (TRC lab)

Flies were reared at 25°C, on molasses-based food, and circadian entrained on 12-hour light-dark cycles. Flies were collected within three hours of eclosion using brief CO2 anesthetization. *D. melanogaster* and *D. yakuba* were raised under identical conditions. Dark-reared flies were put in a dark chamber within 3 hours of eclosion. Flies tested at 0.5 days post eclosion were collected during the first two hours of the light cycle and were exposed to light until they were tested.

Stimulus generation and behavioral turning assays (DAC lab)

Stimuli were presented using DLP Lightcrafter (Texas Instruments, Dallas, TX) projectors (Creamer *et al.*, 2019). Mirrors were used to bounce the projected light onto three screens made of back-projection material, surrounding the fly. The screens covered the front 270 degrees around the fly, and ~45 degrees in elevation above and below the fly. The projectors were set to monochrome mode (green unless otherwise noted), updating at 180 Hz. Stimulus video was generated through a custom MATLAB (Mathworks, Natick, MA) application using PsychToolbox (Kleiner et al., 2007). Stimuli were mapped onto a virtual cylinder around the fly and the MATLAB application generated a viewpoint-corrected video signal.

Behavioral experiments were performed 12-60 hours after staging. For behavioral experiments, we selected female flies, and co-housed them with males after staging. Flies were cold-anesthetized and fixed to needles using UV-cured epoxy (Norland optical adhesive #63, Norland Products, Cranbury, NJ). Flies were then placed above air-suspended polypropylene balls. These balls were 6 mm in diameter and weighed ~120 mg. The balls were painted with two layers of marker coatings- a base silver layer and a red top layer. The motion of balls was detected by either a Parallax mouse sensor board (Parallax, Rocklin, CA) with an MCS-12086 sensor (Unity Opto Technology, Taipei, Taiwan), or a custom board with an ADNS 2080 sensor (Avago Technologies / Broadcom Inc, San Jose, TX). The data from these sensors were transferred to a custom MATLAB application via an Arduino Uno board.

Stimulus generation and behavioral turning assays (TRC lab)

Stimuli were presented using a DLP Lightcrafter (Texas Instruments, Dallas, TX) projector. Three coherent optic fibers were used to direct the projected light onto three screens made of back-projection material, surrounding the fly (Clark *et al.*, 2011; Clark *et al.*, 2014). The screens covered the front 270 degrees around the fly, and ~45 degrees in elevation above and below the fly. The projectors were set to monochrome mode, updating at 120 Hz. Stimulus video was generated through Flystim (https://github.com/ClandininLab/flystim), a custom Python application developed in the Clandinin Lab (Turner et al., 2022). Stimuli were mapped onto a virtual cylinder around the fly and Flystim generated a viewpoint-corrected video signal.

Behavioral experiments were performed 12-48 hours after eclosion, as described in the figures. Flies were cold-anesthetized and fixed to needles using UV-cured adhesive (Bondic, Niagara Falls, NY). Flies were then placed above air-suspended balls made with LAST-A-FOAM FR-4615 polyurethane foam (General Plastics, Tacoma, WA). These balls were 9 mm in diameter

and weighed ~91.7 mg. The motion of balls was detected by a Flea3 FL3-U3-13Y3M camera
(Teledyne Flir, Wilsonville, OR) and Fictrac software (Moore et al., 2014).

Imaging procedures
Two photon imaging (**Fig. 5**) was performed as previously described (Tanaka and Clark, 2022).
Briefly, two-photon images were acquired with a Scientifica microscope at between 6 and 13 Hz
using a 930 nm femtosecond laser (SpectraPhysics, Santa Clara, USA) using ScanImage
(Pologruto et al., 2003). Visual stimuli were presented on three screens occupying 270º of
azimuthal angle about the fly using projectors (Creamer *et al.*, 2019). Optical filters on the
projector and emission filters prevented the visual stimulus light from leaking into the two-
photon images.

Regions of interest (ROIs) were extracted from image timeseries using a watershed algorithm.
Responsive ROIs were included in the analyses. For T4 and T5 neurons, each ROI was identified
as a T4-dominant or T5-dominant ROI by its response to light vs. dark edges, following prior
procedures (Agrochao *et al.*, 2020). For all neuron types, responses were averaged over ROIs
and over trials of each stimulus type to obtain a measurement for each fly; these fly
measurements acted as the independent measurements to compute means and standard error bars
for the figure.

Statistical tests
Throughout the paper, each fly was considered an independent sample for statistical purposes.
Means and standard errors were computed over flies. For imaging experiments, regions of
interest from a specific neuron type were first averaged within each fly, creating a value for each
fly's response. These values were used to calculate means and standard errors over the tested
flies. In the silencing experiments, a 2-sample Student t-test was used to test for significant
differences between the experimental genotype and parental controls.

 Supplementary Figures

Supplementary Figure S1. Individual *D. melanogaster* flies in TRC lab experiments show anti-directional turning.

a) Mean time traces of individual fly responses to the high contrast stimulus, averaged over trials. The flies are those in **Fig. 1d**.
b) Long-timescale responses of individual flies, averaged over the last 1.5 s of the 5-second stimulus in panel (a) (indicated by thick black line). Mean and SEM shown are over the trials presented to that fly.

Supplementary Figure S2. Flies perform anti-directional turning under a wide range of stimulus and growing conditions.

a) Fly turning behavior at different mean screen brightness. We swept brightness from 100 cd/m² to 0.1 cd/m² and measured turning responses to high and low contrast stimuli. Flies performed the most anti-directional behavior in response to high brightness stimuli. At 1 cd/m², flies never turned in the opposite direction of the stimulus, and at 0.1 cd/m², flies turned continuously in the same direction as the stimulus, even in high contrast conditions. We also measured average turning during the last four seconds of stimulation (black bar above time traces). Average fly behavior shown as bars on the right, with individual fly behavior shown as dots. Shaded patches in the time trace plots represent ±1 SEM, as do vertical lines on bar plots. N = 19, 10, 9, 8 flies, top to bottom.

b) Our wildtype flies were Oregon-R strain (Gohl *et al.*, 2011) raised at 20 degrees. They were grown on glucose-based food media with 12-hour light-dark cycles. Experiments were run at high temperature, 12-60 hours after eclosion. We used uniform, red balls to avoid visual feedback from walking. The response of these wildtype flies to a contrast-switching stimulus (as in **Fig. 2c**) is shown in the upper left corner. We also tested different variations of all these parameters. Canton-S flies turned less overall, and showed less anti-directional turning, but still turned in the opposite direction after 5 seconds of

high contrast stimuli. We tested flies walking on highly-visible silver balls with black
dots and saw behavior similar to wildtype. Two-week-old flies showed reduced turning
and much reduced anti-directional behavior. Flies raised at 25 degrees Celsius had
behavior similar to two-week-old flies. When we performed experiments at 25 degrees,
we saw much less optomotor turning, but anti-directional turning persisted. Rearing on
molasses-based media or in the dark did not have strong effects on behavior. N = 22, 8,
12, 12, 24, 19, 19, 13 flies top to bottom, left to right.

c) Other changes to the experimental setup did not cause large differences in behavior. We
compared responses to high contrast stimuli presented with green light (peak wavelength:
525nm) and blue light (peak wavelength: 450), and did not see large differences in
behavior. Head-fixed flies (middle) showed similar behavior to head-free flies (**a**, *top*). N
= 5 and 11 flies, top to bottom.

Supplementary Figure S3. Anti-directional turning behavior occurs when using the optical
filters also employed in the two-photon imaging experiments. High and low contrast
sinusoidal stimuli were presented as in Figure 2c, but using the bandpass filters also used in our
two-photon microscope stimulus presentation. N = 30 flies.

Supplementary Figure S4. *D. yakuba* **lacks plasticity of anti-directional responses in**
adulthood that is observed *D. melanogaster***.**

    a) Adult *yakuba* flies at various ages post eclosion were presented with 5-second, high-
        contrast, rotating sinusoidal gratings as in Fig. 6. Data was acquired in the TRC lab. Anti-
        directional responses stayed consistent from 0.5 days post eclosion (dpe) to 1, 2, and 4
        dpe, although the initial optomotor response became smaller as the flies aged. Shaded
        patches represent ±1 SEM. N = 7-11 flies.

    b) The last 1.5 seconds of the mean turning velocity of each fly was averaged, and the
        population response was plotted.

Citations

Adelson, E., and Bergen, J. (1985). Spatiotemporal energy models for the perception of motion. JOSA A *2*, 284-299.

Agrochao, M., Tanaka, R., Salazar-Gatzimas, E., and Clark, D.A. (2020). Mechanism for analogous illusory motion perception in flies and humans. Proc. Natl. Acad. Sci. *117*, 23044-23053.

Anstis, S.M., and Rogers, B.J. (1975). Illusory reversal of visual depth and movement during changes of contrast. Vision Res. *15*, 957-IN956.

Arenz, A., Drews, M.S., Richter, F.G., Ammer, G., and Borst, A. (2017). The temporal tuning of the Drosophila motion detectors is determined by the dynamics of their input elements. Curr. Biol. *27*, 929-944.

Bahl, A., Ammer, G., Schilling, T., and Borst, A. (2013). Object tracking in motion-blind flies. Nat. Neurosci. *16*, 730-738.

Bahl, A., Serbe, E., Meier, M., Ammer, G., and Borst, A. (2015). Neural mechanisms for Drosophila contrast vision. Neuron *88*, 1240-1252.

Barnhart, E.L., Wang, I.E., Wei, H., Desplan, C., and Clandinin, T.R. (2018). Sequential nonlinear filtering of local motion cues by global motion circuits. Neuron *100*, 229-243. e223.

Bausenwein, B., Dittrich, A., and Fischbach, K.-F. (1992). The optic lobe of Drosophila melanogaster. Cell Tissue Res. *267*, 17-28.

Behnia, R., Clark, D.A., Carter, A.G., Clandinin, T.R., and Desplan, C. (2014). Processing properties of ON and OFF pathways for Drosophila motion detection. Nature *512*, 427-430.

Borst, A., and Weber, F. (2011). Neural Action Fields for Optic Flow Based Navigation: A Simulation Study of the Fly Lobula Plate Network. PLOS ONE *6*, e16303. 10.1371/journal.pone.0016303.

Bosch, D.S., van Swinderen, B., and Millard, S.S. (2015). Dscam2 affects visual perception in Drosophila melanogaster. Frontiers in behavioral neuroscience *9*, 149.

Buchner, E. (1976). Elementary movement detectors in an insect visual system. Biol. Cybern. *24*, 85-101.

Chiappe, M.E., Seelig, J.D., Reiser, M.B., and Jayaraman, V. (2010). Walking modulates speed sensitivity in Drosophila motion vision. Curr. Biol. *20*, 1470-1475.

Clark, D. (1981). Visual responses in developing zebrafish. Brachydanio rerio.

Clark, D.A., Bursztyn, L., Horowitz, M.A., Schnitzer, M.J., and Clandinin, T.R. (2011). Defining the computational structure of the motion detector in Drosophila. Neuron *70*, 1165-1177.

Clark, D.A., Fitzgerald, J.E., Ales, J.M., Gohl, D.M., Silies, M., Norcia, A.M., and Clandinin,
T.R. (2014). Flies and humans share a motion estimation strategy that exploits natural scene
statistics. Nat. Neurosci. *17*, 296-303.

Conway, B.R., Kitaoka, A., Yazdanbakhsh, A., Pack, C.C., and Livingstone, M.S. (2005). Neural
basis for a powerful static motion illusion. J. Neurosci. *25*, 5651-5656.

Creamer, M.S., Mano, O., and Clark, D.A. (2018). Visual Control of Walking Speed in
Drosophila. Neuron *100*, 1460-1473.

Creamer, M.S., Mano, O., Tanaka, R., and Clark, D.A. (2019). A flexible geometry for
panoramic visual and optogenetic stimulation during behavior and physiology. J. Neurosci.
Methods *323*, 48-55.

DeAngelis, B.D., Zavatone-Veth, J.A., and Clark, D.A. (2019). The manifold structure of limb
coordination in walking Drosophila. eLife *8*, e46409.

Drews, M.S., Leonhardt, A., Pirogova, N., Richter, F.G., Schuetzenberger, A., Braun, L., Serbe,
E., and Borst, A. (2020). Dynamic Signal Compression for Robust Motion Vision in Flies. Curr.
Biol.

Duistermars, B., Chow, D., Condro, M., and Frye, M. (2007). The spatial, temporal and contrast
properties of expansion and rotation flight optomotor responses in Drosophila. J. Exp. Biol. *210*,
3218.

Eckert, H., and Dvorak, D.R. (1983). The centrifugal horizontal cells in the lobula plate of the
blowfly, Phaenicia sericata. J. Insect Physiol. *29*, 547-560.

Eichner, H., Joesch, M., Schnell, B., Reiff, D.F., and Borst, A. (2011). Internal structure of the
fly elementary motion detector. Neuron *70*, 1155-1164.

Frazor, R.A., and Geisler, W.S. (2006). Local luminance and contrast in natural images. Vision
Res. *46*, 1585-1598.

Fujiwara, T., Brotas, M., and Chiappe, M.E. (2022). Walking strides direct rapid and flexible
recruitment of visual circuits for course control in Drosophila. Neuron.

Goetz, K.G. (1968). Flight control in Drosophila by visual perception of motion. Biol. Cybern. *4*,
199-208.

Gohl, D.M., Silies, M.A., Gao, X.J., Bhalerao, S., Luongo, F.J., Lin, C.C., Potter, C.J., and
Clandinin, T.R. (2011). A versatile in vivo system for directed dissection of gene expression
patterns. Nat. Methods *8*, 231-237.

Götz, K. (1964). Optomotorische untersuchung des visuellen systems einiger augenmutanten der
fruchtfliege Drosophila. Biol. Cybern. *2*, 77-92.

Götz, K., and Wenking, H. (1973). Visual control of locomotion in the walking fruitfly
Drosophila. J. Comp. Physiol. A *85*, 235-266.

Götz, K.G. (1970). Fractionation of *Drosophila* Populations According to Optomotor Traits.
Journal of Experimental Biology *52*, 419-436.

Götz, K.G. (1975). The optomotor equilibrium of theDrosophila navigation system. Journal of
comparative physiology *99*, 187-210. 10.1007/BF00613835.

Haikala, V., Joesch, M., Borst, A., and Mauss, A.S. (2013). Optogenetic control of fly optomotor
responses. J. Neurosci. *33*, 13927-13934.

Hassenstein, B., and Reichardt, W. (1956). Systemtheoretische Analyse der Zeit-, Reihenfolgen-
und Vorzeichenauswertung bei der Bewegungsperzeption des Rüsselkäfers Chlorophanus. Zeits.
Naturforsch. *11*, 513–524.

Heisenberg, M., and Buchner, E. (1977). The role of retinula cell types in visual behavior
ofDrosophila melanogaster. Journal of Comparative Physiology A: Neuroethology, Sensory,
Neural, and Behavioral Physiology *117*, 127-162.

Henning, M., Ramos-Traslosheros, G., Gür, B., and Silies, M. (2022). Populations of local
direction–selective cells encode global motion patterns generated by self-motion. Science
advances *8*, eabi7112.

Hu, Q., and Victor, J.D. (2010). A set of high-order spatiotemporal stimuli that elicit motion and
reverse-phi percepts. J. Vis. *10*.

Joesch, M., Plett, J., Borst, A., and Reiff, D. (2008). Response properties of motion-sensitive
visual interneurons in the lobula plate of Drosophila melanogaster. Curr. Biol. *18*, 368-374.

Joesch, M., Schnell, B., Raghu, S., Reiff, D., and Borst, A. (2010). ON and OFF pathways in
Drosophila motion vision. Nature *468*, 300-304.

Juusola, M., Dau, A., Song, Z., Solanki, N., Rien, D., Jaciuch, D., Dongre, S.A., Blanchard, F.,
de Polavieja, G.G., and Hardie, R.C. (2017). Microsaccadic sampling of moving image
information provides Drosophila hyperacute vision. Elife *6*, e26117.

Katsov, A., and Clandinin, T. (2008). Motion processing streams in Drosophila are behaviorally
specialized. Neuron *59*, 322-335.

Katsov, A.Y., Freifeld, L., Horowitz, M.A., Kuehn, S., and Clandinin, T.R. (2017). Dynamic
structure of locomotor behavior in walking fruit flies. eLife *6*, e26410.

Ketkar, M.D., Sporar, K., Gür, B., Ramos-Traslosheros, G., Seifert, M., and Silies, M. (2020).
Luminance information is required for the accurate estimation of contrast in rapidly changing
visual contexts. Curr. Biol. *30*, 657-669. e654.

Kim, A.J., Fenk, L.M., Lyu, C., and Maimon, G. (2017). Quantitative predictions orchestrate
visual signaling in Drosophila. Cell *168*, 280-294. e212.

Kiral, F.R., Dutta, S.B., Linneweber, G.A., Hilgert, S., Poppa, C., Duch, C., von Kleist, M.,
Hassan, B.A., and Hiesinger, P.R. (2021). Brain connectivity inversely scales with
developmental temperature in Drosophila. Cell Rep. *37*, 110145.

Kitamoto, T. (2001). Conditional modification of behavior in Drosophila by targeted expression
of a temperature-sensitive shibire allele in defined neurons. J. Neurobiol. *47*, 81-92.

Kleiner, M., Brainard, D., Pelli, D., Ingling, A., Murray, R., and Broussard, C. (2007). What's
new in Psychtoolbox-3. Perception *36*, 1.

Koerner, F., and Schiller, P.H. (1972). The optokinetic response under open and closed loop
conditions in the monkey. Exp. Brain Res. *14*, 318-330.

Krapp, H.G., and Hengstenberg, R. (1996). Estimation of self-motion by optic flow processing in
single visual interneurons. Nature *384*, 463-466.

Leong, J.C.S., Esch, J.J., Poole, B., Ganguli, S., and Clandinin, T.R. (2016). Direction selectivity
in Drosophila emerges from preferred-direction enhancement and null-direction suppression. J.
Neurosci. *36*, 8078-8092.

Leonhardt, A., Ammer, G., Meier, M., Serbe, E., Bahl, A., and Borst, A. (2016). Asymmetry of
Drosophila ON and OFF motion detectors enhances real-world velocity estimation. Nat.
Neurosci. *19*, 706–715.

Livingstone, M., Pack, C., and Born, R. (2001). Two-dimensional substructure of MT receptive
fields. Neuron *30*, 781-793.

Livingstone, M.S., and Conway, B.R. (2003). Substructure of direction-selective receptive fields
in macaque V1. J. Neurophysiol. *89*, 2743-2759.

Maisak, M.S., Haag, J., Ammer, G., Serbe, E., Meier, M., Leonhardt, A., Schilling, T., Bahl, A.,
Rubin, G.M., Nern, A., et al. (2013). A directional tuning map of Drosophila elementary motion
detectors. Nature *500*, 212-216.

Matulis, C.A., Chen, J., Gonzalez-Suarez, A., Behnia, R., and Clark, D.A. (2020).
Heterogeneous temporal contrast adaptation in Drosophila direction-selective circuits. Curr. Biol.

Mauss, A.S., Pankova, K., Arenz, A., Nern, A., Rubin, G.M., and Borst, A. (2015). Neural
circuit to integrate opposing motions in the visual field. Cell *162*, 351-362.

McCann, G.D., and MacGinitie, G. (1965). Optomotor response studies of insect vision.
Proceedings of the Royal Society of London. Series B. Biological Sciences *163*, 369-401.

Mo, C.-H., and Koch, C. (2003). Modeling reverse-phi motion-selective neurons in cortex:
double synaptic-veto mechanism. Neural Comput. *15*, 735-759.

Mongeau, J.-M., and Frye, M.A. (2017). Drosophila spatiotemporally integrates visual signals to
control saccades. Curr. Biol. *27*, 2901-2914. e2902.

Moore, R.J., Taylor, G.J., Paulk, A.C., Pearson, T., van Swinderen, B., and Srinivasan, M.V.
(2014). FicTrac: a visual method for tracking spherical motion and generating fictive animal
paths. J. Neurosci. Methods *225*, 106-119.

Orger, M.B., Smear, M.C., Anstis, S.M., and Baier, H. (2000). Perception of Fourier and non-
Fourier motion by larval zebrafish. Nat. Neurosci. *3*, 1128-1133.

Pologruto, T.A., Sabatini, B.L., and Svoboda, K. (2003). ScanImage: flexible software for
operating laser scanning microscopes. Biomed. Eng. Online *2*, 13.

Rister, J., Pauls, D., Schnell, B., Ting, C., Lee, C., Sinakevitch, I., Morante, J., Strausfeld, N.,
Ito, K., and Heisenberg, M. (2007). Dissection of the peripheral motion channel in the visual
system of Drosophila melanogaster. Neuron *56*, 155-170.

Rivlin-Etzion, M., Wei, W., and Feller, M.B. (2012). Visual stimulation reverses the directional
preference of direction-selective retinal ganglion cells. Neuron *76*, 518-525.

Salazar-Gatzimas, E., Agrochao, M., Fitzgerald, J.E., and Clark, D.A. (2018). The Neuronal
Basis of an Illusory Motion Percept Is Explained by Decorrelation of Parallel Motion Pathways.
Curr. Biol. *28*, 3748-3762. e3748.

Salazar-Gatzimas, E., Chen, J., Creamer, M.S., Mano, O., Mandel, H.B., Matulis, C.A.,
Pottackal, J., and Clark, D.A. (2016). Direct measurement of correlation responses in Drosophila
elementary motion detectors reveals fast timescale tuning. Neuron *92*, 227-239.

Schilling, T., and Borst, A. (2015). Local motion detectors are required for the computation of
expansion flow-fields. Biology open, bio. 012690.

Schnell, B., Raghu, S.V., Nern, A., and Borst, A. (2012). Columnar cells necessary for motion
responses of wide-field visual interneurons in Drosophila. J. Comp. Physiol. A *198*, 389-395.

Schweigart, G., Mergner, T., Evdokimidis, I., Morand, S., and Becker, W. (1997). Gaze
stabilization by optokinetic reflex (OKR) and vestibulo-ocular reflex (VOR) during active head
rotation in man. Vision Res. *37*, 1643-1652.

Seelig, J., Chiappe, M., Lott, G., Dutta, A., Osborne, J., Reiser, M., and Jayaraman, V. (2010).
Two-photon calcium imaging from head-fixed Drosophila during optomotor walking behavior.
Nat. Methods.

Shinomiya, K., Huang, G., Lu, Z., Parag, T., Xu, C.S., Aniceto, R., Ansari, N., Cheatham, N.,
Lauchie, S., Neace, E., et al. (2019). Comparisons between the ON-and OFF-edge motion
pathways in the Drosophila brain. eLife *8*, e40025.

Silies, M., Gohl, D.M., Fisher, Y.E., Freifeld, L., Clark, D.A., and Clandinin, T.R. (2013).
Modular use of peripheral input channels tunes motion-detecting circuitry. Neuron *79*, 111-127.

Strother, J.A., Nern, A., and Reiser, M.B. (2014). Direct observation of ON and OFF pathways
in the Drosophila visual system. Curr. Biol. *24*, 976-983.

Strother, J.A., Wu, S.-T., Rogers, E.M., Eliason, J.L., Wong, A.M., Nern, A., and Reiser, M.B.
(2018). Behavioral state modulates the ON visual motion pathway of Drosophila. Proc. Natl.
Acad. Sci. USA *115*, E102-E111.

Strother, J.A., Wu, S.-T., Wong, A.M., Nern, A., Rogers, E.M., Le, J.Q., Rubin, G.M., and
Reiser, M.B. (2017). The emergence of directional selectivity in the visual motion pathway of
Drosophila. Neuron *94*, 168-182. e110.

Takemura, S.-y., Bharioke, A., Lu, Z., Nern, A., Vitaladevuni, S., Rivlin, P.K., Katz, W.T.,
Olbris, D.J., Plaza, S.M., Winston, P., et al. (2013). A visual motion detection circuit suggested
by Drosophila connectomics. Nature *500*, 175-181.

Tammero, L., Frye, M., and Dickinson, M. (2004). Spatial organization of visuomotor reflexes in
Drosophila. J. Exp. Biol. *207*, 113-122.

Tanaka, R., and Clark, D.A. (2022). Neural mechanisms to exploit positional geometry for
collision avoidance. Curr. Biol. *32*, 2357-2374.e2356.

Theobald, J.C., Duistermars, B.J., Ringach, D.L., and Frye, M.A. (2008). Flies see second-order
motion. Curr. Biol. *18*, R464-R465.

Turner, M.H., Krieger, A., Pang, M.M., and Clandinin, T.R. (2022). Visual and motor signatures
of locomotion dynamically shape a population code for visual features in Drosophila. bioRxiv.

Tuthill, J.C., Chiappe, M.E., and Reiser, M.B. (2011). Neural correlates of illusory motion
perception in Drosophila. Proc. Natl. Acad. Sci. USA *108*, 9685-9690.

Tuthill, J.C., Nern, A., Holtz, S.L., Rubin, G.M., and Reiser, M.B. (2013). Contributions of the
12 neuron classes in the fly lamina to motion vision. Neuron *79*, 128-140.

Vlasits, A.L., Bos, R., Morrie, R.D., Fortuny, C., Flannery, J.G., Feller, M.B., and Rivlin-Etzion,
M. (2014). Visual stimulation switches the polarity of excitatory input to starburst amacrine
cells. Neuron *83*, 1172-1184.

Wei, H., Kyung, H.Y., Kim, P.J., and Desplan, C. (2020). The diversity of lobula plate tangential
cells (LPTCs) in the Drosophila motion vision system. J. Comp. Physiol. A *206*, 139-148.

Wienecke, C.F., Leong, J.C., and Clandinin, T.R. (2018). Linear Summation Underlies Direction
Selectivity in Drosophila. Neuron.

Williamson, W.R., Peek, M.Y., Breads, P., Coop, B., and Card, G.M. (2018). Tools for Rapid
High-Resolution Behavioral Phenotyping of Automatically Isolated Drosophila. Cell Rep. *25*,
1636-1649. e1635.

Wolf, R., and Heisenberg, M. (1990). Visual control of straight flight in Drosophila
melanogaster. Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and
Behavioral Physiology *167*, 269-283.

Yang, H.H., and Clandinin, T.R. (2018). Elementary motion detection in Drosophila: algorithms
and mechanisms. Ann. Rev. Vis. Sci. *4*, 143-163.

York, R.A., Brezovec, L.E., Coughlan, J., Herbst, S., Krieger, A., Lee, S.-Y., Pratt, B., Smart,
A.D., Song, E., and Suvorov, A. (2022). The evolutionary trajectory of drosophilid walking.
Curr. Biol.