Enkurin: A novel marker for myeloproliferative neoplasms from platelet, megakaryocyte, and whole blood specimens.

Sumanth Mosale Seetharam[1#], Yi Liu[2#], Jason Wu[3], Lenn Fechter[4], Kanagavel Murugesan[1], Holden Maecker[4,5], Jason Gotlib[4,6#], James Zehnder[1,4,6#], Ramasamy Paulmurugan[2] and Anandi Krishnan[1,4*]

[1]Department of Pathology, Stanford University, Stanford, CA

[2]Department of Radiology, Stanford University, Stanford, CA

[3]High-Throughput Bioscience Center (HTBC), Stanford University School of Medicine, Stanford, CA

[4]Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA

[5]Department of Microbiology & Immunology, Stanford University School of Medicine, Stanford, CA

[6]Department of Medicine, Stanford University School of Medicine, Stanford, CA

[#]These authors contributed equally

*Correspondence: anandi.krishnan@stanford.edu

Key Points

1. Enkurin, a calcium adaptor protein, is identified as a novel marker of pathogenesis in MPNs.
2. MPN megakaryocyte and platelet expression of enkurin at RNA and protein levels is inversely associated with a cell differentiation cycle gene, CDC20.
3. Likely role for dysregulated calcium homeostasis, and ER and protein folding stress in MPN transformation.

Abstract

Impaired protein homeostasis, though well established in age-related disorders, has been linked in recent research with the pathogenesis of myeloproliferative neoplasms (MPNs). As yet, however, little is known about MPN-specific modulators of proteostasis, thus impeding our ability for increased mechanistic understanding and discovery of additional therapeutic targets. Loss of proteostasis, in itself, is traced to dysregulated mechanisms in protein folding and intracellular calcium signaling at the endoplasmic reticulum (ER). Here, using *ex vivo* and *in vitro* systems (including *CD34+* cultures from patient bone marrow, and healthy cord/peripheral blood specimens), we extend our prior data from MPN patient platelet RNA sequencing, and discover select proteostasis-associated markers at RNA and/or protein levels in each of platelets, parent megakaryocytes, and whole blood specimens. Importantly, we identify a novel role in MPNs for enkurin (*ENKUR*), a calcium mediator protein, implicated originally only in spermatogenesis. Our data reveal consistent *ENKUR* downregulation at both RNA and protein levels across MPN patient specimens and experimental models, with a concomitant upregulation of a cell cycle marker, *CDC20*. Silencing of *ENKUR* by shRNA in CD34+ derived megakaryocytes further confirm this association with *CDC20* at both RNA and protein levels; and indicate a likely role for the *PI3K/Akt* pathway. The inverse association of *ENKUR* and *CDC20* expression was further confirmed upon treatment with thapsigargin (an agent that causes protein misfolding in the ER by selective loss of calcium) in both megakaryocyte and platelet fractions at RNA and protein levels. Together, our work sheds light on enkurin as a novel marker of MPN pathogenesis beyond the genetic alterations; and indicates further mechanistic investigation into a role for dysregulated calcium homeostasis, and ER and protein folding stress in MPN transformation.

Introduction

Myeloproliferative neoplasms (MPNs) are a group of malignant disorders of the bone marrow where a dysregulated balance between proliferation and differentiation gives rise to abnormal numbers of blood cells in circulation. Classical MPNs[1,2] are defined by a combination of clinical, laboratory, morphological and molecular genetic features, and constitute three phenotypic subtypes: essential thrombocythemia (ET), polycythemia vera (PV), and myelofibrosis (MF, primary or secondary if transformed from prior ET/PV). Somatic mutations in one of three driver genes (*JAK2, CALR, MPL*) constitute their shared molecular genetic pathogenesis; causing constitutive JAK/STAT signaling in hematopoietic stem cells (HSC)[3-7]. The mutated *CALR* also induces oncogenic effects by binding and constitutively activating the thrombopoietin receptor and the downstream signaling cascade[7,8] causing cellular transformation and abnormal megakaryopoiesis. To date, therapeutic strategies have largely focused on *JAK2* inhibition (e.g. ruxolitinib[9]) and are effective in alleviating MPN symptoms, but only partially, given their primary role in suppression of inflammation and reduction of circulating pro-inflammatory cytokines[10,11]. Additional therapeutic strategies that target MPN pathogenetic mechanisms[12-17] would be critical in resolving patient disease burden.

A recent study[18,19] assessing mechanisms of MPN pathogenesis identifies dysregulation of proteostasis and protein quality control at the endoplasmic reticulum (ER) as crucial transformative events, and therefore, a strong therapeutic target. However, this study was primarily in mice and remains to be confirmed in independent additional MPN models and patient-derived specimens. Our recent investigation, profiling the blood platelet transcriptome in all three subtypes of MPN patients[20] identified high expression of genes and pathways associated with impaired proteostasis, ER stress and unfolded protein response (UPR) well-recognized in other age-related disorders[21-25]. Here, we extend our prior MPN patient platelet transcriptomic data with a focus on genes associated with proteostasis, cell proliferation, ER stress or calcium signaling at high statistical significance (FDR <0.01), and representing four types of trends in platelet RNA expression across MPN subtypes: progressively downregulated, progressively upregulated, consistently upregulated, and uniquely upregulated in MF alone. Genes selected include i) enkurin, *ENKUR*[26], a little-studied TRPC (transient receptor potential cation channel[27,28]) adaptor protein with known function primarily in the context of sperm motility alone[26,29], ii) *CALR, a* well-known ER chaperone and calcium binding protein[30-34], also relevant in the

context of megakaryocyte to platelet differentiation[35-38], iii) cyclic AMP-responsive element-binding

protein *CREB3L1,* an ER-Golgi stress transducer and transcription protein involved in the UPR

response[39-42] and lastly iv) *CDC20,* involved in cell division and proliferation[43-45], and known to be

upregulated in other cancers[45]. Using MPN patient bone marrow- and healthy cord- and peripheral

blood-derived CD34+ cells differentiated into megakaryocyte and platelet cultures *ex vivo*, we

consistently identify *ENKUR* as a novel peripheral and marrow biomarker in MPNs at both RNA and

protein levels. Our data also demonstrate a negative correlation of *ENKUR* with *CDC20* expression in

the megakaryocyte/platelet cultures, and in independent banked MPN whole blood RNA specimens.

Beyond the utility of enkurin as a potential marker of MPN chronic vs advanced subtypes, more

research is warranted to elucidate the mechanisms of its downregulation in MPNs, and how that may

target cell proliferation pathways.

Methods

106   Please see **Table 1** for a tabulated overview of all RNA and protein measurements in this study.

Ethics Statement

108   All MPN patient and healthy donor samples were obtained under written informed patient consent

109   and were fully anonymized. Study approval was provided by the Stanford University Institutional

110   Review Board. All relevant ethical regulations were followed.

Materials

We collected MPN patient bone marrow specimens from the Stanford Cancer Center, and whole

blood from healthy donors at the Stanford Blood Center. Cell culture supplies including plates, fetal

bovine serum, penicillin, streptomycin, phosphate-buffered saline (PBS), and culture medium were

procured from GIBCO BRL (Frederick, MD). Antibodies for CALR, CDC20, p-PI3K, p-Akt, b-Actin

and anti-rabbit goat IgG were procured from Cell Signaling Technologies (Danvers, MA), for enkurin

from Sigma Aldrich (St. Louis, MO) and CREB3L1 from Thermo Scientific (Rockford, IL). Antibodies

for Alexa647-tagged CD41, PE-CD42b, PE-CD61, and PE-CD45 were from BioLegend (San Diego,

CA). shRNA for ENKUR silencing were synthesized from Protein and Nucleic acid facility at

Stanford University (Stanford, CA). The human HEK 293 FT cells and THP1 human macrophage cells

were purchased from the American Type Culture Collection (ATCC, Manassas, VA) and cultured

123 following supplier instructions. The CD34, CD45 microbeads, LS and MS columns and magnetic

124 separators were purchased from Miltenyi Biotech (Cambridge, MA). Cytokines IL6, thrombopoietin

125 TPO, and Flt3 ligand were from PeproTech Inc. (Cranbury, NJ); SCF and SFEM II media was from

126 Stem Cell Technologies (Kent, WA). Lenti-X™ GoStix™ Plus was purchased from TaKaRa Bio Inc.

127 (San Jose, CA).

Peripheral blood platelet isolation

130 Peripheral blood was collected in acid citrate-dextrose (ACD, 3.2%) sterile yellow-top tubes (Becton,

131 Dickinson and Co.) and was processed within 4 h of collection for all samples. Platelets were isolated

132 by established[46-49] purification protocols. Briefly, the ACD-tube whole blood was first centrifuged at

133 200xg for 20min at room temperature (RT). The platelet-rich plasma (PRP) was removed and

134 Prostaglandin E1 was added to the PRP to prevent exogenous platelet activation. The PRP was then

135 centrifuged at 1000xg for 20min at RT. The platelet pellet was re-suspended in warmed (37 deg C)

136 PIPES saline glucose (PSG). Leukocytes were depleted using $CD45^+$ magnetic beads (Miltenyi Biotec).

137 Isolated platelets were further resuspended in Trizol or LDS buffer for RNA (PCR) and protein

138 (Western Blot) analyses.

RNA extraction and quantification from whole blood (PAXgene tubes)

141 Whole blood from MPN patients and healthy donors was collected into PAXgene tubes (BD

142 Biosciences) containing RBC lysis buffer . RNA was isolated using PAXgene Blood RNA kit following

143 manufacturer instructions (762164, PreAnalytix, Switzerland).

CD34+ cell isolation from MPN bone marrow and healthy peripheral and cord blood cultures

146 $CD34^+$ cells were isolated[50,51]   using microbeads positive selection (Miltenyi Biotec) from source

147 specimens of peripheral blood or cord blood of healthy donors or bone marrow of MPN patients

148 (collected as part of their clinical care with research consent). Briefly, the RBCs were lysed using

149 RBC lysis buffer and the cells pelleted. Cells were suspended in MACS running buffer and incubated

150 with the $CD34^+$ microbeads for 1 hour on ice and then passed through MACS LS columns pre-

151 equilibrated with MACS running buffer. $CD34^+$ cells thus collected were washed using MACS

152 running buffer and resuspended in the SFEM II media ($4\text{-}5\times10^5$ cells/mL) containing TPO (20ng/ml),

SCF (25ng/ml) and Gentamicin (1:1000) and plated/transferred to 12 well plates at 37 °C in 5% CO_2.

On day 3, cells were resuspended in fresh media (SFEM II) containing TPO (20ng/ml), SCF (25ng/ml)

and Gentamicin (1:1000), and on day 6, 9, and 12, in TPO (40ng/ml) and Gentamicin (1:1000). The

cells were harvested on day 15. Cells were collected and centrifuged at 300xg for 10 min to pellet the

larger megakaryocyte fraction and at 3000xg for 30 min for the smaller platelet fraction. With both

peripheral- and cord-blood-derived CD34+ cells, the isolated cells were expanded using hematopoietic

stem cell expansion media (Cell Genix) containing 100ng/ml of IL6, Flt3 ligand, SCF, and TPO for 5

days before starting the culture. The formation of MK cells and platelets were confirmed by surface

marker analysis with CD41, CD42b and CD61.

Cell surface marker analysis using flow cytometry

Following isolation of the megakaryocyte and platelet fractions from the 14-day (+ TPO) CD34+

culture, cells were collected and washed with PBS. For the CD41, CD42b and CD61 stains, cells were

resuspended in MACS running buffer and stained with Alexa 647 anti-human CD41, PE anti-human

CD42b, and PE anti-human CD61, and incubated at RT protected from light for 45 min. Cells were

washed with PBS and resuspended at a final concentration of 2×10^7 cells/ml in PBS, prior to running

the sample on the flow cytometer. All experiments were performed using GUAVA Flow Cytometer

and all flow cytometry analysis was performed in the FlowJo Software. Data from flow cytometry

experiments was acquired by gating for events that were in focus.

Total RNA isolation and PCR with CD34+ derived megakaryocyte and platelet fractions

Total RNA was isolated from the megakaryocyte (MK) and platelet fractions using a mirVana RNA

extraction kit (Life Technologies, Grand Island, NY) in accordance with the manufacturer's

instructions. Briefly, MKs/platelets were homogenized into 300μl lysis buffer, then incubated with

30μl homogenate additive for 10 min. RNA was extracted with acid-phenol and column purified,

then washed three times with washing buffer, and eluted in 40μl of sterile elution buffer. The total

RNA was quantified first using a Nanodrop spectrophotometer, then 100ng of total RNAs were

reverse-transcribed using a Reverse-Transcription Kit (Life Technologies) and RT primers to

synthesize cDNA. Then, qRT-PCR was performed using the TaqMan-PCR primers and probe mix

combined with the cDNA derivates. qRT-PCR was performed through 2 min incubation at 50 °C,

then followed by DNA polymerase activation at 95 °C for 10 min; plus 60-cycles at 95 °C for 15 s, and 60 °C for 60 s in the BioRad CFX96 thermocycler system (BioRad). The qRT-PCR reaction procedure was executed in a 20µl final reaction volume. The expression of target genes was analyzed by the 2–ΔΔCt method.

Thapsigargin-induced ER stress

Isolated human CD34+ cells from the peripheral blood of healthy volunteers were cultured for 15 days with the supplementation of both SCF and TPO for the first 6 days and then with TPO alone till day 15. Fresh media containing either SCF and/ or TPO was added every 3 days. To induce ER stress in these cells, known ER stressor drug Thapsigargin (125nM, as optimum dosage identified via preliminary experiments) was added to the culture on day 7. The culture was maintained till day 15 with media change every 3 days. The cells were harvested and analyzed for gene expression and immunoblotting for the target proteins.

Western Blotting for protein expression in megakaryocytes and platelets

Isolated MKs were aliquoted for flow cytometry, RNA expression and western blotting. For western blotting, an aliquot of the MK cells was pelleted and lysed using cell lysis buffer containing protease inhibitor cocktail, EDTA. The samples were then resolved on a reducing SDS-PAGE (10% acrylamide). Blots were stained using appropriate primary (anti-human rabbit antibody; 1:1000 v/v) and secondary antibodies (anti-rabbit IgG HRP conjugate; 1:2000 v/v).

Protein expression was determined based on the detection of a band. The intensity of the protein bands observed was semi-quantified using IVIS or ImageJ software with normalization of each protein against beta-actin. Blots were visualized using ECL reagent by Amersham Imager 680 or by IVIS imaging systems.

ENKUR gene silencing

a. Construction of the vector

PLKO1 vector was used to construct the ENKUR-silenced cell lines in HEK293T cells (known for their ease of transfection and fast growth rate). The PLKO1 vector was cut using EcoR1 and Age1 to ligate the shRNA using T4 ligase.

The shRNA sequences used were:

shRNA1: PCCGGTCCGGCCAACCTCGATACTCTTATTTCTCGAGAAATAAGAGTATCG

AGGTTGGTTTTTTTTGG

shRNA 2:

PCCGGTCCGGCATGGGAGTGGCTAAAAAGCCCTCGAGGGCTTTTTAGCC

ACTCCCATGTTTTTTTTGG

Scrambled: PCCGGTCCGGGTGCGTTGCTAGTACCAACTCTCGAGAGTTGGTACTA

GCAACGCACTTTTTTTTGG

b. Validation of the construct

The silencing effect of the shRNA construct was assessed using HEK293 FT cells by transfection of

the shRNA containing plasmids (pLKO1-shRNA) using lipofectamine method. Briefly, 100K HEK293

FT cells were seeded in a 12-well plate 24h prior to the transfection. Transfection was performed by

adding a master mix containing lipofectamine and the shRNA in reduced serum media for 4 hours,

before adding the complete media. The cells were harvested 48h after transfection and the silencing

effect assessed by western immunoblotting against enkurin.

Cell culture for ENKUR silencing

HEK293 FT cells were cultured in Dulbecco's Modified Eagle's Medium/high glucose with 10% FBS,

0.1% streptomycin, and 100 U mL^{-1} penicillin at 37 °C, 5% CO_2, and 95% air environment, while

THP-1 cells were cultured in RPMI media containing 10% FBS, 0.1% streptomycin, and 100 U mL^{-1}

penicillin at 37 °C, 5% CO_2, and 95% air environment. The cells were tested for any mycoplasma

contamination using a MycoAlert kit (Lonza, Allendale, NJ), and were maintained at optimum

cultural conditions.

Production of lentiviral particles

After the confirmation of the silencing effect of the shRNA construct, they were packaged along with

the viral plasmids to produce the lentivirus. Briefly, the lentiviral vectors were transfected to HEK293

FT cells using calcium phosphate method. The cells were supplemented with chloroquine and

HEPES. After 48h of incubation, the media was carefully aspirated, and the viral particles were

concentrated by ultracentrifugation. The IFU value was determined by Lenti-X GoStix following

manufacturer instructions (Cat#631280).

Validation of the lentivirus silencing in HEK cells

247 As above, 100K HEK293 FT cells were plated in a 12-well plate 24h prior to lentiviral infection

248 (transduction). The media was aspirated and lentivirus master mix containing polybrene in reduced

249 serum media was added to the cells. After 4h of incubation, the cells were supplemented with

250 complete media and was incubated for 48h before harvesting. The cells were lysed and

251 immunoblotted against enkurin to assess the gene silencing effect.

ENKUR gene silencing in CD34+ derived megakaryocytes

The shRNA expressing lentiviral particles were infected into the CD34+ cells in culture on day 9 (MKs

derived from CD34+ cells) at a ratio of 1:20 (cell: viral particle). The culture was maintained for

another 6 days with supplementation of TPO every 3 days. Cells were harvested on day 15; and

centrifuged at two speeds; low (300xg) and high (3000xg) to separate the MK and platelet fractions

respectively. Each fraction was divided into 3 aliquots for assessing each of cell surface markers (50 –

100K cells), RNA (~50K cells) and protein (100 – 200K cells) expression.

Statistical Analyses

Continuous variables from all experiments were assessed for normality. Data that were normally

distributed were expressed as a mean plus or minus the standard error of the mean. For analyses

involving two groups, a parametric two-tailed student t-test was used. When three or more groups

were analyzed, an ANOVA with a Tukey's post-hoc test was performed. When data were not

normally distributed a Mann-Whitney was used when two groups were analyzed while a Kruskal-

Wallis with a Dunn's multiple comparison post-hoc test was used for analyses of three or more

groups. When appropriate a two-way ANOVA with post-hoc test was used as described. Summary

statistics were used to describe the study cohort and clinical variables were expressed as the mean ±

error of the mean or as a number and percentage (%). Statistical analyses were performed by using

GraphPad Prism (version 9, San Diego, CA), and a p-value < 0.05 was considered statistically

significant.

Results

Validation of RNA and protein expression of 4 MPN candidate markers

Four candidate markers from our prior platelet RNA-seq analyses[20] were identified based on specific

trends in progressive expression across the 3 MPN subtypes vis-à-vis healthy donors (FDR<0.01,

n=120) independent of patient driver mutational status. *ENKUR and CALR*, in particular, were

identified based on opposing trends in progressive expression (**Figure 1A and Figure S1A**), *CDC20* on

its specific high expression in MF patients alone in contrast with healthy donors and ET or PV

patients (**Figure 1A,** top right), and *CREB3L1* based on its substantially increased (> 50-fold)

expression in all MPNs vs healthy (**Figure S1A,** top right). Given that the platelet RNA profile is a

composite of influences from the peripheral circulation as well as of the genetic (and MPN

mutational) profile of megakaryocytes in the bone marrow, we hypothesized *a priori* that our MPN

candidate markers are likely to be variably validated in one versus the other.

First, in whole blood RNA of MPN patients versus healthy donors, expression of *ENKUR* and *CDC20*

was confirmed by qPCR (**Figure 1B**, n=109, $p < 0.005$) as nearly 6-fold downregulated and 3-fold

upregulated respectively in the advanced MPN subtype, MF. Expression of *CREB3L1* was also

confirmed as over 15-fold upregulated across ET, PV, and MF (**Figure S1B,** right panel), whereas

expression of *CALR* (**Figure S1B,** left panel) was variable and not uniformly consistent with the

platelet transcriptome (likely owing to lower resolution for detection of transcripts in whole blood).

Next, in assessing platelet protein levels, trends in expression of ENKUR was significantly different in

PV and MF versus healthy donors, and CDC20 in ET and MF (**Figure 1C**), whereas CALR and

CREB3L1 were not significantly different between patient and healthy donor specimens (**Figure S1C**).

Megakaryocyte and platelet fractions from CD34+ cultures of MPN patient bone marrow

To further delineate RNA expression of the candidate markers as specific to platelets in circulation

alone or also in marrow-derived megakaryocytes, we established an *ex vivo* culture of CD34+ stem

cells from MPN patient bone marrow (n=8, 4 MF) and generated megakaryocyte and platelet

fractions (**Figure 2A-B**, and Methods and **Figure S2A-B** for flow cytometric and histological

confirmation of cultured megakaryocyte and platelet fractions). Given the variable differentiation

states of the nascent platelet fraction and its lower cell quantity than the megakaryocytic fraction, we

focused entirely on the megakaryocytes harvested. Downregulation of *ENKUR* was recapitulated in these patient-derived megakaryocytic RNA (mean approx. 2-fold reduction in MF alone and 5-fold all MPN) with associated upregulation of *CDC20,* CREB3L1 (mean of approx. 1-fold) and *CALR* (~4-fold) in all MPN (n=8, **Figure 2C**).

ENKUR gene silencing

To better evaluate the negative correlation between *ENKUR* and *CDC20/CALR* expression, we sought lentiviral knockdown in primary CD34+ cells (derived from healthy peripheral blood and cord blood specimens) using two distinct *ENKUR* shRNA constructs. **Figure 3A-B** describes our experimental framework. The shRNAs were first expressed in pLKO.1 plasmid that was amplified and isolated from *E.coli*, and the lentiviral plasmids were then transfected into HEK293 FT cells to generate the lentivirus. *ENKUR* shRNA silencing and downregulation of enkurin protein was confirmed at a concentration of 1:20 HEK cell:viral particle. Primary CD34+ cells from healthy peripheral (n=4) and cord blood (n=4) specimens were cultured at four conditions: i) control with no ENKUR silencing, ii) *and* iii) lentiviral knockdown with the two shRNA constructs, and finally iv) scramble control shRNA. Where relevant, shRNA infection was introduced at day 7 of culture and continued for another 7 days (totaling a 15-day culture, **Figure 3B**). Megakaryocyte and platelet fractions thus derived were confirmed by flow cytometric analysis (**Figure 4A**). Downregulated RNA expression of *ENKUR* was significantly associated with concomitant high expression of *CDC20* in both of the megakaryocyte and platelet fractions (**Figure 4B**). Cell quantity in the platelet fraction was just sufficient for RNA qPCR measurements alone, and therefore, protein levels could be assessed only in the megakaryocytes. Inverse correlation between ENKUR and CDC20 was validated even in megakaryocytic protein expression (**Figure 4C**). However, *ENKUR* silencing alone (without the associated patient factors in our *in vitro* experiments, e.g. mutational status) was insufficient to generate a statistically significant differential in *CALR or CREB3L1* expression at both RNA and protein levels (**Figure S3A-B**).

One classical downstream pathway linked previously to ER stress response[18,19] in MPNs[52] and other cancers[53-55] is the PI3K/Akt signaling cascade. Here with shENKUR, we briefly evaluated possible

association with the PI3K/Akt pathway and found that silencing of enkurin correlated with increased
expression of phosphorylated PI3K in megakaryocytes (**Figure 4C**).

Effect of ER stress on CD34+-derived megakaryocytes

Considering that the differential and inversely correlated expression of *ENKUR* and *CDC20* extends
to the CD34+ cell-derived megakaryocytes, and our hypothesis on the role of ER stress in this
response, we assessed RNA and protein levels of these markers following treatment of the cells with
the sarco/endoplasmic reticulum Ca^{2+}-ATPase inhibitor, thapsigargin (125nM). CD34+ cells from
peripheral blood of healthy donors were differentiated into megakaryocyte and platelet fractions,
with thapsigargin introduced at day 7 of a 15-day culture (see Methods and **Figure 5A**; limited cell
quantity of the platelet fraction enabled protein analyses in the megakaryocytic fraction alone).
Thapsigargin evoked a similar expression profile in megakaryocytes **(Figure 5B-C)** as that of MPN
patient bone marrow-derived CD34+ cells with downregulated *ENKUR* and upregulated *CDC20*
expression at both RNA and protein levels.

Discussion

Here, we present the first report, to our knowledge, of enkurin (*ENKUR*) as a potential new
peripheral biomarker and therapeutic strategy in myeloproliferative neoplasms. Our prior study[20]
profiling the platelet transcriptome in patients with chronic MPNs (n=120) identified progressive
association in expression of several proteostasis-associated genes with advancing disease subtype. Our
findings were consistent with other studies demonstrating dysregulated proteostasis as a primary
effector of myeloid transformation[18,56,19,57]. The MPN platelet RNA-seq data also confirmed the limited
impact[58] of treatment by JAK2-inhibitor ruxolitinib (RUX) relative to the substantial disease burden
in myelofibrosis; and urged the need for development of novel candidate drugs to be used alone or in
combination with RUX for the treatment of MPNs.

Four candidate markers from the prior study (*ENKUR, CDC20, CALR, and CREB3L1*) were evaluated
in this work based not only on their significant association with pathways related to proteostasis, cell
proliferation, ER stress or calcium signaling, but also progressive increase or decrease in differential
expression across the MPN chronic vs advanced subtypes (ET/PV to MF versus healthy donors)

regardless of patient *JAK2/CALR* mutational status. Using CD34[+] cells isolated from MPN patient

bone marrow as well as healthy donor peripheral and cord blood, we generated *ex vivo*

megakaryocyte and platelet fractions and discovered at both RNA and protein levels, an inverse

correlation between *ENKUR*, a calmodulin and TRPC channel modulator[26,29] (not previously

associated with MPNs) and *CDC20* (also termed Fizzy[44]), a cell cycle gene and an anaphase promoting

complex activator (with known associations with malignancies broadly[43,59] and select studies in

MPNs[60]). Lentiviral-mediated silencing of *ENKUR* was utilized to further confirm the negative

association between *ENKUR* and *CDC20;* and indicating a potential interaction via the PI3K-Akt

pathway **(Figure 6)**. Our data on the inverse association between enkurin, a membrane calcium influx

adaptor protein, and *CDC20*, a crucial cell cycle and proliferation regulator point to a role for

dysregulated calcium signaling in MPNs; and the possibility for the ratio of expression between the

two to be developed as a potential biomarker of MPN disease.

Next, in evaluating calreticulin (*CALR*), a well-established driver mutation in MPNs[31,61], as well as

CREB3L1, an ER/UPR stress response transcription factor, high expression noted previously across

MPN patient platelet RNA-seq[20] was confirmed in patient-derived specimens (platelet, whole-blood,

and bone-marrow-CD34[+]-cultured megakaryocytes) but only at the RNA (and not protein) levels.

Future investigations applying genome editing in MPN models will be needed to ascertain

overlapping mechanisms of somatic alterations in *CALR* and its effect on function in altered calcium

signaling and dysregulated ER protein folding[62,63] in MPNs.

Taken together, we offer two inversely associated MPN markers, *ENKUR* and *CDC20*, whose whole

blood, platelet and megakaryocyte RNA and protein expression reflect MPN pathobiology

irrespective of patient driver mutational status. These candidate markers also seek to expand our

understanding of MPN pathology beyond the classical inflammatory signatures[64-67].

Limitations of the study

389 There are several limitations to our study. First, higher statistical power and longitudinal prospective

390 data will be necessary to further confirm *ENKUR/CDC20* as markers for MPN disease progression.

391 Second, future investigations evaluating these signatures in patient-derived CD34[+] cells may identify

additional functional aspects of bone marrow and MPN pathology. Third, we recognize that our data are not assessing mechanistic signaling of how loss of *ENKUR* expression or impaired calcium signaling contributes to myeloproliferation; and will need to be specifically interrogated in MPN murine and other models. Follow-on studies evaluating a potential role of the *PI3K/Akt* pathway, in particular the effect of pharmaceutical inhibitors already in use in MF will add significant value to the current study. Studies evaluating platelet function under impaired enkurin expression and calcium modulation are promising immediate future directions. And above all, it will be important to assess how enkurin downregulation might be more broadly relevant to other hematological cancers, such as myelodysplastic syndromes and acute myeloid leukemia, or potentially unique to MPNs alone.

Acknowledgements

This work was funded by the MPN Research Foundation and the US National Institutes of Health
grants 1K08HG010061-01A1 and 3UL1TR001085-04S1 (research re-entry award) to A.K. Other
support included 1S10OD018220 and 1S10OD021763 shared instrumentation grants to the Stanford
Functional Genomics Facility and the Stanford Research Computing Center, and the Charles and Ann
Johnson Foundation to J.G. All authors thank the patients at the Stanford Cancer Center and the
healthy donors at the Stanford Blood Center for their generous participation in this research, and the
Canary Center at Stanford for Cancer Early Detection for access to flow cytometry and microscopy.
Authors also thank Dr. Diwash Jangam for his assistance with the initial Paxgene PCR setup.

Author Contributions AK conceived of the study and the team, secured funding, and wrote the
manuscript. LF and JG provided samples and clinical annotation and reviewed the clinical data. AK
and RP coordinated and oversaw sample acquisition and processing. SS, YL, JW, and KM performed
the experiments and interpreted the analyses with AK, RP, HM, JG, and JZ. All authors critically
reviewed and edited the manuscript. [#]SS and YL contributed equally. [#]JG and JZ also contributed
equally. All authors approved the final manuscript.

Conflict of Interest Disclosures: Authors declare no conflict of interest.

Bibliography
2

1.   Tremblay, D., Yacoub, A., and Hoffman, R. Overview of Myeloproliferative Neoplasms: History,
Pathogenesis, Diagnostic Criteria, and Complications. Hematol Oncol Clin North Am. 2021;35(2):159-
176. Epub 2021/03/02. PubMed PMID: 33641861; PMCID: PMC8669599.
2.   Arber, D.A., Orazi, A., Hasserjian, R., Thiele, J., Borowitz, M.J., Le Beau, M.M., Bloomfield, C.D.,
Cazzola, M., and Vardiman, J.W. The 2016 revision to the World Health Organization classification of
myeloid neoplasms and acute leukemia. Blood. 2016;127(20):2391-2405. PubMed PMID: 27069254.
3.   Rumi, E., and Cazzola, M. Diagnosis, risk stratification, and response evaluation in classical
myeloproliferative neoplasms. Blood. 2017;129(6):680-692. PubMed PMID: 28028026; PMCID:
PMC5335805.
4.   Vainchenker, W., and Kralovics, R. Genetic basis and molecular pathophysiology of classical
myeloproliferative neoplasms. Blood. 2017;129(6):667-679. PubMed PMID: 28028029.
5.   Zoi, K., and Cross, N.C. Genomics of Myeloproliferative Neoplasms. J Clin Oncol. 2017;35(9):947-954.
PubMed PMID: 28297629.
6.   Bao, E.L., Nandakumar, S.K., Liao, X., Bick, A.G., Karjalainen, J., Tabaka, M., Gan, O.I., Havulinna,
A.S., Kiiskinen, T.T.J., Lareau, C.A., de Lapuente Portilla, A.L., Li, B., Emdin, C., Codd, V., Nelson,
C.P., Walker, C.J., Churchhouse, C., de la Chapelle, A., Klein, D.E., Nilsson, B., et al. Inherited
myeloproliferative neoplasm risk affects haematopoietic stem cells. Nature. 2020;586(7831):769-775.
Epub 2020/10/16. PubMed PMID: 33057200; PMCID: PMC7606745.
7.   Marneth, A.E., and Mullally, A. The Molecular Genetics of Myeloproliferative Neoplasms. Cold Spring
Harb Perspect Med. 2020;10(2). Epub 2019/09/25. PubMed PMID: 31548225; PMCID: PMC6996444.
8.   Elf, S., Abdelfattah, N.S., Chen, E., Perales-Patón, J., Rosen, E.A., Ko, A., Peisker, F., Florescu, N.,
Giannini, S., Wolach, O., Morgan, E.A., Tothova, Z., Losman, J.-A., Schneider, R.K., Al-Shahrour, F.,
and Mullally, A. Mutant Calreticulin Requires Both Its Mutant C-terminus and the Thrombopoietin
Receptor for Oncogenic Transformation. Cancer Discovery. 2016;6(4):368-381.
9.   Vannucchi, A.M. Ruxolitinib versus standard therapy for the treatment of polycythemia vera. N Engl J
Med. 2015;372(17):1670-1671. Epub 2015/04/23. PubMed PMID: 25901432.
10.   Kleppe, M., Kwak, M., Koppikar, P., Riester, M., Keller, M., Bastian, L., Hricik, T., Bhagwat, N.,
McKenney, A.S., Papalexi, E., Abdel-Wahab, O., Rampal, R., Marubayashi, S., Chen, J.J., Romanet, V.,
Fridman, J.S., Bromberg, J., Teruya-Feldstein, J., Murakami, M., Radimerski, T., et al. JAK-STAT
pathway activation in malignant and nonmalignant cells contributes to MPN pathogenesis and
therapeutic response. Cancer Discov. 2015;5(3):316-331. Epub 2015/01/13. PubMed PMID: 25572172;
PMCID: PMC4355105.
11.   Deininger, M., Radich, J., Burn, T.C., Huber, R., Paranagama, D., and Verstovsek, S. The effect of long-
term ruxolitinib treatment on JAK2p.V617F allele burden in patients with myelofibrosis. Blood.
2015;126(13):1551-1554. Epub 2015/08/01. PubMed PMID: 26228487; PMCID: PMC4582331.
12.   Jutzi, J.S., Marneth, A.E., Ciboddo, M., Guerra-Moreno, A., Jimenez-Santos, M.J., Kosmidou, A.,
Dressman, J.W., Liang, H., Hamel, R., Lozano, P., Rumi, E., Doench, J.G., Gotlib, J., **Krishnan, A.**, Elf,
S., Al-Shahrour, F., and Mullally, A. Whole-genome CRISPR screening identifies N-glycosylation as a
genetic and therapeutic vulnerability in CALR-mutant MPNs. Blood. 2022;140(11):1291-1304. Epub
2022/06/29. PubMed PMID: 35763665.
13.   Maxson, J.E. A new role for hematoxylin: targeting CALR. Blood. 2021;137(14):1848-1849. Epub
2021/04/09. PubMed PMID: 33830192.
14.   Gerds, A.T. Beyond JAK-STAT: novel therapeutic targets in Ph-negative MPN. Hematology Am Soc
Hematol Educ Program. 2019;2019(1):407-414. Epub 2019/12/07. PubMed PMID: 31808852; PMCID:
PMC6913459 Celgene, Apexx Oncology, CTI Biopharma, and Inctye and research funding from Roche,
Genentech, Incyte, Celgene, Imago Biosciences, Samus Therapeutics, and Gilead.

15. Jayavelu, A.K., Schnoder, T.M., Perner, F., Herzog, C., Meiler, A., Krishnamoorthy, G., Huber, N., Mohr, J., Edelmann-Stephan, B., Austin, R., Brandt, S., Palandri, F., Schroder, N., Isermann, B., Edlich, F., Sinha, A.U., Ungelenk, M., Hubner, C.A., Zeiser, R., Rahmig, S., et al. Splicing factor YBX1 mediates persistence of JAK2-mutated neoplasms. Nature. 2020;588(7836):157-163. Epub 2020/11/27. PubMed PMID: 33239784.

16. Lee, S.S., Verstovsek, S., and Pemmaraju, N. Novel Therapies in Myeloproliferative Neoplasms: Beyond JAK Inhibitor Monotherapy. J Immunother Precis Oncol. 2021;4(3):117-128. Epub 2022/06/07. PubMed PMID: 35663101; PMCID: PMC9138435.

17. How, J., Garcia, J.S., and Mullally, A. Biology and therapeutic targeting of molecular mechanisms in MPN. Blood. 2022. Epub 2022/12/20. PubMed PMID: 36534936.

18. Osorio, F.G., Soria-Valles, C., Santiago-Fernandez, O., Bernal, T., Mittelbrunn, M., Colado, E., Rodriguez, F., Bonzon-Kulichenko, E., Vazquez, J., Porta-de-la-Riva, M., Ceron, J., Fueyo, A., Li, J., Green, A.R., Freije, J.M., and Lopez-Otin, C. Loss of the proteostasis factor AIRAPL causes myeloid transformation by deregulating IGF-1 signaling. Nat Med. 2016;22(1):91-96. Epub 2015/12/23. PubMed PMID: 26692333.

19. LaFave, L.M., and Levine, R.L. Targeting a regulator of protein homeostasis in myeloproliferative neoplasms. Nat Med. 2016;22(1):20-21. Epub 2016/01/07. PubMed PMID: 26735404.

20. Shen, Z., Du, W., Perkins, C., Fechter, L., Natu, V., Maecker, H., Rowley, J., Gotlib, J., Zehnder, J., and Krishnan, A. Platelet transcriptome identifies progressive markers and potential therapeutic targets in chronic myeloproliferative neoplasms. Cell Reports Medicine. 2021;2(10):100425; PMCID: PMC8561315.

21. Jain, K., Tyagi, T., Du, J., Hu, X., Patell, K., Martin, K.A., and Hwa, J. Unfolded Protein Response Differentially Modulates the Platelet Phenotype. Circulation research. 2022;131(4):290-307. Epub 2022/07/22. PubMed PMID: 35862006; PMCID: PMC9357223.

22. Kaushik, S., and Cuervo, A.M. Proteostasis and aging. Nature Medicine. 2015;21(12):1406-1415.

23. Costa-Mattioli, M., and Walter, P. The integrated stress response: From mechanism to disease. Science. 2020;368(6489).

24. Walter, P., and Ron, D. The unfolded protein response: from stress pathway to homeostatic regulation. Science. 2011;334(6059):1081-1086.

25. Lebeaupin, C., Yong, J., and Kaufman, R.J. The impact of the ER unfolded protein response on cancer initiation and progression: therapeutic implications. Adv Exp Med Biol. 2020;1243:113-131; PMCID: PMC7243802.

26. Sutton, K.A., Jungnickel, M.K., Wang, Y., Cullen, K., Lambert, S., and Florman, H.M. Enkurin is a novel calmodulin and TRPC channel binding protein in sperm. Dev Biol. 2004;274(2):426-435. Epub 2004/09/24. PubMed PMID: 15385169.

27. Wright, J.R., Amisten, S., Goodall, A.H., and Mahaut-Smith, M.P. Transcriptomic analysis of the ion channelome of human platelets and megakaryocytic cell lines. Thromb Haemost. 2016;116(2):272-284. Epub 2016/06/10. PubMed PMID: 27277069; PMCID: PMC5080539.

28. Authi, K.S. TRP channels in platelet function. Handb Exp Pharmacol. 2007(179):425-443. Epub 2007/01/16. PubMed PMID: 17217071.

29. Jungnickel, M.K., Sutton, K.A., Baker, M.A., Cohen, M.G., Sanderson, M.J., and Florman, H.M. The flagellar protein Enkurin is required for mouse sperm motility and for transport through the female reproductive tract. Biol Reprod. 2018;99(4):789-797. Epub 2018/05/08. PubMed PMID: 29733335; PMCID: PMC6203878.

30. Fucikova, J., Spisek, R., Kroemer, G., and Galluzzi, L. Calreticulin and cancer. Cell Res. 2021;31(1):5-16. Epub 2020/08/01. PubMed PMID: 32733014; PMCID: PMC7853084.

31. Klampfl, T., Gisslinger, H., Harutyunyan, A.S., Nivarthi, H., Rumi, E., Milosevic, J.D., Them, N.C., Berg, T., Gisslinger, B., Pietra, D., Chen, D., Vladimer, G.I., Bagienski, K., Milanesi, C., Casetti, I.C.,

Sant'Antonio, E., Ferretti, V., Elena, C., Schischlik, F., Cleary, C., et al. Somatic mutations of calreticulin in myeloproliferative neoplasms. N Engl J Med. 2013;369(25):2379-2390. Epub 2013/12/12. PubMed PMID: 24325356.

32. Merlinsky, T.R., Levine, R.L., and Pronier, E. Unfolding the Role of Calreticulin in Myeloproliferative Neoplasm Pathogenesis. Clin Cancer Res. 2019;25(10):2956-2962. Epub 2019/01/19. PubMed PMID: 30655313; PMCID: PMC6522317.

33. Prins, D., González Arias, C., Klampfl, T., Grinfeld, J., and Green, A.R. Mutant Calreticulin in the Myeloproliferative Neoplasms. HemaSphere. 2020;4(1):e333.

34. Michalak, M., Groenendyk, J., Szabo, E., Gold, L.I., and Opas, M. Calreticulin, a multi-process calcium-buffering chaperone of the endoplasmic reticulum. Biochem J. 2009;417(3):651-666. Epub 2009/01/13. PubMed PMID: 19133842.

35. Clapham, D.E. Calcium Signaling. Cell. 2007;131(6):1047-1058.

36. Tonelli, F.M.P., Santos, A.K., Gomes, D.A., da Silva, S.L., Gomes, K.N., Ladeira, L.O., and Resende, R.R. (2012). Stem Cells and Calcium Signaling. In Calcium Signaling, M.S. Islam, ed. (Springer Netherlands), pp. 891-916.

37. Andrea Di Buduo, C., Moccia, F., Battiston, M., De Marco, L., Mazzucato, M., Moratti, R., Tanzi, F., and Balduini, A. The importance of calcium in the regulation of megakaryocyte function. Haematologica. 2014;99(4):769-778.

38. Salati, S., Prudente, Z., Genovese, E., Pennucci, V., Rontauroli, S., Bartalucci, N., Mannarelli, C., Ruberti, S., Zini, R., Rossi, C., Bianchi, E., Guglielmelli, P., Tagliafico, E., Vannucchi, A.M., and Manfredini, R. Calreticulin Affects Hematopoietic Stem/Progenitor Cell Fate by Impacting Erythroid and Megakaryocytic Differentiation. Stem Cells Dev. 2018;27(4):225-236. Epub 2017/12/21. PubMed PMID: 29258411.

39. Chan, C.-P., Kok, K.-H., and Jin, D.-Y. CREB3 subfamily transcription factors are not created equal: Recent insights from global analyses and animal models. Cell Biosci. 2011;1(1):6; PMCID: PMC3116243.

40. Cui, M., Kanemoto, S., Cui, X., Kaneko, M., Asada, R., Matsuhisa, K., Tanimoto, K., Yoshimoto, Y., Shukunami, C., and Imaizumi, K. OASIS modulates hypoxia pathway activity to regulate bone angiogenesis. Sci Rep. 2015;5:16455; PMCID: PMC4642342.

41. García, I.A., Torres Demichelis, V., Viale, D.L., Di Giusto, P., Ezhova, Y., Polishchuk, R.S., Sampieri, L., Martinez, H., Sztul, E., and Alvarez, C. CREB3L1-mediated functional and structural adaptation of the secretory pathway in hormone-stimulated thyroid cells. J Cell Sci. 2017;130(24):4155-4167; PMCID: PMC6518157.

42. Sampieri, L., Di Giusto, P., and Alvarez, C. CREB3 Transcription Factors: ER-Golgi Stress Transducers as Hubs for Cellular Homeostasis. Front Cell Dev Biol. 2019;7:123; PMCID: PMC6616197.

43. Schrock, M.S., Stromberg, B.R., Scarberry, L., and Summers, M.K. APC/C ubiquitin ligase: Functions and mechanisms in tumorigenesis. Semin Cancer Biol. 2020;67(Pt 2):80-91. Epub 2020/03/14. PubMed PMID: 32165320; PMCID: PMC7483777.

44. Visintin, R., Prinz, S., and Amon, A. CDC20 and CDH1: a family of substrate-specific activators of APC-dependent proteolysis. Science. 1997;278(5337):460-463. Epub 1997/10/23. PubMed PMID: 9334304.

45. Yu, H. Cdc20: A WD40 Activator for a Cell Cycle Degradation Machine. Molecular Cell. 2007;27(1):3-16.

46. Campbell, R.A., Franks, Z., Bhatnagar, A., Rowley, J.W., Manne, B.K., Supiano, M.A., Schwertz, H., Weyrich, A.S., and Rondina, M.T. Granzyme A in human platelets regulates the synthesis of proinflammatory cytokines by monocytes in aging. J Immunol. 2018;200(1):295-304; PMCID: PMC5736423.

47. Rowley, J.W., Oler, A.J., Tolley, N.D., Hunter, B.N., Low, E.N., Nix, D.A., Yost, C.C., Zimmerman, G.A., and Weyrich, A.S. Genome-wide RNA-seq analysis of human and mouse platelet transcriptomes. Blood. 2011;118(14):e101-111. Epub 2011/05/21. PubMed PMID: 21596849; PMCID: PMC3193274.

48. Middleton, E.A., Rowley, J.W., Campbell, R.A., Grissom, C.K., Brown, S.M., Beesley, S.J., Schwertz, H., Kosaka, Y., Manne, B.K., Krauel, K., Tolley, N.D., Eustes, A.S., Guo, L., Paine, R., 3rd, Harris, E.S., Zimmerman, G.A., Weyrich, A.S., and Rondina, M.T. Sepsis alters the transcriptional and translational landscape of human and murine platelets. Blood. 2019;134(12):911-923. Epub 2019/08/02. PubMed PMID: 31366617; PMCID: PMC6753624.

49. Amisten, S. A rapid and efficient platelet purification protocol for platelet gene expression studies. Methods in molecular biology (Clifton, N.J.). 2012;788:155-172. Epub 2011/12/02. PubMed PMID: 22130707.

50. Edelstein, L.C., Simon, L.M., Montoya, R.T., Holinstat, M., Chen, E.S., Bergeron, A., Kong, X., Nagalla, S., Mohandas, N., Cohen, D.E., Dong, J.-f., Shaw, C., and Bray, P.F. Racial differences in human platelet PAR4 reactivity reflect expression of PCTP and miR-376c. Nature Medicine. 2013;19(12):1609-1616.

51. Bhatlekar, S., Basak, I., Edelstein, L.C., Campbell, R.A., Lindsey, C.R., Italiano, J.E., Weyrich, A.S., Rowley, J.W., Rondina, M.T., Sola-Visner, M., and Bray, P.F. Anti-apoptotic <i>BCL2L2</i> increases megakaryocyte proplatelet formation in cultures of human cord blood. Haematologica. 2019;104(10):2075-2083.

52. Khan, I., Huang, Z., Wen, Q., Stankiewicz, M.J., Gilles, L., Goldenson, B., Schultz, R., Diebold, L., Gurbuxani, S., Finke, C.M., Lasho, T.L., Koppikar, P., Pardanani, A., Stein, B., Altman, J.K., Levine, R.L., Tefferi, A., and Crispino, J.D. AKT is a therapeutic target in myeloproliferative neoplasms. Leukemia. 2013;27(9):1882-1890. Epub 2013/06/12. PubMed PMID: 23748344; PMCID: PMC4023863.

53. Fruman, D.A., Chiu, H., Hopkins, B.D., Bagrodia, S., Cantley, L.C., and Abraham, R.T. The PI3K Pathway in Human Disease. Cell. 2017;170(4):605-635.

54. Song, M., Bode, A.M., Dong, Z., and Lee, M.-H. AKT as a Therapeutic Target for Cancer. Cancer Research. 2019;79(6):1019-1031.

55. Yang, J., Nie, J., Ma, X., Wei, Y., Peng, Y., and Wei, X. Targeting PI3K in cancer: mechanisms and advances in clinical trials. Mol Cancer. 2019;18(1):26.

56. Liu, L., Inoki, A., Fan, K., Mao, F., Shi, G., Jin, X., Zhao, M., Ney, G., Jones, M.A., Sun, S., Dou, Y., Inoki, K., Qi, L., and Li, Q. ER associated degradation preserves hematopoietic stem cell quiescence and self-renewal by restricting mTOR activity. Blood. 2020. Epub 2020/11/06. PubMed PMID: 33150381.

57. Skov, V., Larsen, T.S., Thomassen, M., Riley, C., Jensen, M.K., Bjerrum, O.W., Kruse, T.A., and Hasselbalch, H.C. Increased Expression of Proteasome-Related Genes In Patients with Primary Myelofibrosis. Blood. 2010;116(21):4117-4117.

58. Deininger, M., Radich, J., Burn, T.C., Huber, R., Paranagama, D., and Verstovsek, S. The effect of long-term ruxolitinib treatment on JAK2p.V617F allele burden in patients with myelofibrosis. Blood. 2015;126(13):1551-1554.

59. VanGenderen, C., Harkness, T.A.A., and Arnason, T.G. The role of Anaphase Promoting Complex activation, inhibition and substrates in cancer development and progression. Aging (Albany NY). 2020;12(15):15818-15855. Epub 2020/08/18. PubMed PMID: 32805721; PMCID: PMC7467358.

60. Spivak, J.L., Jie, C., Williams, D., and Moliterno, A.R. JAK2V671F, Gender and the Gene Expression Profile of Polycythemia Vera. Blood. 2008;112(11):2805-2805.

61. Nangalia, J., Massie, C.E., Baxter, E.J., Nice, F.L., Gundem, G., Wedge, D.C., Avezov, E., Li, J., Kollmann, K., Kent, D.G., Aziz, A., Godfrey, A.L., Hinton, J., Martincorena, I., Van Loo, P., Jones, A.V., Guglielmelli, P., Tarpey, P., Harding, H.P., Fitzpatrick, J.D., et al. Somatic CALR mutations in myeloproliferative neoplasms with nonmutated JAK2. N Engl J Med. 2013;369(25):2391-2405. Epub 2013/12/12. PubMed PMID: 24325359; PMCID: PMC3966280.

62. Di Buduo, C.A., Abbonante, V., Marty, C., Moccia, F., Rumi, E., Pietra, D., Soprano, P.M., Lim, D., Cattaneo, D., Iurlo, A., Gianelli, U., Barosi, G., Rosti, V., Plo, I., Cazzola, M., and Balduini, A. Defective interaction of mutant calreticulin and SOCE in megakaryocytes from patients with myeloproliferative neoplasms. Blood. 2020;135(2):133-144. Epub 2019/11/08. PubMed PMID: 31697806; PMCID: PMC6952826.

63. Mullally, A. Both sides now: losses and gains of mutant CALR. Blood. 2020;135(2):82-83. Epub 2020/01/10. PubMed PMID: 31917838; PMCID: PMC6952827.

64. Geyer, H.L., Dueck, A.C., Scherber, R.M., and Mesa, R.A. Impact of inflammation on myeloproliferative neoplasm symptom development. Mediators Inflamm. 2015;2015:284706; PMCID: PMC4619953.

65. Koschmieder, S., and Chatain, N. Role of inflammation in the biology of myeloproliferative neoplasms. Blood Rev. 2020;42:100711. Epub 2020/06/09. PubMed PMID: 32505517.

66. Vainchenker, W., Leroy, E., Gilles, L., Marty, C., Plo, I., and Constantinescu, S.N. JAK inhibitors for the treatment of myeloproliferative neoplasms and other disorders. F1000Res. 2018;7:82. Epub 2018/02/06. PubMed PMID: 29399328; PMCID: PMC5773931.

67. Woods, B., Chen, W., Chiu, S., Marinaccio, C., Fu, C., Gu, L., Bulic, M., Yang, Q., Zouak, A., Jia, S., Suraneni, P.K., Xu, K., Levine, R.L., Crispino, J.D., and Wen, Q.J. Activation of JAK/STAT Signaling in Megakaryocytes Sustains Myeloproliferation In Vivo. Clin Cancer Res. 2019;25(19):5901-5912. Epub 2019/06/21. PubMed PMID: 31217200; PMCID: PMC6774846.

68. Bruno, S., Ghelli Luserna di Rora, A., Napolitano, R., Soverini, S., Martinelli, G., and Simonetti, G. CDC20 in and out of mitosis: a prognostic factor and therapeutic target in hematological malignancies. J Exp Clin Cancer Res. 2022;41(1):159. Epub 2022/05/01. PubMed PMID: 35490245; PMCID: PMC9055704.

Tables

3 **Table 1:** Summary of all cell types used, their sources and the respective analyses performed.

colspan across	Table 1: Experimental Framework and Components						
#	Cell type	Source	Megakaryocyte (MK) culture	ENKUR silencing	ER stress induction	RNA expression (MK and/or Platelet)	Protein expression (MK and/or Platelet)
1	Platelets	Peripheral blood (n = 49)				X *CALR, CREB3L1, CDC20, ENKUR*	X CALR, CREB3L1, CDC20, ENKUR
2	CD34+	Bone marrow (n = 8)	X			X *CALR, CREB3L1, CDC20, ENKUR*	
3	CD34+	Cord blood (n = 4)	X	X		X *CALR, CREB3L1, CDC20, ENKUR*	X CALR, CREB3L1, CDC20, ENKUR, PI3K and Akt
4	CD34+	Peripheral blood (n = 7)	X	X	X	X *CALR, CREB3L1, CDC20, ENKUR*	X CALR, CREB3L1, CDC20, ENKUR, PI3K and Akt

Figure Legends

Figure 1: RNA expression in platelets and whole blood of MPN patients and protein expression in

platelets

(A) MPN Platelet RNA-seq normalized expression data: MPN patients (total n=99: ET n=24; PV n=33;

MF n=42) vs healthy donors (n=21) (B) whole blood RNA: MPN patients (total n=109: ET n=39; PV

n=38; MF n=32) vs healthy donors (n=10) (C) platelet protein: peripheral blood platelets were isolated

from MPN patients (total n=38: MF n=12; PV n=19; ET n=7) and healthy donors (n=11). Bonferroni

multiple comparisons adjusted p values ****$p<0.0001$, ***$p<0.001$ and **$p<0.01$ when compared with

controls (healthy donors). All RNA expression is normalized to GAPDH and expressed as log2 fold

change.

Figure 2: Culturing of CD34+ cells from the bone marrow of MPN patients and RNA expression

(A) Microscopic images of the CD34+ cells isolated from the bone marrow of MPN patients over the

15-day culture period at 20X magnification. Notice the formation of megakaryocytes indicated by the

red arrow. (B) Flow cytometric analysis of cell surface markers to confirm the production of

megakaryocyte and platelet fractions after 15 days of culture. (C) megakaryocytic RNA: cell culture

from fresh MPN patient bone marrow (n=8) derived CD34+ cells then differentiated over 15 days into

megakaryocytes. All RNA expression is normalized to GAPDH and expressed as log2 fold change.

Bonferroni adjusted p values **$p<0.01$, *$p<0.05$.

Figure 3: Experimental design and workflow of Lentiviral transduction-induced silencing of ENKUR

gene in CD34+ stem cells

(A) Workflow of the shRNA construction and generation of the lentivirus (B) experimental plan for

the silencing of the ENKUR gene in CD34+ stem cells. The CD34+ cells were isolated from both

peripheral and cord blood.

Figure 4: Lentiviral transduction induced silencing of ENKUR gene in CD34+ stem cells

(A) Flow cytometric analysis of cell surface markers to confirm the megakaryocytic and platelet fractions after 15 days of culture. (B) RNA expression levels in the CD34+ cell-derived megakaryocytes and platelets. (C) (i) Western blot of the CD34+ cells derived megakaryocytes (n=4 cord blood and n=4 peripheral blood) and (ii) densitometry for the blots. Densitometry was performed using the IVIS imaging software (****$p<0.0001$, ***$p<0.001$, **$p<0.01$ and *$p<0.05$ when compared with non-silenced controls).

Figure 5: Effect of ER stress on megakaryocytes derived from CD34+ cells from healthy donors

(A) Representative microscopic images of the CD34+ cells isolated from the peripheral blood of healthy donors over the 15-day culture period at 20X magnification. The ER stressor, thapsigargin (125nM) was added to the culture at day 7 and cultured for another 7 days. (B) RNA expression: cell culture from thapsigargin-treated CD34+ derived megakaryocytes (n=7). (C) (i) Western blot of the thapsigargin-treated CD34+ cells derived megakaryocytes (n=3 in triplicate) and (ii) densitometry for the blots. Densitometry was performed using the ImageJ software (****$p<0.0001$,**$p<0.01$ when compared with untreated controls).

Figure 6: Probable mechanism for the role of *ENKUR* downregulation in cell proliferation

We hypothesize that, under normal conditions, enkurin at sufficient physiological levels is bound to PI3K via the SH3 binding domain, thus regulating its activation[26]. However, in the setting of MPNs, downregulation of enkurin likely contributes to free PI3K and its prolonged activation, further activating Akt and potentially other signaling mechanisms prompting overexpression of cell differentiation cycle genes including CDC20[68] and cell proliferation.

Supplementary Figure legends

Figure S1: RNA expression in platelets and whole blood of MPN patients and protein expression in

platelets

(A) MPN Platelet RNA-seq Normalized Expression Data: MPN patients (n=99: ET n=24; PV n=33; MF

n=42) vs healthy donors (n=21) (B) whole blood RNA: MPN patients (n=109: ET n=39; PV n=38; MF

n=32) vs healthy donors (n=10) (C) platelet protein: peripheral blood platelets were isolated from

MPN patients (n=38: MF n=12; PV n=19; ET n=7) and healthy donors (n=11).

Figure S2: Culturing of CD34+ cells from the bone marrow of MPN patients and RNA expression

(A) Confocal microscopy: Microscopic images of the CD34+ cells isolated from the bone marrow of

MPN patients over the 15-day culture period at 20X magnification. The cells were treated with cell

surface markers for CD34+ cells (CD34), Megakaryocytes (CD41, CD42b, and CD61) (B)

Quantification of CD41+ megakaryocytes from CD34+ cells isolated from the bone marrow of MPN

patients over the 15-day culture period (C) Megakaryocytic RNA: cell culture from fresh MPN

patient bone marrow (n=8) derived CD34+ cells then differentiated over 15 days into

megakaryocytes. All RNA expression is normalized to GAPDH and expressed as log2 fold change.

Figure S3: Effect of Lentiviral transduction induced silencing of ENKUR gene on CALR and CREB3L1

gene expression in CD34+ stem cells

(A) RNA expression levels in the CD34+ cell-derived megakaryocytes and platelets. Log2 fold change

was plotted by normalizing with the control (B) (i) Western blot of the CD34+ cell-derived

megakaryocytes (n=4 cord blood and n=4 peripheral blood) and (ii) densitometry for the blots.

Densitometry was performed using IVIS imaging software.

FIGURE 2

FIGURE 5

FIGURE 6

Total: 350 / CD41+ : 97 (27.7%)