Single Molecule Mechanics and Kinetics of Cardiac Myosin Interacting with Regulated Thin Filaments

Sarah R. Clippinger Schulte[1,+], Brent Scott[1,+], Samantha K. Barrick[1], W. Tom Stump[1], Thomas Blackwell[1], Michael J. Greenberg[1,*]

[1] Department of Biochemistry and Molecular Biophysics, Washington University School of Medicine, St. Louis, MO, 63110, USA

[*]Corresponding author:
Michael J. Greenberg
Department of Biochemistry and Molecular Biophysics
Washington University School of Medicine
660 S. Euclid Ave., Campus Box 8231
St. Louis, MO 63110
Phone: (314) 362-8670
Email: greenberg@wustl.edu

[+]Denotes equal contributions from S.R.C.S. and B.S.

Running title: Mechanics of myosin with RTFs

Keywords: optical trapping, single molecule, myosin, thin filament

Abstract

The cardiac cycle is a tightly regulated process wherein the heart generates force to pump blood to the body during systole and then relaxes during diastole. Disruption of this finely tuned cycle can lead to a range of diseases including cardiomyopathies and heart failure. Cardiac contraction is driven by the molecular motor myosin, which pulls regulated thin filaments in a calcium-dependent manner. In some muscle and non-muscle myosins, regulatory proteins on actin tune the kinetics, mechanics, and load dependence of the myosin working stroke; however, it is not well understood whether or how thin filament regulatory proteins tune the mechanics of the cardiac myosin motor. To address this critical gap in knowledge, we used single-molecule techniques to measure the kinetics and mechanics of the substeps of the cardiac myosin working stroke in the presence and absence of thin filament regulatory proteins. We found that regulatory proteins gate the calcium-dependent interactions between myosin and the thin filament. At physiologically relevant ATP concentrations, cardiac myosin's mechanics and unloaded kinetics are not affected by thin filament regulatory proteins. We also measured the load-dependent kinetics of cardiac myosin at physiologically relevant ATP concentrations using an isometric optical clamp, and we found that thin filament regulatory proteins do not affect either the identity or magnitude of myosin's primary load-dependent transition. Interestingly, at low ATP concentrations, thin filament regulatory proteins have a small effect on actomyosin dissociation kinetics, suggesting a mechanism beyond simple steric blocking. These results have important implications for both disease modeling and computational models of muscle contraction.

Significance Statement

Human heart contraction is powered by the molecular motor β-cardiac myosin, which

pulls on thin filaments consisting of actin and the regulatory proteins troponin and

tropomyosin. In some muscle and non-muscle systems, these regulatory proteins tune

the kinetics, mechanics, and load dependence of the myosin working stroke. Despite

having a central role in health and disease, it is not well understood whether the

mechanics or kinetics of β-cardiac myosin are affected by regulatory proteins. We show

that regulatory proteins do not affect the mechanics or load-dependent kinetics of the

working stroke at physiologically relevant ATP concentrations; however, they can affect

the kinetics at low ATP concentrations, suggesting a mechanism beyond simple steric

blocking. This has important implications for modeling of cardiac physiology and

diseases.

Introduction

The human heart is finely tuned to generate the appropriate power necessary to pump blood in response to a wide range of physiological and pathological stimuli. This power output, driven by the interactions between β-cardiac myosin and the thin filament, has exquisite regulatory mechanisms to ensure that the heart generates sufficient power during systole to perfuse the body and then relaxes during diastolic filling (1,2). At the molecular scale, these regulatory mechanisms include the calcium-dependent gating of the interactions between myosin and the thin filament by the proteins troponin and tropomyosin (3), load-dependent effects on myosin kinetics (4,5), and thick filament dependent regulation (6). Dysfunction of these regulatory mechanisms can lead to cardiomyopathy, arrhythmias, and heart failure (7-10), and these regulatory mechanisms have emerged as therapeutic targets (11-13).

β-cardiac myosin (*MYH7*) is the motor that drives cardiac contraction in human ventricles, and it has different mechanochemical properties from α-cardiac myosin (*MYH6*), the motor that drives the contraction of human atrial and mouse ventricular tissues (2,14,15). Recent optical trapping experiments elucidated the mechanical and kinetic parameters that define the interactions between single β-cardiac myosin motors and actin (4,5,16,17). These studies demonstrated that β-cardiac myosin has a two substep working stroke, where the myosin generates ~6 nm total displacement. Moreover, these studies showed that mechanical forces which oppose the β-cardiac myosin working stroke slow the kinetics of ADP release from actomyosin, leading to slowed crossbridge cycling kinetics. This load-dependent slowing of cycling kinetics contributes to the force-velocity relationship in muscle, which is a critical determinant of

power output (4). Importantly, this relationship can be impaired by mutations that cause

disease, and there are compounds in clinical trials for cardiovascular diseases that

target this relationship (16).

While previous studies have defined the mechanical and kinetic properties of β-

cardiac myosin interacting with actin, myosin in the heart interacts with regulated thin

filaments, which are macromolecular complexes consisting of actin and the regulatory

proteins troponin and tropomyosin. It is currently not known whether these thin filament

regulatory proteins affect the mechanics and/or kinetics of the β-cardiac myosin working

stroke. Optical trapping experiments with other myosin isoforms demonstrated that

regulatory proteins on actin can sometimes affect the mechanics and kinetics of the

motor. For example, it was shown that tropomyosin Tm4.2 can increase the force

sensitivity of non-muscle myosin-IIA (18). Moreover, the yeast myosin-V isoform Myo2p

walks processively only in the presence of tropomyosin Tpm1p, since tropomyosin

decorated actin filaments slow the rate of ADP release from the motor (19). Some

studies with skeletal muscle myosin showed that thin filament regulatory proteins

reduce the size of the myosin working stroke by half, potentially through disruption of

the interactions between myosin's two heads (20), while other studies showed no effect

(21,22).

Defining the fundamental parameters underlying heart contraction is critical for

understanding and modeling cardiac physiology and disease. It is essential to determine

how regulated thin filaments, not just actin, tune the mechanics, kinetics, and/or load-

dependent properties of β-cardiac myosin. Therefore, we used high-resolution optical

trapping methods to study these properties in the presence of fully regulated thin

filaments.

Results

Reconstitution of regulated thin filaments

Cardiac actin and β-cardiac myosin were tissue purified from porcine ventricles.

Porcine cardiac actin is identical to human actin, and porcine β-cardiac myosin is 97%

identical to human β-cardiac myosin, with biochemical and biophysical properties that

are indistinguishable from the human isoform (4,5,14,17,23-25). Human troponin and

tropomyosin were expressed recombinantly and reconstituted with purified actin to form

regulated thin filaments. In vitro motility assays demonstrate that these thin filaments

are functional, with no movement at low calcium (pCa 9) and robust movement at high

calcium (pCa 4) (**Fig. 1a**).

For the optical trapping experiments, it is necessary to attach regulated thin

filaments to polystyrene beads via a linkage. In these experiments, we mixed 10%

biotinylated G-actin with unlabeled G-actin during polymerization and then used the

biotinylated linkage to attach the reconstituted thin filament to streptavidin-coated beads

(4). To ensure that the biotinylated actin does not interfere with thin filament regulation

under the same fully activating or inhibiting conditions used in the trapping assays, we

performed in vitro thin filament gliding assays at both low (pCa 9) and high (pCa 4)

calcium (26). Calcium-based regulation of these thin filaments was observed, with

movement at pCa 4 and no movement at pCa 9. There was no significant difference in

the average speed of thin filament translocation at pCa 4 between regulated thin

filaments made from unlabeled actin (0.26 $\pm$ 0.08 μm/s) and those including 10%

biotinylated actin (0.25 $\pm$ 0.08 μm/s) (P = 0.12) (**Fig. 1a**).

Next, we tested whether we could observe calcium-dependent thin filament

regulation in our optical trapping assay. We used the three-bead assay pioneered by

the Spudich lab (27), in which two optically trapped beads are attached to a single actin

filament and then lowered onto a surface-bound bead that is sparsely coated with

myosin. Using reconstituted regulated thin filaments attached to streptavidin-coated

polystyrene beads, we examined binding interactions between myosin and the

regulated thin filaments. At 1 μM ATP, we could clearly resolve binding interactions

between regulated thin filaments and myosin at high calcium; however, these binding

interactions were exceedingly rare at low calcium (**Fig. 1b**). Taken together, our results

demonstrate that we are able to reconstitute functional regulated thin filaments for

optical trapping assays.

Regulatory proteins do not affect the mechanics of the myosin working stroke at low

ATP

To examine the effects of regulatory proteins on the mechanics of the β-cardiac

myosin working stroke, we used the three-bead optical trapping assay. Experiments

were conducted at 1 μM ATP and high calcium (pCa 4) to facilitate observation of

substeps of the myosin working stroke (28). Experiments were conducted in both the

presence and absence of regulatory proteins. Myosin binding to actin causes a

reduction in the variance of the bead position, and individual interactions between

myosin and thin filaments could clearly be resolved (**Figs. 2a-b**). Binding interactions

were resolved using a covariance threshold (29). Cumulative distributions measuring

the total size of the myosin working stroke show that data are well described by a single

Gaussian function (**Fig. 2e;** P = 0.09 for unregulated and P = 0.25 for regulated

filaments)**.** The total size of the β-cardiac myosin working stroke measured with

unregulated actin (5.3 ± 8.6 nm; n = 364 binding events) was consistent with previous

measurements (4,17,24,29) and not significantly different from the size of the working

stroke measured with regulated thin filaments (4.9 ± 9.1 nm; n = 491 binding events; P

= 0.57).

Previous studies have shown that β-cardiac myosin accomplishes its working

stroke in two substeps, with the first substep associated with phosphate release and the

second associated with ADP release (4,17,25,29). To better understand the effects of

regulatory proteins on the mechanics of substeps of the β-cardiac myosin working

stroke, we used ensemble averaging of individual binding interactions, which enables

the detection of subtle substeps that are typically obscured by Brownian motion

(4,30,31). Binding interactions were synchronized either upon the initiation of binding

(time forward averages) or the termination of binding (time reverse averages) using a

changepoint algorithm and then averaged as previously described (29). The difference

between the time forward and time reverse averages is indicative of a two-substep

working stroke, and we see a two-substep working stroke for both the regulated and

unregulated thin filaments (**Figs. 2c-d**), consistent with previous studies of unregulated

thin filaments (4,25,29). The displacement of the time forward and time reverse

averages at detachment gives the size of the total working stroke (29). The difference in

displacement between the end of the time forward averages and the beginning of the

time reverse averages gives the size of the second substep of the working stroke, and

the difference between the total working stroke and the second substep gives the size

of the first substep. Here, we constructed cumulative distributions of substep

displacements from our individual binding interactions, and the data follow the expected

shape of a cumulative distribution for a single Gaussian function (**Figs. 2f**-**g**). We did

not observe a statistically significant difference in the size of either the first (P = 0.36) or

second (P = 0.55) substep of the working stroke in the presence or absence of

regulatory proteins.

Regulatory proteins affect kinetics of detachment at non-physiological, low ATP

concentrations

Optical trapping experiments collected at 1 μM ATP and high calcium provide

information not only on the mechanics of the working stroke but also the kinetics. The

distribution of attachment durations can be used to calculate the actomyosin

dissociation rate. Cumulative distributions of binding interaction durations were well

fitted by single exponential functions to yield the actomyosin detachment rate (**Fig. 3a**).

Interestingly, we observe that the detachment rate measured in the presence of

regulatory proteins (3.9 (-0.3/+0.4) s^{-1}) was slightly slower than the rate measured

without regulatory proteins (5.5 ± 0.4 s^{-1}) (P = 0.002). This is consistent with previous

studies of skeletal muscle myosin conducted at low ATP concentrations which show

slower detachment rates in the presence of regulatory proteins (20).

The assertion that dissociation of regulated thin filaments from myosin is slowed

at low ATP is further supported by kinetic information extracted from the ensemble

averages. The time reverse averages give the rate of transitioning from the second

substep to the detached state, and previous studies have shown that this transition is

related to the biochemical rate of ATP-induced actomyosin dissociation (30,31). The

rates of the time reverse ensemble averages measured here at 1 μM ATP are well fitted

by single exponential functions (**Fig. 2c-d**). The rate of this transition measured in the

presence of regulatory proteins (4 s^{-1}) is slower than the rate measured in the absence

of regulatory proteins (7 s^{-1}). These rates are consistent with the detachment rates

measured from the cumulative distributions of attachment durations, suggesting that this

transition limits the actomyosin detachment rate at low ATP (**Fig. 3a**). Consistent with

the notion that this transition limits detachment, the rates of the time forward averages

collected at low ATP, which report the rate of transitioning from the first substep to the

second (**Figs. 2c-d)**, are much faster than the rates of the time reversed averages (**Fig.**

2c-d) or the detachment rates (**Fig. 3a**). Taken together, our trapping data strongly

suggest that regulatory proteins slow the rate of ATP-induced dissociation at low ATP

concentrations.

To further investigate the basis for the slowed detachment rate in the presence of

regulatory proteins observed in the optical trap at low ATP concentrations, we

measured the rates of ATP-induced actomyosin dissociation and ADP release using

stopped-flow techniques (**Figs. 3b and 3c**). It has previously been shown that at low

concentrations of ATP, the rate of actomyosin dissociation is limited by the rate of ATP-

induced actomyosin dissociation, while at physiologically relevant ATP concentrations,

detachment is limited by the rate of ADP release (14,32,33). We found that the rates of

ADP release in the presence of regulatory proteins (76 ± 5 s^{-1}) and in their absence (78

± 1 s^{-1}) were not significantly different from each other (P = 0.45), and the measured

values are consistent with previous studies (4,9,10,14,34). We also measured the rate

of ATP-induced actomyosin dissociation. Consistent with previous studies, traces were

well fitted by two exponential functions, where the rate of the fast phase (**Fig. 3c**)

reports the rate of ATP-induced actomyosin dissociation, and the rate of the slow phase

(**Supplemental Fig. 1**) measures the rate of a slow ATP-independent isomerization

(see **Methods** for details) (35). The observed fast rate of ATP-induced dissociation was

modeled as the formation of a rapid-equilibrium collision complex (K_1') followed by an

irreversible isomerization and rapid detachment (k_{+2}') (see **Methods** for details). At

saturating ATP, for both the regulated and unregulated thin filaments, the rates of ADP

release were much slower than the rate of ATP-induced dissociation (**Fig. 3d**),

consistent with ADP release limiting detachment at saturating ATP concentrations. At

low ATP concentrations, the rate of ATP binding is given by the second-order ATP

binding rate ($K_1'*k_{+2}'$) (**Figs. 3c inset, and 3d**). Consistent with the optical trapping

measurements at low ATP (**Figs. 2c, 2d and 3a**), the second order rate of ATP binding

is slower for regulated filaments ($2.2 \pm 0.4\ \mu M^{-1}\ s^{-1}$) compared to unregulated filaments

($4.0 \pm 0.8\ \mu M^{-1}\ s^{-1}$, P = 0.04). Taken together, these data demonstrate that regulatory

proteins slow the rate of ATP-induced actomyosin dissociation at low ATP

concentrations; however, at physiologically relevant ATP concentrations, the rate of

ADP release limits dissociation, and this rate is not affected by regulatory proteins.

Regulatory proteins do not affect the loaded kinetics of cardiac myosin at saturating

ATP

To examine the effects of load on myosin's mechanics, we used an isometric

optical clamp, where one optically trapped bead is assigned to be the transducer bead

and the other is assigned as the motor bead (4,36). The position of the transducer bead

is measured, and the motor bead is actively moved by a feedback loop to keep the

transducer bead in the center of the optical trap. When myosin binds to the actin and

pulls, both beads are pulled from the centers of their traps. The motor bead is then

pulled to return the transducer bead back to its original position, exerting a load on the

myosin (**Fig. 4a**). The attachment duration and the load on the myosin are measured.

Using this technique, a range of forces can be measured. Importantly, these

experiments were conducted at physiologically relevant saturating ATP concentrations

(1 mM) to enable the probing of mechanical states that are populated in the healthy

myocardium (4,16).

Using the isometric optical clamp, we obtain a scatter plot where each point

represents the attachment duration and average force of a single binding interaction.

The distribution of attachment durations at a given force are exponentially distributed

(4,37) (**Fig. 4b**). As can be seen, the attachment duration increases gradually as force

is increased, consistent with force-induced slowing of actomyosin dissociation. These

data were fitted with the Bell Equation (38) using maximum likelihood estimation as

previously described (4,37,39) to obtain the force-dependent detachment rate:

$$k(F) = k_o * e^{\frac{-F*d}{k_B*T}}$$ Equation 1

where F is the force, k_o is the rate of the primary force-sensitive transition in the

absence of force, d is the distance to the transition state (i.e., the force sensitivity), and

k_B*T is the thermal energy, which has a value of 4.11 pN*nm at room temperature.

In the absence of regulatory proteins, we found that the rate of the primary force-

sensitive transition, k_o, is 28 (-5/+7) s^{-1}. This is consistent with previous measurements

in the optical trap (4,17,24,25,29) and the rate of the time forward ensemble average

(35 s^{-1} **Fig. 2c**). The distance to the transition state, d, is 0.79 (-0.17/+0.19) nm,

consistent with previous measurements (4,17,24,25,29). Consistent with previous work,

these data demonstrate that for unregulated thin filaments, the detachment rate at

saturating ATP is limited by the force-dependent rate of ADP release (4,17,24,25,29). In

the presence of regulatory proteins (**Fig. 4b**), the rate of the primary force-sensitive

transition, k_0, is 35 (-8/+10) s^{-1}, and the distance to the transition state, d, is 0.64 (± 0.4)

nm. These values are not significantly different when comparing regulated and

unregulated thin filaments (P = 0.16 and P = 0.49 for k_0 and d, respectively). Taken

together, we do not detect any differences in the force-dependent kinetics of cardiac

myosin in the presence or absence of regulatory proteins at physiologically relevant

saturating ATP concentrations.

Discussion

We used high-resolution optical trapping techniques to examine the mechanics and kinetics of the β-cardiac myosin working stroke in both the presence and absence of regulatory proteins. We found that regulatory proteins do not tune the mechanics of the working stroke or the load-dependent kinetics of β-cardiac myosin contraction at physiologically relevant ATP concentrations; however, we did observe slight differences in the kinetics of actomyosin dissociation in the presence of regulatory proteins at low ATP, suggesting that these regulatory proteins have subtle effects beyond just sterically blocking the interactions between myosin and the thin filament.

The molecular role of regulatory proteins in cardiac muscle

In the healthy heart, the ATP concentration is ~8 mM (40,41), and even in heart failure, the ATP concentration remains in the millimolar range (1-4 mM) (42-44). Here, we found that at physiologically relevant millimolar ATP concentrations, regulatory proteins have no appreciable effect on the mechanics or kinetics of the β-cardiac myosin working stroke or the kinetics of actomyosin dissociation (**Fig. 4b**). Consistent with previous studies (7,9,10,45), our stopped-flow measurements show that at physiologically relevant ATP concentrations, the rate of ADP release is much slower than the rate of ATP-induced actomyosin dissociation (**Figs. 3b-c**), and therefore the muscle shortening speed will be limited by the rate of ADP release. We did not observe any differences in the rates of ADP release (**Fig. 3b**) or the rate of actomyosin dissociation measured in the optical trap at saturating ATP (k_0, **Fig. 4b**) in the presence of regulatory proteins. Although we observed subtle changes in the second-order rate of

ATP-induced dissociation between regulated and unregulated filaments (**Fig. 3d**), these

changes are irrelevant at [ATP] > 50 μM where the rate of ADP release limits

detachment. Therefore, in the functioning heart, regulatory proteins do not modulate the

kinetics of the myosin working stroke. Moreover, our data show that the load-dependent

rate of ADP release limits actomyosin dissociation, and this is unchanged by regulatory

proteins (**Figs. 3b-c and 4**). Finally, we show that the mechanics of the working stroke

are unchanged by regulatory proteins (**Fig. 2**). Taken together, our data are consistent

with tropomyosin primarily serving a role in sterically blocking the calcium-dependent

interactions between cardiac myosin and the thin filament under working conditions in

the heart (46,47).

Although not relevant to working conditions in the heart, our results reveal that

regulatory proteins modulate the kinetics of ATP-induced actomyosin dissociation at low

ATP concentrations. To measure the substeps of the working stroke, we conducted

optical trapping experiments at very low ATP concentrations not experienced in the cell,

since this slows mechanical transitions that are too fast to resolve at physiologically

relevant ATP (28). While our results show that regulatory proteins do not change the

size of the working stroke or the coupling between kinetics and mechanics (**Fig. 2**), they

also reveal that regulatory proteins slow the dissociation of actomyosin at very low ATP

concentrations (**Fig. 3a**). The time reverse ensemble averages demonstrate that

actomyosin dissociation is limited by the transition from the second substep to the

detached state (**Figs. 2c-d**), which is similar to the second-order rate of ATP-induced

actomyosin dissociation measured using stopped flow techniques (**Fig. 3b**) (29-31). All

of these rates (actomyosin detachment rate (**Fig. 3a**), time reverse ensemble average

rate (**Fig. 2c-d**), and ATP-induced dissociation rate measured in the stopped flow (**Fig.**

3b)) are similar, consistent with the rate of ATP-induced dissociation limiting

detachment at low ATP. Moreover, for each of these measurements, the rates are

significantly slower in the presence of regulatory proteins than in the absence (**Fig. 3d**).

Taken together, these data demonstrate that at low ATP concentrations, regulatory

proteins slow actomyosin detachment due to slowed ATP-induced dissociation.

While not relevant to physiology, the observation that regulatory proteins can

tune ATP-induced dissociation has interesting implications. The tuning of detachment

kinetics by regulatory proteins cannot be fully explained by a simple steric blocking

mechanism, where tropomyosin only blocks the interactions between myosin and the

thin filament (46). The exact mechanism of this kinetic tuning is not known; however, it

has been proposed that tropomyosin can interact directly with myosin (48), and domain-

specific deletion studies of tropomyosin have demonstrated that myosin can modulate

tropomyosin binding to the thin filament (49,50). Moreover, recent structural studies of

muscle proteins have demonstrated specific interactions between loop-4 of myosin and

tropomyosin (51,52). It is possible that these interactions are allosterically coupled to

changes in the nucleotide binding pocket of myosin, given the complex allosteric

networks between the nucleotide binding site and the actin binding domain (53).

Moreover, our results are consistent with previous optical trapping work examining

skeletal muscle myosin at low ATP concentrations, which demonstrated that regulatory

proteins slow the rate of actomyosin detachment (20). That being said, our data

collected at saturating ATP concentrations demonstrate that kinetic tuning by regulatory

proteins is likely not relevant under working conditions in cardiac muscle, and the

effects of regulatory proteins in cardiac muscle are well described by a steric blocking

model.

The effects of regulatory proteins on actomyosin appear to be isoform specific

Here, we saw that at physiologically relevant ATP concentrations, tropomyosin

and troponin do not appreciably affect the β-cardiac myosin working stroke.

Interestingly, the effects of regulatory proteins on myosin appear to depend on the

specific protein isoforms used. There are many distinct tropomyosin isoforms expressed

in eukaryotic cells, with tissue-specific expression patterns, and even within the same

cell, different tropomyosin isoforms localize to different subcellular actin pools (54).

Moreover, while all myosin isoforms can associate with actin, they preferentially interact

with certain actin structures in an isoform-specific manner (55). The biophysical role of

tropomyosin appears to vary with both the myosin and tropomyosin isoform, where

some tropomyosin isoforms inhibit the interactions of certain myosin isoforms with actin

(55,56), while others promote these interactions (19,57). This isoform specificity has

been proposed to help localize specialized motors to specific regions of the cell (58).

The molecular roles of tropomyosins on myosin motor function can extend

beyond steric effects. For example, decoration of actin with Tpm1p slows the rate of

ADP release from the myosin-V isoform Myo2p, and this slowing of ADP release

kinetics enables Myo2p to processively walk on tropomyosin decorated actin filaments,

something it will not do on bare actin (19). Even within the family of myosin-II motors,

there is evidence that regulatory proteins could tune the load-dependent kinetics of

myosin in an isoform specific manner. For example, non-muscle myosin-IIA's force

sensitivity is increased by tropomyosin Tm4.2 (18). In the case of skeletal muscle

myosin, some studies suggested that tropomyosin decreases the step size of myosin by

inhibiting the ability of both myosin heads to bind to the thin filament (20), while other

studies have not seen this effect (21,22). The working stroke size that we measured for

β-cardiac myosin in the presence of regulatory proteins is indistinguishable from that

measured in the absence of regulatory proteins with both one-headed and two-headed

myosin constructs (4,17,24,25), so we do not believe that our inability to observe a

difference in mechanics is related to the two-headed nature of the construct used here

(59).

Conclusions

Our results clearly demonstrate that under physiologically relevant ATP

concentrations, regulatory proteins do not cause appreciable changes in the mechanics

or kinetics of the β-cardiac myosin working stroke; however, they can tune myosin's

kinetics at low ATP concentrations, suggesting effects beyond a simple steric blocking

mechanism. This has important implications for both our understanding of the

mechanism of muscle regulation and mathematical modeling of muscle contraction,

which relies on accurate, sensitive measurements of these parameters.

Materials and Methods

Protein expression and purification

Cardiac actin and myosin were purified from cryoground porcine ventricles as previously described (10). Human troponin and tropomyosin were expressed recombinantly in E. coli, purified, and complexed as described previously (10). Myosin subfragment-1 (S1) for spectroscopic measurement was prepared by limited proteolysis using chymotrypsin as previously described, and N-(1-Pyrene)Iodoacetamide-labeled actin was prepared as previously described (10). Protein concentrations were determined spectroscopically. For all experiments, at least 2 separate protein preparations were used.

Stopped-flow measurements of ADP release and ATP-induced actomyosin dissociation

Stopped-flow experiments were performed in a SX-20 apparatus (Applied Photophysics). All experiments were conducted at 20°C in high calcium buffer (pCa 4) containing 25 mM KCl, 5 mM free $MgCl_2$, 60 mM MOPS pH 7.0, 2 mM EGTA, 1 mM DTT, and 2.15 mM $CaCl_2$, where the concentration of free calcium was calculated using MaxChelator (60).

ADP release experiments were performed as described previously (7,45,61). Briefly, 1 μM phalloidin-stabilized, pyrene-labeled actin, 1.5 μM tropomyosin (when appropriate), 1.5 μM troponin (when appropriate), 1 μM S1 myosin, and 100 μM Mg*ADP were rapidly mixed with 5 mM Mg*ATP. This caused an increase in fluorescence that was well fit by a single exponential function, where the rate equals the rate of ADP release (35). Each experiment consisted of 5 technically repeated

measurements, and the 3 experiments were used to calculate the mean and standard

deviation. A two-tailed Student's t-test was used for statistical testing.

The rate of ATP-induced actomyosin dissociation was measured as previously

described (61). Briefly, 1 μM phalloidin-stabilized, pyrene-labeled actin, 1.5 μM

tropomyosin (when appropriate), 1.5 μM troponin (when appropriate), 1 μM S1 myosin,

and 0.04 U/mL apyrase were rapidly mixed with varying concentrations of Mg*ATP. The

resultant fluorescence transients were best fit by the sum of two exponential functions,

as previously described (35). The amplitude of the fast phase was fixed to prevent

artifacts due to the dead time of the instrument. As has been shown before, the rates of

the fast and slow phases were hyperbolically related to the concentration of ATP. Data

were interpreted according to the scheme (35):

$$AM \underset{k_{-\alpha}}{\overset{k_{+\alpha}}{\rightleftharpoons}} AM' \overset{K_1'}{\rightleftharpoons} AM(ATP) \overset{k_{+2}'}{\longrightarrow} A^*M.ATP \overset{k_{+3}'}{\longrightarrow} A^* + M.ATP$$

The fast phase of ATP-induced dissociation was modelled as the formation of a collision

complex between actomyosin and ATP that is in rapid equilibrium (K_1') followed by an

irreversible isomerization and rapid dissociation (k_{+2}'). The rate of the fast phase, k_{fast}, is

hyperbolically related to the ATP concentration by:

$k_{fast} = \dfrac{k_{+2}' \cdot [ATP]}{1/K_1' + [ATP]}$ Equation 2

At low [ATP], the second order rate of ATP-induced dissociation is given by K_1' * k_{+2}'.

The concentration of ATP was measured spectroscopically for all experiments. The rate

of ATP-induced dissociation was measured over a full range of concentrations 3 times,

and each time, the fitted parameters were extracted. Reported values are the average

of these 3 trials and the error is the standard deviation. Statistical testing was performed

using a 2-tailed Student's t-test.

In vitro motility assays

In vitro motility assays were performed as described (10). For the experiments

using biotinylated actin, all experimental protocols were identical except 10% biotin-

actin (Cytoskeleton) was added to 2 μM G-actin and stabilized with

tetramethylrhodamine isothiocyanate-labeled phalloidin in KMg25 buffer (60 mM MOPS

pH 7.0, 25 mM KCl, 2 mM EGTA, 4 mM MgCl$_2$, and 1 mM DTT). The concentration of

free calcium was calculated using MaxChelator (60). Reported values are the average

and standard deviations of the speeds from at least 3 separate days of experiments.

Statistical testing was performed using a Mann-Whitney test.

Optical trapping experiments

Experiments were performed on a custom-built, microscope free dual-beam

optical trap described previously (29). These experiments utilized the three-bead

geometry in which a thin filament is held between two optically trapped beads and

lowered on to a surface bound bead that is sparsely coated with myosin (27,28).

Tropomyosin was dialyzed into KMg25 buffer the night before the experiment. Actin was

attached to beads using a biotin-streptavidin linkage, where actin contained 10%

biotinylated actin and polystyrene beads were coated with streptavidin, as previously

described (28,29). Flow cells were coated sparsely with beads as previously described

(28,29). Flow cells were loaded with myosin (1-7 nM in KMg25 with 200 mM KCl to

prevent myosin filament formation) for 5 minutes and the surface was blocked with 1

mg/mL BSA for 5 minutes. This was followed by activation buffer. For low ATP

experiments, the activation buffer contained KMg25 with 1 μM ATP, 192 U/mL glucose

oxidase, 48 μg/mL catalase, 1 mg/mL glucose, and ~25 pM biotin-rhodamine-phalloidin

actin. For the high ATP experiments, conditions were identical, except 1 mM Mg*ATP

was used. When appropriate, troponin and tropomyosin were also included at 200 nM.

The concentration of free calcium was calculated using MaxChelator (60). This was

followed by 4 μL of streptavidin beads. Flow cells were then sealed with vacuum grease

as previously described and data were collected within 90 minutes of sealing (29).

Surface-bound beads were probed for binding interactions using small

oscillations of the stage position that were stopped once data collection began. Data

were collected at 20 kHz and filtered to 10 kHz according to the Nyquist criterion. For

each bead-actin-bead assembly, the trap stiffness was calculated from fitting of the

power spectrum as previously described (28).

Implementation of the isometric optical clamp feedback

Here, we used an all-digital implementation of an isometric feedback clamp (36).

In an isometric optical clamp feedback experiment, the position of one bead (the

transducer) is continuously sampled, and deviations from its original setpoint position

are compensated for by moving the second (motor) bead using acoustic optical

deflectors (AODs, Gooch and Housego). The positions of the beads were recorded

using quadrant photodiodes, the feedback calculations were digitally performed on a

field programmable gate array (FPGA) board (National Instruments PCIe-7852), and the

laser controlling the motor bead was translated using AODs.

The error signal used for the feedback, V_t, is the time filtered positional error for

the current sample period given by:

$$V_t = K_p E_t + K_i \sum_{k=t}^{t-W} E_k$$

where K_p is the user defined proportional gain, K_i is the user defined integral gain, E_t is

the current sample's absolute error from the setpoint, and W is the user defined

integration window (up to a 255-sample memory). The compensating position for the

motor bead in frequency units is then calculated by the FPGA and transmitted to the

parallel port interface of the Digital Frequency Synthesizer (Analog Devices

AD9912A/PCBZ) which controls the beam deflection angle from the AOD. The feedback

loop can run at a maximal speed of 50 kHz, limited by parallel port cable capacitance.

The time constant for the feedback response time was set as previously

described (36). Briefly, a bead-actin-bead dumbbell was held in the dual beam traps, a

square wave was injected into the transducer bead channel, and then movement of the

motor bead by the feedback system was monitored. The proportional and integral gains

were empirically adjusted to give a response time of ~5 ms without introducing

oscillations into the system.

Analysis of single molecule data

All data from optical trapping experiments were analyzed using our custom-built

MATLAB program, SPASM, as previously described (29). Briefly, binding interactions

between myosin and the thin filament were identified using a peak-to-peak covariance

threshold, and the initiation and termination times of binding were identified using a

changepoint algorithm. Data traces were excluded if the separation between the bound

and unbound populations of the covariance histogram was not well defined. To improve

the signal-to-noise ratio, the signal from both beads was summed and divided by 2 as

previously described (29). Ensemble averages and histograms of binding interactions

were generated as previously described (28-30). Ensemble averages were fit by single

exponential functions in MATLAB until the signal plateaued. Cumulative distributions of

step sizes and attachment durations were fit with cumulative functions for the Gaussian

and exponential functions, respectively. Statistical testing for normality was done using

a Shapiro-Wilk test. Statistical testing of the step sizes was done using a 2-tailed

Student's t-test of individual binding interactions. 95% confidence intervals for the

detachment rate were calculated by bootstrapping of individual binding interaction

durations, and statistical significance was calculated according to (62).

In the optical trap, actomyosin remains attached until ADP is released and ATP

induces actomyosin dissociation (35):

$$AM.ADP \xrightarrow{k_{+ADP}} AM \xrightarrow{k_{+ATP}} A + M.ATP$$

The attachment duration, t_{on}, is given by

$$t_{on} = \frac{1}{k_{det}} = \frac{1}{k_{+ADP}} + \frac{1}{k_{+ATP}} = \frac{1}{k_{+ADP}} + \frac{1}{k'_{+2} * [ATP]}$$

At low ATP, the detachment rate, k_{det}, will be dominated by the second order rate of

ATP binding, while at saturating ATP, the detachment rate will be limited by the rate of

ADP release.

For isometric optical clamp experiments, data were collected at saturating ATP

concentrations, where the rate of ADP release limits the rate of actomyosin dissociation

(63). Binding interactions were identified using a variance threshold, set by the position

of the transducer bead, and the force exerted by the motor bead and the attachment

duration were measured. The relationship between the force, F, and the load dependent

detachment rate, k(F) was modeled using the Bell equation (38):

$$k(F) = k_0 * \exp\left(\frac{-F * d}{k_B * T}\right)$$

528 Where k_0 is the rate of the primary force sensitive transition in the absence of force, d is

529 the distance to the transition state, and k_B*T is the thermal energy. The distribution of

530 attachment durations is exponentially distributed at each force, and therefore follows the

531 following probability density distribution (37):

$$k(F, t) = k(F) * \exp\left(-k(F) * t\right)$$

Maximum likelihood estimation (MLE) was used to determine the most likely values of k_0

and d, as previously described (37). 95% confidence intervals for parameter values

fitting were determined using 1000 rounds of bootstrapping simulations, and hypothesis

testing was performed using the difference in the means and the variances of these

distributions as previously described (62).

Data Availability, Reproducibility, and Software

The data and code necessary to generate the figures and statistical tests in this

manuscript are on Github (https://github.com/GreenbergLab/2023-thin-filament-

trapping). Optical trapping data were analyzed with our previously published SPASM

program by running the MATLAB source code from GitHub

(https://github.com/GreenbergLab/SPASM). Similarly, the isometric force clamp data

was analyzed with a modified version of SPASM (in the GitHub repository). Subsequent

figure generation and related analyses were performed in R (version 4.2.2) (64) running

under NixOS 22.05 (Linux). The git repository hosts a "Nix Flake", a reproducible

developmental shell environment with pinned version dependencies. For NixOS users,

users of the Nix package manager on macOS or the Windows Subsystem for Linux

(WSL), an ephemeral shell can be spawned by running `*nix develop*

github:GreenbergLab/2023-thin-filament-trapping?dir=code` in the terminal. Additional R

packages used include the following: "readr_2.1.3", "readxl_1.4.0", "data.table_1.14.6",

"here_1.0.1", "ggpubr_0.5.0", "ggtext_0.1.2", "purrr_0.3.5", "tidyr_1.2.1", "dplyr_1.0.10",

"tibble_3.1.8",   "cowplot_1.1.1",   and   "ggplot2_3.4.0".

Funding Acknowledgements

This work was supported by the National Institutes of Health (R01 HL141086 to M.J.G.)

and the Children's Discovery Institute of Washington University and St. Louis Children's

Hospital (PM-LI-2019-829 to M.J.G.).

Conflict of interest

All experiments were conducted in the absence of any financial relationships that could

be construed as potential conflicts of interest.

Author contributions

S.R.C.S., W.T.S., B.S., and M.J.G. designed the experiments. S.R.C.S., B.S., and

S.K.B. conducted and analyzed the optical trapping experiments. W.T.S. designed and

tested the optical trapping system, including the feedback system. T.B., B.S., and

M.J.G. contributed to tools for data analysis. S.R.C.S. drafted the first draft of the

manuscript with M.J.G. All authors contributed to the analysis of data and writing/editing

of the manuscript. M.J.G. procured funding and oversaw the project.

 Figure Legends

 Figure 1: Reconstitution of thin filaments using biotinylated actin does not affect

 thin filament regulation. a. In vitro motility assay boxplots showing the speed of

 regulated thin filament translocation with and without biotin-labeled actin at pCa 4.

 Individual points show the velocity of >25 thin filaments measured across 3 separate

 experiments. The thick middle line of the boxplot shows the median value, the

 top/bottom of the box are the 1st and 3rd quartiles, and the whiskers extend to 1.5 times

 the interquartile range. There is no difference in speed with or without biotinylated actin

 (P = 0.12, Mann-Whitney test). No movement was seen at pCa 9 with or without

 biotinylated actin. **b.** Representative optical trapping traces of regulated thin filaments

 containing 10% biotinylated actin conducted at low (pCa 9, orange) and high calcium

 (pCa 4, blue). At low calcium, binding events were very rare and at pCa 4 binding

 events were frequent (grey lines denote binding interactions detected by the analysis

 program), demonstrating regulation.

 Figure 2: Optical trapping experiments at 1 μM ATP reveal no changes in working

 stroke mechanics with regulated thin filaments. Data are from 364 interactions

 detected from 10 myosin molecules for the unregulated condition and 491 interactions

 from 9 myosin molecules for the regulated condition. **a-b.** Representative optical

 trapping data for unregulated (black) and regulated (green) thin filaments. Grey lines

 indicate actomyosin interactions identified by automated event detection (see **Methods**

 for details). **c-d.** Ensemble averages of the working stroke for myosin interacting with

 unregulated (left, black) and regulated (right, green) thin filaments. The rates of the

exponential functions fit to the time forward (k_f) and time reverse (k_r) averages are

shown along with the magnitudes of the two substeps. **e-g.** Cumulative distributions

derived from individual binding interactions for the (**e**) total displacement, (**f**) first

substep, and (**g**) second substep. Each plot reports the sample mean, standard

deviation, and P-values from t-tests.

Figure 3: Regulatory proteins affect kinetics at low ATP due to slowing of ATP-

induced dissociation. a. Individual actomyosin attachment durations obtained in the

optical trap are plotted as cumulative distributions for unregulated (black) and regulated

(green) thin filaments. Distributions are fit with single exponential functions (dashed

lines) to obtain the actomyosin detachment rate. Regulatory proteins significantly slow

the rate of detachment (see table in **d**; value is the fitted rate, error is a bootstrapped

95% confidence interval, and P-value is calculated as described in the **Methods**). **b.**

Representative fluorescence transients from stopped-flow experiments measuring the

rate of ADP release from actomyosin. Fits show single exponential fits to the average of

5 transients. There is no difference in the rate of ADP release measured using regulated

or unregulated thin filaments (see table in **d**). **c.** Fast phase of ATP-induced dissociation

of myosin from regulated and unregulated thin filaments (see **Methods** for details).

Each point represents the average of 5 technical repeats collected on one day. 3

separate experimental days were used. Solid lines show fitting of Eq. 2. **d.** Table of

parameters obtained for all kinetic measurements. The row letters indicate from which

figure panel the values are derived. For the stopped-flow transient kinetics (marked "b"

or "c"), the reported values are the average and standard deviations of 3 separate

experimental days and P-values were calculated using a Student's t-test.

Figure 4: Regulatory proteins do not affect myosin's load-dependent kinetics at

physiological ATP. a. Representative traces collected using the isometric optical

clamp. The motor bead ("M") is moved to keep the transducer bead ("T") at an isometric

position. During a binding interaction, the average force and the attachment duration

were recorded. **b.** Attachment durations as a function of force measured for unregulated

(black) and regulated (green) thin filaments. Each point represents a single binding

interaction. 393 binding interactions were observed from 10 myosin molecules for the

unregulated condition and 611 binding interactions were observed from 20 myosin

molecules for the regulated condition. Data were fitted with the Bell equation using

maximum likelihood estimation to obtain k_0 and d. Error bars are the 95% confidence

intervals obtained from 1000 rounds of bootstrapping simulations. These parameters

were not significantly different in the presence of regulatory proteins (P = 0.16 for k_0 and

P = 0.49 for d; see **Methods** for details on statistics and fitting).

Figure 1

640 **Figure 2**

	Term	Unregulated	Regulated	P-Value
a	t_{on} (s^{-1})	5.5 (-0.4/+0.4)	3.9 (-0.3/+0.4)	0.002
b	k_{-ADP} (s^{-1})	79 ± 1	76 ± 5	0.45
c	$1/K_1'$ (µM^{-1})	306 ± 59	496 ± 82	0.008
c	k_{+2}' (s^{-1})	1220 ± 56	1084 ± 49	0.01
c	$K_1' \cdot k_{+2}'$ (µM$^{-1} \cdot$s^{-1})	4.0 ± 0.8	2.2 ± 0.4	0.04

Figure 3

649   **Figure 4**

References

1. Greenberg, M. J., and J. C. Tardiff. 2021. Complexity in genetic cardiomyopathies and new approaches for mechanism-based precision medicine. *J Gen Physiol*. 153(3), doi: 10.1085/jgp.202012662.

2. Barrick, S. K., and M. J. Greenberg. 2021. Cardiac myosin contraction and mechanotransduction in health and disease. *J Biol Chem*. 297(5):101297, doi: 10.1016/j.jbc.2021.101297.

3. McKillop, D. F., and M. A. Geeves. 1993. Regulation of the interaction between actin and myosin subfragment 1: evidence for three states of the thin filament. *Biophys J*. 65(2):693-701, doi: 10.1016/S0006-3495(93)81110-X.

4. Greenberg, M. J., H. Shuman, and E. M. Ostap. 2014. Inherent force-dependent properties of beta-cardiac myosin contribute to the force-velocity relationship of cardiac muscle. *Biophys J*. 107(12):L41-L44, doi: 10.1016/j.bpj.2014.11.005.

5. Sung, J., S. Nag, K. I. Mortensen, C. L. Vestergaard, S. Sutton, K. Ruppel, H. Flyvbjerg, and J. A. Spudich. 2015. Harmonic force spectroscopy measures load-dependent kinetics of individual human beta-cardiac myosin molecules. *Nat Commun*. 6:7931, doi: 10.1038/ncomms8931.

6. Anderson, R. L., D. V. Trivedi, S. S. Sarkar, M. Henze, W. Ma, H. Gong, C. S. Rogers, J. M. Gorham, F. L. Wong, M. M. Morck, J. G. Seidman, K. M. Ruppel, T. C. Irving, R. Cooke, E. M. Green, and J. A. Spudich. 2018. Deciphering the super relaxed state of human beta-cardiac myosin and the mode of action of mavacamten from myosin molecules to muscle fibers. *Proc Natl Acad Sci U S A*. 115(35):E8143-E8152, doi: 10.1073/pnas.1809540115.

7. Barrick, S. K., L. Greenberg, and M. J. Greenberg. 2021. A troponin T variant linked with pediatric dilated cardiomyopathy reduces the coupling of thin filament activation to myosin and calcium binding. *Mol Biol Cell*. 32(18):1677-1689, doi: 10.1091/mbc.E21-02-0082.

8. Ezekian, J. E., S. R. Clippinger, J. M. Garcia, Q. Yang, S. Denfield, A. Jeewa, W. J. Dreyer, W. Zou, Y. Fan, H. D. Allen, J. J. Kim, M. J. Greenberg, and A. P. Landstrom. 2020. Variant R94C in TNNT2-Encoded Troponin T Predisposes to Pediatric Restrictive Cardiomyopathy and Sudden Death Through Impaired Thin Filament Relaxation Resulting in Myocardial Diastolic Dysfunction. *J Am Heart Assoc*. 9(5):e015111, doi: 10.1161/JAHA.119.015111.

9. Clippinger, S. R., P. E. Cloonan, W. Wang, L. Greenberg, W. T. Stump, P. Angsutararux, J. M. Nerbonne, and M. J. Greenberg. 2020. Mechanical dysfunction induced by a hypertrophic cardiomyopathy mutation is the primary driver of cellular adaptation. *bioRxiv*.2020.2005.2004.067181, doi: 10.1101/2020.05.04.067181.

10. Clippinger, S. R., P. E. Cloonan, L. Greenberg, M. Ernst, W. T. Stump, and M. J. Greenberg. 2019. Disrupted mechanobiology links the molecular and cellular phenotypes in familial dilated cardiomyopathy. *Proc Natl Acad Sci U S A*. 116(36):17831-17840, doi: 10.1073/pnas.1910962116.

11. He, H., T. Baka, J. Balschi, A. S. Motani, K. K. Nguyen, Q. Liu, R. Slater, B. Rock, C. Wang, C. Hale, G. Karamanlidis, J. J. Hartman, F. I. Malik, J. D. Reagan, and I. Luptak. 2022. Novel Small-Molecule Troponin Activator Increases Cardiac Contractile Function Without Negative Impact on Energetics. *Circ Heart Fail*. 15(3):e009195, doi: 10.1161/CIRCHEARTFAILURE.121.009195.

12. Malik, F. I., J. J. Hartman, K. A. Elias, B. P. Morgan, H. Rodriguez, K. Brejc, R. L. Anderson, S. H. Sueoka, K. H. Lee, J. T. Finer, R. Sakowicz, R. Baliga, D. R. Cox, M. Garard, G. Godinez, R. Kawas, E. Kraynack, D. Lenzi, P. P. Lu, A. Muci, C. Niu, X. Qian, D. W. Pierce, M. Pokrovskii, I. Suehiro, S. Sylvester, T. Tochimoto, C. Valdez, W. Wang, T. Katori, D. A. Kass, Y. T. Shen, S. F. Vatner, and D. J. Morgans. 2011. Cardiac myosin activation: a potential therapeutic approach for systolic heart failure. *Science*. 331(6023):1439-1443, doi: 10.1126/science.1200113.

697   13.   Green, E. M., H. Wakimoto, R. L. Anderson, M. J. Evanchik, J. M. Gorham, B. C. Harrison, M.
698   Henze, R. Kawas, J. D. Oslob, H. M. Rodriguez, Y. Song, W. Wan, L. A. Leinwand, J. A. Spudich, R.
699   S. McDowell, J. G. Seidman, and C. E. Seidman. 2016. A small-molecule inhibitor of sarcomere
700   contractility suppresses hypertrophic cardiomyopathy in mice. *Science*. 351(6273):617-621, doi:
701   10.1126/science.aad3456.
702   14.   Deacon, J. C., M. J. Bloemink, H. Rezavandi, M. A. Geeves, and L. A. Leinwand. 2012.
703   Identification of functional differences between recombinant human alpha and beta cardiac
704   myosin motors. *Cell Mol Life Sci*. 69(13):2261-2277, doi: 10.1007/s00018-012-0927-3.
705   15.   Palmiter, K. A., M. J. Tyska, D. E. Dupuis, N. R. Alpert, and D. M. Warshaw. 1999. Kinetic
706   differences at the single molecule level account for the functional diversity of rabbit cardiac
707   myosin isoforms. *J Physiol*. 519 Pt 3:669-678, doi: 10.1111/j.1469-7793.1999.0669n.x.
708   16.   Liu, C., M. Kawana, D. Song, K. M. Ruppel, and J. A. Spudich. 2018. Controlling load-dependent
709   kinetics of beta-cardiac myosin at the single-molecule level. *Nat Struct Mol Biol*. 25(6):505-514,
710   doi: 10.1038/s41594-018-0069-x.
711   17.   Woody, M. S., M. J. Greenberg, B. Barua, D. A. Winkelmann, Y. E. Goldman, and E. M. Ostap.
712   2018. Positive cardiac inotrope omecamtiv mecarbil activates muscle despite suppressing the
713   myosin working stroke. *Nat Commun*. 9(1):3838, doi: 10.1038/s41467-018-06193-2.
714   18.   Hundt, N., W. Steffen, S. Pathan-Chhatbar, M. H. Taft, and D. J. Manstein. 2016. Load-dependent
715   modulation of non-muscle myosin-2A function by tropomyosin 4.2. *Sci Rep*. 6:20554, doi:
716   10.1038/srep20554.
717   19.   Hodges, A. R., E. B. Krementsova, C. S. Bookwalter, P. M. Fagnant, T. E. Sladewski, and K. M.
718   Trybus. 2012. Tropomyosin is essential for processive movement of a class V myosin from
719   budding yeast. *Curr Biol*. 22(15):1410-1416, doi: 10.1016/j.cub.2012.05.035.
720   20.   Kad, N. M., S. Kim, D. M. Warshaw, P. VanBuren, and J. E. Baker. 2005. Single-myosin crossbridge
721   interactions with actin filaments regulated by troponin-tropomyosin. *Proc Natl Acad Sci U S A*.
722   102(47):16990-16995, doi: 10.1073/pnas.0506326102.
723   21.   Longyear, T., S. Walcott, and E. P. Debold. 2017. The molecular basis of thin filament activation:
724   from single molecule to muscle. *Sci Rep*. 7(1):1822, doi: 10.1038/s41598-017-01604-8.
725   22.   Unger, M., and E. P. Debold. 2019. Acidosis decreases the Ca(2+) sensitivity of thin filaments by
726   preventing the first actomyosin interaction. *Am J Physiol Cell Physiol*. 317(4):C714-C718, doi:
727   10.1152/ajpcell.00196.2019.
728   23.   Nag, S., R. F. Sommese, Z. Ujfalusi, A. Combs, S. Langer, S. Sutton, L. A. Leinwand, M. A. Geeves,
729   K. M. Ruppel, and J. A. Spudich. 2015. Contractility parameters of human beta-cardiac myosin
730   with the hypertrophic cardiomyopathy mutation R403Q show loss of motor function. *Sci Adv*.
731   1(9):e1500511, doi: 10.1126/sciadv.1500511.
732   24.   Woody, M. S., D. A. Winkelmann, M. Capitanio, E. M. Ostap, and Y. E. Goldman. 2019. Single
733   molecule mechanics resolves the earliest events in force generation by cardiac myosin. *Elife*. 8,
734   doi: 10.7554/eLife.49266.
735   25.   Snoberger, A., B. Barua, J. L. Atherton, H. Shuman, E. Forgacs, Y. E. Goldman, D. A. Winkelmann,
736   and E. M. Ostap. 2021. Myosin with hypertrophic cardiac mutation R712L has a decreased
737   working stroke which is rescued by omecamtiv mecarbil. *eLife*. 10, doi: 10.7554/eLife.63691.
738   26.   Kron, S. J., and J. A. Spudich. 1986. Fluorescent actin filaments move on myosin fixed to a glass
739   surface. *Proc Natl Acad Sci U S A*. 83(17):6272-6276, doi: 10.1073/pnas.83.17.6272.
740   27.   Finer, J. T., R. M. Simmons, and J. A. Spudich. 1994. Single myosin molecule mechanics:
741   piconewton forces and nanometre steps. *Nature*. 368(6467):113-119, doi: 10.1038/368113a0.
742   28.   Greenberg, M. J., H. Shuman, and E. M. Ostap. 2017. Measuring the Kinetic and Mechanical
743   Properties of Non-processive Myosins Using Optical Tweezers. *Methods Mol Biol*. 1486:483-509,
744   doi: 10.1007/978-1-4939-6421-5_19.

29. Blackwell, T., W. T. Stump, S. R. Clippinger, and M. J. Greenberg. 2021. Computational Tool for
Ensemble Averaging of Single-Molecule Data. *Biophys J.* 120(1):10-20, doi:
10.1016/j.bpj.2020.10.047.

30. Veigel, C., L. M. Coluccio, J. D. Jontes, J. C. Sparrow, R. A. Milligan, and J. E. Molloy. 1999. The
motor protein myosin-I produces its working stroke in two steps. *Nature.* 398(6727):530-533,
doi: 10.1038/19104.

31. Chen, C., M. J. Greenberg, J. M. Laakso, E. M. Ostap, Y. E. Goldman, and H. Shuman. 2012.
Kinetic schemes for post-synchronized single molecule dynamics. *Biophys J.* 102(6):L23-25, doi:
10.1016/j.bpj.2012.01.054.

32. Barany, M. 1967. ATPase activity of myosin correlated with speed of muscle shortening. *J Gen
Physiol.* 50(6):Suppl:197-218, doi: 10.1085/jgp.50.6.197.

33. Siemankowski, R. F., M. O. Wiseman, and H. D. White. 1985. ADP dissociation from actomyosin
subfragment 1 is sufficiently slow to limit the unloaded shortening velocity in vertebrate muscle.
Proc Natl Acad Sci U S A. 82(3):658-662, doi: 10.1073/pnas.82.3.658.

34. Papadaki, M., T. Kampaengsri, S. K. Barrick, S. G. Campbell, D. von Lewinski, P. P. Rainer, S. P.
Harris, M. J. Greenberg, and J. A. Kirk. 2022. Myofilament glycation in diabetes reduces
contractility by inhibiting tropomyosin movement, is rescued by cMyBPC domains. *J Mol Cell
Cardiol.* 162:1-9, doi: 10.1016/j.yjmcc.2021.08.012.

35. De La Cruz, E. M., and E. M. Ostap. 2009. Kinetic and equilibrium analysis of the myosin ATPase.
Methods Enzymol. 455:157-192, doi: 10.1016/S0076-6879(08)04206-7.

36. Takagi, Y., E. E. Homsher, Y. E. Goldman, and H. Shuman. 2006. Force generation in single
conventional actomyosin complexes under high dynamic load. *Biophys J.* 90(4):1295-1307, doi:
10.1529/biophysj.105.068429.

37. Woody, M. S., J. H. Lewis, M. J. Greenberg, Y. E. Goldman, and E. M. Ostap. 2016. MEMLET: An
Easy-to-Use Tool for Data Fitting and Model Comparison Using Maximum-Likelihood Estimation.
Biophys J. 111(2):273-282, doi: 10.1016/j.bpj.2016.06.019.

38. Bell, G. I. 1978. Models for the specific adhesion of cells to cells. *Science.* 200(4342):618-627,
doi: 10.1126/science.347575.

39. Laakso, J. M., J. H. Lewis, H. Shuman, and E. M. Ostap. 2008. Myosin I can act as a molecular
force sensor. *Science.* 321(5885):133-136, doi: 10.1126/science.1159419.

40. Stewart, L. C., R. Deslauriers, and V. V. Kupriyanov. 1994. Relationships between cytosolic [ATP],
[ATP]/[ADP] and ionic fluxes in the perfused rat heart: A 31P, 23Na and 87Rb NMR study. *J Mol
Cell Cardiol.* 26(10):1377-1392, doi: 10.1006/jmcc.1994.1156.

41. Wu, F., J. Zhang, and D. A. Beard. 2009. Experimentally observed phenomena on cardiac
energetics in heart failure emerge from simulations of cardiac metabolism. *Proc Natl Acad Sci U
S A.* 106(17):7143-7148, doi: 10.1073/pnas.0812768106.

42. Beer, M., T. Seyfarth, J. Sandstede, W. Landschutz, C. Lipke, H. Kostler, M. von Kienlin, K. Harre,
D. Hahn, and S. Neubauer. 2002. Absolute concentrations of high-energy phosphate metabolites
in normal, hypertrophied, and failing human myocardium measured noninvasively with (31)P-
SLOOP magnetic resonance spectroscopy. *J Am Coll Cardiol.* 40(7):1267-1274, doi:
10.1016/s0735-1097(02)02160-5.

43. Starling, R. C., D. F. Hammer, and R. A. Altschuld. 1998. Human myocardial ATP content and in
vivo contractile function. *Mol Cell Biochem.* 180(1-2):171-177.

44. Yabe, T., K. Mitsunami, T. Inubushi, and M. Kinoshita. 1995. Quantitative measurements of
cardiac phosphorus metabolites in coronary artery disease by 31P magnetic resonance
spectroscopy. *Circulation.* 92(1):15-23, doi: 10.1161/01.cir.92.1.15.

45. Papadaki, M., T. Kampaengsri, S. K. Barrick, S. G. Campbell, D. von Lewinski, P. P. Rainer, S. P.
Harris, M. J. Greenberg, and J. A. Kirk. 2021. Myofilament glycation in diabetes reduces

contractility by inhibiting tropomyosin movement, is rescued by cMyBPC domains. *Journal of*
molecular and cellular cardiology. 162:1-9, doi: 10.1016/j.yjmcc.2021.08.012.

46.   Vibert, P., R. Craig, and W. Lehman. 1997. Steric-model for activation of muscle thin filaments. *J*
Mol Biol. 266(1):8-14, doi: 10.1006/jmbi.1996.0800.

47.   Lehman, W., R. Craig, and P. Vibert. 1994. Ca(2+)-induced tropomyosin movement in Limulus
thin filaments revealed by three-dimensional reconstruction. *Nature*. 368(6466):65-67, doi:
10.1038/368065a0.

48.   Lehrer, S. S. 1994. The regulatory switch of the muscle thin filament: Ca2+ or myosin heads? *J*
Muscle Res Cell Motil. 15(3):232-236, doi: 10.1007/BF00123476.

49.   Hitchcock-DeGregori, S. E., Y. Song, and J. Moraczewska. 2001. Importance of internal regions
and the overall length of tropomyosin for actin binding and regulatory function. *Biochemistry*.
40(7):2104-2112, doi: 10.1021/bi002421z.

50.   Landis, C., N. Back, E. Homsher, and L. S. Tobacman. 1999. Effects of tropomyosin internal
deletions on thin filament function. *J Biol Chem*. 274(44):31279-31285, doi:
10.1074/jbc.274.44.31279.

51.   Doran, M. H., M. J. Rynkiewicz, E. Pavadai, S. M. L. Bodt, D. Rasicci, J. R. Moore, C. M. Yengo, E.
Bullitt, and W. Lehman. 2023. Myosin loop-4 is critical for optimal tropomyosin repositioning on
actin during muscle activation and relaxation. *J Gen Physiol*. 155(2), doi:
10.1085/jgp.202213274.

52.   Doran, M. H., E. Pavadai, M. J. Rynkiewicz, J. Walklate, E. Bullitt, J. R. Moore, M. Regnier, M. A.
Geeves, and W. Lehman. 2020. Cryo-EM and Molecular Docking Shows Myosin Loop 4 Contacts
Actin and Tropomyosin on Thin Filaments. *Biophys J*. 119(4):821-830, doi:
10.1016/j.bpj.2020.07.006.

53.   Porter, J. R., A. Meller, M. I. Zimmerman, M. J. Greenberg, and G. R. Bowman. 2020.
Conformational distributions of isolated myosin motor domains encode their mechanochemical
properties. *Elife*. 9, doi: 10.7554/eLife.55132.

54.   Gunning, P. W., E. C. Hardeman, P. Lappalainen, and D. P. Mulvihill. 2015. Tropomyosin - master
regulator of actin filament function in the cytoskeleton. *Journal of cell science*. 128(16):2965-
2974, doi: 10.1242/jcs.172502.

55.   Ostap, E. M. 2008. Tropomyosins as discriminators of myosin function. *Adv Exp Med Biol*.
644:273-282, doi: 10.1007/978-0-387-85766-4_20.

56.   Kee, A. J., L. Yang, C. A. Lucas, M. J. Greenberg, N. Martel, G. M. Leong, W. E. Hughes, G. J.
Cooney, D. E. James, E. M. Ostap, W. Han, P. W. Gunning, and E. C. Hardeman. 2015. An actin
filament population defined by the tropomyosin Tpm3.1 regulates glucose uptake. *Traffic*.
16(7):691-711, doi: 10.1111/tra.12282.

57.   Pelham, R. J., Jr., J. J. Lin, and Y. L. Wang. 1996. A high molecular mass non-muscle tropomyosin
isoform stimulates retrograde organelle transport. *J Cell Sci*. 109 (Pt 5):981-989, doi:
10.1242/jcs.109.5.981.

58.   Lord, M. 2011. Cytoskeletal regulation: sorting out stress fibers with tropomyosin. *Curr Biol*.
21(7):R255-257, doi: 10.1016/j.cub.2011.02.043.

59.   Tyska, M. J., D. E. Dupuis, W. H. Guilford, J. B. Patlak, G. S. Waller, K. M. Trybus, D. M. Warshaw,
and S. Lowey. 1999. Two heads of myosin are better than one for generating force and motion.
Proc Natl Acad Sci U S A. 96(8):4402-4407, doi: 10.1073/pnas.96.8.4402.

60.   Bers, D. M., C. W. Patton, and R. Nuccitelli. 2010. A practical guide to the preparation of Ca(2+)
buffers. *Methods Cell Biol*. 99:1-26, doi: 10.1016/B978-0-12-374841-6.00001-3.

61.   Greenberg, M. J., T. Lin, Y. E. Goldman, H. Shuman, and E. M. Ostap. 2012. Myosin IC generates
power over a range of loads via a new tension-sensing mechanism. *Proc Natl Acad Sci U S A*.
109(37):E2433-2440, doi: 10.1073/pnas.1207811109.

62.   Barrick, S. K., S. R. Clippinger, L. Greenberg, and M. J. Greenberg. 2019. Computational Tool to
Study Perturbations in Muscle Regulation and Its Application to Heart Disease. *Biophys J.*
116(12):2246-2252, doi: 10.1016/j.bpj.2019.05.002.
63.   Nyitrai, M., and M. A. Geeves. 2004. Adenosine diphosphate and strain sensitivity in myosin
motors. *Philos Trans R Soc Lond B Biol Sci.* 359(1452):1867-1877, doi: 10.1098/rstb.2004.1560.
64.   R Core Team. 2021. R: A Language and Environment for Statistical Computing. *R Foundation for*
Statistical Computing.