Cyclic di-AMP traps proton-coupled K⁺ transporters of the KUP family in an inward-occluded conformation

Michael F. Fuss[#,1], Jan-Philip Wieferig[#,2], Robin A. Corey[#,3], Yvonne Hellmich[1], Igor Tascón[1,5], Joana S. Sousa[2,6], Phillip J. Stansfeld[4], Janet Vonck[*,2], Inga Hänelt[*,1]

[1] Institute of Biochemistry, Goethe University Frankfurt, Frankfurt am Main, Germany
[2] Department of Structural Biology, Max Planck Institute of Biophysics, Frankfurt am Main, Germany
[3] Department of Biochemistry, University of Oxford, Oxford, UK
[4] School of Life Sciences & Department of Chemistry, University of Warwick, Coventry, CV4 7AL, UK
[5] Current address: Instituto Biofisika (UPV/EHU, CSIC), University of the Basque Country, Leioa, Spain. Ikerbasque, Basque Foundation for Science, Bilbao, Spain
[6] Current address: UCB Pharma, UCB Biopharma UK, Slough, SL1 3WE, UK
[#] These authors contributed equally and may change the order of their names
[*] Corresponding authors

Abstract

Cyclic di-AMP is the only known essential second messenger in bacteria and archaea, regulating different proteins indispensable for numerous physiological processes. In particular, it controls various potassium and osmolyte transporters involved in osmoregulation. In *Bacillus subtilis*, the K⁺/H⁺ symporter KimA of the KUP family is inactivated by c-di-AMP. KimA sustains survival at potassium limitation at low external pH by mediating K⁺ ions uptake. However, at elevated intracellular K⁺ concentrations, further K⁺ accumulation would be toxic. In this study, we reveal the molecular basis of how c-di-AMP binding inhibits KimA. We report cryo-EM structures of KimA with bound c-di-AMP in detergent solution and reconstituted in amphipols. By combining structural data with functional assays and molecular dynamics simulations we reveal how c-di-AMP modulates transport. We show that an intracellular loop in the transmembrane domain interacts with c-di-AMP bound to the adjacent cytosolic domain. This reduces the mobility of transmembrane helices at the cytosolic side of the K⁺ binding site and therefore traps KimA in an inward-occluded conformation.

Introduction

In bacteria and archaea, potassium ions are essential, playing roles in osmoregulation[1,2], pH homeostasis[3–5], regulation of protein synthesis[6], enzyme activation[7,8] such as the ribosome[9,10], membrane potential adjustment[11] and electrical signaling[12–14]. K[+] homeostasis must be strictly maintained as deviations and fluctuations of potassium levels have lethal consequences for the cell[15,16]. In Gram-positive bacteria like *Bacillus subtilis* the second messenger cyclic di-AMP (c-di-AMP) is essential for the regulation of channels and transporters that maintain potassium homeostasis[17]. At elevated intracellular K[+] concentrations, c-di-AMP is present at increased concentrations and regulates potassium-transporting proteins in two ways: Firstly, it directly binds its target proteins, leading to the inhibition of potassium uptake systems like K[+] channel KtrAB[18,19] and K[+]/H[+] symporter KimA[17,20] or the activation of potassium exporters like CpaA[21] and KhtTU[22]; secondly, c-di-AMP suppresses gene expression by binding to the *ydaO* riboswitch[23] that among others controls the transcription of the *kimA* and *ktrAB* genes. The ability of c-di-AMP to modulate both the expression and the activity of the same proteins makes it a key player in potassium homeostasis[24]. However, while the control of gene expression is well understood, it remains unclear how c-di-AMP binding controls the activity of potassium transporters and channels.

The c-di-AMP-sensitive, high-affinity K[+]/H[+] symporter KimA from *B. subtilis* is particularly required for the uptake of potassium at low external potassium concentrations in acidic environments. KimA exploits the inward-directed proton gradient to accumulate K[+] against its concentration gradient. As a member of the amino acid-polyamine-organocation (APC) superfamily it has the classical LeuT-fold with the first ten transmembrane helices (TMHs) adopting a 5+5 inverted repeat. TMHs 11 and 12 connect the transmembrane domain (TMD) to the cytosolic domain (CD) of KimA, which consists of four alpha helices and a five-stranded beta sheet[25]. KimA is a homodimer, stabilised by the swapping of the cytosolic domains with respect to the transmembrane domain. A long loop connects the swapped cytosolic domain to the last helix of the TMD. The dimeric cytosolic domains adopt a fold similar to a phosphopantetheine adenylyltransferase (PPAT) domain and have been suggested to bind c-di-AMP[25,26]. However, no structural information is available for the c-di-AMP

binding site and, consequently, the inhibition mechanism of potassium uptake by c-

di-AMP remains elusive. In this study, we report new structural and functional insights

into the binding of c-di-AMP to KimA and how it inhibits the transporter.

Results

C-di-AMP binds to KimA in a cooperative manner and inhibits potassium uptake

The uptake of potassium ions through KimA is known to be impaired by c-di-AMP[16,19].

Here, we confirmed the hypothesis and showed that binding of c-di-AMP leads to a

significant decrease of potassium uptake velocity under *in vivo* conditions. We co-

produced KimA with the inactive adenylate cyclase variant $CdaA_{D171N}$[27] in *Escherichia*

coli LB2003 cells, a strain that lacks all endogenous potassium uptake systems[28], and

determined potassium uptake into the potassium-depleted cells, obtaining a V_{max} of

269.3 ± 24.0 nmol*mg dw^{-1}*min^{-1} and a K_m of 0.49 ± 0.09 mM. Upon the co-

expression of KimA with active, c-di-AMP-synthesising CdaA, the V_{max} was reduced by

~64% to 96.7 ± 10.1 nmol*mg dw^{-1}*min^{-1}, while the K_m did not change

$(0.66 \pm 0.31$ mM) (Figure 1a, Table 1).

81
Figure 1: The effect of c-di-AMP binding to KimA *in vivo* and *in vitro* a) Whole-cell K$^+$ uptake assay in
E. coli LB2003 cells producing KimA with active (CdaA) or inactive ($CdaA_{D171N}$) diadenylate cyclase.
Michaelis-Menten plot shown in graph in graph. Representative experiments shown (n= 3). **b)** Melting
temperatures of KimA purified from cells with CdaA present (KimA WT + CdaA) or absent (KimA WT)
and incubated with an increasing c-di-AMP concentration given in x-fold molar excess over KimA.
Determined with Differential Scanning Fluorometry (DSF). Dashed line indicates T_m of KimA WT w/o c-
di-AMP addition. Data points represent the average and error bars the standard deviation of
measurements from at least biological triplicate.

Table 1: Kinetic parameters of KimA variants in the presence or absence of c-di-AMP (i.e., active (CdaA) or inactive (CdaA$_{D171N}$) diadenylate cyclase). Determined by fitting the Michaelis-Menten plot with Michaelis-Menten equation. Mean and standard deviation from three independent whole-cell potassium uptake assays are shown.

Variant	V_{max} [nmol/(mg dw*min)]	K_m [mM]
KimA$_{WT}$ + CdaA	96.7 ± 10.1	0.66 ± 0.31
KimA$_{WT}$ + CdaA$_{D171N}$	269.3 ± 24.0	0.49 ± 0.09
KimA$_{R337A}$ + CdaA	111.7 ± 25.5	0.54 ± 0.41
KimA$_{R337A}$ + CdaA$_{D171N}$	198.9 ± 35.8	0.39 ± 0.08
KimA$_{Y118A}$ + CdaA	210.6 ± 29.0	18.85 ± 5.82
KimA$_{Y118A}$ + CdaA$_{D171N}$	202.8 ± 19.4	4.8 ± 0.75
KimA$_{N237A}$ + CdaA	207.9 ± 23.6	7.80 ± 2.23
KimA$_{N237A}$ + CdaA$_{D171N}$	161.9 ± 41.8	7.93 ± 4.21
KimA$_{A481WS582W}$ + CdaA	194.2 ± 49.2	0.35 ± 0.15
KimA$_{A481WS582W}$ + CdaA$_{D171N}$	182.0 ± 44.1	0.33 ± 0.08

In our previous structural study[25], when c-di-AMP was added to KimA purified in styrene maleic-acid lipid particles (SMALPs) before EM grid preparation, binding of c-di-AMP to KimA was not observed. In agreement with this we show here that the melting temperature of detergent-purified KimA did not significantly change upon the titration of increasing concentrations of c-di-AMP in differential scanning fluorometry (DSF) measurements (Figure 1b), suggesting that c-di-AMP did not bind in these conditions. To overcome this limitation, we co-produced KimA with the diadenylate cyclase CdaA in *E. coli* LB2003 cells, as this combination showed KimA inhibition *in vivo*, and then purified KimA from this condition. Interestingly, this sample already had a slightly increased melting temperature from 48.1 ± 0.8 °C to 50.7 ± 1.9 °C, prior to the addition of c-di-AMP. The addition of a three-fold molar excess of c-di-AMP led to a further 6.6 °C increase of the melting temperature to 54.7 ± 0.7 °C (Figure 1b). Hence, during purification, a fraction of c-di-AMP appeared to remain bound to KimA, which enabled the binding of further c-di-AMP under *ex vivo* conditions. This suggests a cooperative behaviour of c-di-AMP binding to KimA at least under *in vitro* conditions.

Structural characterisation of the binding of c-di-AMP to KimA

To determine the binding site of c-di-AMP and the effect of c-di-AMP binding on the overall structure of KimA, cryo-EM specimens of purified KimA that had been co-expressed with *cdaA* were prepared in both DDM and amphipols. KimA in DDM was additionally incubated with a ten-fold molar excess of c-di-AMP after purification.

332k and 296k particles were processed to reconstruct cryo-EM maps at 3.3 Å and 3.8

Å resolution, respectively. The overall structure of KimA obtained from the

preparation in DDM resembles the previous structure obtained in SMALPs[25] (RMSDs

of TMD (residues 30-462): 1.31 Å, CD (residues 462-606): 1.65 Å, global RMSD: 1.74

Å). The two TMDs are tilted towards each other, forming a dimer interface at the

extracellular side that would enforce a bending of the membrane by ~130° against its

natural curvature. As speculated for the preparation of KimA in SMALPs, the tilting

likely was caused by a loss of lipids between the TMDs during purification. In contrast,

the map of KimA in amphipols shows the upright dimer architecture of the TMDs

(Supplementary Figure 1a). At the extracellular side an elongated non-protein density

is visible between both protomers, indicating that amphipol molecules wrapped

around both TMDs individually. The structure resembles the upright dimer that was

previously obtained following molecular dynamics (MD) simulations of KimA in a lipid

bilayer[25], with an RMSD of only 3.4 Å (Supplementary Figure 1b). The pivot point for

the tilting of TMDs is located around the C-terminal end of TMH 12, which extends out

of the membrane (residues 459-464). Otherwise, the architectures of the TMDs

(RMSD 0.55 Å for residues 30-462) and of the cytosolic domains (RMSD 0.66 Å for

residues 462-606) of KimA reconstituted in amphipols and in DDM are very similar

(global RMSD 1.26 Å). Also, the dimer interfaces that are mainly formed by the long

connecting loops (residues 462-474) and by a short beta-sheet between the two

cytosolic domains are not affected (Figure 2a). A comparison of the resulting new

structures of KimA in DDM and in amphipols with the previously reported structure

from SMALPs[25] suggests that all of them represent a similar inward-occluded

conformation. In agreement with this, a strong density for the bound substrate

potassium ion is observed in the 3.3 Å cryo-EM map (Figure 2b). The potassium ion is

coordinated by Tyr43 (2.9 Å), the carbonyl and carboxyl of Asp36 (2.8 Å and 3.3 Å,

respectively), hydroxyl and carbonyl of Thr230 (2.9 and 3.2 Å, respectively) and Tyr377

(3.2 Å). The map from KimA obtained in SMALPs suggested that Ser125 is also part of

the K$^+$ binding site, but in MD simulations this was not the case[25]. The higher resolution

map shows that the distance between Ser125 and the potassium ion (4.9 Å) is too long

for coordination (Figure 2b), confirming the MD simulations. In contrast to the

previous map, no extra densities are localized below Asp36 and Tyr377. Asp36 and

Tyr377 are suggested to function as an intracellular gate. Potassium ions below them

were hypothesized to hinder the opening of the gate by a trans-inhibition

mechanism[25]. An explanation for the lack of the inhibitory ions could be the locking of

KimA in the inward-occluded conformation by other means, namely the binding of c-

di-AMP, which could prohibit K$^+$ binding to the trans-inhibitory site from the cytosolic

side. In fact, both cryo-EM maps of KimA in DDM and in amphipols clearly show two

non-protein densities in the CDs, which were modelled as c-di-AMP. Each binding site

is formed mostly by the cytosolic domain of one monomer (Figure 2c): One of the

phosphodiesters of c-di-AMP is coordinated through two hydrogen bonds and a salt

bridge between the non-esterified oxygen atoms and the amide groups of Gln581 and

Ser582 and the guanidinium group of Arg337. One adenosine moiety forms hydrogen

bonds with the backbone carbonyls of Val506 and Pro479. The hydroxyl group of the

ribose of the second adenosine moiety forms hydrogen bonds with His579 and the

backbone amide of Asn580. Only the second adenosine moiety is coordinated by

residues of the neighbouring protomer; it is stacked between Val457 and Leu461 of

TMH 12 that extends out of the membrane and Arg337, located in the loop between

TMHs 8 and 9. Arg337 is the only residue of the TMD that engages in a strong

interaction with c-di-AMP, making it a prime candidate for the transmission of the

inhibitory effect of bound c-di-AMP.

167
Figure 2: Cryo-EM structure of KimA with c-di-AMP bound. a) KimA homodimer with c-di-AMP bound
between the swapped CDs and TMDs of the grey and rainbow-coloured monomers. **b)** K$^+$ binding site
of one TMD. The substrate potassium ion is coordinated by Tyr43, Asp36, Thr230 and Tyr377 with
distances ranging between 2.8 and 3.3 Å. **c)** The c-di-AMP binding pocket is mostly formed by residues
of the cytosolic domain. Only Arg337 of the TMD strongly interacts with c-di-AMP. KimA was purified
after co-production with CdaA and solubilized with DDM. A ten-fold molar excess of c-di-AMP was
added after purification.

Figure 3: Cooperativity network between cytosolic domains of KimA dimer. a) Top view of dimeric CDs highlighting bound c-di-AMP and residues linking both binding sites. Protomer affiliation highlighted in grey. **b)** RMSF (Å) of residues connecting binding pockets in the absence, presence or with asymmetric binding of c-di-AMP. Priming of unoccupied binding site by ligand binding to the opposing protomer is highlighted in grey. Data in top row are from 3 x 2.2 μs simulations, as for the middle row, and data from the bottom row are from 3 x 500 ns simulations. **c)** The presence of c-di-AMP has a considerable effect on the range of X_2 angles sampled by the Trp592 sidechain.

C-di-AMP binding to one binding site primes the other

The structural insights were used to further investigate the binding of c-di-AMP to KimA. In particular, we strove to understand how pre-binding of c-di-AMP facilitates further c-di-AMP binding, as suggested by DSF measurements. To this end, MD simulations were run of the KimA dimer either with or without c-di-AMP, with c-di-AMP remaining stably bound to KimA throughout the simulations where present, with an RMSD relative to the protein of 0.25 ± 0.07 nm over 3 repeats of ca. 2.2 μs. Analysis of the data led to the identification of a cooperativity pathway between the two binding sites (Figure 3a): Ser582, which is in direct contact with c-di-AMP, connects through Phe565 and Arg587 with Trp592. Trp592 links both CDs through a contact with Ile569 across the dimer interface. Ile569 is a direct neighbour of Phe568 which, like Phe565, delimits the c-di-AMP binding pocket. The root mean square fluctuation (RMSF) of all these residues was significantly reduced compared to the apo-protein when c-di-AMP was present in both binding sites, suggesting a stabilising effect of c-di-AMP binding on the CD (Figure 3b, Supplementary Videos 1 and 2). Surprisingly, if

only one binding site was occupied by c-di-AMP, a similar stabilising effect was

observed and all connecting residues of the apo-protomer also showed lower RMSF

as if c-di-AMP was bound (Figure 3b, grey highlighted, Supplementary Video 3). It is

possible that the more stable sidechain of Trp592 (Figure 3c) restricts the motion of

its counterpart Ile569 in the other protomer, which via the described network

facilitates the priming of the second, empty c-di-AMP binding site.

Arg337 increases the affinity for c-di-AMP but is not essential for the inhibition

As Arg337 was the only residue of the TMD identified to strongly interact with c-di-

AMP in the binding pocket, it was suspected to transmit the inhibitory effect of c-di-

AMP binding from the cytosolic domain to the TMD. DSF measurements showed that

the mutated variant KimA$_{R337A}$, which should be unable to link c-di-AMP binding to the

TMD, was still able to bind c-di-AMP when co-expressed with *cdaA;* however, a 20-

fold excess was required for reaching the same thermal stabilisation as with a three-

fold excess for the wildtype (Figure 4a), suggesting a decreased affinity for c-di-AMP.

214
Figure 4: Arg337 is essential for high-affinity binding of c-di-AMP. a) Melting temperatures of KimA
R337A purified from cells with CdaA absent (KimA$_{R337A}$) or present (KimA$_{R337A}$ + CdaA), and incubated
with an increasing c-di-AMP concentration given in x-fold molar excess over KimA. Determined with
Differential Scanning Fluorometry (DSF). Dashed line indicates T$_m$ of KimA WT without c-di-AMP
addition. Data points represent the average and error bars the standard deviation of measurements
from at least biological triplicate. **b)** Whole-cell K$^+$ uptake assay in *E. coli* LB2003 cells producing
KimA$_{R337A}$ with active (CdaA) or inactive (CdaA$_{D171N}$) diadenylate cyclase. Michaelis-Menten plot shown
in graph. Representative experiment shown (n= 3).

To determine how the mutation affects the inhibition by c-di-AMP, *in vivo* potassium
uptake studies were performed. In the presence of active CdaA, the potassium uptake
velocity remained rather high (Figure 4b), but the inhibition by c-di-AMP was not
completely abolished. The V_{max} value for K$^+$ uptake by KimA$_{R337A}$ in the presence of
active CdaA was reduced by ~44% to 111.7 ± 25.5 nmol*mg dw^{-1}*min^{-1} when
compared to the co-expression of KimA$_{R337A}$ with inactive CdaA, where a V_{max} value
for K$^+$ of 198.9 ± 35.8 nmol*mg dw^{-1}*min^{-1} was determined (Table 1). To address
whether the remaining inhibition was related to c-di-AMP binding to KimA or caused
by the production of c-di-AMP in general, variant KimA$_{A481WS582W}$ that should be unable
to bind c-di-AMP because of a sterically blocked binding pocket, was analysed. The
loss of c-di-AMP binding was confirmed by DSF measurements (Supplementary Figure
2a). This loss of binding was accompanied by the insensitivity of the *in vivo* potassium
uptake through KimA$_{A481WS582W}$ to inhibition by c-di-AMP (Supplementary Figure 2b).
The uptake velocities were comparable in the presence of an active and inactive
cyclase, respectively (Table 1). In conclusion, Arg337 appears to be important for high-
affinity c-di-AMP binding but is not necessarily required for communicating c-di-AMP
binding to the TMD for inhibition.

C-di-AMP-induced inhibition of KimA

C-di-AMP does not directly interact with the TMD, and long-range communication is
required because the c-di-AMP binding site is ~37 Å away from the substrate K$^+$
binding site. The question remains how c-di-AMP binding to the CDs of KimA controls
K$^+$ transport at a distance. To address this question, we further evaluated the MD
simulations performed in the presence and absence of c-di-AMP. They showed a
significant stabilisation of the whole intracellular loop between TMH8 and TMH9 upon
c-di-AMP binding, with the largest decrease of the RMSF for Arg337 from 2.3 ± 0.2 Å
to 1.7 ± 0.2 Å. However, KimA with an Arg337 to alanine mutation showed a similar
reduction of flexibility of the loop in the presence of c-di-AMP (Figure 5a). This
suggests that inhibition by c-di-AMP is transmitted to the TMD via this loop.
The MD simulations further revealed that one consequence of c-di-AMP being
removed from the system was a movement of TMH6 away from TMH3, leading to an
opening of the TMD at the cytosolic end. This effect can be seen by measuring the

distance between two residues sitting underneath the intracellular gate, Tyr118
(TMH3) and Asn237 (TMH6). These typically interacted closely (<4 Å) when c-di-AMP
is bound (Figure 5b, Supplementary Figure 3a), which likely contributes to locking the
TMD in an inward-occluded conformation. When c-di-AMP was removed this
interaction was broken (Figure 5b), and the distance frequently shifted from about 4
Å to 8 Å (Supplementary Figure 3b). Distancing of Tyr118 and Asn237 can also be seen
using principal component analysis (PCA) of the MD data (see Methods), which
revealed a rearrangement of the central TMHs in the non-liganded state, with TMH6
moving ca. 4 Å away from TMH3 and TMH8 (Figure 5c, Supplementary Figure 4b).
Rearrangement of TMH6 was not observed in the liganded state (Supplementary
Figure 4a + c). The MD data reveal that, whilst the initial bound K^+ remain tightly bound
in the c-di-AMP-bound state, they are free to rapidly exchange with the bulk solvent
in the c-di-AMP-free state, suggesting a switch to an inward-open conformation. In
agreement with the MD simulations, a mutation of Tyr118 or Asn237 to alanine led to
a loss of inhibition by c-di-AMP in potassium uptake assays, while binding of c-di-AMP
was still possible (Figure 5d, Table 1, Supplementary Figure 5). The mutations seem to
have abolished the observed interaction from the MD simulations, and therefore
reduce the ability of c-di-AMP to lock KimA in an inward-occluded state. The
significantly increased K_m of 19 mM and 8 mM for $KimA_{Y118A}$ and $KimA_{N237A}$,
respectively, is in agreement with the assumption that these residues line the exit
pathway to the cytosol and suggests their involvement in ion release.
In conclusion, the MD simulations together with the functional data suggest that the
binding of c-di-AMP (Figure 5e-1) communicates to the TMD via a stabilisation of the
TMH8-TMH9 loop (Figure 5e-2, blue), which in turn affects the positioning of TMH8
and TMH3 relative to TMH6 (Figure 5e-3 pink, green, yellow). These conformations
would appear to regulate the switching between inward-occluded and inward-open,
thereby presenting a possible mechanism of c-di-AMP inhibition of KimA.

282
Figure 5: Transduction of inhibition from CD to TMD upon c-di-AMP binding. a) RMSF of binding loop
between TMH 8 and 9 determined by MD simulations. **b)** MD simulations reveal an increased likelihood
of interaction (distance ≤ 4 Å) between Tyr118 and Asn237 upon c-di-AMP binding compared to when
no ligand is bound or when Arg337 is changed to alanine. Interactions were analyzed over 3 x 2.2 μs
simulations for the system with and without c-di-AMP. **c)** Views of the KimA TMD in the eigenvector 1
extreme states from simulation data with no ligand bound, showing a move of TMH6 (yellow) away
from TMH3 (green) and TMH8 (pink). The loop connecting TMH8 and 9 is shown in blue. The same data
for the ligand-bound state can be found in Supplementary Figure 4 along with structures for
eigenvector 2 and the relative view from the input model. **d)** Reduced inhibition of potassium uptake
in whole-cell K^+ uptake assays when Tyr118 or Asn237 were mutated to alanine. K^+ uptake for $KimA_{Y118A}$
or $KimA_{N237A}$ variant in the presence of active CdaA or inactive $CdaA_{D171N}$. Added K^+: light blue: 0.1 mM,
blue: 0.5 mM, light green: 1 mM, green: 2 mM, yellow: 5 mM, orange: 7.5 mM, red: 10 mM. Michaelis-
Menten diagram shown in graph. Representative experiment shown (n= 3). **e)** Model for signal
transduction: Upon c-di-AMP binding to CD (1), the binding loop (blue) rigidifies (2), TMH8 and 3
become restricted in their mobility, leading to more constant interaction between the cytosolic ends
of TMH3 and 6, e.g. Tyr118 and Asn237 (3). The opening of the intracellular gate is abolished.

Discussion

Our results provide first structural and functional insights into how c-di-AMP inhibits proteins of the KUP transporter family. Interestingly, the binding mode differs from other c-di-AMP-regulated transport proteins such as potassium channel KtrAB[19], osmolyte transporters OpuC[29] and OpuA[30], and potassium exporter KhtTU[22], where c-di-AMP binds at dimer interfaces in a symmetric binding pocket. In KimA, two c-di-AMP molecules bind to the dimer in an asymmetric binding pocket. Different to the binding sites in the RCK_C domains of KtrA and KhtT and in the CBS domains of OpuA and OpuC, in which two arginine residues, one per protomer, coordinate both phosphate esters, only one of the two phosphate esters of c-di-AMP is coordinated by an arginine (Arg337) in KimA. In fact, not even Arg337 is essential for the binding of c-di-AMP or its inhibitory function, although it appears to increase the affinity. Instead, the major linker to the TMD is the entire binding loop between TMH8 and 9, which becomes more restricted in its movement once c-di-AMP is bound. This stabilisation may limit the freedom of movement of TMH8, which then restrains TMH3 and TMH6, for instance through interactions between Leu325, Tyr118 and Asn237. MD simulations indicate that through these interactions, opening of the inner gate is hindered, arresting KimA in an inward-occluded conformation. Consequently, the rocker switch movement can no longer take place (Figure 6). When c-di-AMP is removed, KimA is once again able to open up to exchange K$^+$ with the bulk solvent, as seen in our simulations.

Figure 6: Schematic model of c-di-AMP-induced inhibition of KimA. a) KimA undergoes a rocker switch movement to fulfil proton-coupled potassium uptake. The binding loop is unrestricted in its movement. Tyr118 and Asn237 are only in loose contact. **b)** C-di-AMP binding rigidifies the binding loop. Helix 8 leans on helix 3, stabilizing the interaction between helix 3 and 6 bringing Tyr 118 and Asn237 closer. KimA is then locked in an inward-occluded conformation, preventing the opening of the inner gate.

To elucidate how conserved the inhibition mechanism is among other KUPs we
performed a structure-based sequence alignment with AlphaFold predictions of four
other KUPs of gram-positive bacteria as well as Kup from the gram-negative *E. coli*,
which have been functionally but not structurally characterized (Supplementary
Figure 6). While Kup from *E. coli* is not regulated by c-di-AMP because the second
messenger is lacking in *E. coli*, KimA from *Listeria monocytogenes* (KimA[Lmo])[26], and
KupA and KupB from *Lactococcus lactis* IL1403[31] showed growth inhibition by c-di-
AMP. Further, c-di-AMP binding was shown for KupA and with lower efficiency for
KupB. The regulation of KimA from *Staphylococcus aureus* (KimA[Sau]) by c-di-AMP
remains unclear[26]. In agreement with the lack of c-di-AMP in *E. coli*, Kup from *E. coli*
does not share any of the characteristic residues Arg337, Tyr118 or Asn237 with KimA
from *B. subtilis*. In contrast, Arg337 is conserved in KimA[Lmo], KimA[Sau] and KupA. KupB
lacks an equivalent amino acid, which could explain the weaker c-di-AMP binding.
KupB is still inhibited by c-di-AMP, supporting our assumption that Arg337 is not
required for transmission of the inhibition to the TMD. Residues Tyr118 and Asn237
are only conserved in KimA[Sau], while KupA, KupB and KimA[Lmo] share a glycine and a
serine, respectively, at those positions. In the binding pocket, KimA[Sau] shows
conservation for Pro479, His579, Asn580, Gln581 and similarity for Val457 and Ser582
(Ile/Thr in KimA[Sau], respectively). The KUPs show conservation only for Gln581 and like
KimA[Sau] have a Thr at the position of Ser582, with the exception of Kup from *E. coli*,
which has a lysine and leucine. Therefore, inhibition of KimA[Sau] by c-di-AMP is likely to
be similar to KimA from *B. subtilis*, while for the other homologs some variations are
to be expected.
349
A surprising observation was that c-di-AMP only bound to KimA *in vitro* when at least
a proportion of c-di-AMP was co-purified along with KimA. Structurally, there is no
obvious reason for this observation. The MD simulations showed that c-di-AMP stably
binds to both the upright- and the tilted-dimer arrangement. The superposition of the
previously solved structure in the absence of c-di-AMP with the c-di-AMP-bound
structures does not show any significant conformational changes in the binding
pocket, apart from a general stabilisation when c-di-AMP is bound. However, binding
of the first c-di-AMP molecule by KimA appears to be unfavourable. A direct hand over

of c-di-AMP from the cyclase could lower the energetic barrier for ligand binding.

Binding of a second molecule then appears to be strongly cooperative within dimeric

KimA. In agreement with this hypothesis, the only membrane-bound diadenylate

cyclase CdaA in *B. subtilis* is described as the major cyclase maintaining the internal c-

di-AMP concentration required for cell growth[32]. In other bacteria like the pathogenic

L. monocytogenes[33,34] and *S. aureus*[35] as well as *L. lactis*[36] CdaA is the only c-di-AMP

cyclase. Since the majority of proteins regulated by c-di-AMP are membrane bound,

the co-localisation of the cyclase appears advantageous if direct interaction is

necessary.

In summary, we revealed key elements for the inhibition of potassium uptake through

KimA by c-di-AMP. We show the first structure of a KUP transporter with bound

secondary messenger c-di-AMP and confirmed its binding *in vivo* and *in vitro*. Based

on our findings we propose a network for the inhibition of KimA by c-di-AMP at a

distance. Structures of outward-open and inward-open KimA are necessary to confirm

the proposed transport cycle and validate the mode of inhibition by c-di-AMP.

Material and Methods

Cloning of KimA variants using site-directed mutagenesis

Point mutations were introduced into pB24KimA using site-directed mutagenesis

(SDM). Primer pairs of 20-40 bp length including a 1-3 bp mismatch were used to

introduce point mutations (Supplementary Table 1). PCR product was digested with

DpnI to remove template DNA and subsequently used to transform *E. coli* DH5α.

Colonies grown on LB agar plates with 100 µg/ml ampicillin were picked and cultured

in 5 ml LB with 100µg/ml ampicillin over day. Plasmids were isolated from cells using

NucleoSpin Plasmid, Mini kit for plasmid DNA (Macherey-Nagel). Correct mutation of

the plasmids was verified by sequencing performed by MicroSynth GmbH Göttingen.

Expression and protein purification of KimA

A colony of *E. coli* LB2003 cells transformed with the expression plasmids for KimA or

variants thereof (pB24KimA) and, if indicated, CdaA variants (pB33CdaA) where grown

at 37°C o/d with 180 rpm shaking in 5 ml KML (1% KCl, 1% tryptone, 0.5% yeast extract

(w/v)) with 100 µg/ml ampicillin and 30 µg/ml chloramphenicol if needed. An

overnight culture of 200 ml KML with respective antibiotics was inoculated from the

o/d culture and incubated at 37°C with 180 rpm. 6 l KML were inoculated with the o/n

culture to an OD_{600} of 0.1 and incubated at 37°C with 180 rpm. Gene expression was

induced with 0.002% arabinose at an OD_{600} of 1. Cells were grown for 1.5 h and then

harvested by centrifugation. Cells were resuspended in buffer containing 420 mM

NaCl, 180 mM KCl, 50 mM Tris-HCl pH 8 supplemented with 1 mM EDTA, 0.1 mM

PMSF, 0.3 mM benzamidine and DNase I. Cells were disrupted by passing through a

cell homogeniser (Stansted Pressure Cell Homogeniser FPG 12800) at 1 kbar. Cell

debris was removed by centrifuging at 15,000 x *g* for 15 min. Membranes were

harvested o/n by centrifuging at 100,000 x *g*. Membranes were resuspended in

aforementioned buffer (100 mg/ml) and homogenised. Membranes were solubilised

for 1 h at 4°C with 1% of DDM (Glycon). Unsolubilised proteins were removed by

centrifuging 30 min at 135,000 x *g*. Supernatant was incubated with 2 ml Ni^{2+}-NTA

resin for 1 h at 4°C. Ni-NTA was washed with 50 column volumes of buffer containing

140 mM NaCl, 60 mM KCl, 20 mM Tris-HCl pH 8 and 0.04% DDM supplemented with

50 mM imidazole. Protein was eluted by using 500 mM imidazole in aforementioned

buffer. Protein was further purified via size exclusion chromatography using a

Superose6 Increase column (GE Healthcare/Cytiva) preequilibrated with

aforementioned buffer without imidazole addition. Protein was concentrated and

incubated with c-di-AMP if needed for DSF, HPLC-MS and cryo-EM experiments.

Preparation of KimA in amphipols

Protein expression and purification was performed as described above until binding

to Ni^{2+}-NTA. The beads were washed with 50 column volumes of buffer (140 mM NaCl,

60 mM KCl, 20 mM Tris-HCl, pH 8) containing 60 mM imidazole and a reduced DDM

concentration of 0.025%. KimA was eluted by making use of the C-terminal HRV-3C-

Protease cleavage site. Therefore, the beads were incubated with 3C-Protease for

1.5 h at 4°C. The beads were washed twice with buffer containing 0.025% DDM. The

elution fraction containing KimA was concentrated and filtered. For the

detergent/amphipol exchange, KimA at a concentration of 12 mg/ml and amphipols

PMAL C8 (10% in water; Anatrace) were incubated at w/w ratio of 1:10 for 1 h at 4°C.

To remove excess detergent, the sample was incubated with biobeads at a detergent-

to-biobeads weight ratio of 1:100 overnight. The biobeads were removed and the

sample was loaded onto a Superose6 Increase 200 10/300 GL column (GE Healthcare)

equilibrated to cryo-EM buffer (50 mM NaCl, 50 mM KCl, 20 mM Tris-HCl, pH 8). KimA-

containing fractions were pooled and concentrated to 2.25 mg/ml for cryo-EM.

In vivo *whole cell potassium uptake*

Protocol was adapted from[25,37]. A colony of *E. coli* LB2003 cells transformed with the

expression plasmids for KimA variants (pB24KimA) and CdaA variants (pB33CdaA) was

grown at 37°C o/d with 180 rpm shaking in 5 ml KML with 100 µg/ml ampicillin and

30 µg/ml chloramphenicol. 75 ml of K_{30} minimal media supplemented with the

respective antibiotics were inoculated with 3 ml of o/d KML culture and incubated o/n

at 37°C. 500 ml of K_{30} minimal media with antibiotics were inoculated to an OD_{600} of

0.15 and incubated at 37°C with 180 rpm. Gene expression was induced at an OD_{600}

of 0.4-0.6 with 0.002% arabinose. Cells were grown 1.5 h before being harvested with

6000 x *g* at 20°C for 10 min. The cell pellet was washed twice in 10 ml 120 mM Tris-

HCl pH 8. Cells were adjusted to an OD_{600} of 30 and incubated for 5 min at 37°C in a

water bath (130 rpm shaking). To permeabilize cells for internal potassium, 1 mM

EDTA was added and cells were incubated at 37°C for 7 min shaking at 130 rpm in a

water bath. Cells were centrifuged with 4500 x *g* for 7 min at 20°C and washed twice

in 200 mM HEPES TEA pH 7.5 to remove EDTA and internal potassium. Cells were

adjusted to an OD_{600} of 30. For uptake, cells were diluted to an OD_{600} of 3 with 200 mM

HEPES TEA pH 7.5 and energized with 0.2% glycerol and 0.002% arabinose. Different

potassium concentrations (0.1, 0.5, 1, 2, 5, 7.5, 10, 15 mM) were added and 1 ml

samples were taken at 0, 1, 2, 4, 7, 10 min after addition. Potassium uptake was

stopped as cells were centrifuged through 200 µl of silicone oil (ρ = 1.04) at 17000 x *g*

for 2 min. Media and oil was removed and the 1.5 ml reaction tube tip containing the

cell pellet was cut off. Cell pellet was resuspended in 1 ml of 5% TCA. Cells were lysed

by freezing at -20°C and subsequent boiling at 95°C for 10 min. Supernatant was

diluted with 3 ml of 6.7 mM CsCl and 4 ml 5 mM CsCl. Cell debris was pelleted by

centrifuging 20 min at 4000 x *g*. Internal potassium concentration was determined

using flame atomic absorption spectroscopy (F-AAS).

Differential Scanning Fluorometry/Thermal Shift Assay (DSF/TSA)

When ligands bind their target, they stabilize or destabilize the protein leading to a shifted protein melting temperature. The melting temperature is detected *via* the increasing fluorescence of the 7-diethylamino-3-(4-maleimidophenyl)-4-methylcoumarin (CPM) dye which binds cysteine residues that become accessible during the unfolding process[38,39]. 0.5 mg/ml purified KimA protein, coexpressed with and without *cdaA*, was incubated with 0-100x molar excess of c-di-AMP over night at 4°C. The next day 16.67 μg/ml CPM dye was added with subsequent incubation for 15 min on ice in the dark. 10 min centrifugation at 17,000 x *g* at 4°C removed aggregates. Melting curves were recorded in a Rotor-Gene Q 5Plex HRM system (Qiagen) using a temperature range of 25-85°C with 1°C steps each 30 s and 90 s prewarm. CPM was excited at 365 nm and emission was detected at 460 nm. Samples were measured in technical triplicates. Melting temperatures were determined at the inflection point of the resulting fluorescence curve.

Cryo-EM specimen preparation and data acquisition

UltrAuFoil R1.2/1.3 400 mesh gold grids were glow-discharged twice before the application of 3 μl of either a 4.0 mg/ml solution of KimA in DDM or 2.25 mg/ml KimA reconstituted in amphipols. The sample was then plunge-frozen using a FEI Vitrobot Mark IV. The chamber and Whatman 595 blotting paper were equilibrated at 4°C and 100% relative humidity.

A Titan Krios (Thermo Scientific) equipped with a Gatan K3 camera in counting mode and energy filter was used for imaging with the software EPU (Thermo Scientific). The fluence over a broken hole was adjusted to 1.1 electrons/Å^2 per frame. Micrographs were acquired as 50-frame movie stacks in 2.7 s or 2.3 s exposures, respectively, at a nominal magnification of 105,000x with a resulting pixel size of 0.83 Å. Defocus values were set in the range of -1.1 to -2.5 μm.

Image processing and model refinement

Micrographs were processed using Relion-3.1[40] (Supplementary Figure 7). The Relion implementation of MotionCor2[41] was used for drift correction and dose weighting.

Gctf[42] was used for the initial CTF estimation. Initially 950k coordinates were picked

from 2,349 micrographs by Topaz[43] after training the neural network with 726

particles that were manually picked from 30 micrographs of KimA in DDM. After two

rounds of 3D classification 332k particles remained that yielded a 3.6 Å map of KimA.

Per particle drift correction and dose-weighting during Bayesian polishing[44] and two

iterations of CTF refinement improved the resolution to 3.5 Å (C1) and 3.3 Å (C2

symmetry applied) (Supplementary Figure 8a).

In the case of KimA in amphipols, 5.18 million coordinates were picked from 4,303

micrographs by Topaz after training the neural network with 530 particles that were

manually selected from 20 micrographs. Bad picks were removed by two consecutive

3D classifications yielding 1.99 million and 861k particles, respectively. A lowpass

filtered map of KimA solubilized in SMA[25] was used as an initial 3D template.

Additional rounds of 3D classification resulted in a set of 296k particles from the 3,417

best micrographs yielding a 4.1 Å map of KimA. The resolution was also improved by

Bayesian polishing and two iterations of CTF-refinement to 4.0 Å (C1) and 3.8 Å (C2)

(Supplementary Figure 8b). Further improvement of the map was achieved through

focused classification of symmetry expanded particles. The particles of the last

refinement were C2 symmetry expanded. Then density outside a wide mask that

contained one TMD and the cytosolic domain of the other protomer was subtracted

from the particle image with a Relion Particle Subtraction job. 3D classification of

these particles with four classes, local searches only and the same mask used for the

particle subtraction as a reference mask yielded one class with 307k particles. 3D-

autorefinement of these particles resulted in a 3.7 Å map of one half of the dimer

(Supplementary Figure 8c).

Individual domains of the model of KimA in SMA (pdb ID 6s3k)[25] were rigid body fitted

into the map of the upright dimer with *Coot*[45] and manually rebuilt. Models were

optimised using Phenix real space refinement[46] (Supplementary Table 2).

Molecular dynamics simulations

Atomistic simulations were built using the coordinates of dimeric KimA bound to c-di-

AMP from this study. The systems were described with the CHARMM36m force

field[47,48] and built into 6:3:1 POPE, POPG, cardiolipin membranes with TIP3P waters

and K^+ and Cl^- to 150 mM, using CHARMM-GUI[49,50]. Three bound K^+ from each subunit

were preserved from the input structure. The c-di-AMP molecules were

parameterised in CHARMM-GUI, and either included in both subunits, a single subunit,

or in neither subunit. Where used, mutations were made in CHARMM-GUI. Each

system was minimized and equilibrated according to the standard CHARMM-GUI

protocol. Production simulations were run in the NPT ensemble, with temperatures

held at 303.5 K using a velocity-rescale thermostat and a coupling constant of 1 ps,

and pressure maintained at 1 bar using a semi-isotropic Parrinello-Rahman pressure

coupling with a coupling constant of 5 ps[51,52]. Short range van der Waals and

electrostatics were cut-off at 1.2 nm. Simulations were run in triplicate, to ca. 2.2 µs

for the wild-type systems with c-di-AMP present or apo, or to 500 ns for the Arg337

to alanine mutation or asymmetric c-di-AMP occupancy systems. In total, ca. 16 µs of

data were gathered.

PCA was carried out on the C-alpha atoms of each KimA monomer. The trajectories

for each subunit and each of the 3 repeats were concatenated before analysis for a

total of 6.6 µs sampling per condition. PCA was performed using the gmx covar and

gmx anaeig programs. For both ligand and no ligand conditions, ca. 1700 eigenvectors

were found, of which eigenvectors 1 and 2 contributed ca. 50% of the total variance

(see Supplementary Figure 9). Projecting these two eigenvectors reveal good overlap

between all 6 KimA monomers, with a generally broader distribution along

eigenvector 1 for the non-liganded state (Supplementary Figure 10).

All simulations were run in Gromacs 2020.1[53]. Data were analysed using Gromacs tools

and VMD[54] Plots were made using Prism 9 (GraphPad).

Data availability

Data supporting the findings of this manuscript are available from the corresponding

authors upon request. A reporting summary for this Article is available as a

Supplementary Information file. The source data of the DSF measurements and the

whole-cell transport assays are provided as a Source Data file. The cryo-EM maps with

amphipols and DDM were deposited in the wwPDB with accession codes EMD-15895

and EMD-15894, respectively, and the models with PDB-ID 8B71 and 8B70.

References

1. Holtmann, G., Bakker, E. P., Uozumi, N. & Bremer, E. KtrAB and KtrCD: two K$^+$
uptake systems in *Bacillus subtilis* and their role in adaptation to hypertonicity.
J Bacteriol **185**, 1289–98 (2003).
2. McLaggan, D., Naprstek, J., Buurman, E. T. & Epstein, W. Interdependence of K$^+$
and glutamate accumulation during osmotic adaptation of *Escherichia coli*. *J*
Biol Chem **269**, 1911–7 (1994).
3. Ochrombel, I., Ott, L., Krämer, R., Burkovski, A. & Marin, K. Impact of improved
potassium accumulation on pH homeostasis, membrane potential adjustment
and survival of *Corynebacterium glutamicum*. *Biochim Biophys Acta* **1807**, 444–
50 (2011).
4. Bakker, E. P. & Mangerich, W. E. The effects of weak acids on potassium uptake
by *Escherichia coli* K-12 inhibition by low cytoplasmic pH. *Biochim Biophys Acta*
730, 379–86 (1983).
5. Booth, I. R. & Kroll, R. G. Regulation of cytoplasmic pH (pH1) in bacteria and its
relationship to metabolism. *Biochem Soc Trans* **11**, 70–2 (1983).
6. Willis, D. B. & Ennis, H. L. Ribonucleic acid and protein synthesis in a mutant of
Bacillus subtilis defective in potassium retention. *J Bacteriol* **96**, 2035–42
(1968).
7. Gohara, D. W. & di Cera, E. Molecular Mechanisms of Enzyme Activation by
Monovalent Cations. *J Biol Chem* **291**, 20840–20848 (2016).
8. Suelter, C. H. Enzymes activated by monovalent cations. *Science* **168**, 789–95
(1970).
9. Rozov, A. *et al.* Importance of potassium ions for ribosome structure and
function revealed by long-wavelength X-ray diffraction. *Nat Commun* **10**, 2519
(2019).
10. Nissen, P., Hansen, J., Ban, N., Moore, P. B. & Steitz, T. A. The structural basis of
ribosome activity in peptide bond synthesis. *Science* **289**, 920–30 (2000).
11. Epstein, W. The roles and regulation of potassium in bacteria. *Prog Nucleic Acid*
Res Mol Biol **75**, 293–320 (2003).
12. Prindle, A. *et al.* Ion channels enable electrical communication in bacterial
communities. *Nature* **527**, 59–63 (2015).
13. Humphries, J. *et al.* Species-Independent Attraction to Biofilms through
Electrical Signaling. *Cell* **168**, 200-209.e12 (2017).
14. Liu, J. *et al.* Coupling between distant biofilms and emergence of nutrient time-
sharing. *Science* **356**, 638–642 (2017).
15. Radchenko, M. v *et al.* Potassium/proton antiport system of *Escherichia coli*. *J*
Biol Chem **281**, 19822–9 (2006).
16. Stautz, J. *et al.* Molecular Mechanisms for Bacterial Potassium Homeostasis. *J*
Mol Biol **433**, 166968 (2021).
17. Gundlach, J. *et al.* Control of potassium homeostasis is an essential function of
the second messenger cyclic di-AMP in *Bacillus subtilis*. *Sci Signal* **10**, eaal3011
(2017).
18. Corrigan, R. M. *et al.* Systematic identification of conserved bacterial c-di-AMP
receptor proteins. *Proc Natl Acad Sci U S A* **110**, 9084–9 (2013).

19.   Kim, H. *et al.* Structural Studies of Potassium Transport Protein KtrA Regulator
of Conductance of K$^+$ (RCK) C Domain in Complex with Cyclic Diadenosine
Monophosphate (c-di-AMP). *J Biol Chem* **290**, 16393–402 (2015).
20.   Gundlach, J. *et al.* Sustained sensing in potassium homeostasis: Cyclic di-AMP
controls potassium uptake by KimA at the levels of expression and activity. *J*
Biol Chem **294**, 9605–9614 (2019).
21.   Chin, K.-H. *et al.* Structural Insights into the Distinct Binding Mode of Cyclic Di-
AMP with SaCpaA_RCK. *Biochemistry* **54**, 4936–51 (2015).
22.   Cereija, T. B., Guerra, J. P. L., Jorge, J. M. P. & Morais-Cabral, J. H. c-di-AMP, a
likely master regulator of bacterial K$^+$ homeostasis machinery, activates a K$^+$
exporter. *Proc Natl Acad Sci U S A* **118**, (2021).
23.   Nelson, J. W. *et al.* Riboswitches in eubacteria sense the second messenger c-
di-AMP. *Nat Chem Biol* **9**, 834–9 (2013).
24.   Stülke, J. & Krüger, L. Cyclic di-AMP Signaling in Bacteria. *Annu Rev Microbiol*
74, 159–179 (2020).
25.   Tascón, I. *et al.* Structural basis of proton-coupled potassium transport in the
KUP family. *Nat Commun* **11**, 626 (2020).
26.   Gibhardt, J. *et al.* c-di-AMP assists osmoadaptation by regulating the *Listeria*
monocytogenes potassium transporters KimA and KtrCD. *J Biol Chem* **294**,
16020–16033 (2019).
27.   Rosenberg, J. *et al.* Structural and biochemical analysis of the essential
diadenylate cyclase CdaA from *Listeria monocytogenes*. *J Biol Chem* **290**, 6596–
606 (2015).
28.   Stumpe, S. & Bakker, E. P. Requirement of a large K$^+$-uptake capacity and of
extracytoplasmic protease activity for protamine resistance of *Escherichia coli*.
Arch Microbiol **167**, 126–36 (1997).
29.   Huynh, T. N. *et al.* Cyclic di-AMP targets the cystathionine beta-synthase
domain of the osmolyte transporter OpuC. *Mol Microbiol* **102**, 233–243 (2016).
30.   Sikkema, H. R. *et al.* Gating by ionic strength and safety check by cyclic-di-AMP
in the ABC transporter OpuA. *Sci Adv* **6**, (2020).
31.   Quintana, I. M. *et al.* The KupA and KupB Proteins of *Lactococcus lactis* IL1403
Are Novel c-di-AMP Receptor Proteins Responsible for Potassium Uptake. *J*
Bacteriol **201**, (2019).
32.   Mehne, F. M. P. *et al.* Cyclic di-AMP homeostasis in *Bacillus subtilis*: both lack
and high level accumulation of the nucleotide are detrimental for cell growth. *J*
Biol Chem **288**, 2004–17 (2013).
33.   Woodward, J. J., Iavarone, A. T. & Portnoy, D. A. c-di-AMP secreted by
intracellular *Listeria monocytogenes* activates a host type I interferon response.
Science **328**, 1703–5 (2010).
34.   Rismondo, J. *et al.* Phenotypes Associated with the Essential Diadenylate
Cyclase CdaA and Its Potential Regulator CdaR in the Human Pathogen *Listeria*
monocytogenes. *J Bacteriol* **198**, 416–26 (2016).
35.   Chaudhuri, R. R. *et al.* Comprehensive identification of essential *Staphylococcus*
aureus genes using Transposon-Mediated Differential Hybridisation (TMDH).
BMC Genomics **10**, 291 (2009).

36.   Zhu, Y. *et al.* Cyclic-di-AMP synthesis by the diadenylate cyclase CdaA is
641   modulated by the peptidoglycan biosynthesis enzyme GlmM in *Lactococcus*
642   *lactis*. *Mol Microbiol* **99**, 1015–27 (2016).
643   37.   Bakker, E. P. & Mangerich, W. E. Interconversion of components of the bacterial
644   proton motive force by electrogenic potassium transport. *J Bacteriol* **147**, 820–
645   6 (1981).
646   38.   Alexandrov, A. I., Mileni, M., Chien, E. Y. T., Hanson, M. A. & Stevens, R. C.
647   Microscale Fluorescent Thermal Stability Assay for Membrane Proteins.
648   *Structure* **16**, 351–359 (2008).
649   39.   Semisotnov, G. v. *et al.* Study of the 'molten globule' intermediate state in
650   protein folding by a hydrophobic fluorescent probe. *Biopolymers* **31**, 119–128
651   (1991).
652   40.   Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure
653   determination in RELION-3. *Elife* **7**, (2018).
654   41.   Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion
655   for improved cryo-electron microscopy. *Nat Methods* **14**, 331–332 (2017).
656   42.   Zhang, K. Gctf: Real-time CTF determination and correction. *J Struct Biol* **193**,
657   1–12 (2016).
658   43.   Bepler, T. *et al.* Positive-unlabeled convolutional neural networks for particle
659   picking in cryo-electron micrographs. *Nat Methods* **16**, 1153–1160 (2019).
660   44.   Zivanov, J., Nakane, T. & Scheres, S. H. W. A Bayesian approach to beam-
661   induced motion correction in cryo-EM single-particle analysis. *IUCrJ* **6**, 5–17
662   (2019).
663   45.   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development
664   of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486–501 (2010).
665   46.   Liebschner, D. *et al.* Macromolecular structure determination using X-rays,
666   neutrons and electrons: recent developments in *Phenix*. *Acta Crystallogr D*
667   *Struct Biol* **75**, 861–877 (2019).
668   47.   Best, R. B. *et al.* Optimization of the additive CHARMM all-atom protein force
669   field targeting improved sampling of the backbone ϕ, ψ and side-chain $\chi(1)$ and
670           $\chi(2)$ dihedral angles. *J Chem Theory Comput* **8**, 3257–3273 (2012).
671   48.   Huang, J. *et al.* CHARMM36m: an improved force field for folded and
672   intrinsically disordered proteins. *Nat Methods* **14**, 71–73 (2017).
673   49.   Jo, S., Kim, T. & Im, W. Automated builder and database of protein/membrane
674   complexes for molecular dynamics simulations. *PLoS One* **2**, e880 (2007).
675   50.   Lee, J. *et al.* CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER,
676   OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive
677   Force Field. *J Chem Theory Comput* **12**, 405–13 (2016).
678   51.   Bussi, G., Donadio, D. & Parrinello, M. Canonical sampling through velocity
679   rescaling. *J Chem Phys* **126**, 014101 (2007).
680   52.   Parrinello, M. & Rahman, A. Polymorphic transitions in single crystals: A new
681   molecular dynamics method. *J Appl Phys* **52**, 7182–7190 (1981).
682   53.   Gromacs 2020.1. https://doi.org/10.5281/zenodo.3685920.
683   54.   Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J Mol*
684   *Graph* **14**, 33–8, 27–8 (1996).
685
686

Acknowledgements

We thank the Central Electron Microscopy Facility of the Max Planck Institute of

Biophysics for cryo-EM infrastructure and technical support, and Juan Castillo-

Hernández and Özkan Yildiz for support in cryo-EM data processing. This work was

supported by the German Research Foundation via SPP1879 to JV and IH (VO 1449/1-

1 and HA 6322/4-1). Research in PJS's lab is also funded by Wellcome (208361/Z/17/Z)

and BBSRC (BB/P01948X/1, BB/R002517/1 and BB/S003339/1). This project made use

of time on ARCHER2 and JADE2 granted via the UK High-End Computing Consortium

for Biomolecular Simulation, HECBioSim (http://hecbiosim.ac.uk), supported by

EPSRC (grant no. EP/R029407/1). This project also used Athena and Sulis at HPC

Midlands+, which were funded by the EPSRC on grants EP/P020232/1 and

EP/T022108/1. We thank the University of Warwick Scientific Computing Research

Technology Platform for computational access.

Author contributions

J.V. and I.H. conceived this study. All authors designed the experiments. M.F.F.

performed in vivo and DSF experiments. M.F.F., Y.H. and I.T. purified KimA for cryo-

EM. J.P.W, J.S.S. and J.V. performed the cryo-EM analysis, and built and validated the

atomic models. R.A.C. performed MD simulations. All authors participated in the data

analysis. M.F.F., J.P.W. and Y.H. wrote the initial draft, all authors participated in

manuscript editing and revision. P.J.S., J.V. and I.H. supervised work and secured the

funding for this work.

Competing interests

The authors declare no competing interests.

Materials & Correspondence

Correspondence to Janet Vonck (janet.vonck@biophys.mpg.de) or Inga Hänelt

(haenelt@biochem.uni-frankfurt.de).