Parental care shapes the evolution of molecular genetic variation

Mashoodh R[1,2]*, Trowsdale A[2], Manica A[2], Kilner RM[2]

[1]Centre for Biodiversity & Environment Research, Department of Genetics, Evolution and
Environment, University College London, United Kingdom
[2]Department of Zoology, University of Cambridge, United Kingdom

*Corresponding author
Email: r.mashoodh@ucl.ac.uk

Abstract

Cooperative social behaviours, such as parental care, have long been hypothesized to relax selection leading to the accumulation of genetic variation in populations. Although the idea has been discussed for decades, there has been relatively little experimental work to investigate how social behaviour contributes to genetic variation in populations. Here, we investigate how parental care can shape molecular genetic variation in the subsocial insect *Nicrophorus vespilloides*. Using whole genome sequencing of populations that had evolved in the presence or absence of parental care for 30 generations, we show that parental care maintains levels of standing genetic variation. In contrast, under a harsh environment without care, strong directional selection caused a reduction in genetic variation. Furthermore, we show that adaptation to the loss of care is associated with genetic divergence between populations at loci related to stress, morphological development and transcriptional regulation. These data reveal how social behaviour is linked to the genetic processes that shape and maintain genetic diversity within populations, and provides rare empirical evidence for an old hypothesis.

Lay Summary

Social behaviours, such as parental care, have long been hypothesized to result in the accumulation of genetic variation in populations. Here, we investigate how parental care can shape molecular genetic variation in a species that perform biparental care, *Nicrophorus vespilloides*. Using genome sequencing of populations that had evolved in the presence or absence of parental care for 30 generations, we show that parental care maintains levels of standing genetic variation. In contrast, under a harsh environment without care, populations lost genetic variation. Furthermore, we show that adaptation to the loss of care is associated with genetic divergence between populations at genes related to stress, morphological development and transcriptional regulation. These data reveal how social behaviour is linked to the genetic processes that shape and maintain genetic diversity within populations.

Keywords: parental care, genetic variation, adaptation, mutation load, population genetics

Introduction

While much recent work has focused on identifying genes that drive social

behaviours [1,2], relatively few studies have examined the longstanding hypothesis that

social behaviour affects the accumulation and maintenance of genetic variation. Yet, social

living is associated with large-scale restructuring and the evolution of genome organization

and architecture [3]. In humans, benevolent social activities, such as modern health care,

are thought to have led to the accumulation of deleterious mutations within populations [4,5].

Therefore, the extent to which genetic variation is shaped by social behaviour has

implications for the health of populations and their capacity to rapidly adapt to environmental

perturbations. However, there have been few empirical tests of how social behaviour might

drive genetic variation in practice. Here we investigate how a cooperative social behaviour,

namely the supply of parental care, contributes to genome-wide levels of genetic variation.

We focus on parental care in a subsocial pair-breeding insect, rather than more elaborate

forms of sociality, to avoid the confounding effects of extreme reproductive skew on genetic

variation, which is common in cooperative insect societies [6].

Cooperative social interactions often function to shield social partners from a harsh

physical environment and the same is true for parental care [7,8]. Without cooperation

generally, and care specifically, individuals would be exposed to strong, frequently

directional, selection pressures from the abiotic environment, which would favour the

evolution of new adaptations and cause an associated reduction in genetic variation. On the

other hand, the presence of parental care relaxes selection from this wider environment,

theoretically allowing genetic variation to accumulate. Indeed, several lines of evidence

suggest that cooperative social behaviours, including care, can relax selection sufficiently to

allow mildly deleterious mutations to accumulate within populations [9–13]. In this way,

parental care, could shift the 'mutation-selection' balance by relaxing selection and

preventing the elimination of new spontaneous mutations. The resulting increase in genetic

variation could emerge in the form of single nucleotide polymorphisms (SNPs) and/or other

structural genetic variants (e.g., indels, transposable elements and/or insertions/deletions)

depending on the natural mutation rate of such variants. Exactly how care might maintain

such variants has been the subject of some speculation [9]. One possibility is that "cryptic"

variants could be maintained in the population with a combination of care-induced genetic

capacitors, epigenetic modifications and/or RNA-mediated signals [14]. Nevertheless,

although the suggestion that cooperative social behavour can shape genetic variation is

relatively longstanding, we still have a poor understanding of how and where it might cause

change at a molecular genetic level.

Here, we use evolving populations of burying beetles (*N. vespilloides*) to explore how

parental care affects levels of standing genetic variation and how populations may adapt in

the face of its loss. In natural populations of this locally abundant subsocial insect, burying

beetle parents raise their young on a carrion nest, formed from a small dead animal, such as

a mouse or songbird. There is continuous variation in the level of parental care supplied,

with around 5% of parents abandoning the brood before their young have even hatched [15].

Offspring can survive without parental care, at least in the laboratory.

We exploited this natural variation in care to establish two types of experimentally

evolving populations in the laboratory, which varied only in the family environment that

larvae experienced during development, and where the same family environment was

created for successive generations within populations. In Full Care populations (FC), parents

remained with their young throughout development; whereas in No Care populations (NC),

parents were removed just prior to hatching. No Care populations rapidly adapted to this

regime (within 14 generations), with adaptive change being detectable through increases in

breeding success and larval density (see Schrader et al., 2017). Moreover, we have

previously shown that No Care populations evolved adaptively [16] and divergently from Full

Care populations in the extent of the pre-hatching care behaviours [17], the extent of sibling

cooperation [18,19] and in their larval morphology [20].

At the 30th generation of experimental evolution, we use pooled whole-genome re-

sequencing of these populations to document genetic variation at the molecular level when

care was present and when it was prevented experimentally (**Figure 1a**). First, we

determined the effect of care on within-population genetic variation (SNP diversity). Second,

we identified the genetic loci that had diverged to the greatest extent following the removal of

care by looking for regions of high genetic differentiation (F_{ST}) between experimental

populations.

Methods

Breeding design & Experimental Evolution

We sampled DNA from experimental populations of *Nicrophorus vespilloides* that had been

evolving under different regimes of parental care, and which were founded from a single

genetically diverse population generated by interbreeding beetles from multiple wild

populations across Cambridgeshire. These populations have been described in detail

previously [19], and comprise a total of 4 populations: two blocks (Block 1 and Block 2;

separated by 1 week) containing two populations evolving with (FC_{POP}) or without parental

care (NC_{POP}). For the first 14 generations, when directional selection was high, an average

of 34 pairs of unrelated beetles were bred at each generation [19]. Thereafter, populations

were maintained with an average of 37 and 49 pairs at each generation for FC and NC

populations, respectively, and were equivalently successful across the generations (**Table**

S1). On the 29th generation (as in every generation previously), we paired sexually mature

males and females within each population. Each pair was placed in a separate breeding box

with moist soil and a thawed carcass (10-12g). We then placed each breeding box in a

cupboard and allowed parents to prepare the carcass and for the female to lay the clutch of

eggs. For the NC_{POP}, after 53h, both parents were removed from the nest just as had

occurred for the prior 29 generations. Approximately 80h after hatching we randomly

selected 2-3 larvae from each family (15-18 families per population) for DNA extraction.

Larval tissue dissection, DNA extraction & Whole-genome sequencing

For each family, DNA from first instar larvae were pooled and extracted using a modified

version of the Qiagen DNEasy Mini Kit. Total DNA quality was checked using gel

electrophoresis and yield was quantified using a Qubit DNA Assay Kit (Thermo Fisher).

Families were pooled in equimolar concentrations such that each individual was represented

equally to generate 4 libraries: FC1, FC2, NC1, and NC2 with pool sizes of 41, 52, 52, 59,

respectively. Whole genome re-sequencing libraries were constructed and sequenced

(150bp paired-end) at a depth of 100x using an Illumina Novaseq 6000 platform by

Novogene (Hong Kong).

Bioinformatic Analyses

Reads were trimmed using TrimGalore (0.5.0; https://github.com/FelixKrueger/TrimGalore)

to remove adaptor sequences, perform quality trimming and discard low-quality reads.

Reads were aligned in paired-end mode using the burrows-wheeler aligner (bwa) to the *N.*

vespilloides reference genome (NCBI Refseq Assembly: GCF_001412225.1) [21,22]. See

Table S2 for read mapping statistics. Duplicates were removed using PicardTools

(http://broadinstitute.github.io/picard/). Pileup files were created using *samtools* [23] from

mapped reads and indels and repeats were filtered using the *Popoolation* toolbox [24].

These pileup files were used to calculate measures of genetic diversity with *Popoolation* (π,

Watterson's θ, synonymous vs non-synonymous rate of π, and Tajima's D). Pileups were

were merged into a single sync file using *Popoolation2* for use with *poolfstat* [25], *Baypass*

[26] and *Popoolation2* to measure between-population divergence (e.g., F_{ST}, Fishers' Exact

Tests and Bayesian auxillary models; described below). All subsequent post-processing and

statistical analyses were performed in *R* version 4.1.2 using the core *R stats* package [27].

Data wrangling and visualisations were performed using the *tidyverse* suite [28].

Intrapopulation genetic variation

We used π and Watterson's θ to measure levels of standing genetic variation within

populations. Watterson's θ represents the expected number of segregating sites observed

between a pair of homologous sequences sampled from a given population, whereas π is

the average number of pairwise difference between all possible pairs of individuals in the

sample. These measures were calculated for non-overlapping 1000bp windows (for sites

with coverage between 40-700 reads) across the genome using tools from *Popoolation*. We

also computed gene-wise synonymous and non-synonymous pi for CDS coordinates of all

genes extracted from the reference annotation using *Popoolation*. To allow comparisons to

F_{ST} windows we computed Tajima's D for 500bp sliding windows with a 250bp overlap. For

all genetic diversity measures, we used non-parametric Kruskal-Wallis tests to test for

differences between Full Care and No Care populations separately for each replicate block

except in the case of Tajima's D. For normally-distributed Tajima's D values we used t-tests

to test for differences between care conditions within each block. Windows were filtered so

that statistics were based on windows that were covered across all replicates of both

populations.

Genetic divergence between populations

To estimate population structure and demographic history, we extracted SNPs from

the population sync file using the R package *poolfstat* [25] using the core model of *BayPass*

version 2.3 [29]. Baypass uses allele-frequencies to estimate a scaled covariance (Ω) matrix,

which can be interpreted as the pairwise estimates of differentiation between the population.

The Ω matrix was converted to a corrrelation matrix in R and visusalised as a tree using the

base R *stats* package.

To further measure the extent of genetic divergence between populations, we used

Popoolation2 [30] to calculate the pairwise fixation index (F_{ST}) for all combinations of

population pairs across 500bp sliding windows (250bp overlap) across the genome. SNPs

were called using sites with read counts between 40 and 700. Hierarchical clustering

indicated that NC1 and NC2 were more closely related to their FC counterparts than to each

other (**Figure S1**). Moreover, inspection of F_{ST} values across the genome indicated that the

overall magnitude of differences between FC and NC differed between the blocks (**Figure**

S2). Therefore, to identify windows where evolving populations may have diverged

consistently, over and above any variation within and between blocks, we computed F_{ST} for

each replicate line separately (i.e., FC1;NC1 and FC2;NC2). We then performed Fisher's

Exact Tests for each of these windows to screen for significant allele frequency differences.

We took the product of the -log(*p*) values for each block (FC1;NC1 *x* FC2;NC2) and selected

the top 0.5% of values as regions of interest. In this way, we selected for loci which diverged

consistently across the blocks, assuming that inconsistent divergence may reflect drift.

Location of windows of interest were annotated using the reference genome and the

intersect command in *bedtools* [31]. A hit was considered only if the window intersected with

the coordinates of the annotation by at least 1bp.

Using the same logic we used the auxillary model in BayPass to identify candidate

SNPs that were consistently associated with the loss of care across both blocks. Using the

covariance structure among the population allele frequencies (Ω), the model explicitly

accounts for the shared history of the populations, rendering the identification of SNPs

potentially subjected to selection less sensitive to the confounding effect of demography

[26,32]. Specifically, the model involves the introduction of a binary auxiliary variable to

classify each locus as being associated or not with the loss of care. This allows the

estimation of posterior inclusion probabilities (and Bayes factors) for each SNP while also

accounting for multiple testing issues. For each SNP, the Bayes factor was expressed in

deciban units (dB) via the transformation 10log10(BF). Significance was assessed based on

the Bayes Factor (BF) between models and SNP markers with strong evidence (BF > 20)

were retained as potential candidates of interest (according to Jeffrey's rule) [33]. We then

examined where these SNPs were located by looking for genes within 500bp of the outlier

SNP (using *bedtools* 'window') making this comparable to our windowed approach.

163
Functional Annotation

Functional enrichment analyses were conducted using the topGO R package version 2.38.1

[34] to identify over-representation of particular functional groups within the diverged genes

in response to the removal of care, based on GO classifications using Fisher's exact test.

GO terms were annotated to the *N. vespilloides* genome using the BLAST2GO (version

5.1.1) workflow to assign homologs to the *Drosophila* non-redundant protein databases [35].

To improve the GO term assignment, *N. vespilloides* genes were further annotated using a

custom script that assigned GO terms from multiple well-annotated insect species (e.g., *A.*

mellifera, B. terrestris, A. cephalotes, N. vitripennis, T. castaneum and *O. taurus*) based on

ortholog assignments obtained using Orthofinder [36] using a custom pipeline

(https://github.com/chriswyatt1/Goatee). To identify transcription factors we searched for the

presence of known Pfam [37] transcription factor domains in the protein sequences of the

gene candidates of interest using Interproscan [38]. Putative promoter regions (5' UTRs)

were classified as the 500bp region upstream of each gene start coordinate [31].

178
Results

Standing genetic variation between populations

First, we determined the effect of care on within-population standing genetic variation by

measuring genetic diversity. We computed both Watterson's theta (θ) and Pi (π) statistics for

each population across 1000bp non-overlapping windows. Populations that evolved under

Full Care (FC1 and FC2) had higher theta values than populations evolved under No Care
(NC1 and NC2) (**Table 1**; Figure **1c**; all *p*'s < 0.001). Similarly, there were higher Pi values in
FC compared to NC, though this effect was not present in Block 1 (**Table 1**; **Figure 1b**).
Together, these results suggest that FC populations maintained more SNP diversity
compared to populations evolving under NC with some detectable variation between blocks
(Table 1). We measured Tajima's D (500 bp overlapping windows) to further characterize
the evolutionary forces shaping genetic diversity between populations. We show that
genome-wide levels of Tajima's D are negative, with both replicates showing a significant
reduction in Tajima's D in NC compared to FC (**Table 2**; **Figure 3**).
193

194
Figure 1. (a) Populations evolved in the presence (Full Care; FC) or in the absence of (No Care; NC)
for 30 generations (two replicates per condition). Larvae were pooled for each replicate population
(see Methods) for whole genome sequencing (WGS). Distribution (top) and median (bottom) of (b) Pi
(π) and (c) Watterson's theta (θ)across 1000bp non-overlapping windows for FC and NC populations
(error bars represent 95% bootstrapped confidence intervals). Block 1 (dashed line) and Block 2 (solid
line) are plotted separately.
201
202
203 **Table 1**. Genetic diversity measures for each population evolving under Full Care (FC) and No Care
204 (NC) for each block. Delta is the difference between FC and NC populations computed separately for
205 each block (* indicates p < 0.05, ** indicates p< 0.001).

		Evolving Population				
	Block	FC	NC	Δ (FC-NC)	χ^2	*p*-value
Watterson's θ	1	0.00790	0.00772	1.85e-04	31.48	2.015e-08**
	2	0.00745	0.00734	1.17e-04	75.61	< 2.2e-16**
π	1	0.00769	0.00768	1e-05	1.53	0.215
	2	0.00772	0.00750	0.00022	132.42	2.2e-16**
π_N/π_S	1	0.0124	0	0.0124	46.81	7.82e-12**
(all genes)	2	0.0108	0	0.0108	83.05	< 2.2e-16**
π_N/π_S	1	0.0244	0	0.0244	4.29	0.038*
(diverged genes)	2	0.0229	0	0.0229	11.23	0.001**

[1]*Non-parametric Kruskal-Wallis rank sum test, df=1*

207
Figure 2. (a) Product of -log(*p*) values for allele frequency differences between Full Care (FC) and No
Care (NC) populations of each block (i.e., FC1;NC1 *x* FC2;NC2) for 500bp sliding windows (250bp
overlap) sorted by position. Dashed red line indicates 99.5th percentile (b) Percent of windows
overlapping with genomic features (ncRNA = non-coding RNA; 5' UTR is defined as 500bp upstream
of gene start position) and the number of genes that correspond to each category (c) Representative
enriched GO terms (biological processes) for the most diverged genes between FC and NC
populations.
215

Genetic differences between populations

Next, we identified the genetic loci that had diverged to the greatest extent following

the removal of care by looking for regions of high genetic differentiation (F_{ST}) between

experimental populations (see **Figure S1** for population structure). We looked for changes

over and above drift by looking for highly consistent divergence across the replicates using a

500bp sliding window approach (see Methods; **Figure 2a**). Highly significant windows

overlapped with both protein-coding and regulatory features of the genome, with 16% of

windows being classified as a regulatory change in contrast to 47.9% of windows in protein

coding genes (**Figure 2b**). Using this approach, we identified 648 differentiated genes

(**Table S3**), with 144 of these windows uniquely intersecting the 5' UTR regions of these

genes only (**Figure 2b**; **Table S4)**. These genes were generally enriched for GO processes

associated with morphogenesis, neural development, immunity, and hormone signaling

(**Figure 2c**; **Table S5**). To test that our approach converged with other methods, we also

identified SNP outliers using a Bayesian approach (see Methods). This method identified

3086 outlier SNPs with consistent allele frequency differences between the NC and FC

populations across both replicate blocks, which fell within 500bp of 1176 genes (**Table S6**).

These SNP outliers broadly converged on our windowed approach (220 genes; **Figure S3**)

with several key genes identified in both methods (**Table S7**).

234
235 **Table 2**. Tajima's D (mean of 500bp sliding windows) for populations evolving under Full Care (FC)
236 and No Care (NC) for each block. Statistics are presented for all windows across the genome
237 (genome-wide) as well as for windows that overlapped with diverged genes (+/- 5kb). Delta is
238 difference between Tajima's D between means of FC and NC populations (** indicates p < 0.001).

		Evolving Population				
	Block	FC	NC	Δ (FC-NC)	t (df)	*p*-value
Tajima's D	1	-0.134	-0.155	0.020	11.63[1]	< 2.2e-16*
(genome-wide)	2	-0.007	-0.053	0.046	26.16[2]	< 2.2e-16*
Tajima's D	1	0.039	0.024	0.086	3.22[3]	0.001*
(diverged genes)	2	0.205	0.119	0.015	17.59[4]	< 2.2e-16*

[1]df=1288045 [2]df=1288388 [3]df=167272 [4]df=167250

To test the hypothesis that genes selected in the No Care lost genetic variation, we

examined the gene-wise ratio of non-synonymous to synonymous π (π_N/π_S; **Table 1** &

Figure S4) and Tajima's D within 5kb of divergent loci (**Table 2; Figure 3**). Both measures

were reduced in No Care populations relative to the Full Care populations, in both replicate

blocks, and this was true genome-wide as well as for the genes identified in our divergence

screens.

246
Figure 3. (a) Genome-wide Tajima's D (mean) for 500bp sliding windows for No Care (NC) and Full
Care (FC) populations (Block 1: dashed; Block 2: solid). Average Tajima's D for FC and NC
populations along gene bodies of two genes that showed extreme Tajima's D values (bottom 5%) and
showed allele frequency divergence (Table S3) (b) cytochrome P450 6k1-like and (c) 5-
hydroxytryptamine receptor-like. See Figure S5 for replicate blocks plotted separately. All error bars
represent 95% bootstrapped confidence intervals.

Discussion

We found that populations with parental care (Full Care) had greater levels of genetic variation, in the form of higher theta and pi diversity, than the populations where care was prevented (No Care). Previous work has suggested that social behaviour contributes to genetic diversity mainly because of its effect on demography, and particularly because of its influence on effective population size (Ne). Population genetic theory predicts that genetic diversity will increase with Ne and mutation rate [39]. Previous empirical work linking behavioural and life-history traits, such as reproductive strategy, fecundity and body size with genetic diversity has suggested that these associations are ultimately mediated by changes in Ne [40–43]. However, our results cannot be explained by demography because populations were maintained at similar population sizes with no differences in fecundity [44] and no possibility of overlapping generations and/or changes in mating structure. Any small deviations in population size between care treatments were biased towards reducing genetic diversity in the Full Care populations – yet we found the opposite result. We suggest instead that the accumulation of genetic variation here is due directly to the effect of parental care in relaxing selection. The founding wild populations were inclined to provide care [20] and likely had already accumulated high levels of standing genetic variation, which was swiftly lost when we exposed populations to selection in a No Care environment.

The majority of this accumulated genetic variation is likely to be either neutral or mildly deleterious, since the majority of new mutations generally fall into either of these two categories [5,45]. Indeed, we have previously demonstrated that inbreeding of these populations resulted in faster extinction of Full Care compared to No Care populations, further suggesting that, at least some, of the variation accumulated in the presence of care was deleterious [12]. Although we measured only SNP variation here, genetic variants (e.g., insertions/deletions, transpositions) that arise through different types of mutation or recombination could also, in theory, be maintained in the population by parental care. Whether care favours particular types of mutants remains to be tested in future studies. In contrast, the harsher No Care environment imposed strong directional selection resulting in rapid adaptation [16,19] and reducing levels of standing genetic variation. We identified genetic divergence at number of loci, which were also associated with the loss of non-synonymous pi (lower π_N/π_S) and reductions in Tajima's D, a pattern that was similar to the genome-wide differences in genetic diversity. Again, this is consistent with the interpretation that No Care populations experienced strong directional selection, whilst Full Care populations harboured more potentially deleterious mutations.

Here, we follow convention in assuming that loci that diverge consistently are likely to represent adaptive genomic change, whereas inconsistent responses to the parental care

treatment (No Care versus Full Care) are due to drift. Yet alternative explanations for
inconsistent responses are also possible, and this might be particularly true for traits under
social selection (as opposed to abiotic selection pressures). Inconsistent patterns of genetic
change across experimental blocks attributed to drift might instead reflect idiosyncratic or
opportunistic responses to selection that arise through subtle variation in founding
populations [46–48]. This is not surprising given that polygenic traits can be genetically
redundant and adaptation can arise through multiple intersecting pathways and unique
combinations of alleles within a population [46,49]. Moreover, the magnitude of these
inconsistent differences, either due to drift or selection, might have been intensified by the
selection regime imposed by the social environment, depending on whether it relaxed
selection or imposed directional selection, for example, or whether the strength of selection
was modulated by genes of social partners, parents or siblings [10,50]. Such effects could
explain variation within and between replicate populations that accumulates over time.
Although we cannot distinguish idiosyncratic adaptive change from drift with our data
currently, future work using a high number of replicated populations measured across
several generations could provide key insights into these evolutionary dynamics [46].
306   Our data suggest that No Care populations diverge from Full Care populations at loci
that could promote immunity, metabolic and behavioural stress resilience in the absence of
care. The loss of care in *N. vespilloides* is likely to be associated with greater levels of
environmental stress during development and heightened exposure to pathogens from the
carrion resource [51,52]. We have previously shown that adaptation to a No Care
environment is associated with gene expression signatures that show blunted stress
responses and compensatory expression in metabolic and developmental pathways [51].
Not surprisingly, many of the genetic differences between the populations are in upstream
regions and/or genes that encode for transcription factors or chromatin modifiers, suggesting
that change in regulatory function is a key component of adaptation to the loss of care. This
is likely to be an underestimate of the extent of regulatory change, as we have yet to
characterise the regulatory landscape of *N. vespilloides* and windows without an annotated
overlap could be in distal promoter and/or enhancer regions. Nevertheless, differences in
regulatory functions could shape levels of gene expression of other genes, further buffering
against stress in the absence of parental care [51]. In this way, the signatures of adaptation
to the loss of parental care are not much different to adaptive genetic responses to other
abiotic stressors in the broader environment. Indeed, a key feature of stress adaptation
across species is that it involves changes in gene regulatory pathways and this is true from
bacteria to plants and animals [46,53–55] . Although we cannot identify a single gene or
master regulator within the regulatory changes, these data do identify candidate regulatory

genes that might play key roles in conferring resilience to the loss of care, and to

environmental stressors more broadly.

Delving more deeply into loci at which we detected the greatest differences, we found

that adaptation to the loss of care involved changes in several genes associated with

immune function (e.g., *CD109 antigen* and *lysozyme c-1*), which could help cope with the

increased exposure to the bacterial pathogens of the carcass nest experienced by No Care

larvae. Previous work on *N. vespilloides* has shown that lysozyme expression is particularly

heightened in parents immediately after the larvae hatch [56,57] and that it is likely to be

particularly important for eliminating pathogenic Gammaproteobacteria [58]. Relatedly, we

found divergence at a number of cytochrome P450 genes (*4ac1, 4ac2, 4c1, 4g15, 9e3 and*

6k1-like; Table S3), which are known to be involved in the metabolism of endogenous

compounds as well as exogenous toxins and disease vectors, and which might also

participate in defensive responses [59].

Cytochrome genes also appear to play a role in mediating the response to social

density in Drosophila. Expression changes at the *Cyp4*, *Cyp6* and *Cyp9* gene families can

be induced by manipulating social density in *Drosophila* and deletions of the *Cyp6a20* gene

have been associated with higher levels of aggression and reduced sociality [60]. This is

particularly interesting given that we have previously shown that larvae from the No Care

populations evolved to show greater levels of sibling cooperation than larvae from the Full

Care populations [13,18]. Cytochrome P450 families tend to share similar functional

domains, and therefore, it is possible that changes in these genes could have effects on

social behaviour *via* its actions on multiple hormonal systems (e.g., pheromones, ecdysone)

[59,61]. Furthermore, P450 genes are also intertwined with juvenile hormone pathways

which are known to be involved in multiple facets of behavioural and morphological

development [62]. This could include adaptations that aid in locating and facilitating the use

of the carrion breeding resource, such as the increase in relative mandible size and reduced

arrival time at the carcass that we also detected in No Care larvae [20,63]. This

interpretation is additionally consistent with most of the genetic hits belonging to cell

signaling and biosynthetic pathways which fall into GO categories associated with

morphological, brain and olfactory development.

We also found changes in neuropeptides that are also involved in metabolic,

homeostatic and feeding pathways (e.g., *orexin, 5-hydroxytryptamine and cholecystokinin*

receptors), raising the possibility that these could represent adaptations in larvae for feeding

and extracting nutrients from the carcass resource in the absence of parents [64]. Previous

work has shown that genes ancestrally associated with metabolic, homeostatic and feeding

pathways can be co-opted to serve new social functions [65,66]. For example, the

oxytocin/vasopressin system is commonly associated with the expression of parental care,

pair bonding and other affiliative social behaviours in mammals [67], but has an ancestral

role associated with promoting water balance [68]. A recent study in the closely related

burying beetle, *N. orbicollis*, suggested that the expression of *inotocin* (the insect homologue

of oxytocin/vasopressin) was correlated with the transition to parenting, an effect that was

more pronounced in males than females [65]. *Takeout* is another gene that, despite being

typically associated with feeding and circadian rhythms, has been shown to be highly

expressed whilst parenting in burying beetles and may be involved in the transition from

infanticide to larval care [66,69,70]. We found divergence in both an *oxytocin receptor* and a

takeout homologue (Table S3), which could explain how male parental care eventually

decayed in the No Care lines [71]. Finally, *angiotensin converting enzyme* was one of the

most diverged genes in our analyses. This gene has varied roles from conferring immunity to

regulating neuropeptide signaling [72]. However, it also appears to be strongly expressed in

insect reproductive tissues and may, therefore, play a role in adaptations in mating and

fecundity between the populations [44].

While we show here that parental care contributes to genetic variation through its

effect on selection, it is possible that the incidence of mutation is itself reduced by the loss of

parental care. Mutation rates have a strong genetic basis and can vary between individuals

and amongst populations [73,74]. Given that the loss of care is a major developmental

stressor, and that stress has been shown to induce mutations, adaptation to the loss of care

could involve genetic mechanisms that dampen and/or buffer the consequences of new

mutations that arise [9,45]. Consistent with this hypothesis, we found high levels of genetic

differentiation amongst genes involved in DNA replication and repair (e.g., *Artemis* and the

PAXIP1 interacting protein; Table S3) [75,76]. These genes, as part of their role in stress

regulation, could facilitate efficient DNA repair, purging new mutations and shaping the

subsequent mutation load of a population. The observation that multiple transfer RNAs

(tRNAs) show divergence is particularly interesting given that variation in tRNAs have been

associated with increased mutation loads *via* transcription assisted mutagenesis [77]. In

other words, both genetic and phenotypic adaptations within each population could favour

an optimal mutation-selection balance, resulting in different levels of standing genetic

variation based on levels of care experienced within each population. Further functional

characterization of these changes would help clarify if parental care facilitates the evolution

of the mutation rate, potentially providing another mechanism for divergence in mutation-

selection balance among populations [45,78].

In short, we have shown that parental care allows genetic variation to accumulate by

relaxing selection. When care is lost, a number of genetic changes quickly follow which may

be adaptive and which result in the loss of standing genetic variation. Better functional

characterization of these gene targets and regulatory regions is now required to understand

the genetic causes and functional consequences of the differences we have found between

populations that are, and are not, exposed to post-hatching care. These are key areas of

future work that will help explain whether the maintenance of standing genetic variation

under parental care is likely to help or hinder adaptation in a rapidly changing world.

Data Access & Code Availability

All raw sequencing data generated have been submitted to the Sequence Read Archive

(SRA) under BioProject ID: PRJNA934336. All code for the analyses contained within this

manuscript can be found at: https://github.com/r-mashoodh/nves_dnaEvol

Acknowledgements: We thank Benjamin Jarrett and Darren Rebar for their help with beetle

population maintenance. The molecular aspects of this project were supported by a

Biotechnology and Biological Sciences Research Council Future Leaders Fellowship

(BB/R01115X/1) and a Royal Society Small Research Grant (RGS\R1\191162) to RM. The

experimental population work was supported by a Consolidator's Grant from the European

Research Council (310785 Baldwinian_Beetles), by a Wolfson Merit Award from the Royal

Society and The Leverhulme Trust (RPG-2018-232) each to RMK. This work was also

supported by a Newton Trust grant (20.40(b)) held jointly by RM and RMK. Finally, we would

like to thank the editors, Allen Moore and two anonymous reviewers for their thoughtful

comments which significantly improved the manuscript.

Author Contributions: RM and RMK conceived of the study. RM and AT conducted the

experimental work. RM and AM analysed the data. RM wrote the paper. All authors

discussed the results and commented on the manuscript.

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Cunningham CB. 2020 Functional genomics of parental care of insects. *Hormones and Behavior* **122**, 104756. (doi:10.1016/j.yhbeh.2020.104756)

2. Toth AL, Robinson GE. 2007 Evo-devo and the evolution of social behavior. *Trends in Genetics* **23**, 334–341. (doi:10.1016/j.tig.2007.05.001)

3. Rubenstein DR *et al.* 2019 Coevolution of Genome Architecture and Social Behavior. *Trends in Ecology & Evolution* **34**, 844–855. (doi:10.1016/j.tree.2019.04.011)

4. Kondrashov AS. 2017 *Crumbling Genome: The Impact of Deleterious Mutations on Humans.* Hoboken, NJ, USA: John Wiley & Sons, Inc. (doi:10.1002/9781118952146)

5. Lynch M. 2016 Mutation and Human Exceptionalism: Our Future Genetic Load. *Genetics* **202**, 869–875. (doi:10.1534/genetics.115.180471)

6. 1979 *Social Insects.* Elsevier. (doi:10.1016/B978-0-12-342201-9.X5001-0)

7. Cornwallis CK, Botero CA, Rubenstein DR, Downing PA, West SA, Griffin AS. 2017 Cooperation facilitates the colonization of harsh environments. *Nat Ecol Evol* **1**, 0057. (doi:10.1038/s41559-016-0057)

8. Royle NJ, Smiseth PT, Kölliker M, editors. 2012 *The evolution of parental care.* First edition. Oxford, United Kingdom: Oxford University Press.

9. Snell-Rood EC, Burger M, Hutton Q, Moczek AP. 2016 Effects of parental care on the accumulation and release of cryptic genetic variation: review of mechanisms and a case study of dung beetles. *Evol Ecol* **30**, 251–265. (doi:10.1007/s10682-015-9813-4)

10. Linksvayer TA, Wade MJ. 2009 GENES WITH SOCIAL EFFECTS ARE EXPECTED TO HARBOR MORE SEQUENCE VARIATION WITHIN AND BETWEEN SPECIES. *Evolution* **63**, 1685–1696. (doi:10.1111/j.1558-5646.2009.00670.x)

11. Pilakouta N, Jamieson S, Moorad JA, Smiseth PT. 2015 Parental care buffers against inbreeding depression in burying beetles. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 8031–8035. (doi:10.1073/pnas.1500658112)

12. Pascoal S, Shimadzu H, Mashoodh R, Kilner RM. 2023 Parental care results in a greater mutation load, for which it is also a phenotypic antidote. *Proc. R. Soc. B.* **290**, 20230115. (doi:10.1098/rspb.2023.0115)

13. Schrader M, Jarrett BJM, Kilner RM. 2018 Parental care and sibling competition independently increase phenotypic variation among burying beetle siblings: BRIEF COMMUNICATION. *Evolution* **72**, 2546–2552. (doi:10.1111/evo.13607)

14. Paaby AB, Rockman MV. 2014 Cryptic genetic variation: evolution's hidden substrate. *Nat Rev Genet* **15**, 247–258. (doi:10.1038/nrg3688)

15. Scott MP. 1998 The ecology and behavior of burying beetles. *Annu. Rev. Entomol.* **43**, 595–618. (doi:10.1146/annurev.ento.43.1.595)

16. Schrader M, Jarrett BJM, Kilner RM. 2015 Using Experimental Evolution to Study Adaptations for Life within the Family. *The American Naturalist* **185**, 610–619. (doi:10.1086/680500)

17. Duarte A, Rebar D, Hallett AC, Jarrett BJM, Kilner RM. 2021 Evolutionary change in the
construction of the nursery environment when parents are prevented from caring for their
young directly. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2102450118.
(doi:10.1073/pnas.2102450118)

18. Rebar D, Bailey NW, Jarrett BJM, Kilner RM. 2020 An evolutionary switch from sibling
rivalry to sibling cooperation, caused by a sustained loss of parental care. *Proc Natl*
Acad Sci USA **117**, 2544–2550. (doi:10.1073/pnas.1911677117)

19. Schrader M, Jarrett BJM, Rebar D, Kilner RM. 2017 Adaptation to a novel family
environment involves both apparent and cryptic phenotypic changes. *Proc. R. Soc. B.*
284, 20171295. (doi:10.1098/rspb.2017.1295)

20. Jarrett BJM, Evans E, Haynes HB, Leaf MR, Rebar D, Duarte A, Schrader M, Kilner RM.
2018 A sustained change in the supply of parental care causes adaptive evolution of
offspring morphology. *Nat Commun* **9**, 3987. (doi:10.1038/s41467-018-06513-6)

21. Cunningham CB *et al.* 2015 The Genome and Methylome of a Beetle with Complex
Social Behavior, *Nicrophorus vespilloides* (Coleoptera: Silphidae). *Genome Biol Evol* **7**,
3383–3396. (doi:10.1093/gbe/evv194)

22. Li H, Durbin R. 2009 Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754–1760. (doi:10.1093/bioinformatics/btp324)

23. Danecek P *et al.* 2021 Twelve years of SAMtools and BCFtools. *GigaScience* **10**,
giab008. (doi:10.1093/gigascience/giab008)

24. Kofler R, Orozco-terWengel P, De Maio N, Pandey RV, Nolte V, Futschik A, Kosiol C,
Schlötterer C. 2011 PoPoolation: A Toolbox for Population Genetic Analysis of Next
Generation Sequencing Data from Pooled Individuals. *PLoS ONE* **6**, e15925.
(doi:10.1371/journal.pone.0015925)

25. Gautier M, Vitalis R, Flori L, Estoup A. 2022 *f*-Statistics estimation and admixture graph
construction with Pool-Seq or allele count data using the R package *poolfstat. Molecular*
Ecology Resources **22**, 1394–1416. (doi:10.1111/1755-0998.13557)

26. Gautier M. 2015 Genome-Wide Scan for Adaptive Divergence and Association with
Population-Specific Covariates. *Genetics* **201**, 1555–1579.
(doi:10.1534/genetics.115.181453)

27. R Core Team. 2019 R: A Language and Environment for Statistical Computing.

28. Wickham H *et al.* 2019 Welcome to the Tidyverse. *JOSS* **4**, 1686.
(doi:10.21105/joss.01686)

29. Olazcuaga L *et al.* 2020 A Whole-Genome Scan for Association with Invasion Success
in the Fruit Fly Drosophila suzukii Using Contrasts of Allele Frequencies Corrected for
Population Structure. *Molecular Biology and Evolution* **37**, 2369–2385.
(doi:10.1093/molbev/msaa098)

30. Kofler R, Pandey RV, Schlotterer C. 2011 PoPoolation2: identifying differentiation
between populations using sequencing of pooled DNA samples (Pool-Seq).
Bioinformatics **27**, 3435–3436. (doi:10.1093/bioinformatics/btr589)

31. Quinlan AR, Hall IM. 2010 BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842. (doi:10.1093/bioinformatics/btq033)

32. Günther T, Coop G. 2013 Robust Identification of Local Adaptation from Allele
Frequencies. *Genetics* **195**, 205–220. (doi:10.1534/genetics.113.152462)

33. Jeffreys H. 1998 *Theory of probability*. 3rd ed. Oxford [Oxfordshire]⬚: New York:
Clarendon Press⬚; Oxford University Press.

34. Alexa A, Rahnenfuhrer J. In press. topGO: Enrichment Analysis for Gene Ontology.

35. Gotz S *et al.* 2008 High-throughput functional annotation and data mining with the
Blast2GO suite. *Nucleic Acids Research* **36**, 3420–3435. (doi:10.1093/nar/gkn176)

36. Emms DM, Kelly S. 2019 OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol* **20**, 238. (doi:10.1186/s13059-019-1832-y)

37. Mistry J *et al.* 2021 Pfam: The protein families database in 2021. *Nucleic Acids*
Research **49**, D412–D419. (doi:10.1093/nar/gkaa913)

38. Blum M *et al.* 2021 The InterPro protein families and domains database: 20 years on.
Nucleic Acids Research **49**, D344–D354. (doi:10.1093/nar/gkaa977)

39. Charlesworth B. 2009 Effective population size and patterns of molecular evolution and
variation. *Nat Rev Genet* **10**, 195–205. (doi:10.1038/nrg2526)

40. Romiguier J *et al.* 2014 Comparative population genomics in animals uncovers the
determinants of genetic diversity. *Nature* **515**, 261–263. (doi:10.1038/nature13685)

41. Bharti DK, Pawar PY, Edgecombe GD, Joshi J. 2023 Genetic diversity varies with
species traits and latitude in predatory soil arthropods (Myriapoda: Chilopoda). *Global*
Ecol Biogeogr , geb.13709. (doi:10.1111/geb.13709)

42. Settepani V, Schou MF, Greve M, Grinsted L, Bechsgaard J, Bilde T. 2017 Evolution of
sociality in spiders leads to depleted genomic diversity at both population and species
levels. *Mol Ecol* **26**, 4197–4210. (doi:10.1111/mec.14196)

43. Chak STC, Harris SE, Hultgren KM, Duffy JE, Rubenstein DR. 2022 Demographic
inference provides insights into the extirpation and ecological dominance of eusocial
snapping shrimps. *Journal of Heredity* **113**, 552–562. (doi:10.1093/jhered/esac035)

44. Schrader M. 2017 Experimental evolution reveals cryptic adaptation to a novel family
environment in burying beetle populations. *Under Review*

45. Baer CF, Miyamoto MM, Denver DR. 2007 Mutation rate variation in multicellular
eukaryotes: causes and consequences. *Nat Rev Genet* **8**, 619–631.
(doi:10.1038/nrg2158)

46. Barghi N *et al.* 2019 Genetic redundancy fuels polygenic adaptation in Drosophila. *PLoS*
Biol **17**, e3000128. (doi:10.1371/journal.pbio.3000128)

47. Schlötterer C. 2023 How predictable is adaptation from standing genetic variation?
Experimental evolution in *Drosophila* highlights the central role of redundancy and
linkage disequilibrium. *Phil. Trans. R. Soc. B* **378**, 20220046.
(doi:10.1098/rstb.2022.0046)

48. Brennan RS, deMayo JA, Dam HG, Finiguerra M, Baumann H, Buffalo V, Pespeni MH.
2022 Experimental evolution reveals the synergistic genomic mechanisms of adaptation
to ocean warming and acidification in a marine copepod. *Proc. Natl. Acad. Sci. U.S.A.*
119, e2201521119. (doi:10.1073/pnas.2201521119)

49. Láruson ÁJ, Yeaman S, Lotterhos KE. 2020 The Importance of Genetic Redundancy in
Evolution. *Trends in Ecology & Evolution* **35**, 809–822. (doi:10.1016/j.tree.2020.04.009)

50. Drown DM, Wade MJ. 2014 RUNAWAY COEVOLUTION: ADAPTATION TO
HERITABLE AND NONHERITABLE ENVIRONMENTS: BRIEF COMMUNICATION.
Evolution **68**, 3039–3046. (doi:10.1111/evo.12470)

51. Mashoodh R, Sarkies P, Westoby J, Kilner R. 2021 Evolved changes in DNA
methylation in response to the sustained loss of parental care in the burying beetle.
(doi:10.1101/2021.03.25.436923)

52. Rozen DE, Engelmoer DJP, Smiseth PT. 2008 Antimicrobial strategies in burying
beetles breeding on carrion. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 17890–17895.
(doi:10.1073/pnas.0805403105)

53. Bhargava S, Sawant K. 2013 Drought stress adaptation: metabolic adjustment and
regulation of gene expression. *Plant Breed* **132**, 21–32. (doi:10.1111/pbr.12004)

54. Conrad TM, Frazier M, Joyce AR, Cho B-K, Knight EM, Lewis NE, Landick R, Palsson
BØ. 2010 RNA polymerase mutants found through adaptive evolution reprogram
Escherichia coli for optimal growth in minimal media. *Proc. Natl. Acad. Sci. U.S.A.* **107**,
20500–20505. (doi:10.1073/pnas.0911253107)

55. De Nadal E, Ammerer G, Posas F. 2011 Controlling gene expression in response to
stress. *Nat Rev Genet* **12**, 833–845. (doi:10.1038/nrg3055)

56. Cotter SC, Kilner RM. 2010 Sexual division of antibacterial resource defence in breeding
burying beetles, *Nicrophorus vespilloides*. *Journal of Animal Ecology* **79**, 35–43.
(doi:10.1111/j.1365-2656.2009.01593.x)

57. Palmer WJ, Duarte A, Schrader M, Day JP, Kilner R, Jiggins FM. 2016 A gene
associated with social immunity in the burying beetle *Nicrophorus vespilloides*. *Proc. R.*
Soc. B. **283**, 20152733. (doi:10.1098/rspb.2015.2733)

58. Duarte A, Welch M, Swannack C, Wagner J, Kilner RM. 2018 Strategies for managing
rival bacterial communities: Lessons from burying beetles. *J Anim Ecol* **87**, 414–427.
(doi:10.1111/1365-2656.12725)

59. Nauen R, Bass C, Feyereisen R, Vontas J. 2022 The Role of Cytochrome P450s in
Insect Toxicology and Resistance. *Annu. Rev. Entomol.* **67**, 105–124.
(doi:10.1146/annurev-ento-070621-061328)

60. Wang L, Dankert H, Perona P, Anderson DJ. 2008 A common genetic target for
environmental and heritable influences on aggressiveness in Drosophila. *Proceedings of*
the National Academy of Sciences **105**, 5657–5663. (doi:10.1073/pnas.0801327105)

61. Iga M, Kataoka H. 2012 Recent Studies on Insect Hormone Metabolic Pathways
Mediated by Cytochrome P450 Enzymes. *Biological & Pharmaceutical Bulletin* **35**, 838–
843. (doi:10.1248/bpb.35.838)

62. Flatt T, Tu M-P, Tatar M. 2005 Hormonal pleiotropy and the juvenile hormone regulation
ofDrosophila development and life history. *Bioessays* **27**, 999–1010.
(doi:10.1002/bies.20290)

63. Jarrett BJM, Rebar D, Haynes HB, Leaf MR, Halliwell C, Kemp R, Kilner RM. 2018
Adaptive evolution of synchronous egg-hatching in compensation for the loss of parental
care. *Proc. R. Soc. B.* **285**, 20181452. (doi:10.1098/rspb.2018.1452)

64. Nässel DR, Zandawala M. 2019 Recent advances in neuropeptide signaling in
Drosophila, from genes to physiology and behavior. *Progress in Neurobiology* **179**,
101607. (doi:10.1016/j.pneurobio.2019.02.003)

65. Potticary AL, Cunningham CB, McKinney EC, Moore PJ, Belay AT, Moore AJ. 2022
Insect homolog of oxytocin/vasopressin associated with parenting of males but not
females in a subsocial beetle. (doi:10.1101/2022.07.06.499045)

66. Potticary AL, McKinney EC, Moore PJ, Moore AJ. 2023 *takeout* gene expression is
associated with temporal kin recognition. (doi:10.1101/2023.06.12.544599)

67. Froemke RC, Young LJ. 2021 Oxytocin, Neural Plasticity, and Social Behavior. *Annu.*
Rev. Neurosci. **44**, 359–381. (doi:10.1146/annurev-neuro-102320-102847)

68. Koto A, Motoyama N, Tahara H, McGregor S, Moriyama M, Okabe T, Miura M, Keller L.
2019 Oxytocin/vasopressin-like peptide inotocin regulates cuticular hydrocarbon
synthesis and water balancing in ants. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 5597–5606.
(doi:10.1073/pnas.1817788116)

69. Moss JB, Cunningham CB, McKinney EC, Moore AJ. 2022 Gene expression underlying
parenting and being parented shows limited plasticity in response to different ambient
temperatures. *Molecular Ecology* **31**, 5326–5338. (doi:10.1111/mec.16649)

70. Sarov-Blat L, So WV, Liu L, Rosbash M. 2000 The Drosophila takeout Gene Is a Novel
Molecular Link between Circadian Rhythms and Feeding Behavior. *Cell* **101**, 647–656.
(doi:10.1016/S0092-8674(00)80876-4)

71. Bladon EK, Pascoal S, Bird N, Mashoodh R, Kilner RM. 2023 The evolutionary demise
of a social interaction: experimentally induced loss of traits involved in the supply and
demand of care. *Evolution Letters* **7**, 168–175. (doi:10.1093/evlett/qrad016)

72. Isaac RE, Ekbote U, Coates D, Shirras AD. 1999 Insect Angiotensin-converting Enzyme:
A Processing Enzyme with Broad Substrate Specificity and a Role in Reproduction.
Annals NY Acad Sci **897**, 342–347. (doi:10.1111/j.1749-6632.1999.tb07904.x)

73. Baer CF *et al.* 2005 Comparative evolutionary genetics of spontaneous mutations
affecting fitness in rhabditid nematodes. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 5785–5790.
(doi:10.1073/pnas.0406056102)

74. Wang Y, McNeil P, Abdulazeez R, Pascual M, Johnston SE, Keightley PD, Obbard DJ.
2022 Variation in mutation, recombination, and transposition rates in *Drosophila*
melanogaster and *Drosophila simulans*. (doi:10.1101/2022.09.12.507595)

75. Kurosawa A, Adachi N. 2010 Functions and Regulation of Artemis: A Goddess in the
Maintenance of Genome Integrity. *JRR* **51**, 503–509. (doi:10.1269/jrr.10017)

76. Muñoz IM, Rouse J. 2009 Control of histone methylation and genome stability by PTIP.
EMBO Reports **10**, 239–245. (doi:10.1038/embor.2009.21)

77. Thornlow BP, Hough J, Roger JM, Gong H, Lowe TM, Corbett-Detig RB. 2018 Transfer
RNA genes experience exceptionally elevated mutation rates. *Proc. Natl. Acad. Sci.*
U.S.A. **115**, 8996–9001. (doi:10.1073/pnas.1801240115)

78. Lynch M, Ackerman MS, Gout J-F, Long H, Sung W, Thomas WK, Foster PL. 2016
Genetic drift, selection and the evolution of the mutation rate. *Nat Rev Genet* **17**, 704–
714. (doi:10.1038/nrg.2016.104)