The Effect of Growth Substrate Availability on Anaerobic Arsenic

Methylation by *Paraclostridium bifermentans* strain EML

Jiangtao Qiao, Hugo Sallet, Karin Lederballe Meibom, Nicolas Jacquemin, Rizlan

Bernier-Latmani[*]

Environmental Microbiology Laboratory, School of Architecture, Civil and

Environmental Engineering, École Polytechnique Fédérale de Lausanne,

Lausanne, Switzerland.

*Corresponding author. E-mail address rizlan.bernier-latmani@epfl.ch. Postal

address EPFL ENAC IIE EML CH A1 375 (Bâtiment CH) Station 6 CH-1015

Lausanne Switzerland. Tel. +41 21 69 35001.

Word number: ~ 7000: Text (including references): 5500; Figures: one small figure:

~300 words; two large figures: ~1200 words.

ABSTRACT

Microbial arsenic (As) methylation is established as a detoxification process under aerobic conditions (converting arsenite) to monomethylarsonic acid (MMAs(V)) but proposed to be a microbial warfare strategy under anoxic conditions due to the toxicity of its main product monomethylarsonous acid (MMAs(III)). Here, we leveraged a recently isolated anaerobic As methylator, *Paraclostridium bifermentans* strain EML to gain insights into this process. Strain EML was inoculated into a series of media that consisted of systematic dilutions of Reinforced Clostridial Broth (RCB) supplied with 25 µM arsenite to assess the impact of growth substrate availability on As methylation. Growth curves evidenced the sensitivity of strain EML to arsenite, and As speciation analysis revealed the production of MMAs(III). Concentrations of MMAs(III) were found to be positively corelated with the RCB medium dilution, suggesting that growth substrate availability plays an important role in the regulation of MMAs(III) production. Further, reverse-transcription quantitative PCR evidenced *arsM* gene transcription increased with RCB dilution. Thus, growth substrate limitation is interpreted to enhance *arsM* gene expression and the associated anaerobic As methylation. We propose that growth substrate competition between microorganisms may also lead to an increase in As methylation, providing support for a microbial warfare function for this process.

Key words: anaerobic As methylation, MMAs(III), *arsM* gene transcript, microbial warfare

41 **SYNOPSIS:** Understanding the controls on anaerobic microbial arsenic

42 methylation is relevant for rice paddy soils and the accumulation of arsenic in rice

43 grains.

INTRODUCTION

Microbial transformations play an important role in the biogeochemical cycling of arsenic (As) in the environment, and include reduction, oxidation, thiolation, methylation, and demethylation of inorganic and organic As.[1-4] These reactions impact the mobility, bioavailability, and toxicity of As compunds.[1-3, 5, 6] In recent years, microbial transformations of As in paddy soils have also drawn increasing attention because of the potential health risk of dietary exposure of As from rice-containing products.[7, 8] For instance, organic As (in particular dimethyl arsenate, DMAs(V)) is commonly detected in rice grains, along with inorganic As.[9, 10] The accumulation of DMAs(V) has been shown to correlate with rice straight-head disease.[11, 12]

Arsenic methylation is a microbially-mediated process involving the transformation of trivalent inorganic As (arsenite, As(III)) into mono-, di-, and trimethylated As compounds and is catalyzed by *S*-adenosyl-methionine methyltransferase (ArsM in prokaryotes).[13-15] Under flooded conditions, the paddy soil is anoxic, favoring the activity of anaerobic microbial As methylators, and, in turn, leading to greater accumulation of methylated As in rice grains as compared to dry cultivation conditions.[16, 17] Generally, As methylation occurring under oxic conditions is proposed as an As(III)-detoxifying process because although more toxic As compounds (monomethylarsonous acid (MMAs(III) and dimethylarsinous acid (DMAs(III)) are produced, they are rapidly oxidized in the presence of O_2 to their less toxic pentavalent counterparts (monomethylarsonic acid (MMAs(V) and dimethylarsinic acid (DMAs(V)).[15] This paradigm is supported by the fact that the heterologous expression of the *arsM* gene conferred As(III) resistance to an As(III)-sensitive *Escherichia coli* strain under aerobic condition.[17, 18]

In contrast, detoxification is unlikely to be the ecological function of As-

methylators inhabiting anoxic environments since organic As products are present

in their trivalent forms. Interestingly, the evolutionary history of the *arsM* gene

predicts its emergence during the anoxic Archaean era, when MMAs(III) would

have been chemically stable.[19, 20] This finding suggests that the original function of

MMAs(III) could have been to serve as a primitive antibiotic.[21, 22] At present, there

is only one confirmed and available anaerobic As-methylating microorganism,

Paraclostridium bifermentans strain EML, a fermenter isolated from a paddy soil in

Vietnam.[23] However, controls over its methylating activity remains unknown.

In microbial communities, competitive phenotypes (including production of

antibiotics) can arise as a consequence of limited resources.[24] Soils represent an

ecosystem in which a large number of microorganisms compete for scarce

resources, thus competition is widespread.[25] We hypothesized that competition for

resources may boost As methylation by *P. bifermentans* strain EML, conferring it

an advantage over its competitors. In this study, we aimed to investigate the effect

of growth substrate limitation on As methylation to test the microbial warfare

hypothesis. From an ecological point of view, it is indeed reasonable to expect

enhanced As methylation under substrate-limiting conditions, as an response to

resource competition. According to our hypothesis, strain EML would increase the

production of toxic MMAs(III) under growth substrate-limited conditions in order to

thwart other microorganisms competing for the same resources.

MATERIALS AND METHODS

Growth Experiment. The anaerobic As-methylating bacterium, *Paraclostridium*

bifermentans strain EML (henceforth strain EML) was previously isolated from an

anaerobic paddy soil enrichment[26] by using a multi-omics approach targeting the

MAGs harboring an expressed *arsM* gene.[23] To investigate how growth substrate

availability affects As methylation activity, dilutions (v/v) of Reinforced Clostridial

Broth (RCB) (Oxoid Ltd) medium in Milli-Q water (100% RCB, 75% RCB, 50%

RCB, and 25% RCB) were prepared in 120 mL serum bottles containing 50 mL

medium (Supporting information Table S1). The medium was brought to a boil for

5 min to remove O_2, then cooled down under a 100% N_2 gas flow to room

temperature, and dispensed into individual culture serum bottle under the same

N_2 atmosphere. The bottles were then sealed with sterile rubber stoppers and

crimped with aluminum caps and the headspace was flushed with 100% N_2 to

ensure anaerobic conditions before autoclaving at 121°C for 15 mins.

A pre-culture of strain EML was grown in RCB anaerobically to mid-

exponential growth phase. Strain EML (~0.5 mL) was inoculated into each RCB

dilution containing 25 μM As(III) as sodium arsenite (or into the equivalent no-

As(III) control) in triplicate using sterile N_2-flushed syringes and needles. The

inoculum represented approximately 1% of the total volume (v/v). All the bottles

were incubated at 30°C in the dark without shaking. A total of eight experimental

conditions were selected (Table S1). At selected time points and for each

condition, triplicate bottles were sampled for growth, which was quantified using

both optical density at 600 nm (OD_{600}) and total protein content estimated using a

BCA protein assay kit (Thermo Scientific, MA, USA). For quantification of the

expression of the *arsM* gene, triplicate cultures were sampled for RNA extraction

at 8 h and 24 h. To test the stability of MMAs(III) in RCB medium, additional

abiotic control experiments (including fresh/spent RCB medium supplemented

with MMAs(III), and chemical reaction between MMAs(III) and sulfide) were

performed in duplicate, see SI.

As Speciation. At each time point and for each condition, aqueous and

intracellular As speciation and the total As were characterized. Samples for

aqueous As species and total As were obtained from 1 mL of culture collected with

sterile, N_2-flushed syringes and needles, filtered through 0.22 µm filters, and

stored in 1 mL 1% HNO_3 (≥ 69 %, Honeywell Fluka). Additionally, to analyze As

species and total As post-sample oxidation, another 1 mL of culture was obtained

as described above and oxidized by adding 10% (v/v) hydrogen peroxide (w/v)

(H_2O_2, 30%, Reactolab SA) and was kept overnight in a 1% HNO_3 solution. To

sample soluble intracellular As species, 1 mL of culture was collected, the cells

were pelleted at 8,000 g for 5 min, and stored at -20°C until use. To release

soluble intracellular As, the cell pellets were lysed in a lysis buffer (0.1% triton X-

100, 0.1% SDS, 10 mM EDTA, and 1 mM Tris-HCl) at 95°C for 15 min by

vortexing every 3 min. The lysed cell suspension was subsequently centrifuged at

8,000 g for 5 min, and the pellet was resuspended in 200 µL 1 x PBS buffer and

used for protein determination in the manner described above. The supernatant

was filtered through 0.22 µm filters and reserved for As speciation and total As

analysis.

Both aqueous and soluble intracellular As speciation were determined by high

performance liquid chromatography and inductively coupled plasma mass

spectrometry (HPLC-ICP-MS) on an Agilent 8900 ICP-QQQ instrument. A

previously described anion exchange protocol using the step-gradient elution

mode with an As Spec anion exchange fast column (50 mm x 4.0 mm, PrinCen,

Guangzhou, China) was used.[23] Six As standards were available: MMAs(III) as

methyldiiodoarsine (Santa Cruz Biotechnology Inc.), TMAs(V)O as trimethyl

arsine oxide (Argus Chemicals Srl., Italy), DMAs(V) as sodium dimethylarsinate

(ABCR, Germany), MMAs(V) as monomethylarsonic acid (Chemservice, PA,

USA), As(III) as sodium arsenite ($NaAsO_2$) (Sigma-Aldrich, MO, USA),and As(V)

as sodium arsenate dibasic heptahydrate ($Na_2HAsO_4 \cdot 7H_2O$) (Sigma-Aldrich, MO,

USA). In addition, monomethylmonothioarsonic acid (MMMTAs(V)) was

synthesized as previously described.[27] Total aqueous and soluble intracellular As

concentrations were measured using the same ICP-MS instruments in stand-

alone mode.[23]

RNA extraction and RT-qPCR. After sampling, each culture was amended with

RNAprotect Bacteria Reagent (Qiagen, Hilden, Germany) following the

manufacturer's recommendations to stabilize RNA and prevent its degradation.

The RNeasy Mini Kit (Qiagen) was used following the manufacturer's instructions

with an initial sample preparation protocol from the Qiagen RNAprotect bacteria

reagent handbook. Protocol 5 (enzymatic lysis, proteinase K digestion, and

mechanical disruption of bacteria) was employed for cell lysis prior to RNA

purification. Genomic DNA digestion was completed during RNA purification using

the RNase-Free DNase set (Qiagen). Reverse transcription was performed with

QuantiTect Rev. Transcription Kit (Qiagen). Detailed information about designing a

specific *arsM* gene primer set, optimizing the PCR amplification condition, and

constructing an *arsM* plasmid to use for standard curve are described in SI. The

RT-qPCR was carried out in a Mic PCR system (Bio Molecular Systems, Mic)

using SYBR Green Master Mix. The reactions (10 µL total volume), contained 5

µL of 2 × SensiFAST™ SYBR No-ROX Kit (Bioline, London, UK), 0.2 µM of each

arsM gene primer, 2.5 µL of cDNA, and 1% (v/v) bovine serum albumin (BSA)

(Sigma). A 10-fold dilution series containing 10^7-10^1 copies of *arsM* plasmid DNA

was used to generate a standard curve. All samples were run in quadruplicates. A

NRT (no-reverse transcriptase control) and a NTC (no template control) were both

included as negative controls.

Quantification of *arsM* gene transcripts. To get a comprehensive

understanding of *arsM* gene expression under variable substrate conditions, both

relative and absolute quantification methods were attempted. For relative

quantification, the specific primer sets for the housekeeping genes considered

and the corresponding amplification conditions are shown in Table S2. However,

these genes are not used in this study because they do not meet the minimum

requirement for a stable reference gene (see Results section for further details).

For absolute quantification, we were concerned that the mRNA yield would be

variable across conditions due both to biological reasons (e.g., rate and extent of

growth) and biases introduced by RNA extraction. To obtain robust results, we

compared two methods of absolute quantification. The first method entailed

adjusting biomass for each culture prior to RNA extraction to ensure that RNA was

extracted from the same amount of biomass (OD_{600}) regardless of conditions. The

same volume of total RNA was used for reverse transcription, and the expression

data (*arsM* copy numbers) were adjusted to OD_{600} (biomass). The second method

consisted of quantifying extracted total RNA and using the same amount of RNA

from all conditions in reverse transcription. The expression data were then

normalized to the corresponding protein concentration.

RESULTS

As(III) inhibits strain EML growth. Anaerobic growth of *Paraclostridium*

bifermentans strain EML was conducted with various concentrations of RCB

(100%, 75%, 50%, or 25% RCB) in the presence and absence of 25 µM As(III).

The growth curves show that strain EML grew rapidly and reached the mid-

exponential phase after about 8 hours in the absence of As(III) (Figure S1 and

Table S3). In contrast, the extent of growth of strain EML was lower in the

presence of As(III). In addition, it reached the stationary and death phases sooner

than in the absence of As(III) (Figure 1), particularly in the low growth substrate

media (50% and 25% RCB) (Figure S1 and Table S4), indicating that the growth

of strain EML is negatively impacted by As(III). While strain EML harbors a gene

encoding an As(III) efflux pump (*acr3*) (Figure S2), we hypothesize that it may not

be efficient enough to pump out intracellular As(III).

MMAs(III) is produced by strain EML. To profile the dynamics of As(III)

transformation during anaerobic growth of strain EML, time-dependent changes in

aqueous and soluble intracellular As speciation were monitored. Analysis of

aqueous As species clearly shows that MMAs(III) was gradually produced by

strain EML and increased during the exponential growth phase (0-12 hours) and

reached a plateau between 24 and 48 hours (post-stationary to death growth

phase) under variable growth substrate conditions (Figures 2 and S3a-d and

Table S5). MMAs(III) also represented the dominant methylated aqueous As

species, and concentrations of MMAs(III) increased concomitantly with the

decrease in aqueous As(III) (Figure S3a-d). RCB amended with 25 µM As(III)

exhibits no transformation of As(III) (data not shown), therefore, our results

demonstrate that this anaerobic As-methylating microorganism produces

MMAs(III).

However, the chemical stability of MMAs(III) is often limited, even under

anoxic conditions. We tested this stability by amending anoxic RCB (100%-25%)

or anoxic spent RCB (in which strain EML had grown) with 3 µM MMAs(III) and

evidenced its significant disappearance from solution (after 24 h) (Figures S4 and

S5 and Tables S6 and S7). Surprisingly, MMAs(III) stability was seemingly

negatively correlated with the medium dilution (Figure S4). Thus, the chemical

stability of MMAs(III) in RCB medium (fresh or spent) is limited, suggesting that, in

fact, MMAs(III) production is underestimated by our measurements in all

conditions but more so in the more diluted RCB medium.

The disappearance of MMAs(III) from the solution was puzzling and we

hypothesized the formation of methylated and thiolated As species (e.g.,

monomethyldithioarsenate, MMDTAs(V)) that are not identifiable analytically in

our system. To probe this possibility, we analyzed As speciation after oxidation of

trivalent As species by H_2O_2, an established strategy.[28-30] Oxidation is expected to

transform MMAs(III) into MMAs(V) quantitatively. As expected, after oxidation, the

concentration of MMAs(V) was greater than that of MMAs(III) measured prior to

oxidation (Figure S3 and Table S5). We propose that compounds such as

MMDTAs(V) are formed chemically through reaction of reduced sulfur compounds

with MMAs(III). These chemical species are retained by the HPLC column but are

eluted after oxidation.[31] Thus, MMAs(V) in the oxidized sample corresponds to the

sum of MMAs(III) and methylated and thiolated As species. Direct evidence of the

formation of unknown, column-retained species is provided by the reaction of

MMAs(III) with sulfide. Indeed, MMAs(III) disappears from solution (Figure S6 and

Table S8), and MMMTAs(V) is formed. On the other hand, in the oxidized samples,

all initial MMAs(III) is recovered in the form of MMAs(V) (Figure S6).

In the oxidized samples, a low concentration of DMAs(V) (Figure S3e-f) and

an unknown As peak (Figure S7) were also detected. Similarly to above, we

propose that a thiolated and/or methylated species may be generated and
retained in the anion-exchange column. However, upon reaction with H_2O_2, it is
partially oxidized to pentavalent methylated As species (DMAs(V)) and readily
eluted and separated. The species corresponding to the abovementioned
unknown peak has not been identified. Arsenic mass balance was evaluated
either by calculating the sum of all As species detected or by measuring total As
(using ICP-MS) (Figure S8 and Table S9). The overall mass balance for As in our
study is around 74%-118% (no oxidation) and 89%-150% (oxidation), suggesting
that chemical oxidation of the sample recovers more of the As chemical species,
albeit in altered form.

Impact of variable substrate concentrations. Strain EML exhibited variable
growth rates for varying RCB dilutions, confounding the interpretation of whether
growth substrate levels affected the extent of As methylation. Normalization of
methylated As (with and without oxidation) to protein concentration (Figure 2 and
Table S10) reveals a trend in normalized MMAs(III) or MMAs(V)/DMAs(V)
concentration as a function of RCB dilution (Figure 2a). Indeed, the normalized
concentration decreased in the following order: 25% RCB > 50% RCB > 75%
RCB > 100% RCB (Figure 2a). The greatest amount of protein-normalized
MMAs(III) was produced by strain EML grown in the highest RCB dilution (25%
RCB, 7,766 ± 919 nmol/g protein), which was about 14 (546 ± 39 nmol/g protein),
34 (224 ± 22 nmol/g protein), and 51 (150 ± 21 nmol/g protein) times higher than
that in the 50%, 75%, and 100% RCB conditions, respectively (Figure 2a). Similar
patterns were also observed in the oxidized samples, with the highest protein-
normalized concentrations of MMAs(V) and DMAs(V) generated in the highest
RCB dilution (25% RCB) (Figure 2b and 2c). Analysis of soluble intracellular As

also clearly shows that strain EML accumulates high amounts of intracellular As(III)

and MMAs(V) during anaerobic As methylation, particularly in the highest RCB

dilution (25% RCB) (Figure S9 and Tables S11 and S12).

arsM gene transcription under variable substrate conditions. In order to

investigate whether the transcription of the gene responsible for As(III)

methylation (*arsM*) responded to growth substrate concentration, gene expression

was quantified using RT-qPCR. We first attempted to use relative expression

analysis and evaluated the expression stability of 8 potential reference genes

(Table S2) with the qBase plus software. Unfortunately, no optimal number of

reference genes could be found due to the relatively high variability amongst

sequential normalization factors (geNorm V > 0.15) and also only medium

expression stability was achieved (0.5 < average geNorm M ≤ 1.0) (Figure S10).

We presume that the considerable variation in growth rate due to the RCB

dilutions markedly impacted gene expression, even for so-called housekeeping

genes. Therefore, we turned to absolute quantification. We adjusted the biomass

(OD_{600}) of strain EML obtained through variable RCB dilutions to approximately

the same value prior to RNA extraction to eliminate possible biomass-related

biases in RNA extraction and reverse transcription. As expected, the transcripts of

strain EML *arsM* gene (adjusted to OD_{600}) were significantly higher ($P < 0.05$) in

the presence of As(III) compared to no As(III) controls (Figure 3a and Table S13).

Amongst the treatments with As(III), we observed that *arsM* gene copy numbers

exhibited an opposing trend to substrate content: 25% RCB > 50% RCB > 75%

RCB > 100% RCB (Figure 3a). The highest number of *arsM* transcripts was

detected in the lowest nutrient media (25% RCB + As(III), 1.04E+05 ± 2.05E+04

copies/OD_{600}), which was approximately 4, 5, and 12 times higher than in the

treatments of 50% RCB + As(III) (2.63E+04 copies/OD_{600}), 75% RCB + As(III)

(2.11E+04 copies/OD_{600}), and 100% RCB + As(III) (8.90E+03 copies/OD_{600}),

respectively (Figure 3a). This result is deemed robust because, in the absence of

As(III), the trend follows the opposite direction, i.e., *arsM* expression is highest in

the no dilution (100% RCB) condition (Figure 3a). We attribute this trend to

imperfect normalization of *arsM* expression (we presume that the expression of

arsM in all dilutions should be the same without As) and biases stemming from the

differences in expression in cells growing in substrate-replete *vs.* substrate-

depleted conditions. However, these biases only strengthen the findings reported

in the presence of As(III) because they would tend to decrease the expression of

arsM in the higher dilution conditions, while the finding reports the highest

transcript number in those conditions.

Furthermore, correlation analysis suggested a significant ($P < 0.05$) positive

correlation between *arsM* gene transcripts and the concentrations of aqueous

MMAs(III) (no-oxidation) (Figure 3b) and aqueous MMAs(V) (post oxidation)

(Figure 3c). Similarly, the correlation between *arsM* gene transcript number and

MMAs(III) content was also supported when using the second experimental

method for absolute transcript quantification, which relies on using the same

amount total RNA for reverse transcripts regardless of biomass amount used for

the extraction (Figure S11 and Table S14). In summary, our findings indicate that

growth substrate availability impacts the expression of *arsM* in the anaerobic As-

methylating strain EML, in turn, affecting the concentration of MMAs(III) produced.

DISCUSSION

In this study, we provide evidence that trivalent monomethylated As, MMAs(III) is

produced as the dominant methylated As species by the recently isolated

anaerobic As-methylating bacterium, *Paraclostridium bifermentans* strain EML,

during anaerobic growth in the presence of As(III) (Figures 1 and 2). The

concentration of MMAs(III) increases during the exponential growth phase and

remains stable in the stationary phase (Figures 1 and 2). This strongly suggests

that anaerobic As methylation resulted from the activity of a functional ArsM from

strain EML, rather than by the fortuitous methylation of As owing to the release of

methyltransferases upon cell lysis as evidenced for methanogens in a previous

study.[32] An *arsM*-containing *ars* operon (*arsM-acr3-MPPE-arsR*1) was identified in

strain EML (Figure S2), supporting this interpretation. A question remains: why

would strain EML generate a product (MMAs(III)) that is more toxic than As(III)? At

first glance, this would appear to be deleterious to the microorganism. However,

this process could result in a beneficial outcome if two conditions are fulfilled. The

first condition is that trivalent methylated As compounds serve as antibiotics to

inhibit other potentially competing microorganisms.[21] Because MMAs(III) and

DMAs(III) are thermodynamically stable under anoxic conditions, they persist

sufficiently long to be effective as antibiotics for anaerobes. The second condition

is that MMAs(III), which is produced intracellularly, is exported to the extracellular

space, precluding self-toxicity. If the rate of efflux of MMAs(III) is greater than that

of As(III), As methylation would represent a net detoxification process.

The conditions propitious for anaerobic As methylation remain elusive. A

comparative study of As methylation across aerobic and anaerobic

microorganisms revealed that, despite encoding a functional ArsM, anaerobes did

not necessarily methylate As(III).[32] This observation was partially attributed to the

efficient efflux of As(III) in the anaerobes and not in the aerobes, precluding

sufficient accumulation of As(III) intracellularly for methylation to occur. Knocking

out the *acr*3 gene, encoding the As(III) efflux pump in the anaerobe *Clostridium*

pasteurianum, resulted in an obvious increase in intracellular As(III) but, overall,

the As methylation efficiency improved little.[32] Therefore, we surmised that in

addition to intracellular As(III), other factors may control anaerobic As methylation.

If the microbial warfare hypothesis for anaerobic As methylation holds, anaerobic

As methylation might be triggered by an environmental signal suggesting

obstacles to optimal growth (e.g., limited resources) or by specific metabolites

produced by other microbial community members signaling their presence as

potential competitors.[23, 32]

Here, we aimed at simulating growth substrate limitation by growing the

anaerobic As methylator, strain EML, in dilutions of RCB medium and probed the

extent of As methylation. We found opposing trends between growth substrate

content and concentration of aqueous MMAs(III) (Figures 1 and 2) or *arsM* gene

transcript numbers (Figure 3). Strain EML grown at lower growth substrate

conditions produced higher (protein-normalized) concentrations of MMAs(III)

(Figure 2), and exhibited higher (protein-normalized) expression of the *arsM* gene

(Figures 3 and S12), suggesting that the cells responded to growth substrate

limitation by increasing As methylation. Taken together, these results demonstrate

an important role for growth substrate availability in regulating anaerobic As

methylation by strain EML. In addition, we observed the accumulation of

intracellular As(III) preferentially in the low-growth substrate conditions, confirming

the close relationship between intracellular As(III) and high As methylation

potential, as previously evidenced.[32] The findings are consistent with a microbial

warfare strategy by which, under growth substrate-limiting conditions, production

of an antibiotic would provide a competitive advantage.

However, efflux of MMAs(III) is required for effective delivery of antibiotics to

other microorganisms and, to avoid self-toxicity (as stated in condition 2 above).

Specific/nonspecific MMAs(III)-pump genes *arsP* [33] and *arsK* [34] are known and the

co-evolution of *arsM* and *arsP* was previously evidenced, suggesting a strategy of

MMAs(III) efflux by MMAs(III)-producing and MMAs(III)-resistant

microorganisms[20-22, 33, 35] The identification of two chromosomally encoded *arsP*

genes in strain EML indicates that it may be capable of effluxing MMAs(III) to the

extracellular space (Figure S12).

In this study, we provide indirect evidence of the role of growth substrate

limitation in anaerobic As methylation. Lower growth substrate levels induced the

upregulation of *arsM* gene transcription, leading to a higher production of

MMAs(III) (and derivatives resulting from chemical reactions). This finding was

interpreted as a strategy to inhibit other microorganisms with an arsenical

antibiotic. Presumably, growth substrate competition would lead to substrate

limitation that, in turn, results in the production of MMAs(III) with its attendant toxic

effect on competitors.

However, direct evidence of microbial warfare under anoxic conditions

remains to be shown. Besides growth substrate availability, other factors such as

metabolites produced by other microbial community might also play a critical role

in controlling anaerobic As methylation. Additionally, the mechanism of regulation

of *arsM* gene expression by the growth substrate concentration remains a

knowledge gap. Further work is required to elucidate other factors controlling

anaerobic As methylation and their mechanism to help develop potential

strategies to limit As methylation in rice paddy soils.

ASSOCIATED CONTENT

Supporting Information

Materials and methods: MMAs(III)-associated abiotic controls; *arsM* gene primer

design; construction of *arsM* plasmid standards.

Figures: Growth curves (OD_{600}); aqueous As species; stability of MMAs(III); As

mass balance; reference genes selected by qBase plus; *arsM* gene transcription.

plasmid *ars* operon, non-normalized aqueous As species with and without

oxidation; the stability of MMAs(III) in fresh/spent RCB; reaction of MMAs(III) with

sulfide; the unknown peak (post-oxidation); As mass balance; normalized

intracellular As; reference genes selected by qBase plus software; transcriptions

of *arsM* genes and correlation analysis; chromosomally encoded *arsP* gene.

Tables: Experimental conditions; reference gene primers; raw data for all

experiments.

AUTHOR INFORMATION

Corresponding Author

Rizlan Bernier-Latmani — EPFL ENAC IIE EML CH A1 375 (Bâtiment CH) Station

6 CH-1015 Lausanne Switzerland. Phone: +41 21 69 35001; E-mail:

rizlan.bernier-latmani@epfl.ch.

Authors

Jiang-Tao Qiao — Environmental Microbiology Laboratory, School of Architecture,

Civil and Environmental Engineering, École Polytechnique Fédérale de Lausanne,

Lausanne, Switzerland

Hugo Sallet — Environmental Microbiology Laboratory, School of Architecture,

Civil and Environmental Engineering, École Polytechnique Fédérale de Lausanne,

Lausanne, Switzerland

Karin Lederballe Meibom — Environmental Microbiology Laboratory, School of

Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale

de Lausanne, Lausanne, Switzerland

Nicolas Jacquemin — Environmental Microbiology Laboratory, School of

Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale

de Lausanne, Lausanne, Switzerland.

Rizlan Bernier-Latmani — Environmental Microbiology Laboratory, School of

Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale

de Lausanne, Lausanne, Switzerland

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors would like to acknowledge the EPFL Central Environmental

Laboratory. This work was funded by the Swiss National Science Foundation

(SNSF) NCCR Microbiomes (grant # 565196).

REFERENCES

(1) Páez-Espino, D.; Tamames, J.; de Lorenzo, V.; Cánovas, D. Microbial responses to
environmental arsenic. *Biometals* **2009,** *22*, 117–130.
(2) Stolz, J. F.; Basu, P.; Santini, J. M.; Oremland, R. S. Arsenic and selenium in microbial
metabolism. *Annu. Rev. Microbiol.* **2006,** *60*, 107–130.
(3) Tamaki, S.; Frankenberger, W. T. Environmental Biochemistry of Arsenic. In *Reviews of*

Environmental Contamination and Toxicology: Continuation of Residue Reviews; Ware, G. W.,
Eds.; Springer New York: New York **1992;** pp 79–110.

(4) Zhu, Y. G.; Xue, X. M.; Kappler, A.; Rosen, B. P.; Meharg, A. A. Linking genes to microbial
biogeochemical cycling: lessons from arsenic. *Environ. Sci. Technol.* **2017,** *51*, 7326-7339.

(5) Zhu, Y. G.; Yoshinaga, M.; Zhao, F. J.; Rosen, B. P. Earth abides arsenic biotransformations.
Annu. Rev. Earth. Planet. Sci. **2014,** *42*, 443–467.

(6) Slyemi, D.; Bonnefoy, V. How prokaryotes deal with arsenic. *Environ. Microbiol. Rep.* **2012,** *4*,
571–586.

(7) Williams, P.; Price, A.; Raab, A.; Hossain, S.; Feldmann, J.; Meharg, A. A. Variation in arsenic
speciation and concentration in paddy rice related to dietary exposure. *Environ. Sci. Technol.* **2005,**
39, 5531–5540.

(8) Davis, M. A.; Signes-Pastor, A. J.; Argos, M.; Slaughter, F.; Pendergrast, C.; Punshon, T.;
Gossai, A.; Ahsan, H.; Karagas, M. R. Assessment of human dietary exposure to arsenic through
rice. *Sci. Total Environ.* **2017,** *586*, 1237–1244.

(9) Zavala, Y. J.; Gerads, R.; Gürleyük, H.; Duxbury, J. M. Arsenic in rice: II. Arsenic speciation in
USA grain and implications for human health. *Environ. Sci. Technol.* **2008,** *42*, 3861-3866.

(10) Meharg, A. A.; Williams, P. N.; Adomako, E.; Lawgali, Y. Y.; Deacon, C.; Villada, A.; Cambell,
R. C.; Sun, G. X.; Zhu, Y. G.; Feldmann, J. Geographical variation in total and inorganic arsenic
content of polished (white) rice. *Environ Sci. Technol.* **2009,** *43*, 1612–1617.

(11) Tang, Z.; Wang, Y.; Gao, A.; Ji, Y.; Yang, B.; Wang, P.; Tang, Z.; Zhao, F. J. Dimethylarsinic
acid is the causal agent inducing rice straighthead disease. *J. Exp. Bot.* **2020,** *71*, 5631-5644.

(12) Chhabra, R.; Goyal, P.; Singh, T.; Vij, L. Understanding straighthead: a complex physiological
disorder of rice (*Oryza sativa L.*). *Acta Physiol. Plant.* **2021,** *43*, 1–14.

(13) Challenger, F. Biological methylation. *Adv. Enzymol. Relat. Subj. Biochem.* **1951,** *12*, 429–491.

(14) Marapakala, K.; Packianathan, C.; Ajees, A. A.; Dheeman, D. S.; Sankaran, B.; Kandavelu, P.;
Rosen, B. P. A disulfide-bond cascade mechanism for arsenic (III) S-adenosylmethionine
methyltransferase. *Acta Crystallogr. D* **2015,** *71*, 505–515.

(15) Qin, J.; Rosen, B. P.; Zhang, Y.; Wang, G.; Franke, S.; Rensing, C. Arsenic detoxification and
evolution of trimethylarsine gas by a microbial arsenite S-adenosylmethionine methyltransferase.
Proc. Natl. Acad. Sci. U.S.A **2006,** *103*, 2075–2080.

(16) Arao, T.; Kawasaki, A.; Baba, K.; Mori, S.; Matsumoto, S. Effects of water management on
cadmium and arsenic accumulation and dimethylarsinic acid concentrations in Japanese rice.
Environ. Sci. Technol. **2009,** *43*, 9361–9367.

(17) Wang, M.; Tang, Z.; Chen, X. P.; Wang, X.; Zhou, W. X.; Tang, Z.; Zhang, J.; Zhao, F. J. Water
management impacts the soil microbial communities and total arsenic and methylated arsenicals
in rice grains. *Environ. Pollut.* **2019,** *247*, 736–744.

(18) Qin, J.; Lehr, C. R.; Yuan, C.; Le, X. C.; McDermott, T. R.; Rosen, B. P., Biotransformation of
arsenic by a Yellowstone thermoacidophilic eukaryotic alga. *Proc. Natl. Acad. Sci. U.S.A* **2009,** *106*,
(13), 5213–5217.

(19) Chen, S. C.; Sun, G. X.; Rosen, B. P.; Zhang, S. Y.; Deng, Y.; Zhu, B. K.; Rensing, C.; Zhu, Y.
G. Recurrent horizontal transfer of arsenite methyltransferase genes facilitated adaptation of life to
arsenic. *Sci. Rep.* **2017,** *7*, 1–11.

(20) Chen, S. C.; Sun, G. X.; Yan, Y.; Konstantinidis, K. T.; Zhang, S. Y.; Deng, Y.; Li, X. M.; Cui, H.
L.; Musat, F.; Popp, D. The Great Oxidation Event expanded the genetic repertoire of arsenic

metabolism and cycling. *Proc. Natl. Acad. Sci. U.S.A* **2020,** *117*, 10414–10421.

(21) Chen, J.; Rosen, B. P. The arsenic methylation cycle: how microbial communities adapted

methylarsenicals for use as weapons in the continuing war for dominance. *Front. Environ. Sci.*

2020, *8*, 43.

(22) Chen, J.; Yoshinaga, M.; Rosen, B. P. The antibiotic action of methylarsenite is an emergent

property of microbial communities. *Mol. Microbiol.* **2019,** *111*, 487–494.

(23) Viacava, K.; Qiao, J. T.; Janowczyk, A.; Poudel, S.; Jacquemin, N.; Meibom, K. L.; Shrestha,

H. K.; Reid, M. C.; Hettich, R. L.; Bernier-Latmani, R. Meta-omics-aided isolation of an elusive

anaerobic arsenic-methylating soil bacterium. *The ISME J.* **2022**, 16, 1740–1749.

(24) Ghoul, M.; Mitri, S. The ecology and evolution of microbial competition. *Trends Microbiol.*

2016, 24, 833–845.

(25) Mallon, C. A.; Poly, F.; Le Roux, X.; Marring, I.; van Elsas, J. D.; Salles, J. F. Resource pulses

can alleviate the biodiversity–invasion relationship in soil microbial communities. *Ecology* **2015**, 96,

915–926.

(26) Reid, M. C.; Maillard, J.; Bagnoud, A.; Falquet, L.; Le Vo, P.; Bernier-Latmani, R. Arsenic

methylation dynamics in a rice paddy soil anaerobic enrichment culture. *Environ. Sci. Technol.*

2017, *51*, 10546–10554.

(27) Kerl, C. F.; Schindele, R. A.; Bru ̈ ggenwirth, L.; Colina Blanco, A. E.; Rafferty, C.; Clemens,

S.; Planer-Friedrich, B. Methylated thioarsenates and monothioarsenate differ in uptake,

transformation, and contribution to total arsenic translocation in rice plants. *Environ. Sci. Technol.*

2019, *53*, 5787-5796.

(28) Scheer, J.; Findenig, S.; Goessler, W.; Francesconi, K. A.; Howard, B.; Umans, J. G.; Pollak,

J.; Tellez-Plaza, M.; Silbergeld, E. K.; Guallar, E. Arsenic species and selected metals in human

urine: validation of HPLC/ICPMS and ICPMS procedures for a long-term population-based

epidemiological study. *Anal. Methods* **2012,** *4*, 406–413.

(29) Currier, J. M.; Saunders, R. J.; Ding, L.; Bodnar, W.; Cable, P.; Matoušek, T.; Creed, J. T.;

Stýblo, M. Comparative oxidation state specific analysis of arsenic species by high-performance

liquid chromatography-inductively coupled plasma-mass spectrometry and hydride generation-

cryotrapping-atomic absorption spectrometry. *J. Anal. At. Spectrom.* **2013,** *28*, 843–852.

(30) Yathavakilla, S. K. V.; Fricke, M.; Creed, P. A.; Heitkemper, D. T.; Shockey, N. V.; Schwegel,

C.; Caruso, J. A.; Creed, J. T. Arsenic speciation and identification of monomethylarsonous acid

and monomethylthioarsonic acid in a complex matrix. *Anal. Chem.* **2008,** *80*, 775–782.

(31) Wang, J.; Kerl, C. F.; Hu, P.; Martin, M.; Mu, T.; Brüggenwirth, L.; Wu, G.; Said-Pullicino, D.;

Romani, M.; Wu, L. Thiolated arsenic species observed in rice paddy pore waters. *Nat. Geosci.*

2020, *13*, 282–287.

(32) Viacava, K.; Meibom, K. L.; Ortega, D.; Dyer, S.; Gelb, A.; Falquet, L.; Minton, N. P.; Mestrot,

A.; Bernier-Latmani, R. Variability in arsenic methylation efficiency across aerobic and anaerobic

microorganisms. *Environ. Sci. Technol.* **2020,** *54*, 14343–14351.

(33) Chen, J.; Madegowda, M.; Bhattacharjee, H.; Rosen, B. P. ArsP: a methylarsenite efflux

permease. *Mol. Microbiol.* **2015,** *98*, 625–635.

(34) Shi, K.; Li, C.; Rensing, C.; Dai, X.; Fan, X.; Wang, G. Efflux transporter ArsK is responsible

for bacterial resistance to arsenite, antimonite, trivalent roxarsone, and methylarsenite. *Appl.*

Environ. Microbiol. **2018,** *84*, e01842-18.

(35) Li, J.; Pawitwar, S. S.; Rosen, B. P. The organoarsenical biocycle and the primordial antibiotic

methylarsenite. *Metallomics* **2016,** *8*, 1047–1055.

FIGURE CAPTIONS

Figure 1. Growth curves (total protein) of *Paraclostridium bifermentans* strain EML in anoxic RCB dilutions (100%, 75%, 50%, or 25% RCB) in the presence of 25 μM As(III). Data are shown as mean values with error bars. Individual values for each biological replicate are included in Supporting Information Table S4.

Figure 2. Time-dependent concentrations of protein-normalized aqueous As species in anoxic RCB dilutions (100%, 75%, 50%, or 25% RCB) inoculated with *Paraclostridium bifermentans* strain EML and 25 μM As(III). Individual values for each biological replicate can be found in Supporting Information Table S10. (a) MMAs(III) (no oxidation), (b) MMAs(V) (post-oxidation) and (c) DMAs(V) (post-oxidation).

Figure 3. (a) Transcripts of *arsM* gene of *Paraclostridium bifermentans* strain EML in anoxic RCB dilutions (100%, 75%, 50%, or 25% RCB) in the presence and absence of 25 μM As(III) at 8 hours of incubation. (b) and (c) Correlation analysis of *arsM* gene transcripts and concentrations of MMAs(III) (no oxidation), and MMAs(V) (post-oxidation) at 8 hours of incubation. Different letters showed significant difference at $P < 0.05$. Individual values for each biological replicate are shown in Supporting Information Table S13.

559   Graphical abstract
560

563 **Figure 1.** Growth curves (total protein) of *Paraclostridium bifermentans* strain

564 EML in anaerobic RCB dilutions (100%, 75%, 50%, or 25% RCB) in the presence

565 of 25 µM As(III). Data are shown as mean values with error bars. Individual values

566 for each biological replicate are included in Supporting Information Table S4.

Figure 2. Time-dependent concentrations of protein-normalized aqueous As species in anaerobic RCB dilutions (100%, 75%, 50%, or 25% RCB) inoculated with *Paraclostridium bifermentans* strain EML and 25 µM As(III). Individual values for each biological replicate can be found in Supporting Information Table S10. (a) MMAs(III) (no oxidation), (b) MMAs(V) (post-oxidation) and (c) DMAs(V) (post-oxidation).

Figure 3. (a) Transcripts of *arsM* gene of *Paraclostridium bifermentans* strain EML in anaerobic RCB dilutions (100%, 75%, 50%, or 25% RCB) in the presence and absence of 25 μM As(III) at 8 hours of incubation. (b) and (c) Correlation analysis of *arsM* gene transcripts and concentrations of MMAs(III) (no oxidation), and MMAs(V) (post-oxidation) at 8 hours of incubation. Different letters showed significant difference at $P < 0.05$. Individual values for each biological replicate are shown in Supporting Information Table S13.