Title: Terrestrial endotherms trade body temperature regulation for the stress response

Authors: Joshua K.R. Tabh[1]*, Mariah Hartjes[2], Gary Burness[2]

Affiliations: [1] Environmental and Life Sciences Graduate Program, Trent University, Peterborough, Ontario, Canada, K9L 0G2

[3] Department of Biology, Trent University, Peterborough, Ontario, Canada, K9L 0G2

Corresponding author

Contact: joshua.k.tabh@gmail.com

Data Availability: All data used for the construction of this study have been submitted alongside the below manuscript. These data will be made available to the public upon publication at: https://github.com/joshuakrobertson/.

Author Contributions: JT and GB conceived the study. MH and JT collected the data, and JT conducted statistical analyses and wrote the first draft of the manuscript. GB contributed to manuscript revisions.

Conflicts of Interest: The authors have no conflicts of interest to declare.

Abstract

In response to stressors, many species limit reproduction, reduce investment in immunity, and even forgo somatic repair to free energy for use toward coping. Often, however, the energetic costs of simply regulating body temperature can exceed those required for each of these processes. Nevertheless, whether thermoregulation is also eschewed in response to stressors is not known. In this study, we used a phylogenetically-controlled meta-regression to test whether changes in body temperature that accompany the stress response – a common phenomenon among terrestrial vertebrates – represent the outcome of an energy allocation trade-off between thermoregulation and the stress response in these animals. Our results, obtained from 65 studies and across 24 species, support this trade-off hypothesis by showing that individuals with high costs of thermoregulation and little energy available for use (i.e those with low body mass and high resting metabolism respectively) tended to become more heterothermic in response to stressors, while those with lower costs of thermoregulation and greater energy availlable for use tended to increase their body temperatures regardless of the ambient environment. Allowing body temperatures to change in these way liberated an average of 5%, and up to 23%, of resting energy expenditure for use toward the stress response, regardless of species. These cross-species findings indicate that terrestrial endotherms trade investment toward thermoregulation for that toward the stress response. With such findings in mind, evaluating how these endotherms may cope with warming climates and concurrent stressors should be key priority in the coming decade.

Main

About 2000 years ago, the Greek physician Galen observed that the body temperatures of his patients tended to rise after psychologically distressing experiences[1]. Given the acute nature of these rises, Galen termed their occurrence "ephemeral fevers"[2] and speculated, like those after him (e.g. Ibn Sina) that they emerged from an abundance of humor[3,4]. "Why" (i.e., functionally) they might emerge at all, however, was largely left unquestioned.

By now, numerous studies have shown that Galen's "ephemeral fever" is not just restricted to humans. Instead, many rodent, non-hominid primate, and even bird species appear to change their body temperatures after exposure to common disturbances, such as transportation or restraint[5,6,7]. Intriguingly, modern evidence also suggests that the term "ephemeral fever" may even be a misnomer, with body temperatures of some individuals sometimes *declining* in response to stressors rather than rising[8,9]. That these stress-induced thermal responses exist across varied taxonomic groups and may differ under certain conditions strongly intimates a possible function, or, perhaps, a conserved and inherent nature (i.e., a constraint) behind their occurrence.

Recently, we proposed that changes in body temperature following stress exposure may be understood as consequences of an allocation trade-off between thermoregulation and the stress response[10,11] (similar to Lewden et al[12]). Under this hypothesis, fluctuations in body temperature that accompany the stress response represent the outcomes of reducing energy expenditure toward thermoregulation in favour of expenditure

toward coping. Unlike previous hypotheses, this particular hypothesis

provides an explanation for why body temperature might either rise *or*

decline after a stress exposure, since relaxing thermoregulation should

merely result in body temperature drifting toward ambient, barring

accumulation of metabolic heat from activation of the stress response. Over

the past few decades, a handful of studies have indeed supported a role of

ambient temperature in shaping the direction and magnitude of stress-

induced thermal responses[13-15]. These studies provide intriguing, albeit

sometimes conflicting, support for our trade-off hypothesis. Nevertheless,

whether an influence of ambient temperature holds across studies, species,

and contexts (i.e. stressors), remains unknown.

In this study, we used data describing body temperature responses to

stressors across terrestrial endotherms to test for an allocation trade-off

between thermoregulation and the stress response (outlined in Fig. 1). Non-

terrestrial endotherms were excluded from our study owing to a lack of

available data, but implications with respect to these species are discussed.

Under our trade-off hypothesis, we made three predictions. First, although

body temperatures should typically increase after a stressor owing to

accumulation of metabolic heat, these increases should be exacerbated in

the warmth and forgone, or even reversed, in the cold. Second, the degree

to which ambient temperature influences stress-induced thermal responses

should depend on body mass, with smaller species/individuals becoming

more heterothermic across observed temperature ranges, owing to higher

surface area to volume ratios and greater susceptibilities to the cold (i.e.

higher lower critical temperatures). Third, how stress-induced thermal

responses change across ambient temperature should depend on the

relative amount of energy available for allocation toward both

thermoregulation and the stress response (as per Stearns[16]). More

specifically, species with relatively high metabolic rates (thus indicating low

energy availability; see Careau[17]) should be more likely to decrease their

body temperature in the cold, and increase their body temperature in the
warmth following a stressor than those with relatively low metabolic rates
(indicating high energy availability).

Results

88 Where provided, positive log-response ratios represent a stress-induced
89 increase in body temperature, while negative log-reponse ratios represent a
90 stress-induced decrease.

Changes in body temperature that follow a stressor depend on air temperature and body mass

Across all species studied, body temperatures increased by an average of
0.71°C in response to stressors (indicated by our model intercept; $\beta \approx$
1.83×10^{-2} [80%: 5.02×10^{-3}, 2.80×10^{-2}; 95%: -1.61×10^{-3}, 3.55×10^{-2}]). As
predicted by our trade-off hypothesis, these stress-induced reseponses were
clearly influenced by ambient temperature in both their magnitudes and
directions ($\beta \approx 7.50 \times 10^{-4}$ [80%: 2.80×10^{-4}, 1.20×10^{-3}; 95%: 3.00×10^{-5},
1.46×10^{-3}]; Fig. 3); at low ambient temperatures, where costs of warming
should be highest, body temperature was more likely to decrease after a
stressor than at high ambient temperatures. By contast, at high ambient
temperatures, where costs of cooling should be highest, body temperature
was more likely to increase after a stressor, and to a greater degree, than at
low ambient temperatures (Fig. 3).
Critically, an influence of ambient temperature on stress-induced
responses was not ubiquitous, and clearly depended on an individual's body
mass (interaction between air temperature and body mass: $\beta = -4.80 \times 10^{-4}$
[80%: -6.60×10^{-4}, -1.40×10^{-4}; 95%: -8.10×10^{-4}, -1.00×10^{-5}]; Fig. 4a). Given
that body mass is capable of dictating both costs of thermoregulation and
susceptibilies to heat and cold among animals (i.e. their zones of
thermoneutrality), this interactive effect between it and ambient

temperature was also expected under a trade-off between thermoregulation
and the stress response. Among individuals with relatively low body masses
(< 222 g, our sample mean), exposure to a stressor tended to induce a state
of heterothermy, with body temperatures decreasing after a stressor in cool
temperatures and increasing after a stressor in the warmth (Fig. 4a). At
ambient temperatures below our sample mean (20.1°C), for example, body
temperatures of these individuals decrease by an average of 0.51°C after a
stressor, yet at ambient temperatures above our sample mean, body
temperatures increased by an average of 0.82°C (Fig. 4b). In direct
contrast, among individuals with relatively high body masses (> 222 g),
body temperature tended to increase after a stressor regardless of the
current ambient temperature (Fig. 4; mean change in body temperature =
+2.10°C at below-average air temperatures, and +1.18°C at above-average
air temperatures). Regardless of body mass and ambient temperature, our
biophysical models predicted that how body temperatures responded to
stressors across these animals reduced energetic costs of maintenance by
on average of 4.6%, and up to 23% of resting metabolic rate (Fig. 4c).

Independent of air temperature, larger animals were also marginally
more likely to increase their body temperature after a stress exposure, and
by a greater degree, than smaller animals (main effect of body mass: β =
4.00×10^{-3} [80%: 9.70×10^{-4}, 7.59×10^{-3}; 95%: -1.72×10^{-3}, 8.98×10^{-3}]; Fig. 3).

Effects of air temperature on thermal responses to stressors are
linked to relative energy expenditure at rest

Overall, body temperatures tended to increase after a stress exposure at
moderate-to-high ambient temperatures (indicated by a positive log
response ratio at ambient temperatures > approximately 10°C), regardless
of whether a species expended relatively low, or relatively high amounts of
energy at rest (i.e. for their given mass; Fig. 5). The magnitude by which
these increases occurred also did not depend upon a species' relative
energy expenditure at rest (main effect of relative resting metabolic rate: β

$\approx$ -3.96×10^{-3} [80%: -1.28×10^{-2}, 4.78×10^{-3}; 95%: -1.83×10^{-2}, 8.56×10^{-3}]; Fig.

3). In the cold (ambient temperatures < approximately 10°C), however,

species expending relatively high amounts of energy at rest (and thus,

having less energy available for allocation elsewhere) tended to decrease

their body temperatures, or allow them to fall, relative to those expending

comparably little energy at rest, as predicted (interaction between air

temperature and relative energy expenditure: β = 1.12×10^{-3} [80%: 2.40×10^{-4},

2.24×10^{-3}; 95%: -3.50×10^{-4}, 2.70×10^{-3}]; Fig. 5a). Since baseline body

temperatures did not differ between species with relatively high, and

relative low energy expenditure, this interactive effect with ambient

temperature could not be explained by body temperatures being near

maximum or minimum potential values alone (e.g. "ceiling" or "floor"

effects; mean baseline body temperature ± s.d.: low energy expenditure =

37.5°C ± 1.71°C; high energy expenditure = 38.2°C ± 1.77°C).

Across all studies where ambient temperatures were below our

sample average (20.1°C), body temperatures of those with relatively high

resting energy expenditure decreased by an average of 0.32°C in response

to a stressor (Fig 5b). Among those with relatively low resting energy

expenditure, however, body temperatures increase by an average of 0.73°C

across the same ambient temperature range (Fig. 5b). In both cases, these

responses to stressors probably contributed to a 5% reduction (mean =

5.45%) in maintenance expenditure (Fig. 5c).

Time, age class, and species identity influence how body

temperatures respond to stress exposures

Beyond air temperature, body mass, and relative energy expenditure, study

methodology had mixed effects on the outcomes of stress-induced thermal

responses. Alone, latency between baseline and stress-induced body

temperature measurements clearly influenced the observed magnitude of

stress-induced thermal responses (first order: β = -7.45×10^{-2} [80%: -

8.38×10^{-2}, -2.00×10^{-2}; 95%: -1.00×10^{-1}, -1.49×10^{-3}]; second order: β = -

4.18×10^{-2} [80%: -7.53×10^{-2}, -1.89×10^{-2}; 95%: -8.79×10^{-2}, -1.16×10^{-3}]; Figs. 3

& 6). Across all studies assessed here, and when holding all other

parameters constant, body temperatures tended to increase immediately

after stress exposure, and then decrease after an average of approximately

35 minutes (2100 seconds; Fig. 6). Whether body temperature was

measured using implanted telemeters or thermocouples, however, did not

appear to influence how body temperatures changed in response to

stressors (β = -4.50×10^{-4} [80%: -5.50×10^{-3}, 9.14×10^{-3}; 95%: -9.84×10^{-3},

1.32×10^{-2}]; Fig. 3).

Although not predicted here, body temperatures were more likely to

decrease after a stress exposure among study populations comprised of

solely adults than among those comprised of solely juveniles (β = 5.78×10^{-3}

[80%: -1.28×10^{-2}, -1.38×10^{-3}; 95%: -1.64×10^{-3}, 9.70×10^{-4}]; Fig. 3). How body

temperature responded to stress exposures was also weakly genetically

linked, with species identity and phylogeny explaining a small degree of

variation in our data (random intercept for species: μ = 2.24×10^{-2} [80%:

7.86×10^{-3}, 3.01×10^{-2}; 95%: 6.10×10^{-4}, 3.50×10^{-2}]; random intercept for

phylogeny: μ = 9.30×10^{-4} [80%: 1.00×10^{-5}, 2.35×10^{-3}; 95%: 0, 3.38×10^{-3}];

Fig. 3).

Discussion

Costs of body temperature regulation dictate body temperature
under stress

Changes in body temperature that accompany the stress response are often

understood as byproducts of the physiological stress response and serve no

function in and of themselves[18-20]. Our results do not support this

understanding. Instead, we provide evidence that stress-induced changes in

body temperature result from attempts to balance energetic expenditure

between body temperature regulation and the stress response. More

specifically, stress-induced thermal responses represent allocation trade-
offs between thermoregulation and stress physiology[10,11].

Supporting our trade-off hypothesis, our findings indicate that, among
terrestrial endotherms, how body temperature changes after a stress
exposure depends on how much energy a species expends on body
temperature regulation (as influenced by their body mass, and thus, their
surface area to volume ratio). In cool environments, for example, species
predicted to have relatively high expenditure toward warming (i.e. those
with low body mass, and thus, high surface area to volume ratios) tended to
decrease their body temperatures, or allow them to fall, following a stress
exposure, while those with relatively low expenditure toward warming, or
even slight expenditure toward cooling (i.e. those with large body masses,
and thus, low surface area to volume ratios) did not (Fig. 4). Under a trade-
off hypothesis, we predicted this to be the case because expenditure toward
the stress response should only necessitate reductions in expenditure
toward thermoregulation when costs of warming or cooling are already
high. By contrast, in the absence of a trade-off, expenditure toward
thermoregulation should be irrelevant to the direction or magnitude of
stress-induced thermal responses.

An interactive effect of ambient temperature and body mass on the
outcome of stress-induced thermal responses is not new to physiological
theory. In birds, Nord and Folkow[21] highlighted that hypothermic responses
to stress exposure seem more likely to occur at low ambient temperatures,
but specifically among species with small body size (e.g. Parids[12,22]). Oka[23]
alluded to a similar trend in small mammals (e.g. rats and domestic
rabbits[13,14,24]). For Nord and Folkow[21], this interaction between body mass
and ambient temperature was interpreted as a byproduct of experimental
methodology, with commonly-used restraint stressors causing greater
disruptions to the retention of body heat in small birds relative to large
birds (i.e. by compression of plumage[21]). Although this interpretation may

be valid in some contexts, it alone does not seem sufficient to explain our
general findings. Numerous studies included in our analysis (n = 28),
employed stressors that did not involve direct contact or handling of
experimental animals at all (e.g. social isolation, exposure to a dominant
individual, introduction to a novel environment etc). Yet despite inclusion of
these studies in our analysis, we still detected a strong interactive effect of
body mass and ambient temperature on the outcome of stress-induced
thermal responses (Fig. 4). Thus, interpreting these responses as passive,
unregulated consequences of experimental design is likely premature.

Under a trade-off hypothesis, we speculated that body mass
influenced stress-induced thermal responses through its effects on heat loss
and, subsequently, energetic demands for heat production (discussed
above). Nevertheless, it remains possible that how body mass influences
thermal responses to stressors occurs though other avenues, for example,
by influencing thermal inertia. Arguably the most direct way to test our
speculation would be to compare how body temperature responds to
stressors at ambient temperatures below, within, and above the zone of
thermoneutrality within a given species, where relative expenditure toward
thermoregulation differs predictably. To the best of our knowledge, very few
studies have sought to conduct this type of comparison and outcomes of
those that have are equivocal[13,14,25].

Relative energy availability also shapes body temperature under stress

Above and beyond body mass, our results indicate that relative energy
expenditure at rest influences how body temperature responds to stressors
in different thermal environment. Specifically, we found that species with
relatively high rates of energy expenditure at rest -- whose energy available
to allocate to both the stress response and thermoregulation should be
limited (see Careau[17]) -- tended to lower their body temperatures after
stressors in the cold, but raise them in the warmth (Fig. 5). By contrast,

species with relatively low rates of energy expenditure at rest -- whose
energy available for allocation toward the stress response and
thermoregulation should abound -- tended to raise their body temperatures
after stressors regardless of the thermal environment (Fig. 5). According to
our trade-off hypothesis, we predicted this combined effect of energy
availability and the thermal environment on stress-induced thermal
responses, since adjustments of expenditure toward body temperature
regulation should predominantly occur when energy is limited[16].

Supporting our findings, food restriction has been shown to reverse
thermal responses to stressors in adults rats, with fasted individuals
decreasing their body temperatures in response to restraint, and fed
individuals increasing them in equivalent conditions[26]. Critically, rats in this
study were held at temperatures slightly below thermoneutrality[27], where
allowing body temperature to decrease in fasted conditions would relax
energetic demands of warming, thus freeing energy for use in the stress
response. Similar trends have also been shown in birds, with socially-
mediated food restriction (i.e. by social subordination) stimulating stress-
induced hypothermia in the cold and hyperthermia in the warmth (core
body temperature proxied by eye region temperature[11]). Collectively, these
findings support that how body temperature changes after a stressor
probably depends on an individual's relative energy balance, and may well
represent active calibration of expenditure toward warming or cooling.

Not surprisingly, our results reiterate a key effect of time on the
magnitude and direction of stress-induced thermal responses. When
controlling for all other parameters, body temperature was more likely to
increase shortly after stress exposure, then decrease to baseline
temperatures, or below, shortly thereafter (Fig. 6). These trends strongly
align with those reported for individual species[28-30] and probably represent
the timeline of physiologically responding to, and recovering from, a
stressor. Perhaps more interestingly, our results also indicated that age

likely influences how body temperature responds to a stressor. When
averaged across species and thermal environments, we found that younger
individuals were more likely to increase their body temperatures after a
stressor, and to a greater extent, than older individuals (refer to negative
slope of age class in Fig. 3). Given that juveniles are expected to allocate
high amounts of energy toward growth[31,32], this result seems surprising.
Although a similar effect of age has been reported in a handful of other
studies[26,33-35], why these differences between juveniles and adults occur is
not known. One possible explanation is that juveniles are comparatively less
averse to wear-and-tear associated with homeostatic overload, once
expenditure toward growth, thermoregulation, and the stress response are
combined (i.e., a "better safe than sorry" strategy[35,36]). Alternatively, the
capacity to change body temperature after a stressor may simply differ
between age groups owing to age-related changes to nervous system
function (reviewed by Oka[23]). In any case, understanding whether these
age-related effects have meaningful implications on energy balance is yet
necessary to determine whether they support or refute our trade-off
hypothesis.

Putative mechanisms

Contextualising stress-induced changes in body temperature as
allocation trade-offs (i.e. between thermoregulation and the stress
response) may suggest that body temperature set-point is actively changed
when a non-thermal stressor is perceived. Thermoregulatory control centres
should therefore be involved in mediating these stress-induced responses.
So far, studies using pharmacological agents to manipulate neural function
in these control centres suggest that they probably does regulate body
temperature responses to stressors[37-39], at least indirectly. Additionally,
their capacity to shape stress-induced thermal responses is also becoming
increasingly clear[40,41]. In an effort to synthesise these findings, Angilletta et
al[42] recently proposed that inhibitory pathways linking the limbic system

with thermoregulatory nuclei of the hypothalamus may be activated during
the stress response, thus temporarily knocking down select
thermoregulatory processes. According to their model, activation of, and
investment in, the acute stress response is therefore actively prioritised
over thermoregulation (representing an "active" allocation trade-off).
Efforts to map inhibitory pathways between limbic structures and
thermoregulatory control centres are limited but generally confirmatory[43,44],
thus providing some early mechanistic support for our trade-off hypothesis.

While changes to body temperature that accompany the stress
response may well represent "active" trade-offs, we still cannot rule out the
possibility that they merely represent passive allocation constraints.
Namely, declines in body temperature in the cold and increases in body
temperature in the warmth might simply indicate a failing to allocate
energy toward responding to the thermal environment experienced during a
stress exposure. In this context, thermal responses to stressors may not
constitute functional responses to balance energetic expenditure between
thermoregulation and the stress response, but rather, may indicate an
inability to regulate body temperature at all under distress (see, for
example, Andreasson et al[35]). Whether these stress-induced changes in body
temperature do indeed represent regulated trade-offs or passive constraints
is an important question that will directly influence whether these
responses may respond to selection or are simply fixed consequences of
how physiological responses to stressors shape body temperature.

Ectotherms and non-terrestrial endotherms

Descriptions of how body temperatures change after stressors among
ectotherms and non-terrestrial endotherms are sparse. We were therefore
unable to formally evaluate whether evidence for an allocation trade-off
between thermoregulation and the stress response extends to these species'
groupings. However, indirect evidence for such a trade-off in ectotherms is

widely supportive. Some species of lizards, grasshoppers, and even larval
newts, for example, reportedly abandon basking and/or heat-seeking
behaviour in the presence of predator cues (reviewed in Angiletta et al[42]). In
the Yarrow's spiny lizard (*Sceloporus jarrovii*), this change in behaviour
results in body temperature both declining toward ambient and away from
preference[45]. While not strictly evidencing an energy allocation trade-off,
these reports do highlight that responding to stressors can, and often does,
take precedence over body temperature regulation in ectotherms with clear
consequences for the latter. Among non-terrestrial endotherms,
thermoregulation may also be relaxed, or even forgone, in the presence of
stressors. In southern bluefin tuna (*Thunnus maccoyii*), for example, body
temperatures can rise dramatically in response to suspected predators[46],
which, in severe cases, is thought to contribute to physiological damage[47] (a
phenomenon known as "burnt meat"). We therefore speculate that our
support for a trade-off between coping was a stressor and regulating body
temperature probably extends beyond the species grouping analysed here.

Conclusions

Together, our results provide strong evidence that changes in body
temperature accompanying the stress response represent allocation trade-
offs between the stress-response and thermoregulation. Across species, we
highlight that by allowing body temperature to drift toward ambient, these
responses contribute to reductions in the energetic costs of
thermoregulation, particularly when such are already high, or when energy
available for allocation to both the stress response and body temperature
regulation are low. These findings, therefore, highlight a new model for
understanding both how and "why" stress-induced changes in body
temperature might occur (outlined in Fig. 1). In doing so, they provide
novel, cross-species evidence that the acute stress response may directly
influence body temperature regulation in endotherms at. Given the clear
implication of these findings for a warming world, further experimental

studies testing our trade-off hypothesis within species should be a priority
in coming years.

Methods

Literature search

We searched primary literature for data describing the magnitude and
context of stress-induced changes in body temperature across endotherms
using Web of Science™. To simplify interpretations of our findings, we
considered only literature pertaining to terrestrial endotherms, excluding
humans (owing to confounding effects of clothing on costs of
thermoregulation).

Query terms for our search were: "stress", "body temperature" and
"NOT heat stress". In total, 3022 studies were returned by our query. From
those returned, we scanned abstracts for evidence that: (1) body
temperature responses to stress exposure were indeed measured, and (2)
the species in which body temperature responses were measured was
endothermic, terrestrial, and non-human; those with abstracts lacking such
evidence were removed from consideration. Next, because we were most
interested in the mechanisms driving thermal responses to stress exposure
at the body core, all studies describing stress-induced responses at surface
body tissues were also omitted. From all remaining studies (Fig. 2), we
scanned reference lists for relevant literature that had not been returned by
Web of Science. Again, abstracts of these studies were reviewed for
evidence that stress-induced changes in core body temperature were indeed
measured and were retained or excluded accordingly (n = 31 studies added;
Fig. 2). Together, all searches and data collection were conducted between
28 April and 27 July, 2020, with publication dates in our data-set ranging
from March, 1958[48] to January, 2020[49] (mean publication year [rounded to
nearest integer] ± sem = 2003 ± 0.879).

Data extraction and inclusion criteria

From each study identified for possible inclusion (n = 136; Fig. 1), we harvested the following data:

(1) The average stress-induced change in body temperature across a sample population (Δ°C; ± s.d. or s.e.m). In some studies, stress-induced thermal responses were monitored and reported across multiple study groups (i.e. per sex, air temperature treatment, discrete time period, etc.). In these cases, we chose to treat data from each group as an independent observation nested within a study. If multiple measurements of change were reported for a given study group, we collected only the value demonstrating the greatest increase or greatest decrease relative to baseline body temperature for that group.

(2) The number of individuals measured.

(3) The study species and family.

(4) The age class of the sample population (factorial; adult, juvenile, or mixed, according to age of reproductive maturity). If age class was not reported, we assumed that the study population was comprised solely of adults.

(5) Sex of individuals in sample population (factorial; female, male, or mixed). If sex was not reported, we assumed it to be mixed.

(6) The season of study, categorised as spring, summer, fall, or winter. If the season of study was not provided, we set it as "not available (NA)".

(7) The average mass of individuals in the study population (grams, ± s.d. or s.e.m; s.e.m. ≈ 0 [0.0001] if missing). If a range of masses was provided (i.e. in n = 220 observations across n = 84 studies), we

calculated the theoretical median and used it in place of average
mass, with the standard deviation equaling the mass range divided by
4. Where average mass of individuals was not reported, we extracted
mean values for the species, sex, and age-class of interest (if provided;
see categories 4-5 above) from primary literature (n = 285
observations across n = 43 species groupings and n = 117 studies).
Categories (4) and (5) describe our approach when data on sex and
age-class were missing.

(8) Air temperature during study (in °C, ± s.d. or s.e.m). For field
studies lacking measures of air temperature, we obtained average
values from nearby weather-stations (< 50 km from the study
location) across the reported duration of study (i.e. according to
specified date and time ranges) where available (n = 13 observations
from n = 3 studies; indicated in data files). Studies reporting stress-
induced thermal responses in media other than air were discarded
and those that did not report variation around air temperature means
were assigned an s.e.m. of ≈ 0 (0.0001; n = 57 observations across n
= 28 studies).

(9) Latency between pre-stress exposure and post-stress exposure
temperature measurements (seconds). Latency values greater than 8
hours (28800 seconds; > 3 s.d. above the mean [3544 seconds]) were
excluded from our analyses (n = 4 observations).

(10) Technique used to measure body temperature (intraperitoneal
telemetry or thermocouple [rectal/cloacal/throat]).

(11) Whether pharmaceuticals other than saline were administered to
study animals before or during stress exposure (binomial; true/false).

For most studies, we extracted these data from the article text or from
article figures using the open-source software FIJI[50]. For all remaining

studies, data were requested from article authors directly. Once data
collection was complete, all studies lacking data from categories (1)-(4) and
(8)-(10), and/or satisfying category (11) were discarded. Together, our final
sample included 165 observations across 68 studies, 24 species, and 30
species strains (n = 12 avian; n = 18 mammalian; average number of
observations/species ± s.d. = 6.98 ± 19.00; phylogeny provided in
supplemental information). Final samples sizes and summaries of sample
exclusions following the preferred reporting items for systematic reviews
and meta-analyses (PRISMA) are outlined in Fig. 2.
Types of stressors applied across our sample were varied and
numerous, ranging from transportation (n = 4 studies), social defeat (n =
10), restraint (mechanically [n = 31 studies] or by handling [n = 43
studies]), exposure to a novel environment (n = 32 studies), and others.

Metabolic rate data

We were interested in testing whether a species' relative energetic
expenditure at rest (and thus, their remaining energy available for
allocation) influences the magnitude of its stress-induced change in body
temperature. To calculate relative energetic expenditure for species or
species strains included in our data set (in publications that met our
inclusion criteria: see "Data extraction and inclusion criteria" above), we
again used Web of Science™ to search for primary literature reporting
either their basal or resting metabolic rates ("BMR" and "RMR"
respectively). Measures of metabolic rate within each study were
categorised as "basal" or "resting" according to whether they satisfied
criteria 1-3 or 1-2 from McKechnie and Wolf[51] respectively; those that did
not satisfy criteria 1-2, at minimum, were discarded. To enhance our search
for metabolic data, we replicated our queries using Google Scholar™.

Since both BMR and RMR likely vary across seasons, we sought to
obtain metabolic data relevant to the season at which stress-induced

changes in body temperature were observed (again, see "Data extraction
and inclusion criteria" above). In most cases, however, the season in which
metabolic expenditure and/or the stress-induced thermal response were
measured were not reported (n = 66 studies of 68). Moreover, to our
knowledge, some species lacked any primary data on basal or resting
metabolism at all. Thus, where necessary, measures of BMR and/or RMR
were obtained from available, but potentially mismatched, seasonal studies,
from the most closely related species studied (n = 5 species; relatedness
determined using a consensus phylogeny approach described below), or
from order-specific allometric relationships (n = 2 species; 1 chiropteran
species and 1 passerine species). Following collection, all metabolic data
were converted to a common metric (watts, "W") using conversion
coefficients described in the appendix.

To account for effects of body mass on energy expenditure across
species, we obtained ordinary residuals from a Bayesian linear mixed effects
model (LMM) predicting natural log-transformed BMR or RMR as a function
of natural log-transformed species mass (mean in g; prior = $\mathcal{N}[0.67, 0.1]$).
Type of measurement was included in this model as a group-level intercept
(i.e. "basal" or "resting"; group-level predictor; prior = $\Gamma[1, 1]$; model
coefficients: Intercept = -3.527 [-4.508, -1.873]; log mass = 0.668 [0.599,
0.734]; measurement type = 0.084 [-0.950, 0.869]) but phylogeny was not
included to avoid redundancy in our subsequent analyses. Residuals from
our model were then categorised as either positive or negative, with
positive values for a given species indicating a relatively high energy
expenditure at rest, and negative values indicating a relatively low energy
expenditure at rest. Categorising data in this way allowed us to broadly and
conservatively evaluate how relatively energy expenditure, and thus, energy
availablily, influenced stress-induced thermal responses while
circumventing over-parameterisation of our statistical model (described
below).

Effect size calculation

To ensure that our data were statistically comparable among studies and species, we converted our estimates of absolute changes in body temperature to relative changes in body temperature. This was achieved by calculating a response ratio as follows:

$$(1) \quad \frac{T_b 1_i}{T_b 0_i}$$

where $T_b 1$ represents an average minimal, or an average maximal body temperature of study population i that was observed after a stress exposure, and $T_b 0$ represents the body temperature of study population i before a stress exposure treatment. For studies that also recorded the body temperatures of control (or "unstressed") individuals throughout experimentation, however, we used the following modified equation:

$$(2) \quad \frac{T_b 1_{Si}}{T_b 1_{Ci}}$$

where $T_b 1_S$ again represents an average minimal, or an average maximal stress-induced body temperature of the stress-exposed population i, and $T_b 1_C$ represents the average body temperature of the control population at the equivalent time point.

To estimate uncertainty around our response ratio values, we calculated a pooled standard error using one of the following equation:

$$(3) \quad \sqrt{\frac{\sigma_{T_b' 1}}{T_b 1} + \frac{\sigma_{T_b' 0}}{T_b 0}}$$

or:

$$(4) \quad \sqrt{\frac{\sigma_{T_b' 1_S}}{T_b 1_S} + \frac{\sigma_{T_b' 1_C}}{T_b 1_C}}$$

where equation (3) applies to studies without a control population, equation
(4) applies to studies with a control population. In these equations, σ_x
represents the standard error of a given subscripted variable x, and all
other variables remain as previously described. In studies where a standard
error around the stress-induced change in body temperature was reported,
we assumed it to be a pooled estimate and used it in place of that derived
from an equation above.

Finally, to both normalise our response ratio estimates and centre them on
zero, we calculated their natural logarithms and used these transformed
values as our final estimates of effect size (i.e. the log-response ratio, or
"lRR"). Effect sizes for studies with and without control populations were
visually compared to confirm that their distributions were similar.

Phylogeny

To estimate phylogenetic relationships among species included in our data-
set, we extracted two sets of 1000 trimmed phylogenetic trees – one for
mammals, and one for birds – from VertLife[52] (Upham et al. 2019) and
BirdTree[50] respectively. These tree-sets were drawn from pseudo-posterior
distributions of relaxed-clock models to infer phylogeny from genetic and
fossil data (number of genes: mammals = 31[52]; birds = 15[53]; avian fossil
backbone from Ericson et al[54]). From each tree-set, we then estimated class-
specific consensus trees using "TreeAnnotator" in Beast 2[55], with a 25%
burn-in, posterior probability limit of 1.0, and with node heights
representing median values. Once our consensus trees were established, we
used them to build variance-covariance matrices approximating the relative
relatedness between species (via the R package "ape"[56]; matrices used in
subsequent analyses). For simplicity, relatedness between all pairs of birds
and mammals was assumed to be negligible and set to zero. Phylogeny
estimation and consensus trees are detailed in the appendix (section 5).

Statistical analyses

All of our statistical analyses were conducted in R version 4.03[57] and figures were produced using the R package "ggplot2"[58]. For each model, we used Hamiltonian Monte Carlo (HMC) sampling in the R package "brms"[59] with 2500 burn-in iterations, 15000 sampling iterations, and a thinning factor of 10 across four HMC chains. Models were validated by visually diagnosing residual distributions and paired posterior coefficient estimates. Gelman-Rubin statistics ($\hat{R}$) for each of our models fell between 0.95 and 1.05, suggesting that chain mixing was consistently adequate, and effective sample size to sample size ratios consistently exceeded 0.75, confirming minimal autocorrelation between HMC chain draws. Versions for all R packages used are reported in the appendix.

To confirm that our results were not biased by small sample sizes among our collated studies, we re-ran of our model using an adjusted log response ratio designed to correct for sample-size related error ("RR$^{\Delta}$"[60]). Results from this model were then compared with our original model at 80% and 95% highest density credible intervals (HDIs). Outcomes of both models were highly similar (see supplemental information; STable 24), suggesting that sample variation in sample size among our collected studies was unlikely to influence our reported findings.

Modeling

We were interested in testing whether the direction and magnitude of stress-induced changes in body temperature were influenced by: (1) ambient temperature, (2) the predicted energetic costs of thermoregulation for a given species and context, and (3) the relative energy available to a species for investment in both thermoregulation and the stress response (here, residual RMR after controlling from body mass, categorised as positive or negative). Clearly, costs of thermoregulation should vary according to air temperature, with the degree of variation being loosely

influenced by a species' body mass. Further, the relative energy available to
a species should be inversely proportional to it's relative expenditure (i.e. at
rest). Accordingly, we tested whether air temperature, body mass, relative
energy expenditure, and the interactions between both air temperature and
body mass, and air temperature and relative energy expenditure predicted
the direction and magnitude of stress-induced changes in core body
temperature. To achieve these ends, we constructed a Bayesian linear
mixed effects model (LMM) with our log response ratio as the response
variable, and air temperature (°C), the natural log of body mass (g), relative
energy expenditure (factorial; positive or negative, indicating high or low
expenditure for a given body mass respectively), the linear interaction
between air temperature and body mass, and the linear interaction between
air temperature and relative energy expenditure as population-level
predictors. Given that both body mass and air temperature can vary
considerably within studies, however, we modeled each parameter as latent,
with their true values being unknown but falling within our observed values
± our observed error. Finally, both air temperature and the log of body mass
were mean-centred to better facilitate interpretation of our model intercept.

To account for recovery or ceiling/floor effects in our analysis, we
included the latency between baseline and stress-induced body temperature
measurements as a population-level predictor in our LMM. Here, we
assumed that the temporal effects of stress exposure on body temperature
were non-linear, with body temperature rapidly changing, then plateauing
or recovering across time[28-30]. For this reason, the latency between baseline
and stress-exposed measurements was modeled as a second-order
polynomial. Similar to air temperature and the log of body mass, latency
between measurements was mean-centred to ease interpretation of our
intercept. To correct for an influence of the method used to measure body
temperature on stress-induced thermal responses, and to correct for the
possibility of age effects, we included measurement technique (factorial;

telemetry or thermocouple) and age of individuals in the study population
(integer; juveniles only = -1, both juveniles and adults = 0, adults only = 1)
in our model as population-level predictors. Finally, species, study identity,
and phylogeny (see section 6) were included as group-level predictors to
control for non-independence between each factor.

For our LMM, we used weakly informative priors that assumed that
log-responses ratios of ⪅ -1 and ⪆ 0.5 (e.g. representing a 24°C decrease
and 10°C increase in body temperature respectively at baseline
temperatures of 38°C) were unlikely. Specifically, priors for air
temperature, log body mass, and the first-order effects of latency to body
temperature measurement were set as $\mathcal{N}(0, 0.1)$, while those for the
interaction between log body mass and air temperature, and the interaction
between relative energy expenditure and air temperature were set as $\mathcal{N}(0,$
$0.05)$. Because we were less certain about the range of possible age-related
and second-order time-related effects, priors for age class and the second-
order effect of time were set as $\mathcal{N}(0, 0.25)$. Next, because both relative
energy expenditure and measurement methodology were factorial, priors
for these parameters was set as skew-normal ($\mathcal{SN}$) with ξ= 0, ω= 0.1, and α
= -2.5. We chose to use a $\mathcal{SN}$ distribution to capture the increased likelihood
of observing negative response ratios than positive response ratios (an
inherent property of natural logarithm transformation). Importantly,
however, a $\mathcal{SN}$ distribution was not assumed for priors of continuous
predictors where absolute slope estimates were expected to be small.
Lastly, priors for all group-level predictors, and σ were weak and flat (-∞ -
∞), or gamma-distributed (group-level predictors: Γ[1.5, 1.0]; σ: Γ[1.0, 1.0]).

Four observations (out of 165; ~2.5% of our data) were removed from
our analysis, owing to their abnormally large effect sizes relative to the
mean (> 3.5 times the sample s.d.; absolute change in body temperature >
10°C). To account for the possibility that posterior distributions for our
model coefficients were non-normal, we calculated modes for all coefficients

in place of means, and highest posterior density credible intervals (80% and
95%) around these modes[61]. All calculated means were marginalised unless
stated otherwise, and absolute changes in body temperature were
calculated from log-response ratios and assumed a baseline body
temperature of 38°C.

Estimating change in energy expenditure attributed to the stress-induced thermal response

A critical assumption of our trade-off hypothesis is that, in response to
a stressor, allowing body temperature to decrease in the cold or increase in
the warmth liberates a meaningful amount of energy that can then be
allocated toward coping with that stressor. With this assumption in mind,
we sought to estimate the relative extent to which stress-induced changes
in body temperature reduced expenditure toward thermoregulation among
our sampled vertebrates. To do so, we first estimated resting metabolic rate
of sampled animals at their known housing- and baseline body temperatures
(henceforth "baseline RMR") following established biophysical models[62],
and by using the R package nicheMapR[63]. These estimates represented
energy expenditure required to maintain thermal balance and assumed that:
(1) baseline body temperature was held constant, (2) loss of body heat to
the ambient environment occured by radiation, forced convection, and free
convection, but not conduction, and (3) the rate at which body heat was lost
to the ambient environment was inversely related to the resistance of both
the core body itself and the insulative layer of plumage or pelage[62]. For our
estimates, we assumed that all animals were elipsoids with volumes (m^3)
equal to their mass in g (body density = 1000 kg/m^3)[63], and with semimajor
axes being 1.2 and 1.3 times longer than semiminor axes for birds and
mammals respectively. Further, we also assumed that: (1) pelage or
plumage depth of sampled animals was equal the logarithm of their body
mass (g) in cm (e.g. pelage depth = 1 and 2 cm for a 10 and 100 g mammal
respectively), (2) thermal conductivity of an animal's pelage or plumage was

0.0272 W/m-°C[62], and (3) wind speed during sampling was negligable (0.1

m/s). Following these estimates, we then re-estimated resting metabolic

rates across the same ambient temperatures, but with body temperatures

representing those measured after stress exposure (henceforth, "stress-

induced RMR"), rather than at baseline. Changes in expenditure toward

thermoregulation were then calculated as the difference between stress-

induced and baseline RMR value. These change values are expected to be

conservative, owing to the exclusion of temperature effects of metabolic

processes themselves (i.e. through q10 effects[64]).

Figures

Figure 1 | Trade-off model explaining the outcome of stress-induced changes in body temperature. "Available energy" indicates surplus energy available for allocation toward biological processes after accounting for maintenance metabolism (i.e. basal metabolism). Thickness of lines indicates the magnitude of influence that one variable exerts on another, in the direction of the arrow.

Figure 2 | PRISMA flowchart detailing the process used to select studies for subsequent data collection.

Effect on Body Temperature
Responses to Stressors

Figure 3 | Factors influencing stress-induced changes in core body temperature among terrestrial endotherms. Dots represent the estimated effect of a given parameter on the magnitude and direction of stress-induced changes in body temperature (measured by a log response ratio, "lRR"). "Time" indicates latency to a maximum or minimum body temperature measurement after stressor onset; "relative resting metabolic rate" indicates the residuals of log-transformed resting metabolic rate (W) regressed against log-transformed body mass (g), then categorised as positive or negative. Negative slopes represent a tendency to reduce core body temperature, while positive slopes represent a tendency to increase core body temperature. Black grey represent 80% highest posterior density intervals credible intervals (HPDIs) and black whiskers represent 95% HPDIs.

Figure 4 | Effect of air temperature and body mass on the magnitude and direction of stress-induced changed in core temperature. A. Negative log response ratios represent stress-induced declines in body temperature, while positive log response ratios represent stress-induced increases in body temperature. Trend lines represent predicted relationships for an animal of a given body mass (g), while holding all other model covariates (shown in Fig. 3) at their averages. Dark ribbons represent 80% highest posterior density intervals (HPDIs) and pale ribbons represent 95% HPDIs. **B.** Large dots indicate average changes in body temperature for species above and below mean body mass for our sample (~222g; beige and green respectively), and at ambient temperatures above and below the average across studies (20.1°C). Mean changes in body temperature are calculated from log-response ratios and assume a baseline body temperature of 38°C. Small dots represent raw data values and error bars indicate 80% and 95% highest posterior density intervals (HPDIs; black and grey respectively). **C.** Large dots display mean changes in resting energy expenditure attributed to stress-induced thermal responses, as predicted by biophysical models. Again, small dots represent raw data values and error bars indicate 80% and 95% highest posterior density intervals (HPDIs; black and grey respectively).

Figure 5 | Effect of relative energy expenditure at rest on the magnitude and direction of stress-induced changed in core temperature. Relative energy expenditure at rest represents the residuals extracted from a linear mixed effects model regressing log-transformed resting metabolic rate (W) against log-transformed body mass (g), then categorised as "low" (i.e. negative residuals) or "high" (i.e. positive residuals). **A.** Negative log response ratios represent stress-induced declines in body temperature, while positive log response ratios represent stress-induced increases in body temperature. Dark ribbons represent 80% highest posterior density intervals (HPDIs) and pale ribbons represent 95% HPDIs. Trend lines and credible intervals are marginalised across all other model predictors. **B.** Large dots represent average changes in body temperature for species with high or low relative energy expenditure (red and navy respectively), and at ambient temperatures above and below the average across studies (20.1°C). Mean changes in body temperature were calculated from log-response ratios and assume a baseline body temperature of 38°C. Small dots represent raw data values and whiskers indicate 80% and 95% highest posterior density intervals (HPDIs) around means (black and grey respectively). **C.** Large dots display mean changes in resting energy expenditure attributed to stress-induced thermal responses, as predicted by biophysical models. Again, small dots represent raw data values and error bars indicate 80% and 95% highest posterior density intervals (HPDIs; black and grey respectively).

Figure 6 | Effect of time post-stress exposure on stress-induced changes in body temperature. Time post stress-exposure is represented by the latency between stress exposure treatment and the measurement of a maximum or minimum body temperature measurement (in seconds; s). Negative log response ratios represent stress-induced declines in body temperature, while positive log response ratios represent stress-induced increases in body temperature. The dark ribbon represents 80% highest posterior density intervals (HPDIs) and the pale ribbons represent 95% HPDIs. The trend lines and credible intervals are marginalised across all other model predictors.

References

1. S. P. Mattern, Galen and his patients. *Lancet* 378, 478-479 (2011).

2. I. Johnston, G. H. R. Horsley, "Galen's method of medicine, volume III. Vol. 518" (Harvard University Press, 2011).

3. I. S. Yeo, Hippocrates in the context of Galen: Galen's commentary on the classification of fevers in Epidemics VI. *Stud. Anc. Med.* 31, 433–43 (2005).

4. M. Parviz, M. Anushiravani, M. Keshavarz, The relationship between vital spirit and fevers in the Canon of Medicine: a probable solution for the controversy over stress-induced hyperthermia. *Iran. J. Public Health* 42, 1073 (2013).

5. R.M. Bush, R. Custer, J. Smeller, L. M. Bush, Physiologic measures of nonhuman primates during physical restraint and chemical immobilization. *J. Am. Vet. Med. Assoc.* 171, 866-869 (1977).

6. D.G. Menon, D. C. Bennett, A. L. Schaefer, K. M. Cheng, Transportation stress and the incidence of exertional rhabdomyolysis in emus (*Dromaius novaehollandiae*). *Poult. Sci.* 93, 273-284 (2014).

7. F. Fazio, S. Casella, E. Giudice, C. Giannetto, G. Piccione, Evaluation of secondary stress biomarkers during road transport in rabbit. *Livest. Sci.* 173, 106-110 (2015).

8. C. Carere, K. Van Ooers, Shy and bold great tits (*Parus major*): body temperature and breath rate in response to handling stress. *Physiol. Behav.* 82, 905-912 (2004).

9. K. A. Muise, A. K. Menzies, C. K. Willis, Stress-induced changes in body temperature of silver-haired bats (*Lasionycteris noctivagans*). *Physiol. Behav.* 194, 356-361 (2018).

10. J. K. Robertson, G. Mastromonaco, G. Burness, Evidence that stress-induced changes in surface temperature serve a thermoregulatory function. *J. Exp. Biol* 223, jeb213421 (2020a)

11. J. K. Robertson, G. F. Mastromonaco, G. Burness, Social hierarchy reveals thermoregulatory trade-offs in response to repeated stressors. *J. Exp. Biol* 223, jeb229047 (2020b).

12. A. Lewden, A. Nord, M. Petit, F. Vézina, Body temperature responses to handling stress in wintering black-capped chickadees (*Poecile atricapilus*). *Physiol. Behav.* 179, 49–54 (2017).

13. Y. Yokoi, Effect of ambient temperature upon emotional hyperthermia and hypothermia in rabbits. *J. Appl. Physiol.* 21, 1795-1798 (1966).

14. A. Amar, A. K. Sanyal, Immobilization stress in rats: effect on rectal temperature and possible role of brain monoamines in hypothermia. *Psychopharmacol.* 73, 157-160 (1981).

15. I. Maggini, F. M. Tahamtani, M, Cardinale, L. Fusani, C. Carere, Body temperature upon mist-netting procedures in three species of migratory songbirds at a stopover site: implications for welfare. *Anim. Welf.* 27, 93-101 (2018).

16. S. C. Stearns, "The Evolution of Life Histories" (Oxford University Press, 1992).

17. V. Careau, Energy intake, basal metabolic rate, and within-individual trade-offs in men and women training for a half marathon: a reanalysis. *Physiol. Biochem. Zool.* 90, 392-398 (2017).

18. N. C. Long, A. J. Vander, M. J. Kluger, Stress-induced rise of body temperature in rats is the same in warm and cool environments. *Physiol. Behav.* 47, 773-775, (1990).

19. D. M. Vianna, P. Carrive, Changes in cutaneous and body temperature during and after conditioned fear to context in the rat. *Euro. J. Neurosci.* 21, 2505-2512 (2005).

20. P. Jerem, S. Jenni-Eiermann, K. Herborn, D. McKeegan, D. J. McCafferty, R. G. Nager, Eye region surface temperature reflects both energy reserves and circulating glucocorticoids in a wild bird. *Sci. Rep.* 8, 1-10 (2018).

21. A. Nord, L. P. Folkow, Ambient temperature effects on stress-induced hyperthermia in Svalbard ptarmigan. *Biol. Open.* 8, bio043497 (2019).

22. F. Andreasson, A. Nord, J. Å. Nilsson, Body temperature responses of great tits *Parus major* to handling in the cold. *Ibis* 162, 836-844 (2020).

23. T. Oka, Stress-induced hyperthermia and hypothermia. *Handb. Clin. Neurol.* 157, 599-621 (2018).

24. R. Grant, Emotional hypothermia in rabbits. *Am. J. Physiol.* 160, 285-290 (1950).

25. E. Briese, Cold increases and warmth diminishes stress-induced rise of colonic temperature in rats. *Physiol. Behav.* 51, 881-883 (1992).

26. L. Volicer, C. West, L. Green, Effect of dietary restriction and stress on body temperature in rats. *J. Gerontol.* 39, 178-182 (1984).

27. C. J. Gordon, Relationship between preferred ambient temperature and autonomic thermoregulatory function in rat. *Am. J. Physiol. Reg. Integr. Comp. Physiol.* 252, R1130-R1137 (1987).

28. E. Briese, M. Cabanac, Stress hyperthermia: Physiological arguments that it is a fever. *Physiol. Behav.* 49, 1153–1157 (1991).

29. M. Cabanac, M. Guillemette. Temperature and heart rate as stress indicators of handled common eider. *Physiol. Behav.* 74, 475–479 (2001).

30. C. H. Vinkers, L. Groenink, M. J. van Bogaert, K. G. Westphal, C. J. Kalkman, R. van Oorschot, R. S. Oosting, B. Olivier, S. M. Korte, Stress-induced hyperthermia and infection-induced fever: Two of a kind? *Physiol. Behav.* 98, 37–43. (2009).

31. F. Vézina, O. P. Love, M. Lessard, T. D. Williams, Shifts in metabolic demands in growing altricial nestlings illustrate context-specific relationships between basal metabolic rate and body composition. *Physiol. Biochem. Zool.* 82, 248-257 (2009).

32. V. Careau, P. Bergeron, D. Garant, D. Réale, J. R. Speakman, M. M. Humphries, The energetic and survival costs of growth in free-ranging chipmunks. *Oecologia* 171, 11-23 (2013).

33. K. D. Foster, C. A. Conn, M. J. Kluger, Fever, tumor necrosis factor, and interleukin-6 in young, mature, and aged Fischer 344 rats. *Am. J. Physiol.* 262, R211-R215 (1992).

34. M. Wachulec, E. Peloso, E. Satinoff, Individual differences in response to LPS and psychological stress in aged rats. *Am. J. Physiol.* 272, R1252-R1257 (1997).

35. F. Andreasson, A. Nord, J. Å. Nilsson, Age-dependent effects of predation risk on night-time hypothermia in two wintering passerine species. *Oecologia* 189, 329-337 (2019).

36. I. Rodriguez-Prieto, E. Fernández-Juricic, J. Martín, To run or to fly: low cost versus low risk escape strategies in blackbirds. *Behav.* 145, 1125-1138 (2008).

37. J. Siemens, G. B. Kamm, Cellular populations and thermosensing mechanisms of the hypothalamic thermoregulatory center. Pflügers Archiv-*Euro. J. Physiol.* 470, 809-822. (2018).

38. Y. S. Pae, H. Lai, A. Horita, Hyperthermia in the rat from handling stress blocked by naltrexone injected into the preoptic-anterior hypothalamus. *Pharmacol. Biochem. Behav.* 22, 337-339 (1985).

39. W. H. Cao, S. F. and Morrison, Brown adipose tissue thermogenesis contributes to fentanyl-evoked hyperthermia. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 288, R723-R732 (2005).

40. J. A. DiMicco, D. V. Zaretsky, The dorsomedial hypothalamus: a new player in thermoregulation. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 292, R47-R63 (2007).

41. N. Kataoka, H. Hioki, T. Kaneko, K. Nakamura, Psychological stress activates a dorsomedial hypothalamus-medullary raphe circuit driving brown adipose tissue thermogenesis and hyperthermia. *Cell Metab.* 20, 346-358 (2014).

42. M. J. Angiletta Jr, J. P. Youngblood, L. K. Neel, J. M. VandenBrooks, The neuroscience of adaptive thermoregulation. *Neurosci. Lett.* 692, 127-136 (2019).

43. N. Pecoraro, H. De Jong, A. B. Ginsberg, M. F. Dallman, Lesions of the medial prefrontal cortex enhance the early phase of psychogenic fever to unexpected sucrose concentration reductions, promote recovery from negative contrast and enhance spontaneous recovery of sucrose-entrained anticipatory activity. *Neurosci.* 153, 901-917 (2008).

44. C. H. Vinkers, E. Y. Bijlsma, L. C. Houtepen, K. G. Westphal, J. G. Veening, L. Groenink, B. Olivier, Medial amygdala lesions differentially influence stress responsivity and sensorimotor gating in rats. *Physiol. Behav.* 99, 395-401 (2010).

45. T. W. Rusch, "Integrating spatial constraints and biotic interactions to assess the costs of thermoregulation by lizards". PhD Thesis, Arizona State University, USA. (2016).

46. S. R. Tracey, K. Hartmann, M. Leef, J. McAllister, Capture-induced physiological stress and postrelease mortality for Southern bluefin tuna (*Thunnus maccoyii*) from a recreational fishery. *Can. J. Fish. Aquat. Sci.* 73, 1547-1556 (2016).

47. Y. Konno, K. Konno, Myosin denaturation in "Burnt" Bluefin tuna meat. *Fish. Sci.* 80, 381-388 (2014).

48. C. G. Wilber, P. F. Robinson, Effect of restraint on body temperature in guinea pigs. *J. Appl. Physiol.* 12, 214-216 (1958).

49. A. M. Lees, H. E. Salvin, I. Colditz, C. Lee, The influence of temperament on body temperature response to handling in Angus cattle. *Animals* 10, 172 (2020).

50. J. I. Schindelin, E. Arganda-Carreras, V. Frise, M. Kaynig, T. Longair, S. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J. Y. Tinevez, Fiji: An open-source platform for biological-image analysis. *Nat. Methods* 9, 676–682 (2012).

51. A. McKechnie, B. Wolf, The allometry of avian basal metabolic rate: Good predictions need good data. *Physiol. Biochem. Zool.* 77, 502–521 (2004).

52. N. Upham, J. Esselstyn, W. Jetz, Inferring the mammal tree: Species-level sets of phylogenies for questions in ecology, evolution, and conservation. *PLoS Biol.* 17, e3000494 (2019).

53. W. Jetz, G. H. Thomas, J. B. Joy, K. Hartmann, A. Mooers, The global diversity of birds in space and time. *Nature* 491, 444–448 (2012).

54. P. Ericson,, C. L. Anderson, T. Britton, A. Elzanowski, U. S. Johansson, M. Källersjö, J. I. Ohlson, T. J. Parsons, D. Zuccon, G. Mayr, Diversification of neoaves: Integration of molecular sequence data and fossils. *Biol. Lett.* 2(1):543–547 (2006).

55. R. Bouckaert, J. Heled, D. Kühnert, T. Vaughan, C.-H. Wu, D. Xie, M. A. Suchard, A. Rambaut, A. J. Drummond, BEAST 2: A software platform for Bayesian evolutionary analysis. *PLoS Comput. Biol.* 10, e1003537 (2014).

56. E. Paradis, K. Schliep, Ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 35, 526–528 (2019).

57. R Core Team. R: "A language and environment for statistical computing" (R Foundation for Statistical Computing, 2021).

58. H. Wickham, "Ggplot2: elegant graphics for data analysis" (Springer, 2016).

59. P. C. Bürkner, brms: an R package for Bayesian multilevel models using Stan. *J. Stat. Soft.* 80, 1-28 (2017).

60. M. J. Lajeunesse, Bias and correction for the log response ratio in ecological meta-analysis. *Ecology* 96, 2056-2063 (2015).

61. A. Gelman, A. Vehtari, J. Carlin, H. S. Stern, D. Dunson, D. Rubin, "Bayesian data analysis" (CRC Press, 1995).

62. W. P. Porter, M. Kearney, Size, shape, and the thermal niche of endotherms. *Proc. Natl. Acad. Sci. U.S.A.* 106, 19666-19672 (2009).

63. M. R. Kearney, W. P. Porter, NicheMapR–an R package for biophysical modelling: the microclimate model. *Ecography* 40, 664-674 (2017).

64. C. R. White, R. S. Seymour, Allometric scaling of mammalian metabolism. *J. Exp. Biol.* 208, 1611-1619 (2005).