HYENA detects oncogenes activated by distal enhancers in cancer

Anqi Yu [1, *], Ali E. Yesilkanal [1, *], Ashish Thakur [2], Fan Wang [1], Yang Yang [1], William Phillips [1], Xiaoyang Wu [1,3], Alexander Muir [1,3], Xin He [2], Francois Spitz [2], Lixing Yang [1, 2, 3,**]

1. Ben May Department for Cancer Research, University of Chicago, Chicago IL, USA
2. Department of Human Genetics, University of Chicago, Chicago IL, USA
3. University of Chicago Comprehensive Cancer Center, Chicago, IL, USA

* these authors contributed equally

** correspondence: lixingyang@uchicago.edu

Abstract

Somatic structural variations (SVs) in cancer can shuffle DNA content in the genome, relocate regulatory elements, and alter genome organization. Enhancer hijacking occurs when SVs relocate distal enhancers to activate proto-oncogenes. However, most enhancer hijacking studies have only focused on protein-coding genes. Here, we develop a computational algorithm "HYENA" to identify candidate oncogenes (both protein-coding and non-coding) activated by enhancer hijacking based on tumor whole-genome and transcriptome sequencing data. HYENA detects genes whose elevated expression is associated with somatic SVs by using a rank-based regression model. We systematically analyze 1,148 tumors across 25 types of adult tumors and identify a total of 192 candidate oncogenes including many non-coding genes. A long non-coding RNA *TOB1-AS1* is activated by various types of SVs in 10% of pancreatic cancers through altered 3-dimension genome structure. We find that high expression of *TOB1-AS1* can promote cell invasion and metastasis. Our study highlights the contribution of genetic alterations in non-coding regions to tumorigenesis and tumor progression.

Introduction

At mega-base-pair scale, linear DNA is organized into topologically associating domains (TADs) [1], and gene expression is regulated by DNA and protein interactions governed by 3D genome organization. Enhancer-promoter interactions are mostly confined within TADs [2–4]. Non-coding somatic single nucleotide variants (SNVs) in promoters and enhancers have been linked to transcriptional changes in nearby genes and tumorigenesis [5]. Structural variations (SVs), including deletions, duplications, inversions, and translocations, can dramatically change TAD organization and gene regulation [6] and subsequently contribute to tumorigenesis. Previously, we discovered that *TERT* is frequently activated in chromophobe renal cell carcinoma by relocation of distal enhancers [7], a mechanism referred to as enhancer hijacking (**Fig. 1a**). In fact, many oncogenes, such as *BCL2* [8], *MYC* [9], *TAL1* [10], *MECOM/EVI1* [11], *GFI1* [12], *IGF2* [13], *PRDM6* [14], and *CHD4* [15], can be activated through this mechanism. These examples demonstrate that genomic architecture plays an important role in cancer pathogenesis. However, the vast majority of the known enhancer hijacking target oncogenes are protein-coding genes, and few non-coding genes have been reported to promote diseases through enhancer hijacking. Here, we refer to non-coding genes as all genes that are not protein-coding. They include long non-coding RNAs (lncRNAs), pseudogenes, and other small RNAs such as microRNAs, small nuclear RNAs (snRNAs), small nucleolar RNAs (snoRNAs), etc. They are known to play important roles in many biological processes [16] and some are known to drive tumorigenesis [17,18]. In this study, we will focus on identifying oncogenes, including oncogenic non-coding genes activated by enhancer hijacking.

Several existing algorithms can detect enhancer hijacking target genes based on patient cohorts, such as CESAM [13] and PANGEA [15]. These two algorithms implemented linear regression and elastic net model (also based on linear regression) to associate elevated gene expression with nearby SVs, respectively. PANGEA also considers the effects of somatic SNVs on gene expression. However, a major drawback of these algorithms is that linear regression is quite sensitive to outliers. Outliers are very common in gene expression data from cancer samples and can seriously impair the performances of these algorithms. In addition, CESAM is optimized for microarray data, while PANGEA depends on annotation of tissue-specific promoter-enhancer pairs, which are not readily available for many tumor types. Cis-X [19] and NeoLoopFinder [20] can detect enhancer hijacking target genes based on individual samples. However, these tools have limitations in detectable genes and input data. Cis-X detects *cis*-activated genes based on allele-specific expression, which requires the genes to carry heterozygous SNVs. NeoLoopFinder takes Hi-C, Chromatin Interaction Analysis with Paired-End Tag (ChIA-PET), or similar data measuring chromatin interactions as input, which remain very limited. Furthermore, identification of recurrent mutational events that result in oncogenic activation requires large patient cohorts. Therefore, tools that use whole-genome and transcriptome sequencing data, which are available at much larger sample sizes, would be more useful in identifying SV-driven oncogene activation. Finally, no non-coding oncogenes have been reported as enhancer hijacking targets by the above algorithms. A recent study on SVs altering gene expression in Pan-Cancer Analysis of Whole Genomes (PCAWG) samples [21] only considered protein-coding genes but not non-coding genes.

Here, we developed <u>Hi</u>jacking of <u>En</u>hancer <u>A</u>ctivity (HYENA) using normal-score regression
and permutation test to detect candidate enhancer hijacking genes (both protein-coding and non-
coding genes) based on tumor whole-genome and transcriptome sequencing data from patient
cohorts. Among the 192 putative oncogenes detected by HYENA, we studied the oncogenic
functions of a lncRNA, *TOB1-AS1*, and demonstrated that it is a regulator of cancer cell invasion
in vitro and tumor metastasis in vivo.
76

Results

HYENA workflow

Conceptually, the SVs leading to elevated gene expression are expression quantitative trait loci (eQTLs). The variants are SVs instead of commonly used germline single nucleotide polymorphisms (SNPs) in eQTL analysis. With somatic SVs and gene expression measured from the same tumors through whole-genome sequencing (WGS) and RNA sequencing (RNA-Seq), we can identify enhancer hijacking target genes by eQTL analysis. However, the complexities of cancer and SVs pose many challenges. For instance, there is tremendous inter-tumor heterogeneity—no two tumors are identical at the molecular level. In addition, there is substantial intra-tumor heterogeneity as tumor tissues are always mixtures of tumor, stromal, and immune cells. Moreover, genome instability is a hallmark of cancer, and gene dosages are frequently altered [22]. Furthermore, gene expression networks in cancer are widely rewired [23], and outliers of gene expression are common.

Here, we developed an algorithm HYENA to overcome the challenges described above (see more details in Methods Section). We used a gene-centric approach to search for elevated expression of genes correlated with the presence of SVs within 500 kb of transcription start sites (**Fig. 1b**). Although promoter-enhancer interaction may occur as far as several mega-bases, mega-base-level long-range interactions are extremely rare. In addition, although duplicated enhancers can upregulate genes [24,25], we do not consider these as enhancer hijacking events since no neo-promoter-enhancer interactions are established. However, small deletions can remove TAD boundaries or repressive elements and lead to neo-promoter-enhancer interactions (**Fig. 1a**). Therefore, small tandem duplications were discarded, and small deletions were retained. For each gene, we annotated SV status (presence or absence of nearby SVs) for all samples. Samples in which the testing genes were highly amplified were discarded since many of these genes are amplified by circular extrachromosomal DNA (ecDNA) [26], and ecDNA can promote accessible chromatin [27] with enhancer rewiring [28]. Only genes with nearby SVs in at least 5% of tumors were further considered. In contrast to CESAM and PANGEA, we did not use linear regression to model the relationships between SV status and gene expression because linear regression is sensitive to outliers and many false positive associations would be detected [29]. Instead, we used a rank-based normal-score regression approach. After quantile normalization of gene expression for both protein-coding and non-coding genes, we added small Gaussian noises to gene expression for tie breaking, ranked the genes based on quantile-normalized noise-added expression, and transformed the ranks to the quantiles of the standard normal distribution. We used the z scores (normal scores) of the quantiles as dependent variables in regression. In the normal-score regression model, tumor purity, copy number of the tested gene, patient age, and sex were included as covariates since these factors confound gene expression. We also included gene expression principal components (PCs) that were not correlated with SV status to model unexplained variations in gene expression. To deduce a better null distribution, we permuted the gene expression 100 times and ran the same regression models. All P values from the permutations were pooled together and used as the null distribution to calculate empirical P values. Then, multiple testing corrections were performed on one-sided P values since we are

only interested in elevated gene expression under the influence of nearby SVs. Finally, genes
were discarded if their elevated expression could be explained by germline eQTLs. The
remaining genes were candidate enhancer hijacking target genes.

Benchmarking performances

There is no gold standard available to comprehensively evaluate the performance of HYENA.
We compared HYENA's performance to two other algorithms—CESAM and PANGEA. All
three algorithms were run on the same somatic SVs and gene expression data from six types of
adult tumors profiled by the PCAWG (**Supplementary Table S1**): malignant lymphoma
(MALY), stomach/gastric adenocarcinoma (STAD), chromophobe renal cell carcinoma (KICH),
colorectal cancer (COAD/READ), thyroid cancer (THCA), and lung squamous cell carcinoma
(LUSC) [21]. Note that PANGEA depends on promoter-enhancer interactions predicted from cell
lines which were not available for thyroid tissue. Therefore, thyroid cancer data were not
analyzed by PANGEA.

To compare the sensitivity of HYENA to the other algorithms, we used eight known enhancer
hijacking target genes including *MYC* [9], *BCL2* [8], *CCNE1* [30], *TERT* [7], *IGF2* [13,30] (in two tumor
types), *IGF2BP3* [31] and *IRS4* [13]. We also expect immunoglobulin genes to be detected as
enhancer hijacking candidates in malignant lymphoma due to V(D)J recombination since the
lymphomas in the PCAWG are B-cell derived Burkitt lymphomas [32]. In B cells, V(D)J
recombination occurs to join different variable (V), joining (J) and constant (C) segments to
produce antibodies with a wide range of antigen recognition ability. Therefore, certain segments
have elevated expression and the recombination events can be detected as somatic SVs. Out of
the eight known enhancer hijacking genes, HYENA detected five (*MYC*, *BCL2*, *TERT*, *IGF2*,
and *IGF2BP3*) (**Fig. 2a** and **Supplementary Fig. S1**), CESAM detected three (*MYC*, *BCL2*, and
TERT), and PANGEA did not detect any (**Fig. 2a**). In the five tumor types analyzed by all three
algorithms, HYENA identified a total of 25 candidate genes, CESAM identified 19, whereas
PANGEA identified 255 genes (**Fig. 2b**, **Supplementary Tables S2**, **S3** and **S4**). Six genes were
detected by both HYENA and CESAM, while PANGEA had little overlap with the other
algorithms (**Fig. 2b**). Of the 16 genes detected by HYENA in malignant lymphoma, there were
two immunoglobulin light chain genes from lambda cluster (*IGLC7* and *IGLJ7*)
(**Supplementary Table S2**). CESAM detected 11 genes with one being immunoglobulin gene
(*IGLC7*) (**Supplementary Table S3**). In contrast, PANGEA detected 30 candidate genes, but
none were immunoglobulin genes (**Supplementary Table S4**).

The ability of the algorithms to detect known target genes seems to be sensitive to sample size.
Both *IGF2* and *IRS4* were initially discovered by CESAM as enhancer hijacking target genes
using copy number variation (CNV) breakpoints profiled by microarray with much larger sample
sizes (378 colorectal cancers and 497 lung squamous cell carcinomas) [13]. In the PCAWG, the
sample sizes with both WGS and RNA-Seq were smaller (51 colorectal cancers and 47 lung
squamous cell carcinomas). HYENA detected *IGF2* in colorectal cancer but not *IRS4*, whereas
CESAM and PANGEA detected neither. In stomach/gastric adenocarcinoma, *IGF2* and *CCNE1*
were identified as enhancer hijacking target genes in a cohort of 208 samples [30]. Neither of these
genes were detected by any of the algorithms because there were only 29 stomach tumors in the

PCAWG. Therefore, known target genes missed by HYENA were likely due to small sample
size. In summary, HYENA had the best sensitivity of the three algorithms.

To evaluate specificity of the algorithms, we ran each algorithm on 20 datasets generated by
randomly shuffling gene expression data in both MALY and breast cancer (BRCA). Since these
gene expression data were random, there should be no associations between SVs and gene
expression, and all genes detected should be false positives. In malignant lymphoma with
observed gene expression, HYENA, CESAM, and PANGEA detected 16, 11, and 30 candidate
genes respectively (**Supplementary Tables S2, S3 and S4**). In the 20 random gene expression
datasets for malignant lymphoma, HYENA detected an average of 0.5 genes per dataset (**Fig.
2c**), and CESAM detected an average of 0.5 genes per dataset, whereas PANGEA detected an
average of 40 genes per dataset (**Supplementary Fig. S2**). In breast cancer with observed gene
expression, HYENA, CESAM, and PANGEA detected 61, 9, and 2,309 candidate genes,
respectively (**Supplementary Tables S2, S3 and S4**). In 20 random gene expression datasets for
breast cancer, HYENA, CESAM, and PANGEA detected 0.35, 0.9 and 2,296 genes on average
(**Fig. 2c** and **Supplementary Fig. S2**). In both tumor types, the numbers of false positives called
by PANGEA in random datasets were comparable to the numbers of genes detected with
observed gene expression (**Supplementary Fig. S2**). In summary, HYENA predicted the least
number of false positives among the three algorithms.

Overall, HYENA has superior sensitivity and specificity in the detection of candidate enhancer
hijacking target genes.

Enhancer hijacking candidate genes in the PCAWG

We used HYENA to analyze a total of 1,146 tumors across 25 tumor types in the PCAWG with
both WGS and RNA-Seq data. When each tumor type was analyzed individually, we identified
192 candidate enhancer hijacking target genes in total (**Supplementary Tables S1** and **S2**), five
of which were known enhancer hijacking targets (**Fig. 3a**). *TERT* was the only gene identified in
two tumor types/cohorts (KICH from the US and renal cell carcinoma [RECA] from Europe).
All other candidate genes were only detected in one tumor type, highlighting high tumor type
specificity of the findings. The number of genes detected in each tumor type also differed
dramatically (**Fig. 3b**). No genes were detected in bladder cancer (BLCA), cervical cancer
(CESC), glioblastoma multiforme (GBM), or low-grade glioma (LGG), probably due to their
small sample sizes. BRCA had the greatest number of candidate genes likely due to the large
sample size as well as the abundance of SVs resulting from homologous recombination
deficiency (HRD) [33]. Although ovarian cancer (OV) also suffers from HRD and had a sample
size comparable with breast cancer, there were many fewer enhancer hijacking target genes
detected. Thyroid cancer genomes were among the most stable genomes in the PCAWG [34].
However, the 15 enhancer hijacking target genes identified in thyroid cancer exceeded the
number of candidate genes in ovarian cancer as well as many other tumor types. Among these 15
genes, *IGF2BP3* was a known oncogene activated by enhancer hijacking [31,35]. There were two
liver cancer cohorts with comparable sample sizes—LIHC from the US and LIRI from Japan.
Interestingly, a total of 18 genes were identified in the US cohort whereas no genes were found
in the Japanese cohort. One possible reason for such a drastic difference could be that hepatitis B

virus (HBV) infection is more common in liver cancer in Japan [36], and virus integration into the
tumor genome can result in oncogene activation [37]. In Chronic Lymphocytic Leukemia (CLLE),
a total of nine genes were detected, and seven were immunoglobulin genes from both lambda
and kappa clusters (**Supplementary Tables S2**). Given that sample size and genome instability
can only explain a small fraction of the variations of enhancer hijacking target genes detected in
different tumor types, the landscape of enhancer hijacking in cancer seems to be mainly driven
by the underlying disease biology. Intriguingly, out of the 192 candidate genes, 73 (38%) were
non-coding genes including lncRNAs and microRNAs (**Fig. 3b**).

Neo-TADs formed through somatic SVs

Next, we focused on the most frequently altered candidate non-coding enhancer-hijacking target
gene in pancreatic cancer: *TOB1-AS1* (**Fig. 4a**), a lncRNA. *TOB1-AS1* was not detected as a
candidate gene by either CESAM (**Supplementary Table S3**) or PANGEA (**Supplementary
Table S4**) using the same input data. Seven (9.6%) out of 74 tumors had some forms of somatic
SVs near *TOB1-AS1* including translocations, deletions, inversions, and tandem duplications
(**Fig. 4b** and **Supplementary Table S5**). For example, tumor 9ebac79d-8b38-4469-837e-
b834725fe6d5 had a translocation between chromosomes 17 and 19 (**Fig. 4c**). The breakpoints
were upstream of *TOB1-AS1* and upstream of *UQCRFS1* (**Fig. 4d**). In tumor 748d3ff3-8699-
4519-8e0f-26b6a0581bff, there was a 19.3 Mb deletion which brought *TOB1-AS1* next to a
region downstream of *KCNJ2* (**Fig. 4c** and **4e**).

We used Akita [38], a convolutional neural network that predicts 3D genome organization, to
assess the 3D architecture of the loci impacted by SVs. While 3D structures are dynamic and
may change with cell-type and gene activity, TAD boundaries are often more stable and remain
similar across different cell-types [1]. TAD boundaries are defined locally by the presence of
binding sites for CCCTC-binding factor (CTCF), a ubiquitously expressed DNA-binding protein
[1,39], and TAD formation arises from the stalling of the cohesin-extruded chromatin loop by
DNA-bound CTCF at these positions [40]. For this reason, one can reliably expect that upon
chromosomal rearrangements, normal TADs can be disrupted, and new TADs can form by
relocations of TAD boundaries. This assumption has been validated with direct experimental
evidence from examining the "neo-TADs" associated with SVs at different loci [41–43]. The
wildtype *TOB1-AS1* locus had a TAD between a CTCF binding site in *RSAD1* and another one
upstream of *SPAG9* (**Fig. 4d** and **Supplementary Fig. S3**). There were TADs spanning
UQCRFS1 and downstream of *KCNJ2* in the two partner regions (**Fig. 4d**, **4e** and
Supplementary Fig. S3). In tumor 9ebac79d-8b38-4469-837e-b834725fe6d5, the translocation
was predicted to lead to a neo-TAD resulting from merging the TADs of *TOB1-AS1* and
UQCRFS1 (**Fig. 4d**). In tumor 748d3ff3-8699-4519-8e0f-26b6a0581bff, another neo-TAD was
predicted to form as a result of the deletion that merged the TADs of *TOB1-AS1* and the
downstream portion of *KCNJ2* (**Fig. 4e**). In both cases, within these predicted neo-TADs, Akita
predicted strong chromatin interactions involving several CTCF binding sites and H3K27Ac
peaks between *TOB1-AS1* and its two SV partners (**Fig. 4d** and **4e** black arrows in the right
panels), indicating newly formed promoter-enhancer interactions. In the vicinity of the *TOB1-
AS1* locus, *TOB1-AS1* was the only gene with significant changes in gene expression. Similar

neo-TADs could be observed in two additional tumors (**Supplementary Fig. S4**). In two tumors harbored tandem duplications of *TOB1-AS1* of 317 kb and 226 kb, the *TOB1-AS1* TADs were expanded (**Supplementary Fig. S5a**). However, not all SVs near *TOB1-AS1* led to alterations in TAD architecture; for example, in tumor a3edc9cc-f54a-4459-a5d0-097879c811e5, *TOB1-AS1* was predicted to remain in its original TAD after a 4 Mb tandem duplication (**Supplementary Fig. S5b**). In summary, at least four out of the seven tumors harboring somatic SVs near *TOB1-AS1* were predicted to result in neo-TADs including *TOB1-AS1*. We then used another deep-learning algorithm called Orca [44] to predict 3D genome structure based on DNA sequences. Orca-predicted 3D genome architectures were very similar to Akita predictions (**Supplementary Fig. S6**) in neo-TAD formation due to SVs in the *TOB1-AS1* locus.

To further study the 3D genome structure of *TOB1-AS1* locus, we performed high-resolution in situ Hi-C sequencing for four pancreatic cancer cell lines. Among these, two cell lines (Panc 10.05 and PATU-8988S) had high expression of *TOB1-AS1*, whereas the other two (PANC-1 and PATU-8988T) had low expression (**Fig. 5a**). At mega-base-pair scale, three cell lines (Panc 10.05, PATU-8988S and PATU-8988T) carried several SVs (black arrows in **Fig. 5b**). In Panc 10.05, a tandem duplication (chr17:43,145,000-45,950,000) was observed upstream of *TOB1-AS1* (**Fig. 5b** black arrow in the left most panel and **Supplementary Table S6**). However, the breakpoint was too far away (2 Mb) from *TOB1-AS1* (chr17:48,944,040-48,945,732) and unlikely to regulate its expression. A neo chromatin loop was detected by NeoLoopFinder [20] near *TOB1-AS1* (chr17:34,010,000-48,980,000) driven by a deletion (chr17:34,460,000-47,450,000) detected by EagleC [45] (**Supplementary Fig. S7a**, **Supplementary Tables S6** and **S7**). The deletion breakpoint was also too far away (1.5 Mb) from *TOB1-AS1* and unlikely to regulate its expression either. No other SVs or neo chromatin loops were detected near *TOB1-AS1* (**Supplementary Tables S6** and **S7**). Interestingly, there was a CNV breakpoint (chr17:48,980,000) 36 kb downstream of *TOB1-AS1* (**Fig. 5c** left most panel) which was also the boundary of the neo chromatin loop. In the high copy region (upstream of the CNV breakpoint), heterozygous SNPs were present with allele ratios of approximately 4:1 (**Supplementary Fig. S8a**), whereas in the low copy region (downstream of the CNV breakpoint), all SNPs were homozygous (**Supplementary Fig. S8b**). These suggested that the DNA copy number changed from five copies to one copy at the CNV breakpoint. The gained copies must connect to some DNA sequences since there should not be any free DNA ends other than telomeres. Given that no off-diagonal 3D genome interactions were observed at chr17:48,980,000, we considered the possibilities that the high copy region was connected to repetitive sequences or to sequences that were not present in the reference genome. If so, reads mapped to the high copy region should have excessive amount of non-uniquely mapped mates or unmapped mates. However, this was not the case (**Supplementary Fig. S9**). The only possible configuration was a foldback inversion in which two identical DNA fragments from the copy gain region were connected head to tail (**Fig. 5d** bottom left panel). As a result, in Panc 10.05, there was a wildtype chromosome 17, two foldback-inversion-derived chromosomes, and a translocation-derived chromosome (**Fig. 5d** bottom left panel and **Supplementary Fig. S7b**). Foldback inversions are very common in cancer. If DNA double strand breaks are not immediately repaired, following replication, the two broken ends of sister chromatids can self-ligate head to tail and sometimes result in dicentric chromosomes [46,47]. Algorithms, such as hic-breakfinder [48] and EagleC [45], rely on off-diagonal 3D

genomic interactions in Hi-C contact matrix to detect SVs. However, foldback inversions do not form any off-diagonal interactions since the two connected DNA fragments have the same coordinates, so they are not detectable by existing algorithms. The 3D genome structure of *TOB1-AS1* locus in Panc 10.05 was quite distinct from the other three cell lines (**Fig. 5c**). The region immediately involved in the foldback inversion had homogeneous 3D interactions (**Fig. 5c** dashed blue triangle in the left most panel) suggesting that a neo-subdomain was formed (**Fig. 5d** right panel). The high expression of *TOB1-AS1* in Panc 10.05 was likely a combined effect of the copy gain and the neo-subdomain. In PATU-8988S and PATU-8988T, a shared SV (chr17:48,880,000-52,520,000) near *TOB1-AS1* was detected (**Fig. 5b** two right panels) since the two cell lines were derived from the same pancreatic cancer patient [49]. This shared SV could not regulate *TOB1-AS1* because it pointed away from *TOB1-AS1* (**Supplementary Fig. S10**). No other SVs were found near *TOB1-AS1* in these two cell lines. The high expression of *TOB1-AS1* in PATU-8988S was likely due to transcription regulation since the promoter of *TOB1-AS1* in PATU-8988S was more accessible than that in PATU-8988T (**Fig. 5e**). This result was consistent with a handful of patient tumors that had high expression of *TOB1-AS1* without any SVs (**Fig. 4a**).

Taken together, our results demonstrated that HYENA can detect genes activated by reorganization of 3D genome architecture.

Oncogenic functions of *TOB1-AS1*

TOB1-AS1 has been reported as a tumor suppressor in several tumor types [50,51]. However, HYENA predicted it to be an oncogene in pancreatic cancers. To test the potential oncogenic functions of *TOB1-AS1* in pancreatic cancer, we performed both in vitro and in vivo experiments. We surveyed pancreatic cancer cell line RNA-Seq data from Cancer Cell Line Encyclopedia (CCLE) and identified that the commonly transcribed isoform of *TOB1-AS1* in pancreatic cancers was ENST00000416263.3 (**Supplementary Fig. S11**). The synthesized *TOB1-AS1* cDNA was cloned and overexpressed in two pancreatic cancer cell lines, PANC-1 and PATU-8988T, both of which had low expression of *TOB1-AS1* (**Fig. 5a** and **Supplementary Fig. S12a**). In both cell lines, overexpression of *TOB1-AS1* (**Fig. 6a**) promoted in vitro cell invasion (**Fig. 6b**). In addition, three weeks after tail vein injection, PANC-1 cells with *TOB1-AS1* overexpression caused higher metastatic burden in immunodeficient mice than the control cells (**Fig. 6c**). Six weeks after orthotopic injection, mice carrying *TOB1-AS1* overexpressing PANC-1 cells showed exacerbated overall tumor burden (**Fig. 6d**), elevated primary tumor burden, and elevated metastatic burden in the spleen (**Fig. 6e** and **Supplementary Fig. S12b**). Liver metastasis was not affected (**Supplementary Fig. S12c**). In addition, we knocked down *TOB1-AS1* in two other pancreatic cancer cell lines Panc 10.05 and PATU-8988S, both of which had high expression of *TOB1-AS1* (**Fig. 5a** and **Supplementary Fig. S12a**), using two antisense oligonucleotides (ASOs) (**Fig. 6f**). *TOB1-AS1* expression was reduced by approximately 50% by both ASOs (**Fig. 6g**). Knockdown of *TOB1-AS1* substantially suppressed cell invasion in vitro (**Fig. 6h**). Note that PATU-8988T and PATU-8988S were derived from the same liver metastasis of a pancreatic cancer patient, and they had drastic difference in *TOB1-AS1* expression (**Fig. 5a** and **Supplementary Fig. S12a**). It was reported that PATU-8988S can form lung metastasis in

vivo with tail vein injection of nude mice, whereas PATU-8988T cannot form any metastasis in
any organ [49]. By altering the expression of *TOB1-AS1*, we were able to reverse the cell invasion
phenotypes in these two cell lines (**Fig. 6b** and **6h**). These results suggested that *TOB1-AS1*
carries important function in regulating cell invasion.

It is possible that *TOB1-AS1*, as an anti-sense lncRNA, transcriptionally regulates the expression
of the sense protein-coding gene *TOB1*. However, we did not find consistent correlations
between *TOB1-AS1* and *TOB1* expression in different pancreatic cancer cohorts and pancreatic
cancer cell lines (**Supplementary Fig. S12d**). Hence, it is unlikely that *TOB1-AS1* functions
through transcriptional regulation of *TOB1*. Although knocking down *TOB1-AS1* resulted in
down regulation of *TOB1* expression, an expected result given that the ASOs also targeted the
introns of *TOB1* (**Fig. 6f**), the decrease in *TOB1* expression was relatively mild at 10-20% (**Fig.
6g**). Overexpression of *TOB1-AS1* did not have major impact on *TOB1* expression (**Fig. 6a**).
Therefore, the oncogenic functions of *TOB1-AS1* that we observed in vitro and in vivo are likely
independent of *TOB1*. To gain further insights into the pathway that *TOB1-AS1* is involved in
and its downstream targets, we performed RNA-Seq on PANC-1-generated mouse tumors with
TOB1-AS1 overexpression and found that the most significantly differentially expressed gene
was *CNNM1* (**Supplementary Fig. S12e**). *CNNM1* is a cyclin and CBS domain divalent metal
cation transport mediator and is predicted to be involved in ion transport [52]. How *TOB1-AS1*
promotes cell invasion and tumor metastasis and whether *CNNM1* plays any roles require further
study.

Our results showed that the lncRNA *TOB1-AS1* is oncogenic and has a pro-metastatic function in
pancreatic cancer, and HYENA is able to detect novel proto-oncogenes activated by distal
enhancers.

Discussion

Here, we report a computational algorithm HYENA to detect candidate oncogenes activated by distal enhancers via somatic SVs. These SV breakpoints fell in the regulatory regions of the genome and caused shuffling of regulatory elements, altering gene expression. The candidate genes we detected were not limited to protein-coding genes but also included non-coding genes. Our in vitro and in vivo experiments showed that a lncRNA identified by HYENA, *TOB1-AS1*, was a potent oncogene in pancreatic cancers.

HYENA detects candidate genes based on patient cohorts rather than individual samples. Genes need to be recurrently rearranged in the cohort to be detectable, and HYENA aims to identify oncogenes recurrently activated by somatic SVs since these events are under positive selection. Therefore, sample size is a major limiting factor. Of the eight ground truth cases, HYENA only detected five (**Fig. 2a**); undetected genes were likely due to small sample size. However, genes detected in individual tumors by tools such as cis-X and NeoLoopFinder may not be oncogenes, and recurrent events would be required to identify candidate oncogenes.

The candidate genes identified by HYENA have statistically significant associations between nearby somatic SVs and elevated expression. However, the relationship may not be causal. It is possible that the presence of SVs and gene expression are unrelated, but both are associated with another factor. We modeled other factors to the best of our ability including gene dosage, tumor purity, patient sex, age, and principal components of gene expression. In addition, it is also possible that the high gene expression caused somatic SVs. Open chromatin and double helix regions unwound during transcription are prone to double-strand DNA breaks which may produce somatic SVs. Therefore, it is possible that some of the candidate genes are not oncogenes. Functional studies are required to determine the disease relevance of the candidate genes.

Note that the predicted 3D genome organization is not cell-type-specific. Akita was trained on five high quality Hi-C and Micro-C datasets (HFF, H1hESC, GM12878, IMR90 and HCT116) [38] and predicts limited cell-type-specific differences. Therefore, the predicted TADs reflect conserved 3D genome structure in the five cell types (foreskin fibroblast, embryonic stem cell, B-lymphocyte, lung fibroblast and colon cancer). There were minor differences between HFF and H1hESC (**Supplementary Fig. S3**) in genome organization. For example, the left boundary of the TAD at the *UQCRFS1* locus was different between HFF and H1hESC (**Supplementary Fig. S3a**). Nonetheless, the translocation between chromosomes 17 and 19 removed the left boundary and merged the right side of the *UQCRFS1* TAD with the *TOB1-AS1* TAD (**Fig. 4d**). Therefore, the cell-type difference likely does not have major impact on our results.

Methods

Datasets

This study used data generated by the Pan-Cancer Analysis of Whole Genomes (PCAWG). We limited our study to a total of 1,146 tumor samples for which both whole-genome sequencing (WGS) and RNA-Seq data were available. The data set was composed of cancers from 25 tumor types including 23 bladder urothelial cancers (BLCA), 88 breast cancers (BRCA), 20 cervical squamous cell carcinomas (CESC), 68 chronic lymphocytic leukemias (CLLE), 51 colorectal cancers (COAD/READ), 20 glioblastoma multiforme (GBM), 42 head and neck squamous cell carcinomas (HNSC), 43 chromophobe renal cell carcinomas (KICH), 37 renal clear cell carcinomas from United States (KIRC), 31 renal papillary cell carcinomas (KIRP), 18 low-grade gliomas (LGG), 51 liver cancers from United States (LIHC), 67 liver cancers from Japan (LIRI), 37 lung adenocarcinomas (LUAD), 47 lung squamous cell carcinomas (LUSC), 95 malignant lymphomas (MALY), 80 ovarian cancers (OV), 74 pancreatic cancers (PACA), 19 prostate adenocarcinomas (PRAD), 49 renal clear cell carcinomas from European Union/France (RECA), 34 sarcomas (SARC), 34 skin cutaneous melanomas (SKCM), 29 stomach adenocarcinomas (STAD), 47 thyroid cancers (THCA), and 42 uterine corpus endometrial carcinomas (UCEC). More detailed information on the sample distribution and annotation can be found in **Supplementary Table S1**.

WGS and RNA-Seq data analysis of tumor and normal samples were performed by the PCAWG consortium as previously described [21]. Somatic and germline SNVs, somatic CNVs, SVs, and tumor purity were detected by multiple algorithms and consensus calls were made. Genome coordinates were based on the hg19 reference genome and GENCODE v19 was used for gene annotation. Gene expression was quantified by HT-Seq (version 0.6.1p1) as fragments per kilobase of million mapped (FPKM). Clinical data such as donor age and sex were downloaded from the PCAWG data portal (https://dcc.icgc.org/pcawg). *TOB1* and *TOB1-AS1* expression data in CCLE pancreatic cancer cell lines were downloaded from DepMap Public 22Q2 version (https://depmap.org/portal/download/all/). Gene expression data of the Cancer Genome Atlas (TCGA) PAAD cohort (TCGA.PAAD.sampleMap/HiSeqV2_PANCAN) and International Cancer Genome Consortium (ICGC) PACA-CA cohort for 45 samples of which "analysis-id" were labeled as "RNA" were downloaded from Xena Data Hubs (https://xenabrowser.net/datapages/) and ICGC data portal (https://dcc.icgc.org/projects/PACA-CA) respectively.

Significant eQTL-gene pairs (v8) were downloaded from the Genotype-Tissue Expression (GTEx) data portal (https://gtexportal.org/home/datasets). Only those eQTLs that had a hg19 liftover variant ID were included in the analysis and hg38 variants with no corresponding hg19 annotation were discarded.

The raw sequencing data for Hi-C and ATAC-Seq were available through NCBI Sequence Read Archive (SRA) with accession number PRJNA1036282. The raw sequencing data for mouse xenograft tumor RNA-Seq were available through NCBI SRA with accession number PRJNA1011356.

HYENA algorithm

First, small tandem duplications (<10 kb) were discarded since they are unlikely to produce new promoter-enhancer interactions. The remaining SVs were mapped to the flanking regions (500 kb upstream and downstream of transcription start sites [TSSs]) of annotated genes. SVs that fall entirely within a gene body were also discarded. The SV status of each gene was defined by the presence or absence of SV breakpoints within the gene or its flanking regions for each tumor. The binary variable SV status was used in the normal-score regression model below. Only genes carrying SVs in at least 5% samples carrying SVs were tested. For each gene, samples with that gene highly amplified (>10 copies) were removed from the regression model.

Gene expression normal scores

Gene expression quantifications (fragments per kilobase per million [FPKM]) were quantile normalized (FPKM-QN) using the *quantile.normalize()* function from the *preprocessCore* R package to enhance cross-sample comparison. To break the ties for genes with identical FPKM-QN values in multiple samples (especially those caused by FPKM of zero) during ranking, very small Gaussian noises were added to all the FPKM-QN values in all samples by *add.Gaussian.noise(mat, mean = 0.000000001, stddev = 0.000000001, symm = F)* from the *RMThreshold* R package. Since the mean and standard deviation of the noises added were small, the rankings of the non-identical values were not affected. For each gene, samples were ranked based on their noised-added expression values, the ranks were mapped to a standard normal distribution and the corresponding z scores were gene expression normal scores. Normal-score conversion forced the expression data into a Gaussian distribution, allowing for parametric comparisons between samples.

Normal-score regression

A generalized linear model was used to test associations between gene expression normal scores and SV status and control for confounding variables such as gene copy number, tumor sample purity, donor age, and sex. To capture unobserved variations in gene expression, the first n principal components (PCs) of the expression data were also included in the regression model, where n was determined as 10% of the sample size of the cohort and up to 20 if the sample size was more than 200. The regression model was as shown below:

Expression_normal_score ~ sv_status + copy_number + purity + age + sex + PC_1 + PC_2 …+ PC_n

For each gene, all PCs were tested for associations with the SV status of that gene, and those PCs that significantly correlate (Mann-Whitney test, $P<0.05$) with SV status were not used in regression.

Calculating empirical P values and model selection

Gene expression data were permuted 100 times by randomly shuffling expression values within the cohort. The normal-score regression was performed in the same way on observed gene expression and permuted expression. *P* values for SV status from permuted expression were

pooled as a null distribution. Then the P values for SV status from observed expression and the
P-value null distribution were used to calculate empirical P values. One-sided P values were
used since we were only interested in elevated gene expression. False discovery rates (FDRs)
were calculated using the Benjamini-Hochberg procedure. Genes with FDR less than 0.1 were
considered candidate genes. For example, in MALY, there were 1,863 genes reaching 5% SV
frequency and 1,863 P values were obtained in each permutation. After 100 permutations,
186,300 P values were generated and should represent the null distribution very well. Empirical
P values were calculated using these 186,300 permuted P values. To test whether more
permutations could be beneficial, we performed 1000 permutations in five benchmarking tumor
types (COAD/READ, KICH, LUSC, MALY, and THCA). A total of 44 candidate genes were
detected in 100 permutations. Four more genes were detected in 1000 permutations and two
genes detected in 100 permutations were missed in 1000 permutations. The FDRs for the shared
candidate genes from 100 and 1000 permutations were nearly identical (**Supplementary Fig.**
S13). Therefore, 100 permutations were sufficient.

The above empirical P value calculation and candidate gene detection were performed iteratively
with no PCs and up to n PCs in the regression model. When different numbers of PCs were
included in the model, the numbers of candidate genes varied. The regression model with the
lowest number of PCs reaching 80% of the maximum number of candidate genes in all
regression models tested was selected as the final model to avoid over fitting. For example, the
sample size for PCAWG BRCA was 88; therefore, we tested from 0 to 9 PCs. Among these, the
model including 8 PCs gave the highest number (82) of candidate genes. Therefore, the model
including 7 PCs with 68 candidate genes was selected as the final model since it had the lowest
number of PCs reaching 80% of 82 candidate genes (**Supplementary Table S8**).

In our normal-score regression, we essentially attempt to model variations in gene expression.
Including confounding factors will improve performance. Tumor purity, gene copy number,
patient age, and sex are factors known to affect gene expression. Therefore, they are included in
the regression model. Unobserved variations may include tumor subtype, tumor stage, patient
ethnicity, smoking status, alcohol consumption, and other unknown factors that may alter gene
expression. Since HYENA is designed for wide applications, we do not require users to provide
information on tumor subtype, tumor stage, patient ethnicity, smoking status, alcohol
consumption, etc. Principle component analysis is a linear decomposition of gene expression
variations. Therefore, including PCs in a regression model is suitable for removing systematic
variations and can better model the effects of SV status. However, some enhancer hijacking
target genes are master transcription factors, such as *MYC*, and have profound impact on gene
expression of multiple pathways. Hence, it is possible that some PCs capture the activities of
transcription factors. If these transcription factors are activated by somatic SVs, the PCs will be
correlated with SV status. Including these PCs will diminish our ability to detect the effects of
SV status. Therefore, we do not include these PCs in the regression model.

Testing eQTL-SV associations

Known germline eQTLs from the matching tissues were obtained from GTEx (**Supplementary**
Table S9). The associations between germline genotypes of eQTLs and SV status of the 213

candidate genes in the PCAWG cohort were tested using a Chi-squared test. Genes with
significant correlations ($P<0.05$) between their SV status and at least one eQTL were removed.
The remaining genes were our final candidate enhancer-hijacking target genes.

Benchmarking

Known enhancer hijacking target genes in PCAWG tumor types were selected to test the
sensitivity of HYENA, CESAM and PANGEA. The genes included *MYC* in malignant
lymphoma, *BCL2* in malignant lymphoma, *CCNE1* in stomach/gastric adenocarcinoma, *TERT* in
chromophobe renal carcinoma, *IGF2* in colorectal cancer, *IGF2* in stomach/gastric
adenocarcinoma, *IGF2BP3* in thyroid cancer, and *IRS4* in lung squamous cell carcinoma. The
same SVs, CNVs, and SNVs were used as input for all three algorithms. For CESAM and
PANGEA, upper-quantile normalized fragments per kilobase per million (FPKM-UQ) were
normalized by tumor purity and gene copy number, and then used as gene expression inputs.
CESAM was run using default parameters, and FDR of 0.1 was used to select significant genes.
PANGEA requires predicted enhancer-promoter (EP) interactions based on ChIP-Seq and RNA-
Seq data. The EP interactions were downloaded from EnhancerAtlas 2.0
(http://www.enhanceratlas.org/) (**Supplementary Table S10**). EP interactions from multiple cell
lines of the same type were merged. PANGEA was run with default parameters as well and
significant genes were provided by PANGEA (multiple testing adjusted *P* value <0.05). To test
false positives for HYENA, CESAM, and PANGEA, 20 random gene expression datasets for
malignant lymphoma and breast cancer were generated by randomly shuffling sample IDs in
gene expression data. HYENA, CESAM, and PANGEA were run with random expression in the
same way as above.

Predicting 3D genome organization

A 1 Mb sequence was extracted from the reference genome centered at each somatic SV
breakpoint and was used as input for Akita [38] to predict the 3D genome organization. Two 500
kb sequences were merged according to the SV orientation to construct the sequence of the
rearranged genome fragments. Akita was used to predict the genome organization for the
rearranged sequence. High-resolution Micro-C data obtained from human H1-ESCs and HFF
cells [53] were used to facilitate TAD annotation together with predicted genome organization.
H3K27Ac and CTCF ChIP-Seq data from the PANC-1 cell line were downloaded from the
ENCODE data portal (https://www.encodeproject.org/). SV breakpoints were provided to Orca [44]
to predict 3D genome structures through its web interface (https://orca.zhoulab.io/).

In situ Hi-C and ATAC-Seq

539 Ten million cells of Panc 10.05, PANC-1, PATU-8988S, and PATU-8988T cell lines were
540 collected to construct Hi-C libraries [39]. The Hi-C libraries were sequenced on Illumina NovaSeq

X Plus platform with 1% phix. About 2 billion reads were obtained from Panc 10.05, PATU-8988S, and PATU-8988T, and 1 billion reads were obtained from PANC-1. The paired-end reads were aligned to chromosomes 1-22, X, Y and M by bwa-mem. SVs were identified by EagleC [45] at 5 kb, 10 kb and 50 kb resolutions. The non-redundant SVs in **Supplementary Table S6** were combined for the three resolutions. Chromatin loops were identified by NeoLoopFinder [20]. A probability threshold of 0.95 was used, and default values were used for all other parameters. Fifty thousand cells of Panc 10.05, PATU-8988S, and PATU-8988T cell lines were harvested to construct ATAC-Seq libraries [54]. The libraries were sequenced using Illumina NovaSeq. About 60 million reads were generated from each library. The paired-end reads were aligned to the reference genome by hisat2. Hi-C and ATAC-Seq read coverages were generated by deepTools with 10 bp bin-size, RPGC normalization, and an effective genome size of 2,864,785,220.

Cell lines

HEK293T, PANC-1, and PATU-8988T cells were obtained from Dr. Alexander Muir (University of Chicago). Panc 10.05 was purchased from ATCC (American Type Culture Collection, USA) (https://www.atcc.org/products/crl-2547) and PATU-8988S was purchased from DSMZ (https://www.dsmz.de/collection/catalogue/details/culture/ACC-204). All cell lines were cultured at 37°C/5% CO_2. HEK293T cells and PANC-1 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) (Gibco, 21041025) containing 10% fetal bovine serum (FBS) (Gibco, A4766), and Panc 10.05 cells were cultured in RPMI-1640 medium (Gibco, 11875093) containing 10% FBS, as per ATCC instructions (https://www.atcc.org/products/crl-3216, https://www.atcc.org/products/crl-1469, https://www.atcc.org/products/crl-2547). PATU-8988T and PATU-8988S cells were cultured with DMEM containing 5% FBS, 5% horse serum (Gibco, 26050088), and 2 mM L-glutamine as recommended by DSMZ (Deutsche Sammlung von Mikroorganismen and Zellkulturen, Germany) (https://www.dsmz.de/collection/catalogue/details/culture/ACC-162). All cell lines have been regularly monitored and tested negative for mycoplasma using a mycoplasma detection kit (Lonza, LT07-218).

TOB1-AS1 **and luciferase overexpression**

A 1,351 bp *TOB1-AS1* cDNA (ENST00000416263.3) was synthesized by GenScript (New Jersey, USA) and subcloned into the lentiviral pCDH-CMV-MCS-EF1-Puro plasmid (SBI, CD510B-1). The cDNA sequence in the plasmid was verified by Sanger sequencing at University of Chicago Medicine Comprehensive Cancer Center core facility. The *TOB1-AS1* overexpression plasmid was amplified by transforming Stellar™ Competent Cells (Takara, 636763) with the plasmid as per instructions and isolated by QIAGEN HiSpeed Plasmid Midi Kit (QIAGEN, 12643). LucOS-Blast vector was obtained from Dr. Yuxuan Phoenix Miao (University of Chicago), cloned, and amplified as described above.

HEK293T cells were plated in T-25 flasks and grown to 75% confluence prior to transfection.
For each T-25 flask, 240μl Opti-MEM (Gibco, 31985070), 1.6μg pCMV-VSV-G, 2.56μg
pMDLg/pRRE, 2.56μg pRSV-Rev, 3.4μg *TOB1-AS1* overexpression vector and 22.8μl TransIT-
LT1 Transfection Reagent (Mirus, MIR 2306) were mixed and incubated at room temperature
for 30 minutes, then added to the plated HEK293T cells with fresh medium. The luciferase
vector was packaged into lentivirus with the same method. Upon 48 hours of incubation,
lentiviral supernatant was collected, filtered through 0.45-μmpolyvinylidene difluoride filter
(Millipore), and mixed with 8μg/ml polybrene. PANC-1 or PATU-8988T cells at 60%
confluence were transduced with the lentiviral supernatant for 48 hours followed by three rounds
of antibiotic selection with 4μg/ml puromycin for *TOB1-AS1* overexpression and 10μg/ml
blasticidin for the luciferase expression. *TOB1-AS1* expression was validated by quantitative
reverse transcription polymerase chain reaction (qRT-PCR), and luciferase expression was
validated by in vitro bioluminescence imaging in black wall 96-well plates (Corning, 3603). D-
luciferin potassium salt (Goldbio, LUCK-100) solution with 0, 1.25, 2.5, 5 and 10μl 15mg/ml
was added into the wells as serial dilutions, and imaging was obtained after 5 minutes. Finally,
TOB1-AS1 overexpression or empty pCDH transduced cell lines with luciferase co-expression
were built for both PATU-8988T and PANC-1 cells.

***TOB1-AS1* transient knock-down using antisense oligonucleotides (ASOs)**

Three Affinity Plus® ASOs were synthesized by Integrated DNA Technologies (IDT), with two
targeting *TOB1-AS1* and one non-targeting negative control. The ASO sequences were:
Non-targeting ASO (NC): 5' -GGCTACTACGCCGTCA- 3'
TOB1-AS1 ASO1: 5' -GCCGATTTGGTAGCTA- 3'
TOB1-AS1 ASO2: 5' -CTGCGGTTTAACTTCC- 3'
The ASOs were transfected into PATU-8988S and Panc 10.05 cells with Lipofecatmine^{TM} 2000
(Invitrogen, 11668019) using reverse-transfection method according to IDT protocol
(https://www.idtdna.com/pages/products/functional-genomics/antisense-oligos) with a final ASO
concentration of 9 nM. Cells were transfected in 6-well plates and incubated for 48 hours to
reach 60% confluence before RNA extraction or Transwell assay.

RNA isolation and qRT-PCR

Cells were plated in 6-well plates and allowed to reach 80% confluence, or transfected by ASOs
as described above, prior to RNA extraction. After cells lysis in 300μl/well TRYzol^{TM}
(Invitrogen, 15596026), RNA samples were prepared following the Direct-zol RNA Miniprep kit
manual (RPI, ZR2052). Reverse transcription was performed using Applied Biosystems High-
Capacity cDNA Reverse Transcription Kit (43-688-14) following manufacturer's instructions.
Quantitative PCR (qPCR) was conducted on StepOnePlus Real-Time PCR System (Applied
Biosystems, 4376600), using PowerUp SYBR Green Master Mix (A25742) following the

manufacturer's instructions with a primer concentration of 300nM in 10μl reaction systems.
Primers were ordered from Integrated DNA Technologies. Primer sequences used in this study
are as follows:
TOB1 forward: 5' -GGCACTGGTATCCTG AAA AGCC- 3'
TOB1 reverse: 5' – GTGGCAGATTGCCACGAACATC- 3'
TOB1-AS1 forward: 5' -GGAGTGGTCAGGTGACTGATT- 3'
TOB1-AS1 reverse: 5' -ATTCCACTCCTGTTTGCAACT- 3'
GAPDH forward: 5' – ACCACAGTCCATGCCATCAC- 3'
GAPDH reverse: 5' -TCCACCACCCTGTTGCTGTA- 3'
Relative expression levels for *TOB1-AS1* and *TOB1* were calculated by the $2^{(-\Delta\Delta C_T)}$ method
based on *GAPDH* expression as an endogenous control.
628

Transwell assay for cell invasion in vitro

Transparent PET membrane culture inserts of 24-well plate (Falcon, 353097) were coated with
Cultrex Reduced Growth Factor Basement Membrane Extract (BME) (R&D Systems, 3533-010-
02) at 50μg per membrane (200μl of 0.25mg/ml BME stock per membrane) at 37°C for an hour.
A total of 100,000 PANC-1 cells/well, 50,000 PATU-8988T cells/well, 50,000 Panc 10.05
cells/well, or 50,000 PATU-8988S cells were resuspended in serum-free, phenol-red free DMEM
medium and seeded into the coated inserts. Phenol-red free DMEM of 500μl (Gibco, A1443001)
with 10% FBS was added to the bottom of the wells and the cells were allowed to invade for 16
hours. Additional wells with 500μl serum-free, phenol-red free DMEM medium without FBS in
the bottom chamber were seeded with the same number of cells as indicated above as a negative
control. At the end of the assay, the membranes were stained with 500μl 4μg/ml Calcein AM
(CaAM) (Corning, 354216) for one hour at 37°C. The cells that failed to invade were removed
from the top chamber with a cotton swab and all inserts were transferred into 1x Cell
Dissociation Solution (Bio-Techne, 3455-05-03) and shaken at 150rpm for an hour at 37°C.
Finally, CaAM signal from the invaded cells was measured by a plate reader (Perkin Elmer
Victor X3) at 465/535nm.

Tumor metastasis in vivo

All animal experiments for this study were approved by the University of Chicago Institutional
Animal Care and Use Committee (IACUC) prior to execution. Male NSG mice were ordered
from the Jackson Laboratory (strain#005557). For tail vein inoculation, mice were injected
intravenously through the tail vein with luciferase-expressing at 400,000 cells/mouse for PANC-
1 cells in cold phosphate buffered saline (PBS) (Gibco, 10010-023). For orthotopic inoculation,
mice were injected with 200,000 PANC-1 cells/mouse into the pancreas under general
anesthesia. Cells were resuspended in cold PBS containing 5.6mg/mL Cultrex Reduced Growth
Factor BME (R&D Systems, 3533-010-02). Primary tumor and metastatic tumor burdens were
measured weekly for 4 and 6 weeks for tail vein injection models and orthotopic models,

respectively, via bioluminescence imaging using Xenogen IVIS 200 Imaging System
(PerkinElmer) at the University of Chicago Integrated Small Animal Imaging Research Resource
(iSAIRR) Facility. Each mouse was weighed and injected intra-peritoneally with D-luciferin
solution at a concentration of 150µg/g of body weight 14 minutes prior to image scanning ventral
side up.

Ex vivo IVIS imaging

Ex vivo imaging was done for the PANC-1 orthotopic injection mice after 8 weeks of orthotopic
inoculation. Mice were injected intra-peritoneally with D-luciferin solution at a concentration of
150µg/g of body weight immediately before euthanasia. Immediately after necropsy, mice were
dissected, and tissues of interest (primary tumors, livers and spleens) were placed into individual
wells of 6-well plates covered with 300 µg/mL D-luciferin. Tissues were imaged using Xenogen
IVIS 200 Imaging System (PerkinElmer) and analysis was performed (Living Image Software,
PerkinElmer) maintaining the regions of interest (ROIs) over the tissues as a constant size.

Tumor RNA sequencing and gene expression analysis

RNA was isolated from mouse subcutaneous tumors (six *TOB1-AS1* overexpression and six
control mice) after 6 weeks of PANC-1 cell subcutaneous injection using Direct-zol RNA
Miniprep kit (RPI, ZR2052). Quality and quantity of the RNA was assessed using Qubit.
Sequencing was performed using the Illumina NovaSeq 6000. About 40 million reads were
sequenced per sample. The pair-end reads were aligned to mouse genome (mm10) and human
genome (hg19) with hisat2, and the reads mapped to mouse or human genomes were
disambiguated using AstraZeneca-NGS disambiguate package. Gene counts were generated with
htseq-count. Differential gene expression was analyzed using DESeq2. Differentially expressed
genes were defined as genes with a FDR smaller than 0.1 and a fold change greater than 1.5.

Code availability

683 The HYENA package is available at https://github.com/yanglab-
684 computationalgenomics/HYENA.

Acknowledgements

We thank the Center for Research Informatics at the University of Chicago for providing the
computing infrastructure, Matthew Stephens for helpful suggestions, Marsha Rosner for
assistance in lentiviral experiments, and Ani Solanki for assistance in animal experiments. The
work was supported by the Goldblatt Endowment (A.Y.), the National Institutes of Health grant
K22CA193848 (L.Y.), R01CA269977 (L.Y.) and University of Chicago and UChicago
Comprehensive Cancer Center (L.Y.).

Disclosure

The authors have no competing interests to declare.

 Figure Legends

Figure 1. Outline of enhancer hijacking and HYENA algorithm. a, Mechanisms of gene
activation by SVs. SVs can activate genes by recruiting distal active enhancers (top panel) and
by removing TAD boundaries and forming de novo enhancer-promoter interactions (bottom
panel). **b**, HYENA workflow. Green and purple boxes denote input and output files, respectively.
Orange boxes denote intermediate steps. Numbers in parentheses represent default values of
HYENA.

Figure 2. Benchmarking HYENA. a, Comparison of HYENA, CESAM, and PANGEA in
detecting oncogenes known to be activated by enhancer hijacking in six tumor types from the
PCAWG cohort. **b**, UPSET plot demonstrating candidate genes identified and shared among the
three tools in five tumor types of PCAWG. The numbers of candidate genes predicted by three
algorithms are shown on the bottom left (19, 25, and 255). On the bottom right, individual dots
denote genes detected by one tool, and dots connected by lines denote genes detected by multiple
tools. The numbers of genes detected are shown above the dots and lines. For example, the dot
immediately on the right of "PANGEA" shows there are 254 candidate genes detected only by
PANGEA but not CESAM and HYENA. The left most line connecting two dots indicates that
there are six genes detected by both CESAM and HYENA but not by PANGEA. **c**, Number of
genes detected by HYENA in two PCAWG tumor types using observed gene expression and
randomized expression. Genes detected in random expression datasets are false positives.

Figure 3. Enhancer hijacking candidate genes in PCAWG. a, Candidate genes detected by
HYENA in individual tumor types of PCAWG. *TERT* is plotted twice since it is detected in two
cancer types. Genes labelled as red are known enhancer hijacking targets. **b**, Diverse types of
candidate genes identified by HYENA in PCAWG. Numbers after tumor type names denote
sample size in the corresponding tumor types.

Figure 4. *TOB1-AS1* activated by various types of SVs in pancreatic cancer. a, Normalized
expression of *TOB1-AS1* in samples with (n=7) and without (n=67) nearby SVs in pancreatic
cancers. The boxplot shows median values (thick black lines), upper and lower quartiles (boxes),
and 1.5× interquartile range (whiskers). Individual tumors are shown as black dots. **b**, Circos plot
summarizing intrachromosomal SVs (blue, n=5) and translocations (red, n=3) near *TOB1-AS1*. **c**,
Diagrams depicting putative enhancer hijacking mechanisms that activate *TOB1-AS1* in one
tumor with a 17:19 translocation (left panel) and another tumor with a large deletion (right
panel). **d**, Predicted 3D chromatin interaction maps of *TOB1-AS1* (left panel), *UQCRFS1*
(middle panel), and the translocated region in tumor 9ebac79d-8b38-4469-837e-b834725fe6d5
(right panel). The downstream fragment of the chromosome 19 SV breakpoint was flipped in
orientation and linked to chromosome 17. H3K27Ac and CTCF ChIP-Seq data of PANC-1 cell
line are shown at the bottom. The expected level of 3D contacts depends on linear distance
between two genomic locations. Longer distances correlate with fewer contacts. Akita predicts
3D contacts based on DNA sequences. The heatmaps are showing the ratio between predicted
and expected contacts. The darkest red represent regions having 100 times more contacts than
expected given the distance between the regions. **e**, Predicted 3D chromatin interaction maps of
TOB1-AS1 (left panel) and *KCNJ2* (middle panel) loci without deletion as well as the same
region following deletion in tumor 748d3ff3-8699-4519-8e0f-26b6a0581bff (right panel).

Figure 5. 3D genome structures in the *TOB1-AS1* locus in pancreatic cancer cell lines. a,
TOB1-AS1 expression in pancreatic cancer cell lines in CCLE. The cell lines in red are selected
for further studies. **b** and **c**, 3D genomic interactions in four pancreatic cancer cell lines. Black
arrows represent SVs with off-diagonal interactions. The locations of *TOB1-AS1* are marked by
blue lines. In Panc 10.05, the blue arrow points to the CNV breakpoint and the dashed blue
triangle represents the neo-subdomain formed due to the foldback inversion. **d**, The reference
chromosome 17 and derived chromosomes in Panc 10.05. The chromosomes are not to scale.
TOB1-AS1 is shown as small blue boxes in the chromosomes. **e**, Open chromatin measured by
ATAC-Seq in PATU-8988S and PATU-8988T at the *TOB1-AS1* locus.

Figure 6. *TOB1-AS1* promotes cell invasion and tumor metastasis. a, *TOB1-AS1* and *TOB1*
relative expression levels in PATU-8988T and PANC-1 cells transduced with *TOB1-AS1*
overexpression vector (n=3) or control vector (n=3). **b**, *TOB1-AS1* overexpression in PATU-
8988T (4 biological replicates) and PANC-1 (3 biological replicates) promoted in vitro cell
invasion using Transwell assay. Each biological replicate was an independent experiment with 7
technical replicates per experimental group. The average fold change of cell invasion was
calculated after the background invasion measured in the absence of any chemotactic agent was
subtracted from each technical replicate. *P* values were calculated by two-sided student t test. **c**,
TOB1-AS1 overexpression in PANC-1 cells promoted in vivo tumor metastasis in the tail vein
injection model. **d**, *TOB1-AS1* overexpression in PANC-1 cells exacerbated in vivo tumor
growth and spontaneous metastasis in the orthotopic tumor model. Images of radiance in
immunodeficient mice are shown on the left while the quantifications of radiance are shown on
the right. Eight mice were used in both overexpression group and the empty vector control. The
images were analyzed by setting the regions of interest (ROIs) to mouse torsos and measuring
the average radiance level (in p/sec/cm^2/sr). **e**, Primary tumor burden and spleen metastatic
burden were higher in the mice that were orthotopically injected with *TOB1-AS1* overexpression
PANC-1 cells. The bar plots show quantified total radiance with a set area (in p/sec). **f**, Targeting
TOB1-AS1 by two ASOs. **g**, *TOB1-AS1* knockdown in Panc 10.05 and PATU-8988S cells
transduced with ASO1 (n=3), ASO2 (n=3) or non-targeting control ASO (NC) (n=3). **h**, *TOB1-
AS1* knockdown suppressed Panc 10.05 (3 biological replicates) and PATU-8988S (3 biological
replicates) cell invasion in vitro. Cell invasion fold change calculation is the same as in **b**. Two-
sided student t test was used. Error bars in all panels indicate standard error of the mean.

References

1. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of
chromatin interactions. *Nature* **485**, 376–380 (2012).

2. Krijger, P. H. L. & De Laat, W. Regulation of disease-associated gene expression in the
3D genome. *Nat. Rev. Mol. Cell Biol.* **17**, 771–782 (2016).

3. Symmons, O. *et al.* Functional and topological characteristics of mammalian regulatory
domains. *Genome Res.* **24**, 390–400 (2014).

4. Lupiáñez, D. G. *et al.* Disruptions of topological chromatin domains cause pathogenic
rewiring of gene-enhancer interactions. *Cell* **161**, 1012–1025 (2015).

5. Zhang, W. *et al.* A global transcriptional network connecting noncoding mutations to
changes in tumor gene expression. *Nat. Genet. 2018 504* **50**, 613–620 (2018).

6. Weischenfeldt, J., Symmons, O., Spitz, F. & Korbel, J. O. Phenotypic impact of genomic
structural variation: insights from and for human disease. *Nat. Rev. Genet.* **14**, 125–138
(2013).

7. Davis, C. F. *et al.* The somatic genomic landscape of chromophobe renal cell carcinoma.
Cancer Cell **26**, 319–330 (2014).

8. Bakhshi, A. *et al.* Cloning the chromosomal breakpoint of t(14;18) human lymphomas:
clustering around Jh on chromosome 14 and near a transcriptional unit on 18. *Cell* **41**,
899–906 (1985).

9. Gostissa, M. *et al.* Long-range oncogenic activation of Igh-c-myc translocations by the
Igh 3′ regulatory region. *Nature* **462**, 803–807 (2009).

10. Hnisz, D. *et al.* Activation of proto-oncogenes by disruption of chromosome
neighborhoods. *Science* **351**, 1454–1458 (2016).

11. Gröschel, S. *et al.* A single oncogenic enhancer rearrangement causes concomitant EVI1
and GATA2 deregulation in leukemia. *Cell* **157**, 369–381 (2014).

12. Northcott, P. A. *et al.* Enhancer hijacking activates GFI1 family oncogenes in
medulloblastoma. *Nature* **511**, 428–434 (2014).

13. Weischenfeldt, J. *et al.* Pan-cancer analysis of somatic copy-number alterations implicates
IRS4 and IGF2 in enhancer hijacking. *Nat. Genet.* **49**, 65–74 (2016).

14. Northcott, P. A. *et al.* The whole-genome landscape of medulloblastoma subtypes. *Nature*
547, 311–317 (2017).

15. He, B. *et al.* Diverse noncoding mutations contribute to deregulation of cis-regulatory
landscape in pediatric cancers. *Sci. Adv.* **6**, eaba3064 (2020).

16. Kopp, F. & Mendell, J. T. Functional Classification and Experimental Dissection of Long
Noncoding RNAs. *Cell* **172**, 393–407 (2018).

17. Lin, C.-P. & He, L. Noncoding RNAs in Cancer Development. *Annu. Rev. Cancer Biol.* **1**,
163–184 (2017).

18. Liu, S. J., Dang, H. X., Lim, D. A., Feng, F. Y. & Maher, C. A. Long noncoding RNAs in
cancer metastasis. *Nat. Rev. Cancer 2021 217* **21**, 446–460 (2021).

19. Liu, Y. *et al.* Discovery of regulatory noncoding variants in individual cancer genomes by
using cis-X. *Nat. Genet.* **52**, 811–818 (2020).

20. Wang, X. *et al.* Genome-wide detection of enhancer-hijacking events from chromatin
interaction data in rearranged genomes. *Nat. Methods* **18**, 661–668 (2021).

21. Campbell, P. J. *et al.* Pan-cancer analysis of whole genomes. *Nature* **578**, 82–93 (2020).

22. Beroukhim, R. *et al.* The landscape of somatic copy-number alteration across human
cancers. *Nature* **463**, 899–905 (2010).

23. Uhlen, M. *et al.* A pathology atlas of the human cancer transcriptome. *Science* **357**,
eaan2507 (2017).

24. Zhang, X. *et al.* Identification of focally amplified lineage-specific super-enhancers in
human epithelial cancers. *Nat. Genet.* **48**, 176–182 (2015).

25. Takeda, D. Y. *et al.* A Somatically Acquired Enhancer of the Androgen Receptor Is a
Noncoding Driver in Advanced Prostate Cancer. *Cell* **174**, 422-432.e13 (2018).

26. Turner, K. M. *et al.* Extrachromosomal oncogene amplification drives tumour evolution
and genetic heterogeneity. *Nature* **543**, 122–125 (2017).

27. Wu, S. *et al.* Circular ecDNA promotes accessible chromatin and high oncogene
expression. *Nature* **575**, 699–703 (2019).

28. Morton, A. R. *et al.* Functional Enhancers Shape Extrachromosomal Oncogene
Amplifications. *Cell* **179**, 1330-1341.e13 (2019).

29. Li, Y., Ge, X., Peng, F., Li, W. & Li, J. J. Exaggerated false positives by popular
differential expression methods when analyzing human population samples. *Genome Biol.*
23, 79 (2022).

30. Ooi, W. F. *et al.* Integrated paired-end enhancer profiling and whole-genome sequencing
reveals recurrent CCNE1 and IGF2 enhancer hijacking in primary gastric
adenocarcinoma. *Gut* **69**, 1039–1052 (2020).

31. Yun, J. W. *et al.* Dysregulation of cancer genes by recurrent intergenic fusions. *Genome*
Biol. **21**, 166 (2020).

32. Richter, J. *et al.* Recurrent mutation of the ID3 gene in Burkitt lymphoma identified by
integrated genome, exome and transcriptome sequencing. *Nat. Genet.* **44**, 1316–1320
(2012).

33. Nik-Zainal, S. *et al.* Landscape of somatic mutations in 560 breast cancer whole-genome
sequences. *Nature* **534**, 47–54 (2016).

34. Li, Y. *et al.* Patterns of somatic structural variation in human cancer genomes. *Nature* **578**,
112–121 (2020).

35. Panebianco, F. *et al.* THADA fusion is a mechanism of IGF2BP3 activation and IGF1R

signaling in thyroid cancer. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 2307–2312 (2017).

36.   Zapatka, M. *et al.* The landscape of viral associations in human cancers. *Nat. Genet. 2020*
847   *523* **52**, 320–330 (2020).

848   37.   Neuveut, C., Wei, Y. & Buendia, M. A. Mechanisms of HBV-related
849   hepatocarcinogenesis. *J. Hepatol.* **52**, 594–604 (2010).

850   38.   Fudenberg, G., Kelley, D. R. & Pollard, K. S. Predicting 3D genome folding from DNA
851   sequence with Akita. *Nat. Methods* **17**, 1111–1117 (2020).

852   39.   Rao, S. S. P. *et al.* A 3D Map of the Human Genome at Kilobase Resolution Reveals
853   Principles of Chromatin Looping. *Cell* **159**, 1665–1680 (2014).

854   40.   Fudenberg, G. *et al.* Formation of Chromosomal Domains by Loop Extrusion. *Cell Rep.*
855   **15**, 2038–2049 (2016).

856   41.   Franke, M. *et al.* Formation of new chromatin domains determines pathogenicity of
857   genomic duplications. *Nature* **538**, 265–269 (2016).

858   42.   Melo, U. S. *et al.* Complete lung agenesis caused by complex genomic rearrangements
859   with neo-TAD formation at the SHH locus. *Hum. Genet.* **140**, 1459–1469 (2021).

860   43.   de Bruijn, S. E. *et al.* Structural Variants Create New Topological-Associated Domains
861   and Ectopic Retinal Enhancer-Gene Contact in Dominant Retinitis Pigmentosa. *Am. J.*
862   *Hum. Genet.* **107**, 802–814 (2020).

863   44.   Zhou, J. Sequence-based modeling of three-dimensional genome architecture from
864   kilobase to chromosome scale. *Nat. Genet.* **54**, 725–734 (2022).

865   45.   Wang, X., Luan, Y. & Yue, F. EagleC: A deep-learning framework for detecting a full
866   range of structural variations from bulk and single-cell contact maps. *Sci. Adv.* **8**, 9215
867   (2022).

868   46.   Li, Y. *et al.* Constitutional and somatic rearrangement of chromosome 21 in acute
869   lymphoblastic leukaemia. *Nature* **508**, 98–102 (2014).

870   47.   Maciejowski, J., Li, Y., Bosco, N., Campbell, P. J. & de Lange, T. Chromothripsis and
871   Kataegis Induced by Telomere Crisis. *Cell* **163**, 1641–1654 (2015).

872   48.   Dixon, J. R. *et al.* Integrative detection and analysis of structural variation in cancer
873   genomes. *Nat. Genet.* **50**, 1388–1398 (2018).

874   49.   Elsässer, H. P., Lehr, U., Agricola, B. & Kern, H. F. Establishment and characterisation of
875   two cell lines with different grade of differentiation derived from one primary human
876   pancreatic adenocarcinoma. *Virchows Arch. B. Cell Pathol. Incl. Mol. Pathol.* **61**, 295–
877   306 (1992).

878   50.   Yao, J. *et al.* Long noncoding RNA TOB1-AS1, an epigenetically silenced gene,
879   functioned as a novel tumor suppressor by sponging miR-27b in cervical cancer. *Am. J.*
880   *Cancer Res.* **8**, 1483 (2018).

881   51.   Shangguan, W. *et al.* TOB1-AS1 suppresses non-small cell lung cancer cell migration and

882   invasion through a ceRNA network. *Exp. Ther. Med.* **18**, (2019).

883   52.   Wang, C. Y. *et al.* Molecular cloning and characterization of a novel gene family of four
884   ancient conserved domain proteins (ACDP). *Gene* **306**, 37–44 (2003).

885   53.   Krietenstein, N. *et al.* Ultrastructural Details of Mammalian Chromosome Architecture.
886   *Mol. Cell* **78**, 554-565.e7 (2020).

887   54.   Grandi, F. C., Modi, H., Kampman, L. & Corces, M. R. Chromatin accessibility profiling
888   by ATAC-seq. *Nat. Protoc. 2022 176* **17**, 1518–1552 (2022).

a CCLE RNA-Seq

TOB1-AS1 expression level (TPM)

Pancreatic cancer cell lines: SUIT-2, Panc 08.13, DAN-G, PANC-1, SNU-324, SW 1990, PATU-8988T, Panc 03.27, AsPC-1, KCI-MOH1, PK-1, PACADD-119, L3.3, PACADD-137, Panc 02.13, PK-59, SNU-213, Panc 05.04, CFPAC-1, Hs 766T, PACADD-159, PK-8, SNU-410, BxPC-3, HPAC, T3M-4, PK-45H, MIA PaCa-2, PACADD-188, PACADD-165, 950-5-BIK, PSN1, Panc 04.03, PA-TU-8902, HPAF-II, HUP-T4, HUP-T3, Capan-2, TCC-PAN2, PACADD-135, KP4, QGP-1, KP-2, JOPACA-1, Panc 02.03, KP-3, Capan-1, PATU-8988S, SU.86.86, YAPC, Panc 10.05

b

Panc 10.05 *TOB1-AS1* high | PANC-1 *TOB1-AS1* low | PATU-8988S *TOB1-AS1* high | PATU-8988T *TOB1-AS1* low

⇆ SVs

chr17 45,000,000 — 50,000,000 TOB1-AS1

c

Panc 10.05 | PANC-1 | PATU-8988S | PATU-8988T

chr17 48,900,000 — 48,950,000 — 49,000,000 TOB1-AS1 — CNV breakpoint

Read cov

d

Ref chr17 — *TOB1-AS1* — CNV breakpoint
Copy number: 5 1 2

Panc 10.05:
WT chr
Foldback inversion
Foldback inversion
t(6,17)
→ neo-subdomain

e ATAC-Seq read coverage

PATU-8988S *TOB1-AS1* high
PATU-8988T *TOB1-AS1* low

chr17 48,938,000 48,941,000 48,944,000 48,947,000

TOB1 — TOB1-AS1