HYENA detects non-coding genes activated by distal enhancers in cancer

Ali E. Yesilkanal [1,*], Anqi Yu [1,*], Ashish Thakur [2], Fan Wang [1], Xiaoyang Wu [1,3], Alexander Muir [1,3], Xin He [2], Francois Spitz [2], Lixing Yang [1,2,3,**]

1. Ben May Department for Cancer Research, University of Chicago, Chicago IL, USA
2. Department of Human Genetics, University of Chicago, Chicago IL, USA
3. University of Chicago Comprehensive Cancer Center, Chicago, IL, USA

* these authors contributed equally

** correspondence: lixingyang@uchicago.edu

Abstract

Somatic structural variations (SVs) in cancer can shuffle DNA content in the genome, relocate regulatory elements, and alter genome organization. Enhancer hijacking occurs when SVs relocate distal enhancers to activate gene expression. However, most enhancer hijacking studies have only focused on protein-coding genes. Here, we develop a computational algorithm "HYENA" to identify candidate oncogenes (both protein-coding and non-coding) activated by enhancer hijacking based on tumor whole-genome and transcriptome sequencing data. HYENA detects genes whose elevated expression is associated with somatic SVs by using a rank-based regression model. We systematically analyze 1,148 tumors across 25 types of adult tumors and identify a total of 192 candidate oncogenes including many non-coding genes. A long non-coding RNA *TOB1-AS1* is activated by various types of SVs in 10% of pancreatic cancers through altered 3-dimension genome structure. We find that high expression of *TOB1-AS1* can promote cell invasion and metastasis. Our study highlights the contribution of genetic alterations in non-coding regions to tumorigenesis and tumor progression.

Introduction

At mega-base-pair scale, linear DNA is organized into topologically associating domains (TADs) [1], and gene expression is regulated by DNA and protein interactions governed by 3D genome organization. Enhancer-promoter interactions are mostly confined within TADs [2–4]. Structural variants (SVs), including deletions, duplications, inversions and translocations, can dramatically change the TAD organization and gene regulation [5], and subsequently contribute to tumorigenesis. Previously, we discovered that *TERT* is frequently activated in chromophobe renal cell carcinoma by the relocation of distal enhancers [6], a mechanism referred to as enhancer hijacking (**Fig. 1a**). In fact, many oncogenes, such as *BCL2* [7], *MYC* [8], *TAL1* [9], *MECOM/EVI1* [10], *GFI1* [11], *IGF2* [12], *PRDM6* [13], and *CHD4* [14], are activated through this mechanism. These examples demonstrate that genomic architecture plays an important role in pathogenesis. However, the vast majority of the known enhancer hijacking target genes are protein-coding genes, and few non-coding genes have been reported to promote diseases through enhancer hijacking. Non-coding genes are known to play important roles in many biological processes [15] and some are known to drive tumorigenesis [16,17]. In this study, we will focus on identifying oncogenic non-coding genes activated by enhancer hijacking.

Several existing algorithms can detect enhancer hijacking target genes based on patient cohorts, such as CESAM [12] and PANGEA [14]. These two algorithms perform linear regression and elastic net model (also based on linear regression) to associate elevated gene expression with nearby SVs, respectively. PANGEA also considers the effects of somatic single nucleotide variants (SNVs) on gene expression. However, a major drawback of these algorithms is that linear regression is quite sensitive to outliers. Outliers are very common in gene expression data from cancer samples and can seriously impair the performances of these algorithms. In addition, CESAM is optimized for microarray data, while PANGEA depends on annotation of tissue-specific promoter-enhancer pairs which are not readily available for many tumor types. Cis-X [18] and NeoLoopFinder [19] can detect enhancer hijacking target genes based on individual samples. However, these tools have limitations in detectable genes and input data. Cis-X detects *cis-*activated genes based on allele-specific expression, which requires the genes to carry heterozygous SNVs. NeoLoopFinder takes Hi-C, Chromatin Interaction Analysis with Paired-End Tag (ChIA-PET), or similar data measuring chromatin interactions as input. Furthermore, to identify oncogenes, recurrent events in patient cohorts will still be required. Finally, no non-coding genes have been reported as enhancer hijacking targets by the above algorithms. A recent study on SVs altering gene expression in Pan-Cancer Analysis of Whole Genomes (PCAWG) samples [20] only considered protein-coding genes but not non-coding genes.

Here, we developed Hijacking of Enhancer Activity (HYENA) using normal-score regression and permutation test to detect candidate enhancer hijacking genes (both protein-coding and non-coding genes) based on tumor whole-genome and transcriptome sequencing data from patient cohorts. HYENA identified a non-coding gene, *TOB1-AS1*, as a regulator of cancer cell invasion in vitro and metastasis in vivo.

Results

HYENA workflow

Conceptually, the SVs leading to elevated gene expression are expression quantitative trait loci (eQTLs). The variants are SVs instead of commonly used germline single nucleotide polymorphisms (SNPs) in eQTL analysis. With somatic SVs and gene expression measured from the same tumors through whole-genome sequencing (WGS) and RNA sequencing (RNAseq), we can identify enhancer hijacking target genes by eQTL analysis. However, the complexities of cancer and SVs pose many challenges. For instance, there is tremendous inter-tumor heterogeneity—no two tumors are identical at the molecular level. In addition, there is substantial intra-tumor heterogeneity as tumor tissues are always mixtures of tumor, stromal and immune cells. Moreover, genome instability is a hallmark of cancer, and gene dosages are frequently altered [21]. Furthermore, gene expression networks are widely rewired [22], and outliers of gene expression are common.

Here, we developed an algorithm HYENA to overcome the challenges described above (see more details in Methods Section). We used a gene-centric approach to search for elevated expression of genes correlated with the presence of nearby SVs (**Fig. 1b**). Although duplicated enhancers can upregulate genes [23,24], we do not consider these as enhancer hijacking events since no neo-promoter-enhancer interactions are established. However, small deletions can remove TAD boundaries or repressive elements and lead to neo-promoter-enhancer interactions (**Fig. 1a**). Therefore, small tandem duplications were discarded, and small deletions were retained. For each gene, we annotated SV status (presence or absence of nearby SVs) for all samples. Samples in which the testing genes were highly amplified were discarded since many of these genes are amplified by circular extrachromosomal DNA (ecDNA) [25] and ecDNA can promote accessible chromatin [26] with enhancer rewiring [27]. Only genes with nearby SVs in at least 5% of tumors were further considered. In contrast to CESAM and PANGEA, we did not use linear regression to model the relationships between SV status and gene expression because linear regression is sensitive to outliers and many false significant associations would be detected [28]. Instead, we used a rank-based normal-score regression approach. After quantile normalization of gene expression for both protein-coding and non-coding genes, we added small Gaussian noises to gene expression for tie breaking, ranked the genes based on quantile-normalized noise-added expression, and transformed the ranks to the quantiles of standard normal distribution. We used the z scores (normal scores) of the quantiles as dependent variables in regression. In the normal-score regression model, tumor purity, copy number of the testing gene, patient age, and sex were included as covariates since these factors confound gene expression. We also included gene expression principal components (PCs) that were not correlated with SV status to model unexplained variations in gene expression. To deduce a better null distribution, we permuted the gene expression 100 times and ran the same regression models. All *P* values from the permutations were pooled together and used as the null distribution to calculate empirical *P* values. Then, multiple test corrections were performed on one-sided *P* values since we are only interested in elevated gene expression under the influence of nearby SVs. Finally, genes were

108 discarded if their elevated expression could be explained by germline eQTLs. The remaining
109 genes were candidate enhancer hijacking target genes.

Benchmarking performances

There is no gold standard available to comprehensively evaluate the performance of HYENA.
Here, we used eight known enhancer hijacking target genes including *MYC* [8], *BCL2* [7], *CCNE1* [29],
TERT [6], *IGF2* [12,29], *IGF2BP3* [30] and *IRS4* [12] from six types of adult tumors profiled by PCAWG
[20] to assess the sensitivity of HYENA (**Supplementary Table S1**). The six tumor types were
malignant lymphoma (MALY), stomach/gastric cancer (STAD), chromophobe renal cell
carcinoma (KICH), colorectal cancer (COAD/READ), thyroid cancer (THCA) and lung
squamous cell carcinoma (LUSC). We also expect immunoglobulin genes to be detected as
enhancer hijacking candidates in malignant lymphoma due to V(D)J recombination since the
lymphomas in PCAWG are B-cell derived Burkitt lymphomas [31]. In B cells, V(D)J
recombination occurs to join different variable (V), joining (J) and constant (C) segments to
produce antibodies with a wide range of antigen recognition ability. Therefore, certain segments
have elevated expression, and the recombination events can be detected as somatic SVs. In
addition to HYENA, we also tested two other algorithms—CESAM and PANGEA. All three
algorithms were run on the same somatic SVs and gene expression data from the PCAWG
cohort. Note that PANGEA depends on promoter-enhancer interactions predicted from cell lines
which were not available for thyroid tissue. Therefore, thyroid cancer data were not analyzed by
PANGEA.

Out of the eight known genes, HYENA detected five, CESAM detected three and PANGEA did
not detect any (**Fig. 2a**). In the five tumor types analyzed by all three algorithms, HYENA
identified a total of 25 candidate genes, CESAM identified 19, whereas PANGEA identified 255
genes (**Fig. 2b, Supplementary Tables S2, S3 and S4**). Six genes were detected by both
HYENA and CESAM. However, PANGEA had little overlap with the other algorithms (**Fig.
2b**). Both *IGF2* and *IRS4* were initially discovered by CESAM as enhancer hijacking target
genes using copy number variation (CNV) breakpoints profiled by microarray with much larger
sample sizes (378 colorectal cancers and 497 lung squamous cell carcinomas) [12]. In PCAWG, the
sample sizes with both WGS and RNAseq were smaller (51 colorectal cancers and 47 lung
squamous cell carcinomas). HYENA detected *IGF2* in colorectal cancer but not *IRS4*, whereas
CESAM and PANGEA detected neither. In stomach/gastric cancer, *IGF2* and *CCNE1* were
identified as enhancer hijacking genes in a cohort of 208 samples [29]. Neither of these genes were
detected by any of the algorithms because there were only 29 stomach tumors in PCAWG.
Therefore, known genes missed by HYENA were likely due to small sample size. Of the 16
genes detected by HYENA in malignant lymphoma, there were two immunoglobulin light chain
genes from lambda cluster (*IGLC7* and *IGLJ7*) (**Supplementary Table S2**). CESAM detected
11 genes with one being immunoglobulin gene (*IGLC7*) (**Supplementary Table S3**). In contrast,
PANGEA detected 30 candidate genes, but none were immunoglobulin genes (**Supplementary
Table S4**). In summary, HYENA had the best sensitivity of the three algorithms.

To evaluate specificity of the algorithms, we generated 20 random datasets by shuffling gene
expression data in both malignant lymphoma and breast cancer and ran all three algorithms

again. Since these gene expression data were random, there should be no associations between
SVs and gene expression, and all genes detected should be false positives. In malignant
lymphoma with observed gene expression, HYENA, CESAM and PANGEA detected 16, 11 and
30 candidate genes respectively (**Supplementary Tables S2, S3 and S4**). In the 20 random gene
expression datasets, HYENA detected an average of 0.5 genes per dataset (**Fig. 2c**), and CESAM
detected 0.5 genes, whereas PANGEA detected 40 genes (**Supplementary Fig. S1**). In breast
cancer with observed gene expression, HYENA, CESAM and PANGEA detected 61, 9 and
2,309 candidate genes, respectively (**Supplementary Tables S2, S3 and S4**). In 20 random gene
expression datasets, HYENA, CESAM and PANGEA detected 0.35, 0.9 and 2,296 genes on
average (**Fig. 2c** and **Supplementary Fig. S1**). In both tumor types, the numbers of false
positives called by PANGEA in random datasets were comparable to the numbers of genes
detected with observed gene expression (**Supplementary Fig. S1**). In summary, HYENA
predicted the least number of false positives among the three algorithms.

Overall, HYENA has superior sensitivity and specificity in the detection of enhancer hijacking
candidate genes.

Enhancer hijacking candidate genes in PCAWG

We used HYENA to analyze a total of 1,146 tumors across 25 tumor types in PCAWG with both
WGS and RNAseq data. When each tumor type was analyzed individually, we identified 192
candidate enhancer hijacking genes in total (**Supplementary Tables S1** and **S2**), five of which
were known enhancer hijacking targets (**Fig. 3a**). *TERT* was the only gene identified in two
tumor types/cohorts (chromophobe renal cell carcinoma [KICH] from the US and renal cell
carcinoma [RECA] from Europe). Almost all genes were only detected in one tumor type,
highlighting high tumor type specificity of the findings. The number of genes detected in each
tumor type also differed dramatically (**Fig. 3b**). No genes were detected in bladder cancer
(BLCA), cervical cancers (CESC), glioblastoma multiforme (GBM), and low-grade gliomas
(LGG) probably due to their small sample sizes. Breast cancers (BRCA) had the greatest number
of candidate genes likely due to the large sample size as well as the abundance of SVs resulting
from homologous recombination deficiency (HRD) [32]. Although ovarian cancers (OV) also
suffer from HRD and their sample size was comparable with breast cancers, there were much
fewer enhancer hijacking genes detected. Thyroid cancer (THCA) genomes were among the
most stable genomes in PCAWG [33]. However, the 15 enhancer hijacking genes identified in
thyroid cancers exceeded the number of candidate genes in ovarian cancer as well as many other
tumor types. Among these 15 genes, *IGF2BP3* was a known oncogene activated by enhancer
hijacking [30,34]. Interestingly, there were two liver cancer cohorts with comparable sample sizes—
LIHC from the US and LIRI from Japan. A total of 18 genes were identified in the US cohort
whereas no genes were found in the Japanese cohort. One possible reason for such a drastic
difference could be that hepatitis B virus (HBV) infection is more common in liver cancers in
Japan [35], and virus integration in the tumor genome can result in oncogene activation [36]. In
Chronic Lymphocytic Leukemia (CLLE), a total of nine genes were detected, and seven were
immunoglobulin genes from both lambda and kappa clusters. Given that sample size and genome
instability can only explain a small fraction of variations of enhancer hijacking genes detected in

different tumor types, the landscape of enhancer hijacking in cancer seems to be mainly driven
by the underlying disease biology. Intriguingly, out of the 192 candidate genes, 73 (38%) were
non-coding genes including long non-coding RNAs (lncRNAs) and microRNAs (**Fig. 3b**).

Neo-TADs formed through somatic SVs

Next, we focus on the most frequently altered enhancer hijacking candidate non-coding gene in
pancreatic cancer *TOB1-AS1* (**Fig. 4a**), a long non-coding RNA (lncRNA). Seven (9.6%) out of
74 tumors had a variety of somatic SVs near *TOB1-AS1* including translocations, deletions,
inversions and tandem duplications (**Fig. 4b** and **Supplementary Table S5**). For example, tumor
9ebac79d-8b38-4469-837e-b834725fe6d5 had a translocation between chromosomes 17 and 19
(**Fig. 4c**). The breakpoints were upstream of *TOB1-AS1* and upstream of *UQCRFS1* (**Fig. 4d**). In
tumor 748d3ff3-8699-4519-8e0f-26b6a0581bff, there was a 19.3 Mb deletion which brought
TOB1-AS1 next to a region downstream of *KCNJ2* (**Fig. 4c** and **4e**).

We used Hi-C maps to investigate the genome organization of the loci in their normal context
and Akita [37], a convolutional neural network that predicts 3D genome organization, to estimate
the 3D architecture of the loci impacted by SVs. While 3D structures are dynamic and may
change with cell-type and gene activity, TAD boundaries are often more stable and remain
similar between different cell-types [1]. TAD boundaries are defined locally by the presence of
binding sites for CCCTC-binding factor (CTCF), a ubiquitously expressed DNA-binding protein
[1,38], and their formation arises from the stalling of the cohesin-extruded chromatin loop by CTCF
at these positions [39]. For this reason, one can reliably expect that upon chromosomal
rearrangement, new TADs will be formed as directed by the position of relocated TAD
boundaries. This assumption has been validated by direct experimental evidences examining the
"neo-TADs" associated with SVs at different loci [40–42]. The wildtype *TOB1-AS1* locus had a
TAD between a CTCF binding site in *RSAD1* and another one upstream of *SPAG9* (**Fig. 4d** and
Supplementary Fig. S2). There were TADs spanning *UQCRFS1* and downstream of *KCNJ2* in
the two partner regions (**Fig. 4d**, **4e** and **Supplementary Fig. S2**). In tumor 9ebac79d-8b38-
4469-837e-b834725fe6d5, the translocation was predicted to lead to a neo-TAD resulting from
merging the TADs of *TOB1-AS1* and *UQCRFS1* into one (**Fig. 4d**). In tumor 748d3ff3-8699-
4519-8e0f-26b6a0581bff, another neo-TAD was predicted to form as a result of the deletion that
merged the TADs of *TOB1-AS1* and the downstream portion of *KCNJ2* into one (**Fig. 4e**).
Within these predicted neo-TADs, Akita predicted new strong interactions involving several
CTCF binding sites across two SV partner regions (**Fig. 4d** and **4e** pink dots in the right panels).
Similar neo-TADs could be observed in two additional tumors (**Supplementary Fig. S3**). In the
two tumors with 317 kb and 226 kb tandem duplications, the *TOB1-AS1* TAD was expanded
(**Supplementary Fig. S4a**). However, in the last tumor, *TOB1-AS1* was predicted to remain in its
original TAD after translocation (**Supplementary Fig. S4b**). In summary, at least four out of the
seven tumors harboring somatic SVs near *TOB1-AS1* were predicted to result in neo-TADs
including *TOB1-AS1*.

These results demonstrated that HYENA can detect genes activated by reorganization of 3D
genome architecture.

Oncogenic functions of *TOB1-AS1*

TOB1-AS1 has been reported as a tumor suppressor in several tumor types [44,45]. However, it was predicted to be an oncogene by HYENA in pancreatic cancers. To test if *TOB1-AS1* has oncogenic functions in pancreatic cancer, we performed both in vitro and in vivo experiments. We surveyed pancreatic cancer cell line RNAseq data from Cancer Cell Line Encyclopedia (CCLE) and identified that the commonly transcribed isoform of *TOB1-AS1* was ENST00000416263.3 in pancreatic cancers (**Supplementary Fig. S5**). The synthesized *TOB1-AS1* cDNA was cloned and overexpressed in two pancreatic cancer cell lines, PANC-1 and PATU-8988T, both of which had low expression of *TOB1-AS1* (**Supplementary Fig. S6**). In both cell lines, overexpression of *TOB1-AS1* promoted in vitro cell invasion (**Fig. 4f**). In addition, three weeks after tail vein injection, PANC-1 cells with *TOB1-AS1* overexpression caused higher metastatic burden in immunodeficient mice than the control cells (**Fig. 4g**). Our results showed that the lncRNA *TOB1-AS1* is oncogenic and has pro-metastatic function in pancreatic cancer, and HYENA is able to detect novel proto-oncogenes activated by distal enhancers.

Discussion

Here, we report a computational algorithm HYENA to detect candidate oncogenes activated by distal enhancers via somatic SVs. These SV breakpoints fell in the non-coding regions of the genome and caused shuffling of regulatory elements and altered gene expression. The candidate genes we detected were not limited to protein-coding genes but also included non-coding genes. Our in vitro and in vivo experiments showed that a lncRNA identified by HYENA, *TOB1-AS1*, was a potent oncogene in pancreatic cancers.

HYENA detects candidate genes based on patient cohorts rather than individual samples. Genes need to be recurrently rearranged in the cohort to be detectable. Therefore, sample size is a major limiting factor. Of the eight ground truth cases, HYENA only detected five (**Fig. 2a**). The undetected ones were likely due to small sample size. HYENA aims to identify oncogenes recurrently activated by somatic SVs since these events are under positive selection. However, genes detected in individual tumors by tools such as cis-X and NeoLoopFinder may not be oncogenes. To identify candidate oncogenes among them, recurrence would still be required.

The candidate genes identified by HYENA have statistically significant associations between nearby somatic SVs and elevated expression. However, the relationship may not be causal. It is possible that the presence of SVs and gene expression are unrelated, but both are associated with another factor. We modeled other factors to the best of our ability including gene dosage, tumor purity, patient gender, patient age and principal components of gene expression. In addition, it is also possible that the high gene expression caused somatic SVs. Open chromatin and unwounded double helix regions during transcription are prone to double-strand DNA breaks which may produce somatic SVs. Therefore, it is possible that some of the candidate genes are not oncogenes. Functional studies are required to determine the disease relevance of the candidate genes.

Note that the predicted 3D genome organization is not cell-type-specific. Akita was trained on five high quality Hi-C and Micro-C datasets (HFF, H1hESC, GM12878, IMR90 and HCT116) [37] and predicts limited cell-type-specific differences. Therefore, the predicted TADs reflect conserved 3D genome structure in the five cell types (foreskin fibroblast, embryonic stem cell, B-lymphocyte, lung fibroblast and colon cancer). There were minor differences between HFF and H1hESC (**Supplementary Fig. S2**) in genome organization. For example, the left boundary of the TAD at the *UQCRFS1* locus was different between HFF and H1hESC (**Supplementary Fig. S2a**). Nonetheless, the translocation between chromosomes 17 and 19 removed the left boundary and the right side of the *UQCRFS1* TAD merged with the *TOB1-AS1* TAD (**Fig. 4d**). Therefore, the cell-type difference likely does not have major impact on our results.

Methods

Datasets

This study used data generated by the Pan-Cancer Analysis of Whole Genomes (PCAWG). We limited our study to a total of 1,146 tumor samples for which both whole-genome sequencing (WGS) and RNAseq data were available. The data set was composed of cancers from 25 tumor types including 23 bladder urothelial cancers (BLCA), 88 breast cancers (BRCA), 20 cervical squamous cell carcinomas (CESC), 68 chronic lymphocytic leukemias (CLLE), 51 colorectal cancers (COAD/READ), 20 glioblastoma multiforme (GBM), 42 head and neck squamous cell carcinomas (HNSC), 43 chromophobe renal cell carcinomas (KICH), 37 renal clear cell carcinomas from United States (KIRC), 31 renal papillary cell carcinomas (KIRP), 18 low-grade gliomas (LGG), 51 liver cancers from United States (LIHC), 67 liver cancers from Japan (LIRI), 37 lung adenocarcinomas (LUAD), 47 lung squamous cell carcinomas (LUSC), 95 malignant lymphomas (MALY), 80 ovarian cancers (OV), 74 pancreatic cancers (PACA), 19 prostate adenocarcinomas (PRAD), 49 renal clear cell carcinomas from European Union/France (RECA), 34 sarcomas (SARC), 34 skin cutaneous melanomas (SKCM), 29 stomach adenocarcinomas (STAD), 47 thyroid cancers (THCA), and 42 uterine corpus endometrial carcinomas (UCEC). More detailed information on the sample distribution and annotation can be found in **Supplementary Table S1**.

WGS and RNA-seq data analysis of tumor and normal samples were performed by the PCAWG consortium as previously described [20]. Somatic and germline SNVs, somatic CNVs, SVs and tumor purity were detected by multiple algorithms and consensus calls were made. Genome coordinates were based on hg19 reference genome and GENCODE v19 was used for gene annotation. Gene expression was quantified by HT-Seq (version 0.6.1p1) as fragments per kilobase of million mapped (FPKM). Clinical data such as donor age and sex were downloaded from the PCAWG data portal (https://dcc.icgc.org/pcawg).

Significant eQTL-gene pairs (v8) were downloaded from the Genotype-Tissue Expression (GTEx) data portal (https://gtexportal.org/home/datasets). Only those eQTLs that had a hg19 liftover variant ID were included in the analysis and hg38 variants with no corresponding hg19 annotation were discarded.

HYENA algorithm

First, small tandem duplications (<10kb) were discarded since they are unlikely to produce new promoter-enhancer interactions. The remaining SVs were mapped to the flanking regions (500kb upstream and downstream of transcription start sites [TSSs]) of annotated genes. SVs that fall entirely within a gene body were also discarded. The SV status of each gene was defined by the presence or absence of SV breakpoints within the gene or its flanking regions for each tumor. The binary variable SV status was used in the normal-score regression model below. Only genes with at least 5% samples carrying SVs were tested. For each gene, samples with that gene highly amplified (>10 copies) were removed from the regression model.

Gene expression normal scores

Gene expression quantifications (fragments per kilobase per million [FPKM]) were quantile
normalized (FPKM-QN) using *quantile.normalize()* function from the *preprocessCore* R
package to enhance cross-sample comparison. To break the ties for genes with identical FPKM-
QN values in multiple samples (especially those caused by FPKM of zero) during ranking, very
small Gaussian noises were added to all the FPKM-QN values in all samples by
add.Gaussian.noise(mat, mean = 0.000000001, stddev = 0.000000001, symm = F) from the
RMThreshold R package. Since the mean and standard deviation of the noises added were small,
the rankings of the non-identical values were not affected. For each gene, samples were ranked
based on their noised-added expression values, the ranks were mapped to a standard normal
distribution and the corresponding z scores were gene expression normal scores. Normal score
conversion forced the expression data into a Gaussian distribution, allowing for parametric
comparisons between samples.

Normal-score regression

Generalized linear model was used to test associations between gene expression normal scores
and SV status and control for confounding variables such as gene copy number, tumor sample
purity, donor age and sex. To capture unobserved variations in gene expression, the first n
principal components (PCs) of the expression data were also included in the regression model,
where n was determined as 10% of sample size of the cohort and up to 20 if the sample size was
more than 200. The regression model was shown as below:

$\text{Expression\_normal\_score} \sim \text{sv\_status} + \text{copy\_number} + \text{purity} + \text{age} + \text{sex} + PC_1 + PC_2 \ldots + PC_n$

For each gene, all PCs were tested for associations with the SV status of that gene and those PCs
that significantly correlate (Mann-Whitney test, $P<0.05$) with SV status were not used in
regression.

Calculating empirical p values and model selection

Gene expression data were permuted 100 times by randomly shuffling expression values within
the cohort. The normal-score regression was performed the same way on observed gene
expression and permuted expression. *P* values for SV status from permuted expression were
pooled as a null distribution. Then the *P* values for SV status from observed expression and the
P-value null distribution were used to calculate empirical *P* values. One-sided *P* values were
used since we were only interested in elevated gene expression. False discovery rates (FDRs)
were calculated using Benjamini-Hochberg procedure. Genes with FDR less than 0.1 were
considered candidate genes.

The above empirical *P* value calculation and candidate gene detection were performed iteratively
with no PC and up to n PCs in the regression model. When different numbers of PCs were
included in the model, the numbers of candidate genes varied. The regression model with the
least number of PCs reaching 80% of maximum number of candidate genes in all regression
models tested was selected as final model to avoid over fitting. For example, the sample size for
PCAWG BRCA was 88, therefore, we tested from 0 to 9 PCs. Among these, the model including

8 PCs gave the highest number (82) of candidate genes. Therefore, the model including 7 PCs

with 68 candidate genes was selected as final model since it had the lowest number of PCs

reaching 80% of 82 candidate genes (**Supplementary Table S6**).

Testing eQTL-SV associations

Known germline eQTLs from the matching tissues were obtained from GTEx (**Supplementary**

Table S7). The associations between germline genotypes of eQTLs and SV status of the 213

candidate genes in PCAWG cohort were tested. Genes with significant correlations (P<0.05)

between their SV status and at least one eQTL were removed. The remaining genes were our

final enhancer-hijacking candidate genes.

Benchmarking

Known enhancer hijacking target genes in PCAWG tumor types were selected to test the

sensitivity of HYENA, CESAM and PANGEA. The genes included *MYC* in malignant

lymphoma, *BCL2* in malignant lymphoma, *CCNE1* in stomach/gastric cancer, *TERT* in

chromophobe renal carcinoma, *IGF2* in colorectal cancer, *IGF2* in stomach/gastric cancer,

IGF2BP3 in thyroid cancer, and *IRS4* in lung squamous cell carcinoma. The same SVs, CNVs,

and SNVs were used as input for all three algorithms. For CESAM and PANGEA, upper-

quantile normalized fragments per kilobase per million (FPKM-UQ) were normalized by tumor

purity and gene copy number, and then used as gene expression inputs. CESAM was run using

default parameters, and FDR of 0.1 was used to select significant genes. PANGEA requires

predicted enhancer-promoter (EP) interactions based on ChIP-seq and RNA-seq data. The EP

interactions were downloaded from EnhancerAtlas 2.0 (http://www.enhanceratlas.org/)

(**Supplementary Table S8**). EP interactions from multiple cell lines of the same type were

merged. PANGEA was run with default parameters as well and significant genes were provided

by PANGEA (multiple testing adjusted P value <0.05). To test false positives for HYENA,

CESAM and PANGEA, 20 random gene expression datasets for malignant lymphoma and breast

cancer were generated by randomly shuffling sample IDs in gene expression data. HYENA,

CESAM and PANGEA were run with random expression in the same way as above.

Predicting 3D genome organization

A 1 Mb sequence was extracted from the reference genome centered at each somatic SV

breakpoint and was used as input for Akita [37] to predict the 3D genome organization. Two 500

kb sequences were merged according to the SV orientation to construct the sequence of

rearranged genome. Akita was used to predict the genome organization for the rearranged

sequence. High-resolution Micro-C data obtained from human H1-ESCs and HFF cells [43] were

used to facilitate TAD annotation together with predicted genome organization. H3K27Ac and

CTCF ChIP-seq data from PANC-1 cell line were downloaded from ENCODE data portal

(https://www.encodeproject.org/).

Cell lines

HEK293T, PANC-1, and PATU-8988T cells were obtained from Dr. Alexander Muir
(University of Chicago). All cell lines were cultured at 37°C/5% CO_2. HEK293T cells and
PANC-1 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) (Gibco,
21041025) containing 10% fetal bovine serum (FBS) (Gibco, A4766) as per ATCC (American
Type Culture Collection, USA) instructions (https://www.atcc.org/products/crl-3216,
https://www.atcc.org/products/crl-1469). PATU-8988T cells were cultured with DMEM
containing 5% FBS, 5% horse serum (Gibco, 26050088), and 2 mM L-glutamine as
recommended by DSMZ (Deutsche Sammlung von Mikroorganismen und Zellkulturen,
Germany) (https://www.dsmz.de/collection/catalogue/details/culture/ACC-162). All cell lines
have been regularly monitored and tested negative for mycoplasma using mycoplasma detection
kit (Lonza, LT07-218).

TOB1-AS1 and luciferase overexpression

A 1,351 bp *TOB1-AS1* cDNA (ENST00000416263.3) was synthesized by GenScript (New
Jersey, USA) and subcloned into the lentiviral pCDH-CMV-MCS-EF1-Puro plasmid (SBI,
CD510B-1). The cDNA sequence in the plasmid was verified by Sanger sequencing at
University of Chicago Medicine Comprehensive Cancer Center core facility. The *TOB1-AS1*
overexpression plasmid was amplified by transforming Stellar™ Competent Cells (Takara,
636763) with the plasmid as per instructions and isolated by QIAGEN HiSpeed Plasmid Midi
Kit (QIAGEN, 12643). LucOS-Blast vector was obtained from Dr. Yuxuan Phoenix Miao
(University of Chicago), cloned, and amplified as described above.

HEK293T cells were plated in T-25 flasks and grown to 75% confluence prior to transfection.
For each T-25 flask, 240µl Opti-MEM (Gibco, 31985070), 1.6µg pCMV-VSV-G, 2.56µg
pMDLg/pRRE, 2.56µg pRSV-Rev, 3.4µg *TOB1-AS1* overexpression vector and 22.8µl TransIT-
LT1 Transfection Reagent (Mirus, MIR 2306) were mixed and incubated at room temperature
for 30 minutes, then added to the plated HEK293T cells with fresh medium. The luciferase
vector was packaged into lentivirus with the same method. Upon 48 hours of incubation,
lentiviral supernatant was collected, filtered through 0.45-µmpolyvinylidene difluoride filter
(Millipore), and mixed with 8µg/ml polybrene. PANC-1 or PATU-8988T cells at 60%
confluence were transduced with the lentiviral supernatant for 48 hours followed by three rounds
of antibiotic selection with 4µg/ml puromycin for *TOB1-AS1* overexpression and 10µg/ml
blasticidin for the luciferase expression. *TOB1-AS1* expression was validated by quantitative
reverse transcription polymerase chain reaction (qRT-PCR), and luciferase expression was
validated by in vitro bioluminescence imaging in black wall 96-well plates (Corning, 3603). D-
luciferin potassium salt (Goldbio, LUCK-100) solution with 0, 1.25, 2.5, 5 and 10µl 15mg/ml
was added into the wells in a serial dilution way, and imaging was obtained after 5 minutes.

435 Finally, *TOB1-AS1* overexpression or empty pCDH transduced cell lines with luciferase co-
436 expression were built for both PATU-8988T and PANC-1 cells.

RNA isolation and qRT-PCR

Cells were plated in 6-well plates and allowed to reach 80% confluence prior to RNA extraction.
After cells being lysed in 300µl/well TRYzol^{TM} (Invitrogen, 15596026), RNA samples were
prepared following the manual of Direct-zol RNA Miniprep kit (RPI, ZR2052). Reverse
transcription was performed using Applied Biosystems High-Capacity cDNA Reverse
Transcription Kit (43-688-14) following manufacturer's instructions. Quantitative PCR (qPCR)
was conducted on StepOnePlus Real-Time PCR System (Applied Biosystems, 4376600), using
PowerUp SYBR Green Master Mix (A25742) following the manufacturer's instructions with a
primer concentration of 300nM in 10µl reaction systems. Primers were ordered from Integrated
DNA Technologies. Primer sequences used in this study are as follows:
TOB1 forward: 5' -GGCACTGGTATCCTG AAA AGCC- 3'
TOB1 reverse: 5' – GTGGCAGATTGCCACGAACATC- 3'
TOB1-AS1 forward: 5' -GGAGTGGTCAGGTGACTGATT- 3'
TOB1-AS1 reverse: 5' -ATTCCACTCCTGTTTGCAACT- 3'
GAPDH forward: 5' – ACCACAGTCCATGCCATCAC- 3'
GAPDH reverse: 5' -TCCACCACCCTGTTGCTGTA- 3'
Relative expression levels for *TOB1-AS1* and *TOB1* were calculated by the $2^{\wedge}(-\Delta\Delta C_T)$ method
based on *GAPDH* expression.
456

Transwell assay for cell invasion in vitro

Transparent PET membrane culture inserts of 24-well plate (Falcon, 353097) were coated with
Cultrex Reduced Growth Factor Basement Membrane Extract (BME) (R&D Systems, 3533-010-
02) at 50µg per membrane (200µl of 0.25mg/ml BME stock per membrane) at 37°C for an hour.
A total of 100,000 PANC-1 cells/well and 50,000 PATU-8988T cells/well were resuspended in
serum-free, phenol-red free DMEM media and seeded into the coated inserts. Phenol red free
DMEM of 500µl (Gibco, A1443001) with 10% FBS was added to the bottom of the wells and
the cells were allowed to invade for 16 hours. Additional wells with 500µl serum free, phenol
red free DMEM media without FBS in the bottom chamber were seeded with the same number
of cells as indicated above as negative control. At the end of the assay, the membranes were
stained with 500µl 4µg/ml Calcein AM (CaAM) (Corning, 354216) for an hour at 37°C. The
cells that failed to invade were removed from the top chamber with a cotton swab and all inserts
were transferred into 1x Cell Dissociation Solution (Bio-Techne, 3455-05-03) and shaken at
150rpm for an hour at 37°C. Finally, CaAM signal from the invaded cells was measured by a
plate reader (Perkin Elmer Victor X3) at 465/535nm.

Tumor metastasis in vivo

All animal experiments for this study were approved by the University of Chicago Institutional Animal Care and Use Committee (IACUC) prior to execution. Male NSG mice were ordered from the Jackson Laboratory (strain#005557). Mice were injected intravenously through the tail vein with luciferase-expressing cells at 250,000 cells/mouse for PATU-8988T and at 400,000 cells/mouse for PANC-1 cells. Metastatic burden was measured weekly for 4 weeks via bioluminescence imaging using Xenogen IVIS 200 Imaging System (PerkinElmer) at the University of Chicago Integrated Small Animal Imaging Research Resource (iSAIRR) Facility. Each mouse was weighed and injected intra-peritoneally with D-luciferin solution at a concentration of 150μg/g of body weight, 14 minutes prior to image scanning ventral side up.

Code availability

The HYENA package is available at https://github.com/yanglab-computationalgenomics/HYENA.

Acknowledgements

We thank the Center for Research Informatics at the University of Chicago for providing the computing infrastructure, Matthew Stephens for helpful suggestions, Marsha Rosner for assistance in lentiviral experiments, and Ani Solanki for assistance in animal experiments. The work was supported by the Goldblatt Endowment (A.Y.), the National Institutes of Health grant K22CA193848 (L.Y.), R01CA269977 (L.Y.) and University of Chicago and UChicago Comprehensive Cancer Center (L.Y.).

495 **Disclosure**

496 The authors have no competing interests to declare.

 Figure Legends

Figure 1. Outline of enhancer hijacking and HYENA algorithm. a, Mechanisms of gene
activation by SVs. SVs can activate genes by recruiting distal active enhancers (top panel), by
removing TAD boundaries and forming de novo enhancer-promoter interactions (middle panel),
or by removing repressive elements that block gene expression (bottom panel). **b**, HYENA
workflow. Green and purple boxes denote input and output files, respectively. Orange boxes
denote intermediate steps. Numbers in parentheses represent default values of HYENA.

Figure 2. Benchmarking HYENA. a, Comparison of HYENA, CESAM, and PANGEA in
detecting oncogenes known to be activated by enhancer hijacking in six tumor types from the
PCAWG cohort. **b**, UPSET plot demonstrating candidate genes identified and shared among the
three tools in five tumor types of PCAWG. Individual dots denote genes detected by one tool,
and dots connected by lines denote genes detected by multiple tools. **c**, Number of genes
detected by HYENA in two PCAWG tumor types using observed gene expression and
randomized expression. Genes detected in random expression datasets are false positives.

Figure 3. Enhancer hijacking candidate genes in PCAWG. a, Candidate genes detected by
HYENA in individual tumor types of PCAWG. Genes that are plotted twice (e.g. *TERT*) are
detected in more than one cancer types. Genes labelled as red are known enhancer hijacking
targets. **b**, Diverse types of candidate genes identified by HYENA in PCAWG. Numbers in
parentheses denote sample size in the corresponding tumor types.

Figure 4. *TOB1-AS1* activated by various types of SVs in pancreatic cancer. a, Normalized
expression of the *TOB1-AS1* in samples with (n=7) and without nearby SVs (n=67) in pancreatic
cancers. Boxplot shows median values (thick black lines), upper and lower quartiles (boxes),
1.5× interquartile range (whiskers) and outliers (black dots). **b**, Circos plot summarizing
intrachromosomal SVs (blue, n=5) and translocations (red, n=3) near *TOB1-AS1*. **c**, Diagrams
depicting putative enhancer hijacking mechanisms that activate *TOB1-AS1* in one tumor with a
17:19 translocation (left panel) and another tumor with a large deletion (right panel). **d**, Predicted
3D chromatin interaction maps of the *TOB1-AS1* (left panel), *UQCRFS1* (middle panel), and the
translocated region in tumor 9ebac79d-8b38-4469-837e-b834725fe6d5 (right panel). The
downstream fragment of the chromosome 19 SV breakpoint was flipped in orientation and linked
to chromosome 17. H3K27Ac and CTCF ChIP-seq data of PANC-1 cell line are shown at the
bottom. **e**, Predicted 3D chromatin interaction maps of *TOB1-AS1* (left panel) and *KCNJ2*
(middle panel) loci without deletion as well as the region after deletion in tumor 748d3ff3-8699-
4519-8e0f-26b6a0581bff (right panel). **f**, *TOB1-AS1* overexpression in PATU-8988T (4
biological replicates) and PANC-1 (3 biological replicates) promoted in vitro cell invasion using
Transwell assay. Each biological replicate was an independent experiment with 7 technical
replicates per experimental group. The average fold change of cell invasion was calculated after
the background invasion measured in the absence of any chemotactic agent being subtracted
from each technical replicate. *P* values were calculated by two-sided student t test. **g**, *TOB1-AS1*
overexpression in PANC-1 cells promoted in vivo tumor metastasis. Images of radiance in
immunodeficient mice are shown on the left while the quantifications of radiance are shown on
the right. Eight mice were used in both overexpression group and empty vector control. The
images were analyzed by setting the regions of interest (ROIs) to mouse torsos and measuring

the average radiance level (in p/sec/cm^2/sr). Two-side student t test was used. *, **, ***
represent $P<.05, <.01$, and $<.001$, respectively. Error bars in **f** and **g** indicate standard error of the
mean.
543

References

1. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of
chromatin interactions. *Nature* **485**, 376–380 (2012).

2. Krijger, P. H. L. & De Laat, W. Regulation of disease-associated gene expression in the
3D genome. *Nat. Rev. Mol. Cell Biol.* **17**, 771–782 (2016).

3. Symmons, O. *et al.* Functional and topological characteristics of mammalian regulatory
domains. *Genome Res.* **24**, 390–400 (2014).

4. Lupiáñez, D. G. *et al.* Disruptions of topological chromatin domains cause pathogenic
rewiring of gene-enhancer interactions. *Cell* **161**, 1012–1025 (2015).

5. Weischenfeldt, J., Symmons, O., Spitz, F. & Korbel, J. O. Phenotypic impact of genomic
structural variation: insights from and for human disease. *Nat. Rev. Genet.* **14**, 125–138
(2013).

6. Davis, C. F. *et al.* The somatic genomic landscape of chromophobe renal cell carcinoma.
Cancer Cell **26**, 319–330 (2014).

7. Bakhshi, A. *et al.* Cloning the chromosomal breakpoint of t(14;18) human lymphomas:
clustering around Jh on chromosome 14 and near a transcriptional unit on 18. *Cell* **41**,
899–906 (1985).

8. Gostissa, M. *et al.* Long-range oncogenic activation of Igh-c-myc translocations by the
Igh 3′ regulatory region. *Nature* **462**, 803–807 (2009).

9. Hnisz, D. *et al.* Activation of proto-oncogenes by disruption of chromosome
neighborhoods. *Science* **351**, 1454–1458 (2016).

10. Gröschel, S. *et al.* A single oncogenic enhancer rearrangement causes concomitant EVI1
and GATA2 deregulation in leukemia. *Cell* **157**, 369–381 (2014).

11. Northcott, P. A. *et al.* Enhancer hijacking activates GFI1 family oncogenes in
medulloblastoma. *Nature* **511**, 428–434 (2014).

12. Weischenfeldt, J. *et al.* Pan-cancer analysis of somatic copy-number alterations implicates
IRS4 and IGF2 in enhancer hijacking. *Nat. Genet.* **49**, 65–74 (2016).

13. Northcott, P. A. *et al.* The whole-genome landscape of medulloblastoma subtypes. *Nature*
547, 311–317 (2017).

14. He, B. *et al.* Diverse noncoding mutations contribute to deregulation of cis-regulatory
landscape in pediatric cancers. *Sci. Adv.* **6**, eaba3064 (2020).

15. Kopp, F. & Mendell, J. T. Functional Classification and Experimental Dissection of Long
Noncoding RNAs. *Cell* **172**, 393–407 (2018).

16. Lin, C.-P. & He, L. Noncoding RNAs in Cancer Development. *Annu. Rev. Cancer Biol.* **1**,
163–184 (2017).

17. Liu, S. J., Dang, H. X., Lim, D. A., Feng, F. Y. & Maher, C. A. Long noncoding RNAs in
cancer metastasis. *Nat. Rev. Cancer 2021 217* **21**, 446–460 (2021).

18. Liu, Y. *et al.* Discovery of regulatory noncoding variants in individual cancer genomes by
using cis-X. *Nat. Genet.* **52**, 811–818 (2020).

19. Wang, X. *et al.* Genome-wide detection of enhancer-hijacking events from chromatin
interaction data in rearranged genomes. *Nat. Methods* **18**, 661–668 (2021).

20. Campbell, P. J. *et al.* Pan-cancer analysis of whole genomes. *Nature* **578**, 82–93 (2020).

21. Beroukhim, R. *et al.* The landscape of somatic copy-number alteration across human
cancers. *Nature* **463**, 899–905 (2010).

22. Uhlen, M. *et al.* A pathology atlas of the human cancer transcriptome. *Science* **357**,
eaan2507 (2017).

23. Zhang, X. *et al.* Identification of focally amplified lineage-specific super-enhancers in
human epithelial cancers. *Nat. Genet.* **48**, 176–182 (2015).

24. Takeda, D. Y. *et al.* A Somatically Acquired Enhancer of the Androgen Receptor Is a
Noncoding Driver in Advanced Prostate Cancer. *Cell* **174**, 422-432.e13 (2018).

25. Turner, K. M. *et al.* Extrachromosomal oncogene amplification drives tumour evolution
and genetic heterogeneity. *Nature* **543**, 122–125 (2017).

26. Wu, S. *et al.* Circular ecDNA promotes accessible chromatin and high oncogene
expression. *Nature* **575**, 699–703 (2019).

27. Morton, A. R. *et al.* Functional Enhancers Shape Extrachromosomal Oncogene
Amplifications. *Cell* **179**, 1330-1341.e13 (2019).

28. Li, Y., Ge, X., Peng, F., Li, W. & Li, J. J. Exaggerated false positives by popular
differential expression methods when analyzing human population samples. *Genome Biol.*
23, 79 (2022).

29. Ooi, W. F. *et al.* Integrated paired-end enhancer profiling and whole-genome sequencing
reveals recurrent CCNE1 and IGF2 enhancer hijacking in primary gastric
adenocarcinoma. *Gut* **69**, 1039–1052 (2020).

30. Yun, J. W. *et al.* Dysregulation of cancer genes by recurrent intergenic fusions. *Genome*
Biol. **21**, 166 (2020).

31. Richter, J. *et al.* Recurrent mutation of the ID3 gene in Burkitt lymphoma identified by
integrated genome, exome and transcriptome sequencing. *Nat. Genet.* **44**, 1316–1320
(2012).

32. Nik-Zainal, S. *et al.* Landscape of somatic mutations in 560 breast cancer whole-genome
sequences. *Nature* **534**, 47–54 (2016).

33. Li, Y. *et al.* Patterns of somatic structural variation in human cancer genomes. *Nature* **578**,
112–121 (2020).

34. Panebianco, F. *et al.* THADA fusion is a mechanism of IGF2BP3 activation and IGF1R
signaling in thyroid cancer. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 2307–2312 (2017).

35. Zapatka, M. *et al.* The landscape of viral associations in human cancers. *Nat. Genet. 2020*

523 **52**, 320–330 (2020).

36.   Neuveut, C., Wei, Y. & Buendia, M. A. Mechanisms of HBV-related
620   hepatocarcinogenesis. *J. Hepatol.* **52**, 594–604 (2010).

621   37.   Fudenberg, G., Kelley, D. R. & Pollard, K. S. Predicting 3D genome folding from DNA
622   sequence with Akita. *Nat. Methods* **17**, 1111–1117 (2020).

623   38.   Rao, S. S. P. *et al.* A 3D Map of the Human Genome at Kilobase Resolution Reveals
624   Principles of Chromatin Looping. *Cell* **159**, 1665–1680 (2014).

625   39.   Fudenberg, G. *et al.* Formation of Chromosomal Domains by Loop Extrusion. *Cell Rep.*
626   **15**, 2038–2049 (2016).

627   40.   Franke, M. *et al.* Formation of new chromatin domains determines pathogenicity of
628   genomic duplications. *Nature* **538**, 265–269 (2016).

629   41.   Melo, U. S. *et al.* Complete lung agenesis caused by complex genomic rearrangements
630   with neo-TAD formation at the SHH locus. *Hum. Genet.* **140**, 1459–1469 (2021).

631   42.   de Bruijn, S. E. *et al.* Structural Variants Create New Topological-Associated Domains
632   and Ectopic Retinal Enhancer-Gene Contact in Dominant Retinitis Pigmentosa. *Am. J.*
633   *Hum. Genet.* **107**, 802–814 (2020).

634   43.   Krietenstein, N. *et al.* Ultrastructural Details of Mammalian Chromosome Architecture.
635   *Mol. Cell* **78**, 554-565.e7 (2020).

636   44.   Yao, J. *et al.* Long noncoding RNA TOB1-AS1, an epigenetically silenced gene,
637   functioned as a novel tumor suppressor by sponging miR-27b in cervical cancer. *Am. J.*
638   *Cancer Res.* **8**, 1483 (2018).

639   45.   Shangguan, W. *et al.* TOB1-AS1 suppresses non-small cell lung cancer cell migration and
640   invasion through a ceRNA network. *Exp. Ther. Med.* **18**, (2019).

a *TOB1-AS1* FDR=0.07

b Intra-chr SVs / Inter-chr SVs

c 9ebac79d-8b38-4469-837e-b834725fe6d5 (translocation)

748d3ff3-8699-4519-8e0f-26b6a0581bff (deletion)

enhancers ● ● ● — X SV breakpoints — unexpressed genes — expressed genes — Deleted region (19.3 Mb)

d Wildtype *TOB1-AS1* locus — Wildtype *UQCRFS1* locus — 9ebac79d-8b38-4469-837e-b834725fe6d5 (translocation)

X SV breakpoints
➤ SV partners
➤ SV partners
■ Interacting loci

log(predicted/expected)

TADs

XYLT2 ACSF2 EPN3 ABCC3 SPAG9 NME1-NME2
RSAD1 CACNA1G LUC7L3 *TOB1-AS1*

AC005307.5 UQCRFS1 VSTM2B-DT

UQCRFS1 *TOB1-AS1* SPAG9 NME1-NME2

H3K27Ac
CTCF

chr17:48,363,541-49,281,045   chr19:28,958,642-29,876,146   chr19:29,417,394-29,876,146 chr17:48,822,293-49,281,045

e Wildtype *TOB1-AS1* locus — Wildtype *KCNJ2* locus — 748d3ff3-8699-4519-8e0f-26b6a0581bff (deletion)

TADs

EPN3 ABCC3 SPAG9 MBTD1
CACNA1G LUC7L3 *TOB1-AS1* NME1-NME2 UTP18

KCNJ16 KCNJ2

EPN3 ABCC3 *TOB1-AS1*
CACNA1G LUC7L3

H3K27Ac
CTCF

chr17:48,570,928-49,488,432   chr17:67,852,922-68,770,426   chr17:48,570,928-49,029,680 chr17:68,311,674-68,770,426

f PANC-1 / PATU-8988T

Cell invasion fold change — Over expression / Empty vector

*** **

g Over-expression / Vector

Radiance (x10⁵ p/sec/cm²/sr)

PANC-1 metastasis

Average radiance in torso (p/sec/cm²/sr) — Over expression / Empty vector

*