1 **Dopamine and norepinephrine differentially mediate the exploration-exploitation tradeoff**

2 Cathy S. Chen[1], Dana Mueller[1], Evan Knep[1], R. Becket Ebitz[2], Nicola M. Grissom[+1]

3 [1] Department of Psychology, University of Minnesota, Minneapolis MN 55455
4 [2] Department of Neurosciences, Université de Montréal, Quebec, Canada
5
6
+to whom correspondence should be addressed:
Nicola Grissom
Department of Psychology
University of Minnesota
75 East River Rd
Minneapolis, MN 55455
ngrissom@umn.edu
14

Abstract

The catecholamines dopamine (DA) and norepinephrine (NE) have been repeatedly implicated in

neuropsychiatric vulnerability, in part via their roles in mediating the decision making processes.

Although the two neuromodulators share a synthesis pathway and are co-activated under states

of arousal, they engage in distinct circuits and roles in modulating neural activity across the

brain. However, in the computational neuroscience literature, they have been assigned similar

roles in modulating the latent cognitive processes of decision making, in particular the

exploration-exploitation tradeoff. Revealing how each neuromodulator contributes to this

explore-exploit process will be important in guiding mechanistic hypotheses emerging from

computational psychiatric approaches. To understand the differences and overlaps of the roles of

these two catecholamine systems in regulating exploration and exploitation, a direct comparison

using the same dynamic decision making task is needed. Here, we ran mice in a restless two-

armed bandit task, which encourages both exploration and exploitation. We systemically

administered a nonselective DA receptor antagonist (flupenthixol), a nonselective DA receptor

agonist (apomorphine), a NE beta-receptor antagonist (propranolol), and a NE beta-receptor

agonist (isoproterenol), and examined changes in exploration within subjects across sessions. We

found a bidirectional modulatory effect of dopamine receptor activity on the level of exploration.

Increasing dopamine activity decreased exploration and decreasing dopamine activity increased

exploration. Beta-noradrenergic receptor activity also modulated exploration, but the modulatory

effect was mediated by sex. Reinforcement learning model parameters suggested that dopamine

modulation affected exploration via decision noise and norepinephrine modulation affected

exploration via outcome sensitivity. Together, these findings suggested that the mechanisms that

govern the transition between exploration and exploitation are sensitive to changes in both

catecholamine functions and revealed differential roles for NE and DA in mediating exploration.

Introduction

Dysfunctions in cognitive processes, particularly within the domain of executive function, are

implicated in numerous neuropsychiatric disorders [1–3]. One essential aspect of executive

function is value-based decision making. Decision making involving exploring and learning in a

dynamic environment has been a focus in computational neuroscience, which has developed

approaches for discovering latent structures in decision making strategies [4–7]. Many

computational models for value-based decision making distinguish two essential latent

processes: exploration and exploitation [8–12]. Exploration is a ubiquitous learning process

across species that allows the discovery of the best action to take in an uncertain environment

[6,9,13]. Once a favorable action or option is discovered, exploitation, i.e. repeating a rewarding

action, is necessary to obtain the best rates of reward. However, exploitation must be balanced

with continued exploration as environments and their reward probabilities change. Dysregulation

in this exploration-exploitation tradeoff can be observed in the phenotypes of numerous

neuropsychiatric disorders and challenges, including schizophrenia, autism spectrum disorders,

addictions, and chronic stress [14–18]. Understanding the fundamental mechanisms impacting

the computations in our brain that underlie the balance between exploration and exploitation

could help identify critical circuits that are associated with differential risk factors of

neuropsychiatric disorders and open avenues for novel interventions for executive function

challenges.

Many neuropsychiatric challenges are associated with dysfunction in the catecholamines

dopamine and norepinephrine [16,19–24]. These neuromodulators are well-positioned to carry

information about the state of the environment and influence behavior outputs via downstream

control of action selection [9,25]. Indeed, dopamine and norepinephrine have each separately

been implicated in mediating the exploration-exploitation tradeoff by promoting exploration

[9,23–28]. While dopamine and norepinephrine share biosynthetic pathways and are co-activated

under states of arousal, they act through partially separable circuits and have distinct

pharmacological profiles [29]. It is notable, therefore, that dopamine and norepinephrine have

largely been ascribed similar or even identical roles in mediating the explore-exploit tradeoff in

the computational neuroscience literature.

We have previously shown that multiple latent cognitive processes, such as the reinforcement

learning model-derived parameters of learning rate and decision noise, can describe differences

in the rates of exploration between groups [4]. Dopamine has often been ascribed a role in

mediating reward prediction errors and value-based learning [10,30–32] while norepinephrine in

modulating arousal, attention, and value assessment [23,33,34], but more recently both

neuromodulators have been implicated in the process of action selection [35,36]. Therefore, it is

unknown whether dopamine and norepinephrine govern this exploration-exploitation tradeoff via

distinct or similar latent cognitive processes because they have been largely studied in this

context in isolation [26].

To uncover the distinct roles of dopamine and norepinephrine in the exploration-exploitation

tradeoff, we conducted within-subjects pharmacological manipulations for these two

neuromodulators on exploration, and the latent cognitive parameters that influence exploration,

in the same restless bandit task. We pharmacologically up- and down-regulated dopaminergic

and noradrenergic receptor activity and compared the modulatory effects on exploration in the

same dynamic decision making task, a spatial restless two-armed bandit [4,6]. We found a

bidirectional modulatory effect of dopamine receptor activity on the level of exploration.

Increasing dopamine activity decreased exploration and decreasing dopamine activity increased

exploration. Beta-noradrenergic receptor activity also modulated exploration but the modulatory

effect was mediated by sex. Contrary to dopamine modulating choice solely by changing value-

based learning for outcomes, via reinforcement learning modeling we find that dopamine

mediates exploration by changing decision noise, i.e. the precision of value-based choice

selection. In contrast, noradrenergic activity surprisingly influenced exploration by changing

value-based learning for outcomes in a sex dependent manner. These data suggest differential

roles of dopamine and norepinephrine in mediating exploration, and complex roles for both

catecholamines in signaling reward.

Results

Aged-matched adult male and female wildtype mice (n = 32, 16 males and 16 females, strain B6129SF1/J) were trained to perform a restless two-armed spatial bandit task in touch-screen operant chambers (**Figure 1A**). In this task, animals were presented with two identical visual targets (squares) on the left and right side of the screen during each trial. They indicated their choices by nose poking at one of the two target locations, each of which provided some probability of reward that changed independently and randomly over time (**Figure 1B)**. Animals had to infer which location offered better payoff by constantly sampling the two locations and learning from the reward outcomes. The dynamic reward contingency of this task naturally encourages the animal to balance between exploiting a favorable option when it is found and exploring to gain information about potential better alternatives. This task has been employed in rodents and primates as well as human subjects to understand learning and exploration and have successfully revealed divergent exploration strategies [4,6,37]. In this study, we adopted this task to understand the modulatory effect of catecholamine receptor activity on the transition between exploration and exploitation.

To examine the distinct and/or overlapping roles of dopamine and norepinephrine in modulating exploration, we tested the systemic effect of a non-selective dopamine receptor agonist apomorphine (0.1mg/kg), a non-selective dopamine receptor antagonist flupenthixol (0.03mg/kg), a beta-noradrenergic receptor agonist isoproterenol (0.3mg/kg), and a beta-noradrenergic receptor antagonist propranolol (5mg/kg) on exploration. The dosages of the drugs were based on previous studies on the role of dopamine and norepinephrine in cognitive processes [38–44]. Animals were intraperitoneally (IP) administered either saline (control) or treatment with 5 mg/kg of propranolol, 0.3mg/kg of isoproterenol, or 0.03 mg/kg of flupenthixol prior to the bandit task or 0.1mg/kg of apomorphine 30 mins prior to the bandit task. Saline and drug treatments were administered interleaved by session (**Figure 1C, 2A**). Mice performed 12 sessions under each pharmacological condition (6 control and 6 drug treatment sessions). The experiment used a within-subject design, where animals received each of the four drugs tested, allowing us to directly compare the effect of drugs within individual animals. Each session consisted of 300 trials.

132 To examine whether mice had learned to perform the task and were able to sustain reward, we

133 first calculated the average probability of obtaining reward across sessions in all control and

134 treatment groups. It's worth noting that in this task, there is no asymptotic performance because

135 the animals need to constantly adapt to the changing reward contingency and any one option is

136 not always the best choice to make. At any given time, an animal could decide to repeat a

137 favorable option or to explore the options. The performance is best measured by the amount of

138 reward acquired. Because reward schedules were dynamic and stochastic, sessions could differ

139 slightly in the amount of reward that was available. We therefore compared the probability of

140 actual obtained reward against the probability of reward if chosen randomly (chance). In the

141 dopamine modulation condition, both control groups (saline) and treatment groups (apomorphine

142 and flupenthixol) were able to obtain reward more frequently than chance (**Figure 1D, E**, one-

143 sample t test, vehicle control (apomorphine): $t(31) = 13.96$, $p<0.0001$; apomorphine: $t(31)=$

144 17.24, $p<0.0001$; vehicle control (flupenthixol): $t(31)=12.36$, $p<0.0001$; flupenthixol:

145 $t(31)=16.48$, $p<0.0001$). When modulating beta-noradrenergic receptor activity, both control

146 groups (saline) and treatment groups (isoproterenol and propranolol) were able to obtain reward

147 more frequently than chance (**Figure 2B, D**, one-sample t test, vehicle control (isoproterenol):

148 $t(31)=10.01$, $p<0.0001$; isoproterenol: $t(31)=13.84$, $p<0.0001$; vehicle control (propranolol):

149 $t(31)=10.60$, $p<0.0001$; propranolol: $t(31)=20.08$, $p<0.0001$). These results suggested that

150 animals were not choosing randomly. The animals demonstrated a strong understanding of the

151 task and were able to effectively obtain rewards above chance under baseline (vehicle control)

152 and all drug treatments.

154 Next, we asked whether manipulating dopaminergic receptor activity influenced their reward

155 acquisition performance. We used a generalized linear mixed model (GLMM) to estimate effects

156 and interactions of drug treatments and sex, with session and animal identity as random effects

157 (**Equation 1**, see methods). Increasing dopamine receptor activity with apomorphine

158 administration did not significantly alter their reward acquisition performance (**Figure 1D**,

159 GLMM, main effect of drug, $p= 0.11$, $\beta 1 = -0.017$ **)**. Antagonizing dopamine receptor activity

160 with flupenthixol administration also did not significantly alter reward acquisition performance

161 (**Figure 1E**, GLMM, main effect of drug, $p= 0.28$, $\beta 1 = -0.008$). Increasing beta-noradrenergic

receptor activity with isoproterenol administration did not affect the amount of reward obtained

(**Figure 2B**, GLMM, main effect of drug, p= 0.93, $\beta1$ = -0.0005). However, when decreasing

beta-noradrenergic receptor activity with propranolol administration, animals obtained

significantly more reward (**Figure 2D**, GLMM, main effect of drug, p < 0.0001, $\beta1$ = -0.497).

Even though manipulating dopamine receptor activity and increasing beta-noradrenergic receptor

activity did not affect how much overall obtained reward, we considered the possibility that the

tuning of catecholamine receptor activity affect latent exploration strategies underlying choice

patterns, but not an observable change in the level of accuracy. Our previous study demonstrated

that similar reward acquisition performance can be achieved via divergent explore strategies [4].

Pharmacological manipulations of both dopaminergic and noradrenergic receptor activity have

been shown to have a dose-dependent effect on motor functions [44,45]. In this study, dosage

selection was based on previous literature examining cognitive functions and a low dose was

used to have influence on cognition but avoid motor function impairment [38,41]. One

commonly used behavioral metric to examine motor function in a cognitive task is response

time. Therefore, we calculated the response time, which was the time elapsed between the onset

of choices and nose poke response to make a choice, as recorded by the touchscreen operant

chamber. If the administration of drugs impaired motor function, we would expect to see longer

response time in the drug group compared to vehicle control. Upregulating dopamine receptor

activity with apomorphine did not significantly influence the response time (GLMM, main effect

of drug, p=0.77, $\beta1$ = 0.10). Similarly, downregulating dopamine receptor activity with

flupenthixol did not significantly alter animals' response time (GLMM, main effect of drug,

p=0.96, $\beta1$ = -0.01). Modulating beta-noradrenergic receptor activity resulted in bidirectional

changes in response time. When increasing beta-noradrenergic receptor activity with

isoproterenol, both males and females took significantly longer to respond (**Figure 2C,** GLMM,

main effect of drug, p<0.0001, $\beta1$ = -3.51). Decreasing beta-noradrenergic receptor activity with

propranolol significantly decreased the response time (**Figure 2E,** GLMM, main effect of drug,

p=0.04, $\beta1$ = 0.51), primarily driven by response time reduction in females under propranolol

compared to vehicle controls (GLMM, interaction term, p<0.009, $\beta3$ = -0.57). Faster response

time under propranolol could be associated with more reward obtained - propranolol was the

only drug treatment that altered reward acquisition performance among the four drugs tested.

Previous studies have linked response time with the complexity of the strategy used [4,46,47].

Faster response time under propranolol might also point to the possibility that decreasing beta-

noradrenergic receptor activity brought on changes in strategies that took shorter time to execute.

Isoproterenol was a highly significant predictor for lower response time in both male and female

mice, implicating the potential drug effect on motor behaviors.

Opposing modulatory effect of dopamine and norepinephrine receptor activity on win-stay

lose-shift behaviors

The role of dopamine has been heavily studied and shown to be a key contributor to value-based

decision making and reinforcement learning [27,28,31,48]. Although modulating dopamine

receptor activity did not significantly influence the overall amount of reward obtained or the

response time, it is possible that apomorphine and flupenthixol exerted influence on how animals

adapted their choices to reward outcome. To understand how increasing and decreasing

dopamine activity influenced animals' reward-driven choice behaviors, we examined the

probability of win-stay (repeating a choice when it was rewarded on the previous trial) and the

probability of lose-shift (switching to the other choice when not rewarded). We found that when

increasing dopamine receptor activity with apomorphine administration, both males and females

had increased win-stay behavior (**Figure 1F,** GLMM, main effect of drug, p=0.011, $\beta 1$= -0.02)

and decreased lose-shift behavior (**Figure 1G,** GLMM, main effect of drug, p=0.016, $\beta 1$=

0.034). This result suggested that apomorphine increased the overall stickiness of choice and

more stay behaviors regardless of the outcome.

There was no significant change in the probability of win-stay when the animals were

administered flupenthixol (**Figure 1H,** GLMM, main effect of drug, p=0.27, $\beta 1$= -0.001). It's

worth noting that there was baseline sex difference in the probability of win-stay, with females

having a higher probability of win-stay (GLMM, main effect of sex, apomorphine: p=0.065, $\beta 2$ =

-0.054; flupenthixol: p=0.01, $\beta 2$ =-0.05). The results also revealed an opposing effect from

dopamine agonist that antagonizing dopamine receptor activity with flupenthixol administration

increased the probability of lose-shift (**Figure 1I,** GLMM, main effect of drug, p=0.04, $\beta 1$ = -

0.014).

To control for the total amount of switching, we calculated outcome sensitivity (**Equation 2,** see

Methods) to understand how much of the switching behavior was outcome-sensitive. The result

suggests that there were no significant changes in outcome sensitivity when animals were

administered apomorphine (GLMM, main effect of drug, p=0.22, β1 = -0.08). However,

administration of flupenthixol significantly increased outcome sensitivity across both sexes

(**Figure S1A,** GLMM, main effect of drug, p=0.006, β1= -0.06) and there was more increase in

outcome sensitivity in males (GLMM, interaction term, p=0.027, β3= -0.10).

Modulating beta-noradrenergic receptor activity also influenced win-stay and lose-shift

behaviors. Increasing beta-noradrenergic activity with isoproterenol administration decreased the

probability of win-stay compared to vehicle control (**Figure 2F,** GLMM, main effect of drug,

0.009, β1=0.007). Isoproterenol administration also decreased the probability of lose-shift in

both sexes (**Figure 2G,** GLMM, main effect of drug, p=0.0007, β1=0.083), primarily driven by

significant decrease in lose-shift in females under isoproterenol (**Figure 2G inset,** GLMM,

interaction term, p=0.004, β3=-0.077). Outcome sensitivity analysis revealed a significant

decreased in outcome sensitivity when administer isoproterenol compared to vehicle control

(**Figure S1B,** GLMM, main effect of drug, p=0.021, β1 = 0.134). In contrast, decreasing beta-

noradrenergic activity with propranolol administration increased the probability of win-stay

compared to vehicle control (**Figure 2H,** GLMM, main effect of drug, p<0.0001, β1 = -0.023).

There was also a significant interaction between sex and drug that males showed a greater

increase in win-stay on propranolol compared to females (**Figure 2H inset,** GLMM, interaction

term, 0.02, β3=-0.036). Interestingly, propranolol administration also decreased the probability

of lose-shift but only in males (**Figure 2I,** GLMM, main effect of drug, 0.45, β1 = -0.01,

interaction term, p= 0.002, β3 =0.08). Outcome sensitivity analysis revealed a significant

increase in outcome sensitivity with propranolol administration (**Figure S1C,** GLMM, main

effect of drug, 0.003, β1 = -0.186). Together, these results suggested an opposing role of

dopamine and norepinephrine in modulating reward-driven decision making. Increasing

dopamine or decreasing norepinephrine activity resulted in increased win-stay and decreased

lose-shift but this effect is more sex-different in the noradrenergic manipulations.

A Hidden Markov model (HMM) identifies distinct patterns of exploration and
exploitation associated with modulation of dopamine and norepinephrine activity.
256

The changes in the reward-driven behaviors such as win-stay and lose-shift could be a
manifestation of strategic changes in how animals explored the changing environment. To
understand how modulation of dopamine and norepinephrine affect the balance between
exploration and exploitation, we used a Hidden Markov model (HMM) that modeled exploration
and exploitation as two latent goal states to infer which choices were exploratory or exploitative
(**Figure 3A**). Figure 3A showed an example of HMM labeling of choices of an animal in the
restless bandit task. The shaded area indicates exploratory trials inferred by the HMM. The
animal displayed mixtures of exploratory bouts where choices were distributed across two
choices and exploitative bouts where one choice was repeated. The HMM model has previously
been used in rodents, primates and humans [4,6,49] to infer explore-exploit state from choices
and has shown robust correlation with neural activity as well as other behavioral metrics,
including response time, value function, and reinforcement learning.
269

Essentially, when a HMM model is fitted to animals' choices, it estimates a transition matrix,
which is the mapping between every state at one time point and every state at some future point
of time. The HMM model fitting is the process of estimating how the distribution over states will
evolve over time. The transition matrix estimates the probability of transition between states -
that is the probability of staying in the explore state, the probability of transitioning from explore
state to exploit state, the probability of transitioning from exploit state to explore state, and the
probability of staying in exploit state. Note that exploration is modeled as a uniform distribution
over choices. This is because the uniform distribution over choices is the maximum entropy
distribution of categorical variables. We want to make as few assumptions as we possibly can
about the choices they make during exploration. With the HMM transition matrix optimized
from each animal's choice sequences, we then used the Viterbi algorithm to decode latent states
from choices, allowing us to label each choice as either exploratory or exploitative.
282

To examine how much animals were exploring, we calculated the average number of exploratory
choices inferred from HMM. The results revealed a bidirectional modulatory effect of dopamine

receptor activity on the level of exploration. When animals were administered apomorphine, we

found that animals on average had fewer exploratory trials than when they were on vehicle

control, with 55.5% ± 12.6% STD of trials labeled as exploratory on apomorphine and 72.3% ±

13.6% STD of trials labeled as exploratory on vehicle (**Figure 3B, D,** GLMM, main effect of

drug, p<0.0001, $\beta1 = 0.174$). This is also consistent with our findings in the win-stay lose-shift

analysis that apomorphine increased the stickiness or repetitiveness of choice behaviors. In

contrast, when decreasing dopamine receptor activity with flupenthixol administration, animals

explore more compared to vehicle control, with 58.2% ± 12.0% STD of trials being exploratory

on flupenthixol and 52.1% ± 12.3% STD of trials being exploratory on vehicle control (**Figure**

3B, D, GLMM, main effect of drug, 0.002, $\beta1 = -0.062$). This result suggested that manipulating

dopamine receptor activity bidirectionally affected the level of exploration across both sexes

(**Figure 3C, Figure S2)** - increasing dopamine receptor activity decreased exploratory choices

and decreasing dopamine receptor activity increased exploratory choices.

Modulating beta-noradrenergic receptor activity also influenced the probability of exploration

and the modulatory effect on exploration was associated with sex. Decreasing beta-noradrenergic

receptor activity with propranolol significantly decreased the level of exploration (**Figure 4A, C,**

D) but only in males (**Figure 4F, Figure S2**, GLMM, interaction term, 0.0001, $\beta1 = 0.138$).

Increasing beta-noradrenergic receptor activity with isoproterenol administration significantly

decreased exploration in females (**Figure 3G, 3E, Figure S2**, interaction term, p= 0.002, $\beta3=-$

0.103). Together these results suggested a sex-linked neuromodulatory effect of beta-

noradrenergic receptor on exploration - increasing beta-noradrenergic activity decreased

exploration in males and decreasing beta-noradrenergic activity decreased exploration in

females. One possibility is that this sex-differentiated modulatory effect reflected sex-dependent

ceiling/floor effects of noradrenergic signaling. Another possibility is that there is a potential

inverted U-shape relationship between beta-noradrenergic receptor activity level and exploration

(**Figure 4H**). Previous studies have suggested higher tonic norepinephrine (NE) levels in females

due to estrogen-mediated enhanced NE synthesis and reduced NE clearance [50]. The difference

in baseline NE activity across sexes may contribute to the differential modulatory effect on

exploration when increasing or decreasing in noradrenergic receptor activity.

Reinforcement learning (RL) models revealed changes in distinct parameters under
modulation of dopamine and epinephrine
318
The results of the HMM suggested that both dopamine and norepinephrine modulation
influenced the level of exploration. However, it is unclear whether changes in exploration were
due to changes in similar or distinct latent cognitive parameters. Does dopamine and
norepinephrine modulate exploration via distinct or overlapping mechanisms? To address this
question, we fitted a series of reinforcement learning (RL) models to understand the effect of
pharmacological manipulation on the latent cognitive parameters that could influence exploration
and exploitation [10,51–53].
326
To make inferences about changes in the RL model parameters, we first identified the best-fitting
RL model for the animals' behaviors. The majority of RL model parameters can be categorized
in three ways: value-dependent learning terms, value-independent bias terms, and decision
noise/randomness terms [54]. Previous studies demonstrated the effect of various RL parameters
on value-based decision making, including value-dependent terms such as learning rate and
asymmetrical learning rate [4,38,55,56], value-independent terms such as choice bias [4,54,57],
and noise terms such as inverse temperature and lapse rate [38,57,58]. Here, we compared six
reinforcement learning models that incorporated one or more of the above parameters (learning
rate, bias, noise) and had different assumptions about the latent cognitive processes that animals
might employ during exploration (**Figure 5A, Equations 3-7)**. These models included: (1) a
"random" model with some overall bias for one choice over the other, (2) a "noisy win-stay lose-
shift" model (WSLS) that assumes a win-stay lose-shift policy with some level of randomness,
(3) a two-parameter "RL" model with a consistent learning rate and some inverse temperature
that captures decision noise, (4) a three-parameter "RLCK" model that captures both value-based
and value-independent decision with two parameters for learning rate and choice bias, and an
overall decision noise parameter, (5) a three-parameter "RLγ" model that captures asymmetrical
learning with a learning rate parameter and a scaling parameter for negative outcome learning
and a decision noise parameter, and (6) a four-parameter "RLCKγ" asymmetrical learning and
bias model that includes a choice bias term on top of the "RLγ" (see methods). We fitted these

models to each individual animal and compared the likelihood of each of the six models under

four vehicle control conditions and four drug conditions.

In dopamine modulation conditions, the RLCK model (learning + choice kernel,) and the

RLCKγ (asymmetrical learning and bias) model best fit the behaviors (**Figure 5B,** average AIC

across DA conditions: model 1 random: 73838.03; model 2 WSLS: 55843.26; model 3 RL:

58133.45; model 4 RLCK: 54652.02; model 5 RLγ: 57545.47; model 6 RLCKγ: 54612.56).

Since the RLCKγ model did not significantly improve the model fitting, we decided to use the

more parsimonious RLCK model that has three parameters (learning rate, choice bias, and

decision noise). Then we examined how the reinforcement learning model parameters changed

with dopamine modulation. Previous study has shown that both learning rate and inverse

temperature parameters could induce changes in the overall level of exploration [4]. Therefore,

we asked whether the bidirectional changes in exploration when up- or down-regulating

dopamine receptor activity were due to changes in learning rate or decision noise. The results

revealed a bidirectional effect of dopamine modulation on the inverse temperature parameter

(**Figure 5C-E**). Because RL model data can be non-normally distributed, the Shapiro-Wilk test

of normality was conducted to determine whether the RL parameters were normally distributed.

The result suggested that the parameters were not always normally distributed (decision noise:

vehicle: p= 0.85, APO: p<0.0001; vehicle: p<0.0001, FLU: p=0.09), therefore we will report

both parametric and non-parametric statistics results for RL parameters. Increasing dopamine

receptor activity with apomorphine administration increased inverse temperature and decreased

random noise (**Figure 5C**, Wilcoxon matched-pairs test, p= 0.0165; paired t-test, p= 0.053).

Decreasing dopamine receptor activity with flupenthixol administration decreased inverse

temperature and increased random noise (**Figure 5D**, Wilcoxon matched-pairs test, p= 0.0253;

paired t-test, p= 0.053). Neither dopamine agonist (apomorphine) nor dopamine antagonist

(flupenthixol) significantly influenced learning rate or choice kernel compared to vehicle

controls.

In norepinephrine modulation conditions, the RLCK model (learning + choice kernel,) and the

RLCKγ (asymmetrical learning and bias) model also best fit the behavioral data (**Figure 5F, G,**

average AIC across NE conditions: model 1 random: 66898.83; model 2 WSLS: 56061.3; model

3 RL: 53194.59; model 4 RLCK: 50035.02; model 5 RLγ: 52664.48; model 6

RLCKγ:50027.82). However, since the RLCKγ model did not significantly improve the model

fitting, we decided to use the more parsimonious RLCK model for the NE condition as well. We

found that modulating beta-noradrenergic receptor activity also resulted in changes in the

reinforcement learning model parameter. Up-regulating norepinephrine activity with

isoproterenol significantly decreased the learning rate (**Figure 5H**, Wilcoxon matched-pairs test,

p= 0.0001; paired t-test, p= 0.0008). This result is consistent with the decreased win-stay and

decreased lose-shift behaviors when administered isoproterenol because the lower learning rate

under isoproterenol resulted in reduced outcome sensitivity (**Supplementary figure 1C**), which

means learning less from either a rewarded or a non-rewarded outcome. There were no

significant changes in other parameters when administered isoproterenol compared to vehicle

control. We did not find any significant changes in any reinforcement learning parameters when

administered propranolol compared to vehicle control. Together, these results suggested distinct

roles of dopaminergic drugs in modulating exploration via decision noise and isoproterenol in

modulating exploration via outcome sensitivity.

Discussion

In this study, we pharmacologically up- and down-regulated dopamine and beta-noradrenergic

receptor activity and examined the modulatory effects on exploration in an explore/exploit task.

We used a combination of computational modeling approaches to characterize changes in

exploration and latent cognitive variables that could contribute to exploration. Modulating

dopamine receptor activity revealed a bi-directional modulatory effect on exploration -

increasing dopamine receptor activity decreased the level of exploration and decreasing

dopamine receptor activity increased the level of exploration. Modulation of beta-noradrenergic

receptor activity also influenced how much animals explored - beta-noradrenergic receptor

antagonist (propranolol) decreased the level of exploration. However, when examining the

modulatory effect of beta-noradrenergic receptor activity on exploration across sexes, we found

that modulatory effect was sex-different: isoproterenol selectively decreased exploration in

females whereas propranolol decreased exploration in males. Previously, the role of dopamine

and norepinephrine in mediating exploration were often examined in isolation with different task

designs. This study allowed the direct comparison of the modulatory effect of dopamine and

norepinephrine on the transition between exploration and exploitation in the same task across the

same animals.

Dopamine's function as a neuromodulator has been described in two key ways. First, the phasic

activation of midbrain dopamine neurons is thought to be a key contributor to reinforcement

learning [30,59–62]. Numerous studies across species have shown that dopamine neurons

encoded the reward prediction error (RPE) [30,31]. Ventral tegmental dopamine neuronal

activity and release in ventral striatum is proposed to reflect this error, which is used to update

action values [30–32]. Consistent with this view, some previous pharmacological investigations

in human decision making have observed changes in model-derived learning rates that directly

correlate with dopamine synthesis agonism via L-DOPA [63,64] and dopamine receptor D2/D3

antagonism via amisulpride [26]. However, this framework does not necessarily account for a

large body of literature describing a second way to view dopamine's neuromodulatory function,

as tonic levels influencing motivation, vigor, and cognitive flexibility [48,65], especially when

pharmacologically targeted in frontal cortex. Because pharmacological agents are by definition

modulating receptor activity over long periods of time, this role for tonic dopamine is poised to

be computationally described. Indeed, when pharmacologically or genetically modulating tonic

dopamine function, many studies have noted profound changes in the decision "noise" (inverse

temperature) or preservative errors - consistent with a role for dopamine modulation in cognitive

flexibility in earlier work [35,38,66–69]. Two prior examples combining pharmacological

manipulation and computational modeling in animal tasks explicitly testing the explore-exploit

tradeoff particularly highlight this. Cinotti et al. found that systemic dopamine blockade with

flupenxithol increased the amount of random exploratory choices without affecting learning rate

in a rat three-arm bandit task [38], particularly in situations where uncertainty was manipulated.

Similarly, Ebitz et al. reported that systemic cocaine, which blocks dopamine reuptake, reduced

flexibility and regulated tonic exploration in rhesus macaques [69]. Likewise, we find that

systemic activation of dopamine receptors significantly decreased exploration as defined by our

Hidden Markov model and decision noise as described by our reinforcement learning model,

while antagonism of these receptors increased these computational parameters measuring

flexible behavior. Our data thus provide additional support for the idea that dopamine

modulation across the brain provides more than a learning signal, but also a key signal tuning the

precision or flexibility of behavior.

As noted in our introduction, the roles of dopamine and norepinephrine are sometimes conflated

in the computational neuroscience literature. This is in part because modulation of these systems

can be correlated, and noninvasive measures of catecholamine function, for example blink rate

and pupil diameter, could be ascribed to the endogenous activity of either neuromodulator

(although pharmacological action may target one or the other) [26,53,70–72]. This is further

complicated by the fact that some studies examining the relationship between dopamine and

exploration have used pharmacological manipulations which simultaneously influence both tonic

dopamine and norepinephrine levels, such as psychostimulants or L-DOPA [63,69,73–75]. This

makes it challenging to dissociate the modulatory roles of dopamine and norepinephrine in

exploration i.e. how dopamine and norepinephrine distinctively contribute to changes in

reinforcement learning model parameters, and addressing this challenge was a major goal of our

study.

Surprisingly, we found that when modulating beta-noradrenergic receptor activity changed the

level of exploration, it did so via changes in outcome sensitivity. Previous studies have proposed

the locus coeruleus-norepinephrine (LC-NE) system to regulate exploration by changing decision

noise. Kane et al. chemogenetically stimulated locus coeruleus activity in rats and found that

increased LC tonic activity increased in increased exploration, best explained by an increase in

decision noise [9,76]. Other studies have echoed the findings that norepinephrine levels predicted

the level of random exploration, or decision noise, in the system [53,71,77]. However, in this

study we did not find any changes in the noise parameter (inverse temperature, or how closely

you adhere to your value rule) associated with up- and down-regulation of beta-noradrenergic

activity. Instead, the results revealed some learning rate changes when increasing beta-

noradrenergic receptor activity. Model-free analysis on outcome sensitivity suggested that

increasing or decreasing beta-noradrenergic receptor activity bi-directionally changed outcome

sensitivity. Propranolol increased outcome sensitivity and decreased exploration, which means

the animals were more likely to switch choice after loss and more likely to repeat rewarded

choices. In contrast isoproterenol decreased outcome sensitivity, as seen in reduced win-stay and

lose-shift behavior. This is consistent with the decreased learning rate when administered

isoproterenol because a decrease in learning rate is associated with decreased value learning

from each outcome. Where might this strange association between noradrenergic modulation,

exploration, and outcome sensitivity emerge from? We have previously shown that exploration

as defined by our Hidden Markov model can be driven by either changes in learning rate or

changes in inverse temperature [4], supporting the idea that the exploration we see here could be

driven by long-term changes in the influence of outcomes. Work focused on beta noradrenergic

signaling in the context of aversive learning has highlighted a role for noradrenergic signaling in

extinction [78], over a phasic timeframe of minutes to hours. This suggests the intriguing

possibility that the effect of noradrenergic blockade here may be to prevent extinction of learned

values. In short, although the most common associations in the computational neuroscience

literature are between dopamine and reward learning versus norepinephrine and decision noise,

our current data suggests a more nuanced role for each of these modulators.

Adding an additional layer of nuance to these findings, we find that sex differences are a major

axis of variability in the effect of beta-noradrenergic receptor modulation on the explore-exploit

tradeoff. Increasing NE receptor activity with the beta agonist isoproterenol decreased

exploration only in males, while decreasing NE receptor activity with the beta antagonist

propranolol decreased exploration especially in females. One possible explanation is that the

relationship between exploration and NE receptor activity was nonlinear but instead, followed an

inverted U-shape. Such inverted U-shape correlation has been suggested between cognitive

performance and NE level [23,27,79]. Since females have been reported to have higher tonic NE

levels due to both function and structural differences [50,80], it is possible that the baseline

difference in tonic NE level across sexes could posit individuals on different parts of the inverted

U-curve, where increasing or decreasing NE could affect exploration in one sex without

impacting the other.

Taken together, this study demonstrates the distinct role of dopamine and norepinephrine in

tuning the exploration-exploitation tradeoff when learning about an uncertain environment.

Dopamine modulated exploration via decision noise and norepinephrine modulated exploration

via outcome sensitivity. The systemic manipulation of dopamine and norepinephrine receptor

activity did not allow for more precise modulation of phasic changes in catecholamine levels.

However, the computational tool that we presented here can complement higher resolution

neural recording techniques to examine for phasic changes in dopamine and norepinephrine

activity on a trial-by-trial basis. Finally, this study also highlighted that the modulatory effects of

catecholamine on exploration-exploitation tradeoff are strongly influenced by sex and sex-linked

mechanisms. Therefore, future research should take into account sex-correlated individual

variability when examining the neural correlates of exploration.

Figures and figure legends

Figure 1

Figure 1. Modulation of dopamine receptor activity affected reward-directed behaviors.

Up-regulating dopamine activity increased stickiness in choice behaviors regardless of

outcomes.

A) Schematic of the mouse touchscreen chamber and the trial structure of the two-armed spatial

restless bandit task. **B)** An example of the dynamic reward contingency showing the changing

reward probabilities associated with each option. **C)** The dopaminergic drug administration

schedule and drug dosage used to modulate dopamine receptor activity. A non-selective

dopamine receptor agonist apomorphine (0.1mg/kg) and a non-selective dopamine receptor

antagonist flupenthixol (0.03mg/kg) were systemically administered to respectively up- and

down-regulate dopamine receptor activity. 0.9% saline was used as the vehicle control. Control

and drug sessions were interleaved and repeated for six sessions each. **D)** (left) Average

probability of obtaining reward compared to the chance level probability of reward for each

animal under apomorphine (APO) and vehicle (dot). Error bars depict mean ± SEM across

sessions for each animal. (right) probability of obtaining reward over chance level across APO

(dark green) and vehicle (gray). **E)** (left) Average probability of obtaining reward compared to

the chance level probability of reward for each animal under flupenthixol (FLU) and vehicle

(dot). Error bars depict mean ± SEM across sessions for each animal. (right) probability of

obtaining reward over chance level across FLU (light green) and vehicle (gray). **F)** Average

probability of win-stay on vehicle control or APO. **G)** Average lose-shift on vehicle control or

APO. **H)** Average probability of win-stay on vehicle control or FLU. **I)** Average lose-shift on

vehicle control or FLU. * indicates $p < 0.05$. Graphs depict mean ± SEM across animals unless

specified otherwise.

Figure 2

Figure 2. The modulatory effects of beta-noradrenergic receptor activity on reward-

directed behavior was also influenced by sex.

A) The noradrenergic drug administration schedule and drug dosage used to modulate

noradrenergic receptor activity. A beta-noradrenergic receptor agonist isoproterenol (0.3mg/kg)

and a beta-noradrenergic receptor antagonist propranolol (5mg/kg) were systemically

administered to respectively up- and down-regulate norepinephrine activity. 0.9% saline was

used as the vehicle control. Control and drug sessions were interleaved and repeated for six

sessions each. **B)** (left) Average probability of obtaining reward compared to the chance level

probability of reward for each animal under isoproterenol (ISP) and vehicle (dot). Error bars

depict mean ± SEM across sessions for each animal. (right) probability of obtaining reward over

chance level across ISP (dark brown) and vehicle (gray). **C)** Average response time under ISP

and vehicle. ISP significantly increased response time. **D)** (left) Average probability of obtaining

reward compared to the chance level probability of reward for each animal under propranolol

(PRO) and vehicle (dot). Error bars depict mean ± SEM across sessions for each animal. (right)

probability of obtaining reward over chance level across PRO (light brown) and vehicle (gray).

E) Average response time under PRO and vehicle. PRO significantly decreased response time.

F) Average probability of win-stay under ISP and vehicle. **G)** Average probability of lose-shift

under ISP and vehicle. (inset) probability of lose-shift across treatments across sexes. Decrease

in lose-shift under ISP was primarily driven by changes in females (interaction term). **H)**

Average probability of win-stay under PRO and vehicle. (inset) probability of win-stay across

treatments across sexes. Increase in win-stay under PRO was primarily driven by changes in

males (interaction term). **I)** Average probability of lose-shift under PRO and vehicle. (inset)

probability of lose-shift across treatments across sexes. Decrease in lose-shift under PRO was

primarily driven by changes in males (interaction term). * indicates $p < 0.05$. Graphs depict mean

± SEM across animals unless specified otherwise.

Figure 3

Figure 3. Up- and down-regulation of dopamine activity had bidirectional effects on the

level of exploration.

A) (left) Structure of a Hidden Markov model (HMM) that modeled exploration and exploitation

as latent goal states undering observed choices. This model incorporates two exploit states for

two choices and one explore state where choices were uniformly distributed across options.

(right) Reward probabilities, choices, and HMM labels for an example session of 300 trials for a

given mouse. Shaded areas demonstrated HMM-labeled exploratory choices. **B)** Distribution of

the percentage of HMM-labeled exploratory choices under flupenthixol (FLU)/ vehicle (top) and

apomorphine (APO)/vehicle (bottom). **C)** Probability of exploration for dopamine antagonist

(FLU) and agonist (APO), normalized by their vehicle control. Decreasing dopamine activity

increased exploration and increasing dopamine activity decreased exploration. **D)** Probability of

exploration by session with vehicle and drug session interleaved (flupenthixol: top;

apomorphine: bottom). Drug administration sessions are in colored shades. * indicates p < 0.05.

Graphs depict mean ± SEM across animals.

Figure 4

Figure 4. Downregulating norepinephrine activity influenced exploration but the effect was
modulated by sex.

A) Distribution of the percentage of HMM-labeled exploratory choices under propranolol
(PRO)/ vehicle. **B)** Distribution of the percentage of HMM-labeled exploratory choices under
isoproterenol (ISP)/vehicle. **C)** Probability of exploration for beta-noradrenergic antagonist
(PRO) and agonist (ISP), normalized by their vehicle control. Propranolol decreased the level of
exploration. **D)** Probability of exploration by session with vehicle and PRO session interleaved.
Drug administration sessions are in colored shades. **E)** Probability of exploration by session with
vehicle and ISP session interleaved. Drug administration sessions are in colored shades. **F)**
Probability of exploration across ISP and vehicle across sexes. ISP significantly decreased the
level of exploration in males **G)** Probability of exploration across PRO and vehicle across sexes.
The effect of PRO on exploration was primarily driven by a significant decrease in exploration
under PRO in females. **H)** Cartoon of proposed inverted U-shape relationship between tonic

norepinephrine (NE) level and level of exploration. Sex difference in baseline NE level may

contribute to differential modulation of NE agonist and antagonist across sexes. * indicates $p <$

0.05. Graphs depict mean $\pm$ SEM across animals.

Figure 5

Figure 5. Modulation of dopamine and noradrenergic receptor activity led to changes in

different reinforcement learning (RL) model parameters.

A) A diagram of reinforcement learning (RL) parameters that capture learning rate (α), asymmetric learning (γ), choice bias (α_c), inverse temperature/decision noise (β). The RL models tested included one or more of the above parameters. **B)** Model agreement for six RL models using log likelihood for vehicle and apomorphine (APO) condition (top), and vehicle and flupenthixol condition (bottom). The three-parameter RLCK model was the best fit and most parsimonious model for behaviors. **C)** Average inverse temperature (β) across apomorphine and vehicle. **D)** Average inverse temperature (β) across flupenthixol and vehicle. **E)** Agonizing (apomorphine) and antagonizing (flupenthixol) dopamine activity revealed bidirectional effect on inverse temperature. **F)** Model agreement for six RL models using log likelihood for vehicle and isoproterenol (ISP) condition **G)** Model agreement for six RL models using log likelihood for vehicle and propranolol (PRO) condition. **H**) Average learning rate (α) across isoproterenol and vehicle. * indicates $p < 0.05$. Graphs depict mean ± SEM across animals.

Supplementary figure 1

Figure S1. Effect of dopamine and norepinephrine modulation on outcome sensitivity.

Related to Figure 1.

A) Average outcome sensitivity across flupenxithol (FLU) and vehicle. **B)** Average outcome

sensitivity across isoproterenol (ISP) and vehicle. **C)** Average outcome sensitivity across

propranolol (PRO) and vehicle. * indicates $p < 0.05$. Graphs depict mean ± SEM across animals.

Supplementary figure 2

Figure S2. Effect of dopamine and norepinephrine modulation on exploration in female and male mice. Related to Figure 3,4.

Methods

Animals. Thirty-two BL6129SF1/J mice (16 males and 16 females) were obtained from Jackson

Laboratories (stock #101043). Mice arrived at the lab at 7 weeks of age and adapted to 0900-

2100 hours reversed light cycle throughout the testing phase. Mice were housed in groups of four

with ad libitum access to water while being mildly food restricted (85-95% of free feeding

weight) before the start of the experiment (12 weeks) and during the experiment. All animals

were cared for according to the guidelines of the National Institution of Health and experimental

protocols were as approved by the Institutional Animal Care and Use Committee (IACUC) of the

University of Minnesota.

Apparatus. Sixteen identical triangular touchscreen operant chambers (Lafayette Instrument Co.,

Lafayette, IN) were used for training and testing. Two walls were black acrylic plastic. The third

wall housed the touchscreen and was positioned directly opposite the magazine. The magazine

provided liquid reinforcers (50% Ensure) delivered by a peristaltic pump, typically 7ul (280 ms

pump duration). ABET-II software (Lafayette Instrument Co., Lafayette, IN) was used to

program operant schedules and to analyze all data from training and testing.

Behavioral task. *Two-armed spatial restless bandit task.* Animals were trained to perform a two-

armed spatial restless bandit task in the touchscreen operant chamber. Each trial, animals were

presented with two identical squares on the left and right side of the screen. Nose poke to one of

the target locations on the touchscreen was required to register a response. Each location is

associated with some probability of reward, which changes independently over time. For every

trial, there is a 10% chance that the reward probability of a given arm will increase or decrease

by 10%. All the walks were generated randomly with a few criteria: 1) the overall reward

probabilities of two arms are within 20% of each other, preventing one arm being overly better

than the other, 2) the reward probability cannot go down to 0% or go up to 100%, 3) there are no

30 consecutive trials where the reward probabilities of both arms are lower than 20% to ensure

motivation.

Animals ran a simple deterministic schedule on Monday to re-adapt to the operant chamber after

weekends off and ran a different restless bandit walk each day from Tuesday to Friday. Animals

ran for 2 rounds of 4 consecutive days and within each day, animals completed either 300 trials

or spent a maximum of two hours in the operant chamber. On average across all sessions,

animals performed 276.5 trials with a standard deviation of 8.6 trials (male average: 253.7 trials,

std = 15.4; female average 299.3 trials, std = 0.74. Data was recorded by the ABET II system and

was exported for further analysis. All computational modeling was conducted using python.

Drug administration. To assess the effect of norepinephrine and dopamine receptor activity on

explore behaviors, we used the following four different drugs to increase or decrease beta-

noradrenergic or dopaminergic receptor activity systemically: a beta-noradrenergic receptor

agonist isoproterenol (0.3mg/kg), a beta-noradrenergic receptor antagonist propranolol (5mg/kg),

a non-selective dopamine receptor agonist apomorphine (0.1mg/kg), and a non-selective

dopamine receptor antagonist flupenthixol (0.03mg/kg).

All drugs were fully dissolved in 0.9% saline and protected from light. Animals received

intraperitoneal (IP) injection of drug or saline (control) at an injection volume of 5ml/kg right

before the experiment with the exception of apomorphine being administered 20 minutes before

the experiment due to the immediate drug influence of locomotor behaviors. Drug and saline

were administered in alternating sessions using a within-subject so that every animal received

each of the four drugs. The dopamine receptor agonist and antagonist were administered

counterbalanced across animals. Half of the animals (n = 8) were randomly selected to receive

apomorphine first, the other received flupenthixol first. Dosage chosen was based on previous

studies on the effect of drugs on cognitive functions [38–44]. Doses for each drug were chosen

from previous studies as the lowest doses necessary to produce alterations in cognitive functions

including decision making, learning and exploration with minimal influence on locomotion.

Data Analysis

General analysis techniques. Data was analyzed with custom PYTHON, and Prism 8 scripts.

Generalized linear mixed models (GLMMs), ANOVA, and t-test were used to determine the

fixed effects of drug, sex, and interaction term, accounting for random effects of animal identity

and session, unless otherwise specified. P values were compared against the standard $\alpha = 0.05$

threshold. The sample size is n = 32 (16 males and 16 females) for all statistical tests. No animal

was excluded from the experiment. All statistical tests used and statistical details were reported

in the results. All figures depict mean ± SEM.

Generalized Linear Mixed Models (GLMMs). In order to determine whether drug and sex

predicted the reward acquisition performance, response time, probability of win-stay, probability

of lose-shift, outcome sensitivity, probability of exploration, we fitted a series of generalized

linear mixed model to examine the fixed effects of drug, sex, and interaction term, with animal

identity and session number as random effects.

$Y = \beta_0 + \beta_1(drug) + \beta_2(sex) + \beta_3(drug)(sex) + u_1(animal\ ID) + u_2(session)$ **[1]**

Where Y is the dependent variable (obtained reward, response time, win-stay, lose-shift,

outcome sensitivity, exploration). In this model, β1 describes the main effect of drug and β2

describes the main effect of sex. β3 captures any interaction effect between drug and sex. u1 and

u2 capture the random effect of animal identity and session respectively.

Outcome Sensitivity. In order to examine how much of the switching behavior was sensitive to

reward outcome, we calculated the difference between switching given no reward and switching

given reward control by the total amount of switching.

$$outcome\ sensitivity = \frac{p(switch|no\ reward) - p(switch|reward)}{p(switch)}$$ **[2]**

Hidden Markov model (HMM). In order to identify when animals were exploring, we fit a

Hidden Markov model (HMM) to the animals' choice sequence. Our model consisted of two

hidden states - explore state and exploit state, which are defined by the probability of making

each choice (k, out of N_k possible options) and the probability of transitioning from one state to

another. The HMM models exploration as a uniform distribution over choices, i.e. the emissions

model for the explore state was uniform across options. This is the maximum entropy

distribution of a categorical variable, which makes the few assumptions of true distribution of

choices during exploration and therefore does not bias the model towards or away from any

particular type of high-entropy choice period. In the exploitation state, subjects repeatedly

sample the same choice, therefore the exploit states only permit one choice, i.e. exploit-left state

only emits left choices and exploit-right state only emits right choice.

The latent states in this model are Markovian, meaning that they are time-independent. They

depend only on the most recent state. the HMM model estimates a transition matrix to fit the

behavior by mapping of past and future states. This matrix is a system of stochastic equations

describing the 1-time-step probability of transitioning between every combination of states. In

our model, there were 3 possible states (2 exploit states and 1 explore state). The parameters

were tied across exploit states such that each exploit state had the same probability of beginning

(from exploring) and of sustaining itself. Transitions out of the exploration, into exploitative

states, were also tied. The model also assumed that the mice had to pass through exploration in

order to start exploiting a new option, even if only for a single trial. Through fixing the emissions

model, constraining the structure of the transmission matrix, and tying the parameters, the final

HMM had only two free parameters: one corresponding to the probability of exploring, given

exploration on the last trial, and one corresponding to the probability of exploiting, given

exploitation on the last trial.

The model was fit via expectation-maximization using the Baum Welch algorithm [81]. This

algorithm finds a (possibly local) maxima of the complete-data likelihood. A complete set of

parameters θ includes the emission and transition models, discussed already, but also initial

distribution over states, typically denoted as PI. Because the mice had no knowledge of the

environment at the first trial of the session, the initial distribution started with p (explore state) =

1. The algorithm was reinitialized with random seeds 10 times, and the model that maximized

the observed (incomplete) data log likelihood across all the sessions for each animal was

ultimately taken as the best. To decode latent states from choices, we used the Viterbi algorithm

to discover the most probable posteriori sequence of latent states.

Reinforcement learning (RL) models. We fitted five reinforcement learning (RL) models that

could potentially characterize animals' choice behaviors, with the structure of each RL model

detailed below. AIC weights were calculated from AIC values across all treatment groups and

compared across models to determine the best model with the highest relative likelihood.

The first model (random) assumes that animals choose between two arms randomly with

some overall bias for one side over the other. This choice bias for choosing left side over right

side is captured with a parameter *b*. The probability of choosing left side on trial *t* is:

764                           $p_t^L = b$ [3]   *"random"*

The second model is a noisy win-stay lose-shift (WSLS) model that adapts choices with

regards to outcomes. This model assumes a win-stay lose-shift policy that is to repeat a rewarded

choice and to switch to the other choice if not rewarded. Furthermore, this model includes a

parameter ϵ that captures the level of randomness, allowing a stochastic application of the win-

stay lose-shift policy. The probability of choosing arm k on trial t is:

$$p_t^k = \begin{cases} 1 - \frac{\epsilon}{2}, if\ (c_{t-1} = k\ and\ r_{t-1} = 1\ OR\ c_{t-1} \neq k\ and\ r_{t-1} = 0) \\ \frac{\epsilon}{2}, if\ (c_{t-1} \neq k\ and\ r_{t-1} = 1\ OR\ c_{t-1} = k\ and\ r_{t-1} = 0) \end{cases} \quad \text{[4] } \textit{"noisy WSLS"}$$

c_t indicates the choice on trial t and r_t is a binary variable that indicates whether or not trial t was

rewarded.

The third model (RL) is a basic delta-rule reinforcement learning (RL) model. This two-

parameter model assumes that animals learn by consistently updating Q values, which are values

defined for options (left and right side). These Q values, in turn, dictate what choice to make

next. For example, in a multi-armed bandit task, Q_t^k is the value estimation of how good arm k at

trial t, and is updated based on the reward outcome of each trial:

$$Q_{t+1}^k = Q_t^k + \alpha(r_t - Q_t^k) \quad \text{[5] } \textit{"RL"}$$

In each trial, $r_t - Q_t^k$ captures the reward prediction error (RPE), which is the diffe-rence

between expected outcome and the actual outcome. The parameter a is the learning rate, which

determines the rate of updating RPE. With Q values defined for each arm, choice selection on

each trial was performed based on a Softmax probability distribution:

$$p(a_{t+1} = k) = \frac{e^{\beta Q_t^k}}{\sum_j e^{\beta Q_t^j}}$$

, where inverse temperature β determines the level of random decision noise.

The fourth model (*RLCK*) incorporates a choice updating rules in addition to the value

updating rule in model 3. The model assumes that choice kernel, which captures the outcome-

independent tendency to repeat a previous choice, also influences decision making. The choice

kernel updating rule is similar to the value-updating rule:

$$CK_{t+1}^k = CK_t^k + \alpha_c(a_t^k - CK_t^k) \quad \text{[6] } \textit{"RLCK"}$$

, where a_t^k is a binary variable that indicates whether or not arm k was chosen on trial t and a_t is

choice kernel updating rate, characterizing choice persistence. The value and choice kernel term

were combined to compute the probability of choosing arm k on trial t:

793
$$p_t^k = \frac{e^{\beta(Q_t^k + CK_t^k)}}{\sum_j e^{\beta(Q_t^j + CK_t^j)}}$$

794 , where β is the inverse temperature, capturing the decision noise outside value and choice bias.

795   The fifth model (*RLγ*) is an asymmetrical learning model that incorporates an asymmetric

796 learning scalar parameter that scales the learning rate on the trials where there is no reward. The

797 choice selection is also based on a Softmax probability distribution where the inverse

798 temperature determines the decision noise in the system.

799
$$Q_{t+1}^k = \begin{cases} Q_t^k + \alpha(r_t - Q_t^k), r_t = 1 \\ Q_t^k + \gamma \times \alpha(r_t - Q_t^k), r_t = 0 \end{cases} \qquad [7] \text{ ``RL}\gamma\text{''}$$

The sixth model (*RLCKγ*) is the combination of the fourth and the fifth model,

incorporating both asymmetrical learning and choice kernel. This model included a learning rate,

an asymmetric learning scalar, a choice kernel, and an inverse temperature parameter.

Competing Interests

The authors have declared no competing interests.

Acknowledgements

Thank you to Dr. David Redish and Madison J Merfeld for comments improving this manuscript.

Thank you to Anila Bano for helping with the experiment.

Author contributions

Conceptualization - N.M.G., R.B.E., and C.S.C.; Methodology - C.S.C., and N.M.G.;

Investigation - C.S.C., D.M, and E.K.; Data analyses - C.S.C.; Data interpretation - C.S.C,

R.B.E, and N.M.G; Writing - Original Draft, C.S.C.; Writing - Review & Editing C.S.C., R.B.E.,

D.M., E.K., and N.M.G.; Funding Acquisition - N.M.G. and R.B.E.; Supervision - N.M.G.

Funding

This work was supported by NIMH R01 MH123661 (N.M.G), NIMH P50 MH119569 (N.M.G),

and NIMH T32 training grant MH115886 (C.S.C), startup funds from the University of

Minnesota (N.M.G.), a Young Investigator Grant from the Brain and Behavior Foundation

(R.B.E.), an Unfettered Research Grant from the Mistletoe Foundation (R.B.E.), and Fonds de

Recherche du Québec Santé, Chercheur-Boursier Junior 1, #284309 (R.B.E).

References

1. Grissom NM, Reyes TM. Let's call the whole thing off: evaluating gender and sex differences in executive function. Neuropsychopharmacology. 2019. 2019.
2. Malloy-Diniz LF, Miranda DM, Grassi-Oliveira R. Editorial: Executive Functions in Psychiatric Disorders. Front Psychol. 2017;8:1461.
3. Elliott R. Executive functions and their disorders. Br Med Bull. 2003;65:49–59.
4. Chen CS, Knep E, Han A, Ebitz RB, Grissom N. Sex differences in learning from exploration. Elife. 2021;10.
5. Chen CS, Ebitz RB, Bindas SR, Redish AD, Hayden BY, Grissom NM. Divergent Strategies for Learning in Males and Females. Curr Biol. 2021;31:39–50.e4.
6. Ebitz RB, Albarran E, Moore T. Exploration Disrupts Choice-Predictive Signals and Alters Dynamics in Prefrontal Cortex. Neuron. 2018;97:475.
7. Rothenhoefer KM, Costa VD, Bartolo R, Vicario-Feliciano R, Murray EA, Averbeck BB. Effects of Ventral Striatum Lesions on Stimulus-Based versus Action-Based Reinforcement Learning. J Neurosci. 2017;37:6902–6914.
8. Sutton RS, Barto AG. Reinformcent Learning: An Introduction. 1998.
9. Wilson RC, Bonawitz E, Costa VD, Ebitz RB. Balancing exploration and exploitation with information and randomization. Curr Opin Behav Sci. 2021;38:49–56.
10. Daw ND, O'Doherty JP, Dayan P, Seymour B, Dolan RJ. Cortical substrates for exploratory decisions in humans. Nature. 2006;441:876–879.
11. Niv Y, Joel D, Meilijson I, Ruppin E. Evolution of reinforcement learning in uncertain environments: A simple explanation for complex foraging behaviors. Adapt Behav. 2002;10:5–24.
12. Jepma M, Schaaf JV, Visser I, Huizenga HM. Uncertainty-driven regulation of learning and exploration in adolescents: A computational account. PLoS Comput Biol. 2020;16:e1008276.
13. Gershman SJ. Uncertainty and Exploration. Decision (Wash D C). 2019;6:277–286.
14. Kim H, Lee D, Shin Y-M, Chey J. Impaired strategic decision making in schizophrenia. Brain Res. 2007;1180:90–100.
15. Mussey JL, Travers BG, Klinger LG, Klinger MR. Decision-making skills in ASD: performance on the Iowa Gambling Task. Autism Res. 2015;8:105–114.
16. Addicott MA, Pearson JM, Schechter JC, Sapyta JJ, Weiss MD, Kollins SH. Attention-deficit/hyperactivity disorder and the explore/exploit trade-off. Neuropsychopharmacology. 2020. 11 October 2020. https://doi.org/10.1038/s41386-020-00881-8.
17. Morris LS, Baek K, Kundu P, Harrison NA, Frank MJ, Voon V. Biases in the Explore-Exploit Tradeoff in Addictions: The Role of Avoidance of Uncertainty. Neuropsychopharmacology. 2016;41:940–948.
18. Kaske EA, Chen CS, Meyer C, Yang F, Ebitz B, Grissom N, et al. Prolonged physiological stress is associated with a lower rate of exploratory learning that is compounded by depression. Biol Psychiatry Cogn Neurosci Neuroimaging. 2022. December 2022. https://doi.org/10.1016/j.bpsc.2022.12.004.
19. Kobayashi K. Role of catecholamine signaling in brain and nervous system functions: new insights from mouse molecular genetic study. J Investig Dermatol Symp Proc. 2001;6:115–121.
20. Money KM, Stanwood GD. Developmental origins of brain disorders: roles for dopamine.

Front Cell Neurosci. 2013;7:260.

21. Ressler KJ, Nemeroff CB. Role of norepinephrine in the pathophysiology of
neuropsychiatric disorders. CNS Spectr. 2001;6:663–666, 670.

22. Williams OOF, Coppolino M, George SR, Perreault ML. Sex Differences in Dopamine
Receptors and Relevance to Neuropsychiatric Disorders. Brain Sci. 2021;11.

23. Aston-Jones G, Cohen JD. An integrative theory of locus coeruleus-norepinephrine
function: adaptive gain and optimal performance. Annu Rev Neurosci. 2005;28:403–450.

24. Bouret S, Sara SJ. Network reset: a simplified overarching theory of locus coeruleus
noradrenaline function. Trends Neurosci. 2005;28:574–582.

25. Cohen JD, McClure SM, Yu AJ. Should I stay or should I go? How the human brain
manages the trade-off between exploitation and exploration. Philosophical Transactions of
the Royal Society B: Biological Sciences. 2007;362:933–942.

26. Cremer A, Kalbe F, Müller JC, Wiedemann K, Schwabe L. Disentangling the roles of
dopamine and noradrenaline in the exploration-exploitation tradeoff during human decision-
making. Neuropsychopharmacology. 2022. 15 December 2022.
https://doi.org/10.1038/s41386-022-01517-9.

27. Redish AD, Jensen S, Johnson A, Kurth-Nelson Z. Reconciling reinforcement learning
models with behavioral extinction and renewal: implications for addiction, relapse, and
problem gambling. Psychol Rev. 2007;114:784–805.

28. Seamans JK, Yang CR. The principal features and mechanisms of dopamine modulation in
the prefrontal cortex. Prog Neurobiol. 2004;74:1–58.

29. Ranjbar-Slamloo Y, Fazlali Z. Dopamine and Noradrenaline in the Brain; Overlapping or
Dissociate Functions? Front Mol Neurosci. 2019;12:334.

30. Schultz W. Predictive reward signal of dopamine neurons. J Neurophysiol. 1998;80:1–27.

31. Niv Y. Reinforcement learning in the brain. J Math Psychol. 2009;53:139–154.

32. Schultz W. Updating dopamine reward signals. Curr Opin Neurobiol. 2013;23:229–238.

33. Yu AJ, Dayan P. Uncertainty, neuromodulation, and attention. Neuron. 2005;46:681–692.

34. Amemiya S, Kubota N, Umeyama N, Nishijima T, Kita I. Noradrenergic signaling in the
medial prefrontal cortex and amygdala differentially regulates vicarious trial-and-error in a
spatial decision-making task. Behav Brain Res. 2016;297:104–111.

35. Humphries MD, Khamassi M, Gurney K. Dopaminergic Control of the Exploration-
Exploitation Trade-Off via the Basal Ganglia. Front Neurosci. 2012;6:9.

36. Warren CM, Wilson RC, van der Wee NJ, Giltay EJ, van Noorden MS, Cohen JD, et al. The
effect of atomoxetine on random and directed exploration in humans. PLoS One.
2017;12:e0176034.

37. Grossman CD, Bari BA, Cohen JY. Serotonin neurons modulate learning rate through
uncertainty. bioRxiv. 2020:2020.10.24.353508.

38. Cinotti F, Fresno V, Aklil N, Coutureau E, Girard B, Marchand AR, et al. Dopamine
blockade impairs the exploration-exploitation trade-off in rats. Sci Rep. 2019;9:6770.

39. Grigoryan GA. Serotonin and Impulsivity (animal experiments). Neurosci Behav Physiol.
2012;42:885–894.

40. Ichihara K, Nabeshima T, Kameyama T. Opposite effects induced by low and high doses of
apomorphine on single-trial passive avoidance learning in mice. Pharmacol Biochem Behav.
1988;30:107–113.

41. Nakamura K, Kurasawa M, Tanaka Y. Apomorphine-induced hypoattention in rats and
reversal of the choice performance impairment by aniracetam. Eur J Pharmacol.

1998;342:127–138.

42. Fernandez-Tome MP, Sanchez-Blazquez P, del Rio J. Impairment by apomorphine of one-
trial passive avoidance learning in mice: the opposing roles of the dopamine and
noradrenaline systems. Psychopharmacology . 1979;61:43–47.

43. Cabib S, Puglisi-Allegra S, Oliverio A. Chronic stress enhances apomorphine-induced
stereotyped behavior in mice: involvement of endogenous opioids. Brain Res.
1984;298:138–140.

44. Goldschmidt PL, Frances H, Simon P. Stimulation of beta-adrenergic receptors and
spontaneous motor activity in mice. Pharmacol Biochem Behav. 1984;21:177–180.

45. Weed MR, Gold LH. The effects of dopaminergic agents on reaction time in rhesus
monkeys. Psychopharmacology . 1998;137:33–42.

46. Filipowicz ALS, Levine J, Piasini E, Tavoni G, Kable JW, Gold JI. The complexity of
model-free and model-based learning strategies. 2019. 2019.
https://doi.org/10.1101/2019.12.28.879965.

47. Kool W, McGuire JT, Rosen ZB, Botvinick MM. Decision making and the avoidance of
cognitive demand. J Exp Psychol Gen. 2010;139:665–682.

48. Floresco SB. Prefrontal dopamine and behavioral flexibility: shifting from an 'inverted-U'
toward a family of functions. Front Neurosci. 2013;7:62.

49. Ebitz RB, Tu JC, Hayden BY. Rules warp feature encoding in decision-making circuits.
PLoS Biol. 2020;18:e3000951.

50. Bangasser DA, Wiersielis KR, Khantsis S. Sex differences in the locus coeruleus-
norepinephrine system and its regulation by stress. Brain Res. 2016;1641:177–188.

51. Pearson JM, Hayden BY, Raghavachari S, Platt ML. Neurons in posterior cingulate cortex
signal exploratory decisions in a dynamic multioption choice task. Curr Biol. 2009;19:1532–
1537.

52. Ishii S, Yoshida W, Yoshimoto J. Control of exploitation-exploration meta-parameter in
reinforcement learning. Neural Netw. 2002;15:665–687.

53. Jepma M, Nieuwenhuis S. Pupil Diameter Predicts Changes in the Exploration–Exploitation
Trade-off: Evidence for the Adaptive Gain Theory. J Cogn Neurosci. 2011;23:1587–1596.

54. Katahira K. The statistical structures of reinforcement learning with asymmetric value
updates. J Math Psychol. 2018;87:31–45.

55. Gershman SJ. Empirical priors for reinforcement learning models. J Math Psychol.
2016;71:1–6.

56. Frank MJ, Moustafa AA, Haughey HM, Curran T, Hutchison KE. Genetic triple dissociation
reveals multiple roles for dopamine in reinforcement learning. Proc Natl Acad Sci U S A.
2007;104:16311–16316.

57. Wilson R, Collins A. Ten simple rules for the computational modeling of behavioral data.
2019. 2019.

58. Economides M, Kurth-Nelson Z, Lübbert A, Guitart-Masip M, Dolan RJ. Model-Based
Reasoning in Humans Becomes Automatic with Training. PLoS Comput Biol.
2015;11:e1004463.

59. Barto AG. Adaptive critics and the basal ganglia. Models of Information Processing in the
Basal Ganglia. 1995;382:215–232.

60. Montague PR, Dayan P, Person C, Sejnowski TJ. Bee foraging in uncertain environments
using predictive hebbian learning. Nature. 1995;377:725–728.

61. Montague PR, Dayan P, Sejnowski TJ. A framework for mesencephalic dopamine systems

based on predictive Hebbian learning. J Neurosci. 1996;16:1936–1947.

62. Schultz W, Dayan P, Montague PR. A neural substrate of prediction and reward. Science.
1997;275:1593–1599.

63. Rutledge RB, Lazzaro SC, Lau B, Myers CE, Gluck MA, Glimcher PW. Dopaminergic
drugs modulate learning rates and perseveration in Parkinson's patients in a dynamic
foraging task. J Neurosci. 2009;29:15104–15114.

64. Frank MJ, Seeberger LC, O'reilly RC. By carrot or by stick: cognitive reinforcement
learning in parkinsonism. Science. 2004;306:1940–1943.

65. Beeler JA, Cools R, Luciana M, Ostlund SB, Petzinger G. A kinder, gentler dopamine…
highlighting dopamine's role in behavioral flexibility. Front Neurosci. 2014;8:4.

66. Eisenegger C, Naef M, Linssen A, Clark L, Gandamaneni PK, Müller U, et al. Role of
dopamine D2 receptors in human reinforcement learning. Neuropsychopharmacology.
2014;39:2366–2375.

67. Lee E, Seo M, Dal Monte O, Averbeck BB. Injection of a dopamine type 2 receptor
antagonist into the dorsal striatum disrupts choices driven by previous outcomes, but not
perceptual inference. J Neurosci. 2015;35:6298–6306.

68. Beeler JA, Daw N, Frazier CRM, Zhuang X. Tonic dopamine modulates exploitation of
reward learning. Front Behav Neurosci. 2010;4:170.

69. Ebitz RB, Sleezer BJ, Jedema HP, Bradberry CW, Hayden BY. Tonic exploration governs
both flexibility and lapses. PLoS Comput Biol. 2019;15:e1007475.

70. Sescousse G, Ligneul R, van Holst RJ, Janssen LK, de Boer F, Janssen M, et al.
Spontaneous eye blink rate and dopamine synthesis capacity: preliminary evidence for an
absence of positive correlation. Eur J Neurosci. 2018;47:1081–1086.

71. Ebitz RB, Moore T. Both a Gauge and a Filter: Cognitive Modulations of Pupil Size. Front
Neurol. 2018;9:1190.

72. Findling C, Skvortsova V, Dromnelle R, Palminteri S, Wyart V. Computational noise in
reward-guided learning drives behavioral variability in volatile environments. Nat Neurosci.
2019;22:2066–2077.

73. Chakroun K, Mathar D, Wiehler A, Ganzer F, Peters J. Dopaminergic modulation of the
exploration/exploitation trade-off in human decision-making. Elife. 2020;9.

74. Pessiglione M, Seymour B, Flandin G, Dolan RJ, Frith CD. Dopamine-dependent prediction
errors underpin reward-seeking behaviour in humans. Nature. 2006;442:1042–1045.

75. Orsini CA, Willis ML, Gilbert RJ, Bizon JL, Setlow B. Sex differences in a rat model of
risky decision making. Behav Neurosci. 2016;130:50–61.

76. Kane GA, Vazey EM, Wilson RC, Shenhav A, Daw ND, Aston-Jones G, et al. Increased
locus coeruleus tonic activity causes disengagement from a patch-foraging task. Cogn Affect
Behav Neurosci. 2017;17:1073–1083.

77. Joshi S, Gold JI. Pupil Size as a Window on Neural Substrates of Cognition. Trends Cogn
Sci. 2020;24:466–480.

78. Iordanova MD, Yau JO-Y, McDannald MA, Corbit LH. Neural substrates of appetitive and
aversive prediction error. Neurosci Biobehav Rev. 2021;123:337–351.

79. Mäki-Marttunen V, Andreassen OA, Espeseth T. The role of norepinephrine in the
pathophysiology of schizophrenia. Neurosci Biobehav Rev. 2020;118:298–314.

80. Pinos H, Collado P, Rodríguez-Zafra M, Rodríguez C, Segovia S, Guillamón A. The
development of sex differences in the locus coeruleus of the rat. Brain Res Bull.
2001;56:73–78.

81. Bilmes J. A Gentle Tutorial of the EM Algorithm and its Application to Parameter
Estimation for Gaussian Mixture and Hidden Markov Models. 1998. 1998.