Title: Dopamine and norepinephrine differentially mediate the exploration-exploitation tradeoff

Abbreviated title: Dopamine and norepinephrine mediate exploration

Authors: Cathy S. Chen[+1], Dana Mueller[1], Evan Knep[1], R. Becket Ebitz[2], Nicola M. Grissom[+1]

Affiliations: [1] Department of Psychology, University of Minnesota, Minneapolis MN 55455

[2] Department of Neurosciences, Université de Montréal, Quebec, Canada

Corresponding author[+]: Cathy Chen (chen5388@umn.edu)

Nicola Grissom (ngrissom@umn.edu)

Number of pages: 59

Number of figures: 5

Word count: Abstract (249), Introduction (772), Discussion (1476).

Competing Interests:

The authors have declared no competing interests.

Acknowledgements

This work was supported by NIMH R01 MH123661 (N.M.G), NIMH P50 MH119569 (N.M.G),

and NIMH T32 training grant MH115886 (C.S.C), startup funds from the University of

Minnesota (N.M.G.), a Young Investigator Grant from the Brain and Behavior Foundation

(R.B.E.), an Unfettered Research Grant from the Mistletoe Foundation (R.B.E.), and Fonds de

Recherche du Québec Santé, Chercheur-Boursier Junior 1, #284309 (R.B.E). Thank you to Dr.

David Redish and Madison J Merfeld for comments improving this manuscript. Thank you to

Anila Bano for helping with the experiment.

Abstract

The catecholamines dopamine (DA) and norepinephrine (NE) have been implicated in

neuropsychiatric vulnerability, in part via their roles in mediating the decision making processes.

Although the two neuromodulators share a synthesis pathway and are co-activated, they engage

in distinct circuits and roles in modulating neural activity across the brain. However, in the

computational neuroscience literature, they have been assigned similar roles in modulating the

exploration-exploitation tradeoff. Revealing how each neuromodulator contributes to this

explore-exploit process is important in guiding mechanistic hypotheses emerging from

computational psychiatric approaches. To understand the differences and overlaps of the roles of

dopamine and norepinephrine in mediating exploration, a direct comparison using the same

dynamic decision making task is needed. Here, we ran mice in a restless bandit task, which

encourages both exploration and exploitation. We systemically administered a nonselective DA

antagonist (flupenthixol), a nonselective DA agonist (apomorphine), a NE beta-receptor

antagonist (propranolol), and a NE beta-receptor agonist (isoproterenol), and examined changes

in exploration within subjects across sessions. We found a bidirectional modulatory effect of

dopamine receptor activity on exploration - increasing dopamine activity decreased exploration

and decreasing dopamine activity increased exploration. The modulation of exploration via beta-

noradrenergic activity was mediated by sex. Computational model parameters revealed that

dopamine modulation affected exploration via decision noise and norepinephrine modulation via

outcome sensitivity. Together, these findings suggested that the mechanisms that govern the

transition between exploration and exploitation are sensitive to changes in both catecholamine

functions and revealed differential roles for NE and DA in mediating exploration.

Significance Statement

Both dopamine (DA) and norepinephrine (NE) has been implicated in the decision making

process. Although these two catecholamines have shared aspects of their biosynthetic pathways

and projection targets, they are thought to exert many core functions via distinct neural targets

and receptor subtypes. However, the computational neuroscience literature often ascribes similar

roles to these catecholamines, despite the above evidence. Resolving this discrepancy is

important in guiding mechanistic hypotheses emerging from computational psychiatric

approaches. This study examines the role of dopamine and norepinephrine on the explore-exploit

tradeoff. By testing mice, we were able to compare multiple pharmacological agents within

subjects, and examine source of individual differences, allowing direct comparison between the

effects of these two catecholamines in modulating decision making.

Introduction

Dysfunctions in cognitive processes, particularly within the domain of executive function, are implicated in numerous neuropsychiatric disorders (Elliott, 2003; Malloy-Diniz et al., 2017; Grissom and Reyes, 2019). One essential aspect of executive function is value-based decision making. Decision making involving exploring and learning in an uncertain environment has been a focus in computational neuroscience, which has developed approaches for discovering latent structures in decision making strategies (Rothenhoefer et al., 2017; Ebitz et al., 2018; Chen et al., 2021a, 2021b). Many computational models for value-based decision making distinguish two essential latent processes: exploration and exploitation (Sutton and Barto, 1998; Niv et al., 2002; Daw et al., 2006; Jepma et al., 2020; Wilson et al., 2021). Exploration is the ubiquitous learning process across species that allows the discovery of the best action to take in an uncertain environment (Ebitz et al., 2018; Gershman, 2019; Chen et al., 2021b; Wilson et al., 2021). Once a favorable action or option is discovered, exploitation, i.e. repeating a rewarding action, is necessary to obtain the best rates of reward. However, exploitation must be balanced with continued exploration as environments and their reward probabilities change. Dysregulation in this exploration-exploitation tradeoff is implicated in the phenotypes of numerous neuropsychiatric disorders and challenges, including schizophrenia, autism spectrum disorders, addictions, and chronic stress (Kim et al., 2007; Mussey et al., 2015; Morris et al., 2016; Addicott et al., 2020; Kaske et al., 2022). Understanding the fundamental mechanisms impacting the computations in our brain that underlie the balance between exploration and exploitation could help identify critical circuits that are associated with differential risk factors for neuropsychiatric disorders and open avenues for novel interventions for executive function challenges.

Many neuropsychiatric challenges are associated with dysfunction in the catecholamines

dopamine and norepinephrine (Kobayashi, 2001; Ressler and Nemeroff, 2001; Aston-Jones and

Cohen, 2005; Bouret and Sara, 2005; Money and Stanwood, 2013; Addicott et al., 2020;

Williams et al., 2021). These neuromodulators are well-positioned to carry information about the

state of the environment and influence behavior outputs via downstream control of action

selection (Cohen et al., 2007; Wilson et al., 2021). Indeed, dopamine and norepinephrine have

each separately been implicated in mediating the exploration-exploitation tradeoff (Seamans and

Yang, 2004; Aston-Jones and Cohen, 2005; Bouret and Sara, 2005; Cohen et al., 2007; Redish et

al., 2007; Wilson et al., 2021; Cremer et al., 2022). While dopamine and norepinephrine share

biosynthetic pathways and are co-activated under states of arousal, they act through partially

separable circuits and have distinct neural targets (Ranjbar-Slamloo and Fazlali, 2019). However,

in the computational neuroscience literature, dopamine and norepinephrine have largely been

ascribed similar or even identical roles in mediating the explore-exploit tradeoff despite of the

distinct pharmacological profiles.

We have previously shown that multiple latent cognitive processes, such as the

reinforcement learning model-derived parameters of learning rate and decision noise, can

describe differences in the level of exploration between groups (Chen et al., 2021b). Dopamine

has often been implicated in mediating reward prediction errors and value-based learning

(Schultz, 1998, 2013; Daw et al., 2006; Niv, 2009) while norepinephrine in modulating arousal,

attention, and value assessment (Aston-Jones and Cohen, 2005; Yu and Dayan, 2005; Amemiya

et al., 2016), but more recently both neuromodulators have been implicated in the process of

action selection (Humphries et al., 2012; Warren et al., 2017). Therefore, it is unknown whether

dopamine and norepinephrine govern this exploration-exploitation tradeoff via distinct or similar

latent cognitive processes because they have been largely studied in this context in isolation

(Cremer et al., 2022).

To uncover the distinct roles of dopamine and norepinephrine in the exploration-

exploitation tradeoff, we conducted within-subjects pharmacological manipulations for these two

neuromodulators to examine their effect on exploration and the latent cognitive parameters that

influence exploration, in the same restless bandit task. We pharmacologically up- and down-

regulated dopaminergic and noradrenergic receptor activity and compared the modulatory effects

on exploration in the same dynamic decision making task, a spatial restless two-armed bandit

(Ebitz et al., 2018; Chen et al., 2021b). We found a bidirectional modulatory effect of dopamine

receptor activity on the level of exploration. Increasing dopamine activity decreased exploration

and decreasing dopamine activity increased exploration. Beta-noradrenergic receptor activity

also modulated exploration but the modulatory effect was mediated by sex. Contrary to

dopamine modulating choice solely by changing value-based learning for outcomes, via

reinforcement learning modeling we find that dopamine mediates exploration by changing

decision noise, i.e. the precision of value-based choice selection. In contrast, noradrenergic

activity surprisingly influenced exploration by changing value-based learning for outcomes in a

sex dependent manner. These data suggest differential roles of dopamine and norepinephrine in

mediating exploration, and complex roles for both catecholamines in signaling reward.

Materials and Methods

<u>Animals.</u> Thirty-two BL6129SF1/J mice (16 males and 16 females) were obtained from Jackson

Laboratories (stock #101043). Mice arrived at the lab at 7 weeks of age and adapted to 0900-

2100 hours reversed light cycle throughout the testing phase. Mice were housed in groups of four

with ad libitum access to water while being mildly food restricted (85-95% of free feeding

weight) before the start of the experiment (12 weeks) and during the experiment. All animals

were cared for according to the guidelines of the National Institution of Health and experimental

protocols were approved by the Institutional Animal Care and Use Committee (IACUC) of the

University of Minnesota.

<u>Apparatus.</u> Sixteen identical triangular touchscreen operant chambers (Lafayette Instrument Co.,

Lafayette, IN) were used for training and testing. Two walls were black acrylic plastic. The third

wall housed the touchscreen and was positioned directly opposite the magazine. The magazine

provided liquid reinforcers (50% Ensure) delivered by a peristaltic pump, typically 7ul (280 ms

pump duration). ABET-II software (Lafayette Instrument Co., Lafayette, IN) was used to

program operant schedules and to analyze all data from training and testing.

<u>Behavioral task</u>. *Two-armed spatial restless bandit task.* Animals were trained to perform a two-

armed spatial restless bandit task in the touchscreen operant chamber. Each trial, animals were

presented with two identical squares on the left and right side of the screen. Nose poke to one of

the target locations on the touchscreen was required to register a response. Each location is

associated with some probability of reward, which changes independently over time. For every

trial, there is a 10% chance that the reward probability of a given arm will increase or decrease

by 10%. All the walks were generated randomly with a few criteria: 1) the overall reward

probabilities of two arms are within 20% of each other, preventing one arm being overly better

than the other, 2) the reward probability cannot go down to 0% or go up to 100%, 3) there are no

30 consecutive trials where the reward probabilities of both arms are lower than 20% to ensure

motivation.

Animals ran for 6 control sessions and 6 drug sessions for each pharmacological

manipulation. The wash-out period between each drug condition is 3 days. Each session

consisted of 300 trials. Within each session, animals completed either 300 trials or spent a

maximum of two hours in the operant chamber. On average across all sessions, animals

performed 276.5 trials with a standard deviation of 8.6 trials (male average: 253.7 trials, std =

15.4; female average 299.3 trials, std = 0.74. Data were recorded by the ABET II system and was

exported for further analysis. All computational modeling was conducted using python.

<u>Drug administration.</u> To assess the effect of norepinephrine and dopamine receptor activity on

explore behaviors, we used the following four different drugs to increase or decrease beta-

noradrenergic or dopaminergic receptor activity systemically: a beta-noradrenergic receptor

agonist isoproterenol (0.3mg/kg), a beta-noradrenergic receptor antagonist propranolol (5mg/kg),

a non-selective dopamine receptor agonist apomorphine (0.1mg/kg), and a non-selective

dopamine receptor antagonist flupenthixol (0.03mg/kg).

All drugs were fully dissolved in 0.9% saline and protected from light. Animals received

intraperitoneal (IP) injection of drug or saline (control) at an injection volume of 5ml/kg right

before the experiment with the exception of apomorphine being administered 20 minutes before

the experiment due to the immediate drug influence of locomotor behaviors. Drug and saline

were administered in alternating sessions using a within-subject so that every animal received

each of the four drugs. The dopamine receptor agonist and antagonist were administered

counterbalanced across animals. Half of the animals (n = 8) were randomly selected to receive

apomorphine first, the other received flupenthixol first. Since this experiment prioritized using a

within-subject design to understand the effect of each drug within individual animals and the

source of individual differences, the length of study only limited us to test one dose for each of

the four drugs. Therefore, dosage chosen was based on previous studies on the effect of drugs on

cognitive functions (Fernandez-Tome et al., 1979; Cabib et al., 1984; Goldschmidt et al., 1984;

Ichihara et al., 1988; Nakamura et al., 1998; Grigoryan, 2012; Cinotti et al., 2019). Doses for

each drug were chosen from previous studies as the lowest doses necessary to produce alterations

in cognitive functions including decision making, learning and exploration with minimal

influence on locomotion.

Data Analysis

General analysis techniques. Data were analyzed with custom PYTHON, and Prism 8 scripts.

Generalized linear mixed models (GLMMs), ANOVA, and t-test were used to determine the

fixed effects of drug, sex, and interaction term, accounting for random effects of animal identity

and session, unless otherwise specified. P values were compared against the standard $\alpha = 0.05$

threshold. The sample size is n = 32 (16 males and 16 females) for all statistical tests. No animal

was excluded from the experiment. All statistical tests used and statistical details were reported

in the results. All figures depict mean ± SEM.

Generalized Linear Mixed Models (GLMMs). In order to determine whether drug and sex

predicted the reward acquisition performance, response time, probability of win-stay, probability

of lose-shift, outcome sensitivity, probability of exploration, we fitted a series of generalized

linear mixed model to examine the fixed effects of drug, sex, and interaction term, with animal

identity and sessions as random effects.

$Y = \beta_0 + \beta_1(drug) + \beta_2(sex) + \beta_3(drug)(sex) + u_1(animal\ ID) + u_2(session)$ **[1]**

,where Y is the dependent variable (obtained reward, response time, win-stay, lose-shift,

outcome sensitivity, exploration). In this model, β1 describes the main effect of drug and β2

describes the main effect of sex. β3 captures any interaction effect between drug and sex. u1 and

u2 capture the random effect of animal identity and session respectively.

Outcome Sensitivity. In order to examine how much of the switching behavior was sensitive to

reward outcome, we calculated the difference between switching given no reward and switching

given reward control by the total amount of switching. If switching behavior was not affected by

the outcome at all, the outcome sensitivity should be close to zero.

$outcome\ sensitivity = \frac{p(switch|no\ reward) - p(switch|reward)}{p(switch)}$ **[2]**

Hidden Markov model (HMM).

In order to identify when animals were exploring, we fit a Hidden Markov model (HMM) to the

animals' choice sequence. Our model consisted of two hidden states - explore state and exploit

state, which are defined by the probability of making each choice (k, out of N_k possible options)

and the probability of transitioning from one state to another. When an HMM model is fitted to

animals' choices, it estimates a transition matrix, which is the mapping between every state at

one time point and every state at some future point of time. The HMM model fitting is the

process of estimating how the distribution over states will evolve over time. The transition

matrix estimates the probability of each type of transition between states. The HMM models

exploration as a uniform distribution over choices, i.e. the emissions model for the explore state

was uniform across options. This is the maximum entropy distribution of a categorical variable,

which makes the few assumptions of true distribution of choices during exploration and therefore

does not bias the model towards or away from any particular type of high-entropy choice period.

In the exploitation state, subjects repeatedly sample the same choice, therefore the exploit states

only permit one choice, i.e. exploit-left state only emits left choices and exploit-right state only

emits right choice.

The latent states in this model are Markovian, meaning that they are time-independent.

They depend only on the most recent state. The HMM model estimates a transition matrix to fit

the behavior by mapping of past and future states. This matrix is a system of stochastic equations

describing the 1-time-step probability of transitioning between every combination of states. In

our model, there were 3 possible states (2 exploit states and 1 explore state). The parameters

were tied across exploit states such that each exploit state had the same probability of beginning

(from exploring) and of sustaining itself. Transitions out of the exploration, into exploitative

states, were also tied. The model also assumed that the mice had to pass through exploration in

order to start exploiting a new option, even if only for a single trial. Through fixing the emissions

model, constraining the structure of the transmission matrix, and tying the parameters, the final

HMM had only two free parameters: one corresponding to the probability of exploring, given

exploration on the last trial, and one corresponding to the probability of exploiting, given

exploitation on the last trial.

The model was fit via expectation-maximization using the Baum Welch algorithm

(Bilmes, 1998). This algorithm finds a (possibly local) maxima of the complete-data likelihood.

A complete set of parameters θ includes the emission and transition models, discussed already,

but also initial distribution over states, typically denoted as PI. Because the mice had no

knowledge of the environment at the first trial of the session, the initial distribution started with p

(explore state) = 1. The algorithm was reinitialized with random seeds 10 times, and the model

that maximized the observed (incomplete) data log likelihood across all the sessions for each

animal was ultimately taken as the best. To decode latent states from choices, we used the

Viterbi algorithm to discover the most probable posteriori sequence of latent states.

Reinforcement learning (RL) models. We fitted five reinforcement learning (RL) models that

could potentially characterize animals' choice behaviors, with the structure of each RL model

detailed below. AIC weights were calculated from AIC values across all treatment groups and

compared across models to determine the best model with the highest relative likelihood. Model

agreement was calculated as the probability of choice given model to demonstrate how well the

best fit model captures actual choices.

The first model (random) assumes that animals choose between two arms randomly with

some overall bias for one side over the other. This choice bias for choosing left side over right

side is captured with a parameter *b*. The probability of choosing left side on trial *t* is:

$$p_t^L = b \qquad\qquad [3] \ \textit{"random"}$$

The second model is a noisy win-stay lose-shift (WSLS) model that adapts choices with

regards to outcomes. This model assumes a win-stay lose-shift policy that is to repeat a rewarded

choice and to switch to the other choice if not rewarded. Furthermore, this model includes a

parameter ϵ that captures the level of randomness, allowing a stochastic application of the win-

stay lose-shift policy. The probability of choosing arm k on trial t is:

$$p_t^k = \begin{cases} 1 - \frac{\epsilon}{2}, if \ (c_{t-1} = k \ and \ r_{t-1} = 1 \ OR \ c_{t-1} \neq k \ and \ r_{t-1} = 0) \\ \frac{\epsilon}{2}, if \ (c_{t-1} \neq k \ and \ r_{t-1} = 1 \ OR \ c_{t-1} = k \ and \ r_{t-1} = 0) \end{cases}$$ [4] *"noisy WSLS"*

, c_t indicates the choice on trial t and r_t is a binary variable that indicates whether or not trial t

was rewarded.

The third model (RL) is a basic delta-rule reinforcement learning (RL) model. This two-

parameter model assumes that animals learn by consistently updating Q values, which are values

defined for options (left and right side). These Q values, in turn, dictate what choice to make

next. For example, in a multi-armed bandit task, Q_t^k is the value estimation of how good arm k at

trial t, and is updated based on the reward outcome of each trial:

$$Q_{t+1}^k = Q_t^k + \alpha(r_t - Q_t^k)$$ [5] *"RL"*

In each trial, $r_t - Q_t^k$ captures the reward prediction error (RPE), which is the diffe-rence

between expected outcome and the actual outcome. The parameter a is the learning rate, which

determines the rate of updating RPE. With Q values defined for each arm, choice selection on

each trial was performed based on a Softmax probability distribution:

$$p(a_{t+1} = k) = \frac{e^{\beta Q_t^k}}{\sum_j e^{\beta Q_t^j}}$$

, where inverse temperature β determines the level of random decision noise.

The fourth model (*RLCK*) incorporates a choice updating rules in addition to the value

updating rule in model 3. The model assumes that choice kernel, which captures the outcome-

independent tendency to repeat a previous choice, also influences decision making. The choice

kernel updating rule is similar to the value-updating rule:

$$CK_{t+1}^k = CK_t^k + \alpha_c(a_t^k - CK_t^k)$$ [6] *"RLCK"*

, where a_t^k is a binary variable that indicates whether or not arm k was chosen on trial t and a_t is

choice kernel updating rate, characterizing choice persistence. The value and choice kernel term

were combined to compute the probability of choosing arm k on trial t:

282
$$p_t^k = \frac{e^{\beta(Q_t^k + CK_t^k)}}{\sum_j e^{\beta(Q_t^j + CK_t^j)}}$$

283   , where β is the inverse temperature, capturing the decision noise outside value and choice bias.

284   The fifth model ($RL\gamma$) is an asymmetrical learning model that incorporates an asymmetric

285   learning scalar parameter that scales the learning rate on the trials where there is no reward. The

286   choice selection is also based on a Softmax probability distribution where the inverse

287   temperature determines the decision noise in the system.

288
$$Q_{t+1}^k = \begin{cases} Q_t^k + \alpha(r_t - Q_t^k), r_t = 1 \\ Q_t^k + \gamma \times \alpha(r_t - Q_t^k), r_t = 0 \end{cases} \qquad [7] \text{ ``}RL\gamma\text{''}$$

The sixth model ($RLCK\gamma$) is the combination of the fourth and the fifth model,

incorporating both asymmetrical learning and choice kernel. This model included a learning rate,

an asymmetric learning scalar, a choice kernel, and an inverse temperature parameter.

Results

Aged-matched adult male and female wildtype mice (n = 32, 16 males and 16 females,

strain B6129SF1/J) were trained to perform a restless two-armed spatial bandit task in touch-

screen operant chambers (**Figure 1A**). In this task, animals were presented with two identical

visual targets (squares) on the left and right side of the screen during each trial. They indicated

their choices by nose poking at one of the two target locations, each of which provided some

probability of reward that changed independently and randomly over time (**Figure 1B**). The

dynamic reward contingency of this task naturally encourages the animal to balance between

exploiting a favorable option when it is found and exploring to gain information about potential

better alternatives. This task has been employed in rodents and primates as well as human

subjects to understand learning and exploration and have successfully revealed divergent

exploration strategies (Ebitz et al., 2018; Grossman et al., 2020; Chen et al., 2021b). In this

study, we adopted this task to understand the modulatory effect of catecholamine receptor

activity on the transition between exploration and exploitation.

To examine the distinct and/or overlapping roles of dopamine and norepinephrine in

modulating exploration, we tested the systemic effect of a non-selective dopamine receptor

agonist apomorphine (0.1mg/kg), a non-selective dopamine receptor antagonist flupenthixol

(0.03mg/kg), a beta-noradrenergic receptor agonist isoproterenol (0.3mg/kg), and a beta-

noradrenergic receptor antagonist propranolol (5mg/kg) on exploration. The experiment used a

within-subject design, where all animals received each of the four drugs tested. This design

allowed us to directly compare the effect of each drug within individual animals and dissociate

drug effects from potential cohort differences. Since each animal received all four drugs, the

length of study did not allow us to test multiple doses for each drug. Therefore, we selected the

dosages of the drugs based on numerous previous studies on the role of dopamine and

norepinephrine in cognitive processes (Fernandez-Tome et al., 1979; Cabib et al., 1984;

Goldschmidt et al., 1984; Ichihara et al., 1988; Nakamura et al., 1998; Grigoryan, 2012; Cinotti

et al., 2019). Animals were intraperitoneally (IP) administered either saline (control) or treatment

with 5 mg/kg of propranolol, 0.3mg/kg of isoproterenol, or 0.03 mg/kg of flupenthixol

immediately prior to the bandit task or 0.1mg/kg of apomorphine 30 minutes prior to the bandit

task. Saline and drug treatments were administered interleaved by session (**Figure 1C, 2A**). Mice

performed 12 sessions under each pharmacological condition (6 control and 6 drug treatment

sessions). Each drug effect was compared to its own vehicle control collected in the interleaved

sessions.

To examine whether mice had learned to perform the task and were able to sustain

reward, we first calculated the average probability of obtaining reward across sessions in all

control and drug groups. It's worth noting that in this task, there is no asymptotic performance

because the animals need to constantly adapt to the changing reward contingency and any one

option is not always the best choice to make. At any given time, an animal could decide to repeat

a favorable option or to explore the options. The performance is best measured by the amount of

reward acquired. Because reward schedules were dynamic and stochastic, sessions could differ

slightly in the amount of reward that was available. We therefore compared the probability of

actual obtained reward against the probability of reward if chosen randomly (chance). In the

dopamine modulation condition, both control groups (saline) and treatment groups (apomorphine

and flupenthixol) were able to obtain reward more frequently than chance (**Figure 1D, E**, one-

sample t test, vehicle control (apomorphine): $t(31) = 13.96$, $p<0.0001$; apomorphine: $t(31)=$

17.24, $p<0.0001$; vehicle control (flupenthixol): $t(31)=12.36$, $p<0.0001$; flupenthixol:

t(31)=16.48, p<0.0001). When modulating beta-noradrenergic receptor activity, both control

groups (saline) and treatment groups (isoproterenol and propranolol) were able to obtain reward

more frequently than chance (**Figure 2B, D**, one-sample t test, vehicle control (isoproterenol):

t(31)=10.01, p<0.0001; isoproterenol: t(31)=13.84, p<0.0001; vehicle control (propranolol):

t(31)=10.60, p<0.0001; propranolol: t(31)=20.08, p<0.0001). These results suggested that

animals were not choosing randomly. The animals demonstrated a strong understanding of the

task and were able to effectively obtain rewards above chance under baseline (vehicle control)

and all drug treatments.

Next, we asked whether manipulating dopaminergic receptor activity influenced their

reward acquisition performance. We used a generalized linear mixed model (GLMM) to estimate

effects and interactions of drug treatments and sex, with session and animal identity as random

effects (**Equation 1**, see methods). Increasing dopamine receptor activity with apomorphine

administration did not significantly alter their reward acquisition performance (**Figure 1D**,

GLMM, main effect of drug, p= 0.11, $\beta1$ = -0.017). Antagonizing dopamine receptor activity

with flupenthixol administration also did not significantly alter reward acquisition performance

(**Figure 1E**, GLMM, main effect of drug, p= 0.28, $\beta1$ = -0.008). Increasing beta-noradrenergic

receptor activity with isoproterenol administration did not affect the amount of reward obtained

(**Figure 2B**, GLMM, main effect of drug, p= 0.93, $\beta1$ = -0.0005). However, when decreasing

beta-noradrenergic receptor activity with propranolol administration, animals obtained

significantly more reward (**Figure 2D**, GLMM, main effect of drug, p < 0.0001, $\beta1$ = -0.497).

Even though manipulating dopamine receptor activity and increasing beta-noradrenergic receptor

activity did not affect how much overall obtained reward, we considered the possibility that the

tuning of catecholamine receptor activity affected latent exploration strategies underlying choice

patterns, but not an observable change in the level of accuracy. Our previous study demonstrated

that similar reward acquisition performance can be achieved via divergent explore strategies

(Chen et al., 2021b).

Pharmacological manipulations of both dopaminergic and noradrenergic receptor activity

have been shown to have a dose-dependent effect on motor functions (Goldschmidt et al., 1984;

Weed and Gold, 1998). In this study, dosage selection was based on previous literature

examining cognitive functions and a low dose was used to have influence on cognition but avoid

motor function impairment (Nakamura et al., 1998; Cinotti et al., 2019). One commonly used

behavioral metric to examine motor function in a cognitive task is response time. Therefore, we

calculated the response time, which was the time elapsed between the onset of choices and nose

poke response to make a choice, as recorded by the touchscreen operant chamber. If the

administration of drugs impaired motor function, we would expect to see longer response time in

the drug group compared to vehicle control. Upregulating dopamine receptor activity with

apomorphine did not significantly influence the response time (GLMM, main effect of drug,

$p=0.77$, $\beta1 = 0.10$). Similarly, downregulating dopamine receptor activity with flupenthixol did

not significantly alter animals' response time (GLMM, main effect of drug, $p=0.96$, $\beta1 = -0.01$).

Modulating beta-noradrenergic receptor activity resulted in bidirectional changes in response

time. When increasing beta-noradrenergic receptor activity with isoproterenol, both males and

females took significantly longer to respond (**Figure 2C,** GLMM, main effect of drug, $p<0.0001$,

$\beta1 = -3.51$). Decreasing beta-noradrenergic receptor activity with propranolol significantly

decreased the response time (**Figure 2E,** GLMM, main effect of drug, $p=0.04$, $\beta1 = 0.51$),

primarily driven by response time reduction in females under propranolol compared to vehicle

controls (GLMM, interaction term, $p<0.009$, $\beta3 = -0.57$). Faster response time under propranolol

could be associated with more reward obtained - propranolol was the only drug treatment that

altered reward acquisition performance among the four drugs tested. Previous studies have

linked response time with the complexity of the strategy used (Kool et al., 2010; Filipowicz et

al., 2019; Chen et al., 2021b). Faster response time under propranolol might also point to the

possibility that decreasing beta-noradrenergic receptor activity brought on changes in strategies

that took shorter time to execute. Isoproterenol was a highly significant predictor for lower

response time in both male and female mice, implicating the potential drug effect on motor

behaviors.

Opposing modulatory effect of dopamine and norepinephrine receptor activity on win-stay

lose-shift behaviors

The role of dopamine has been heavily studied and shown to be a key contributor to

value-based decision making and reinforcement learning (Seamans and Yang, 2004; Redish et

al., 2007; Niv, 2009; Floresco, 2013). Although modulating dopamine receptor activity did not

significantly influence the overall amount of reward obtained or the response time, it is possible

that apomorphine and flupenthixol exerted influence on how animals adapted their choices to

reward outcome. To understand how increasing and decreasing dopamine activity influenced

animals' reward-driven choice behaviors, we examined the probability of win-stay (repeating a

choice when it was rewarded on the previous trial) and the probability of lose-shift (switching to

the other choice when not rewarded). We found that when increasing dopamine receptor activity

with apomorphine administration, both males and females had increased win-stay behavior

(**Figure 1F,** GLMM, main effect of drug, p=0.011, $\beta1$= -0.02) and decreased lose-shift behavior

(**Figure 1G,** GLMM, main effect of drug, p=0.016, $\beta1$= 0.034). This result suggested that

apomorphine increased the overall "stickiness" or inflexibility of choice and more stay behaviors

regardless of the outcome.

There was no significant change in the probability of win-stay when the animals were

administered flupenthixol (**Figure 1H,** GLMM, main effect of drug, p=0.27, β1= -0.001). It's

worth noting that there was main effect of sex in the probability of win-stay, with females having

a higher probability of win-stay, in both DA agonist and antagonist manipulations (GLMM, main

effect of sex, apomorphine: p=0.065, β2 = -0.054; flupenthixol: p=0.01, β2 =-0.05). The results

also revealed an opposing effect from dopamine agonist that antagonizing dopamine receptor

activity with flupenthixol administration increased the probability of lose-shift (**Figure 1I,**

GLMM, main effect of drug, p=0.04, β1 = -0.014).

To control for the total amount of switching, we calculated outcome sensitivity

(**Equation 2,** see Methods) to understand how much of the switching behavior was outcome-

sensitive. The result suggests that there were no significant changes in outcome sensitivity when

animals were administered apomorphine (GLMM, main effect of drug, p=0.22, β1 = -0.08).

However, administration of flupenthixol significantly increased outcome sensitivity across both

sexes (**Figure 1-1A,** GLMM, main effect of drug, p=0.006, β1= -0.06) and there was more

increase in outcome sensitivity in males (GLMM, interaction term, p=0.027, β3= -0.10).

Modulating beta-noradrenergic receptor activity also influenced win-stay and lose-shift

behaviors. Increasing beta-noradrenergic activity with isoproterenol administration decreased the

probability of win-stay compared to vehicle control (**Figure 2F,** GLMM, main effect of drug,

0.009, β1=0.007). Isoproterenol administration also decreased the probability of lose-shift in

both sexes (**Figure 2G,** GLMM, main effect of drug, p=0.0007, β1=0.083), primarily driven by

significant decrease in lose-shift in females under isoproterenol (**Figure 2G inset,** GLMM,

interaction term, p=0.004, β3=-0.077). Outcome sensitivity analysis revealed a significant

decrease in outcome sensitivity when administer isoproterenol compared to vehicle control

(**Figure 1-1B,** GLMM, main effect of drug, p=0.021, β1 = 0.134). In contrast, decreasing beta-

noradrenergic activity with propranolol administration increased the probability of win-stay

compared to vehicle control (**Figure 2H,** GLMM, main effect of drug, p<0.0001, β1 = -0.023).

There was also a significant interaction between sex and drug that males showed a greater

increase in win-stay on propranolol compared to females (**Figure 2H inset,** GLMM, interaction

term, 0.02, β3=-0.036). Interestingly, propranolol administration also decreased the probability

of lose-shift but only in males (**Figure 2I,** GLMM, main effect of drug, 0.45, β1 = -0.01,

interaction term, p= 0.002, β3 =0.08). Outcome sensitivity analysis revealed a significant

increase in outcome sensitivity with propranolol administration (**Figure 1-1C,** GLMM, main

effect of drug, 0.003, β1 = -0.186). Together, these results suggested an opposing role of

dopamine and norepinephrine in modulating reward-driven decision making. Increasing

dopamine or decreasing norepinephrine activity resulted in increased win-stay and decreased

lose-shift but this effect is more sex-different in the noradrenergic manipulations.

A Hidden Markov model (HMM) identifies distinct patterns of exploration and

exploitation associated with modulation of dopamine and norepinephrine activity.

The changes in the reward-driven behaviors such as win-stay and lose-shift could be a

manifestation of strategic changes in how animals explored the changing environment. To

understand how modulation of dopamine and norepinephrine affect the balance between

exploration and exploitation, we used a Hidden Markov model (HMM) that modeled exploration

and exploitation as two latent goal states to infer which choices were exploratory or exploitative

(Figure 3A). The HMM model has previously been used in rodents, primates and humans (Ebitz

et al., 2018, 2020; Chen et al., 2021b) to infer explore-exploit state from choices and has shown

robust correlation with neural activity as well as other behavioral metrics, including response

time, value function, and reinforcement learning. We have previously established fast-switching

(putatively explore) and slow-switching (putatively exploit) regimes in mice choice behavior

(Chen et al., 2021b). This HMM method produces explore/exploit labels that better track neural

activity and choice behavior across species than reinforcement learning (RL) models (Ebitz et

al., 2018; Chen et al., 2021b). Figure 3A showed an example of HMM labeling of choices of an

animal in the restless bandit task. The shaded area indicates exploratory trials inferred by the

HMM. The animal displayed mixtures of exploratory bouts where choices were distributed

across two choices and exploitative bouts where one choice was repeated.

The HMM allows us to make statistical inferences about the probability that each choice

was due to exploration or exploitation by modeling explore/exploit as the distinct latent goal

states underlying choices. To evaluate the face validity of the HMM labels, we next examined

whether HMM-labeled exploration matched the normative definition of exploration. First, by

definition, exploration is a pattern of non-reward-maximizing choices whose purpose is to gain

information. A non-reward maximizing goal would produce choices that are orthogonal to

reward value. Therefore, we examined the probability of choosing a choice with regards to its

relative value. In all control and dopamine treatment conditions, HMM-labeled exploratory

choices were non-reward-maximizing: they were orthogonal to reward value, whereas exploit

choices were correlated with relative reward value (**Figure 3B, E)**. The probability of choosing a

target choice across bins of reward value was not different from chance ($t(43) = 0.39$, $p = 0.70$),

which means that choice behavior was not driven by reward value during exploration. During

exploration, animals chose both high-value choices and low-value choices at around chance level

(mean = 49.8% ± 3.4% STD across all value bins). During exploitation, animals were much

more likely to choose a high value choice over a low value choice (mean probability of choosing

high value choice = 75.8% ± 10.0% STD, mean probability of choosing low value choice =

23.3% ± 10.1% STD). Second, prior research has shown that exploratory choices are more

computationally expensive thus take longer than exploitative choices (Kool et al., 2010;

Filipowicz et al., 2019; Chen et al., 2021b). We calculated the response time across HMM-

inferred states and demonstrated that across all treatment conditions, HMM-inferred exploratory

choices had longer response time than exploit choices (**Figure 3C, D, F, G,** paired t test, vehicle

control (apomorphine): $t(31) = 4.48$, $p<0.0001$; apomorphine: $t(31)= 4.42$, $p<0.0001$; vehicle

control (flupenthixol): $t(31)=2.27$, $p=0.03$; flupenthixol: $t(31)=4.49$, $p<0.0001$). Together, these

results suggested that HMM produced meaningful and robust labeling of explore/exploit choices

that match the normative definition of exploration and were able to explain variances in value

selection and response time. Because this approach to infer exploration is agnostic to the

generative models and depends only on the temporal statistics of choices (Ebitz et al., 2018,

2020; Chen et al., 2021b), it is particularly ideal for circumstances like this one, where we

suspect that the generative computations may differ across treatment groups.

Modulating dopamine receptor activity bidirectionally affected the level of exploration

To examine how much animals were exploring, we calculated the average number of

exploratory choices inferred from HMM. The results revealed a bidirectional modulatory effect

of dopamine receptor activity on the level of exploration. When animals were administered

apomorphine, we found that animals on average had fewer exploratory trials than when they

were on vehicle control, with 55.5% ± 12.6% STD of trials labeled as exploratory on

apomorphine and 72.3% ± 13.6% STD of trials labeled as exploratory on vehicle (**Figure 3H, J,**

GLMM, main effect of drug, p<0.0001, β1 = 0.174). This is also consistent with our findings in

the win-stay lose-shift analysis that apomorphine increased the stickiness or repetitiveness of

choice behaviors. In contrast, when decreasing dopamine receptor activity with flupenthixol

administration, animals explore more compared to vehicle control, with 58.2% ± 12.0% STD of

trials being exploratory on flupenthixol and 52.1% ± 12.3% STD of trials being exploratory on

vehicle control (**Figure 3H, J**, GLMM, main effect of drug, 0.002, β1 = -0.062). This result

suggested that manipulating dopamine receptor activity bidirectionally affected the level of

exploration across both sexes (**Figure 3I, Figure 3-1)** - increasing dopamine receptor activity

decreased exploratory choices and decreasing dopamine receptor activity increased exploratory

choices.

The effect of beta-noradrenergic receptor activity on exploration was sex-modulated

First, we conducted an HMM model validation by examining the value selection and

response time during HMM-labeled explore and exploit state. Same as our findings in the

dopamine treatment conditions, we found that in all norepinephrine conditions, HMM-inferred

exploratory choices were also orthogonal to reward and exploit choices proportional to relative

reward value (**Figure 4A, D)**. The probability of choosing a target choice across bins of reward

value was not different from chance (t(43) = 0.85, p = 0.398). During exploration, animals chose

both high-value choices and low-value choices at around chance level (mean = 49.8% ± 1.8%

STD across all value bins). During exploitation, animals were much more likely to choose a high

value choice over a low value choice (mean probability of choosing high value choice = 74.5% ±

8.6% STD, mean probability of choosing low value choice = 24.2% ± 9.3% STD). Response

time during HMM-labeled exploratory choices were also longer than exploitative choices across

all control conditions and propranolol condition (**Figure 4B, C, E, F,** paired t test, vehicle

control (isoproterenol): $t(31)=3.01$, $p=0.005$; isoproterenol: $t(31)=1.48$, $p=0.15$; vehicle control

(propranolol): $t(31)=1.93$, $p=0.06$; propranolol: $t(30)=5.70$, $p<0.0001$). With confidence in the

approach to infer exploration, we next examined the effect of propranolol and isoproterenol on

the level of exploration.

Modulating beta-noradrenergic receptor activity also influenced the probability of

exploration and the modulatory effect on exploration was associated with sex. Decreasing beta-

noradrenergic receptor activity with propranolol significantly decreased the level of exploration

(**Figure 4G-I**) but only in males (**Figure 4J, Figure 3-1**, GLMM, interaction term, 0.0001, $\beta1 =$

0.138). Increasing beta-noradrenergic receptor activity with isoproterenol administration

significantly decreased exploration in females (**Figure 3K, Figure 3-1**, interaction term, $p=$

0.002, $\beta3=-0.103$). Together these results suggested a sex-linked neuromodulatory effect of beta-

noradrenergic receptor on exploration - increasing beta-noradrenergic activity decreased

exploration in males and decreasing beta-noradrenergic activity decreased exploration in

females. One possibility is that this sex-differentiated modulatory effect reflected sex-dependent

ceiling/floor effects of noradrenergic signaling.

Reinforcement learning (RL) models revealed changes in distinct parameters under

modulation of dopamine and epinephrine

The results of the HMM suggested that both dopamine and norepinephrine modulation

influenced the level of exploration. However, it is unclear whether changes in exploration were

due to changes in similar or distinct latent cognitive parameters. Does dopamine and

norepinephrine modulate exploration via distinct or overlapping mechanisms? To address this

question, we fitted a series of reinforcement learning (RL) models to understand the effect of

pharmacological manipulation on the latent cognitive parameters that could influence exploration

and exploitation (Ishii et al., 2002; Daw et al., 2006; Pearson et al., 2009; Jepma and

Nieuwenhuis, 2011).

To make inferences about changes in the RL model parameters, we first identified the

best-fitting RL model for the animals' behaviors. The majority of RL model parameters can be

categorized in three ways: value-dependent learning terms, value-independent bias terms, and

decision noise/randomness terms (Katahira, 2018). Previous studies demonstrated the effect of

various RL parameters on value-based decision making, including value-dependent terms such as

learning rate and asymmetrical learning rate (Frank et al., 2007; Gershman, 2016; Cinotti et al.,

2019; Chen et al., 2021b), value-independent terms such as choice bias (Katahira, 2018; Wilson

and Collins, 2019; Chen et al., 2021b), and noise terms such as inverse temperature and lapse

rate (Economides et al., 2015; Cinotti et al., 2019; Wilson and Collins, 2019). Here, we

compared six reinforcement learning models that incorporated one or more of the above

parameters (learning rate, bias, noise) and had different assumptions about the latent cognitive

processes that animals might employ during exploration (**Figure 5A, Equations 3-7**). These

models included: (1) a "random" model with some overall bias for one choice over the other, (2)

a "noisy win-stay lose-shift" model (WSLS) that assumes a win-stay lose-shift policy with some

level of randomness, (3) a two-parameter "RL" model with a consistent learning rate and some

inverse temperature that captures decision noise, (4) a three-parameter "RLCK" model that

captures both value-based and value-independent decision with two parameters for learning rate

and choice bias, and an overall decision noise parameter, (5) a three-parameter "RLγ" model that

captures asymmetrical learning with a learning rate parameter and a scaling parameter for

negative outcome learning and a decision noise parameter, and (6) a four-parameter "RLCKγ"

asymmetrical learning and bias model that includes a choice bias term on top of the "RLγ" (see

methods). We fitted these models to each individual animal and compared the likelihood of each

of the six models under four vehicle control conditions and four drug conditions.

In dopamine modulation conditions, the RLCK model (learning + choice kernel,) and the

RLCKγ (asymmetrical learning and bias) model best fit the behaviors (**Figure 5B,** average AIC

across DA conditions: model 1 random: 73838.03; model 2 WSLS: 55843.26; model 3 RL:

58133.45; model 4 RLCK: 54652.02; model 5 RLγ: 57545.47; model 6 RLCKγ: 54612.56).

Since the RLCKγ model did not significantly improve the model fitting, we decided to use the

more parsimonious RLCK model that has three parameters (learning rate, choice bias, and

decision noise). To examine how well the best fit RLCK model was at actually predicting

animals' choices, we measured the model agreement for each model, which was calculated as the

probability of choice correctly predicted by the optimized model parameters for each model

(**Figure 5C**). In dopamine agonist condition, the RLCK model predicted over 66% of animals'

actual choices (vehicle: 66.1% ± 5.3% STD, APO: 67.6% ± 5.0%). In the dopamine antagonist

condition, the RLCK model also predicted over 66% of animals' actual choices (vehicle: 66.4%

± 6.4%, FLU: 67.3% ± 5.4%). Figure 5D and 5E showed the model simulation, actual choice,

and reward probability of the same animal. This also visually demonstrated that the RLCK

model was able to accurately predict animal's changing choice behaviors. Additionally, this also

demonstrated that animal choice behaviors were following the reward contingency.

Then we examined how the reinforcement learning model parameters changed with

dopamine modulation. Previous study has shown that both learning rate and inverse temperature

parameters could induce changes in the overall level of exploration (Chen et al., 2021b).

Therefore, we asked whether the bidirectional changes in exploration when up- or down-

regulating dopamine receptor activity were due to changes in learning rate or decision noise. The

results revealed a bidirectional effect of dopamine modulation on the inverse temperature

parameter (**Figure 5F-H**). Because RL model data can be non-normally distributed, the Shapiro-

Wilk test of normality was conducted to determine whether the RL parameters were normally

distributed. The result suggested that the parameters were not always normally distributed

(decision noise: vehicle: $p= 0.85$, APO: $p<0.0001$; vehicle: $p<0.0001$, FLU: $p=0.09$), therefore

we will report both parametric and non-parametric statistics results for RL parameters.

Increasing dopamine receptor activity with apomorphine administration increased inverse

temperature and decreased random noise (**Figure 5F**, Wilcoxon matched-pairs test, $p= 0.0165$;

paired t-test, $p= 0.053$). Decreasing dopamine receptor activity with flupenthixol administration

decreased inverse temperature and increased random noise (**Figure 5G**, Wilcoxon matched-pairs

test, $p= 0.0253$; paired t-test, $p= 0.053$). Neither dopamine agonist (apomorphine) nor dopamine

antagonist (flupenthixol) significantly influenced learning rate or choice kernel compared to

vehicle controls.

In norepinephrine modulation conditions, the RLCK model (learning + choice kernel,)

and the RLCKγ (asymmetrical learning and bias) model also best fit the behavioral data (**Figure**

5I, average AIC across NE conditions: model 1 random: 66898.83; model 2 WSLS: 56061.3;

model 3 RL: 53194.59; model 4 RLCK: 50035.02; model 5 RLγ: 52664.48; model 6

RLCKγ:50027.82). However, since the RLCKγ model did not significantly improve the model

fitting, we decided to use the more parsimonious RLCK model for the NE condition as well. The

model agreement was calculated for all NE modulations. In the noradrenergic agonist condition,

the RLCK model predicted over 65% of animals' actual choices (**Figure 5J,** vehicle: 65.7% ±

5.9% STD, ISP: 66.4% ± 5.7%). In the noradrenergic antagonist condition, the RLCK model also

predicted over 65% of animals' actual choices (vehicle: 65.5% ± 3.9% STD, PRO: 69.8% ±

4.5%). Figure 5K and 5L showed the model simulation, actual choice, and reward probability of

the same animal as Figure 5D and 5E.

We found that modulating beta-noradrenergic receptor activity also resulted in changes in

the reinforcement learning model parameter. Up-regulating norepinephrine activity with

isoproterenol significantly decreased the learning rate (**Figure 5M**, Wilcoxon matched-pairs test,

p= 0.0001; paired t-test, p= 0.0008). This result is consistent with the decreased win-stay and

decreased lose-shift behaviors when administered isoproterenol because the lower learning rate

under isoproterenol resulted in reduced outcome sensitivity (**Supplementary figure 1C**), which

means learning less from either a rewarded or a non-rewarded outcome. There were no

significant changes in other parameters when administered isoproterenol compared to vehicle

control. We did not find any significant changes in any reinforcement learning parameters when

administered propranolol compared to vehicle control. Together, these results suggested distinct

roles of dopaminergic drugs in modulating exploration via decision noise and isoproterenol in

modulating exploration via outcome sensitivity.

Discussion

In this study, we pharmacologically up- and down-regulated dopamine and beta-

noradrenergic receptor activity and examined the modulatory effects of the manipulations on

exploration in an explore/exploit task. We used a combination of computational modeling

approaches to characterize changes in the level of exploration and latent cognitive variables that

could contribute to exploration. Modulating dopamine receptor activity revealed a bi-directional

modulatory effect on exploration - increasing dopamine receptor activity decreased the level of

exploration and decreasing dopamine receptor activity increased the level of exploration.

Modulation of beta-noradrenergic receptor activity also influenced how much animals explored -

beta-noradrenergic receptor antagonist (propranolol) decreased the level of exploration.

However, when examining the modulatory effect of beta-noradrenergic receptor activity on

exploration across sexes, we found that modulatory effect was sex-different: isoproterenol

selectively decreased exploration in females whereas propranolol decreased exploration in males.

Previously, the role of dopamine and norepinephrine in mediating exploration were often

examined in isolation with different task designs. This study allowed the direct comparison of

the modulatory effect of dopamine and norepinephrine on the transition between exploration and

exploitation in the same task across the same animals.

Dopamine's function as a neuromodulator has been described in two key ways. First, the

phasic activation of midbrain dopamine neurons is thought to be a key contributor to

reinforcement learning (Barto, 1995; Montague et al., 1995, 1996; Schultz et al., 1997; Schultz,

1998). Numerous studies across species have shown that dopamine neurons encoded the reward

prediction error (RPE) (Schultz, 1998; Niv, 2009). Ventral tegmental dopamine neuronal activity

and release in ventral striatum is proposed to reflect this error, which is used to update action

values (Schultz, 1998, 2013; Niv, 2009). Consistent with this view, some previous

pharmacological investigations in human decision making have observed changes in model-

derived learning rates that directly correlate with dopamine synthesis agonism via L-DOPA

(Frank et al., 2004; Rutledge et al., 2009) and dopamine receptor D2/D3 antagonism via

amisulpride (Cremer et al., 2022). However, this framework does not necessarily account for a

large body of literature describing a second way to view dopamine's neuromodulatory function,

as tonic levels influencing motivation, vigor, and cognitive flexibility (Floresco, 2013; Beeler et

al., 2014), especially when pharmacologically targeted in the frontal cortex. Because

pharmacological agents are by definition modulating receptor activity over long periods of time,

this role for tonic dopamine is poised to be computationally described. Indeed, when

pharmacologically or genetically modulating tonic dopamine function, many studies have noted

profound changes in the decision "noise" (inverse temperature) or perseverative errors -

consistent with a role for dopamine modulation in cognitive flexibility in earlier work (Beeler et

al., 2010; Humphries et al., 2012; Eisenegger et al., 2014; Lee et al., 2015; Cinotti et al., 2019;

Ebitz et al., 2019). Two prior examples combining pharmacological manipulation and

computational modeling in animal tasks explicitly testing the explore-exploit tradeoff

particularly highlight this. Cinotti et al. found that systemic dopamine blockade with flupenxithol

increased the amount of random exploratory choices without affecting learning rate in a rat three-

arm bandit task (Cinotti et al., 2019), particularly in situations where uncertainty was

manipulated. Similarly, Ebitz et al. reported that systemic cocaine, which blocks dopamine

reuptake, reduced flexibility and regulated tonic exploration in rhesus macaques (Ebitz et al.,

2019). Likewise, we find that systemic activation of dopamine receptors significantly decreased

exploration as defined by our Hidden Markov model and decision noise as described by our

reinforcement learning model, while antagonism of these receptors increased these

computational parameters measuring flexible behavior. Our data thus provide additional support

for the idea that dopamine modulation across the brain provides more than a learning signal, but

also a key signal tuning the precision or flexibility of behavior.

As noted in our introduction, the roles of dopamine and norepinephrine are sometimes

conflated in the computational neuroscience literature. This is in part because modulation of

these systems can be correlated, and noninvasive measures of catecholamine function, for

example blink rate and pupil diameter, could be ascribed to the endogenous activity of either

neuromodulator (although pharmacological action may target one or the other) (Jepma and

Nieuwenhuis, 2011; Ebitz and Moore, 2018; Sescousse et al., 2018; Findling et al., 2019; Cremer

et al., 2022). This is further complicated by the fact that some studies examining the relationship

between dopamine and exploration have used pharmacological manipulations which

simultaneously influence both tonic dopamine and norepinephrine levels, such as

psychostimulants or L-DOPA (Pessiglione et al., 2006; Rutledge et al., 2009; Orsini et al., 2016;

Ebitz et al., 2019; Chakroun et al., 2020). This makes it challenging to dissociate the modulatory

roles of dopamine and norepinephrine in exploration i.e. how dopamine and norepinephrine

distinctively contribute to changes in reinforcement learning model parameters, and addressing

this challenge was a major goal of our study.

Surprisingly, we found that when modulating beta-noradrenergic receptor activity

changed the level of exploration, it did so via changes in outcome sensitivity. Previous studies

have proposed the locus coeruleus-norepinephrine (LC-NE) system to regulate exploration by

changing decision noise. Kane et al. chemogenetically stimulated locus coeruleus activity in rats

and found that increased LC tonic activity increased in increased exploration, best explained by

an increase in decision noise (Kane et al., 2017; Wilson et al., 2021). Other studies have echoed

the findings that norepinephrine levels predicted the level of random exploration, or decision

noise, in the system (Jepma and Nieuwenhuis, 2011; Ebitz and Moore, 2018; Joshi and Gold,

2020). However, in this study we did not find any changes in the noise parameter (inverse

temperature, or how closely you adhere to your value rule) associated with up- and down-

regulation of beta-noradrenergic activity. Instead, the results revealed some learning rate changes

when increasing beta-noradrenergic receptor activity. Model-free analysis on outcome sensitivity

suggested that increasing or decreasing beta-noradrenergic receptor activity bi-directionally

changed outcome sensitivity. Propranolol increased outcome sensitivity and decreased

exploration, which means the animals were more likely to switch choice after loss and more

likely to repeat rewarded choices. In contrast isoproterenol decreased outcome sensitivity, as

seen in reduced win-stay and lose-shift behavior. This is consistent with the decreased learning

rate when administered isoproterenol because a decrease in learning rate is associated with

decreased value learning from each outcome. Where might this strange association between

noradrenergic modulation, exploration, and outcome sensitivity emerge from? We have

previously shown that exploration as defined by our Hidden Markov model can be driven by

either changes in learning rate or changes in inverse temperature (Chen et al., 2021b), supporting

the idea that the exploration we see here could be driven by long-term changes in the influence

of outcomes. Work focused on beta noradrenergic signaling in the context of aversive learning

has highlighted a role for noradrenergic signaling in extinction (Iordanova et al., 2021), over a

phasic timeframe of minutes to hours. This suggests the intriguing possibility that the effect of

noradrenergic blockade here may be to prevent extinction of learned values. In short, although

the most common associations in the computational neuroscience literature are between

dopamine and reward learning versus norepinephrine and decision noise, our current data

suggests a more nuanced role for each of these modulators.

Adding an additional layer of nuance to these findings, we find that sex differences are a

major axis of variability in the effect of beta-noradrenergic receptor modulation on the explore-

exploit tradeoff. Increasing NE receptor activity with the beta agonist isoproterenol decreased

exploration only in males, while decreasing NE receptor activity with the beta antagonist

propranolol decreased exploration especially in females. One possible explanation is that the

relationship between exploration and NE receptor activity was nonlinear but instead, followed an

inverted U-shape. Such inverted U-shape correlation has been suggested between cognitive

performance and NE level (Aston-Jones and Cohen, 2005; Redish et al., 2007; Mäki-Marttunen

et al., 2020). Since females have been reported to have higher tonic NE levels due to both

function and structural differences (Pinos et al., 2001; Bangasser et al., 2016), it is possible that

the baseline difference in tonic NE level across sexes could posit individuals on different parts of

the inverted U-curve, where increasing or decreasing NE could affect exploration in one sex

without impacting the other.

Taken together, this study demonstrates the distinct role of dopamine and norepinephrine

in tuning the exploration-exploitation tradeoff when learning about an uncertain environment.

Dopamine modulated exploration via decision noise and norepinephrine modulated exploration

via outcome sensitivity. The systemic manipulation of dopamine and norepinephrine receptor

activity did not allow for more precise modulation of phasic changes in catecholamine levels.

However, the computational tool that we presented here can complement higher resolution

neural recording techniques to examine for phasic changes in dopamine and norepinephrine

activity on a trial-by-trial basis. Finally, this study also highlighted that the modulatory effects of

catecholamine on exploration-exploitation tradeoff are strongly influenced by sex and sex-linked

mechanisms. Therefore, future research should take into account sex-correlated individual

variability when examining the neural correlates of exploration.

References

Addicott MA, Pearson JM, Schechter JC, Sapyta JJ, Weiss MD, Kollins SH (2020) Attention-

deficit/hyperactivity disorder and the explore/exploit trade-off. Neuropsychopharmacology

Available at: http://dx.doi.org/10.1038/s41386-020-00881-8.

Amemiya S, Kubota N, Umeyama N, Nishijima T, Kita I (2016) Noradrenergic signaling in the

medial prefrontal cortex and amygdala differentially regulates vicarious trial-and-error in a

spatial decision-making task. Behav Brain Res 297:104–111 Available at:

http://dx.doi.org/10.1016/j.bbr.2015.09.002.

Aston-Jones G, Cohen JD (2005) An integrative theory of locus coeruleus-norepinephrine

function: adaptive gain and optimal performance. Annu Rev Neurosci 28:403–450

Available at: http://dx.doi.org/10.1146/annurev.neuro.28.061604.135709.

Bangasser DA, Wiersielis KR, Khantsis S (2016) Sex differences in the locus coeruleus-

norepinephrine system and its regulation by stress. Brain Res 1641:177–188 Available at:

http://dx.doi.org/10.1016/j.brainres.2015.11.021.

Barto AG (1995) Adaptive critics and the basal ganglia. Models of information processing in the

basal ganglia 382:215–232 Available at: https://psycnet.apa.org/fulltext/1995-97527-

007.pdf.

Beeler JA, Cools R, Luciana M, Ostlund SB, Petzinger G (2014) A kinder, gentler dopamine…

highlighting dopamine's role in behavioral flexibility. Front Neurosci 8:4 Available at:

http://dx.doi.org/10.3389/fnins.2014.00004.

Beeler JA, Daw N, Frazier CRM, Zhuang X (2010) Tonic dopamine modulates exploitation of

reward learning. Front Behav Neurosci 4:170 Available at:

http://dx.doi.org/10.3389/fnbeh.2010.00170.

Bilmes J (1998) A Gentle Tutorial of the EM Algorithm and its Application to Parameter

Estimation for Gaussian Mixture and Hidden Markov Models. Available at:

http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.28.613 [Accessed April 23, 2021].

Bouret S, Sara SJ (2005) Network reset: a simplified overarching theory of locus coeruleus

noradrenaline function. Trends Neurosci 28:574–582 Available at:

http://dx.doi.org/10.1016/j.tins.2005.09.002.

Cabib S, Puglisi-Allegra S, Oliverio A (1984) Chronic stress enhances apomorphine-induced

stereotyped behavior in mice: involvement of endogenous opioids. Brain Res 298:138–140

Available at: http://dx.doi.org/10.1016/0006-8993(84)91156-9.

Chakroun K, Mathar D, Wiehler A, Ganzer F, Peters J (2020) Dopaminergic modulation of the

exploration/exploitation trade-off in human decision-making. Elife 9 Available at:

http://dx.doi.org/10.7554/eLife.51260.

Chen CS, Ebitz RB, Bindas SR, Redish AD, Hayden BY, Grissom NM (2021a) Divergent

Strategies for Learning in Males and Females. Curr Biol 31:39–50.e4 Available at:

http://dx.doi.org/10.1016/j.cub.2020.09.075.

Chen CS, Knep E, Han A, Ebitz RB, Grissom N (2021b) Sex differences in learning from

exploration. Elife 10 Available at: http://dx.doi.org/10.7554/eLife.69748.

Cinotti F, Fresno V, Aklil N, Coutureau E, Girard B, Marchand AR, Khamassi M (2019)

Dopamine blockade impairs the exploration-exploitation trade-off in rats. Sci Rep 9:6770

Available at: http://dx.doi.org/10.1038/s41598-019-43245-z.

Cohen JD, McClure SM, Yu AJ (2007) Should I stay or should I go? How the human brain

manages the trade-off between exploitation and exploration. Philosophical Transactions of

the Royal Society B: Biological Sciences 362:933–942 Available at:

http://dx.doi.org/10.1098/rstb.2007.2098.

Cremer A, Kalbe F, Müller JC, Wiedemann K, Schwabe L (2022) Disentangling the roles of

dopamine and noradrenaline in the exploration-exploitation tradeoff during human decision-

making. Neuropsychopharmacology Available at: http://dx.doi.org/10.1038/s41386-022-

01517-9.

Daw ND, O'Doherty JP, Dayan P, Seymour B, Dolan RJ (2006) Cortical substrates for

exploratory decisions in humans. Nature 441:876–879 Available at:

http://dx.doi.org/10.1038/nature04766.

Ebitz RB, Albarran E, Moore T (2018) Exploration Disrupts Choice-Predictive Signals and

Alters Dynamics in Prefrontal Cortex. Neuron 97:475 Available at:

http://dx.doi.org/10.1016/j.neuron.2018.01.011.

Ebitz RB, Moore T (2018) Both a Gauge and a Filter: Cognitive Modulations of Pupil Size.

Front Neurol 9:1190 Available at: http://dx.doi.org/10.3389/fneur.2018.01190.

Ebitz RB, Sleezer BJ, Jedema HP, Bradberry CW, Hayden BY (2019) Tonic exploration governs

both flexibility and lapses. PLoS Comput Biol 15:e1007475 Available at:

http://dx.doi.org/10.1371/journal.pcbi.1007475.

Ebitz RB, Tu JC, Hayden BY (2020) Rules warp feature encoding in decision-making circuits.

PLoS Biol 18:e3000951 Available at: http://dx.doi.org/10.1371/journal.pbio.3000951.

Economides M, Kurth-Nelson Z, Lübbert A, Guitart-Masip M, Dolan RJ (2015) Model-Based

Reasoning in Humans Becomes Automatic with Training. PLoS Comput Biol 11:e1004463

Available at: http://dx.doi.org/10.1371/journal.pcbi.1004463.

Eisenegger C, Naef M, Linssen A, Clark L, Gandamaneni PK, Müller U, Robbins TW (2014)

Role of dopamine D2 receptors in human reinforcement learning.

821   Neuropsychopharmacology 39:2366–2375 Available at:

822   http://dx.doi.org/10.1038/npp.2014.84.

823 Elliott R (2003) Executive functions and their disorders. Br Med Bull 65:49–59 Available at:

824   http://dx.doi.org/10.1093/bmb/65.1.49.

825 Fernandez-Tome MP, Sanchez-Blazquez P, del Rio J (1979) Impairment by apomorphine of one-

826   trial passive avoidance learning in mice: the opposing roles of the dopamine and

827   noradrenaline systems. Psychopharmacology 61:43–47 Available at:

828   http://dx.doi.org/10.1007/BF00426809.

829 Filipowicz ALS, Levine J, Piasini E, Tavoni G, Kable JW, Gold JI (2019) The complexity of

830   model-free and model-based learning strategies. Available at:

831   http://dx.doi.org/10.1101/2019.12.28.879965.

832 Findling C, Skvortsova V, Dromnelle R, Palminteri S, Wyart V (2019) Computational noise in

833   reward-guided learning drives behavioral variability in volatile environments. Nat Neurosci

834   22:2066–2077 Available at: http://dx.doi.org/10.1038/s41593-019-0518-9.

835 Floresco SB (2013) Prefrontal dopamine and behavioral flexibility: shifting from an "inverted-

836   U" toward a family of functions. Front Neurosci 7:62 Available at:

837   http://dx.doi.org/10.3389/fnins.2013.00062.

838 Frank MJ, Moustafa AA, Haughey HM, Curran T, Hutchison KE (2007) Genetic triple

839   dissociation reveals multiple roles for dopamine in reinforcement learning. Proc Natl Acad

840   Sci U S A 104:16311–16316 Available at: http://dx.doi.org/10.1073/pnas.0706111104.

841 Frank MJ, Seeberger LC, O'reilly RC (2004) By carrot or by stick: cognitive reinforcement

842   learning in parkinsonism. Science 306:1940–1943 Available at:

843   http://dx.doi.org/10.1126/science.1102941.

Gershman SJ (2016) Empirical priors for reinforcement learning models. J Math Psychol 71:1–6

Available at: https://www.sciencedirect.com/science/article/pii/S0022249616000080.

Gershman SJ (2019) Uncertainty and Exploration. Decision (Wash D C) 6:277–286 Available

at: http://dx.doi.org/10.1037/dec0000101.

Goldschmidt PL, Frances H, Simon P (1984) Stimulation of beta-adrenergic receptors and

spontaneous motor activity in mice. Pharmacol Biochem Behav 21:177–180 Available at:

http://dx.doi.org/10.1016/0091-3057(84)90210-7.

Grigoryan GA (2012) Serotonin and Impulsivity (animal experiments). Neurosci Behav Physiol

42:885–894 Available at: https://doi.org/10.1007/s11055-012-9654-3.

Grissom NM, Reyes TM (2019) Let's call the whole thing off: evaluating gender and sex

differences in executive function. Neuropsychopharmacology Available at:

https://www.nature.com/articles/s41386-018-0179-5.

Grossman CD, Bari BA, Cohen JY (2020) Serotonin neurons modulate learning rate through

uncertainty. bioRxiv:2020.10.24.353508 Available at:

https://www.biorxiv.org/content/10.1101/2020.10.24.353508v1 [Accessed September 29,

2021].

Humphries MD, Khamassi M, Gurney K (2012) Dopaminergic Control of the Exploration-

Exploitation Trade-Off via the Basal Ganglia. Front Neurosci 6:9 Available at:

http://dx.doi.org/10.3389/fnins.2012.00009.

Ichihara K, Nabeshima T, Kameyama T (1988) Opposite effects induced by low and high doses

of apomorphine on single-trial passive avoidance learning in mice. Pharmacol Biochem

Behav 30:107–113 Available at: http://dx.doi.org/10.1016/0091-3057(88)90431-5.

Iordanova MD, Yau JO-Y, McDannald MA, Corbit LH (2021) Neural substrates of appetitive

and aversive prediction error. Neurosci Biobehav Rev 123:337–351 Available at:

http://dx.doi.org/10.1016/j.neubiorev.2020.10.029.

Ishii S, Yoshida W, Yoshimoto J (2002) Control of exploitation-exploration meta-parameter in

reinforcement learning. Neural Netw 15:665–687 Available at:

http://dx.doi.org/10.1016/s0893-6080(02)00056-4.

Jepma M, Nieuwenhuis S (2011) Pupil Diameter Predicts Changes in the Exploration–

Exploitation Trade-off: Evidence for the Adaptive Gain Theory. J Cogn Neurosci 23:1587–

1596 Available at: https://doi.org/10.1162/jocn.2010.21548.

Jepma M, Schaaf JV, Visser I, Huizenga HM (2020) Uncertainty-driven regulation of learning

and exploration in adolescents: A computational account. PLoS Comput Biol 16:e1008276

Available at: http://dx.doi.org/10.1371/journal.pcbi.1008276.

Joshi S, Gold JI (2020) Pupil Size as a Window on Neural Substrates of Cognition. Trends Cogn

Sci 24:466–480 Available at: http://dx.doi.org/10.1016/j.tics.2020.03.005.

Kane GA, Vazey EM, Wilson RC, Shenhav A, Daw ND, Aston-Jones G, Cohen JD (2017)

Increased locus coeruleus tonic activity causes disengagement from a patch-foraging task.

Cogn Affect Behav Neurosci 17:1073–1083 Available at: http://dx.doi.org/10.3758/s13415-

017-0531-y.

Kaske EA, Chen CS, Meyer C, Yang F, Ebitz B, Grissom N, Kapoor A, Darrow DP, Herman AB

(2022) Prolonged physiological stress is associated with a lower rate of exploratory learning

that is compounded by depression. Biol Psychiatry Cogn Neurosci Neuroimaging Available

at: https://linkinghub.elsevier.com/retrieve/pii/S2451902222003391.

Katahira K (2018) The statistical structures of reinforcement learning with asymmetric value

updates. J Math Psychol 87:31–45 Available at:

https://www.sciencedirect.com/science/article/pii/S0022249617302407.

Kim H, Lee D, Shin Y-M, Chey J (2007) Impaired strategic decision making in schizophrenia.

Brain Res 1180:90–100 Available at: http://dx.doi.org/10.1016/j.brainres.2007.08.049.

Kobayashi K (2001) Role of catecholamine signaling in brain and nervous system functions: new

insights from mouse molecular genetic study. J Investig Dermatol Symp Proc 6:115–121

Available at: http://dx.doi.org/10.1046/j.0022-202x.2001.00011.x.

Kool W, McGuire JT, Rosen ZB, Botvinick MM (2010) Decision making and the avoidance of

cognitive demand. J Exp Psychol Gen 139:665–682 Available at:

http://dx.doi.org/10.1037/a0020198.

Lee E, Seo M, Dal Monte O, Averbeck BB (2015) Injection of a dopamine type 2 receptor

antagonist into the dorsal striatum disrupts choices driven by previous outcomes, but not

perceptual inference. J Neurosci 35:6298–6306 Available at:

http://dx.doi.org/10.1523/JNEUROSCI.4561-14.2015.

Mäki-Marttunen V, Andreassen OA, Espeseth T (2020) The role of norepinephrine in the

pathophysiology of schizophrenia. Neurosci Biobehav Rev 118:298–314 Available at:

http://dx.doi.org/10.1016/j.neubiorev.2020.07.038.

Malloy-Diniz LF, Miranda DM, Grassi-Oliveira R (2017) Editorial: Executive Functions in

Psychiatric Disorders. Front Psychol 8:1461 Available at:

http://dx.doi.org/10.3389/fpsyg.2017.01461.

Money KM, Stanwood GD (2013) Developmental origins of brain disorders: roles for dopamine.

Front Cell Neurosci 7:260 Available at: http://dx.doi.org/10.3389/fncel.2013.00260.

Montague PR, Dayan P, Person C, Sejnowski TJ (1995) Bee foraging in uncertain environments

using predictive hebbian learning. Nature 377:725–728 Available at:

http://dx.doi.org/10.1038/377725a0.

Montague PR, Dayan P, Sejnowski TJ (1996) A framework for mesencephalic dopamine

915   systems based on predictive Hebbian learning. J Neurosci 16:1936–1947 Available at:

916   https://www.ncbi.nlm.nih.gov/pubmed/8774460.

917   Morris LS, Baek K, Kundu P, Harrison NA, Frank MJ, Voon V (2016) Biases in the Explore-

918   Exploit Tradeoff in Addictions: The Role of Avoidance of Uncertainty.

919   Neuropsychopharmacology 41:940–948 Available at:

920   http://dx.doi.org/10.1038/npp.2015.208.

921   Mussey JL, Travers BG, Klinger LG, Klinger MR (2015) Decision-making skills in ASD:

922   performance on the Iowa Gambling Task. Autism Res 8:105–114 Available at:

923   http://dx.doi.org/10.1002/aur.1429.

924   Nakamura K, Kurasawa M, Tanaka Y (1998) Apomorphine-induced hypoattention in rats and

925   reversal of the choice performance impairment by aniracetam. Eur J Pharmacol 342:127–

926   138 Available at: http://dx.doi.org/10.1016/s0014-2999(97)01457-x.

927   Niv Y (2009) Reinforcement learning in the brain. J Math Psychol 53:139–154 Available at:

928   https://www.princeton.edu/~yael/Publications/Niv2009.pdf.

929   Niv Y, Joel D, Meilijson I, Ruppin E (2002) Evolution of reinforcement learning in uncertain

930   environments: A simple explanation for complex foraging behaviors. Adapt Behav 10:5–24

931   Available at: https://psycnet.apa.org/fulltext/2003-02884-002.pdf.

932   Orsini CA, Willis ML, Gilbert RJ, Bizon JL, Setlow B (2016) Sex differences in a rat model of

933   risky decision making. Behav Neurosci 130:50–61 Available at:

934   http://dx.doi.org/10.1037/bne0000111.

Pearson JM, Hayden BY, Raghavachari S, Platt ML (2009) Neurons in posterior cingulate cortex

signal exploratory decisions in a dynamic multioption choice task. Curr Biol 19:1532–1537

Available at: http://dx.doi.org/10.1016/j.cub.2009.07.048.

Pessiglione M, Seymour B, Flandin G, Dolan RJ, Frith CD (2006) Dopamine-dependent

prediction errors underpin reward-seeking behaviour in humans. Nature 442:1042–1045

Available at: http://dx.doi.org/10.1038/nature05051.

Pinos H, Collado P, Rodríguez-Zafra M, Rodríguez C, Segovia S, Guillamón A (2001) The

development of sex differences in the locus coeruleus of the rat. Brain Res Bull 56:73–78

Available at: http://dx.doi.org/10.1016/s0361-9230(01)00540-8.

Ranjbar-Slamloo Y, Fazlali Z (2019) Dopamine and Noradrenaline in the Brain; Overlapping or

Dissociate Functions? Front Mol Neurosci 12:334 Available at:

http://dx.doi.org/10.3389/fnmol.2019.00334.

Redish AD, Jensen S, Johnson A, Kurth-Nelson Z (2007) Reconciling reinforcement learning

models with behavioral extinction and renewal: implications for addiction, relapse, and

problem gambling. Psychol Rev 114:784–805 Available at: http://dx.doi.org/10.1037/0033-

295X.114.3.784.

Ressler KJ, Nemeroff CB (2001) Role of norepinephrine in the pathophysiology of

neuropsychiatric disorders. CNS Spectr 6:663–666, 670 Available at:

http://dx.doi.org/10.1017/s1092852900001358.

Rothenhoefer KM, Costa VD, Bartolo R, Vicario-Feliciano R, Murray EA, Averbeck BB (2017)

Effects of Ventral Striatum Lesions on Stimulus-Based versus Action-Based Reinforcement

Learning. J Neurosci 37:6902–6914 Available at:

http://dx.doi.org/10.1523/JNEUROSCI.0631-17.2017.

Rutledge RB, Lazzaro SC, Lau B, Myers CE, Gluck MA, Glimcher PW (2009) Dopaminergic

drugs modulate learning rates and perseveration in Parkinson's patients in a dynamic

foraging task. J Neurosci 29:15104–15114 Available at:

http://dx.doi.org/10.1523/JNEUROSCI.3524-09.2009.

Schultz W (1998) Predictive reward signal of dopamine neurons. J Neurophysiol 80:1–27

Available at: http://dx.doi.org/10.1152/jn.1998.80.1.1.

Schultz W (2013) Updating dopamine reward signals. Curr Opin Neurobiol 23:229–238

Available at: http://dx.doi.org/10.1016/j.conb.2012.11.012.

Schultz W, Dayan P, Montague PR (1997) A neural substrate of prediction and reward. Science

275:1593–1599 Available at: http://dx.doi.org/10.1126/science.275.5306.1593.

Seamans JK, Yang CR (2004) The principal features and mechanisms of dopamine modulation

in the prefrontal cortex. Prog Neurobiol 74:1–58 Available at:

http://dx.doi.org/10.1016/j.pneurobio.2004.05.006.

Sescousse G, Ligneul R, van Holst RJ, Janssen LK, de Boer F, Janssen M, Berry AS, Jagust WJ,

Cools R (2018) Spontaneous eye blink rate and dopamine synthesis capacity: preliminary

evidence for an absence of positive correlation. Eur J Neurosci 47:1081–1086 Available at:

http://dx.doi.org/10.1111/ejn.13895.

Sutton RS, Barto AG (1998) Reinformcent Learning: An Introduction.

Warren CM, Wilson RC, van der Wee NJ, Giltay EJ, van Noorden MS, Cohen JD, Nieuwenhuis

S (2017) The effect of atomoxetine on random and directed exploration in humans. PLoS

One 12:e0176034 Available at: http://dx.doi.org/10.1371/journal.pone.0176034.

Weed MR, Gold LH (1998) The effects of dopaminergic agents on reaction time in rhesus

monkeys. Psychopharmacology 137:33–42 Available at:

http://dx.doi.org/10.1007/s002130050590.

Williams OOF, Coppolino M, George SR, Perreault ML (2021) Sex Differences in Dopamine

Receptors and Relevance to Neuropsychiatric Disorders. Brain Sci 11 Available at:

http://dx.doi.org/10.3390/brainsci11091199.

Wilson RC, Bonawitz E, Costa VD, Ebitz RB (2021) Balancing exploration and exploitation

with information and randomization. Curr Opin Behav Sci 38:49–56 Available at:

http://dx.doi.org/10.1016/j.cobeha.2020.10.001.

Wilson R, Collins A (2019) Ten simple rules for the computational modeling of behavioral data.

Available at:

https://pdfs.semanticscholar.org/91b9/d3ab7532ea24ae70cd726355f25235b1fe8b.pdf.

Yu AJ, Dayan P (2005) Uncertainty, neuromodulation, and attention. Neuron 46:681–692

Available at: http://dx.doi.org/10.1016/j.neuron.2005.04.026.

Figures and figure legends

Figure 1. Modulation of dopamine receptor activity affected reward-directed behaviors.

Up-regulating dopamine activity increased stickiness in choice behaviors regardless of

outcomes.

A) Schematic of the mouse touchscreen chamber and the trial structure of the two-armed spatial

restless bandit task. **B)** An example of the dynamic reward contingency showing the changing

reward probabilities associated with each option over 300 trials. **C)** The dopaminergic drug administration schedule and drug dosage used to modulate dopamine receptor activity. A non-selective dopamine receptor agonist apomorphine (0.1mg/kg) and a non-selective dopamine receptor antagonist flupenthixol (0.03mg/kg) were systemically administered to respectively up- and down-regulate dopamine receptor activity. 0.9% saline was used as the vehicle control. Control and drug sessions were interleaved and repeated for six sessions each. **D)** (left) Average probability of obtaining reward compared to the chance level probability of reward for each animal under apomorphine (APO) and vehicle (dot). Error bars depict mean ± SEM across sessions for each animal. (right) probability of obtaining reward over chance level across vehicle and APO condition. **E)** (left) Average probability of obtaining reward compared to the chance level probability of reward for each animal under flupenthixol (FLU) and vehicle (dot). Error bars depict mean ± SEM across sessions for each animal. (right) probability of obtaining reward over chance level across vehicle and FLU condition. **F)** Average probability of win-stay on vehicle and APO condition. **G)** Average lose-shift on vehicle and APO condition. **H)** Average probability of win-stay on vehicle and FLU condition. **I)** Average lose-shift on vehicle and FLU condition. * indicates $p < 0.05$. Graphs depict mean ± SEM across animals unless specified otherwise.

Figure 2. The modulatory effects of beta-noradrenergic receptor activity on reward-directed behavior was also influenced by sex.

A) The noradrenergic drug administration schedule and drug dosage used to modulate

noradrenergic receptor activity. A beta-noradrenergic receptor agonist isoproterenol (0.3mg/kg)

and a beta-noradrenergic receptor antagonist propranolol (5mg/kg) were systemically

administered to respectively up- and down-regulate norepinephrine activity. 0.9% saline was

used as the vehicle control. Control and drug sessions were interleaved and repeated for six

sessions each. **B)** (left) Average probability of obtaining reward compared to the chance level

probability of reward for each animal under isoproterenol (ISP) and vehicle (dot). Error bars

depict mean ± SEM across sessions for each animal. (right) probability of obtaining reward over

chance level across vehicle and ISP. **C)** Average response time under ISP and vehicle. ISP

significantly increased response time. **D)** (left) Average probability of obtaining reward

compared to the chance level probability of reward for each animal under vehicle and

propranolol (PRO). Error bars depict mean ± SEM across sessions for each animal. (right)

probability of obtaining reward over chance level across PRO (light brown) and vehicle (gray).

E) Average response time under PRO and vehicle. PRO significantly decreased response time.

F) Average probability of win-stay under ISP and vehicle. **G)** Average probability of lose-shift

under ISP and vehicle. (Inset) probability of lose-shift across treatments across sexes. Decrease

in lose-shift under ISP was primarily driven by changes in females (interaction term). **H)**

Average probability of win-stay under PRO and vehicle. (Inset) probability of win-stay across

treatments across sexes. Increase in win-stay under PRO was primarily driven by changes in

males (interaction term). **I)** Average probability of lose-shift under PRO and vehicle. (inset)

probability of lose-shift across treatments across sexes. Decrease in lose-shift under PRO was

primarily driven by changes in males (interaction term). * indicates $p < 0.05$. Graphs depict mean

± SEM across animals unless specified otherwise.

1045

1046

Figure 3. Up- and down-regulation of dopamine activity had bidirectional effects on the

level of exploration.

A) (left) Structure of a Hidden Markov model (HMM) that modeled exploration and exploitation

as latent goal states underlying observed choices. This model incorporates two exploit states for

left and right choice respectively, and one explore state where choices were uniformly distributed

across options. (right) Reward probabilities, choices, and HMM labels for an example session of

300 trials for an example mouse. Shaded areas demonstrated HMM-labeled exploratory choices.

B) Probability of choice as a function of value differences between choices for exploratory and

exploitative states under vehicle (left) and flupenthixol (FLU) (right). **C, D)** Difference in

response time for explore and exploit choices under vehicle and flupenthixol (FLU). **E)**

Probability of choice as a function of value differences between choices for exploratory and

exploitative states under vehicle (left) and apomorphine (APO) (right). **F, G)** Difference in

response time for explore and exploit choices under vehicle and apomorphine (APO). **H)**

Distribution of the percentage of HMM-labeled exploratory choices under flupenthixol (FLU)/

vehicle (top) and apomorphine (APO)/vehicle (bottom). **I)** Probability of exploration for

dopamine antagonist (FLU) and agonist (APO), normalized by their vehicle control. Decreasing

dopamine activity increased exploration and increasing dopamine activity decreased exploration.

J) Probability of exploration by session with vehicle and drug session interleaved (flupenthixol:

top; apomorphine: bottom). Drug administration sessions are in colored shades. * indicates p <

0.05. Graphs depict mean ± SEM across animals.

1067

Figure 4. Downregulating norepinephrine activity influenced exploration but the effect was modulated by sex.

A) Probability of choice as a function of value differences between choices for exploratory and exploitative states under vehicle (left) and propranolol (PRO) (right). **B, C)** Difference in response time for explore and exploit choices under vehicle and propranolol (PRO). **D)** Probability of choice as a function of value differences between choices for exploratory and exploitative states under vehicle (left) and isoproterenol (ISP) (right). **E, F)** Difference in

response time for explore and exploit choices under vehicle and isoproterenol (ISP).

G) Distribution of the percentage of HMM-labeled exploratory choices under propranolol

(PRO)/ vehicle (top) and under isoproterenol (ISP)/vehicle (bottom). **H)** Probability of

exploration for beta-noradrenergic antagonist (PRO) and agonist (ISP), normalized by their

vehicle control. Propranolol decreased the level of exploration. **I)** Probability of exploration by

session with vehicle and drug session interleaved. Drug administration sessions are in colored

shades. Top: propranolol (PRO) condition; bottom: isoproterenol (ISP) condition. **J)** Probability

of exploration under PRO and vehicle across sexes. The effect of PRO on exploration was

primarily driven by a significant decrease in exploration under PRO in males. **K)** Probability of

exploration under ISP and vehicle across sexes. ISP significantly decreased exploration in

females but not males. * indicates $p < 0.05$. Graphs depict mean ± SEM across animals.

1087

1088

Figure 5. Modulation of dopamine and noradrenergic receptor activity led to changes in

different reinforcement learning (RL) model parameters.

A) A diagram of reinforcement learning (RL) parameters that capture learning rate (α),

asymmetric learning (γ), choice bias (α_c), inverse temperature/decision noise (β). The RL models

tested included one or more of the above parameters. **B)** Model likelihood for six RL models

using log likelihood for vehicle and apomorphine (APO) condition (top), and vehicle and

flupenthixol condition (bottom). The three-parameter RLCK model was the best fit and most

parsimonious model for behaviors. **C)** Model agreement for the best fit model (RLCK), which

measures the probability of the model predicting the actual choices of animals for vehicle and

apomorphine (APO) condition (top), and vehicle and flupenthixol condition (bottom). **D, E)**

Actual choices (purple) and simulated choices (green) from the best fit model (RLCK) for the

same animal under vehicle/apomorphine and vehicle/flupenthixol. Grey line represents the

reward probability of the left choice. **F)** Average inverse temperature (β) across apomorphine

and vehicle. (inset) inverse temperature across conditions after removing two outliers. **G)**

Average inverse temperature (β) across flupenthixol and vehicle. (inset) inverse temperature

across conditions after removing two outliers. **H)** Agonizing (apomorphine) and antagonizing

(flupenthixol) dopamine activity revealed bidirectional effect on inverse temperature, controled

by their own vehicle control. **I)** Model likelihood for six RL models using log likelihood for

vehicle and isoproterenol (ISP) condition (top) and vehicle and propranolol (PRO) condition

(bottom). **J)** Model agreement for the best fit model (RLCK), which measures the probability of

the model predicting the actual choices of animals for vehicle and isoproterenol (ISP) condition

(top) and vehicle and propranolol (PRO) condition (bottom). **K, L)** Actual choices (purple) and

simulated choices (green) from the best fit model (RLCK) for the same animal under

vehicle/propranolol and vehicle/isoproterenol . Grey line represents the reward probability of the

left choice. **M)** Average learning rate (α) across isoproterenol and vehicle. * indicates $p < 0.05$.

Graphs depict mean $\pm$ SEM across animals.

1115

1116   **Supplementary figure 1**

1117

1118

1119

Figure 1-1. Effect of dopamine and norepinephrine modulation on outcome sensitivity.

Related to Figure 1.

A) Average outcome sensitivity across flupenxithol (FLU) and vehicle. **B)** Average outcome

sensitivity across isoproterenol (ISP) and vehicle. **C)** Average outcome sensitivity across

propranolol (PRO) and vehicle. * indicates $p < 0.05$. Graphs depict mean ± SEM across animals.

Supplementary figure 2

1126

1127

1128 **Figure 3-1. Effect of dopamine and norepinephrine modulation on exploration in female**

1129 **and male mice. Related to Figure 3,4.**

1130