Genome-wide analysis of anterior-posterior mRNA localization in *Stentor coeruleus* reveals a role for the microtubule cytoskeleton

Ashley R. Albright[1,2], David Angeles-Albores[3], Wallace Marshall[1,2,4,*]

[1]Department of Biochemistry and Biophysics, University of California, San Francisco, San Francisco, CA, USA

[2]Center for Cellular Construction, University of California, San Francisco, San Francisco, CA, USA

[3]Altos Labs, San Francisco, CA, USA

[4]Twitter: @WallaceUCSF

*Correspondence: wallace.marshall@ucsf.edu

Abstract

Cells have complex and beautiful structures that are important for their function, but understanding the molecular mechanisms that produce these structures is a challenging problem due to the gap in size scales between molecular interactions and cellular structures. The giant ciliate *Stentor coeruleus* is a unicellular model organism whose large size, reproducible structure, and ability to heal wounds and regenerate has historically allowed the formation of structure in a single cell to be addressed using methods of experimental embryology. Such studies have shown that specific cellular structures, such as the oral apparatus, always form in specific regions of the cell, which raises the question: what is the source of positional information within this organism? By analogy with embryonic development, in which localized mRNA is often used to mark position, we asked whether position along the anterior-posterior axis of Stentor might be marked by specific regionalized mRNAs. By physically bisecting cells and conducting half-cell RNA sequencing, we were able to identify sets of messages enriched in either the anterior or posterior half. We repeated this analysis in cells in which a set of longitudinal microtubule bundles running down the whole length of the cell, known as KM-fibers, were disrupted by RNAi of β-tubulin. We found that many messages either lost their regionalized distribution or switched to an opposite distribution, such that anterior-enriched messages in control became posterior-enriched in the RNAi cells, or vice versa. This study indicates that mRNA can be regionalized within a single giant cell

and that microtubules may play a role, possibly by serving as tracks for the movement of the messages.

Introduction

Understanding how complex patterns arise within individual cells is a challenging biological problem due to the vast difference in spatial scales from molecules to whole cells [1,2]. The questions that arise in understanding cellular morphogenesis are essentially the same as those that arise in understanding the development of embryos - how does the system know which structures to make and how to position them relative to an overall body plan? A classic model organism that facilitates the study of developmental biology in a single cell is the giant ciliate *Stentor coeruleus* [3]. *Stentor* cells are 1 millimeter long and covered with blue stripes of pigment, which alternate with longitudinal rows of cilia consisting of basal bodies associated with microtubule bundles (KM-fibers) that run the length of the cell. The cone-shaped cell has an array of cilia known as the oral apparatus (OA) at its anterior end and a holdfast at the posterior. Orthogonal to the anterior-posterior (AP) axis defined by these structures, the pigment stripes show a graded distribution of width, defining a circumferential axis. Starting with the widest stripes, the stripe width gradually decreases as one moves counter-clockwise around the circumference, until eventually, the narrowest stripes meet the widest stripes, creating a discontinuity known as the contrast zone.

Part of the reason that *Stentor* was first used as a model system is its ability to regenerate after surgical manipulations. If any part of the *Stentor* cell is cut off, the missing piece regenerates in a matter of hours to yield a normal cell. If a cell is cut in half, each half regenerates its normal structure [4,5]. Part of the reason that *Stentor* can regenerate from these surgical operations is that it contains an elongated macronucleus that contains many copies of every gene. Another reason Stentor can regenerate is its ability to employ a range of mechanical wound-closure mechanisms to keep its cytoplasm from leaking out while it patches over a wound [6]. *Stentor's* huge size, easily visible patterning, and amazing powers of regeneration attracted many developmental

biologists during the turn of the last century, including Thomas Hunt Morgan [5]. For almost a century, up until the 1970s, microsurgeries were used to analyze morphogenetic processes in *Stentor*, resulting in a wealth of information about how cells respond to geometrical re-arrangements [7,8]. But *Stentor* was never developed as a molecular model system, and we do not currently know the molecular basis for how it can regenerate its structures. To exploit the unique opportunities presented by this organism, we sequenced the *Stentor* genome [9], developed methods for RNAi by feeding [10], and analyzed the transcriptional program of regeneration [11].

Perhaps the most fundamental question about pattern formation in any organism is what is the source of positional information. That such information exists in *Stentor* can be inferred from the fact that cellular structures always form in defined and reproducible positions [7]. For example, during regeneration or normal cell division, the new OA always forms halfway down the AP axis at the contrast zone between narrow and wide stripes [12]. It is generally true in ciliates that specific surface structures form at positions that can be defined with respect to a cylindrical coordinate system [13–15]. In this system, position is defined by two orthogonal axes - an AP axis that runs along the length of the cell, and a circumferential axis that runs around the equator, which in Stentor is manifest by the gradient of pigment stripe width. We have previously analyzed the distribution of the proteome relative to the AP axis, by cutting individual Stentor cells into pieces and analyzing the fragments by mass spectrometry [16]. This study found that approximately 25% of the proteome was polarized relative to this axis.

The role of localized mRNA as a source of positional information in embryonic development is well established, particularly in *Drosophila*. We anticipate that regionalized RNAs in different parts of the *Stentor* cell will be important sources of positional information, just like they are in embryos. Unlike the case of *Drosophila* development, where many important genes had already been identified in genetic experiments, we do not currently have a list of candidate positional information molecules to map. Inspired by prior work in cryo-slicing *Drosophila* embryos to obtain

positional information [17,18], we can use a similar unbiased approach to map the entire transcriptome to identify genes that might have a regionalized distribution.

Modern studies address sub-cellular localization of transcripts with single-molecule (smFISH [19]) or multiplexed error-robust fluorescence *in situ* hybridization (merFISH [20]). Many studies that survey message localization by FISH do so in cells that lack an obvious polarity axis, so collecting physical fragments of cells representing specific regions and mapping fragments back to any reference frame would not be possible. Consequently, in those cases, each message must be visualized sequentially by imaging, which is time-consuming and expensive. The giant size of *Stentor*, combined with the reproducible cell geometry and easily visualized body axis markers, makes it possible to use a sequencing-based method to map RNA localization relative to the body axis coordinates. The first question is whether genes are regionalized in *Stentor*. If it turns out that RNAs are localized to different parts of the *Stentor* cell, the next question is: how might this be achieved?

The surface of a *Stentor* is covered with parallel microtubule bundles with the plus end at the anterior and the minus end at the posterior. Studies in other systems from yeast to human neurons show that the cytoskeleton is involved in active RNA transport[21,22], therefore we hypothesize that transcripts in this giant cell may be polarized by trafficking specific messages along the cell's polarized cytoskeletal tracks. RNAi knockdown of β-tubulin, a major component of the microtubule cytoskeleton, causes *Stentor* to lose their characteristic trumpet shape (Fig 1A-C [10]) and lose their orderly microtubule bundles (Fig 1D-E [10]), but such cells are still alive and retain an OA at one end and a holdfast at the other. It is thus still possible to recognize the AP axis in such cells and ask whether disruption of the microtubule tracks is hypothesized to carry messages that have any effect on localization.

Figure 1 – β-tubulin knockdown results in loss of typical cell morphology and orderly microtubule bundles Brightfield images of (A) control and (B, C) β-tubulin knockdown cells showing disruption of typical cell morphology. Immunofluorescence of α-tubulin stained (green) in (D) control and (E) knockdown cells show that cells have broken (blue arrow) and discontinuous (white arrow) rows of microtubule bundles. Cells are oriented with the anterior at the top and posterior at the bottom. All images in this figure are adapted from Slabodnick et al 2014 [10].

Here, we investigate both the disruption of messages and the microtubule cytoskeleton in transcript polarity by conducting half-cell RNA sequencing in paired anterior and posterior halves of *Stentor coeruleus* following lateral bisection. Overall, we find evidence for a global disruption of transcript polarity as well as identify 117 candidate transcripts with an anti-correlated shift in AP skew.

Methods

Stentor culture, RNAi, and bisection

Stentor coeruleus were obtained from Carolina Biological Supply and maintained in small Pyrex dishes at room temperature in filter-sterilized pasteurized spring water (Carolina Biological Supply – Burlington, NC) subsequently referred to as Carolina Spring Water (CSW). Twice a week, a lentil-sized pellet of *Chlamydomonas reinhardtii* grown on TAP plates was resuspended in CSW and fed to *Stentor*. Before RNAi, *Stentor* were isolated from the main culture in 6-well tissue culture plates containing 2 mL CSW, starved for 3 days, and washed 3 times with CSW to eliminate residual *Chlamydomonas.*

β-tubulin RNAi was conducted by feeding as previously described[10] with minor changes. The sequenced target was amplified with the following oligos: forward - 5' ATGGTGACTTAAATCACTTGGTTAGTGC 3', reverse – 5' TTAAGCAGCTTCCTCTTCCTCATCG 3'. Transformed HT115 *E. coli* were grown to log phase and expression of dsRNA induced with 1 mM IPTG overnight at room temperature. Overnight 50 mL cultures were spun down, washed with CSW, and resuspended in 30 mL of CSW. Aliquots of 1 mL were pelleted, with as much supernatant as possible removed, and frozen at -80° for future use. For feeding, pellets were resuspended in 1 mL CSW. *Stentor* were fed 500 uL control (empty vector) or β-tubulin RNAi bacteria once a day for 7 days.

Individual *Stentor* were washed and transferred to a 2 uL droplet of 1.5% methylcellulose in CSW and bisected laterally using a glass needle. Halves were pipetted up and down in CSW to remove residual methylcellulose before processing.

Half-cell RNA-sequencing

Within 20 minutes of bisection, cells were lysed and prepped for sequencing with the NEBNext Single Cell/Low Input RNA Library Prep kit for Illumina (Cat. No. E6420) according to the manufacturer's protocol. Samples were pooled and submitted for sequencing. Sequencing on an Illumina NovaSeq6000 SP200 was performed at the UCSF CAT, supported by UCSF PBBR, RRP IMIA, and NIH 1S10OD028511-01 grants.

Data processing and analysis

Raw data will be available on Dryad following approval here:
https://doi.org/10.6078/D1WT6W

Data were processed and analyzed with custom scripts in both Python and R, which are available here: https://github.com/aralbright/2022_AADAWM

Adapter trimming was performed using flexbar [23] as instructed by NEB for the NEBNext Single Cell Library Prep Kit: https://github.com/nebiolabs/nebnext-single-cell-rna-seq. We then used kallisto [24] to generate a reference index and quantify transcript abundance. Using sleuth [25], we obtained normalized transcripts per million (TPM). TPM values were log(1+x) transformed. We performed a principal components analysis (PCA) to visualize the relatedness of the samples. To reduce the effects of outlying transcript abundance on significance, we regularized mean log-transformed TPM values (subsequently referred to as log1 Regularized Transcripts Per Million or log1RPM). Anterior-posterior skew is then defined as the difference over the sum of log1RPM.

$$Skew = \frac{A_{log1RPM} - P_{log1RPM}}{A_{log1RPM} + P_{log1RPM}}$$

Genes were selected for further analysis that passed a minimal cutoff of abundance, to ensure sufficiently robust statistical comparisons, and it was also required that genes had a coefficient of variation (variance divided by the mean) less than 3, to remove genes with such variable distributions that comparisons might be unreliable. For genes meeting these criteria, skews were compared for both control and β-tubulin RNAi samples.

We were particularly interested in genes with anti-correlated skews, the most interesting candidates for disrupted localization. Such genes were selected with feature selection

by PCA, in which the genes were treated as factors and the samples data points, to identify linear combinations of genes that give the highest variation among the samples. Feature selection was then followed by a t-test for significance.

Results

To ask whether any genes have localized distribution relative to the AP axis, we manually cut *Stentor* cells in half, using the OA as a marker to define the anterior end, and cutting approximately mid-way between the OA and the other end of the cell. For each dissected cell, the anterior half and posterior half cells were used to build libraries for RNA sequencing. We then define skew in terms of the difference in abundance of transcripts between the two half-cells in each sample. To quantify polarization along the AP axis, we log(1+x) transformed and regularized normalized transcripts per million. Then, we calculated the difference over the sum of AP values for each gene within each sample to obtain a skew value(see Methods for details). We removed genes with highly variable expression between samples or expression levels at the extreme end of high or low to ensure that these characteristics are not affecting our overall calculation of skew. We expect that skews will be variable to some extent even in normally polarized cells, given that the precision of bisection is subject to human error. With this analysis, each gene is assigned an average skew, taken over all samples, with positive values reflecting anterior enrichment and negative values reflecting posterior enrichment.

Among the 100 genes showing the largest absolute skew, 73 had a negative skew, suggesting that there may be a higher degree of regionalization in the posterior half of the cell. Interesting genes include SteCoe_9217, which encodes a cytoplasmic dynein light chain, and SteCoe_19711, which encodes an Ank_2 domain protein. Most of these top 100 skewed genes encode hypothetical proteins, including SteCoe_14908, which corresponds to a gene upregulated during regeneration of the OA. Understanding what functions these uncharacterized proteins may play in Stentor pattern formation is an interesting future goal, but here our primary interest is in the mechanisms by which the localization is achieved.

Given that the surface of Stentor is covered with parallel, oriented microtubule bundles, an obvious hypothesis is that motors might move transcripts to one end or the other of these bundles. To understand the role of the microtubule cytoskeleton in cellular patterning, we conducted half-cell RNA-sequencing of paired anteriors and posteriors in control and β-tubulin knockdown *Stentor coeruleus*. As expected, PCA shows a clear separation between control and β-tubulin samples on the first principal component (Fig 2A). Interestingly, anterior and posterior (AP) control samples are separated by the first component while AP β-tubulin samples are not. These differences at the whole-sample level suggest that the microtubule cytoskeleton is broadly important for proper RNA localization, such that anterior and posterior samples become more similar to each other in the absence of the microtubule tracks.

We observed that the mean skew of all genes is more variable across control than β-tubulin samples (Fig 2B). This might be expected if there was a general loss of polarized localization in the β-tubulin RNAi, such that variation in the exact position of cutting the cell would have less of an effect on the average skew.

Figure 2 – Polarity disruption suggested by PCA and average skew in control vs. β-tubulin knockdown samples (A) PCA of normalized transcripts per million (TPM) across entire samples. Control labeled in green and β-tubulin knockdown in magenta. Anteriors are represented by circles and posteriors by 'x.' (B) Average skew of all genes within each sample.

If microtubule tracks worked to separate a subset of transcripts along the AP axis, one would expect a loss of skew in the RNAi cells, which is certainly seen for many genes.

For other genes, in which the skews remain correlated, it suggests that some degree of regionalization is still possible without intact microtubule tracks. The track model can thus account for loss of correlation or retention of positive correlation. A more surprising outcome would be anti-correlation, such that a gene switches from anterior enriched in the control to posterior enriched in the knockdown, or vice versa. To look for such genes, we performed feature selection by PCA to detect genes with anti-correlated skews (see Methods). Although changes in skew that remain correlated may be significant (i.e. cases where an anterior-expressed gene still has an anterior skew, but perhaps more extreme), candidates with anti-correlated skews upon β-tubulin knockdown are the most interesting for further analysis as these account for the largest differences in skew between control and β-tubulin cells. Following feature selection, we found 117 candidates out of 500 tested with a significant anti-correlated skew (Fig 3A, Supplementary Table 1).

Figure 3B-F shows 5 representative examples of candidates found in our study. Of all 117 candidates, SteCoe_32460 has the most positive difference in skew (Figure 3B, Table 1), is differentially expressed specifically during tail regeneration [11], and is part of a family of calcium-binding EF-hand proteins thought to be important for contraction of the posterior tail in *Stentor* [26,27]. SteCoe_23664 has the second most positive difference in skew (Fig 3C, Table 1) and is a MORN repeat or possible Histone H3K4 specific methyltransferase protein. SteCoe_19711 shifts significantly more anterior upon tubulin knockdown (Fig 3D) and as mentioned previously, is one of the most posterior skewed genes in the control. SteCoe_19711 encodes an Ank_2 domain protein, suggesting it is part of the Ankyrin protein family. Members of the Ankyrin protein family tether proteins within and to the cell surface, for example the actin cytoskeleton, although the function of these proteins in ciliates is not known. Ankyrin proteins are also associated with ciliopathies [28] and in some cases have RNA-binding activity [29]. SteCoe_18056, which encodes a Toprim_2 domain protein, shifts more posterior upon tubulin knockdown (Fig 3E) and is also differentially expressed during regeneration [11]. SteCoe_13508 shifts significantly more anteriorly upon tubulin knockdown (Fig 3F) and is also differentially expressed during tail regeneration [11]. SteCoe_19508 is a possible

homolog of yeast PAB1, which interacts with poly (A) tails and functions in mRNA export [30]. Other potential candidates for further study include the most positive (Table 1) and most negative (Table 2) differences in skew between control and β-tubulin RNAi.

Figure 3 – Significant anti-correlated skews between anterior and posterior halves yield 117 candidates for transcripts with disrupted localization (A) Average skew for each gene across all control versus β-tubulin samples after filtering. Significantly skewed candidates are colored in pink. (B-E) Representative examples of genes with significantly different skews between control and β-tubulin knockdown cells.

Table 1. Top 10 positive-difference skews. Positive-difference identifies transcripts that shift posterior in β-tubulin knockdown.

Gene	Skew	Protein Domain
SteCoe_32460	0.043	EF-hand_7 domain protein
SteCoe_23664	0.042	MORN domain protein
SteCoe_3417	0.034	Ank_2 domain protein
SteCoe_17314	0.033	NIF domain protein
SteCoe_27943	0.029	Mito_carr domain protein
SteCoe_18611	0.026	JAB domain protein
SteCoe_18056	0.023	Toprim_2 domain protein
SteCoe_28593	0.022	JAB domain protein
SteCoe_26480	0.021	Ribosomal_L24e domain protein
SteCoe_28782	0.020	cNMP_binding domain protein

Table 2. Top 10 negative-difference skews. Negative-difference identifies transcripts that shift anterior in β-tubulin knockdown.

Gene	Skew	Protein Domain
SteCoe_9817	-0.063	CK1 kinase
SteCoe_18211	-0.054	ZZ domain protein
SteCoe_13508	-0.052	RRM_1 domain protein
SteCoe_20037	-0.048	Kelch_4 domain protein
SteCoe_14279	-0.046	IBR domain protein
SteCoe_3635	-0.044	Carb_anhydrase domain protein
SteCoe_13813	-0.043	CK1 kinase
SteCoe_24682	-0.042	Lectin_leg-like domain protein
SteCoe_20403	-0.042	Aldo_ket_red domain protein
SteCoe_26355	-0.041	Ion_trans domain protein

Discussion

Here, we provide evidence that the microtubule cytoskeleton is broadly important for proper RNA localization in the giant ciliate, *Stentor coeruleus.* Knockdown of β-tubulin, a major component of the microtubule cytoskeleton, disrupts normal cell morphology and causes cells to lose their orderly microtubule bundles (Fig 1, [10]). Half-cell RNA-sequencing in paired anteriors and posteriors of the cell also reveals global disruptions in polarity by PCA and in a more variable skew present in control versus β-tubulin knockdown cells (Fig 2). Overall, we find many candidates with significant and anti-correlated skews (Fig 3). We speculate that these candidates may be important for cellular patterning given the different enrichment as well as significantly different skew in expression upon β-tubulin knockdown where cell polarity is disrupted.

This work shows that transcript patterning is disrupted upon the loss of β-tubulin and disruption of the microtubule cytoskeleton, but raises several additional questions: (1) Are the candidates we found directly involved in cellular patterning or the regeneration of new patterned structures? (2) Is the microtubule cytoskeleton essential for regeneration given its role in patterning the cell? (3) Does protein enrichment in one half over the other arise from differential transcript localization or possibly translation? As we work on developing additional molecular tools in *Stentor,* we plan to conduct RNAi of the 5 representative candidates mentioned among others to assay for morphological and/or regeneration defects. Other candidates include for study include those listed in Table 1, Table 2, and all 117 in Supplementary Table 1. One limitation of this study is the percent pseudoalignment is low ranging from 10-20%, or 10-23 million reads per sample. We do however expect lower than typical pseudoalignment as ribosomal DNA is present at over 1 million ploidy in *Stentor* [9]. Regardless, we are encouraged by the results we found despite a relatively low number of reads per sample.

RNA localization could be a key to understanding pattern formation and regeneration in *Stentor*. Our results suggest a potential model for how this localization is achieved. Given that disruption of the cortical microtubule bundles led to a loss or reversal of

polarized localization for many messages, we propose that messages associate with microtubule motors via RNA-binding adaptors, such as BicD, and then move to the plus or minus ends of the bundles (Figure 4). Future work will be needed to test this model by interfering with motors and potential adaptor proteins.

Figure 4. Model for RNA localization in *Stentor* via motor-directed transport.

As this is a preprint-in-progress, we welcome input on additional interesting candidates for further analysis, as well as any feedback or other suggestions for this manuscript.

Acknowledgements

We would like to acknowledge Mark Slabodnick for prior work on *Stentor* as well as sharing images from Slabodnick et al 2014 for use in this manuscript, Connie Yan for assistance with *Stentor* bisection, and the Marshall Lab for feedback. A.A. was supported by NIH 5K12GM081266. This work was supported by NIH R35 GM130327, on which W.F.M. is the principal investigator.

Declaration of Competing Interests

The authors declare that they have no competing interests.

Supplementary Information

Supplementary Table 1. All 117 candidates and associated protein domains.

References

1. Kirschner M, Gerhart J, Mitchison T. Molecular "vitalism." Cell. 2000;100: 79–88.

2. Shulman JM, St Johnston D. Pattern formation in single cells. Trends Cell Biol. 1999;9: M60-4.

3. Marshall WF. Regeneration in Stentor coeruleus. Front Cell Dev Biol. 2021;9: 753625.

4. Lillie FR. On the smallest parts of stentor capable of regeneration; a contribution on the limits of divisibility of living matter. J Morphol. 1896;12: 239–249.

5. Morgan TH. REGENERATION OF PROPORTIONATE STRUCTURES IN STENTOR. Biol Bull. 1901;2: 311–328.

6. Zhang KS, Blauch LR, Huang W, Marshall WF, Tang SKY. Microfluidic guillotine reveals multiple timescales and mechanical modes of wound response in Stentor coeruleus. BMC Biol. 2021;19: 63.

7. Tartar V. The biology of Stentor. New York: Pergammon Press; 1961.

8. De Terra N. Evidence for cortical control of macronuclear behavior in Stentor. J Cell Physiol. 1971;78: 377–386.

9. Slabodnick MM, Ruby JG, Reiff SB, Swart EC, Gosai S, Prabakaran S, et al. The macronuclear genome of Stentor coeruleus reveals tiny introns in a giant cell. Curr Biol. 2017;27: 569–575.

10. Slabodnick MM, Ruby JG, Dunn JG, Feldman JL, DeRisi JL, Marshall WF. The kinase regulator mob1 acts as a patterning protein for stentor morphogenesis. PLoS Biol. 2014;12: e1001861.

11. Sood P, Lin A, Yan C, McGillivary R, Diaz U, Makushok T, et al. Modular, cascade-like transcriptional program of regeneration in Stentor. Elife. 2022;11. doi:10.7554/eLife.80778

12. Tartar V. Grafting experiments concerning primordium formation in Stentor coeruleus. J Exp Zool. 1956;131: 75–121.

13. Frankel J. Pattern Formation. New York, NY: Oxford University Press; 1990.

14. Wloga D, Frankel J. From molecules to morphology: cellular organization of Tetrahymena thermophila. Methods Cell Biol. 2012;109: 83–140.

15. Cole E, Gaertig J. Anterior-posterior pattern formation in ciliates. J Eukaryot Microbiol. 2022;69: e12890.

16. Lin A, Piehowski PD, Tsai C-F, Makushok T, Yi L, Diaz U, et al. Determining protein polarization proteome-wide using physical dissection of individual Stentor coeruleus cells. Curr Biol. 2022;32: 2300-2308.e4.

17. Combs PA, Eisen MB. Genome-wide measurement of spatial expression in patterning mutants of Drosophila melanogaster. F1000Res. 2017;6: 41.

18. Haines JE, Eisen MB. Patterns of chromatin accessibility along the anterior-posterior axis in the early Drosophila embryo. PLoS Genet. 2018;14: e1007367.

19. Raj A, van den Bogaard P, Rifkin SA, van Oudenaarden A, Tyagi S. Imaging individual mRNA molecules using multiple singly labeled probes. Nat Methods. 2008;5: 877–879.

20. Xia C, Fan J, Emanuel G, Hao J, Zhuang X. Spatial transcriptome profiling by MERFISH reveals subcellular RNA compartmentalization and cell cycle-dependent gene expression. Proc Natl Acad Sci U S A. 2019;116: 19490–19499.

21. Takizawa PA, Sil A, Swedlow JR, Herskowitz I, Vale RD. Actin-dependent localization of an RNA encoding a cell-fate determinant in yeast. Nature. 1997;389: 90–93.

22. Litman P, Barg J, Ginzburg I. Microtubules are involved in the localization of tau mRNA in primary neuronal cell cultures. Neuron. 1994;13: 1463–1474.

23. Dodt M, Roehr JT, Ahmed R, Dieterich C. FLEXBAR-flexible barcode and adapter processing for next-generation sequencing platforms. Biology (Basel). 2012;1: 895–905.

24. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. Nat Biotechnol. 2016;34: 525–527.

25. Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. Differential analysis of RNA-seq incorporating quantification uncertainty. Nat Methods. 2017;14: 687–690.

26. Huang B, Pitelka DR. The contractile process in the ciliate, Stentor coeruleus. I. The role of microtubules and filaments. J Cell Biol. 1973;57: 704–728.

27. Maloney MS, McDaniel WS, Locknar SA, Torlina HM. Identification and localization of a protein immunologically related to Caltractin (Centrin) in the Myonemes and Membranelles of the Heterotrich ciliate Stentor coeruleus. J Eukaryot Microbiol. 2005;52: 328–338.

28. Rothé B, Gagnieux C, Leal-Esteban LC, Constam DB. Role of the RNA-binding protein Bicaudal-C1 and interacting factors in cystic kidney diseases. Cell Signal. 2020;68: 109499.

29. Estrada Mallarino L, Engel C, Ilık İA, Maticzka D, Heyl F, Müller B, et al. Nephronophthisis gene products display RNA-binding properties and are recruited to stress granules. Sci Rep. 2020;10: 15954.

30. Brambilla M, Martani F, Bertacchi S, Vitangeli I, Branduardi P. The Saccharomyces cerevisiae poly (A) binding protein (Pab1): Master regulator of mRNA metabolism and cell physiology. Yeast. 2019;36: 23–34.