A class-specific effect of dysmyelination on the excitability of hippocampal interneurons

Delphine Pinatel[1], Edouard Pearlstein[1], Giulia Bonetto[1,4], Laurence Goutebroze[2], Domna Karagogeos[3], Valérie Crépel[1] and Catherine Faivre-Sarrailh[1*]

[1]Aix Marseille Université, INSERM UMR1249, Institut de Neurobiologie de la Méditerranée, F-13273 Marseille, France.
[2]INSERM UMR-S 1270, Institut du Fer à Moulin, Sorbonne Université, Faculté des Sciences et Ingénierie, F-75005, Paris, France.
[3]Department of Basic Sciences, University of Crete Medical School and Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology, University of Crete, Heraklion, Greece.
[4]Present address: AstraZeneca Ltd, UK.

Correspondence: catherine.sarrailh@univ-amu.fr

Number of words: 11,950

Number of Figures: Figures: 10, Supplementary Figures: 4

Key-words: protein 4.1B, juxtaparanodes, myelin, parvalbumin, somatostatin, interneuron, hippocampus

Running title: dysmyelination of hippocampal interneurons

Acknowledgments

We are grateful to Dr Thomas Marissal for helpful discussions. We thank the University of California Davis/National Institutes of Health NeuroMab Facility and Developmental Studies Hybridoma Bank of the University of Iowa. This work was supported by the Association pour la Recherche sur la Sclérose en Plaques (ARSEP) to DP, CFS and DK.

Abstract

The role of myelination for axonal conduction is well-established in projection neurons but little is known about its significance in GABAergic interneurons. Myelination is discontinuous along interneuron axons and the mechanisms controlling myelin patterning and segregation of ion channels at the nodes of Ranvier have not been elucidated. Protein 4.1B is implicated in the organization of the nodes of Ranvier as a linker between paranodal and juxtaparanodal membrane proteins to the spectrin cytoskeleton. In the present study, 4.1B KO mice are used as a genetic model to analyze the functional role of myelin in Lhx6-positive parvalbumin and somatostatin neurons, two major classes of GABAergic neurons in the hippocampus. We show that deletion of 4.1B induces disruption of juxtaparanodal K^+ channel clustering and mislocalization of nodal or heminodal Na^+ channels. Strikingly, 4.1B-deficiency causes loss of myelin in GABAergic axons in the hippocampus. In particular, stratum oriens O-LM cells display severe axonal dysmyelination and a reduced excitability. This reduced excitability is associated with a decrease in occurrence probability of small amplitude synaptic inhibitory events on pyramidal cells. In contrast, stratum pyramidale fast-spiking basket cells do not appear affected. The aberrant myelination of hippocampal interneurons is also correlated with impairment of spatial memory in 4.1B KO mice. In conclusion, our results indicate a class-specific effect of dysmyelination on the excitability of hippocampal interneurons associated with a functional alteration of inhibitory drive and impairment of spatial memory.

Introduction

Recent reports have established that subtypes of GABAergic neurons could be myelinated. Indeed, a substantial fraction of myelin, both in mouse and human neocortex, belongs to inhibitory neurons by comparison with pyramidal cells (Micheva et al., 2016; Stedehouder et al., 2017). Long-range projecting hippocampal GABAergic neurons are myelinated and play a role in the synchronization among distant brain areas (Jinno et al., 2007; Melzer et al., 2012). Surprisingly, locally-projecting interneurons such as the parvalbumin (PV) cells are frequently myelinated, independently of their morphological subtypes (i.e. basket or bi-stratified) in the mouse hippocampus. A fraction of somatostatin (SST) interneurons, which include both long-range and Oriens-Lacunosum Moleculare (O-LM) or bistratified local projecting neurons has been also reported to be myelinated (Micheva et al., 2016; Stedehouder et al., 2017, 2019). Myelin optimizes action potential (AP) propagation by forming electrical insulation and restricting ion channel distribution, but its significance is still elusive in local projecting interneurons. Oligodendrocytes wrapping of GABAergic axons may be essential for providing axonal metabolic support (Krasnow and Attwell, 2016; Philips and Rothstein, 2017). Recent reports indicate that myelination of PV interneurons is modulated by neuronal activity or sensory experience and is able to shape the inhibitory circuitry (Benamer et al., 2020; Stedehouder et al., 2018; Yang et al., 2020). However, how myelination could modulate the function of the different subtypes of interneurons remains poorly evaluated.

In the neocortex and hippocampus, GABAergic interneurons present the particularity to be covered by myelin sheaths that display a patchy distribution along the axons (Benamer et al., 2020; Micheva et al., 2016; Stedehouder et al., 2017).The discontinuous myelination of inhibitory ramified axons implies the presence of heminodal structures along mature axons. The mechanisms regulating myelin coverage as well as segregation of ion channels at heminodes and nodes of Ranvier remain to be elucidated. Remarkably, hippocampal PV and SST neurons display clusters of Na^+ channels forming prenodes along the axons before myelination (Bonetto et al., 2019; Freeman et al., 2016). These prenodes can be induced by oligodendrocyte-secreted matrix molecules including Contactin (Dubessy et al., 2019). Such Na^+ channel clusters along pre-myelinated inhibitory axons are associated with increased conduction velocity (Freeman et al., 2015). We also showed that the axons of hippocampal PV and SST interneurons are highly enriched before myelination in Kv1 channels that may regulate firing during development. The Kv1 channels are associated with the cell adhesion molecules Contactin2/TAG-1, Caspr2, and ADAM22 and the scaffolding protein 4.1B forming complexes in the juxtaparanodal domains of the nodes of Ranvier after myelination (Bonetto et al., 2019). It is however so far unknown whether deficiency in any of the juxtaparanodal components Contactin2/TAG-1, Caspr2 or 4.1B could affect the myelination of hippocampal interneurons. Here, we focused our study on the consequences of 4.1B deficiency.

Protein 4.1B belongs to a superfamily of proteins that share a conserved FERM domain (4.1/ ezrin/radixin/moesin) and serve as membrane cytoskeleton linkers and participate in a wide variety of cellular events such as motility and cell adhesion (Denisenko-Nehrbass et al., 2003). In myelinated axons, protein 4.1B is implicated in the organization of the nodes of Ranvier as a linker between

paranodal and juxtaparanodal membrane proteins to the actin/spectrin cytoskeleton. At juxtaparanodes, 4.1B is linked to the Caspr2/Contactin2 cell adhesion complex and is required for the clustering of the Kv1 channels as reported in the PNS and CNS (Buttermore et al., 2011; Cifuentes-Diaz et al., 2011; Einheber et al., 2013; Horresh et al., 2010). At paranodes, 4.1B mediates the linkage between the Caspr/Contactin adhesion complex and ßII-spectrin and participates to the boundary with the nodal complex enriched in Nav channels and Neurofascin186 anchored to ßIV-spectrin/AnkyrinG scaffold (Brivio et al., 2017; Zonta et al., 2008). In addition to its role in the organization of ion channel domains at the nodes of Ranvier, 4.1B linked with Caspr at paranodes has been also reported to promote internodal elongation during development (Brivio et al., 2017).

We discovered that *4.1B*$^{-/-}$ mice display selective aberrant myelination of GABAergic interneurons in the hippocampus. An intriguing question is whether hippocampal interneurons require paranodal cytoskeleton as an instructive cue regulating the extent of myelin sheath coverage. We study the cellular features of SST and PV cells in the hippocampus by examining ion channel distribution, axon initial segment (AIS) topography and electrophysiological intrinsic properties. We show that 4.1B-deficiency induces a dramatic and selective loss of myelin sheath along SST and PV axons in the stratum radiatum of CA1 hippocampus. Notably, the excitability of O-LM cells located in the stratum oriens is specifically reduced in contrast to PV basket cells not affected in the stratum pyramidale. This study reveals the functional role of myelin in affecting the excitability of subtypes of hippocampal interneurons and modulating spatial-related behavior.

Results

Deletion of the scaffolding protein 4.1B causes severe loss of myelination in inhibitory axons of the stratum radiatum in the hippocampus

We used the transgenic *Lhx6-tdTomato* reporter line (Jackson laboratory) to fluorescently label subtypes of GABAergic neurons. Lhx6 is a transcription factor expressed in inhibitory neuron subclasses originating from the medial ganglionic eminence, which includes all hippocampal PV and SST neurons (Liodis et al., 2007). Double-immunostaining for MBP as a myelin marker, and Caspr as a paranodal marker was performed and we evaluated that 75 ± 5 % of the myelinated axons were Lhx6-positive in the stratum radiatum at P35 (Fig. S1A, D).

We showed previously that cell adhesion and scaffolding molecules associated in complex with juxtaparanodal Kv1 are highly expressed in hippocampal GABAergic interneurons (Bonetto et al., 2019). Contactin2 is selectively expressed in SST cells in the stratum oriens and PV cells in the stratum pyramidale of the rat hippocampus (Bonetto et al., 2019). Here we ask whether deficiency in any of the juxtaparanodal components Contactin2/TAG-1, Caspr2 or 4.1B could induce dysmyelination of hippocampal interneurons. Strikingly, we observed that 4.1B KO mice displayed a severe loss of myelin sheaths in the hippocampus as shown in Fig.1. The myelin pattern was selectively altered in the hippocampus of the 4.1B KO mice, markedly in the stratum radiatum both at P35 and P70 (Fig. 1B, F

Figure 1: 4.1B KO mice show strong alteration of myelin in the adult hippocampus

A-B: Hippocampal vibratome sections from P35 wild-type and *4.1B⁻/⁻* mice immunostained for MBP as a myelin marker. Note the lack of myelin sheaths in the CA1 stratum radiatum of the *4.1B⁻/⁻* hippocampus whereas myelin of the corpus callosum is preserved (compare B with A). C-F: Hippocampus from P70 wild-type, *Contactin2⁻/⁻*, *Caspr2⁻/⁻*, and *4.1B⁻/⁻* mice immunostained for MBP. Only the *4.1B⁻/⁻* mice show a selective and massive loss of myelin in the stratum radiatum. G, H: Double-staining for MBP (blue) and Caspr (green) as a paranodal marker. Maximum intensity of confocal images (5-z steps of 2 μm). F: Quantification of the number of paranodes/10^4μm² in the different layers of the CA1 hippocampus. G: Quantification of the total myelin length/10^4μm² in the sp (bin1) and sr divided in 5 bins (40 x300 μm). Mean ± SEM of 3-4 mice for each genotype. Significant difference by comparison with wild-type: * P<0.05 and ** P<0.01 using the Student's t-test. CC: corpus callosum, so: stratum oriens, sp: stratum pyramidale, sr: stratum radiatum, slm: stratum lacunosum moleculare. Bar: 200 μm in A, B; 50 μm in C-F; 25 μm in G, H.

and S1H) as an indication that myelination of inhibitory interneurons may be severely affected. We did
not observe any massive loss of myelin outside the hippocampus, as shown in myelinated tracts like the
corpus callosum (Fig. 1B) or in the layers of the somato-sensory cortex (Fig. S2). In contrast, Contactin2-
deficiency did not markedly impair myelination in the CA1 hippocampus as illustrated at P70 (Fig. 1D).
In *Contactin2-/-* mice crossed with the *Lhx6-tdTomato* reporter line, the total length of myelinated axons
and the percentage of Lhx6-positive myelinated axons in the stratum radiatum were similar to control
mice as quantified at P35 (Fig. S1A-D). As reported in several studies, Caspr2 KO mice show a reduced
density of PV-positive interneurons in the hippocampus associated with decreased inhibitory synaptic
transmission in CA1 pyramidal neurons (Paterno et al., 2021; Penagarikano et al., 2011). However, we
did not observe any major alteration of myelinated PV inhibitory axons in the hippocampus at P70 (Fig.
1E and S1E, F).

In the hippocampus of *4.1B-/-* mice, the severe reduction of myelin sheaths occurred already at
P25 in the stratum radiatum indicating that it originated from a developmental defect, taking place at an
early stage of hippocampal myelination (Fig. S3). Since the extent of myelination was difficult to be
quantified in regions with a dense and intermingled network of myelinated axons, the number of
internodes was estimated from the number of paranodes immunolabeled for Caspr in the different layers
of the CA1 hippocampus at P70 (Fig. 1G-I). The density of paranodes was significantly reduced by 88%
(P=0.0147, Student's t test) in the stratum radiatum of 4.1B-deficient mice (n=3-4 mice/genotype). The
density of paranodes tended also to decrease in the stratum oriens (-18%; P=0.370), stratum pyramidale
(-30%; P=0.140) and stratum lacunosum-moleculare (-31%; P=0.051). The total length of myelinated
axonal segments of the stratum radiatum was measured at different levels starting from the pyramidal
layer towards the stratum lacunosum-moleculare in the CA1 hippocampus (Fig. 1J). Myelination was
significantly decreased by 75% (P=0.0015, Student's t test) in the stratum radiatum of 4.1B-deficient
mice, with a 94 % reduction in the middle region of the layer (P=0.0015). Therefore, the fact that myelin
mostly belongs to Lhx6-positive axons in the stratum radiatum, suggests that myelination of inhibitory
interneurons may be selectively altered in the hippocampus of 4.1B-deficient mice.

It was first critical to evaluate whether the lack of myelin could be due to a loss of GABAergic
interneurons extending their axons throughout the stratum radiatum. Distinct interneurons subdivide
their connections to sub-compartments of pyramidal neurons with the PV basket cells connecting the
soma, the bistratified cells connecting basal and apical dendrites in the stratum oriens and radiatum,
and the SST O-LM cells projecting to the distal apical dendrites in the stratum lacunosum-moleculare
(Somogyi and Klausberger, 2005) (see schematic illustration Fig. 7A, H). The 4.1B KO mice did not
show a reduced density of PV cells in the stratum pyramidale (Fig. 2A, B) as quantified at P35 in the
CA1 region (Fig. 2E). A high density of PV axons was present in the stratum radiatum of the 4.1B-
deficient mice (Fig. 2H) as observed in the wild-type (Fig. 2G), albeit they were only partly myelinated
in the region bordering the stratum pyramidale (arrow in Fig. 2H) and the total length of myelinated PV
axons was significantly reduced (-63%, P=0.0317, Mann-Whitney test) (Fig. 2K). The density of SST
interneurons in the stratum oriens was also preserved (Fig. 2C, D, F). The axons of O-LM interneurons

Figure 2

Figure 2: Parvalbumin and somatostatin axons are dysmyelinated in the _4.1B^{-/-}_ hippocampus

A-D: Hippocampal sections of wild-type and _4.1B^{-/-}_ mice immunostained for PV (A, B) or SST (C, D) at P35. E, F: Quantification of the cell density/10^5 μm^2 in the CA1 stratum oriens (SO) and stratum pyramidale (SP). No statistical difference in cell density between the genotypes (PV cell density: SO, P=0.108; SP, P=0.412 and SST cell density: SO, P=0.413; SP, P=0.057; Student's t test; n=10-12 ROIs in hippocampal sections at 2 levels in the antero-posterior axis, 3 mice/genotype). G, H: Double-staining for PV (red) and MBP (green) at P70 showing that PV axons are present in the stratum radiatum and poorly myelinated in the _4.1B^{-/-}_ mice (H). The arrow in H points to a PV axon, which is partly myelinated. K: Quantification of the total length of PV myelinated axon/10^5 μm^2 in the stratum radiatum at P70 (n=5 ROIs, 3 mice/genotype). I, J: Double-staining for SST (green) and MBP (blue) at P35 showing O-LM SST cells in the stratum oriens extending their axon through the stratum radiatum to project into the stratum lacunosum moleculare. Note that SST axons in the stratum radiatum are fully myelinated in the WT (asterisks in I) and partly myelinated in _4.1B^{-/-}_ mice (arrows in J). L, M Quantitative analyses of the total length of SST myelinated axon/10^5 μm^2 in the stratum radiatum at P35 (n=10 ROIs, 3 mice/genotype) and the percentage of myelin coverage of individual SST axons (n=29-35 axons, 3 mice/genotype; the mean length of selected axons is 174±9 µm in WT and 197±12 µm in _4.1B^{-/-}_ mice). Significant difference by comparison with wild-type: * P<0.05; *** P<0.001; Mann-Whitney test.

Figure 3: The density of oligodendroglial cells is preserved in the *4.1B$^{-/-}$* hippocampus
Hippocampal sections of wild-type and *4.1B$^{-/-}$* mice at P35 immunostained for MBP (blue), SST (red) and Olig2 (green) as a marker of the oligodendrocyte lineage. Bar: 50 µm. Quantitative analysis of the density in Olig2-positive cells/10^5 µm^2 in the different layers of the CA1 hippocampus (n=11-12 ROIs, 3 mice/genotype). No significant difference between the genotypes (so: P=0.320; sp: P=0.090; sr: P=0.192; slm, P=0.829; Student's t test).

crossing the stratum radiatum and projecting to the stratum lacunosum-moleculare were observed in
the 4.1B-deficient mice (Fig. 2J) as in the wild-type (Fig. 2I). However, O-LM axons were only partly
myelinated in the lower and upper regions of the stratum radiatum (arrows in Fig. 2J) in the 4.1B KO
mice with a myelin coverage of 42.3 ± 4.0% (Fig. 2M) while myelin coverage was 79.7 ± 3.4% for wild-
type mice (Fig. 2I and M). The total length of myelinated SST axons was significantly reduced (-58%,
P<0.0001, Mann-Whitney test) in the stratum radiatum (Fig. 2L). Thus, 4.1B-deficiency strongly reduces
myelin ensheathing of PV and SST axons in the stratum radiatum.
199
Next, we asked whether such myelin loss could be due to a reduced number of oligodendrocytes
in the hippocampus using immunostaining for Olig2 as a marker for the oligodendrocyte lineage (Fig.
3A, B). The density of Olig2-positive cells in the stratum radiatum was 22.3 ± 1.6/10^5 μm^2 in the 4.1B
KO mice versus 25.3 ± 2.2/10^5 μm^2 in wild-type animals at P35 (Fig. 3C). We did not observe any
decrease in the number of Olig2-positive cells in the CA1 layers that could explain the dramatic loss of
myelin in 4.1B KO mice.
206

4.1B-deficient GABAergic axons display reduced internodal length
Protein 4.1B mediates interactions between membrane adhesion complexes and the spectrin
cytoskeleton and binds Caspr at paranodes as schematized in Fig. 4A. We previously showed that 4.1B
promotes via its interaction with Caspr, the growth of oligodendroglial processes and myelin sheath
convergence in the developing spinal cord (Brivio et al., 2017). Immunofluorescence staining for 4.1B
in the hippocampus indicated that the scaffolding protein was expressed by GABAergic PV interneurons
and not by pyramidal neurons in the CA1 pyramidal layer at P70 (Fig. 4C). As a control for specificity,
no staining was observed on hippocampal sections from *4.1B*$^{-/-}$ mice (Fig. 4D). Protein 4.1B
immunostaining was present around the soma of PV interneurons, along the internodes and enriched
at paranodes whereas the nodal gap was unlabeled (Fig. 4E, E'). We asked whether the elongation of
internodes could be affected in the 4.1B-deficient CA1 hippocampus at P70 and analyzed the distribution
of the length of internodes (Fig. 4B). Analysis of the cumulative frequency indicates a significant
difference between the genotypes (P<0.0001; Kolmogorov-Smirnov test). The mean internodal length
was significantly reduced (P=0.0286, Mann-Whitney test) in 4.1B KO (34.8 ± 2.9 μm; n=191) by
comparison with wild-type (43.8 ± 0.3 μm; n= 275) mice. The maximum length was similarly decreased
to 93.8 μm in the 4.1B KO versus 114.5 μm in the wild-type mice. The reduction of internodal length
strongly indicates that protein 4.1B may be implicated in myelin sheath elongation.
224

4.1B-deficiency causes misdistribution of nodal and heminodal Nav channels in hippocampal
GABAergic interneurons
At the paranodal junction, the axonal cytoskeleton including 4.1B and ßII-spectrin also organizes the
boundary between the internodes and the nodal Nav channels (Fig. 4A) (Brivio et al., 2017; Buttermore
et al., 2011; Zhang et al., 2013; Zonta et al., 2008). Accordingly, the Nav complex is mislocalized distant
from the paranodes in the developing spinal cord of the 4.1B KO mice before the convergence of
internodes (Brivio et al., 2017). In the adult spinal cord, this phenotype is no longer observed and the

Figure 4: 4.1B is enriched at paranodes of GABAergic axons and involved in internodal elongation

A: Schematic drawing to depict the role of 4.1B at paranodes mediating the anchorage between Caspr and the ßII-spectrin at the boundary with the nodal ßIV-spectrin. B: Distribution of internodal lengths as percentage. Quantitative analysis was performed on hippocampal sections of *Lhx6-Cre;tdTomato* control and *4.1B⁻ᐟ⁻* mice at P70 immunostained for MBP and Caspr. Confocal imaging of 25 to 50-μm width z-stack were used to measure the length of myelin sheaths between two paranodes in the CA1 region (n= 275 in control and n=191 in *4.1B⁻ᐟ⁻* mice). Significant difference by comparison with control: * P<0.05; Mann-Whitney test (Means ± SEM of 4 ROIs from 2 mice/genotype). C-E: Hippocampal sections of wild-type and *4.1B⁻ᐟ⁻* mice at P35 immunostained for MBP (blue), PV (red) and 4.1B (green). Note that 4.1B is detected at the soma (C) and along the axons of PV interneurons, being enriched at paranodes and not detected at the nodal gap (E, E'). As a control for specificity, no staining is detected in *4.1B⁻ᐟ⁻* hippocampus (D). Bar: 10 μm in C, D; 5 μm in E, E'.

nodal Nav clusters become properly juxtaposed to the paranodes in mature nodes. Hippocampal
inhibitory axons are covered by discontinuous myelin with persistence of heminodal structures in the
adult, and myelination of ramified axons implies the presence of nodes at branch points so that it was
particularly interesting to analyze whether the distribution of Nav channels may be disturbed in the 4.1B-
deficient axons.

We first analyzed the architecture of the nodes of Ranvier in the GABAergic interneurons of the
CA1 hippocampus of wild-type mice at P30 and P70 by performing panNav channel immunostaining.
As illustrated in Fig. 5A, myelination along Lhx6-positive axons was partly discontinuous in the stratum
radiatum at P70, leaving uncovered axonal segments (asterisks). Nav channels were detected at nodes
(Fig. 5B, B', G) and branch points (Fig. 5C, C'). Heminodal clustering of Nav channels was observed for
interspaced internodes (Fig. 5E). We noticed that the lengths of nodal Nav immunostaining measured
between two paranodes were higher at P30 than at P70. Analysis of the cumulative frequency indicates
that the distribution of nodal Nav immunostaining lengths was significantly different with age (P=0.0264;
Kolmogorov-Smirnov test). At the age of P30, 44% of the nodes displayed a length > 2 µm versus only
22 % at P70, indicating that the internodes may still elongate during the late developmental stage (Fig.
5D).

We then investigated whether 4.1B deficiency may induce alteration of the distribution of nodal
and heminodal Nav channels in the CA1 hippocampus at P70. In 4.1B-deficient mice, a number of
inhibitory PV or SST axons display interrupted myelination when entering the stratum radiatum showing
heminodes (Fig. 2H, J and S4B). Interestingly, we observed at heminodes of 4.1B-deficient inhibitory
axons, a gap between the site of Nav clustering and the paranode stained for Caspr (Fig. 5F and Fig.
S4B) as compared to control mice (Fig. 5E and S4A). The gap could reach 10 µm and displayed a mean
value of 2.25 ± 0.51 µm significantly different from the control value (0.04 ± 0.02 µm) (n=25 heminodes,
2 mice/genotype; P<0.0001, Mann-Whitney test) (Fig. 5I). By contrast, the length of heminodal Nav
clusters was similar in control (1.74 ± 0.19 µm) and in 4.1B KO (1.84 ± 0.12 µm) mice (P=0.427, Mann-
Whitney test) (Fig. 5J). A gap could also be observed between nodal Nav and convergent internodes
stained for MBP (Fig. 5H) in 4.1B-deficient mice by comparison with control ones (P=0.0029, Mann-
Whitney test) (Fig. 5I), together with an increase in the length of nodal Nav clusters (Fig. 5J). The mean
length of nodal Nav in 4.1B KO (1.92 ± 0.25 µm, n=26) was significantly increased by comparison with
control mice (1.59 ± 0.14 µm, n=87) (P=0.0068, Mann-Whitney test). Thus, the positioning of Nav
clusters is significantly disturbed at the nodes and heminodes of dysmyelinated GABAergic interneurons
of 4.1B-deficient hippocampus.

Juxtaparanodal alterations in hippocampal interneurons of Contactin2-, Caspr2- and 4.1B null
mutant mice
Finally, we analyzed the distribution of the juxtaparanodal Kv1 channels in the stratum radiatum at P70.
As expected, immunostaining for Kv1.2 was detected at the juxtaparanodes or hemi-juxtaparanodes
under the compact myelin including at branch points both in Lhx6- and PV-positive axons (Fig. 6A, B).

Figure 5: The positioning of heminodal and nodal of Nav channels is altered in 4.1B-deficient GABAergic axons

Hippocampal sections of P70 *Lhx6-Cre;tdTomato* control and *4.1B⁻/⁻* mice were immunostained for MBP (blue), Caspr (white) and panNav (green). tdTomato expressed in Lhx6-positive cells or PV immunostaining (C) is in red. A: shows a branched myelinated Lhx6 axon with contiguous (arrowhead) and spaced (asterisks) internodes. Immunostaining for panNav at nodes with different lengths (B, B') and at branch points (C, C'). D: Distribution of nodal lengths in Lhx6-positive axons at P30 and P70 in CA1 expressed as percentages. The length of nodal Nav was measured between two paranodes (n=41 at P30 and n=64 at P70). E-J: Clustering of Nav channels at the heminodes (E, F) and nodes (G, H) of control and *4.1B⁻/⁻* mice in the CA1 hippocampus. Nav clusters are close to paranodes both at heminodes (E) and nodes (G) in control GABAergic axons (arrows). In contrast, a gap is frequently observed between Nav clusters and paranodes at heminodes (F) and nodes (H) in 4.1B-deficient GABAergic axons (arrows). Confocal images with maximum intensity of 2-z or 3-z steps of 1 µm. Bar: 10 µm in A; 5 µm in B, C, E-H. I: Quantitative analysis of the distance between Nav cluster and paranode at heminodes and nodes. J: Quantitative analysis of the length of nodal Nav cluster at heminodes and nodes. Significant difference by comparison with wild-type: ** P<0.01; *** P<0.001; Mann-Whitney test (heminodes: n= 25/genotype; nodes: n=87 in control and n=26 in 4.1B KO mice).

Figure 6: The clustering of juxtaparanodal Kv1 channels is reduced in *Contactin2-/-, Caspr2-/-*, and *4.1B-/-* GABAergic axons

Immunostaining for Kv1.2 channels in GABAergic axons of the stratum radiatum in CA1 at P70. A, H: Juxtaparanodal clustering of Kv1 channels in *Lhx6-Cre;tdTomato* control mouse immunostained for MBP (blue), Caspr (white) and Kv1.2 (green). tdTomato expressed in Lhx6-positive cells is in red. B: Juxtaparanodal clustering of Kv1 channels (green) in wild-type at PV axon branch point (red) with paranodes stained for Caspr (blue). C: Quantification of the percentage of paranodes associated with Kv1.2 clustering in the PV axons of the CA1 stratum radiatum. Mutant mice are compared with their respective controls. Means ± SEM of 7-11 ROIs from 3 mice/genotype. Significant difference by comparison with wild-type: * P<0.05; *** P<0.001 using the Mann-Whitney test. D-G: Immunostaining for Kv1.2 (green) in wild-type (D), *Contactin2-/-* (E), *Caspr2-/-* (F), and *4.1B-/-* (G) in PV axons (red) with paranodes stained for Caspr (blue). Yellow arrows point to juxtaparanodal clustering of Kv1 in control or mutant mice and white arrows indicate the lack of proper Kv1 clustering in mutants. Alteration of Kv1 clustering is also observed in *4.1B-/-; Lhx6-Cre;tdTomato* mice (I compared to control in H). Bar: 5 μm.

We investigated whether the deficiency in Kv1-associated CAMs may induce alteration of the
juxtaparanodes in hippocampal interneurons like it was observed in myelinated tracts of the corpus
callosum, optic nerves, or spinal cord for Contactin2 KO and Caspr2 KO mice (Poliak et al., 2003;
Savvaki et al., 2008; Traka et al., 2003). As analyzed in double-blind experiments, Contactin2- or
Caspr2- deficiency induced a strong decrease in Kv1 channel clustering at the juxtaparanodes of PV
myelinated axons in the CA1 hippocampus at P70 (Fig. 6D-F) although a residual expression of Kv1
was still present at some juxtaparanodes in both mutants. The percentage of paranodes bordered by
Kv1.2 immunostaining in PV axons was significantly reduced by 30% in Contactin2 KO and 53% in
Caspr2 KO mice (P= 0.014 and P<0.0001, Mann-Whitney test, respectively) (Fig. 6C). The scaffolding
protein 4.1B is known to bind Caspr2 and to be required for the proper assembly of the juxtaparanodes
both in the PNS and CNS (Buttermore et al., 2011; Cifuentes-Diaz et al., 2011; Einheber et al., 2013).
We observed a dramatic decrease (-61%; P<0,0001, Mann-Whitney test, n=9 ROIs, 3 mice/genotype)
in the Kv1 clustering at juxtaparanodes of myelinated PV axons of 4.1B KO mice at P70 (Fig. 6G). *4.1B*
$^{-/-}$ mice crossed with the *Lhx6-Cre;tdTomato* line showed similar alteration of Kv1 clustering at
juxtaparanodes in Lhx6-positive axons as compared with control mice (Fig. 6H and I). In summary,
deficiency in each of the Kv1-associated proteins, Contactin2, Caspr2, and 4.1B induced alteration of
juxtaparanodal Kv1 clustering along myelinated GABAergic axons in the hippocampus.
289
290   The excitability of Lhx6-positive interneurons is differently affected in the stratum oriens and
291   stratum pyramidale of 4.1B KO mice
We showed that the 4.1B KO mice exhibited a dramatic and selective loss of myelin coverage of the
inhibitory axons crossing the stratum radiatum. In particular the SST O-LM interneurons showed a lack
of myelin. The PV axons crossing the stratum radiatum that may belong to bistratified interneurons were
also dysmyelinated whereas PV axons of basket cells seemed to be more preserved in the stratum
pyramidale. These observations led us to further question whether the phenotypes observed could be
associated with physiological changes in hippocampal interneurons. We first recorded the activity of
Lhx6Cre;tdTomato-positive cells in the stratum pyramidale of control and 4.1B KO mice (Figure 7A-G).
Different subtypes of Lhx6-interneurons are located in the stratum pyramidale as schematized in Fig.
7A, including the fast-spiking basket cells innervating the soma of pyramidal cells and bistratified cells
contacting their basal and apical dendrites. We performed whole-cell patch-clamp in current clamp
configuration recordings on P30 acute slices and selected fast-spiking interneurons displaying a typical
continuous non-stuttering discharge (Hu et al., 2014) with a frequency above 50 Hz under injection of a
200 pA current intensity (Fig. 7B). As shown by the *F-I* curve during injection of current steps of 100 pA
increments, the firing frequencies were unchanged in *4.1B$^{-/-}$* compared to control mice (P=0.5505, two-
way ANOVA) (Fig. 7B and C). The resting membrane potential (Vrest) (-61.38 ± 2.40 mV in *4.1B$^{-/-}$* vs -
58.63 ± 3.53 mV in controls, P=0.7391 using Mann-Whitney test) and the input resistance (Rinput)
(339.25 ± 40.15 MΩ in *4.1B$^{-/-}$* vs 381.13 ± 54.47 MΩ in controls, P=0.5737 using Mann-Whitney test)
were not changed (Fig. 7D and E). Similarly, the rheobase (25.63 ± 5.30 pA in *4.1B$^{-/-}$* vs 37.40 ± 5.59
pA in controls, P=0.0998, Mann-Whitney test) (Fig. 7F) and afterhyperpolarization (AHP) area (-309.07
± 47.69 mV.ms in *4.1B$^{-/-}$* vs -178.40 ± 60.73 mV.ms in controls, P=0.1304, Mann-Whitney test) were not

Figure 7: The excitability of O-LM interneurons in the stratum oriens is selectively decreased in *4.1B*^{-/-} mice A-G: Patch-clamp recordings of Lhx6-positive PV fast-spiking interneurons in the stratum pyramidale (SP). A: Schematic illustration of basket and bistratified (BiC) cells in CA1 located in the SP and innervating the soma and dendrites of a pyramidal cell (PC). B: Typical fast-spiking activity evoked by a 200 pA stimulation during 500 ms showing similar firing patterns of 4.1B-deficient and control basket cells. C: *F-I* relationship of mean spike frequency depending on current intensities is not different between the genotypes. The resting membrane potential (Vrest) (D), input resistance (Rinput) (E), rheobase (F) and afterhyperpolarization (AHP) area (G) are not affected in *4.1B*^{-/-} mice. Means ± SEM. Detailed statistical analysis using Mann-Whitney test, see Table 1. H-N: Patch-clamp recordings of Lhx6-positive SST interneurons in the stratum oriens (SO) from control and *4.1B*^{-/-} mice. H: O-LM and bistratified cells in CA1 located in the SO and innervating PC dendrites. I: Example of continuous spike train discharge elicited by a 120 pA depolarizing current pulse during 500 ms and voltage response to a -70 pA hyperpolarizing or in control and *4.1B*^{-/-} mice. J: *F-I* curves. The mean frequency is significantly reduced in *4.1B*-deficient inhibitory neurons compared to controls (two-way ANOVA, ***P<0.0001, **P<0.01, *P<0.05). K: The resting membrane potential is not affected. L: The Rinput is significantly decreased and the rheobase (M) is not changed. N: The AHP area of individual AP is significantly increased (*P<0.05, Mann-Whitney test). For a detailed statistical summary of intrinsic parameters, see Table 2.

Table 1. Intrinsic electrophysiological properties of fast-spiking Lhx6-interneurons in the stratum pyramidale

	Control (mean±SEM, n=8)	4.1B$^{-/-}$ (mean±SEM, n=8)	MW test, *p* value
Passive properties			
Resting membrane potential (mV)	-58.63±3.53	-61.38±2.40	MW, *0.7391*
Input resistance (MΩ)	381.13±54.47	339.25±40.15	MW, *0.5737*
Rheobase (pA)	37.40±5.59	25.63±5.30	MW, 0.0998
Action potential parameters			
Amplitude (mV)	59.33±3.73	55.22±2.51	MW, *0.4256*
Area (mV.s)	60.64±6.43	58.34±5.19	MW, *0.3823*
Half-width (ms)	0.95±0.08	0.98±0.07	MW, *0.9397*
Threshold (mV)	-46.46±2.56	-46.45±1.97	MW, *>0.9999*
AHP area (mV.ms)	-178.40±60.73	-309.07±47.69	MW, *0.1304*
AHP peak (mV)	-9.06±1.50	-9.73±0.83	MW, *0.5737*

Statistical differences between groups were calculated with a Mann-Whitney test (MW). Control: n=8 mice; *4.1B$^{-/-}$*: n=5 mice.

Table 2. Intrinsic electrophysiological properties of Lhx6-interneurons in the stratum oriens

	Control (mean±SEM, n=12)	4.1B$^{-/-}$ (mean±SEM, n=17)	MW test, *p* value
Passive properties			
Resting membrane potential (mV)	-62.17±2.93	-57.59±1.65	MW, *0.2527*
Input resistance (MΩ)	357.33±30.30	278.94±21.12	MW, *0.0236**
Rheobase (pA)	38.25±5.73	52.65±5.79	MW, 0.1221
Action potential parameters			
Amplitude (mV)	65.34±1.84	64.06±1.94	MW, *0.6788*
Area (mV.s)	111.44±17.87	104.80±8.93	MW, *0.8788*
Half-width (ms)	1.61±0.25	1.63±0.16	MW, *0.5929*
Threshold (mV)	-45.69±1.83	-43.65±1.43	MW, *0.5558*
AHP area (mV.ms)	-351.29±58.99	-526.59±53.50	MW, *0.0380**
AHP peak (mV)	-12.57±1.79	-16.24±1.25	MW, *0.1280*

Statistical differences between groups were calculated with a Mann-Whitney test (MW). Control: n=9 mice; *4.1B$^{-/-}$*: n=8 mice.

312   significantly affected (Fig. 7G). Overall, we found no significant modification in the intrinsic
313   electrophysiological properties of fast-spiking basket cells (Table 1). Thus, the PV basket cells, which
314   axons only displayed mild alteration of myelin in the stratum pyramidale, did no exhibit alteration of their
315   excitability.
316
To investigate the electrophysiological properties of O-LM cells, we next recorded
Lhx6Cre;tdTomato-positive interneurons in the stratum oriens. As shown in the schematic illustration,
different subtypes of Lhx6-interneurons are located in the stratum oriens including SST O-LM cells
innervating distal dendrites and bistratified cells connecting basal and apical dendrites of pyramidal cells
(Fig. 7H). The O-LM cells are the most abundant subtype of Lhx6-interneurons in the stratum oriens and
display continuous discharge activity in contrast to the bistratified cells (Tricoire et al., 2011). Here, we
selected Lhx6-positive O-LM cells that displayed a non-stuttering and continuous pattern of discharge
during spike trains (500 ms) as illustrated by traces for control and $4.1B^{-/-}$ mice upon 120 pA current
injection (Fig. 7I). We observed a marked decrease in the mean discharge frequency of $4.1B^{-/-}$ cells as
indicated by the *F-I* curve depending on injected current steps of 10 pA increments (P<0.0001, two-way
ANOVA). Notably, the discharge frequency was significantly decreased upon 100, 110, 120 and 130 pA
stimulation (P=0.037, P=0.018, P=0.002, P=0.014, respectively; two-way ANOVA and Sidack post-test)
(Fig. 7J). The resting membrane potential (Vrest) was unchanged (-57.59 ± 1.65 mV in $4.1B^{-/-}$ vs -62.
17 ± 2.93 mV in controls, P=0.2527, Mann-Whitney test) (Fig. 7K). Hyperpolarizing current steps were
injected for analyzing input resistance (Rinput) that was significantly decreased by 22% (278.94 ± 21.12
MΩ in $4.1B^{-/-}$ vs 357.33 ± 30.30 MΩ in controls, P=0.0236, Mann-Whitney test) (Fig. 7L). The rheobase
was not significantly changed (52.65 ± 5.79 pA in $4.1B^{-/-}$ vs 38.25 ± 5.73 pA in controls, P=0.1221, Mann-
Whitney test) (Fig. 7M). The intrinsic parameters of APs were also analyzed showing an increased AHP
area (-526.59 ± 53.50 mV.ms in $4.1B^{-/-}$ vs -351.29 ± 58.99 mV.ms in controls, P=0.0380, Mann-Whitney
test) (Fig. 7I, N and Table 2). These data indicated that the dysmyelinating phenotype of $4.1B^{-/-}$ mice in
O-LM cells was associated with a selective decrease in their excitability.
338

Structural modification of the Axon Initial Segment of O-LM interneurons in $4.1B^{-/-}$ mice
The efficacy of spike generation is related to the structural organization of AIS, which could be modified
in dysmyelinated axons. We investigated whether the decreased excitability in SST O-LM cells may be
due to differences in the localization of AP initiation site. We first used the method of multiple derivatives
of the somatic AP signal that allows to visualize a separation of the axonal and somatodendritic APs
(Fig. 8A and B). Our results showed that 58% of control stratum oriens interneurons including O-LM
cells had a stationary inflection in the rising phase of the AP (Fig. 8A2 and C top pie chart). We calculated
the second (d^2V/dt^2) and third (d^3V/dt^3) derivatives to better illustrate separation of two inflection points
(Fig. 8A3, arrow in A4) as an indication of the distant site of the AP initiation from the soma. In contrast,
64% of $4.1B^{-/-}$ interneurons did not display stationary inflection (Fig. 8B2-B4 and C bottom pie chart).
This statistically significant difference (P=0.0017, Fisher's exact test) suggests that the AP was initiated
closer to the soma in $4.1B^{-/-}$ neurons than in control ones. This could be due to a modification of AIS
geometry such as a shortening or a displacement relative to the soma.

Figure 8: Structural modification of the AIS of O-LM interneurons in 4.1B[-/-] mice

A-C: Analysis of somatic action potentials (APs). Representative examples illustrating the voltage waveform of the AP (A1 and B1; in mV) from a control (A1) and a *4.1B[-/-]* neurons (B1). Below are presented the first (A2 and B2; dV/dt, in mV/ms), second (A3 and B3; d^2V/dt^2, in mV/ms^2), and third derivatives (A4 and B4; d^3V/dt^3, in mV/ms^3). The stationary inflection in the d^2V/dt^2 trace is indicated with asterisks. The third derivative allows to detect the second inflection point when the trace reaches or goes under zero (arrow). C: We observed a second inflection point in 58% of control (top pie graph) and 36% of *4.1B[-/-]* cells (bottom pie graph). This difference in the number of cells displaying an inflection point (asterisks in A2 and A3) is significant (P=0.017, Fisher's exact test). This could be due to the AIS shortening in *4.1B[-/-]* O-LM interneurons. D: Hippocampal sections of wild-type and *4.1B[-/-]* mice at P35 immunostained for MBP (blue), SST (red) and AnkyrinG (green). E: Length of AIS measured for AnkyrinG immunostaining of SST interneurons in the stratum oriens (SO). Significant difference by comparison with wild-type: *** P<0.001 using Mann-Whitney test. F: Distance of AIS onset from the soma is not changed between wild-type and *4.1B[-/-]* mice (P=0.099, Mann-Whitney test; n=22 in wild-type and n=32 cells in *4.1B[-/-]* mice; 3 mice/genotype). Bar: 10 µm in D.

Figure 9: Inhibitory inputs onto CA1 pyramidal neurons are affected in *4.1B-/-* mice
A: Schematic illustration of inhibitory inputs onto the soma, proximal and distal dendrites of pyramidal cells. B: Example traces of sIPSCs onto CA1 pyramidal neurons of control and *4.1B-/-* mice. C: The amplitude of sIPSCs is significantly increased in *4.1B-/-* mice (**P=0.0022) and the frequency is not affected (P=0.1320); 15 to 400 pA events analyzed during 3 min. Statistical values were performed with a Mann-Whitney test. D: Cumulative probability for the amplitude of sIPSCs (3 min, 6 cells and 4-5 mice/genotype) showing a significant different distribution between genotypes (Kolmogorov-Smirnov test, ***P<0.0001). Note that the probability of small amplitude events is reduced in *4.1B-/-* mice suggesting that distal dendritic inhibition by O-LM cells could be altered.

We thus evaluated whether the loss of myelin in hippocampal O-LM interneurons could be associated with remodeling of the AIS using immunostaining for AnkyrinG. We observed that the AIS of SST interneurons in the stratum oriens was significantly shorter (Fig. 8D, E) in 4.1B KO by comparison with wild-type mice (18.6 ± 0.7 µm, n=29 in *4.1B$^{-/-}$* vs 23.0 ± 0.9 µm, n=29 in wild-type; P=0.0004, Mann-Whitney test). The AIS onset measured from the soma or the primary dendrite was not modified (3.8 ± 0.5 µm in *4.1B$^{-/-}$* vs 4.8 ± 0.5 µm in wild-type; P=0.099, Mann-Whitney test) (Fig. 8F). Taken together, our results indicated that dysmyelination in *4.1B$^{-/-}$* mice caused a shortening of AIS and a net decrease in the intrinsic excitability of O-LM cells.

Inhibitory inputs onto CA1 pyramidal neurons are influenced by dysmyelinated *4.1B$^{-/-}$* phenotype
We then asked the question of the role of interneuron myelination in the proper inhibitory synaptic transmission on pyramidal cells. First, we recorded spontaneous inhibitory postsynaptic currents (sIPSCs) on pyramidal cells in APV (40 µM) and CNQX (10 µM) condition (Figure 9A and B). We showed a significant increase in sIPSCs peak amplitude in the *4.1B* KO mice (45.61 ± 2.93 pA in *4.1B$^{-/-}$* vs 31.51 ± 2.93 pA in controls; P=0.0022, Mann-Whitney test). The frequency of sIPSCs also tended to be increased (4.97 ± 1.02 Hz in *4.1B$^{-/-}$* vs 2.76 ± 1.13 Hz in controls, P=0.1320, Mann-Whitney test) (Fig. 9C). Interestingly, the occurrence probability of small amplitude sIPSCs was decreased in *4.1B* KO mice (P<0.0001, Kolmogorov-Smirnov test) (Fig. 9D). To further examine the properties of inhibitory synaptic inputs, we recorded miniature inhibitory postsynaptic currents (mIPSCs) on pyramidal cells as AP-independent inhibition. We found that neither the mean amplitude (28.29 ± 2.33 pA in *4.1B$^{-/-}$* mice vs 24.57 ± 2.68 pA in controls, P=0.2224, Mann-Whitney test) nor the mean frequency were affected in *4.1B$^{-/-}$* mice (2.41 ± 0.47 Hz in *4.1B$^{-/-}$* mice vs 2.73 ± 0.51 Hz in controls, P=0.5457, Mann-Whitney test). This data on mIPSCs indicated that the *4.1B$^{-/-}$* mutation may not modify the inhibitory pre-synaptic terminals connecting pyramidal cells.

The increased amplitude of sIPSCs in *4.1B$^{-/-}$* mice could reflect an enhanced proximal and a reduced distal GABAergic inhibition onto pyramidal cells. Taken together, these results suggest that the distal inhibitory inputs onto pyramidal cells may be reduced as a consequence of the dysmyelination and decreased excitability of O-LM cells.

Spatial memory is altered in *4.1B$^{-/-}$* mice
The CA1 inhibitory interneurons have been implicated in spatial learning and memory (Jeong and Singer, 2022). We tested whether the severe loss of myelin in interneurons may induce alteration of hippocampus-based place learning and memory using the Y-maze and Barnes maze tests. The spatial working memory was affected in *4.1B$^{-/-}$* mice as tested through the Y-maze (Fig. 10A). Spontaneous alternation was measured during 8 min and the percentage of alternation was significantly decreased in 4.1B KO (53.5 ± 2.9 %) versus control (64.0 ± 2.1) mice (n=8 per genotype, P=0.0068, Student's t test). The number of total entries in maze arms did not differ between the genotypes (33.6 ± 3.3 in 4.1B KO vs 35.1 ± 4.0 in control mice, P=0.7623, Student's t test) as an indication that the locomotor activity is

Figure 10: Alteration of working and spatial memory in 4.1B-deficient mice

Control and *4.1B⁻/⁻* mice with *Lhx6-Cre;tdTomato* genotype were tested for Y-maze and Barnes maze. A: Spontaneous alternation in Y-maze was measured during 8 min and the percentage of alternation was significantly decreased in 4.1B-deficient mice (n=8 per genotype, P<0.01 using the Student's t-test). B: Barnes maze task performance. Acquisition learning trials were performed during 4 days (2 trials/day) and the probe test the day after during which the escape box was removed. The latency to locate or to enter the escape box was noted and the mean performance of two trials was expressed as mean ± SEM. C-I: Exploration paths were analyzed and classified into five strategies including spatial (direct, corrected, long-correction) and non-spatial (serial or random) search strategies. Examples of serial (C) or random (F) on day 1 and direct (D) or corrected (G) strategies during the probe test are provided. The percentage of each strategies is shown for control (E) and *4.1B⁻/⁻* mice (H). I: Strategies were scaled from direct=1 (best strategy), corrected=0.75, long-correction=0.5, serial=0.25 and random=0 (worst strategy) to give a cognitive score for each trial. Significant difference of *4.1B⁻/⁻* mice by comparison with controls using two-ways ANOVA, * P<0.05 and **** P<0.0001; controls n=14, *4.1B⁻/⁻* n=11).

not affected. Next, we analyzed the Barnes maze task performance. Acquisition learning trials were
performed during 4 days (2 trials/day) and the probe test the day after during which the escape box was
removed. Decrease in latency to locate target was observed during training in each genotype as an
indication that spatial learning occurred in *4.1B$^{-/-}$* mice. We observed that the latency to target or the
latency to enter the escape box was significantly increased in the 4.1B KO mice by comparison with
controls (two-way ANOVA, genotype effect: P=0.0157 and P=0.0153, respectively; WT n=14; *4.1B$^{-/-}$*
n=11) (Fig. 10B). Since the simple quantification of latency describing performance in the Barnes maze
may result in only a partial understanding of mouse behavior and cognitive capacity, we analyzed and
classified exploration paths into five strategies including spatial (direct, corrected, long-correction) and
non-spatial (serial or random) search strategies as illustrated in Fig. 10C, D, F, G) (Cheng et al., 2019;
Illouz et al., 2016). The *4.1B$^{-/-}$* mice appeared to adopt distinct search strategies by comparison with
control mice during the learning trials. Control mice used spatial (57%), serial (29%) or random (14%)
strategies during the first acquisition day of training. The frequency of the spatial search strategy
increased to 86% for the probe test with only 14% of mice using of a serial approach (Fig. 10E). In
contrast, in 4.1B KO mice, the random approach was by far the most dominant strategy (73 %) on the
first acquisition day, the spatial search strategy being used 18% of the time. There was an effective
evolution of the *4.1B$^{-/-}$* mice to use most frequently a spatial search strategy (64%) for the probe test,
whereas 9% used serial and 27% random exploration paths (Fig. 10H). Scaling of search strategies
from direct=1 (best strategy) to random=0 (worst strategy) gives a cognitive score that was significantly
(P<0.0001, two-way ANOVA) reduced in the 4.1B KO mice (Fig. 10I). In conclusion, the selective
dysmyelination of hippocampal interneurons was associated with impairment of working and spatial
memories and cognitive capacity in the 4.1B-deficient mice.
414
415 **Discussion**
416
Our results describe for the first time a genetic model of selective myelin loss in the hippocampus. We
show that the extent of myelination of PV and SST axons is severely reduced in the stratum radiatum.
Deficiency in the scaffolding protein 4.1B also impairs Kv1 clustering at juxtaparanodes and proper
localization of Nav at the nodes of Ranvier. The excitability of the O-LM cells in the stratum oriens is
decreased while the properties of the fast-spiking interneurons in the stratum pyramidale are
unchanged. Our data suggest that this excitability defect reduces the distal inhibitory drive onto
pyramidal cells. Last but not least, behavioral tests indicate that alteration of spatial memory and
cognitive performance are related phenotypes to the *4.1B$^{-/-}$* genotype.
425
4.1B is implicated in the myelin coverage of hippocampal inhibitory axons
Despite the well-established physiological importance of myelination to speed axonal conduction,
relatively little is known about its significance for GABAergic local interneurons. Recent reports have
highlighted the fact that the different subtypes of GABAergic neurons could be myelinated and
suggested that oligodendrocytes can recognize the class identity of individual types of interneurons that
they target (Zonouzi et al., 2019). PV- and to a lesser extent SST-expressing cells in the CA1 mouse

hippocampus are frequently myelinated on their proximal axonal segments, independently of their

subtype identity (Micheva et al., 2016; Stedehouder et al., 2017) The sparse myelination of inhibitory

ramified axons implies the presence of heminodes along mature axons. One inherent feature that limits

myelin sheath growth and determines node position is the branching pattern. An intriguing question is

whether axonal cues may be instructive to determine the extent of myelin sheath coverage. It is known

that hippocampal inhibitory axons can form pre-nodal clusters of Nav channels along premyelinated

axons (Bonetto et al., 2019; Freeman et al., 2015). At a later stage of development, such prenodes of

Nav/Neurofascin/AnkyrinG may indicate positions for limiting myelin sheath extension (Malavasi et al.,

2021; Thetiot et al., 2020; Vagionitis et al., 2022). During development, components of the nodes of

Ranvier accumulate adjacent to the ends of myelin sheaths to form heminodes prior to forming nodes.

The developing myelin sheaths may stop growing when they meet a prenodal cluster, which would be

instructive for the position of the node. Indeed, mutation of the nodal CAM Neurofascin-186 has been

reported to increase internodal distance (Vagionitis et al., 2022). Here, we observed that deletion of

4.1B reduces the mean internodal length indicating that myelin sheath elongation is restricted in the

mutant mice. 4.1B-deficiency strongly reduces myelin ensheathment of both PV and SST axons in the

stratum radiatum, by 58 and 63%, respectively. The scaffolding 4.1B, which mediates the linkage

between Caspr and the ßII-spectrin may be implicated as a paranodal cue modulating the myelin

coverage of GABAergic axons. The association of 4.1 proteins with their transmembrane receptors and

the actin-spectrin cytoskeleton can be negatively modulated by phosphorylation (Wang et al., 2014) so

that 4.1B may act as a clutch to promote myelin sheath elongation depending on axonal signaling.

Whether the paranodal cytoskeleton including 4.1B may be implicated in adaptive myelination deserves

further investigation.

The localized effect of 4.1B-deficiency on the myelination of hippocampal interneurons is

intriguing. The stratified organization of the hippocampus likely reveals the selective role of 4.1B in

promoting myelin sheath elongation in inhibitory interneurons. In the stratum radiatum, we evaluated

that as much as 75 % of the myelinated axons are Lhx6-positive. This is in accordance with other studies

indicating that the fraction of myelin ensheathing GABAergic neurons is reaching up 80 % in the stratum

pyramidale and stratum radiatum (Stedehouder et al. 2017). We did not observe any dramatic loss of

myelin immunostaining in other brain areas of 4.1B KO mice as illustrated in the somato-sensory cortex

(Fig. S2). However, a precise quantitative study would be required to examine whether selective

dysmyelination of GABAergic interneurons is induced by 4.1B deficiency throughout cortical layers and

in particular, in layer2/3, in which nearly half of the myelin ensheaths inhibitory neurons (Micheva et al.,

2016; Stedehouder et al., 2017). On the other hand, the band 4.1 transcripts undergo extensive

alternative splicing and belongs to a family of proteins including 4.1G, 4.1N, and 4.1R proteins

expressed in brain (Buttermore et al., 2011; Ivanovic et al., 2012; Kang et al., 2009). Therefore, we can

hypothesize that the cell-type specific effect of 4.1B deletion may rely on compensatory mechanisms

such as upregulation of 4.1G or 4.1R, as shown in the PNS (Einheber et al., 2013; Horresh et al., 2010).

4.1B is required for the assembly of nodal and juxtaparanodal domains in myelinated inhibitory axons

In addition to the severe reduction of myelin sheath coverage, 4.1B deficiency induces alteration of
juxtaparanodal and nodal ion channels. The scaffolding protein 4.1B is linked to the Caspr2/Contactin2
cell adhesion complex at juxtaparanodes (Brivio et al., 2017; Buttermore et al., 2011; Zhang et al., 2013;
Zonta et al., 2008). As previously reported in the PNS and CNS (Buttermore et al., 2011; Cifuentes-Diaz
et al., 2011; Einheber et al., 2013; Horresh et al., 2010), juxtaparanodal Kv1 clustering is strongly
reduced in 4.1B-deficient GABAergic axons of the hippocampus. We observed a similar alteration in
Contactin2$^{-/-}$ and *Caspr2$^{-/-}$* GABAergic axons, although some residual expression of Kv1.2 can be
detected at the juxtaparanodes of the mutant mice. The Kv1 channels are localized under the compact
myelin and their function is still elusive under physiological conditions (Pinatel and Faivre-Sarrailh,
2021). On the other hand, protein 4.1B binds Caspr at paranodes where it organizes the boundary with
the nodal Nav channels. Strikingly, the nodal and heminodal Nav channels are no longer juxtaposed to
the paranodes in the adult *4.1B$^{-/-}$* mice, as previously reported in the developing spinal cord and likely
as a consequence of the disorganization of the spectrin cytoskeleton (Brivio et al., 2017).

Whether the loss of myelin in the 4.1B mutant influences ion channel expression at the AIS is
another intriguing question. We observed structural modification of the AIS of SST cells in the stratum
oriens. The length of the AIS as measured with AnkyrinG is significantly shorter in 4.1B KO by
comparison with wild-type mice, whereas its onset from the soma or the primary dendrite is not modified.
Protein 4.1B is not required for the stabilization of AnkyrinG at the AIS as shown in cultured hippocampal
interneurons from *4.1B$^{-/-}$* mice (Bonetto et al., 2019). However, 4.1B interacting with NuMA1 inhibits
endocytosis of AIS membrane proteins and is required for AIS assembly, but not maintenance, as shown
by shRNA silencing 4.1B expression at early stage of hippocampal culture (Torii et al., 2020). Therefore,
we can propose that the AIS shortening in 4.1B-deficient SST interneurons may not be a primary event
but may result from an adaptation to the reduction of myelin axonal coverage. Adaptation of the AIS has
been reported in the cuprizone chemical model of demyelination (Hamada and Kole, 2015).

Class-specific effect of dysmyelination on the excitability of hippocampal interneurons
A defining feature of inhibitory interneuron subsets is their precise axonal arborisation whereby inhibitory
synapses target specific subdomains of pyramidal neurons. PV basket cells innervate the perisomatic
region and the SST O-LM interneurons connect the distal dendrites of pyramidal neurons. We observed
a dramatic loss of myelin in the stratum radiatum whereas myelination appeared slightly affected in the
stratum pyramidale. Such a layer-specific alteration may indicate that the O-LM and bistratified
interneurons may be more affected than the basket cells. The myelinated pattern of O-LM SST
interneurons is highly visible since they extend myelinated axons from the stratum oriens straight across
the stratum radiatum. The sheath coverage of O-LM axons is dramatically reduced by 55% in the 4.1B
KO by comparison with wild-type mice.

We asked whether dysmyelination may induce changes in the neuronal excitability of
interneuron subtypes innervating the different layers of the CA1 hippocampus. In this context, we
showed that PV basket cells displayed a similar pattern of fast-spiking discharge in *4.1B$^{-/-}$* than in control

mice. Even if the rheobase tended to decrease suggesting that *4.1B^-/-* basket cells could be more

excitable than controls, we found no change in the other intrinsic parameters detected at the soma. In

contrast, the dysmyelinated phenotype of O-LM interneurons is associated with lower excitability in *4.1B^-*

^-/- mice. This was correlated with a significant decrease in the input resistance and a significant increase

in the AHP area. Such electrophysiological modifications may reflect changes in ion channel expression

at the AIS. In cortical pyramidal neurons, demyelination induced by cuprizone causes restructuring of

AIS and reduces excitability at this site. These acutely demyelinated axons displayed Kv7.3 channel

expression extended at the distal AIS (Hamada and Kole, 2015). The I_M current comprised of K_V7

channels is known to control the interevent intervals and is particularly effective in influencing firing

frequency of O-LM cells (Lawrence et al., 2006). Therefore, one can hypothesize that the increase of

AHP area in *4.1B^-/-* O-LM cells could be due to a change in Kv7 expression at the AIS. In addition, we

showed that the length of the AIS is significantly reduced without any change in its distance from the

soma, which could lead to a decrease in excitability. Indeed, structural characteristics of the AIS, such

as the length and/or distance to the soma, strongly affect the excitability and firing behavior of neurons

(Kole and Stuart, 2012; Kuba, 2012; Yamada and Kuba, 2016). Along this line, the derivative analysis

suggests that the AP is initiated closer to the soma in *4.1B^-/-* O-LM neurons than in controls. In addition,

protein 4.1B is associated in a complex with the Kv1 channels along the axon. The clustering of Kv1

channels is altered at juxtaparanodes of 4.1B-deficient mice and as a consequence, Kv1 can be diffuse

along the axon to participate in the modulation of intrinsic excitability (Rama et al., 2017).

Functionally, it is well established that PV interneurons are involved in perisomatic inhibition

while SST interneurons take part in dendritic inhibition (Somogyi and Klausberger, 2005; Udakis et al.,

2020). Thus, we investigated whether the inhibitory inputs could be modified onto the pyramidal cells.

We showed that the occurrence probability of small amplitude sIPSCs is reduced in *4.1B^-/-* mice

suggesting that the distal and not the proximal inhibitory inputs onto pyramidal cells may be decreased

as a consequence of the reduced excitability of O-LM cells. The mIPSCs recorded in pyramidal cells did

not differ in amplitude and frequency between *4.1B^+/+* and *4.1B^-/-* neurons suggesting that the number of

presynaptic terminals is similar between both genotypes. This last result indicates that dysmyelination

of inhibitory axons in CA1 may not be associated with structural synaptic alterations, but a reduction of

distal inhibitory inputs.

An important point to keep in view is that myelin sheath, apart from its insulator role, provides

metabolic and trophic support in the central nervous system (Kann et al., 2014; Krasnow and Attwell,

2016). As a possible consequence of altered trophic support, it has been reported that in cortical neurons

from lysophosphatidylcholine demyelinated mice, there is a decrease in mIPSC amplitude and

frequency consistent with a selective loss of PV inhibitory synapses (Zoupi et al., 2021). Similarly,

demyelination of cortical PV basket cells from cuprizone treated mice strongly reduces the density of

inhibitory presynaptic terminals onto the soma of pyramidal cells (Dubey et al., 2022). In contrast, we

did not evidence synaptic alterations associated with dysmyelinated interneurons in 4.1B KO mice. As

for long projecting neurons, it is also hypothesized that myelin optimizes the propagation of APs along

inhibitory axons. Myelination defects of cortical PV fast-spiking cells are associated with reduced firing
frequency and delayed predicted conduction velocity of their APs (Benamer et al., 2020). Indeed, a
positive correlation between the degree of axonal myelination and conduction velocity has been reported
for cortical PV interneurons (Micheva et al., 2021). It has been also shown that the topography of
interneuron myelination depends on the axon diameter and interbranch distance (Stedehouder et al.,
2019). Here, we observed that the unramified axonal segment of O-LM cells crossing the stratum
radiatum is ensheathed by myelin throughout. The strong reduction of myelin coverage along O-LM cell
axons in 4.1B KO mice should decrease AP conduction velocity. In acute demyelinated cortical axons,
conduction velocity of APs is reduced and ectopic APs appear; this is associated with a redistribution of
Na$^+$ channels at branch points with increased nodal expression length of Nav1.6 (Hamada and Kole,
2015). Here, we show mislocalization and increased length of nodal Nav channels that could induce
modifications in membrane capacitance and axial resistance to current flow from the node into the
internode (Arancibia-Cárcamo et al., 2017). The misdistribution of ion channels implies that AP
propagation should be altered along dysmyelinated O-LM axons in *4.1B$^{-/-}$* mice.

Selective myelin alteration of hippocampal interneurons is associated with impairment of spatial working
memory
Myelination of interneurons may play a critical role in brain functions such as learning and memory
(Bonetto et al., 2021; Yang et al., 2020). A dysfunction of hippocampal inhibitory neurons can alter theta
and gamma oscillations involved in navigation performance (Kalemaki et al., 2018; Kunz et al., 2019;
White et al., 2012). The hippocampal SST cells have been reported to contribute to learning-induced
persistent plasticity and spatial memory consolidation (Honoré et al., 2021). For example, the selective
loss of mTORC1 function in SST cells impaired spatial long-term memory as tested through the Barnes
maze with a decrease in search strategy and performance (Artinian et al., 2019). In the present study,
we observed that the spatial working memory is affected in 4.1B KO mice as tested through the Y-maze
and Barnes maze tests. In particular, using the Barnes maze, we observed that 4.1B-deficient mice
displayed altered performance and spatial search strategy during learning trials and probe test. The O-
LM cells are modulating the afferents onto pyramidal cell dendrites, facilitating through disinhibition the
Schaffer collateral pathway from CA3 and down-regulating the temporo-ammonic pathway from the
entorhinal cortex (Leão et al., 2012). The severe loss of myelin in interneurons of the hippocampus could
be associated with a dysfunction in the inhibitory network. Our results highlight the role of myelin in O-
LM interneurons in fine tuning the hippocampal inhibitory drive that may be involved in spatial exploration
and memory.

This study also addresses the contribution of dysmyelination in interneuropathies described in
multiple sclerosis. Demyelinated lesions have been reported in the hippocampus of multiple sclerosis
patients correlated with cognitive alterations (Geurts et al., 2007). GABAergic interneurons are
selectively vulnerable to demyelination in multiple sclerosis (Zoupi et al., 2021). Inhibitory network
modifications could lead to an increased incidence of epileptic seizures in multiple sclerosis patients
(Kelley and Rodriguez, 2009) and are associated with diminished cognitive function (Nicholas et al.,

2016). Here we show a subtype-specific role of myelin in hippocampal GABAergic interneurons. This
raises the intriguing question of whether subtypes of interneurons may be specifically affected in
demyelinating diseases.
594
595   **METHODS**
596
Animals
The care and use of mice in all experiments were carried out according to the European and Institutional
guidelines for the care and use of laboratory animals and approved by the local authority (laboratory's
agreement number D13-055-8, Préfecture des Bouches du Rhône). The following mouse strains were
used in this study: C57bl/6 mice (Janvier Breeding Centre), previously described *Tag-1/Contactin2* KO
mice (Traka et al., 2003), *4.1B* KO mice (Cifuentes-Diaz et al., 2011) and *Cntnap2/Caspr2* KO mice
(Poliak et al., 2003). Both *Contactin2* and *4.1B* KO mice were crossed with *Lhx6-Cre* (strain 026555)
and *Ai14* (strain 007908) mouse strains (Jackson Laboratory, Bar Harbor, ME, USA) to obtain mice with
the following genotypes *Contactin2$^{-/-}$;Lhx6-Cre;Ai14* and *4.1B$^{-/-}$;Lhx6-Cre;Ai14* expressing tdTomato.
Fixed brains from *Caspr2* KO obtained from the Jackson laboratory (strain 017482) were a gift from Dr.
N. Noraz (NeuroMyogene Institute, Lyon, France).
608
Immunofluorescence staining, confocal microscopy and image analysis
Rabbit antiserum against protein 4.1B (Cifuentes-Diaz et al., 2011) and rabbit anti-Caspr (Bonnon et al.,
2007) were described previously. The following commercial primary antibodies were used: rabbit anti-
Olig2 antibody and mouse anti-panNav mAb (clone K58/35) purchased from Sigma-Aldrich Chimie,
(Saint-Quentin-Fallavier, France), rat anti-MBP mAb (ab7349) from Abcam (Cambridge, UK), goat anti-
PV antibody (PVG-214) from Swant (Burgdorf, Switzerland), goat anti-SST mAb (sc55565) and mouse
anti-SST (sc55565) from Santa-Cruz Biotechnology (Heidelberg, Germany). Mouse anti-ankyrinG
(clone N106/36), anti-Kv1.2 (clone K14/16), anti-Kv3.1b (clone N16B/8) mAbs were obtained from
NeuroMab (UC Davis/NIH NeuroMab Facility). AlexaFluor-405, 488, -568 and -647-conjugated
secondary antibodies were purchased from Molecular Probes (Life Technologies, Courtaboeuf, France).
619
P25, P30, P35, and P70 mice were deeply anesthetized with a mix of Zoletyl/Domitor and then
transcardially perfused with PBS followed by 2% paraformaldehyde in PBS. Brains were removed and
placed in the same fixative overnight. Eighty micron-thick vibratome sections were permeabilized and
blocked for 1 h in PBS containing 5% horse serum and 0.3% Triton X-100. Antigen retrieval in 10 mM
citrate buffer, pH6, at 90°C for 30 min was performed before immunostaining with anti-panNav and anti-
AnkyrinG antibodies. Floating sections were incubated for 2 days at 4°C with combinations of the
following primary antibodies: goat anti-PV (1:1000), mouse anti-SST (1:200), goat anti-SST (1:500),
rabbit anti-Caspr (1:2000), mouse anti-Kv1.2 (1:400), mouse anti-Kv3.1b (1:100), mouse anti-ankyrinG
(1:100), mouse anti-panNav (1:200), rat anti-MBP (1:200), rabbit anti-Olig2 (1:2500). Sections were then
washed and incubated with the appropriate AlexaFluor-conjugated secondary antibodies (1:500)
overnight, washed and mounted on slides with Vectashield mounting medium (Vector Laboratory,

Eurobio Scientific, Les Ulis, France). Image acquisition was performed on a Zeiss (Carl-Zeiss, Iena,
Germany) laser-scanning microscope LSM780 equipped with 40X 1.32 NA oil-immersion objective or
20X objective. Images of AlexaFluor-stained cells were obtained using the 488 nm band of an Argon
laser and the 405 nm, 568 nm and 647 nm bands of a solid-state laser for excitation. Fluorescence
images were collected automatically with an average of two-frame scans at airy 1. Maximum intensity
projection of images was carried out using ImageJ software (NIH). Images are single confocal sections
unless the number of z-steps is indicated.

The density of paranodes immunolabeled for Caspr was estimated in the different layers
(stratum oriens, stratum pyramidale, stratum radiatum and stratum lacunosum-moleculare) of the CA1
hippocampus at P70. The density of single paranodes (1 doublet of paranodes=2 single paranodes)
was quantified in control and 4.1B-deficient mice using maximum intensity projection images (5-z steps
of 2 µm) (ROI of 10^5 µm^2/animal, n=3-4 mice/genotype). The total length of myelinated axons was
measured in the stratum pyramidale (1 bin) and stratum radiatum divided in 5 bins (40x300 µm) on
single confocal sections, n=3-4 mice/genotype. We also measured the total length of Lhx6- or PV-
positive myelinated axon in the stratum radiatum at P35 or P70 (5 ROIs of 10^5 µm^2 from 3
mice/genotype). The length of myelinated SST axons was measured in the stratum radiatum of control
and 4.1B KO mice at P35 (stacks of 2 µm 5-8z steps; 10 ROIs of 10^5 µm^2 from 3 mice/genotype). To
analyze the extent of myelin coverage of individual SST-positive axons across the stratum radiatum,
only axons with a length>100µm were taken into account (n=29-35 axons from 3 mice/genotype). The
distribution of the lengths of internodes was evaluated in the CA1 hippocampus of control and *4.1B$^{-/-}$*
Lhx6-Cre;tdTomato mice at P70. Double-immunostaining for MBP and Caspr and confocal imaging
were performed with analysis on maximum intensity projection images (25 to 50 µm-width z-stacks). We
measured the length of myelin sheaths bordered at each end by a paranode in 4 ROIs of 10^5 µm^2 from
2 mice/genotype (n= 275 in control and n=191 in *4.1B$^{-/-}$* mice). The length of nodal Nav was measured
in the CA1 hippocampus of control *Lhx6-Cre;tdTomato* at P30 and P70. Triple-immunostaining for
Caspr, MBP and panNav was performed and Lhx6-positive axons expressing tdTomato were detected
with the red channel. The distribution of nodal length was analyzed on confocal images (1 µm 2-5z-
steps) from 2 mice/age (P30: n=41, P70: n=64). Next the length of nodal or heminodal Nav channel
clusters and the gap between Nav clusters and paranode were measured in the CA1 hippocampus of
4.1B$^{+/+}$ and *4.1B$^{-/-}$ Lhx6-Cre;tdTomato* mice at P70 (n>25/genotype from 2-3 mice/genotype). The
clustering of juxtaparanodal Kv1 channels was estimated in the CA1 hippocampus of wild-type, *4.1B$^{-/-}$*,
Caspr2$^{-/-}$, and *Contactin2$^{-/-}$* mice at P70. The percentage of single paranodes bordered by Kv1.2
immunostaining in PV-positive axons was quantified on 5-10 µm z-stack confocal images. More than
150 single paranodes were analyzed in each genotype in 7-11 ROIs from 3 mice/genotype). Mutant
mice were compared with their respective controls in double-blind experiments.

Electrophysiological recordings
Mice were deeply anesthetized with Zoletil/Domitor and subsequently decapitated. Brains were quickly
removed from the skull and placed into ice-cold oxygenated (95% O2 and 5% CO2) cutting solution

containing the following (in mM): 132 choline, 2.5 KCl, 1.25 NaH2PO4, 25 NaHCO3, 7 MgCl2, 0.5

CaCl2, and 8 D-glucose (pH: 7.4, 290–300 mOsm/L). 300 µm-thick coronal brain slices were prepared

using a VT1200S microtome (Leica Microsystems, Wetzlar, Germany). Slices were left at room

temperature in oxygenated (95% O2 and 5% CO2) solution of artificial cerebrospinal fluid (aCSF)

containing the following (in mM): 126 NaCl, 3.5 KCl, 1.2 NaH2PO4, 26 NaHCO3, 1.3 MgCl2, 2.0 CaCl2,

and 10 D-glucose, pH 7.4. Then, slices were transferred into a submerged recording chamber and

perfused with oxygenated aCSF at a flow rate of 2 to 3 ml/min. Recordings were performed under visual

guidance using infrared differential interference contrast microscopy (SliceScope Pro 3000M,

Scientifica, Uckfield, UK). tdTomato in neurons was excited by a UV lamp and the fluorescence was

visualized using a CCD camera (Hamamatsu, Japan). Whole-cell patch-clamp recordings were made

using a Multiclamp 700B amplifier (Molecular Devices, CA, USA), filtered at 2 kHz using the built-in

Bessel filter of the amplifier. Data were digitized at 20 kHz with a Digidata 1440A (Molecular Devices)

to a personal computer, and acquired using Clampex 10.1 software (PClamp, Molecular Devices). For

current clamp recording mode patch pipettes were pulled from borosilicate glass tubing with resistances

of 6–8 MΩ and filled with an internal solution containing the following (in mM): 130 KMeSO$_4$, 5 KCl, 10

4-(2-hydroxyethyl)-1-piperazi-methanesulfonic   acid,   2.5   MgATP,   0.3   NaGTP,   0.2

ethyleneglycoltetraacetic acid, 10 phosphocreatine. Access resistance ranged between 15 and 50 MΩ,

and the results were discarded if the access resistance changed by >20%. Neurons recorded in

hippocampal CA1 region were held at a holding potential of −70 mV. Series resistance (between 5–21

MΩ) was monitored throughout each experiment and neurons with a change in series resistance of more

than 25% were excluded from the analysis.

To study intrinsic electrophysiological properties of CA1 inhibitory neurons, we analyzed the

voltage responses to a series of hyperpolarizing and depolarizing square current pulses of 500 ms

duration of amplitudes between −70 pA and 130 pA with 10 pA step intervals from a holding potential of

−70 mV in each cell. The voltage responses of neurons upon the current injections also helped to

distinguish interneurons of a fast-spiking character having no or small 'sag' response to a

hyperpolarizing current pulse from cells with lower spiking rate and a larger 'sag', typical of O-LM cells

(Maccaferri and McBain, 1996) (Table 1 and 2). Input resistance (Rinput) was determined by calculating

the slope of the plot of the membrane potential variation induced by a hyperpolarizing 500 ms step of

current (I: from -70 pA to 0 pA). Firing frequency was studied by injecting 500 ms pulses of depolarizing

current (I: from 10 pA up to 130 pA) into the cell and plotting the spike frequency (f) as a function of the

current intensity (f/I plot). For presumed fast-spiking neurons, a second protocol was used were current

steps varied from +100 pA to +400 pA with 100 pA steps, in order to analyze possible differences in

high-speed discharge. Both analyses were done using Prism 8 software (GraphPad Software, La Jolla,

CA). For the analysis of AP, the first AP evoked by a suprathreshold depolarizing current pulse

(considered as the Rheobase) was selected. The following AP features were calculated using

Minianalysis software: peak amplitude, area, half-width, threshold, AHP area and AHP peak (Table 1

and 2). We further analyzed the shape of the AP rising phase using multiple derivatives of the raw

membrane potential signal using Clampfit software (PClamp, Molecular Devices). In some dV/dt curves

appeared a shouldering. A "double bump" in the d^2V/dt^2 curve and a value equal to or less than zero of

the d^3V/dt^3 trace between the two bumps indicated that the shouldering corresponded to a stationary

inflection in the rising slope of the AP. This indicated that somatic and axonal components of the AP

could be temporally differentiated (Kress et al., 2008; Paterno et al., 2021).

GABA-induced spontaneous inhibitory post synaptic currents (sIPSCs) were recorded in

pyramidal cells in voltage-clamp mode using patch pipettes of 8-10 MΩ filled with an internal solution

containing the following (in mM): 140 CsCl, 1 MgCl2, 10 HEPES, 4 NaCl, 2 Mg-ATP, 0.3 Na-GTP, 0.1

EGTA. The GABA-mediated current was pharmacologically isolated in the presence of AMPA and

NMDA receptor antagonists (10 µM CNQX, 40 µM D-APV respectively). For recording of miniature

IPSCs (mIPSCs) TTX (1µM) was added to the ACSF. CNQX, D-APV and TTX were purchased from

Tocris Bioscience (Bristol, UK).

Behavioral analysis

We examined spatial learning memory through Y-maze and Barnes maze from 3-month mice.

Heterozygous mice (*4.1B+/-; Lhx6-Cre; Ai14*) were crossed to generate the following genotypes *4.1B+/+;*

Lhx6-Cre; Ai14 and *4.1B-/-; Lhx6-Cre; Ai14*. Animals were first handled 2 minutes per day for 4 days.

Spontaneous Y-maze test: Animals (4 males and 4 females of each genotype) were placed in the center

of a Y-maze (Noldus apparatus, Wageningen, The Netherlands) and no external cues were visible from

inside the maze. Spontaneous alternation was recorded during 8 min (d'Isa et al., 2021; Deacon and

Rawlins, 2006). *Barnes maze test:* Animals of each genotype were tested in the maze in random order

(controls: 11 males and 3 females; 4.1B KO: 7 males and 3 females). Visual cues were placed around

the maze, and the ceiling was equipped with high illumination (120 lux) and background sound (85 dB).

The training for the test was performed for 4 days. Each animal adapted to an escape box for 2 min on

the first day, was placed in the center of the maze and explored the maze to find an escape hole, which

is connected to the escape box. Once the animal entered the escape box, it was left there for 30 s, and

the trial ended. Each animal was given 2 daily trials with a 30 min inter-trial interval for the 4 consecutive

days. The animals who did not find the escape box after 120 s were gently guided toward the escape

hole and left there for 30 s. The retention test was performed at 1 day after the training. The escape box

was removed, and the latency time to reach the entry zone, where the escape box was previously

located, was recorded during 120 s for each animal. Using an automatic tracking system, the trajectory

was recorded (Ethovision, Noldus). The path length, average speed, latency to reach or to enter the

escape hole were quantified. Decreases in latency to escape hole with increased training is indicating

that spatial learning occurred. We estimated the percentage of trials using spatial (direct, corrected, with

long-correction) or non-spatial (random or serial) search strategies (Cheng et al., 2019; Illouz et al.,

2016).

Statistical analysis

All values are given as means ± SEM. All statistical tests were performed using Prism 8 (GraphPad

Software). For comparisons of two independent groups, we used Mann–Whitney test or Student's t-test

when values lie in a normal distribution using Shapiro-Wilk test. For multiple group comparisons, we used two-way ANOVA followed by the Sidak's multiple comparison test. To quantify the probability that two sets of samples were drawn from the same probability distribution, we used the Kolmogorov–Smirnov test. To compare the rate of SST cells with stationary inflection to those without, we used Fisher's exact test.

References

Arancibia-Cárcamo, I.L., Ford, M.C., Cossell, L., Ishida, K., Tohyama, K., Attwell, D., 2017. Node of Ranvier length as a potential regulator of myelinated axon conduction speed. eLife 6, e23329. https://doi.org/10.7554/eLife.23329

Artinian, J., Jordan, A., Khlaifia, A., Honoré, E., La Fontaine, A., Racine, A.-S., Laplante, I., Lacaille, J.-C., 2019. Regulation of Hippocampal Memory by mTORC1 in Somatostatin Interneurons. J Neurosci 39, 8439–8456. https://doi.org/10.1523/JNEUROSCI.0728-19.2019

Benamer, N., Vidal, M., Angulo, M.C., 2020. The cerebral cortex is a substrate of multiple interactions between GABAergic interneurons and oligodendrocyte lineage cells. Neurosci Lett 715. https://doi.org/ARTN 134615 10.1016/j.neulet.2019.134615

Benamer, Najate, Vidal, M., Balia, M., Angulo, M.C., 2020. Myelination of parvalbumin interneurons shapes the function of cortical sensory inhibitory circuits. Nat Commun 11, 5151. https://doi.org/10.1038/s41467-020-18984-7

Bonetto, G., Belin, D., Káradóttir, R.T., 2021. Myelin: A gatekeeper of activity-dependent circuit plasticity? Science 374, eaba6905. https://doi.org/10.1126/science.aba6905

Bonetto, G., Hivert, B., Goutebroze, L., Karagogeos, D., Crepel, V., Faivre-Sarrailh, C., 2019. Selective Axonal Expression of the Kv1 Channel Complex in Pre-myelinated GABAergic Hippocampal Neurons. Front Cell Neurosci 13, 222. https://doi.org/10.3389/fncel.2019.00222

Bonnon, C., Bel, C., Goutebroze, L., Maigret, B., Girault, J.A., Faivre-Sarrailh, C., 2007. PGY repeats and N-glycans govern the trafficking of paranodin and its selective association with contactin and neurofascin-155. Mol Biol Cell 18, 229–41.

Brivio, V., Faivre-Sarrailh, C., Peles, E., Sherman, D.L., Brophy, P.J., 2017. Assembly of CNS Nodes of Ranvier in Myelinated Nerves Is Promoted by the Axon Cytoskeleton. Curr Biol 27, 1068–1073. https://doi.org/10.1016/j.cub.2017.01.025

Buttermore, E.D., Dupree, J.L., Cheng, J., An, X., Tessarollo, L., Bhat, M.A., 2011. The cytoskeletal adaptor protein band 4.1B is required for the maintenance of paranodal axoglial septate junctions in myelinated axons. J Neurosci 31, 8013–24.

Cheng, W.H., Martens, K.M., Bashir, A., Cheung, H., Stukas, S., Gibbs, E., Namjoshi, D.R., Button, E.B., Wilkinson, A., Barron, C.J., Cashman, N.R., Cripton, P.A., Wellington, C.L., 2019. CHIMERA repetitive mild traumatic brain injury induces chronic behavioural and neuropathological phenotypes in wild-type and APP/PS1 mice. Alzheimers Res Ther 11, 6. https://doi.org/10.1186/s13195-018-0461-0

Cifuentes-Diaz, C., Chareyre, F., Garcia, M., Devaux, J., Carnaud, M., Levasseur, G., Niwa-Kawakita, M., Harroch, S., Girault, J.A., Giovannini, M., Goutebroze, L., 2011. Protein 4.1B contributes to the organization of peripheral myelinated axons. PLoS One 6, e25043.

d'Isa, R., Comi, G., Leocani, L., 2021. Apparatus design and behavioural testing protocol for the evaluation of spatial working memory in mice through the spontaneous alternation T-maze. Sci Rep 11, 21177. https://doi.org/10.1038/s41598-021-00402-7

Deacon, R.M.J., Rawlins, J.N.P., 2006. T-maze alternation in the rodent. Nat Protoc 1, 7–12.
https://doi.org/10.1038/nprot.2006.2

Denisenko-Nehrbass, N., Oguievetskaia, K., Goutebroze, L., Galvez, T., Yamakawa, H., Ohara, O.,
Carnaud, M., Girault, J.A., 2003. Protein 4.1B associates with both Caspr/paranodin and Caspr2
at paranodes and juxtaparanodes of myelinated fibres. Eur J Neurosci 17, 411–6.

Dubessy, A.L., Mazuir, E., Rappeneau, Q., Ou, S., Abi Ghanem, C., Piquand, K., Aigrot, M.S., Thetiot,
M., Desmazieres, A., Chan, E., Fitzgibbon, M., Fleming, M., Krauss, R., Zalc, B., Ranscht, B.,
Lubetzki, C., Sol-Foulon, N., 2019. Role of a Contactin multi-molecular complex secreted by
oligodendrocytes in nodal protein clustering in the CNS. Glia 67, 2248–2263.
https://doi.org/10.1002/glia.23681

Dubey, M., Pascual-Garcia, M., Helmes, K., Wever, D.D., Hamada, M.S., Kushner, S.A., Kole, M.H.P.,
2022. Myelination synchronizes cortical oscillations by consolidating parvalbumin-mediated
phasic inhibition. Elife 11, e73827. https://doi.org/10.7554/eLife.73827

Einheber, S., Meng, X., Rubin, M., Lam, I., Mohandas, N., An, X., Shrager, P., Kissil, J., Maurel, P.,
Salzer, J.L., 2013. The 4.1B cytoskeletal protein regulates the domain organization and sheath
thickness of myelinated axons. Glia 61, 240–53.

Freeman, S.A., Desmazieres, A., Fricker, D., Lubetzki, C., Sol-Foulon, N., 2016. Mechanisms of sodium
channel clustering and its influence on axonal impulse conduction. Cell Mol Life Sci 73, 723–
35. https://doi.org/10.1007/s00018-015-2081-1

Freeman, S.A., Desmazieres, A., Simonnet, J., Gatta, M., Pfeiffer, F., Aigrot, M.S., Rappeneau, Q.,
Guerreiro, S., Michel, P.P., Yanagawa, Y., Barbin, G., Brophy, P.J., Fricker, D., Lubetzki, C.,
Sol-Foulon, N., 2015. Acceleration of conduction velocity linked to clustering of nodal
components precedes myelination. Proc Natl Acad Sci U S A 112, E321-8.
https://doi.org/10.1073/pnas.1419099112

Geurts, J.J.G., Bö, L., Roosendaal, S.D., Hazes, T., Daniëls, R., Barkhof, F., Witter, M.P., Huitinga, I.,
van der Valk, P., 2007. Extensive hippocampal demyelination in multiple sclerosis. J
Neuropathol Exp Neurol 66, 819–827. https://doi.org/10.1097/nen.0b013e3181461f54

Hamada, M.S., Kole, M.H., 2015. Myelin loss and axonal ion channel adaptations associated with gray
matter neuronal hyperexcitability. J Neurosci 35, 7272–86.
https://doi.org/10.1523/JNEUROSCI.4747-14.2015

Honoré, E., Khlaifia, A., Bosson, A., Lacaille, J.-C., 2021. Hippocampal Somatostatin Interneurons,
Long-Term Synaptic Plasticity and Memory. Front Neural Circuits 15, 687558.
https://doi.org/10.3389/fncir.2021.687558

Horresh, I., Bar, V., Kissil, J.L., Peles, E., 2010. Organization of myelinated axons by Caspr and Caspr2
requires the cytoskeletal adapter protein 4.1B. J Neurosci 30, 2480–9.

Hu, H., Gan, J., Jonas, P., 2014. Interneurons. Fast-spiking, parvalbumin(+) GABAergic interneurons:
from cellular design to microcircuit function. Science 345, 1255263.
https://doi.org/10.1126/science.1255263

Illouz, T., Madar, R., Clague, C., Griffioen, K.J., Louzoun, Y., Okun, E., 2016. Unbiased classification of
spatial strategies in the Barnes maze. Bioinformatics 32, 3314–3320.
https://doi.org/10.1093/bioinformatics/btw376

Ivanovic, A., Horresh, I., Golan, N., Spiegel, I., Sabanay, H., Frechter, S., Ohno, S., Terada, N., Mobius,
W., Rosenbluth, J., Brose, N., Peles, E., 2012. The cytoskeletal adapter protein 4.1G organizes
the internodes in peripheral myelinated nerves. J Cell Biol 196, 337–44.
https://doi.org/10.1083/jcb.201111127

Jeong, N., Singer, A.C., 2022. Learning from inhibition: Functional roles of hippocampal CA1 inhibition
in spatial learning and memory. Curr Opin Neurobiol 76, 102604.
https://doi.org/10.1016/j.conb.2022.102604

Jinno, S., Klausberger, T., Marton, L.F., Dalezios, Y., Roberts, J.D., Fuentealba, P., Bushong, E.A.,
Henze, D., Buzsaki, G., Somogyi, P., 2007. Neuronal diversity in GABAergic long-range
projections from the hippocampus. J Neurosci 27, 8790–804.
https://doi.org/10.1523/JNEUROSCI.1847-07.2007

Kalemaki, K., Konstantoudaki, X., Tivodar, S., Sidiropoulou, K., Karagogeos, D., 2018. Mice With
Decreased Number of Interneurons Exhibit Aberrant Spontaneous and Oscillatory Activity in the
Cortex. Front Neural Circuit 12. https://doi.org/ARTN 96 10.3389/fncir.2018.00096

Kang, Q., Wang, T., Zhang, H., Mohandas, N., An, X., 2009. A Golgi-associated protein 4.1B variant is
required for assimilation of proteins in the membrane. J Cell Sci 122, 1091–1099.
https://doi.org/10.1242/jcs.039644

Kann, O., Papageorgiou, I.E., Draguhn, A., 2014. Highly energized inhibitory interneurons are a central
element for information processing in cortical networks. J Cerebr Blood F Met 34, 1270–1282.
https://doi.org/10.1038/jcbfm.2014.104

Kelley, B.J., Rodriguez, M., 2009. Seizures in patients with multiple sclerosis: epidemiology,
pathophysiology and management. CNS Drugs 23, 805–15. https://doi.org/10.2165/11310900-
000000000-00000

Kole, M.H., Stuart, G.J., 2012. Signal processing in the axon initial segment. Neuron 73, 235–47.

Krasnow, A.M., Attwell, D., 2016. NMDA Receptors: Power Switches for Oligodendrocytes. Neuron 91,
3–5. https://doi.org/10.1016/j.neuron.2016.06.023

Kress, G.J., Dowling, M.J., Meeks, J.P., Mennerick, S., 2008. High threshold, proximal initiation, and
slow conduction velocity of action potentials in dentate granule neuron mossy fibers. J
Neurophysiol 100, 281–291. https://doi.org/10.1152/jn.90295.2008

Kuba, H., 2012. Structural tuning and plasticity of the axon initial segment in auditory neurons. J Physiol
590, 5571–9.

Kunz, L., Maidenbaum, S., Chen, D., Wang, L., Jacobs, J., Axmacher, N., 2019. Mesoscopic Neural
Representations in Spatial Navigation. Trends Cogn Sci 23, 615–630.
https://doi.org/10.1016/j.tics.2019.04.011

Lawrence, J.J., Saraga, F., Churchill, J.F., Statland, J.M., Travis, K.E., Skinner, F.K., McBain, C.J.,
2006. Somatodendritic Kv7/KCNQ/M channels control interspike interval in hippocampal
interneurons. J Neurosci 26, 12325–12338. https://doi.org/10.1523/JNEUROSCI.3521-06.2006

Leão, R.N., Mikulovic, S., Leão, K.E., Munguba, H., Gezelius, H., Enjin, A., Patra, K., Eriksson, A., Loew,
L.M., Tort, A.B.L., Kullander, K., 2012. OLM interneurons differentially modulate CA3 and
entorhinal inputs to hippocampal CA1 neurons. Nat Neurosci 15, 1524–1530.
https://doi.org/10.1038/nn.3235

Liodis, P., Denaxa, M., Grigoriou, M., Akufo-Addo, C., Yanagawa, Y., Pachnis, V., 2007. Lhx6 activity is
required for the normal migration and specification of cortical interneuron subtypes. J Neurosci
27, 3078–3089. https://doi.org/10.1523/JNEUROSCI.3055-06.2007

Maccaferri, G., McBain, C.J., 1996. The hyperpolarization-activated current (Ih) and its contribution to
pacemaker activity in rat CA1 hippocampal stratum oriens-alveus interneurones. J Physiol 497
(Pt 1), 119–130. https://doi.org/10.1113/jphysiol.1996.sp021754

Malavasi, E.L., Ghosh, A., Booth, D.G., Zagnoni, M., Sherman, D.L., Brophy, P.J., 2021. Dynamic early
clusters of nodal proteins contribute to node of Ranvier assembly during myelination of
peripheral neurons. Elife 10, e68089. https://doi.org/10.7554/eLife.68089

Melzer, S., Michael, M., Caputi, A., Eliava, M., Fuchs, E.C., Whittington, M.A., Monyer, H., 2012. Long-
range-projecting GABAergic neurons modulate inhibition in hippocampus and entorhinal cortex.
Science 335, 1506–10. https://doi.org/10.1126/science.1217139

Micheva, K.D., Kiraly, M., Perez, M.M., Madison, D.V., 2021. Conduction Velocity Along the Local Axons
of Parvalbumin Interneurons Correlates With the Degree of Axonal Myelination. Cereb Cortex
31, 3374–3392. https://doi.org/10.1093/cercor/bhab018

Micheva, K.D., Wolman, D., Mensh, B.D., Pax, E., Buchanan, J., Smith, S.J., Bock, D.D., 2016. A large
fraction of neocortical myelin ensheathes axons of local inhibitory neurons. Elife 5.
https://doi.org/10.7554/eLife.15784

Nicholas, R., Magliozzi, R., Campbell, G., Mahad, D., Reynolds, R., 2016. Temporal lobe cortical
pathology and inhibitory GABA interneuron cell loss are associated with seizures in multiple
sclerosis. Mult Scler 22, 25–35. https://doi.org/10.1177/1352458515579445

Paterno, R., Marafiga, J.R., Ramsay, H., Li, T., Salvati, K.A., Baraban, S.C., 2021. Hippocampal gamma
and sharp-wave ripple oscillations are altered in a Cntnap2 mouse model of autism spectrum
disorder. Cell Rep 37, 109970. https://doi.org/10.1016/j.celrep.2021.109970

Penagarikano, O., Abrahams, B.S., Herman, E.I., Winden, K.D., Gdalyahu, A., Dong, H., Sonnenblick,
L.I., Gruver, R., Almajano, J., Bragin, A., Golshani, P., Trachtenberg, J.T., Peles, E., Geschwind,
D.H., 2011. Absence of CNTNAP2 leads to epilepsy, neuronal migration abnormalities, and core
autism-related deficits. Cell 147, 235–46.

Philips, T., Rothstein, J.D., 2017. Oligodendroglia: metabolic supporters of neurons. J Clin Invest 127,
3271–3280. https://doi.org/10.1172/JCI90610

Pinatel, D., Faivre-Sarrailh, C., 2021. Assembly and Function of the Juxtaparanodal Kv1 Complex in
Health and Disease. Life-Basel 11. https://doi.org/ARTN 8 10.3390/life11010008

Poliak, S., Salomon, D., Elhanany, H., Sabanay, H., Kiernan, B., Pevny, L., Stewart, C.L., Xu, X., Chiu,
S.Y., Shrager, P., Furley, A.J., Peles, E., 2003. Juxtaparanodal clustering of Shaker-like K+
channels in myelinated axons depends on Caspr2 and TAG-1. J Cell Biol 162, 1149–60.

Rama, S., Zbili, M., Fékété, A., Tapia, M., Benitez, M.J., Boumedine, N., Garrido, J.J., Debanne, D.,
2017. The role of axonal Kv1 channels in CA3 pyramidal cell excitability. Sci Rep 7, 315.
https://doi.org/10.1038/s41598-017-00388-1

Savvaki, M., Panagiotaropoulos, T., Stamatakis, A., Sargiannidou, I., Karatzioula, P., Watanabe, K.,
Stylianopoulou, F., Karagogeos, D., Kleopa, K.A., 2008. Impairment of learning and memory in
TAG-1 deficient mice associated with shorter CNS internodes and disrupted juxtaparanodes.
Mol Cell Neurosci 39, 478–90. https://doi.org/10.1016/j.mcn.2008.07.025

Somogyi, P., Klausberger, T., 2005. Defined types of cortical interneurone structure space and spike
timing in the hippocampus. J Physiol 562, 9–26. https://doi.org/10.1113/jphysiol.2004.078915

Stedehouder, J., Brizee, D., Shpak, G., Kushner, S.A., 2018. Activity-Dependent Myelination of
Parvalbumin Interneurons Mediated by Axonal Morphological Plasticity. J Neurosci 38, 3631–
3642. https://doi.org/10.1523/JNEUROSCI.0074-18.2018

Stedehouder, J., Brizee, D., Slotman, J.A., Pascual-Garcia, M., Leyrer, M.L., Bouwen, B.L., Dirven,
C.M., Gao, Z., Berson, D.M., Houtsmuller, A.B., Kushner, S.A., 2019. Local axonal morphology
guides the topography of interneuron myelination in mouse and human neocortex. Elife 8,
e48615. https://doi.org/10.7554/eLife.48615

Stedehouder, J., Couey, J.J., Brizee, D., Hosseini, B., Slotman, J.A., Dirven, C.M.F., Shpak, G.,
Houtsmuller, A.B., Kushner, S.A., 2017. Fast-spiking Parvalbumin Interneurons are Frequently
Myelinated in the Cerebral Cortex of Mice and Humans. Cereb Cortex 27, 5001–5013.
https://doi.org/10.1093/cercor/bhx203

Thetiot, M., Freeman, S.A., Roux, T., Dubessy, A.L., Aigrot, M.S., Rappeneau, Q., Lejeune, F.X.,
Tailleur, J., Sol-Foulon, N., Lubetzki, C., Desmazieres, A., 2020. An alternative mechanism of
early nodal clustering and myelination onset in GABAergic neurons of the central nervous
system. Glia 68, 1891–1909. https://doi.org/10.1002/glia.23812

Torii, T., Ogawa, Y., Liu, C.H., Ho, T.S., Hamdan, H., Wang, C.C., Oses-Prieto, J.A., Burlingame, A.L.,
Rasband, M.N., 2020. NuMA1 promotes axon initial segment assembly through inhibition of
endocytosis. J Cell Biol 219. https://doi.org/10.1083/jcb.201907048

Traka, M., Goutebroze, L., Denisenko, N., Bessa, M., Nifli, A., Havaki, S., Iwakura, Y., Fukamauchi, F.,
Watanabe, K., Soliven, B., Girault, J.A., Karagogeos, D., 2003. Association of TAG-1 with
Caspr2 is essential for the molecular organization of juxtaparanodal regions of myelinated
fibers. J Cell Biol 162, 1161–72.

Tricoire, L., Pelkey, K.A., Erkkila, B.E., Jeffries, B.W., Yuan, X., McBain, C.J., 2011. A blueprint for the
spatiotemporal origins of mouse hippocampal interneuron diversity. J Neurosci 31, 10948–
10970. https://doi.org/10.1523/JNEUROSCI.0323-11.2011

Udakis, M., Pedrosa, V., Chamberlain, S.E.L., Clopath, C., Mellor, J.R., 2020. Interneuron-specific
plasticity at parvalbumin and somatostatin inhibitory synapses onto CA1 pyramidal neurons
shapes hippocampal output. Nat Commun 11, 4395. https://doi.org/10.1038/s41467-020-
18074-8

Vagionitis, S., Auer, F., Xiao, Y., Almeida, R.G., Lyons, D.A., Czopka, T., 2022. Clusters of neuronal
neurofascin prefigure the position of a subset of nodes of Ranvier along individual central
nervous system axons in vivo. Cell Rep 38, 110366.
https://doi.org/10.1016/j.celrep.2022.110366

Wang, Z., Zhang, J., Ye, M., Zhu, M., Zhang, B., Roy, M., Liu, J., An, X., 2014. Tumor suppressor role
of protein 4.1B/DAL-1. Cell Mol Life Sci 71, 4815–4830. https://doi.org/10.1007/s00018-014-
1707-z

White, D.J., Congedo, M., Ciorciari, J., Silberstein, R.B., 2012. Brain oscillatory activity during spatial
navigation: theta and gamma activity link medial temporal and parietal regions. J Cogn Neurosci
24, 686–697. https://doi.org/10.1162/jocn_a_00098

Yamada, R., Kuba, H., 2016. Structural and Functional Plasticity at the Axon Initial Segment. Front Cell
Neurosci 10, 250. https://doi.org/10.3389/fncel.2016.00250

Yang, S.M., Michel, K., Jokhi, V., Nedivi, E., Arlotta, P., 2020. Neuron class-specific responses govern
adaptive myelin remodeling in the neocortex. Science 370.
https://doi.org/10.1126/science.abd2109

Zhang, C., Susuki, K., Zollinger, D.R., Dupree, J.L., Rasband, M.N., 2013. Membrane domain
organization of myelinated axons requires betaII spectrin. J Cell Biol 203, 437–43.
https://doi.org/10.1083/jcb.201308116

Zonouzi, M., Berger, D., Jokhi, V., Kedaigle, A., Lichtman, J., Arlotta, P., 2019. Individual
Oligodendrocytes Show Bias for Inhibitory Axons in the Neocortex. Cell Rep 27, 2799-2808 e3.
https://doi.org/10.1016/j.celrep.2019.05.018

Zonta, B., Tait, S., Melrose, S., Anderson, H., Harroch, S., Higginson, J., Sherman, D.L., Brophy, P.J.,
2008. Glial and neuronal isoforms of Neurofascin have distinct roles in the assembly of nodes
of Ranvier in the central nervous system. J Cell Biol 181, 1169–77.
https://doi.org/10.1083/jcb.200712154

Zoupi, L., Booker, S.A., Eigel, D., Werner, C., Kind, P.C., Spires-Jones, T.L., Ben Newland, Williams,
A.C., 2021. Selective vulnerability of inhibitory networks in multiple sclerosis. Acta Neuropathol
141, 415–429. https://doi.org/10.1007/s00401-020-02258-z