Photosynthetic and water transport strategies of plants along a tropical forest aridity gradient:

a test of optimality theory

Corresponding author:

Huanyuan Zhang-Zheng, (0000-0003-4801-8771) (huanyuan.zhangzheng@gmail.com)

Author list:

Huanyuan Zhang-Zheng1, (0000-0003-4801-8771)

Yadvinder Malhi1, (0000-0002-3503-4783)

Agne Gvozdevaite1,

Theresa Peprah4,

Mickey Boackye3,

Kasia Ziemińska2, (0000-0001-7750-4758)

Stephen Adu-Bredu4,

Jesús Aguirre-Gutiérrez 1,(0000-0001-9190-3229)

David Sandoval 5,

Colin Prentice 5, (0000-0002-1296-6764)

Imma Oliveras1,2, (0000-0001-5345-2236)

1 Environmental Change Institute, School of Geography and the Environment, University of
Oxford, Oxford, United Kingdom,

2 AMAP (Botanique et Modelisation de l'Architecture des Plantes et des Végétations), CIRAD,

CNRS, INRA, IRD,Université de Montpellier, Montpellier, France

3 Department of Environmental Science, Policy, and Management, University of California,

Berkeley, CA, 94720, USA

4 Forestry Research Institute of Ghana, Council for Scientific and Industrial Research, Kumasi,

Ghana

5 Georgina Mace Centre for the Living Planet, Department of Life Sciences, Imperial College

London, Silwood Park Campus, Buckhurst Road, Ascot, SL5 7PY, UK

Word Count

Total word count for the main body of the text: 5426

Summary: 200

Introduction: 707

Materials and Methods: 2491

Results: 1029

Discussion: 999 (18.4% of total word count)

Number of figures:2 (all in colour)

Number of table: 1 (all in colour)

Supporting information is provided in another document.

Keywords

42   Optimality; photosynthesis; aridity; plant functional traits; plant hydraulics; xylem hydraulics

Summary

(1) The research conducted, including the rationale

The direct effect of aridity on photosynthetic and water-transport strategies is not easy to discern
in global analyses because of large-scale correlations between precipitation and temperature. We
analyze tree traits collected along an aridity gradient in Ghana, West Africa, that shows little
temperature variation, in an attempt to disentangle thermal and hydraulic influences on plant traits.

(2) Methods

Predictions derived from optimality theory of the variation of key plant traits along the gradient
are tested with field measurements.

(3) Results

Most photosynthetic traits show trends consistent with optimality-theory predictions, including
higher photosynthetic rates in the drier sites, and an association of higher photosynthetic rates with
greater respiration rates and greater water transport. Leaf economic and hydraulic traits show less
consistency with theory or global-scale pattern, especially predictions based on xylem efficiency-
sfatety trafeoff. Nonetheless, the link between photosynthesis and water transport still holds:
species (predominantly deciduous species found in drier sites) with both higher sapwood-to-leaf
area ratio (AS/AL) and potential hydraulic conductivity (Kp), implying higher transpiration, tend
to have both higher photosynthetic capacity, and lower leaf-internal CO_2.

(4) Conclusions

These results indicate that aridity is an independent driver of spatial patterns of photosynthetic
traits, while plants show a diversity of water-transport strategies along the aridity gradient.

Plain language summary: Along an aridity gradient in Ghana, West-Africa, we used optimality theory to explain why higher photosynthetic rates should be found at drier places and how plants arrange water transportation to support quicker photosynthesis at the drier site. We also reported surprising data-theory inconsistency for some hydraulic traits along the aridity gradient where further research is needed.

Introduction

Three key photosynthetic processes are frequently considered when seeking to understand plants photosynthesis strategies: light availability and electron transport; aridity and water transport; and CO_2 concentration and carboxylation (Farquhar et al., 2001). Plants capacities in these processes vary considerably along environmental gradients (Wang *et al.*, 2017a; Bahar *et al.*, 2017; Yang *et al.*, 2019; Oliveras *et al.*, 2020). Recently, many efforts have been made to propose universal rules to explain worldwide plant photosynthetic strategies, frequently cited as 'optimality theories', which could serve as a basic theoretical framework for vegetation carbon modelling and enable quantitative predictions of key photosynthetic traits (Franklin *et al.*, 2020; Harrison *et al.*, 2021).

One of the main challenges confronting these universal theories is to explain the 'pure' effect of aridity on photosynthesis (Rogers *et al.*, 2017). Such challenges become particularly pressing in the context of climate as greater atmospheric dryness (water vapour deficit, VPD) is predicted for most places (Neelin *et al.*, 2006; Grossiord *et al.*, 2020a; Bauman *et al.*, 2022), which may strongly influence photosynthesis and hence the carbon cycle (Canadell *et al.*, 2021). However, the theoretical expectation for the impact of aridity on plant traits based on optimality theories has not been summarized and tested. Most current earth systems models predict a negative relationship between photosynthesis (denoted by CO_2 assimilation rate per leaf area, A_{area}) and VPD simply due to the closing of stomata without incorporating the dynamics of photosynthetic capacity (denoted by electron-transport capacity, J_{max25} and Rubisco carboxylation capacity standardized to 25 degree celsius, V_{cmax25}) (Wang *et al.*, 2017a; Green *et al.*, 2020). On the contrary, a study focusing on Amazonia argued that higher VPD increases photosynthetic capacity which

counteracts the reduced conductivity, leading to higher photosynthesis under drier climates

(Restrepo-Coupe *et al.*, 2013; Green *et al.*, 2020). However, a global study found the above pattern

only exists in wet ecosystems and higher VPD reduces photosynthesis overall (Fu *et al.*, 2022)

although globally higher V_{cmax25} was indeed found for plants grown in drier sites (Peng *et al.*, 2021).

Under experimental conditions, CO_2 assimilation was found to be lower under high VPD (Long &

Woolhouse, 1978; Dai *et al.*, 1992; Cunningham, 2005).

With such mixed evidence in the literature, the stand-alone effect of aridity on photosynthesis still

remains unclear. There are two particular challenges. First, on a large spatial scale, aridity can be

confounded with temperature (especially when VPD is used as a metric of aridity). Temperature

is a stronger driver of photosynthesis than aridity (Smith *et al.*, 2019; Peng *et al.*, 2021), but few

studies try to disentangle aridity from temperature (Grossiord *et al.*, 2020a). Second, the effect of

VPD can be confounded with soil water availability. Optimality theory predicted higher Vcmax

and A_{area} under higher VPD (Smith *et al.*, 2019) but it is unclear how plants in drier environments

arrange water transportation to support higher $A_{area.}$ A comprehensive theoretical framework is

lacking to incorporate the effect of VPD on all leaf-level photosynthesis processes (light, water

and CO_2) with full consideration of water delivery to leaves (Mencuccini *et al.*, 2019a).

Here, we examine a dataset of detailed traits measurements along an aridity gradient in West

African forests to disentangle the effect of aridity on photosynthesis from temperature and to

explain the effect with optimality theory. The key questions we address are: (1) do drier

environments have higher photosynthesis rates and how do aridity and photosynthesis interact? (2)

If photosynthetic rates are higher in arid environments, as predicted by optimality theory, how do

plants arrange greater water transportation, given greater water stress in drier places? To answer

these questions, we adopted a theory-data comparison approach where we first review the

expectation of recent 'universal' theories and deduced 16 testable predictions (some of which have

previously been tested but with confounding results). We then examined the consistency between

each prediction and field measurement along the aridity gradient (Table 1). Consistency would

give field-observed patterns a mechanistic explanation and reinforce the stand-alone impact of

aridity on the corresponding trait, while inconsistency would imply weakness of the theory and

possible confounding interactions between aridity and other climate variables. Before closing the

paper, we summarize the consistency and inconsistency with an integrated theoretical framework
to address and explore the 'pure' effect of aridity on photosynthesis.

Description of theory and hypotheses development

'Optimality theory' was developed recently with the assumption that plants can optimize
photosynthesis and minimize maintenance costs according to their living environments by
optimizing investment in the above processes, which provides a universal explanation of the
variation of photosynthetic strategies under different growing environments (Prentice *et al.*, 2014;
Wang *et al.*, 2017b; Mencuccini *et al.*, 2019a; Stocker *et al.*, 2020; Xu *et al.*, 2021). Although the
above-cited studies have tested the theories on global scales and along elevation gradients,
discussion and validation of these theories along other abiotic gradients, such as aridity gradients,
are still lacking. Therefore, we first review the implication of such theories on plants
photosynthetic strategies along aridity gradients.

In drier sites, higher VPD increases potential transpiration per unit leaf area. As predicted by the
'least-cost hypothesis' (Wright *et al.*, 2001, 2003; Medlyn *et al.*, 2011; Prentice *et al.*, 2014), plants
can compensate for high water costs in dry climates by keeping stomata relatively closed. Thus, in
drier sites, plants are expected to have lower leaf internal CO_2 concentration (c_i), lower internal-
to-external CO_2 ratio (c_i/c_a) and lower stomatal conductance (g_s). The 'coordination hypothesis'
(Beerling & Quick, 1995; Maire *et al.*, 2012; Walker *et al.*, 2014) assumes equilibrium between
Rubisco-limited photosynthesis rates (A_C) (depending on V_{cmax25} and c_i) and electron transport-
limited photosynthesis rates (A_J) (depending on J_{max25} and leaf absorbed PPFD) (see the
quantitative expression in (Wang *et al.*, 2017b; Smith *et al.*, 2019; Stocker *et al.*, 2020)). To
maintain such an equilibrium, plants in drier sites are expected to have larger V_{cmax25} to compensate
for the lower c_i. Otherwise, A_C would be lower than A_J resulting in unused capacity for electron
transport (J_{max25}). To sum up, lower c_i but higher V_{cmax25} is expected toward drier sites if J_{max25}
stays constant (in which case A_J would be slightly lower due to smaller c_i).

In reality, toward drier sites, it is common to see higher leaf absorbed photosynthetic photon flux
density (I_{abs}) because of less cloud cover and more open canopies. Considering an additional
optimality criterion that J_{max25} is acclimated to I_{abs} (Smith *et al.*, 2019), supported by multiple

experiments (Björkman, 1981; Ögren, 1993), we would expect higher J_{max25} in drier sites, which

further encourages higher V_{cmax25} (see above paragraph). Higher J_{max25} and V_{cmax25} would give rise

to higher A_C and A_J. All the above would lead to higher leaf photosynthetic protein cost, hence

higher leaf dark respiration (Rd), and higher transpiration stream maintenance cost, hence higher

stem respiration per leaf area (Rs) (Prentice *et al.*, 2014). Note that R_s is stem respiration per leaf

area (R_{stem_leaf} hereafter), different from the commonly reported stem respiration per stem area

(R_{stem_stem}). Some of the above predictions have been seen on global scale; for example, higher R_d

has been found in drier sites (Wright *et al.*, 2001; Atkin *et al.*, 2015) and higher assimilation rate

has been reported from drier sites (Maire *et al.*, 2015).

It is worth noting that V_{cmax25}, g_s and c_i here are discussed as an overall value for a forest stand,

disregarding diurnal variation and intraspecific variation (Stangl *et al.*, 2019; Han *et al.*, 2022). At

the species or individual scales, there is a positive correlation between A_{sat} (light-saturated

assimilation rate at 400 ppm), V_{cmax25}, g_s and c_i (Wright *et al.*, 2003; Prentice *et al.*, 2014), instead

of the opposite trend of V_{cmax25} and c_i/c_a as discussed above regarding spatial variation only.

Photosynthesis strategies predicted by the optimality theory above can be linked with xylem water

transportation strategies via stomatal behaviour, as given by Fick's law,

$g_s = A_{area} / (c_a - c_i)$ (1)

Where g_s is stomatal conductance (umol CO_2 m-2 s-1), A_{area} is CO_2 assimilation rate per leaf area

(umol CO2 m2 s-1), and leaf internal (c_i, ppm) and external (c_a, ppm) CO_2 concentration

We focus on daytime conditions that produce maximum rates of transpiration and photosynthesis,

when water loss through stomata must equal water transport through xylem (Brodribb *et al.*, 2002;

Xu *et al.*, 2021):

$E = 1.6 \, g_s \, VPD / P_{atm} = K_s \, \Delta\Psi_{max} \, A_S/A_L \, / \, h$ (2)

Where Patm is atmospheric pressure (Mpa), KS is the sapwood-specific hydraulic conductivity

(mol m–1s–1 MPa–1); AS/AL is the ratio of sapwood to leaf area (m2m–2) $\Delta\Psi_{max}$ is the

maximum decrease in water potential from soil to leaves (MPa) h is the transpiration stream

pathlength (m), roughly equivalent to plant height, and E is the transpiration rate (mol m–2s–1),

Combining the above two equations we obtain a link between water transportation and

photosynthesis:

$K_s \ \Delta\Psi max \ A_S/A_L \ / \ h = 1.6 \ VPD \ A_{area} \ / \ (c_a - c_i) \ /P_{atm} = E$ (3)

Note that here we focus on forest-stand scale as an average across time. Relationships could be

very different at other scales (Mencuccini *et al.*, 2019a).

In drier sites with higher VPD, despite less open stomata (less g_s and less c_i), there should

inevitably be a larger E (Granier *et al.*, 1996) and more negative $\Delta\Psi max$ (Gleason *et al.*, 2013);

therefore smaller maximum tree height, and more negative turgor loss point (TLP, Mpa) in drier

sites to increase hydraulic resistance (note that TLP must be more negative than $\Delta\Psi max$) (Ryan &

Yoder, 1997; Bartlett *et al.*, 2012). Equation 3 implies that in drier sites with high VPD, plants

require a larger A_S/A_L and/or larger K_s in order to support the same amount of photosynthesis

with enhanced transpiration. Following the xylem safety–efficiency trade-off (Manzoni *et al.*,

2013; Gleason *et al.*, 2016; Bittencourt *et al.*, 2016; Grossiord *et al.*, 2020b) (although arguments

against this trade-off exist, here we present testable hypotheses expected by the trade-off), plants

at drier sites would be expected to have lower hydraulic efficiency and lower hydraulic

conductivity. Lower hydraulic conductivity is often associated with smaller vessel diameter, higher

vessel density and higher wood density (Poorter *et al.*, 2010; Schuldt *et al.*, 2013; Hoeber *et al.*,

2014). Such patterns have been observed along an Australian aridity gradient (Gleason *et al.*, 2013;

Pfautsch *et al.*, 2016), but no effect of aridity on vessel diameter was reported (Olson & Rosell,

2013; Olson *et al.*, 2014). Plants in drier sites should have increased hydraulic safety - more

negative TLP and more negative P50 (Hacke *et al.*, 2001; Martínez-Vilalta *et al.*, 2009; Gleason

et al., 2013; Togashi *et al.*, 2015; Liu *et al.*, 2019; López *et al.*, 2021). In short, toward drier sites,

we would expect to see, higher A_S/A_L and more negative TLP. The safety-efficiency trade-off

implies lower K_s, smaller vessel diameter, higher vessel density and higher wood density.

The trade-off between Ks and AS/AL is also embedded in the variance of traits in equation 3. Ks
and AS/AL could vary by two orders of magnitude (100-fold variation) (Mencuccini *et al.*, 2019b)
on a global scale, while ci/ca and A_{area} vary much less (ci/ca: 2 fold; A_{area}: 10 fold) (Wright *et al.*,
2004; Wang *et al.*, 2017b). This leads to a trade-off between Ks and AS/AL (i.e. KS x AS/AL
should vary less than either of them). However, given that there are also variations of ci/ca, A_{area},
h and ΔΨmax, it is possible that different species range along a spectrum from high A_{area} and E to
low A_{area} and E while always satisfying equation 3 (Prentice *et al.*, 2014).

In short, the above review leads to an integrated hypothesis that plants in drier (normally also
brighter) sites tend to develop a photosynthesis strategy with less stomatal conductance and hence
lower ci, stronger photosynthetic capacities (larger V_{cmax25}, J_{max25} and A_{area}) with more
maintenance cost (higher Rd and Rs), higher maximum carbon assimilation rate and larger
maximum evapotranspiration which the water transport system would adjust to with higher AS/AL,
lower Ks, lower tree height and more negative TLP. Information on leaf economic traits is
provided in Appendix 3. We break the above prediction down into 16 testable hypotheses (Table
1) and test each of them along a forest aridity gradient.

Materials and Methods

Study sites - the aridity gradient

This study presents and analyses physiological traits data collected from seven one-hectare forest
and savanna plots distributed along a wet to dry rainfall gradient across three sites, Ankasa (ANK,
moist rainforest), Bobiri (BOB, semi-deciduous forest) and Kogyae (KOG, dry forest and mesic
savanna), in Ghana, West Africa (Figure S1 S2) (Moore *et al.*, 2018; Oliveras *et al.*, 2020), as part
of the Global Ecosystem Monitoring (GEM) network (Malhi *et al.*, 2021). These sites share very
similar mean annual temperature but span a steep gradient of aridity (Figure 1), which provided a
"natural laboratory" to disentangle the hydraulic aspect of plant traits variation from temperature.

Although one-hectare plots (e.g. BOB-02) within the same site (e.g. BOB) share very similar air
temperature and precipitation, they can differ in terms of soil moisture supply due to small-scale
variations in soil properties and topography (Table S1). Along the aridity gradient, there are also
variations in soil and vegetation type, with vegetation seasonality and deciduousness increasing
considerably towards drier sites (Moore *et al.*, 2018). More information about the soil properties
and climate of all three sites can be found in (Domingues *et al.*, 2010; Chiti *et al.*, 2010; Moore *et*
al., 2018). Moreover, the swampy rainforest (ANK03) is partly inundated during the wet season
unlike ANK01, which is located on a hill and never inundated. From KOG02 (dry forest), KOG04
to KOG05 (savanna), forest plots become more deciduous with a smaller number of trees. Within
any site, there are many common species between plots but species composition (e.g., top five
abundant species) could still be very different. There is almost no common abundant species
between the three sites (ANK, BOB and KOG).

Aridity index and soil moisture stress

At site scales, we calculated an aridity index as the ratio of annual potential evapotranspiration
(PET) to mean annual precipitation (MAP). To understand soil moisture stress experienced by
plants, we reported not only measured surface (12 cm depth) soil volumetric water content, but
also hydraulic simulations on plot scales with SPLASH v2.0 (Sandoval & Prentice, 2020). This
model requires three sets of input data: (1) field observed climate data at site scale during 2011-
2016 (2) soil properties measured following the RAINFOR protocols (Quesada *et al.*, 2010); (3)
terrain data: root zone was assumed 2m, while upslope drainage area, slope inclination and
orientation were extracted from (Yamazaki *et al.*, 2019). We considered two modelled indices: the
relative soil moisture saturation (Θ) was defined as the volumetric water content (θ) normalized
by the volumetric water content at saturation (θ_{SAT}); a vegetation water stress index (α) was
estimated as the ratio of annual actual evapotranspiration (AET) to PET.

Functional trait data measurements

Leaf traits field campaigns to measure leaf traits were conducted using a standardized protocol between October 2015 and September 2016 in all plots (Oliveras *et al.*, 2020), covering both dry and wet seasons (see Appendix 1 for sampling protocol). To ensure consistent comparison along the aridity gradient, only sun leaves were sampled. Sampled leaves were chosen from individuals that correspond to the most dominant species in each plot. To determine the target species, species contributing to up to 80% of the total basal area of the plot were ranked by basal area. As equation 3 focuses on daytime conditions with maximum transpiration, we use Kp (potential specific hydraulic conductivity) as a proxy of Ks and plant stature (H_{max}) as a proxy of path length (h). Kp was calculated from vessel density and vessel diameter following (Poorter *et al.*, 2010).

Following (Prentice *et al.*, 2014), we calculated stem respiration per leaf area (Rs_leaf) instead of the commonly presented Rs_stem, as a 'maintenance cost of photosynthesis' (See Appendix 1). To our knowledge, Rs_leaf has not previously been presented with in-situ data in the literature. Here we argue the importance to understand stem respiration from per leaf area perspectives because (1) higher stem respiration per stem area has been found in wetter sites (Yang *et al.*, 2016), contradictory to theoretical expectation (Prentice *et al.*, 2014) (2) consistency with photosynthetic traits which were commonly reported per leaf area.

All data reported in this study were field-measured except for wood density, which was obtained from a global species database (Zanne *et al.*, 2009). Global scale sapwood-to-leaf-area ratio in Figure 3 is sourced from (Mencuccini *et al.*, 2019b). Global scale vessel diameter in Figure S4 is sourced from (Choat *et al.*, 2012).

Statistical analysis

Hypotheses 1-14 (Table 1) were tested by significant differences between wet and dry sites. Principal component analysis (PCA) and standardized major axis regression are used to understand the relationship between Ks, As/AL and photosynthesis traits (Hypothesis 15-16).

We performed a plot-to-plot comparison in answering Hypotheses 1 to 14 as follows: (1) The distribution was visually inspected with histograms and transformations were applied if necessary.

(2) Outliers were checked with the R package *outliers::scores*, 'iqr' method with threshold 1.5.

This removed leaves/trees with extreme values but made the community-weighted mean a better

representation of the whole plot. (3) Community-weighted means were calculated based on the

basal area of each species. Standard error was calculated with the same weights (Madansky &

Alexander, 2017). (4) Significance of differences in plot-to-plot community-weighted means were

then tested with *lm*(), *glht*(), and *cld*(). (See Rscript *community_weighted_mean_Ghana_log_YS.R*

for a full description). In testing Hypotheses 1-14, a hypothesis was accepted if KOG (dry region)

was significantly different to ANK (wet region) while BOB (middle aridity) sat in between (Figure

1). (5) Variance partitioning was done with *vegan::varpart()*, following the 'RDA' method with

the expression: *varpart* (Trait, ~ Plot, ~ Species, data = Trait). Variance partitioning reveals

whether the change of traits along the aridity gradient was driven by intraspecific or interspecific

variation. Variance partitioning is also used to diagnose whether the intra-specific variation or

measurement errors are overwhelming. To double-check the impact of intraspecific variation, we

recalculated a community-weighted mean by assuming that the same species share the same value

of trait (i.e. remove intraspecific variation) and extrapolated traits value to forest plots without trait

measurements (Appendix 5)

For hypothesis 16, we applied Principal Component Analysis (PCA) with *FactoMineR::PCA*(),

(See Rscript ~/link_hydraulic_photosynthesis/PCA.R). Asat, Kp, AS/AL and V_{cmax25} were log10

transformed. We avoid standardization by setting 'scale.unit' as False in function PCA() so that

the variance of a trait was reflected by the length of an arrow in Figure 2. Trait-trait correlations

(bivariate plot) were calculated (Rscript *Compare_with_Others.R*). For hypothesis 15, the slopes

and significance of correlation were calculated using Standard Major Axis Regression (*function*

smatr::sma()), which is preferable to ordinary least squares regression in summarizing the

relationship between two plant traits (Wright *et al.*, 2005; Warton *et al.*, 2012) as it considers

uncertainties of both axes. All analyses were done at the species level (i.e. each point in Figure 2

represents a species) to compare with other studies and join among datasets. Hypothesis 15 was

also tested at the global scale because Ks was reported to negatively correlate with AS/AL but

there is no report on the global correlation between Kp and AS/AL (Appendix 4).

Results

Aridity gradient

The values of the aridity index (PET/MAP) (site scale) revealed a clear aridity gradient from ANK (moist rainforest site) to BOB (mid) and KOG (dry) (Appendix 2). The same order could be arrived at with VPD or maximum cumulative water deficit (MCWD). The standard deviation of monthly VPD also suggested that the seasonality was weaker in ANK and increased towards KOG.

On the other hand, the simulations of relative soil moisture saturation (Θ) and vegetation water stress index (α) (plot scale) showed that BOB was the least soil moisture stressed site, followed by ANK and KOG. BOB-02 had the highest values in these two metrics. The model reported the highest runoff at ANK-03, capturing to some degree the seasonal flooding, as also observed in the field. The different patterns of Θ (or α) to aridity index along the aridity gradient were caused by the soil characteristics which in turn define water holding capacity (WHC) and hydraulic conductivity; for example, the plots in BOB were atmospherically drier (higher PET/MAP) than in ANK but they could hold more water (higher Θ). Especially in BOB-02, the infiltration rate was strongly reduced (60 mm/hr, less than half of ANK plots), and hence water can stay more time in the root zone while percolating. This acts as a buffer against the evaporative demand, maintaining water availability during dry months. The hydrological modelling outputs also matched with field observation of plot vegetation characteristics (see Methods).

For presentation (Figure 1), we rank sites by aridity index and then plots within sites by soil moisture stress.

The effect of aridity on traits

From a photosynthesis perspective, along the aridity gradient, we saw consistency between theoretical prediction and field measurements (Table 1) for all traits: ci/ca (0.71 to 0.85), V_{cmax25}

(21.58 to 46.48 umol CO_2 m2 s-1), J_{max25} (38.48 to 91.44 umol CO_2 m2 s-1), Rd (-1.66 to -2.41

umol CO_2 m2 s-1), Rs (3.76 to 11.70 umol CO_2 m2 s-1), Asat (4.56 to 7.72 umol CO_2 m2 s-1) and

Amax (15.88 to 22.86 umol CO_2 m2 s-1). However, we saw less consistency between theory and

field measurements regarding leaf economic traits (Appendix 3). Nmass was expected to increase

from wet to dry but this is not backed by our field measurements. LMA was slightly higher in the

dry sites than in the wet sites as theoretically expected, but not a gradual increase. Pmass increased

gradually from wet to dry plots, (from 0.94 to 1.67 g kg-1) along the gradient in accord with global

observation. Although the link among soil nitrogen, leaf nitrogen and photosynthesis was

frequently made (Walker *et al.*, 2014; Gvozdevaite *et al.*, 2018), we found that such a link is rather

ambiguous along the aridity gradient on site scale.

From a water transpiration perspective, the hypotheses were consistent with field measurements

for leaf traits. AS/AL was higher in drier sites (359.62 to 901.66 cm2 m-2) and TLP was more

negative in drier sites (-1.33 to -1.63 Mpa). However, no consistency was found between

theoretical expectations and field measurements for any xylem-related traits. Along the aridity

gradient, there was an increasing trend of field Kp toward drier sites (from 28.62 to 59.29 kg m-1

Mpa-1 s-1), against the xylem safety-efficiency trade-off. Behind the above trend, vessel diameter

and vessel density also contradicted the hypotheses. Vessel diameter did not change along the

aridity gradient, while vessel density increased toward drier sites (from 49.14 to 82.07 micron).

The drier sites (KOG) had higher Kp, higher twig density and higher wood density than the wetter

sites on site scales, but we also found Kp negatively correlated with twig density on species scales

(see Appendix 4). ANK-01 had very high wood density and twig density which breaks the

increasing trend formed by other plots. To conclude, the trends of all photosynthetic traits were

successfully predicted by theories based on VPD alone. As leaf economy traits, soil nutrients and

soil moisture (Θ or α) overall did not have a clear trend along the gradient, considering nutrient

and water deliveries to leaves does not seem to aid the prediction of variation of photosynthetic

traits along the aridity gradient.

With variance partitioning, we found that the plot-to-plot trends of all traits were dominated by

inter-specific rather than intra-specific variation (i.e., components [a] are smaller than [b] in

Appendix 5). Such a finding was expected as there are few common species between plots. The

analogous patterns between twig and wood density along the aridity gradient also supported
species turnover since twig density was field measured and wood density was parsed from a global
database by species (Zanne *et al.*, 2009). To double-check the conclusion of predominant
interspecific variation, we recalculated community weighted means by assuming that the same
species share the same value of trait (i.e. remove intraspecific variation) and extrapolated trait
values to forest plots without trait measurements. We found that the conclusions in Table 1 still
hold after extrapolation (Appendix 5). The trend of trait variation from plot to plot could be well
re-constructed on a species basis, which hints at the possibility of extrapolation to other sites with
species composition information or upscaling to larger scales. Nonetheless, for within plot
variance, intraspecific variation or measurement errors (component [d]) were large for most traits:
accounting for 95% of turgor loss point variance, followed by V_{cmax25} (74%) and J_{max25} (66%).

The coordination between photosynthesis and water transportation

Data from our West African aridity gradient reveal a weak positive correlation between Kp and
AS/AL, contradictory toHypothesis 15, and inconsistent with the negative correlation that emerged
on global scales (Appendix 4). AS/AL for the Ghanaian aridity gradient was higher than the
pantropical average (Appendix 4). For hypothesis 16, we further explore the link between AS/AL,
Kp and photosynthetic trait using PCA. Species with both high AS/AL and Kp tend to have higher
V_{cmax25} and lower ci/ca. Such species tend to be deciduous and appear more in drier plots (Figure
2). There was a larger variance of hydraulic traits compared to photosynthetic traits (Figure 2).
The pattern is consistent if we redo the above PCA with Asat instead of V_{cmax25} (Appendix 4). This
finding supports hypothesis 16 (Table 1) as well as equation 3.

Discussion

The trend of traits along the aridity gradient

Although most hypotheses (Table 1) have been tested with spatially varying aridity at multiple
scales (Harrison *et al.*, 2021), testing them along the Ghana aridity gradient helps to scrutinize the

pattern in the absence of temperature variation. The patterns of all photosynthetic traits measured
along the aridity gradient (ci/ca, J_{max25}, V_{cmax25}, Dresp, Rs, Asat, Amax, namely hypotheses 1-7)
are consistent with the theoretical expectations, which underscores that aridity is a direct and
critical driver of photosynthetic traits, in absence of confounding effect with temperature. The
increase of photosynthetic capacity towards drier sites is useful in explaining multiple previous
observations, including that (1) savanna has higher photosynthesis rates than wet evergreen forest
(Gvozdevaite, 2018; Oliveras *et al.*, 2020) (2) woody savanna has sparse canopy but similar net
primary productivity to wet evergreen forest (Moore *et al.*, 2018) (3) for wet Amazonia forests,
leaves flushed in dry season have higher photosynthetic capacities which increase forest
productivity (Wu *et al.*, 2020; Green *et al.*, 2020).

From a water transportation perspective, forests in drier sites have higher TLP, lower H_{max} and
higher AS/AL (agreed with hypotheses 8-10), in support of a greater mid-day transpiration stream.
However, hypotheses derived from the safety-efficiency trade-off (hypotheses for Kp and vessel
diameter) were not in agreement with measurements along the aridity gradient. It is possible that
the trade-off may work well for single-species studies (Pritzkow *et al.*, 2020) and become weak
on large scales and across species (Gleason *et al.*, 2016; Grossiord *et al.*, 2020b). Much higher
deciduousness in KOG (dry site) than in the wet sites may play a role as higher hydraulic efficiency
was reported from deciduous species or more deciduous forests (Choat *et al.*, 2005; Chen *et al.*,
2008; Liu *et al.*, 2021). We reported a negative correlation between AS/AL and Kp at global scales
but a positive correlation along the aridity gradient (Appendix 4). One of the reasons for these
contrasting opposite correlations may lie in a geographical sampling bias – the global dataset with
scarce data points from West Africa compared with the Ghanaian dataset. The other possibility
could be a confounding effect by temperature or vegetation type at the global scale (a Simpson's
paradox); for example, a negative correlation between AS/AL and Ks was reported globally
(Mencuccini *et al.*, 2019b) and on continental (Australia) scales (Gleason *et al.*, 2012), but an
insignificant correlation was also reported for tropical forest stands on local scales without varying
temperature (Poorter *et al.*, 2010; Schuldt *et al.*, 2013; Hoeber *et al.*, 2014).

By assuming that traits with a clear and strong trend along the aridity gradient are more tightly
bound with aridity (Figure 1), ci/ca, TLP and AS/AL was found to be the most aridity-driven traits.

The runners-up are Rd, Rs, $J_{max25,}$ and V_{cmax25}, which was thought acclimated to ci/ca and light

intensity (Wang *et al.*, 2017b). Although ci/ca, Vcmax25, Kp and AS/AL all vary from wet to dry

sites, the PCA (Figure 2) further illustrates that, surprisingly, it is photosynthetic traits instead of

hydraulic traits that contrast species from wet to dry sites (also from evergreen to deciduous).

Given that large photosynthetic traits variation from wet to dry plot was induced by species

turnover (Appendix 5), our studies hint that facing a drier climate, if allowed time, West African

forests photosynthesis could adapt to a drier climate by changing species abundance with possibly

more deciduousness and higher photosynthesis capacity albeit less stomatal openness (Aguirre-

Gutiérrez *et al.*, 2019). Without consideration of the positive effect of aridity on photosynthetic

capacity, models could possibly underestimate forest productivity under future drier climates.

Combining photosynthesis and hydraulic hypotheses

Our analysis unifies photosynthesis and hydraulic hypotheses to explain plant strategies along the

aridity gradient. Namely, species in drier sites (with more deciduousness) tend to develop a

photosynthesis strategy with less stomata openness (ci/ca), stronger photosynthetic capacities

(J_{max25} and V_{cmax25}) with more maintenance cost (higher Rd and Rs), quicker photosynthesis rate

(Asat) and larger maximum transpiration, supported by large Kp and large AS/AL. The product of

AS/AL and Kp is a proxy of water delivery per leaf area, which was previously found well

correlated with proxies of photosynthesis rate: Asat (Santiago *et al.*, 2004), quantum yield of

electron transport (Brodribb & Feild, 2000) and electron transfer rate (Brodribb *et al.*, 2002). The

large variance of wood traits (way larger than leaf traits) (Mencuccini *et al.*, 2019b), hints that

plants might have a wide range of choices of traits combinations to provide adequate water

transportation (Sperry *et al.*, 2002; Prentice *et al.*, 2014) in drier sites to support faster

photosynthesis. Further investigations into xylem functioning are required to understand how

larger water transportation was achieved in drier sites. Notably, we successfully predicted plants

photosynthesis strategies along the aridity gradient (hypothesis 1-7) based solely on VPD without

incorporating leaf economic traits nor soil moisture. Nonetheless, our theoretical deduction

implicitly assumes that plants in the drier site could arrange water transportation (e.g., high AS/AL

and high Kp in Figure 2) and have adequate access to belowground water. Regarding the ongoing

discussion on the impact of VPD versus soil moisture, soil moisture may be playing a role at other

temporal scales (e.g., daily) (Liu *et al.*, 2020; Fu *et al.*, 2022) or under extreme soil drought (Sperry
et al., 2002).

Conclusion

Along the aridity gradient, we found that species with both higher AS/AL and Kp (greater potential
mid-day transpiration steam) tend to have higher Vcmax25 and lower ci/ca, which appears more
in drier sites with more deciduous species. With such a working example in West Africa, the study
not only underscores the importance of incorporating the positive effect of aridity on
photosynthesis capacity in carbon modelling but also simplifies the coupling between carbon and
water cycle: future modelling studies, following our theoretical framework, could estimate site-
time-averaged leaf-level photosynthesis simply from VPD only and consider soil moisture (or
water availability) in simulating forest stands dynamic such as drought-induced mortality or
seasonality (Hubau *et al.*, 2020; Bauman *et al.*, 2022).

Data availability

458 Figures could be downloaded from

459 https://github.com/Hzhang-ouce/Ghana_rainfall_trait_variation_optimality_github. To reproduce
460 figures, data and R codes mentioned in the main text could also be found in the above repository.

Acknowledgements

We thank Natascha Luijken, Ya-Jun Chen, Yu-Heng Sun, Maurizio Mencuccini and Akwasi
DGyamfi, Guillaume Delhaye for valuable discussion and assistance with data processing. Y.M.
is supported by the Jackson Foundation. H.Z. received Henfrey Scholarship (by St Catherine's
College, Oxford) and Tang Scholarship (by China-Oxford Schorlarship Fund). The field data
collection was funded by a grant award (ERC GEM-TRAIT, grant no. ERC-2012-ADG_20120216)
to Y.M., with additional support for the fieldwork from a Royal Society-Leverhulme Africa

Capacity Building Award and Marie Curie Fellowship to I.O. (FP7-2012-IEF-327990-
TipTropTrans).

Conflict of interest

474   The authors have no conflicts of interest to declare that are relevant to the content of this article.

Author contributions

IO, YM, ICP, and HZ designed the research and interpreted the results. DS did the hydraulic
modelling. YM, AG, TP, MB, KZ, SAB, JAG, IO, HZ contributed to data collection. HZ carried
out the analyses and wrote the paper with inputs and revisions from all co-authors.

Table 1 Traits name, unit, hypotheses and findings from field measurements along the rainfall
gradient, Green color denotes consistency between theory and our field data. Orange color
denote inconsistency.

#	Hypotheses	Data	Consistent
	Variables associated with photosynthesis and respiration (Optimality theory)		
1	Toward drier sites, the ratio between leaf-internal and ambient CO2 (ci/ca, %) (from 13C) **decreases**.	Decrease	✓
2	Toward drier sites, Rubisco carboxylation capacity at 25 C° (Vc_{max25}, umol CO2 m2 s-1) **increases**.	Slight increase	✓
3	Toward drier sites, electron transport capacity at 25 C° (J_{max25}, umol CO2 m2 s-1) **increases**.	Slight increase	✓
4	Toward drier sites, light saturated assimilation rate at 400 ppm (Asat, umol CO2 m2 s-1) **increases**.	Increases	✓
5	Toward drier sites, light saturated assimilation rate at 2000 ppm (Amax, umol CO2 m2 s-1) **increases.**	Increase	✓

6	Toward drier sites, leaf dark respiration (Rd, umol CO_2 m2 s-1) **increases**.	Increase	✓
7	Toward drier sites, specific stem respiration (Rs, umol CO_2 m2 s-1) **increases.**	Increase	✓
	Variables associated with water transportation		
8	Toward drier sites, Sapwood to leaf area ratio (Huber value) (AS/AL, cm2 m-2) **increases.**	Increase	✓
9	Toward drier sites, turgor loss point (TLP, MPa) becomes more negative.	More negative	✓
10	Toward drier sites, plant stature, calculated as maximum tree height of a species (H_{max}, m) **decreases**.	Slight decrease	✓
11	Toward drier sites, wood density (g cm-3) and twig density (g cm-3) **increase** (if following the safety-efficiency trade-off).	Slight increase	✓
12	Toward drier sites, potential specific hydraulic conductivity (Kp, kg m-1 Mpa-1 s-1) **decreases** (if following the safety-efficiency trade-off).	Slight increase	
13	Toward drier sites, vessel diameter (micron) **decreases.** (if following the safety-efficiency trade-off).	No trend	
14	Toward drier sites, vessel density (mm-2) **decreases.** (if following the safety-efficiency trade-off).	Increase	
15	AS/AL and Kp are negatively correlated. (if following safety-efficiency trade-off, and global scale analysis – see introduction)	Positive correlation	
16	For species with high A_S/A_L and Kp, there is high Vcmax (or high Asat)	See Figure 3	✓

'Data' column summarize patterns in Figure 1. A trend of trait is qualitatively recognized if KOG (dry region) is significantly different to ANK (wet region) while BOB ranks between. 'Slight increase' suggest that the pattern fits the above criteria broadly albeit one plot behave inconsistently. Colours indicate results that are consistent (green), weakly consistent (light green) and inconsistent (orange) with theoretical expectations. Ticks in the column 'consistent' indicate consistency between hypothesies and data

MAT, MAP and soil moisture

Variables associated with photosynthesis and respiration

Figure 1 Plot scale community weighted mean (with standard error) from the wettest (left) to the

driest (right) plot. Mean annual air temperature, precipitation and soil volumetric water content

at 12 cm depth were also shown. Forest plots are arrayed from left to right in order of increasing

aridity according to the aridity gradient description described in the text. The number denotes the

number of samples, which could be a leaf, a branch, a tree or a species depending on the variable.

The letters denote significance (P<0.05) in plot-to-plot difference.

Figure 2 Principal components analysis for Huber value (AS/AL), the ratio between leaf internal and ambient CO_2 (ci/ca), Rubisco carboxylation capacity at 25 degree (Vcmax25) and potential specific conductivity (Kp) on species scale. Values are transformed to achieve normal distribution but not standardized to equal variance; therefore the length of arrows denotes the variance of the specific trait. The ellipses for each site are confidence ellipses around group mean points. The PCA axes in all figures are identical. Note that the three figures display the same PCA, but with a different classification of scatter points.

References

Aguirre-Gutiérrez J, Oliveras I, Rifai S, Fauset S, Adu-Bredu S, Affum-Baffoe K, Baker TR, Feldpausch TR, Gvozdevaite A, Hubau W, *et al.* **2019**. Drier tropical forests are susceptible to functional changes in response to a long-term drought. *Ecology Letters* **22**: 855–865.

Atkin OK, Bloomfield KJ, Reich PB, Tjoelker MG, Asner GP, Bonal D, Bönisch G, Bradford MG, Cernusak LA, Cosio EG, *et al.* **2015**. Global variability in leaf respiration in relation to climate, plant functional types and leaf traits. *New Phytologist* **206**: 614–636.

Bahar NHA, Ishida FY, Weerasinghe LK, Guerrieri R, O'Sullivan OS, Bloomfield KJ, Asner GP, Martin RE, Lloyd J, Malhi Y, *et al.* **2017**. Leaf-level photosynthetic capacity in lowland Amazonian and high-elevation Andean tropical moist forests of Peru. *New Phytologist* **214**: 1002–1018.

Bartlett MK, Scoffoni C, Sack L. **2012**. The determinants of leaf turgor loss point and prediction of drought tolerance of species and biomes: A global meta-analysis. *Ecology Letters* **15**.

Bauman D, Fortunel C, Delhaye G, Malhi Y, Cernusak LA, Bentley LP, Rifai SW, Aguirre-Gutiérrez J, Menor IO, Phillips OL, *et al.* **2022**. Tropical tree mortality has increased with rising atmospheric water stress. *Nature*.

Beerling DJ, Quick WP. **1995**. A new technique for estimating rates of carboxylation and electron
transport in leaves of C3 plants for use in dynamic global vegetation models. *Global Change*
Biology **1**: 289–294.

Bittencourt PRL, Pereira L, Oliveira RS. **2016**. On xylem hydraulic efficiencies, wood space-
use and the safety–efficiency tradeoff. *New Phytologist* **211**: 1152–1155.

Björkman O. **1981**. Responses to Different Quantum Flux Densities. *Physiological Plant Ecology*
I: 57–107.

Brodribb TJ, Feild TS. **2000**. Stem hydraulic supply is linked to leaf photosynthetic capacity:
evidence from New Caledonian and Tasmanian rainforests. *Plant, Cell & Environment* **23**: 1381–
1388.

Brodribb TJ, Holbrook NM, Gutierrez M v. **2002**. Hydraulic and photosynthetic co-ordination
in seasonally dry tropical forest trees. *Plant, Cell & Environment* **25**: 1435–1444.

Canadell JG, Monteiro PMS, Costa MH, Cunha LC da, Cox PM, Eliseev A v., Henson S,
Ishii M, Jaccard S, Koven C, *et al.* **2021**. Global carbon and other biogeochemical cycles and
feedbacks. In: Climate Change 2021: The Physical Science Basis. Contribution of Working Group
I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.

Chen J-W, Zhang Q, Cao K-F. **2008**. Inter-species variation of photosynthetic and xylem
hydraulic traits in the deciduous and evergreen Euphorbiaceae tree species from a seasonally
tropical forest in south-western China. *Ecological Research* **24**: 65.

Chiti T, Certini G, Grieco E, Valentini R. **2010**. The role of soil in storing carbon in tropical
rainforests: The case of Ankasa Park, Ghana. *Plant and Soil* **331**: 453–461.

Choat B, Ball MC, Luly JG, Holtum JAM. **2005**. Hydraulic architecture of deciduous and
evergreen dry rainforest tree species from north-eastern Australia. *Trees* **19**: 305–311.

Choat B, Jansen S, Brodribb TJ, Cochard H, Delzon S, Bhaskar R, Bucci SJ, Feild TS,
Gleason SM, Hacke UG, *et al.* **2012**. Global convergence in the vulnerability of forests to drought.
Nature 2012 491:7426 **491**: 752–755.

Cunningham SC. **2005**. Photosynthetic responses to vapour pressure deficit in temperate and
tropical evergreen rainforest trees of Australia. *Oecologia* **142**: 521–528.

Dai Z, Edwards GE, Ku MSB. **1992**. Control of photosynthesis and stomatal conductance in
Ricinus communis L.(castor bean) by leaf to air vapor pressure deficit. *Plant physiology* **99**: 1426.

Domingues TF, Meir P, Feldpausch TR, Saiz G, Veenendaal EM, Schrodt F, Bird M,
Djagbletey G, Hien F, Compaore H, *et al.* **2010**. Co-limitation of photosynthetic capacity by
nitrogen and phosphorus in West Africa woodlands. *Plant, Cell & Environment* **33**: 959–980.

Franklin O, Harrison SP, Dewar R, Farrior CE, Brännström Å, Dieckmann U, Pietsch S,
Falster D, Cramer W, Loreau M, *et al.* **2020**. Organizing principles for vegetation dynamics.
Nature Plants **6**: 444–453.

Fu Z, Ciais P, Prentice IC, Gentine P, Makowski D, Bastos A, Luo X, Green JK, Stoy PC,
Yang H, *et al.* **2022**. Atmospheric dryness reduces photosynthesis along a large range of soil water
deficits. *Nature Communications 2022 13:1* **13**: 1–10.

Gleason SM, Butler DW, Waryszak P. **2013**. Shifts in leaf and stem hydraulic traits across aridity
gradients in eastern Australia. *International Journal of Plant Sciences* **174**: 1292–1301.

Gleason SM, Butler DW, Ziemińska K, Waryszak P, Westoby M. **2012**. Stem xylem
conductivity is key to plant water balance across Australian angiosperm species. *Functional*
Ecology **26**: 343–352.

Gleason SM, Westoby M, Jansen S, Choat B, Hacke UG, Pratt RB, Bhaskar R, Brodribb TJ,
Bucci SJ, Cao KF, *et al.* **2016**. Weak tradeoff between xylem safety and xylem-specific hydraulic
efficiency across the world's woody plant species. *New Phytologist* **209**: 123–136.

Granier A, Hue R, Barigah ST. **1996**. Transpiration of natural rain forest and its dependence on
climatic factors. *Agricultural and Forest Meteorology* **78**: 19–29.

Green JK, Berry J, Ciais P, Zhang Y, Gentine P. **2020**. Amazon rainforest photosynthesis
increases in response to atmospheric dryness. *Science Advances* **6**.

Grossiord C, Buckley TN, Cernusak LA, Novick KA, Poulter B, Siegwolf RTW, Sperry JS, McDowell NG. **2020a**. Plant responses to rising vapor pressure deficit. *New Phytologist* **226**: 1550–1566.

Grossiord C, Ulrich DEM, Vilagrosa A. **2020b**. Controls of the hydraulic safety–efficiency trade-off. *Tree Physiology* **40**: 573–576.

Gvozdevaite A. **2018**. The role of economic, venation and morphological leaf traits in plant and ecosystem function along forest-savanna gradients in the tropics.

Gvozdevaite A, Oliveras I, Domingues TF, Peprah T, Boakye M, Afriyie L, da Silva Peixoto K, de Farias J, Almeida de Oliveira E, Almeida Farias CC, *et al.* **2018**. Leaf-level photosynthetic capacity dynamics in relation to soil and foliar nutrients along forest–savanna boundaries in Ghana and Brazil. *Tree Physiology* **38**: 1912–1925.

Hacke UG, Sperry JS, Pockman WT, Davis SD, McCulloh KA. **2001**. Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia 2000 126:4* **126**: 457–461.

Han T, Feng Q, Yu T, Yang X, Zhang X, Li K. **2022**. Characteristic of Stomatal Conductance and Optimal Stomatal Behaviour in an Arid Oasis of Northwestern China. *Sustainability 2022, Vol. 14, Page 968* **14**: 968.

Harrison SP, Cramer W, Franklin O, Prentice IC, Wang H, Brännström Å, de Boer H, Dieckmann U, Joshi J, Keenan TF, *et al.* **2021**. Eco-evolutionary optimality as a means to improve vegetation and land-surface models. *New Phytologist* **231**: 2125–2141.

Hoeber S, Leuschner C, Köhler L, Arias-Aguilar D, Schuldt B. **2014**. The importance of hydraulic conductivity and wood density to growth performance in eight tree species from a tropical semi-dry climate. *Forest Ecology and Management* **330**: 126–136.

Hubau W, Lewis SL, Phillips OL, Affum-Baffoe K, Beeckman H, Cuní-Sanchez A, Daniels AK, Ewango CEN, Fauset S, Mukinzi JM. **2020**. Asynchronous carbon sink saturation in African and Amazonian tropical forests. *Nature* **579**: 80–87.

Liu H, Gleason SM, Hao G, Hua L, He P, Goldstein G, Ye Q. **2019**. Hydraulic traits are
coordinated with maximum plant height at the global scale. *Science Advances* **5**: eaav1332.

Liu L, Gudmundsson L, Hauser M, Qin D, Li S, Seneviratne SI. **2020**. Soil moisture dominates
dryness stress on ecosystem production globally. *Nature Communications* **11**: 4892.

Liu H, Ye Q, Gleason SM, He P, Yin D. **2021**. Weak tradeoff between xylem hydraulic efficiency
and safety: climatic seasonality matters. *New Phytologist* **229**: 1440–1452.

Long SP, Woolhouse HW. **1978**. The responses of net photosynthesis to vapour pressure deficit
and CO2 concentration in Spartina townsendii (sensu lato), a C4 species from a cool temperate
climate. *Journal of Experimental Botany* **29**: 567–577.

López R, Cano FJ, Martin-Stpaul NK, Cochard H, Choat B. **2021**. Coordination of stem and
leaf traits define different strategies to regulate water loss and tolerance ranges to aridity. *New*
Phytologist **230**: 497–509.

Maire V, Martre P, Kattge J, Gastal F, Esser G, Fontaine S, Soussana J-F. **2012**. The
coordination of leaf photosynthesis links C and N fluxes in C3 plant species. *PloS one* **7**: e38345.

Maire V, Wright IJ, Prentice IC, Batjes NH, Bhaskar R, van Bodegom PM, Cornwell WK,
Ellsworth D, Niinemets Ü, Ordonez A. **2015**. Global effects of soil and climate on leaf
photosynthetic traits and rates. *Global Ecology and Biogeography* **24**: 706–717.

Malhi Y, Girardin C, Metcalfe DB, Doughty CE, Aragão LEOC, Rifai SW, Oliveras I,
Shenkin A, Aguirre-Gutiérrez J, Dahlsjö CAL, *et al.* 2021. The Global Ecosystems Monitoring
network: Monitoring ecosystem productivity and carbon cycling across the tropics. *Biological*
Conservation **253**: 108889.

Manzoni S, Vico G, Katul G, Palmroth S, Jackson RB, Porporato A. **2013**. Hydraulic limits
on maximum plant transpiration and the emergence of the safety–efficiency trade-off. *New*
Phytologist **198**: 169–178.

Martínez-Vilalta J, Cochard H, Mencuccini M, Sterck F, Herrero A, Korhonen JFJ, Lloréns
P, Nikinmaa E, Nolè A, Poyatos R, *et al.* 2009. Hydraulic adjustment of Scots pine across Europe.
The New phytologist **184**: 353–364.

Medlyn BE, Duursma RA, Eamus D, Ellsworth DS, Prentice IC, Barton CVM, Crous KY,
de Angelis P, Freeman M, Wingate L. **2011**. Reconciling the optimal and empirical approaches
to modelling stomatal conductance. *Global Change Biology* **17**: 2134–2144.

Mencuccini M, Manzoni S, Christoffersen B. **2019a**. Modelling water fluxes in plants: from
tissues to biosphere. *New Phytologist* **222**: 1207–1222.

Mencuccini M, Rosas T, Rowland L, Choat B, Cornelissen H, Jansen S, Kramer K, Lapenis
A, Manzoni S, Niinemets Ü, *et al.* 2019b. Leaf economics and plant hydraulics drive leaf : wood
area ratios. *New Phytologist* **224**: 1544–1556.

Moore S, Adu-Bredu S, Duah-Gyamfi A, Addo-Danso SD, Ibrahim F, Mbou AT, de
Grandcourt A, Valentini R, Nicolini G, Djagbletey G, *et al.* 2018. Forest biomass, productivity
and carbon cycling along a rainfall gradient in West Africa. *Global Change Biology* **24**: e496–
e510.

Neelin JD, Münnich M, Su H, Meyerson JE, Holloway CE. **2006**. Tropical drying trends in
global warming models and observations. *Proceedings of the National Academy of Sciences* **103**:
6110–6115.

Ögren E. **1993**. Convexity of the Photosynthetic Light-Response Curve in Relation to Intensity
and Direction of Light during Growth. *Plant physiology* **101**: 1013–1019.

Oliveras I, Bentley L, Fyllas NM, Gvozdevaite A, Shenkin AF, Peprah T, Morandi P, Peixoto
KS, Boakye M, Adu-Bredu S, *et al.* 2020. The Influence of Taxonomy and Environment on Leaf
Trait Variation Along Tropical Abiotic Gradients. *Frontiers in Forests and Global Change* **3**: 18.

Olson ME, Anfodillo T, Rosell JA, Petit G, Crivellaro A, Isnard S, León-Gómez C, Alvarado-
Cárdenas LO, Castorena M. **2014**. Universal hydraulics of the flowering plants: vessel diameter
scales with stem length across angiosperm lineages, habits and climates. *Ecology Letters* **17**: 988–
997.

Olson ME, Rosell JA. **2013**. Vessel diameter–stem diameter scaling across woody angiosperms
and the ecological causes of xylem vessel diameter variation. *New Phytologist* **197**: 1204–1213.

Peng Y, Bloomfield KJ, Cernusak LA, Domingues TF, Colin Prentice I. **2021**. Global climate
and nutrient controls of photosynthetic capacity. *Communications Biology 2021 4:1* **4**: 1–9.

Pfautsch S, Harbusch M, Wesolowski A, Smith R, Macfarlane C, Tjoelker MG, Reich PB,
Adams MA. **2016**. Climate determines vascular traits in the ecologically diverse genus Eucalyptus.
Ecology Letters **19**: 240–248.

Poorter L, McDonald I, Alarcón A, Fichtler E, Licona JC, Peña-Claros M, Sterck F, Villegas
Z, Sass-Klaassen U. **2010**. The importance of wood traits and hydraulic conductance for the
performance and life history strategies of 42 rainforest tree species. *New Phytologist*.

Prentice IC, Dong N, Gleason SM, Maire V, Wright IJ. **2014**. Balancing the costs of carbon
gain and water transport: testing a new theoretical framework for plant functional ecology. *Ecology*
Letters **17**: 82–91.

Pritzkow C, Williamson V, Szota C, Trouvé R, Arndt SK. **2020**. Phenotypic plasticity and
genetic adaptation of functional traits influences intra-specific variation in hydraulic efficiency
and safety. *Tree Physiology* **40**: 215–229.

Quesada CA, Lloyd J, Schwarz M, Patiño S, Baker TR, Czimczik C, Fyllas NM, Martinelli
L, Nardoto GB, Schmerler J, *et al.* **2010**. Variations in chemical and physical properties of
Amazon forest soils in relation to their genesis. *Biogeosciences* **7**.

Restrepo-Coupe N, da Rocha HR, Hutyra LR, da Araujo AC, Borma LS, Christoffersen B,
Cabral OMR, de Camargo PB, Cardoso FL, da Costa ACL, *et al.* **2013**. What drives the
seasonality of photosynthesis across the Amazon basin? A cross-site analysis of eddy flux tower
measurements from the Brasil flux network. *Agricultural and Forest Meteorology* **182–183**: 128–
144.

Rogers A, Medlyn BE, Dukes JS, Bonan G, von Caemmerer S, Dietze MC, Kattge J, Leakey
ADB, Mercado LM, Niinemets Ü, *et al.* **2017**. A roadmap for improving the representation of
photosynthesis in Earth system models. *New Phytologist* **213**: 22–42.

Ryan MG, Yoder BJ. **1997**. Hydraulic Limits to Tree Height and Tree Growth. *BioScience* **47**.

Sandoval D, Prentice IC. **2020**. Simple process-led algorithms for simulating habitats (SPLASH
v.2.0): robust calculations of water and energy fluxes in complex terrain. *EGU General Assembly*
2020 .

Santiago LS, Goldstein G, Meinzer FC, Fisher JB, Machado K, Woodruff D, Jones T. **2004**.
Leaf photosynthetic traits scale with hydraulic conductivity and wood density in Panamanian
forest canopy trees. *Oecologia* **140**: 543–550.

Schuldt B, Leuschner C, Brock N, Horna V. **2013**. Changes in wood density, wood anatomy
and hydraulic properties of the xylem along the root-to-shoot flow path in tropical rainforest trees.
Tree physiology **33**: 161–174.

Smith NG, Keenan TF, Prentice IC, Wang H, Wright IJ, Niinemets U, Crous KY, Domingues
TF, Guerrieri R, Ishida FY, ***et al.*** **2019**. Global photosynthetic capacity is optimized to the
environment.

Sperry JS, Hacke UG, Oren R, Comstock JP. **2002**. Water deficits and hydraulic limits to leaf
water supply. *Plant, Cell & Environment* **25**: 251–263.

Stangl ZR, Tarvainen L, Wallin G, Ubierna N, Räntfors M, Marshall JD. **2019**. Diurnal
variation in mesophyll conductance and its influence on modelled water-use efficiency in a mature
boreal Pinus sylvestris stand. *Photosynthesis Research* **141**: 53–63.

Stocker BD, Wang H, Smith NG, Harrison SP, Keenan TF, Sandoval D, Davis T, Prentice
IC. **2020**. P-model v1.0: An optimality-based light use efficiency model for simulating ecosystem
gross primary production. *Geoscientific Model Development* **13**: 1545–1581.

Togashi HF, Prentice IC, Evans BJ, Forrester DI, Drake P, Feikema P, Brooksbank K,
Eamus D, Taylor D. **2015**. Morphological and moisture availability controls of the leaf area-to-
sapwood area ratio: analysis of measurements on A ustralian trees. *Ecology and Evolution* **5**: 1263–
1270.

Walker AP, Beckerman AP, Gu L, Kattge J, Cernusak LA, Domingues TF, Scales JC,
Wohlfahrt G, Wullschleger SD, Woodward FI. **2014**. The relationship of leaf photosynthetic
traits – Vcmax and Jmax – to leaf nitrogen, leaf phosphorus, and specific leaf area: a meta-analysis
and modeling study. *Ecology and Evolution* **4**: 3218–3235.

Wang H, Prentice IC, Davis TW, Keenan TF, Wright IJ, Peng C. **2017a**. Photosynthetic
responses to altitude: an explanation based on optimality principles. *New Phytologist* **213**: 976–
982.

Wang H, Prentice IC, Keenan TF, Davis TW, Wright IJ, Cornwell WK, Evans BJ, Peng C.
2017b. Towards a universal model for carbon dioxide uptake by plants. *Nature Plants 2017 3:9* **3**:
734–741.

Warton DI, Duursma RA, Falster DS, Taskinen S. **2012**. smatr 3-an R package for estimation
and inference about allometric lines. *Methods in ecology and evolution* **3**: 257–259.

Wright IJ, Reich PB, Cornelissen JHC, Falster DS, Groom PK, Hikosaka K, Lee W, Lusk
CH, Niinemets Ü, Oleksyn J. **2005**. Modulation of leaf economic traits and trait relationships by
climate. *Global Ecology and Biogeography* **14**: 411–421.

Wright IJ, Reich PB, Westoby M. **2001**. Strategy shifts in leaf physiology, structure and nutrient
content between species of high- and low-rainfall and high- and low-nutrient habitats. *Functional*
Ecology **15**: 423–434.

Wright IJ, Reich PB, Westoby M. **2003**. Least-cost input mixtures of water and nitrogen for
photosynthesis. *The American naturalist* **161**: 98–111.

Wright IJ, Reich PB, Westoby M, Ackerly DD, Baruch Z, Bongers F, Cavender-Bares J,
Chapin T, Cornellssen JHC, Diemer M, *et al.* 2004. The worldwide leaf economics spectrum.
Nature 2004 428:6985 **428**: 821–827.

Wu J, Serbin SP, Ely KS, Wolfe BT, Dickman LT, Grossiord C, Michaletz ST, Collins AD,
Detto M, McDowell NG, *et al.* 2020. The response of stomatal conductance to seasonal drought
in tropical forests. *Global Change Biology* **26**: 823–839.

Xu H, Wang H, Prentice IC, Harrison SP, Wright IJ. **2021**. Coordination of plant hydraulic
and photosynthetic traits: confronting optimality theory with field measurements. *New Phytologist*
232: 1286–1296.

Yamazaki D, Ikeshima D, Sosa J, Bates PD, Allen GH, Pavelsky TM. **2019**. MERIT Hydro: A
high-resolution global hydrography map based on latest topography dataset. *Water Resources*
Research **55**: 5053–5073.

Yang J, He Y, Aubrey DP, Zhuang Q, Teskey RO. **2016**. Global patterns and predictors of stem
CO2 efflux in forest ecosystems. *Global Change Biology* **22**: 1433–1444.

Yang Y, Wang H, Harrison SP, Prentice IC, Wright IJ, Peng C, Lin G. **2019**. Quantifying
leaf-trait covariation and its controls across climates and biomes. *New Phytologist* **221**: 155–168.

Zanne AE, Lopez-Gonzalez G, Coomes DA, Ilic J, Jansen S, Lewis SL, Miller RB, Swenson
NG, Wiemann MC, Chave J. **2009**. Data from: Towards a worldwide wood economics spectrum.

745