On the physiological and structural contributors to the dynamic balance of excitation and inhibition in local cortical networks *

Farshad Shirani†   Hannah Choi†

January 10, 2023

Abstract

Overall balance of excitation and inhibition in cortical networks is central to their functionality and normal operation. Such orchestrated temporal co-evolution of excitation and inhibition is established through local connections, and is homeostatically regulated to keep the networks in a vigilant state, so that they readily respond to their inputs and rapidly transition between different states without losing their stability. Several homeostatic plasticity mechanisms have been identified that adjust the strength and kinetics of synaptic activity in cortical networks to restore their balance when it is perturbed from a normal level. However, it is not yet well-understood how a balanced state is established, how it is dynamically regulated, and how it can accurately be measured, partially due to lack of a universally accepted definition of excitation-inhibition balance. Here, we use biologically plausible mathematical models to demonstrate how some of the key physiological factors that control the kinetics of synapses and structural factors that determine the overall topology of a cortical network contribute to establishment of an overall balance of excitation and inhibition throughout the network. We characterize a network's baseline balanced state by a set of functional properties, and show that deviations from this reference state can be continuously quantified by measuring the ratio between mean excitatory and mean inhibitory synaptic conductances in the network. We show how variations in physiological and structural parameters of a network dynamically change the level of excitation-inhibition balance and result in transitions in the spontaneous activity of the network to high-amplitude slow oscillatory regimes. Our results particularly show that the commonly observed ratio between the number of inhibitory and excitatory neurons in local cortical networks is nearly optimal, and the values of inhibitory synaptic decay time constants and the density of inhibitory-to-inhibitory connectivity are critical to network balance and stability.

Author summary

Excitatory neurons largely dominate the microcircuitry of cortical networks and are central to their information processing endeavor. The recurrent excitation that they provide to each other allows the entire network to sustain its activity. Inhibitory neurons then play a crucial role in stabilizing and regulating this recurrent excitation. Otherwise, the recurrent

---

*This work was supported by the National Eye Institute of the National Institutes of Health under Award Number R00 EY030840. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

†School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332, USA

excitation will lead to saturation of the excitatory activity. This dynamic self-regulated coordination of overall neuronal activity is a fundamental property of cortical networks. It results in an intrinsic balance of excitation and inhibition throughout a network, which sets the operating condition of the network's spontaneous activity at a highly excitable yet fully stabilized state.The importance of this excitation-inhibition balance to functionality of cortical networks has motivated numerous theoretical and experimental studies, aiming to understand how such a balance is established locally in cortical networks, and under what conditions it is disrupted. To further inform such studies, we use biologically plausible models of spontaneous neocortical activity to study how some of the key physiological and structural properties of local cortical networks contribute to establishing excitation-inhibition balance, and how this balance is dynamically affected by variations in those properties.

Introduction

A cortical network of one cubic millimeter in volume in mammalian neocortex is composed of tens of thousands of neurons, the majority of which being excitatory. Each of the neurons in this network receives excitatory and inhibitory synaptic inputs from over a thousand other neurons, both through long-range corticocortical fibers coming from neurons residing outside the network and through intracortical fibers coming from the neurons inside the network [1], [2, Figure 1]. Within the network, neurons are highly interconnected via all types of excitatory-to-excitatory, excitatory-to-inhibitory, inhibitory-to-excitatory and inhibitory-to-inhibitory connections. Such a massively interconnected network of dynamically interacting neurons must have intrinsic mechanisms to control the level of overall excitation and inhibition that is generated in the network at every instances of time. If the recurrent excitation provided by the population of excitatory neurons on itself—which is necessary for self-sustaining activity of the network—is not sufficiently balanced by the inhibition it receives from inhibitory neurons, then the overall level of excitation in the network can rapidly rise to an extreme level at which the spiking rates of the neurons saturate. Oppositely, if the inhibitory neurons impose excessive inhibition on the excitatory population, the network loses the level of excitability that it needs to effectively respond to inputs coming from other cortical areas. Hence, maintaining an overall balance of excitation and inhibition is crucial for the functionality of a cortical network.

Theoretical and experimental studies have confirmed the existence of a dynamically regulated balance of excitation and inhibition in local cortical networks at multiple states of wakefulness and sleep [3–18]. It has been hypothesized that balance of excitation and inhibition is essential for controlling network-level information transmission [19, 20], efficient, high-precision, and high-dimensional representations and processing of sensory information [5,11,13], enabling cortical computations by enhancing the range of network sensitivity to sensory inputs [8], selective amplification of specific activity patterns in unstructured inputs [21], maintaining information in working memory [22], and, importantly, preserving network stability [10,23]. Pathological conditions resulting in deviations from normal levels of excitation-inhibition balance, hence hypo- or hyper-excitation in cortical networks, have been associated with several neurological disorders, such as Autism Spectrum Disorders, schizophrenia, mood disorders, Alzheimer's disease, Rett Syndrome, and epilepsy [7, 9, 19, 23–27]. Nevertheless, neuromodulatory mediated deviations from finely balanced network states, as long as they do not result in pathological dysfunction, are also thought to be important in enabling certain network operations, such as performing long-term changes in sensory receptive fields [8,9,24].

Despite the importance of excitation-inhibition balance in cortical functionality, and numerous homeostatic mechanisms that have been proposed as possible regulatory processes involved in maintaining this balance [8, 10, 19, 23, 28–33], little is known about how balance of excitation and inhibition is established in cortical networks, what cellular and network properties are homeostatically adjusted to maintain it, and how it can be accurately and meaningfully measured [16, 28, 34]. However, it can be conceived that the balance is most likely established and regulated locally, that means through internal recurrent interactions within a local cortical network, since global interactions between different regions of the cortex is predominantly excitatory and cannot be effectively balanced at a global scale by short-range activity of inhibitory neurons [2, 7, 11, 18, 35].

One source of ambiguity in interpreting currently available experimental and theoretical results, which also prevents developing measurement techniques that are generally applicable to different studies, is that in fact there is still no universally accepted definition of excitation-inhibition balance in the literature. For instance, *loose* balance of excitation and inhibition in a network has been defined as a state during which temporal variations in excitatory and inhibitory input currents to neurons are correlated on a slow time scale, whereas on a faster time scale they exhibit uncorrelated fluctuations. When faster fluctuations are also strongly correlated, with possibly a small time lag between them, the balance has been called *tight* [11, 31, 36]. From a related but mostly spatial point of view, *global* balance has been referred to a state at which, each neuron of the network receives approximately equal amounts of excitation and inhibition, so that on average the overall levels of excitation and inhibition in the network are the same. The balance is called *fine-scale* or *detailed* when, in addition to having global balance, the amounts of excitation and inhibition to neurons also balance each other at finer spatial resolution; that is on subsets of synaptic inputs corresponding to specific signaling pathways [8, 20, 31]. In these definitions, however, it remains unclear how to accurately quantify 'how well' the activity of a network is balanced, or how critical the current level of imbalance in a network is.

Here, we computationally investigate overall balance of excitation and inhibition in a local cortical network, utilizing an interpretation of network balance closely related to the global balance described above. We identify a state of overall balance by a set of qualitative network properties, which are expected to be present in a normally functioning network. In particular, under this overall balance, spontaneous spiking activity of the neurons within the network is predominately asynchronous and irregular—an activity regime commonly observed in globally and tightly balanced networks [7, 10, 11, 13, 15, 20, 36, 37]. Such a balanced network is highly responsive to inputs, and can rapidly switch between different operating states. We show how deviations from the balanced state can be quantified, and how critical levels of imbalance in a network can result in a phase transition in network dynamics.

We primarily aim to demonstrate how different physiological parameters of synapses, and some key structural parameters of a local network, contribute to establishing and maintaining overall excitation-inhibition balance in the network. Synaptic properties of a network can be dynamically adjusted through homeostatic plasticity mechanisms to compensate for changes in excitatory and inhibitory activity in the network, and thereby regulate the network's balance. Structural organization of a network determines the types and amounts of interactions between neuronal populations, and hence is central to establishing the overall balance of activity in the network. Therefore, knowing how changes in each of the key synaptic and structural parameters of a network affect its overall balance, and whether or not such changes can create a critical state in the dynamics of the network, is important in understanding the homeostatic mechanisms that regulate excitation-inhibition balance in local

cortical networks, and in identifying the sources of pathological conditions that may arise during cortical development.

We base or study on a biologically reasonable mean-field (neural mass) model of a local cortical network, so that we can take advantage of the simplicity of a mean-field framework to comprehensively investigate how the overall balance of excitation and inhibition is affected by continuous changes in each of the network parameters over a wide range of biologically plausible values. Performing such an extensive study experimentally is not practical, nor is it using detailed spiking neuronal network models. We use standard numerical bifurcation analysis techniques to demonstrate how the long-term collective activity of excitatory and inhibitory neurons in the model, and hence the overall balance of excitation and inhibition, changes with each of the key physiological and structural network parameters being taken as a bifurcation parameter. We show that, in most cases, the changes in parameters that result in over-excitation in the network can eventually become critical and lead to a transition in network's dynamics to a slow oscillatory bursting regime. Moreover, we analyze instantaneous correlations between mean excitatory and inhibitory activities in the network, and argue that a tight correlation between the temporal variations of these activities is not necessarily an indication of a functional level of balance in the network. We conclude our study by verifying some of the main findings of our mean-field-based analysis using a more detailed spiking neuronal network model.

In what follows, we first provide, in the Materials and methods section, an overview of our bifurcation analysis framework and a detailed description of the models we use to perform our analyses. In the Results section, we demonstrate the results we obtain on establishing a baseline balanced state in the network, and how this balanced state is affected by changes in each of the main synaptic and structural factors incorporated into the models. We summarize and discuss the key findings of our study in the Discussion section.

Materials and methods

We perform our analysis using a biologically plausible mean-field description of the collective dynamics of a local cortical network, developed in a sequence of works described in [38–42]. This model has been shown to fairly accurately predict mean spontaneous activity of the neurons of the network in asynchronous irregular regimes, as well as their responses to external stimuli. We set the values of the biophysical parameters of the model based on realistic values of cortical neurons in the mouse and rat brain [37, 43], so that the model presents a balanced state of excitation and inhibition, with mean firing activity of the neurons being very close to that observed in detailed models of rat cortical microcircuitry [37].

We use MatCont, version 7.3, [44] to perform numerical bifurcation analysis on the mean-field model. We choose as bifurcation parameters each of the physiological synaptic parameters of the neurons, as well as network structure parameters such as connectivity density and the percentage of the total number of neurons that are inhibitory. We perform codimension-one continuation of an stable equilibrium state of the network with respect to variations in each of the bifurcation parameters. For each of the synaptic and structural bifurcation parameters considered, we demonstrate how the equilibrium values of network quantities such as mean firing rates, mean membrane conductances, mean membrane potentials, and mean synaptic currents change as the bifurcation parameter varies within its entire range of biologically plausible values. When a phase transition to an oscillatory behavior is detected at a bifurcation point on the curves of equilibria, we also perform codimension-one continuation

of the emerging limit cycles to observe how the frequency of the oscillations changes with respect to variations in the bifurcation parameters.

Additionally, we perform codimension-two continuation of the network equilibrium by simultaneously considering both excitatory and inhibitory synaptic parameters as bifurcation parameters, to see how joint modulation of the kinetics of excitatory and inhibitory synaptic activity contributes to maintaining or disturbing the balance of excitation and inhibition throughout the network. We perform similar analysis to also investigate certain joint contribution of synaptic and structural factors.

To verify whether the results predicted by the mean-field model still remain qualitatively valid when the simplifying assumptions of the mean-field description are removed, we also construct a spiking neuronal network model with equivalent structural, synaptic, and cellular parameters to those in the mean-field model. We simulate the network for two different sets of parameter values based on the predictions of the mean-field model, one corresponding to the network's baseline balanced state with asynchronous and irregular neuronal firing activity, and the other corresponding to an imbalanced oscillatory state. We compare the results obtained from the two models using the same quantitative measures as those previously computed for the mean-field model.

Detailed description of the mean-field and spiking neuronal network models are presented below, along with a discussion on the choice of their realistic parameter values. The numerical analysis results and their biological implications are discussed in the Results section.

Mean-field neuronal population model

The mean-field model we use here has been developed based on the Markovian master equations provided in [38], which describe the overall activity of a randomly connected balanced network of neurons in an asynchronous irregular spiking regime. Specifically, these equations present the temporal evolution of the mean and variance of the firing rates of neurons within the excitatory and inhibitory populations of the network, as well as the covariance of the firing rates between the two populations. The master equations given in [38] have been extended in [41] by including an additional equation that allows for spike frequency adaptation of the neurons.

Application of the master equations given in [38] requires developing neuronal transfer functions that characterize the stationary firing rate of neurons in response to their stationary presynaptic excitatory and inhibitory spiking activity. Such transfer functions provide population-level description of the neuronal activity and capture the specific properties of the single-neuron models and synaptic interactions that are considered in building the mean-field model of the entire network. Due to the nonlinearities involved in incorporating such properties, analytical derivation of the transfer functions is often quite challenging. Here, as in [41], we use the semi-analytically calculated transfer functions that are proposed in [39] and [40] under the assumption that the characterization of the transfer functions depends only on the statistical properties of subthreshold membrane potential fluctuations. These transfer functions rely on an effective membrane potential threshold for each excitatory and inhibitory population. This effective threshold is expressed as a second-degree polynomial on the moments of the subthreshold membrane potentials within each population, namely, on the mean, standard deviation, and autocorrelation time constant of the membrane potential fluctuations. The coefficients of this second-degree polynomial are obtained by fitting it to the dynamics of numerically simulated single neurons at different excitatory and inhibitory presynaptic spiking frequencies [39–41].

Below, we briefly provide the formulation of the mean-field model as given in [41], with several notational changes, some modifications for incorporating external inputs to the network, and a correction on the equations governing neuronal adaptation.

Mean-field model equations

To present the equations of the mean-field model, let E and I denote, respectively, the excitatory and inhibitory neuronal populations of a local cortical network composed of a total number of N neurons. Note that $N = N_E + N_I$, where N_E and N_I denote the total number of excitatory and inhibitory neurons in the network, respectively. For all time $t \in [0, T]$, $T > 0$, and population types X and Y, where $X, Y \in \{E, I\}$, the modeled neuronal activity is represented by the following variables:

- $p_X(t)$, measured in Hz, denoting the mean firing rate of neurons in the X population at time t,

- $q_{XY}(t)$, measured in Hz^2, denoting the covariance of the firing rates between the X and Y populations at time t,

- $w_X(t)$, measured in pA, denoting the mean adaptation current of neurons in the X population at time t,

- $r_{XY}^{Ext}(t)$, measured in Hz, denoting the average rate of spikes received by neurons of X population at time t through each of the afferent fibers arriving from external neurons of type Y. Although in general some fraction of the afferent fibers can arrive from external inhibitory neurons, throughout this paper we assume all these external input spikes to the network are received only from excitatory neurons.

The system of differential equations that governs the time evolution of the state variables p_X, q_{XY}, and w_X for a given external drive r_{XY}^{Ext} are provided by the master equations (13)–(15) below. Note that $q_{EI} = q_{IE}$, hence it is sufficient to solve (13)–(15) only for one of these two quantities. As stated above, these master equations require calculation of the transfer functions that relate the firing rate of neurons in each population to their presynaptic excitatory and inhibitory spike rates. The conductance-based internal interactions that yield the derivation of these transfer functions are modeled as follows [39, 40, 42].

Let K_{XY}^{Int}, with $X, Y \in \{E, I\}$, denote the average number of presynaptic connections that neurons within the X population in the network receive internally from the neurons of the Y population of the network. Similarly, let K_{XY}^{Ext} denote the average number of presynaptic connections that neurons within the X population receive from external neurons of type Y residing outside the network. Then, the average number of presynaptic connections that neurons within the X population receive in total from both internal and external neurons of type Y is given as

$$K_{XY} := K_{XY}^{Int} + K_{XY}^{Ext}, \qquad X, Y \in \{E, I\}. \tag{1}$$

As stated above, throughout this paper we assume all external cortical connections to the local network under the study are of excitatory type, that is, $K_{XI}^{Ext} = 0$, $X \in \{E, I\}$. Moreover, letting P_{XY} denote the connection probabilities between neurons of X and Y populations, as described in Table 1, the average number of internal connections are given as

$$K_{XY}^{Int} = P_{XY}N_Y, \qquad X, Y \in \{E, I\}. \tag{2}$$

Next, let r_{XE} and r_{XI} denote, respectively, the average rate of excitatory and inhibitory presynaptic spikes that neurons in a population of type X receive both internally from other neurons within the network and externally from neurons residing outside the network. That is,

$$r_{\mathrm{XE}}\left(p_{\mathrm{E}}, p_{\mathrm{I}}, r_{\mathrm{XE}}^{\mathrm{Ext}}\right) = \frac{1}{\mathrm{K}_{\mathrm{XE}}}\left[\mathrm{K}_{\mathrm{XE}}^{\mathrm{Int}}\, p_{\mathrm{E}} + \mathrm{K}_{\mathrm{XE}}^{\mathrm{Ext}}\, r_{\mathrm{XE}}^{\mathrm{Ext}}\right], \qquad \mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}, \tag{3}$$

$$r_{\mathrm{XI}}\left(p_{\mathrm{E}}, p_{\mathrm{I}}, r_{\mathrm{XI}}^{\mathrm{Ext}}\right) = \frac{1}{\mathrm{K}_{\mathrm{XI}}}\left[\mathrm{K}_{\mathrm{XI}}^{\mathrm{Int}}\, p_{\mathrm{I}} + \mathrm{K}_{\mathrm{XI}}^{\mathrm{Ext}}\, r_{\mathrm{XI}}^{\mathrm{Ext}}\right] = p_{\mathrm{I}}, \qquad \mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}, \tag{4}$$

where the second equality for r_{XI} is due to the assumption $\mathrm{K}_{\mathrm{XI}}^{\mathrm{Ext}} = 0$.

The train of spikes received by a neuron from its presynaptic neurons dynamically change the neuron's membrane conductance. Let $M_{G_{\mathrm{XE}}^{\mathrm{Syn}}}$ and $M_{G_{\mathrm{XI}}^{\mathrm{Syn}}}$ denote, respectively, the mean value of the total excitatory and inhibitory synaptic conductances of a neuron of type X. Note that, throughout the paper, we will use M_A and S_A to denote the mappings that give the mean and standard deviation of the quantity A, respectively. The total excitatory/inhibitory synaptic conductance of a neuron of type X is the conductance resulting cumulatively from all the synapses that are made on this neuron by other excitatory/inhibitory neurons in the network. The effective value of these synaptic components depend on the rate of input spikes to the neurons. As in [45] and [40], the mean conductances can approximately be given as

$$M_{G_{\mathrm{XE}}^{\mathrm{Syn}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}) = \mathrm{K}_{\mathrm{XE}}\mathrm{Q}_{\mathrm{XE}}^{\mathrm{Syn}}\tau_{\mathrm{XE}}^{\mathrm{Syn}}r_{\mathrm{XE}}, \qquad \mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}, \tag{5}$$

$$M_{G_{\mathrm{XI}}^{\mathrm{Syn}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}) = \mathrm{K}_{\mathrm{XI}}\mathrm{Q}_{\mathrm{XI}}^{\mathrm{Syn}}\tau_{\mathrm{XI}}^{\mathrm{Syn}}r_{\mathrm{XI}}, \qquad \mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}, \tag{6}$$

which then allow for the calculation of $M_{G_{\mathrm{X}}}$, the mean value of the total membrane conductance of the neurons within the X population, as

$$M_{G_{\mathrm{X}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}) = M_{G_{\mathrm{XE}}^{\mathrm{Syn}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}) + M_{G_{\mathrm{XI}}^{\mathrm{Syn}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}) + \mathrm{G}_{\mathrm{X}}^{\mathrm{Leak}}, \qquad \mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}. \tag{7}$$

The parameter $\mathrm{G}_{\mathrm{X}}^{\mathrm{Leak}}$ in (7) denotes the leak conductance of the neurons. The parameters $\mathrm{Q}_{\mathrm{XY}}^{\mathrm{Syn}}$ and $\tau_{\mathrm{XY}}^{\mathrm{Syn}}$ in (5) and (6), with $\mathrm{X}, \mathrm{Y} \in \{\mathrm{E}, \mathrm{I}\}$, denote the quantal conductance and decay time constant of the synaptic connections, as described in Table 1. Note that these synaptic parameters correspond to the first-order model of synaptic conductance kinetics given by equation (18) below. Note also that neurons typically receive multiple synapses per single connection. The parameters $\mathrm{Q}_{\mathrm{XY}}^{\mathrm{Syn}}$ in (5) and (6) denote synaptic quantal conductances *per connection*, and hence they are equal to the sum of the per-synapse quantal conductances generated by each of the synapses of a neuronal connection.

An approximation of the mean membrane potential of the X population based on the mean conductances calculated above has been provided in [45] and [40] as

$$M_{V_{\mathrm{X}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}, w_{\mathrm{X}}) = \frac{1}{M_{G_{\mathrm{X}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}})}\Big[\mathrm{V}_{\mathrm{E}}^{\mathrm{Syn}}M_{G_{\mathrm{XE}}^{\mathrm{Syn}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}) + \mathrm{V}_{\mathrm{I}}^{\mathrm{Syn}}M_{G_{\mathrm{XI}}^{\mathrm{Syn}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}})$$
$$+ \mathrm{V}_{\mathrm{X}}^{\mathrm{Leak}}\mathrm{G}_{\mathrm{X}}^{\mathrm{Leak}} - w_{\mathrm{X}}\Big], \qquad \mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}, \tag{8}$$

which can be roughly understood as a steady-state current law on the different currents flowing through the membrane of a neuron, with the neuron's membrane potential and conductances being replaced with their mean values across the population. The parameters $\mathrm{V}_{\mathrm{E}}^{\mathrm{Syn}}$ and $\mathrm{V}_{\mathrm{I}}^{\mathrm{Syn}}$ denote the excitatory and inhibitory synaptic reversal potentials as described

in Table 1. We assume these parameters do not depend on the type of the postsynaptic neurons.

The following approximations for the standard deviation $S_{V_{\mathrm{x}}}$ and a global autocorrelation time constant (approximate speed) $T_{V_{\mathrm{x}}}$ of membrane potential fluctuations have been obtained in [39] and [40] as

$$S_{V_{\mathrm{x}}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}}) = \left[\sum_{\mathrm{H} \in \{\mathrm{E}, \mathrm{I}\}} \frac{\mathrm{K}_{\mathrm{xH}} r_{\mathrm{xH}} \left[\tau_{\mathrm{xH}}^{\mathrm{Syn}} Z_{\mathrm{xH}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}}) \right]^2}{2 \left[\hat{\tau}_{\mathrm{x}}^{\mathrm{Mem}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}) + \tau_{\mathrm{xH}}^{\mathrm{Syn}} \right]} \right]^{\frac{1}{2}}, \qquad \mathrm{x} \in \{\mathrm{E}, \mathrm{I}\} \qquad (9)$$

$$T_{V_{\mathrm{x}}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}}) = \frac{\sum_{\mathrm{H} \in \{\mathrm{E}, \mathrm{I}\}} \mathrm{K}_{\mathrm{xH}} r_{\mathrm{xH}} \left[\tau_{\mathrm{xH}}^{\mathrm{Syn}} Z_{\mathrm{xH}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}}) \right]^2}{2 S_{V_{\mathrm{x}}}^2(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}})}, \qquad \mathrm{x} \in \{\mathrm{E}, \mathrm{I}\}. \qquad (10)$$

where

$$Z_{\mathrm{xH}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}}) := \frac{\mathrm{Q}_{\mathrm{xH}}^{\mathrm{Syn}}}{M_{G_{\mathrm{x}}}(r_{\mathrm{xE}}, r_{\mathrm{xI}})} \left[\mathrm{V}_{\mathrm{H}}^{\mathrm{Syn}} - M_{V_{\mathrm{x}}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}}) \right], \qquad \mathrm{x}, \mathrm{H} \in \{\mathrm{E}, \mathrm{I}\},$$

and $\hat{\tau}_{\mathrm{x}}^{\mathrm{Mem}}$ denotes the effective membrane time constant of the x population, given by

$$\hat{\tau}_{\mathrm{x}}^{\mathrm{Mem}} := \frac{\mathrm{C}_{\mathrm{x}}^{\mathrm{Mem}}}{M_{G_{\mathrm{x}}}(r_{\mathrm{xE}}, r_{\mathrm{xI}})}, \qquad \mathrm{x} \in \{\mathrm{E}, \mathrm{I}\}.$$

The parameter $\mathrm{C}_{\mathrm{x}}^{\mathrm{Mem}}$ denotes the membrane capacitance of the neurons as described in Table 1.

The transfer functions F_{x} of neurons within each population x can now be characterized using the membrane potential moments $M_{V_{\mathrm{x}}}$, $S_{V_{\mathrm{x}}}$, and $T_{V_{\mathrm{x}}}$ given by (8), (9), and (10), respectively. For this, a semi-analytic form has been proposed in [39] as

$$F_{\mathrm{x}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}}) = \frac{1}{2 T_{V_{\mathrm{x}}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}})} \, \mathrm{erfc} \left(\frac{\Theta_{\mathrm{x}}(M_{V_{\mathrm{x}}}, S_{V_{\mathrm{x}}}, \tilde{T}_{V_{\mathrm{x}}}) - M_{V_{\mathrm{x}}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}})}{\sqrt{2} S_{V_{\mathrm{x}}}(r_{\mathrm{xE}}, r_{\mathrm{xI}}, w_{\mathrm{x}})} \right),$$
$$\mathrm{x} \in \{\mathrm{E}, \mathrm{I}\}, \qquad (11)$$

where erfc is the complementary error function and $\tilde{T}_{V_{\mathrm{x}}} := T_{V_{\mathrm{x}}} / \tau_{\mathrm{x}}^{\mathrm{Mem}}$, with $\tau_{\mathrm{x}}^{\mathrm{Mem}} := \mathrm{C}_{\mathrm{x}}^{\mathrm{Mem}} / \mathrm{G}_{\mathrm{x}}^{\mathrm{Leak}}$ denoting the membrane time constant of the neurons of type x.

The effective membrane potential threshold Θ_{x} in (11), which accounts for the nonlinearities in the neuronal dynamics, is expressed in [41] as the following second-degree polynomial

$$\Theta_{\mathrm{x}}(M_{V_{\mathrm{x}}}, S_{V_{\mathrm{x}}}, \tilde{T}_{V_{\mathrm{x}}}) = \theta_{\mathrm{x}}^0 + \sum_{A \in \mathcal{D}} \theta_{\mathrm{x}}^{(A)} \left(\frac{A - \overline{A}}{\Delta A} \right)$$
$$+ \sum_{(A, B) \in \mathcal{D}^2} \theta_{\mathrm{x}}^{(A, B)} \left(\frac{A - \overline{A}}{\Delta A} \right) \left(\frac{B - \overline{B}}{\Delta B} \right), \qquad (12)$$

where $\mathcal{D} := \left\{ M_{V_{\mathrm{x}}}, S_{V_{\mathrm{x}}}, \tilde{T}_{V_{\mathrm{x}}} \right\}$ and $\mathcal{D}^2 := (\mathcal{D} \times \mathcal{D}) \setminus \left\{ (S_{V_{\mathrm{x}}}, M_{V_{\mathrm{x}}}), (\tilde{T}_{V_{\mathrm{x}}}, M_{V_{\mathrm{x}}}), (\tilde{T}_{V_{\mathrm{x}}}, S_{V_{\mathrm{x}}}) \right\}$. The normalization parameters $\overline{A}$ and ΔA, $A \in \mathcal{D}$, in (12) are not population type specific and their values are given in Table 2. It should be noted that some of these parameter values appear to be misreported in [41]. Here, we used their values as originally given in [39]. The coefficients θ_{x}^0, $\theta_{\mathrm{x}}^{(A)}$, and $\theta_{\mathrm{x}}^{(A, B)}$, $A \in \mathcal{D}$, $(A, B) \in \mathcal{D}^2$, are given in Table 3. They are calculated in [41] by fitting the effective thresholds and the resulting transfers function to numerically

calculated dynamics of adaptive exponential integrate and fire (AdEx) neurons. Note that the fit parameters in Table 3 are provided for both regular-spiking and fast-spiking neurons. Unless otherwise stated, we assume throughout this paper that all excitatory neurons are regular-spiking and all inhibitory neurons are fast-spiking.

Finally, calculation of the semi-analytic transfer functions (11) allows us to complete the presentation of the mean-field model by providing the masters equations that govern the evolution of the population-level neuronal activity in the network. For $X \in \{E, I\}$, $Y \in \{E, I\}$, and $t \in [0, T]$, $T > 0$, the equations are given as

$$
T^{\text{Mod}} \frac{dp_X}{dt} = F_X(r_{XE}, r_{XI}, w_X) - p_X
$$
$$
+ \frac{1}{2} \sum_{J \in \{E,I\}} \sum_{J' \in \{E,I\}} q_{JJ'} \frac{\partial^2}{\partial p_J \partial p_{J'}} F_X(r_{XE}, r_{XI}, w_X), \tag{13}
$$

$$
T^{\text{Mod}} \frac{dq_{XY}}{dt} = \left(F_X(r_{XE}, r_{XI}, w_X) - p_X \right) \left(F_Y(r_{YE}, r_{YI}, w_Y) - p_Y \right)
$$
$$
+ \sum_{J \in \{E,I\}} \left[q_{YJ} \frac{\partial}{\partial p_J} F_X(r_{XE}, r_{XI}, w_X) + q_{XJ} \frac{\partial}{\partial p_J} F_Y(r_{YE}, r_{YI}, w_X) \right]
$$
$$
- 2q_{XY} + \delta_{XY} \frac{1}{N_X} \left(\frac{1}{T^{\text{Mod}}} - F_Y(r_{YE}, r_{YI}, w_X) \right) F_X(r_{XE}, r_{XI}, w_X), \tag{14}
$$

$$
\tau_{w_X} \frac{dw_X}{dt} = -w_X + \tau_{w_X} \gamma_X p_X + \eta_X \left[M_{V_X}(r_{EE}, r_{EI}, w_X) - V_X^{\text{Leak}} \right], \tag{15}
$$

where T^{Mod} is the modeling time scale of the Markovian description of the network activity considered in [38], and τ_{w_X}, γ_X, and η_X are adaptation parameters of the neurons as described in Table 1. Note that, for simplicity of the exposition, the dependence of variables p_X, q_{XY}, and w_X on t is not explicitly shown in (13)–(15). Moreover, note that r_{XY} depends on the variables p_E, p_I and the external inputs r_{XY}^{Ext}, as in (3) and (4).

Equations (13) and (14), which give the temporal evolution of the mean and covariance of the firing rates have been developed in [38]. The additional equation (15), which captures the dynamics of sub-threshold and spike-triggered neuronal adaptation corresponding to the adaptation equation of a single-neuron AdEx model [46], has been provided in the set of equations in [41]. Note that (15) includes a correction on the original equation [41, Equ. (2.6)], as this equation suffers from a unit mismatch between the physical quantities and does not correctly correspond to the adaptation equation of the network's constitutive AdEx neurons.

We numerically solve (13)–(15) and analyze bifurcations in the equilibrium solutions of these equations to obtain the solution curves and bifurcation diagrams presented in the Results section. The distinction between the dynamics of the excitatory and inhibitory neuronal populations is made through inclusion or exclusion of the adaptation equation (15) and corresponding choice of the fit parameters given in Table 3. We consider all excitatory neurons to be regular-spiking, meaning that their dynamics undergoes adaptation. Unless otherwise stated, we consider all inhibitory neurons to be fast-spiking, with no adaptation. Therefore, for the fast-spiking inhibitory population we set $w_I = 0$ and exclude (15) for $X = I$ from the numerical computations.

In the numerical analysis presented in the Results section, we also investigate dynamic variations in the mean value of the synaptic currents flowing through the membrane of the neurons in each population. We use the approximate steady-state current law (8) to derive

the following approximations:

$$M_{I_{\mathrm{XE}}^{\mathrm{Syn}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}, w_{\mathrm{X}}) = (\mathrm{V}_{\mathrm{E}}^{\mathrm{Syn}} - M_{V_{\mathrm{X}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}, w_{\mathrm{X}}))M_{G_{\mathrm{XE}}^{\mathrm{Syn}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}), \qquad \mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}, \qquad (16)$$

$$M_{I_{\mathrm{XI}}^{\mathrm{Syn}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}, w_{\mathrm{X}}) = (\mathrm{V}_{\mathrm{I}}^{\mathrm{Syn}} - M_{V_{\mathrm{X}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}, w_{\mathrm{X}}))M_{G_{\mathrm{XI}}^{\mathrm{Syn}}}(r_{\mathrm{XE}}, r_{\mathrm{XI}}), \qquad \mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}, \qquad (17)$$

where $M_{I_{\mathrm{XE}}^{\mathrm{Syn}}}$ and $M_{I_{\mathrm{XI}}^{\mathrm{Syn}}}$ denote the mean excitatory and inhibitory synaptic currents in the x population, respectively.

Mean-field model parameters

The description and value of the biophysical parameters of the mean-field model presented above are given in Table 1. The parameter values are chosen to be in the range of realistic parameter values reported in the literature for the mouse and rat cortex. With these parameter values, which hereafter we refer to as *baseline parameter values*, the dynamics of the model shows a proper balance of excitation and inhibition wherein the mean firing rates of the excitatory and inhibitory populations closely coincide with those of the rat neocortex in an asynchronous irregular regime—as observed through both *in vivo* measurements and biophysically detailed *in silico* reconstruction of the rat neocortical microcircuitry [37]. Correspondingly, we refer to this state of balanced activity in the network as *baseline balanced state*.

The values of the passive membrane parameters $\mathrm{C}_{\mathrm{X}}^{\mathrm{Mem}}$, $\mathrm{G}_{\mathrm{X}}^{\mathrm{Leak}}$, and $\mathrm{V}_{\mathrm{X}}^{\mathrm{Leak}}$, $\mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}$, are chosen to be approximately equal to the median values given in [43, Suplementary Table 2], with the membrane leak conductance being reciprocal to the membrane resistance given in there. The range of values provided in [43] are obtained by tuning the parameters of a family of generalized leaky integrate-and-fire models so that they reproduce the spiking activity of a large number of recorded neurons in the primary visual cortex of adult mouse. The associated neuronal recordings are available in the Allen Cell Types Database [47].

The values of the adaptation parameters $\tau_{w_{\mathrm{E}}}$, η_{E}, and γ_{E} given in Table 1 are equal to the values chosen in [41] for these parameters. These values are used in [41] in the calculations resulting in the fit parameters given in Table 3 for regular-spiking excitatory neurons. Note that, as in [41], we assume all inhibitory neurons in the baseline model are non-adapting. Moreover, note that the values of the adaption parameters chosen here are also comparable to the values provided in [46] by fitting an AdEx model to dynamics of a biophysically detailed conductance-based model of a regular-spiking neuron.

The values of the synaptic reversal potentials given in Table 1 are fairly typical. We choose the baseline values for the other synaptic parameters to be equal to their mean values as given in [37]. The parameter values provided in [37] are used for a detailed digital reconstruction of the juvenile rat somatosensory microcircuitry as part of the Blue Brain Project [48]. Specifically, we set the values of $\tau_{\mathrm{XY}}^{\mathrm{Syn}}$, $\mathrm{X}, \mathrm{Y} \in \{\mathrm{E}, \mathrm{I}\}$, equal to the values given in [37, Table S6]. For synaptic quantal conductances, however, an adjustment is made on the values provided in [37]. The average quantal conductance value of 0.85 nS stated in [37, Page 471] for excitatory synapses, and the average value of 0.84 nS stated for inhibitory synapses, are specified as per-synapse conductances. Whereas, as stated in the model description above, the parameters $\mathrm{Q}_{\mathrm{XY}}^{\mathrm{Syn}}$ in the mean-field model we use here are associated with synaptic quantal conductances per connection. Therefore, to make adjustment for this difference, we scale the average quantal conductances in [37] by the average number of synapses per connection. These scaling values are also provided in [37, Page 464], as 3.6 synapses per connection for

Table 1: **Cellular, synaptic, and structural parameters of the mean-field and spiking network models.**

	Parameter	Description	Value	Unit
Cellular	C_E^{Mem}	Membrane capacitance of the excitatory neurons	110	pF
	C_I^{Mem}	Membrane capacitance of the inhibitory neurons	65	pF
	G_E^{Leak}	Leak conductance of the excitatory neurons	6	nS
	G_I^{Leak}	Leak conductance of the inhibitory neurons	5	nS
	V_E^{Leak}	Leak reversal potential of the excitatory neurons	-75	mV
	V_I^{Leak}	Leak reversal potential of the inhibitory neurons	-72	mV
	τ_{w_E}	Adaptation time constant of the excitatory neurons	500	ms
	τ_{w_I}	Adaptation time constant of the inhibitory neurons	500	ms
	η_E	Subthreshold adaptation current of the excitatory neurons	4	nS
	η_I	Subthreshold adaptation current of the inhibitory neurons	0	nS
	γ_E	Spike-triggered adaptation current increment of the excitatory neurons	60	pA
	γ_I	Spike-triggered adaptation current increment of the inhibitory neurons	0	pA
Synaptic	V_E^{Syn}	Reversal potential of the excitatory synapses	0	mV
	V_I^{Syn}	Reversal potential of the inhibitory synapses	-80	mV
	Q_{EE}^{Syn}	Quantal conductance of the excitatory-to-excitatory synaptic connections	3*	nS
	Q_{IE}^{Syn}	Quantal conductance of the excitatory-to-inhibitory synaptic connections	3*	nS
	Q_{EI}^{Syn}	Quantal conductance of the inhibitory-to-excitatory synaptic connections	12*	nS
	Q_{II}^{Syn}	Quantal conductance of the inhibitory-to-inhibitory synaptic connections	12*	nS
	τ_{EE}^{Syn}	Decay time constant of the excitatory-to-excitatory synapses	1.7	ms
	τ_{IE}^{Syn}	Decay time constant of the excitatory-to-inhibitory synapses	1.7	ms
	τ_{EI}^{Syn}	Decay time constant of the inhibitory-to-excitatory synapses	8.3	ms
	τ_{II}^{Syn}	Decay time constant of the inhibitory-to-inhibitory synapses	8.3	ms
Structural	N_E	Number of excitatory neurons in the network	8700	—
	N_I	Number of inhibitory neurons in the network	1300	—
	P_{EE}	Probability of excitatory-to-excitatory cortical connections	0.05	—
	P_{IE}	Probability of excitatory-to-inhibitory cortical connections	0.05	—
	P_{EI}	Probability of inhibitory-to-excitatory cortical connections	0.05	—
	P_{II}	Probability of inhibitory-to-inhibitory cortical connections	0.05	—
	K_{EE}^{Ext}	Number of external excitatory connections made to excitatory neurons	1200	—
	K_{IE}^{Ext}	Number of external excitatory connections made to inhibitory neurons	1200	—
	K_{EI}^{Ext}	Number of external inhibitory connections made to excitatory neurons	0	—
	K_{II}^{Ext}	Number of external inhibitory connections made to inhibitory neurons	0	—
	T^{Mod}	Time scale of the mean-field model	20	ms
Spiking Network	V_E^{Thr}	Spiking threshold potential of the excitatory neurons	-50	mV
	V_I^{Thr}	Spiking threshold potential of the inhibitory neurons	-50	mV
	Λ_E	Sharpness factor of the spike initiation term in AdEx model of excitatory neurons	2	mV
	Λ_I	Sharpness factor of the spike initiation term in AdEx model of inhibitory neurons	0.5	mV
	T_E^{Ref}	Refractory period of the excitatory neurons	5	ms
	T_I^{Ref}	Refractory period of the inhibitory neurons	5	ms
	N_E^{Ext}	Number of external excitatory input channels	1000	—
	P_{EE}^{Ext}	Probability of connections between external input channels and excitatory neurons	0.05	—
	P_{IE}^{Ext}	Probability of connections between external input channels and inhibitory neurons	0.05	—

*The values of synaptic quantal conductances given here are quantal conductances per connection. These values are calculated using the estimates provided in [37] as follows: $Q_{EE}^{Syn} = Q_{IE}^{Syn} = 3$ nS is approximately equal to the per-synapse excitatory quantal conductance of 0.85 nS times 3.6 synapses per excitatory connection, and $Q_{EI}^{Syn} = Q_{II}^{Syn} = 12$ nS is approximately equal to the per-synapse inhibitory quantal conductance of 0.84 nS times 13.9 synapses per inhibitory connection.

excitatory connections and 13.9 synapses per connection for inhibitory neurons. The adjusted conductances are then given in Table 1 as values of Q_{XY}^{Syn}, $X, Y \in \{E, I\}$.

Table 2: **Normalization parameters of the semi-analytic transfer functions [39].**

Cell type	$\overline{M_{V_{\mathrm{x}}}}$	$\overline{S_{V_{\mathrm{x}}}}$	$\overline{\tilde{T}_{V_{\mathrm{x}}}}$	$\Delta M_{V_{\mathrm{x}}}$	$\Delta S_{V_{\mathrm{x}}}$	$\Delta \tilde{T}_{V_{\mathrm{x}}}$
X = E, I	-60 mV	4 mV	0.5	10 mV	6 mV	1

We consider a randomly connected network composed of a total number of N = 10000 neurons, out of which $\mathrm{N_E}$ = 8700 neurons are excitatory and the remaining $\mathrm{N_I}$ = 1300 are inhibitory. This relatively large size of the network allows for the mean field approximation described above, and is also comparable to the size of the neuronal populations in layers 2/3, 5, and 6 of the rat neocortical microcircuitry investigated in [37]. The 87% excitatory versus 13% inhibitory proportions of the neurons are chosen according to the overall estimates provided in [37, Page 461].

Choosing the parameter values for internal and external connectivity requires some considerations and simulation-based adjustments. The local connectivity density of neuronal networks varies across cortical regions and layers. Moreover, more than 80% of synapses in a local network of nearly 200 micrometer in diameter come from external neurons residing outside of the network [37,49]. As a result, it is estimated that when a slice of cortical tissue with a typical thickness of 300 micrometer is cut from the cortex, only about 10% of excitatory synapses and about 38% of inhibitory synapses remain intact [49]. Therefore, the balance of excitation and inhibition in such cortical slice is largely deviated toward over-inhibition. Taking these into account, we choose the connectivity parameters of the mean-field model so that the resulting network, driven by a biologically reasonable rate of background spikes from external neurons, presents a balance of excitation and inhibition with mean excitatory and inhibitory firing rates being consistent with those measured in a detailed digital reconstruction of the rat cortical microcircuitry in an asynchronous irregular spiking regime [37].

We set the connection probability of all types of internal network connections to be equal to 0.05. For excitatory-to-excitatory connections, this value is quite comparable to the values reported in the literature for different cortical layers [37,50,51]. For other types of connections, this value appears to be almost half the typical values estimated experimentally. However, experimental estimations of neuronal connectivity are often obtained using cortical slices. Hence—due to the uneven reduction in the number of excitatory and inhibitory connections during slicing, as described above—using such estimates of inhibitory connection probabilities for a structurally simplified network such as the one we consider here can potentially result in an over-inhibited network. Our preliminary simulations of the mean-field model with larger values of inhibitory connection probabilities shows an imbalance of activity as expected, with the model requiring excessive amount of external excitatory drive in order to produce a reasonable firing activity. Therefore, to achieve a reasonable balance of activity, we choose inhibitory connection probabilities smaller than the experimentally obtained estimates.

We assume neurons of the network do not receive any external inhibitory inputs, whereas they each receive an average number of $\mathrm{K_{EE}^{Ext}} = \mathrm{K_{IE}^{Ext}} = 1200$ excitatory connections from external neurons. This number of external inputs is comparable with the estimates provided in [51, Table 3]. Moreover, with this number of external connections, along with the population size and internal connection probability chosen above, each neuron receives more than 60% of its synapses from external neurons, a percentage comparable to the estimates given in [37] and [49] as discussed above. Additionally, with the chosen number of external connections and internal connectivity, and with an average external spiking rate of $r_{\mathrm{EE}}^{\mathrm{Ext}} = r_{\mathrm{IE}}^{\mathrm{Ext}} = 1$ Hz, the numerical simulation of the model results in excitatory and inhibitory

Table 3: **Fit parameters of the semi-analytic transfer functions for populations of regular-spiking (RS) and fast-spiking (FS) neurons [41].**

Cell type	θ_x^0	$\theta_x^{(M_{V_x})}$	$\theta_x^{(S_{V_x})}$	$\theta_x^{(\bar{T}_{V_x})}$	$\theta_x^{(M_{V_x},M_{V_x})}$	$\theta_x^{(S_{V_x},S_{V_x})}$	$\theta_x^{(\bar{T}_{V_x},\bar{T}_{V_x})}$	$\theta_x^{(M_{V_x},S_{V_x})}$	$\theta_x^{(M_{V_x},\bar{T}_{V_x})}$	$\theta_x^{(S_{V_x},\bar{T}_{V_x})}$
X = E (RS)	-49.8	5.06	-25	1.4	-0.41	10.5	-36	7.4	1.2	-40.7
X = I (FS)	-51.4	4	-8.3	0.2	-0.5	1.4	-14.6	4.5	2.8	-15.3

firing rates which are very close to those obtained in reconstructed rat microcircuitry [37]. The details of the simulations are provided in the Results section.

Finally, note that in order for the Markovian assumption used in the derivation of the mean-field model to be valid, the time scale of the model, T^{Mod}, must be small enough so that neurons fire a maximum of one spike at each Markovian time step of length T^{Mod}, and must be large enough so that the network dynamics can be considered memoryless over the timescale of T^{Mod}; see [40–42]. Here, we set the value of T^{Mod} to be approximately equal to the time constant of the neurons, as suggested in [41] and [42]. This completes the discussion of the parameter values for the mean-field model. The rest of the parameters given at the bottom of Table 1 belong to the spiking network model described below.

Spiking neuronal network model

We construct a network of randomly connected AdEx spiking neurons, in direct correspondence to the mean-field description presented above. For this, first let $\mathcal{N}_E \subset \mathbb{N}$ and $\mathcal{N}_I \subset \mathbb{N}$, such that $\mathcal{N}_E \cap \mathcal{N}_I = \varnothing$, be two ordered sets that index neurons of the excitatory and inhibitory populations, respectively. Note that $\mathcal{N}_E$ is of cardinality N_E, and $\mathcal{N}_I$ is of cardinality N_I. The total number of neurons in the network is then indexed by the set $\mathcal{N} := \mathcal{N}_E \cup \mathcal{N}_I$. Also, let $\mathcal{N}_E^{\text{Ext}} \subset \mathbb{N}$, such that $\mathcal{N}_E^{\text{Ext}} \cap \mathcal{N} = \varnothing$, be an ordered set that indexes the external excitatory neurons that project onto at least one neuron inside the network. Note that, similar to the mean-field description, we assume no external inhibitory inputs to the network. With $n \in \mathcal{N}$, the neuronal activity of the spiking network is then represented by the following variables:

- $v_n(t)$, measured in mV, denoting the membrane potential of the nth neuron at time t,

- $w_n(t)$, measured in pA, denoting the adaptation current of the nth neuron at time t.

A neuron in the network fires a spike at a time $t = t^\star$ when its membrane potential exceeds its spiking threshold potential denoted by V_n^{Thr}, that is $v_n(t^\star) > V_n^{\text{Thr}}$. Let $\mathscr{S}_n^t$, $n \in \mathcal{N}$, denote a set that stores all spike times of the nth neuron up to time t. Let, moreover, $\mathscr{P}_m^t$, $m \in \mathcal{N}_E^{\text{Ext}}$, denote a set that stores all spike times of the mth excitatory external neuron up to time t. As described below, we assume these external spike times are Poission-distributed.

Now, let $g_{nm}(t)$ denote the membrane synaptic conductance of the nth postsynaptic neuron generated through its synaptic connection with the mth presynaptic neuron. Let the set $\{c_{nm}\}_{n \in \mathcal{N}, m \in \mathcal{N} \cup \mathcal{N}_E^{\text{Ext}}}$ capture the connectivity of the network, including connections from external neurons, such that $c_{nm} = 1$ if there is a connection from the mth neuron to the nth neuron, and $c_{nm} = 0$ otherwise. Then, approximating kinetics of the synaptic conductance at each presynaptic spike time $t^\star$ by an instantaneous rise to a peak (quantal) conductance Q_{nm}^{Syn} followed by an exponential decay with time constant τ_{nm}^{Syn}, the membrane

synaptic conductances are given as

$$g_{nm}(t) = \begin{cases} \sum_{t^\star \in \mathscr{T}_{nm}^t} \mathrm{Q}_{nm}^{\mathrm{Syn}} \exp\left(-\frac{t-t^\star}{\tau_{nm}^{\mathrm{Syn}}}\right) H(t-t^\star) & \text{if } c_{nm} = 1, \\ 0 & \text{if } c_{nm} = 0, \end{cases} \tag{18}$$

where H denotes the Heaviside step function and

$$\mathscr{T}_{nm}^t = \begin{cases} \mathscr{S}_m^t & \text{if } c_{nm} = 1 \text{ and } m \in \mathcal{N}, \\ \mathscr{P}_m^t & \text{if } c_{nm} = 1 \text{ and } m \in \mathcal{N}_{\mathrm{E}}^{\mathrm{Ext}} \\ \varnothing & \text{if } c_{nm} = 0. \end{cases}$$

Next, the total synaptic current I_n^{Syn} to the nth postsynaptic neuron can be computed as a function of v_n and t, by adding together all fractions of current coming from each individual synapse that the neuron receives. Let I_n^{Syn} be decomposed into its excitatory and inhibitory components as $I_n^{\mathrm{Syn}} = I_{n,\mathrm{E}}^{\mathrm{Syn}} + I_{n,\mathrm{I}}^{\mathrm{Syn}}$. Then,

$$I_{n,\mathrm{E}}^{\mathrm{Syn}}(v_n, t) = \sum_{m \in \mathcal{N}_{\mathrm{E}} \cup \mathcal{N}_{\mathrm{E}}^{\mathrm{Ext}}} g_{nm}(t)\left(\mathrm{V}_{nm}^{\mathrm{Syn}} - v_n\right), \tag{19}$$

$$I_{n,\mathrm{I}}^{\mathrm{Syn}}(v_n, t) = \sum_{m \in \mathcal{N}_{\mathrm{I}}} g_{nm}(t)\left(\mathrm{V}_{nm}^{\mathrm{Syn}} - v_n\right). \tag{20}$$

where $\mathrm{V}_{nm}^{\mathrm{Syn}}$ denotes the reversal potential of the synapse between the nth postsynaptic and mth presynaptic neurons when $c_{nm} = 1$.

Finally, representing each neuron by an AdEx model [46], the subthreshold activity of the network is given by the following system of differential equations for all $n \in \mathcal{N}$ and $t \in [0, T]$, $T > 0$,

$$\mathrm{C}_n^{\mathrm{Mem}} \frac{\mathrm{d}v_n}{\mathrm{d}t} = -\mathrm{G}_n^{\mathrm{Leak}}\left(v_n - \mathrm{V}_n^{\mathrm{Leak}}\right) + \mathrm{G}_n^{\mathrm{Leak}} \Lambda_n \exp\left(\frac{v_n - \mathrm{V}_n^{\mathrm{Thr}}}{\Lambda_n}\right) - w_n + I_n^{\mathrm{Syn}}(v_n, t), \tag{21}$$

$$\tau_{w_n} \frac{\mathrm{d}w_n}{\mathrm{d}t} = -w_n + \eta_n\left(v_n - \mathrm{V}_n^{\mathrm{Leak}}\right). \tag{22}$$

The parameters $\mathrm{C}_n^{\mathrm{Mem}}$ and $\mathrm{G}_n^{\mathrm{Leak}}$, and $\mathrm{V}_n^{\mathrm{Leak}}$ in (21) denote the membrane capacitance, leak conductance, and leak reversal potential of the nth neuron, respectively. Moreover, $\mathrm{V}_n^{\mathrm{Thr}}$ and Λ_n denote, respectively, the spiking threshold potential of the nth neuron and its sharpness factor for spike initiation. The parameters τ_{w_n} and η_n in (22) denote the adaptation time constant and the subthreshold adaptation current of the nth neuron, respectively.

The spiking activity of the network is captured by performing the following updates at each time instance $t = t^\star$ for all neurons that fire a spike at $t = t^\star$, that is, for all $n \in \mathcal{N}$ such that $v_n(t^\star) > \mathrm{V}_n^{\mathrm{Thr}}$:

- v_n is reset to the resting potential, denoted by $\mathrm{V}_n^{\mathrm{Rest}}$, and is kept at this value for a duration of time equal to the refractory period of the nth neuron, $\mathrm{T}_n^{\mathrm{Ref}}$,

- w_n is incremented by a constant amount γ_n,

- $t^\star$ is added to the set $\mathscr{S}_n^t$.

For the numerical analysis presented in the Results section, we set the parameters of all excitatory neurons and synapses, as well as all inhibitory neurons and synapses, to be the same, and equal to the baseline parameter values given in Table 1. That is, for all $n \in \mathcal{N}_X$ and $X \in \{E, I\}$, we set $C_n^{Mem} = C_X^{Mem}$, $G_n^{Leak} = G_X^{Leak}$, $V_n^{Leak} = V_X^{Leak}$, $\tau_{w_n} = \tau_{w_X}$, $\eta_n = \eta_X$, and $\gamma_n = \gamma_X$, in accordance with the baseline parameters used for the mean-field model. We also set the specific parameters of AdEx models as $V_n^{Thr} = V_X^{Thr}$, $\Lambda_n = \Lambda_X$, and $T_n^{Ref} = T_X^{Ref}$, for all $n \in \mathcal{N}_X$ and $X \in \{E, I\}$. Note that all inhibitory neurons are considered to be non-adapting, that is, $\eta_n = 0$ and $\gamma_n = 0$ for all $n \in \mathcal{N}_I$. Similarly, for all synapses of type Y-to-X, where $X, Y \in \{E, I\}$, that means for all $n \in \mathcal{N}_X$ and $m \in \mathcal{N}_Y$, we set $Q_{nm}^{Syn} = Q_{XY}^{Syn}$, $\tau_{nm}^{Syn} = \tau_{XY}^{Syn}$, and $V_{nm}^{Syn} = V_Y^{Syn}$. Note that, similar to the mean-field model, we assume that the synaptic reversal potentials do not depend on the type of the postsynaptic neurons.

We generate the elements of the internal network connectivity, $\{c_{nm}\}_{n \in \mathcal{N}, m \in \mathcal{N}}$, using the same connection probabilities used for the mean-field model. That is, we set $c_{nm} = 1$ with probability P_{XY}, as given in Table 1, when $n \in \mathcal{N}_X$ and $m \in \mathcal{N}_Y$, with $X, Y \in \{E, I\}$. To set the elements of external connectivity, $\{c_{nm}\}_{n \in \mathcal{N}, m \in \mathcal{N}_E^{Ext}}$, as well as the cardinality of $\mathcal{N}_E^{Ext}$, denoted by N_E^{Ext}, we first note that an *in vivo* network of size $10,000$ neurons can receive over $100,000$ afferent fibers from neurons outside of the network. The activities of the external neurons, however, are correlated. To lower the computational cost of simulating the network, and also to approximately take into account the correlation between external inputs, we assume that our spiking network receives input from only 1000, but independently spiking, input channels. That is, we set $N_E^{Ext} = 1000$. These input channels are then randomly connected to the neurons inside the network, with a connection probability of P_{XE}^{Ext}, $X \in \{E, I\}$. Therefore, $c_{nm} = 1$ with probability P_{XE}^{Ext} when $n \in \mathcal{N}_X$ and $m \in \mathcal{N}_E^{Ext}$, with $X \in \{E, I\}$.

We assume that spikes in each external input channel arrive randomly at Poisson-distributed time instances. To be able to compare the activity of the two models, we adjust the rate of spiking in each channel so that the average excitatory drive to neurons of the spiking network model becomes equal to the average drive to the neurons of the mean-field model. For this, note that each neuron of type X in the mean-field model receives, on average, inputs from K_{XE}^{Ext} external excitatory neurons, each firing at an average rate of r_{XE}^{Ext} spikes per second. Correspondingly, each neuron in the spiking network receives inputs from an average number of $1000 \times P_{XE}^{Ext}$ channels. If $r_{EE}^{Ext}(t)$ is set equal, or linearly proportional to $r_{IE}^{Ext}(t)$ for all $t \in [0, T]$, then we can choose the values for P_{EE}^{Ext} and P_{IE}^{Ext} such that

$$\frac{K_{EE}^{Ext}}{1000 \, P_{EE}^{Ext}} r_{EE}^{Ext}(t) = \frac{K_{IE}^{Ext}}{1000 \, P_{IE}^{Ext}} r_{IE}^{Ext}(t) =: \rho(t), \qquad \text{for all } t \in [0, T]. \tag{23}$$

The scaling factors $K_{XE}^{Ext}/(1000 \, P_{XE}^{Ext})$, $X \in \{E, I\}$, in (23) make the required adjustments on the external drive to the mean-field model so that $\rho(t)$, as defined in (23), provides an equivalent drive to the spiking network. Therefore, we set the spike rate of each Poisson channel equal to $\rho(t)$. For the numerical analysis presented in the Results section, we set $r_{EE}^{Ext} = r_{IE}^{Ext}$ as in the mean-field model, and $P_{EE}^{Ext} = P_{IE}^{Ext} = 0.05$ as given in Table 1.

Results

The mean-field model (13)–(15) and the spiking network model (21)–(22), with biophysical parameter values given in Table 1, are used here to investigate how the dynamic balance of excitation and inhibition in cortical networks is affected by variations in some of the important physiological and structural factors of the network. We first show that the mean-field model with baseline parameter values is balanced with an activity rate typically observed

in asynchronous irregular regimes, and is very responsive to changes in external inputs. We then perform the numerical analysis described in the Materials and methods section and present how the network balance is affected by changes in key synaptic and structural parameters. We discuss the results and their biological implications in the Discussion section.

The baseline balanced state

In order to investigate the effect of different synaptic and network parameters on the overall balance of excitation and inhibition, as described in next sections, it is important to first establish a reference balanced state for the network. Here, we demonstrate that the choice of baseline biophysical parameter values discussed in the Materials and methods section results in such a baseline balanced state in the mean-field network activity. For this, we solve the mean-field equations (13)–(15) with the baseline parameter values given in Table 1. The transfer functions used in (13)–(15) are given by (11), with their arguments being computed using (1)–(10), (12), and the fit parameters given in Tables 2 and 3. We drive the network with constant excitatory inputs of mean frequency $r_{\mathrm{EE}}^{\mathrm{Ext}}(t) = r_{\mathrm{IE}}^{\mathrm{Ext}}(t) = 1$ Hz, which presents a background activity at a level typically observed in irregularly spiking excitatory neurons at a cortical resting state [37].

Simulation of the mean-field model with the baseline setup described above identifies a stable equilibrium in the activity of the network, at which the excitatory neurons fire at the mean rate $p_{\mathrm{E}} = 1.15$ Hz, and the inhibitory neurons fire at the mean rate $p_{\mathrm{I}} = 5.71$ Hz. These rates of activity are close to the average excitatory firing rate of 1.09 Hz and inhibitory firing rate of 6.00 Hz obtained in a detailed simulation of the rat neocortical microcircuitry, when the constructed network is presenting balanced activity in an asynchronous irregular regime [37, Figure 17]; see also [36].

It is shown in the literature that excitatory and inhibitory synaptic conductances in the intact neocortex are well balanced and proportional to each other [3]. Using (5) and (6), the mean value of excitatory and inhibitory synaptic conductances at the network equilibrium described above are calculated for the excitatory population as $M_{G_{\mathrm{EE}}^{\mathrm{Syn}}} = 8.7$ nS and $M_{G_{\mathrm{EI}}^{\mathrm{Syn}}} = 37$ nS, which are also equal to the values obtained for the inhibitory population. These mean conductance values are comparable to the experimentally measured values provided in [3]. They give an excitatory to inhibitory conductance ratio of $M_{G_{\mathrm{EE}}^{\mathrm{Syn}}}/M_{G_{\mathrm{EI}}^{\mathrm{Syn}}} = 0.235$, which is consistent with the experimental findings that imply inhibitory conductances are much larger than excitatory conductances [30, 52]. The results shown in [3], however, indicate a balanced conductance ratio of approximately equal to one. The discrepancy between these results, as hinted in [3], may be because of the dominance of inhibitory synapses that can bias the results of experimental measurements. As we stated before in our discussion on the choice of parameter values for synaptic quantal conductances, the average number of inhibitory synapses per connection in our model is approximately four times larger than the average number of excitatory synapses per connection. If we assume this difference in the number of synapses proportionally biases experimentally measured conductances by a factor four, then the ratio $M_{G_{\mathrm{EE}}^{\mathrm{Syn}}}/M_{G_{\mathrm{EI}}^{\mathrm{Syn}}} = 0.235$ that we obtained in our simulation should be scaled by a factor of four to be comparable to the approximate ratio of 1 reported in [3]. We avoid such adjustment though, and in our analysis provided in the next sections we consider the ratio $M_{G_{\mathrm{EE}}^{\mathrm{Syn}}}/M_{G_{\mathrm{EI}}^{\mathrm{Syn}}} = 0.235$ as a reference for the steady-state value of the balanced conductance ratio.

Experimental observations suggest that the dynamic balance of excitation and inhibition in local cortical networks keeps the neurons of the network in a depolarized state near their

Figure 1: **Steady-state mean-field activity with respect to variations in the mean frequency of the external inputs.** All parameter values of the mean-field model are set to their baseline values given in Table 1. The model is driven by external inputs of different mean frequency $r_{\mathrm{EE}}^{\mathrm{Ext}} = r_{\mathrm{IE}}^{\mathrm{Ext}}$, and the resulting steady-state values of different network quantities are shown in the graphs. The points marked by dots in the graphs correspond to the baseline mean input frequency of 1 Hz. (a) Mean excitatory firing rate p_{E}, mean inhibitory firing rate p_{I}, ratio between the mean firing rates $p_{\mathrm{E}}/p_{\mathrm{I}}$, and the mean excitatory adaptation current w_{E}. (b) Excitatory membrane potential V_{E} and inhibitory membrane potential V_{I} of the neurons. Solid lines indicate the mean values $M_{V_{\mathrm{E}}}$ and $M_{V_{\mathrm{I}}}$ of the membrane potentials, and shaded areas indicate variations in the membrane potentials within a range of one-standard deviation ($S_{V_{\mathrm{E}}}$ and $S_{V_{\mathrm{I}}}$) from the mean values. (c) Mean synaptic conductances $M_{G_{\mathrm{EX}}^{\mathrm{Syn}}}$, $\mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}$ of the excitatory population, and the ratio between the two conductances. Mean synaptic conductances of the inhibitory population take the same values as those of the neurons of the excitatory population.

firing threshold, so that the network can be rapidly activated by external excitatory inputs and become involved in specific computational tasks [3, 53] To ensure that the baseline balanced state in the mean-field model indeed corresponds to such a state of highly responsive network activity, we simulate the model with the same baseline parameter values as before, but with different values for the constant mean frequency of the excitatory inputs, $r_{\mathrm{EE}}^{\mathrm{Ext}} = r_{\mathrm{IE}}^{\mathrm{Ext}}$. The resulting steady-sate values for different biophysical quantities of the model, obtained at the stable equilibrium of the equations for each input frequency value, are shown in Figure 1.

It can be seen in Figure 1 that all biophysical quantities of the model, such as the mean firing rates, mean excitatory adaptation current, mean value and standard deviation of membrane potentials, and mean synaptic conductances take biologically reasonable values as the input frequency varies over a wide ranges of values. The variation profile of the mean firing rates p_{E} and p_{I} shown in Figure 1a indicates that the overall activity of the neurons at the baseline background input frequency of 1 Hz is indeed close ot the firing threshold of the neurons, and the network is sufficiently responsive to rises in the frequency of excitatory inputs.

The inhibitory activity in the model rises in parallel to the excitatory activity as the external input frequency increases to larger values. However, the results shown in Figure 1 imply that the neuronal interactions throughout the network control the overall level of inhibition in the network at a level that still allows for an elevated level of overall excitation—which is necessary for the network to be able to perform the processing task in demand. This results in a change in the overall balance of excitation and inhibition toward higher excitation, as observed through the increase in the ratio between mean excitatory and inhibitory synaptic conductances, shown in Figure 1c. Nevertheless, the level of inhibition in the network remains sufficiently strong to prevent network instability and hyperactivity when the network receives an excessive amount of excitatory inputs from other cortical regions.

The observations made above suggest that the mean-field model with the baseline parameter values and background external inputs of mean frequency 1 Hz presents a well-balanced state of overall excitatory and inhibitory activity—at which the network stays in an asynchronous irregular regime and is highly responsive to external cortical inputs, without undergoing internal instabilities when the level of external excitation increases. Therefore, we choose this state as the baseline balanced state of the mean-field model, and use it as a reference state to study how such a balanced state is disturbed by changes in the values of physiological and structural parameters of the network.

Synaptic contributors to the balance of excitation and inhibition

The kinetics of synaptic activity in the conductance-based model we use here is governed by three main physiological factors, namely, synaptic decay time constants, synaptic quantal conductances, and synaptic reversal potentials. Variations in these physiological factors directly change the efficacy of synaptic communications between the neurons and hence has substantial impact on the dynamic balance of excitation and inhibition across the network. We investigate such impacts by showing how the steady-state balance between excitatory and inhibitory synaptic conductances and currents—obtained at the stable equilibrium of the baseline mean-field activity described above—is affected by variations in each of these synaptic factors. In particular, we identify critical states of imbalanced activity which result in the loss of stability of the network equilibrium and lead to a transition of the network dynamics to an oscillatory regime.

Dynamics of the mean-field model (13)–(15) is highly dependent on the profiles of the transfer functions F_E and F_I, which can change significantly if the physiological parameters of the synapses change. Hence, demonstrating the effects of variations in synaptic parameters on the profile of transfer functions (neuronal response curves) helps our understanding of how such variations affect the balance of mean-field activity in the network. Figure 2 illustrates how neuronal response curves vary with respect to changes in each of the three synaptic parameters we consider here. Variations in the decay time constants of inhibitory synapses, as shown in Figure 2a, changes the gain (sensitivity) of both inhibitory and excitatory neurons by changing the slope of their response curves. Additionally, such variations also change the excitability of the neurons by horizontally shifting their response curves. Similar effects are observed when the decay time constants of excitatory neurons change, however, with changes in neuronal excitability being less pronounced in this case. Figure 2b shows that changes in synaptic quantal conductances similarly affect gain and excitability of the neurons, with a higher sensitivity of the response curves being observed with respect to variations in excitatory quantal conductances. Changes in the synaptic reversal potentials, as shown in Figure 2c, significantly alter the gain of the neurons but have a lesser impact on their excitability.

Figure 2: **Variations in the response curves of neurons as a result of changes in synaptic parameters.** For all graphs, the mean inhibitory spike rates received by both populations are fixed at the typical value $r_{\mathrm{EI}} = r_{\mathrm{II}} = 6$ Hz. The mean excitatory spike rate received by each population is varied over a plausible range of values to obtain each neuronal response curves F_{X}, $\mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}$, which are calculated using (11) and the fit parameters given in Table 3. All parameter values involved in the calculation of F_{X}, except those specified on each graph, take their baseline values as given in Table 1. Thick curves in each graph show the response curves obtained at baseline synaptic parameter values. Other curves in each graph illustrate variations in the shape of the response curves as a synaptic parameter changes. Arrows indicate variations corresponding to 10 evenly distributed incremental changes in the parameter values specified in each graph. (a) Response curves with respect to variations in inhibitory (shown on the left side of the panel) and excitatory (shown on the right side of the panel) synaptic decay time constants. The value of $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$ is varied from 5 ms to 18 ms, and the value of $\tau_{\mathrm{IE}}^{\mathrm{Syn}} = \tau_{\mathrm{EE}}^{\mathrm{Syn}}$ is varied from 1 ms to 3 ms. (b) Response curves with respect to variations in inhibitory and excitatory synaptic quantal conductances. The value of $\mathrm{Q}_{\mathrm{EI}}^{\mathrm{Syn}} = \mathrm{Q}_{\mathrm{II}}^{\mathrm{Syn}}$ is varied from 1 nS to 25 nS, and the value of $\mathrm{Q}_{\mathrm{IE}}^{\mathrm{Syn}} = \mathrm{Q}_{\mathrm{EE}}^{\mathrm{Syn}}$ is varied from 1 nS to 8 nS. (c) Response curves with respect to variations in inhibitory and excitatory synaptic reversal potentials. The value of $\mathrm{V}_{\mathrm{I}}^{\mathrm{Syn}}$ is varied from -103 mV to -53 mV, and the value of $\mathrm{V}_{\mathrm{E}}^{\mathrm{Syn}}$ is varied from -30 mV to 20 mV.

Changes in the gain of the neurons when the excitatory synaptic reversal potentials are increased or decreased from their baseline values are pronounced, and occur monotonically.

However, gain changes with respect to variations in inhibitory reversal potentials appear to be non-monotonic. At lower output frequency values, both increasing $V_{\mathrm{I}}^{\mathrm{Syn}}$ above its baseline value, and decreasing it below its baseline value, result in an increase in the gain of the neurons.

Local inhibitory sub-networks are known to play a key role in stabilizing the dynamics of local networks and coordinating the flow of activity across cortical areas [8, 12, 31, 32, 35, 54]. Therefore, in what follows, we initially perform our analysis based on codimension-one continuation of the baseline equilibrium state with respect to variations in each of the inhibitory synaptic parameters. Then, we extend the analysis to codimension-two by additionally considering variations in excitatory synaptic parameters.

Effect of synaptic decay time constants

The decay time constant of a synapse determines how long the activity initiated in the synapse by an incoming action potential (spike) will last. Hence, the decay time constant has a significant impact on the efficacy of the synapse, which can also be directly implied from the mean synaptic conductance equations (5) and (6). Moreover, changes in synaptic decay time constants also changes the standard deviation and autocorrelation time constants of membrane potential fluctuations, as implied from (9) and (10). As a result, the transfer functions (response curves) of the neurons is highly impacted by variations in the decay time constants, which is confirmed by the results shown in Figure 2a. To show how the decay time constants of the synapses then impact the global dynamics of the network, and how they contribute to maintaining or disturbing the overall balance of excitation and inhibition, we first continue the stable equilibrium of the baseline balanced network with respect to variations in inhibitory decay time constants $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$. The results are shown in Figure 3.

The response curves given in Figure 2a show that increasing the value of $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$ decreases the gain and excitability of inhibitory and excitatory neurons. As a result, the mean firing rates of both excitatory and inhibitory populations decrease, concurrently, with increasing $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$, as shown in Figure 3a. This paired variation in the mean firing activity of the populations, however, does not necessarily imply that the overall balance of excitation and inhibition remains unchanged with respect to changes in inhibitory decay time constants. The ratio of the mean excitatory to mean inhibitory synaptic conductances in the excitatory population, $M_{G_{\mathrm{EE}}^{\mathrm{Syn}}}/M_{G_{\mathrm{EI}}^{\mathrm{Syn}}}$, shown in Figure 3b, implies that the excitation-inhibition balance moves toward over-inhibition as the efficacy of inhibitory synapses grows with increasing $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$. Indeed, unlike the profile of the inhibitory firing rate p_{I}, at large values of $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$ the mean inhibitory synaptic conductance $M_{G_{\mathrm{EI}}^{\mathrm{Syn}}}$ shown in Figure 3b rises slightly as $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$ increases. It should be noted that mean synaptic conductances in the inhibitory population, $M_{G_{\mathrm{IE}}^{\mathrm{Syn}}}$ and $M_{G_{\mathrm{II}}^{\mathrm{Syn}}}$, which are not shown in Figure 2a, take the same values as those in the excitatory population due to the same choices of connection probabilities in our model.

On the opposite direction, decreasing the value of inhibitory synaptic decay time constants increases the ratio $M_{G_{\mathrm{EE}}^{\mathrm{Syn}}}/M_{G_{\mathrm{EI}}^{\mathrm{Syn}}}$ and moves the excitation-inhibition balance toward more excitation. The excessive excitation in the network then drives the network's dynamics to a critical state, at which further reduction in $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$ results in an overall level of excitation that cannot be effectively balanced by the increased level of inhibition. At this critical state, the network's dynamics undergoes a phase transition to an oscillatory state, identified by a Hopf bifurcation point in the codimension-one continuations shown in Figure 3. The limit

(a) (b)

(c)

(d) (e) (f)

Figure 3: **Effect of variations in synaptic decay time constants on the long-term mean-field activity of the network.** All parameter values of the mean-field model, except for the synaptic decay time constants that are explicitly specified in each graph, are set to their baseline values given in Table 1. The model is driven by external inputs of constant mean frequency $r_{\mathrm{EE}}^{\mathrm{Ext}} = r_{\mathrm{IE}}^{\mathrm{Ext}} = 1$ Hz. In graphs (a)–(e), the decay time constants of inhibitory synapses made on neurons of both excitatory and inhibitory populations are set to take the same value, which is varied as the bifurcation parameter. The resulting codimension-one continuation of the network equilibrium and emerging limit cycles are shown for different quantities. Curves of equilibria are shown by solid lines, and the minimum and maximum values that each quantity takes on the limit cycles are shown by dotted lines. Dark-colored segments of each curve indicate stable equilibria/limit cycles, whereas light-colored segments indicate unstable equilibria/limit cycles. Unlabeled dots in each graph correspond to the baseline parameter values $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}} = 8.3$ ms. Points labeled by H are Hopf bifurcation points. We show bifurcation diagrams for (a) mean excitatory firing rate p_{E} and mean inhibitory firing rate p_{I}, (b) mean synaptic conductances $M_{G_{\mathrm{EX}}^{\mathrm{Syn}}}$, $\mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}$, of the excitatory population, and the ratio between the two conductances, (c) absolute value of the mean excitatory synaptic currents, $|M_{I_{\mathrm{XE}}^{\mathrm{Syn}}}|$, $\mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}$, as well as the ratio between mean excitatory and inhibitory synaptic currents, and (d) mean excitatory and inhibitory synaptic driving forces $|M_{v_{\mathrm{E}}} - \mathrm{V}_{\mathrm{X}}^{\mathrm{Syn}}|$, $\mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}$, in the excitatory population, as well as such driving forces $|M_{v_{\mathrm{I}}} - \mathrm{V}_{\mathrm{X}}^{\mathrm{Syn}}|$, $\mathrm{X} \in \{\mathrm{E}, \mathrm{I}\}$, in the inhibitory population. (e) Frequency of the limit cycles originating from the Hopf bifurcation point. (f) Codimension-two continuation of the Hopf bifurcation point when the excitatory synaptic decay time constants are also allowed to vary as a bifurcation parameter. The point marked by a dot has the baseline excitatory parameter value of $\tau_{\mathrm{EE}}^{\mathrm{Syn}} = \tau_{\mathrm{IE}}^{\mathrm{Syn}} = 1.7$ ms. Arrows indicate transition from stable (dark) to unstable (light) equilibria as the parameters are varied across the curve.

cycle originated from the Hopf bifurcation point has also been continued, and the curves of minimum and maximum values taken by different biophysical quantities of the model on the resulting cycles are shown in Figure 3. When the inhibitory synaptic decay time constants decrease below the critical value $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}} = 7.06$ ms, which is the value at which the Hopf bifurcation occurs, the previously stable equilibrium of the model becomes unstable. As a result, the orbits of the system depart the vicinity of the equilibrium and converge to a stable limit cycle on the curves of cycles originated from the Hopf bifurcation. Figure 3e shows that the stable oscillations on this cycle lie in the delta frequency band (1 - 4 Hz). It should be noted that further continuation of the curves of limit cycles, as partially shown in Figure 3, detects a fold bifurcation of limit cycles and emergence of unstable limit cycles at low values of $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$. However, analyzing the dynamics of the model at such values is not pertinent to the purpose of this paper. In fact at such values the mean-field model loses its validity in accurately predicting the emerging dynamic regimes, since the quantity under the square root in (9) approaches negative values.

The results shown in Figure 3d imply that the mean membrane potential of both excitatory and inhibitory populations decreases with increases in the value of $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$. This means that, on average, neurons in the network become more hyperpolarized when the level of inhibition in the network is enhanced by increasing the efficacy of the inhibitory synapses. Since in this study the values of synaptic reversal potentials are kept fixed, the reduction in the mean membrane potential of the neurons due to an increase in $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$ yields a reduction in the mean electrochemical driving force of the inhibitory synapses in both populations, $|M_{V_{\mathrm{x}}} - \mathrm{V}_{\mathrm{I}}^{\mathrm{Syn}}|$, $\mathrm{x} \in \{\mathrm{E}, \mathrm{I}\}$, and a rise in the mean driving force of the excitatory synapses, $|M_{V_{\mathrm{x}}} - \mathrm{V}_{\mathrm{E}}^{\mathrm{Syn}}|$, $\mathrm{x} \in \{\mathrm{E}, \mathrm{I}\}$. The curves of the equilibrium driving forces shown in Figure 3d illustrate such profiles of variations.

The absolute values of the mean synaptic currents at the equilibrium, calculated using (16) and (17), are shown in Figure 3c. As discussed above and illustrated in Figures 3b and 3d, the steady-state values of the two contributing components of these currents, the mean synaptic conductances and the mean synaptic driving forces, change in opposite directions with increases in $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$. The combination of these two components, however, results in steady-state mean synaptic currents which decrease in absolute value as $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$ is increased, as we can see in the curves of equilibria shown in Figure 3c; note that inhibitory currents are not shown separately in Figure 3c as their profile is similar to that of excitatory currents. In particular, despite the increase in mean driving forces of the excitatory synapses, the mean excitatory synaptic currents decrease as $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$ is increased. This implies that the effect of decreasing excitatory conductances, shown in Figure 3b, dominates the effect of increasing mean excitatory driving forces, hence resulting in the net decrease of excitatory synaptic currents.

Figure 3c also shows that the steady-state ratio between the absolute values of mean excitatory and mean inhibitory synaptic currents in the inhibitory population, $|M_{I_{\mathrm{IE}}^{\mathrm{Syn}}}|/|M_{I_{\mathrm{II}}^{\mathrm{Syn}}}|$, does not significantly change with respect to $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$, whereas the ratio between these currents in the excitatory population, $|M_{I_{\mathrm{EE}}^{\mathrm{Syn}}}|/|M_{I_{\mathrm{EI}}^{\mathrm{Syn}}}|$, decreases with increasing $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}}$. Therefore, similar to the ratio $M_{G_{\mathrm{EE}}^{\mathrm{Syn}}}/M_{G_{\mathrm{EI}}^{\mathrm{Syn}}}$ between synaptic conductances, here the ratio $|M_{I_{\mathrm{EE}}^{\mathrm{Syn}}}|/|M_{I_{\mathrm{EI}}^{\mathrm{Syn}}}|$ between synaptic currents also indicates shifts in the level of excitation-inhibition balance, toward over-inhibition or over-excitation.

The results provided above describe only the effect of inhibitory synaptic decay time constants on the network balance. Nevertheless, the excitation-inhibition balance in the network can also be affected by changes in the excitatory synaptic decay time constants. It is

implied from the the neuronal response curves shown in Figure 2a that changes in excitatory and inhibitory decay time constants have opposite impacts on the gain of neurons. Therefore, with fixed values of $\tau_{\text{EI}}^{\text{Syn}} = \tau_{\text{II}}^{\text{Syn}}$, the mean firing rates of both inhibitory and excitatory populations increase concurrently with increases in $\tau_{\text{EE}}^{\text{Syn}} = \tau_{\text{IE}}^{\text{Syn}}$. However, similar to changes in inhibitory decay time constants described above, it is not intuitively possible to accurately predict whether the network balance will be maintained during such paired changes in the mean firing rates of the populations; and if not, to which direction, either over-inhibition or over-excitation, the balance will shift. In fact, the relative changes between the magnitude of the mean excitatory and inhibitory response curves, and the varying sensitivity of these curves to changes in synaptic decay time constants—which results from complicated interactions in the network between adaptive and non-adaptive dynamic neurons—can result in non-intuitive changes in the network balance.

To demonstrate the joint effect of both excitatory and inhibitory synaptic decay time constants on network balance, we perform a codimension-two continuation of the Hopf bifurcation point that was detected in the codimension-one bifurcation analysis described before. We consider the same values for the decay time constants of excitatory synapses in both populations, $\tau_{\text{EE}}^{\text{Syn}} = \tau_{\text{IE}}^{\text{Syn}}$, and vary it as a second bifurcation parameter. The resulting curve of Hopf points is shown in Figure 3f.

It is observed that, when $\tau_{\text{EE}}^{\text{Syn}} = \tau_{\text{IE}}^{\text{Syn}}$ is reduced from its baseline value, the critical transition point (Hopf bifurcation) in the dynamics of the network occurs at lower values of $\tau_{\text{EI}}^{\text{Syn}} = \tau_{\text{II}}^{\text{Syn}}$. This implies that lower values of excitatory synaptic decay time constants reduce the overall level of excitation in the network, so that this reduction in excitation effectively compensates for a rise in excitation caused by decreases in inhibitory decay time constants $\tau_{\text{EI}}^{\text{Syn}} = \tau_{\text{II}}^{\text{Syn}}$. Therefore, due to this compensatory effect, lower values of excitatory synaptic decay time constants allow for the network to maintain a stable balanced state at much lower values of inhibitory decay time constants.

The upper part of the curve of Hopf points in Figure 3f shows a less intuitive dynamics for larger values of $\tau_{\text{EE}}^{\text{Syn}} = \tau_{\text{IE}}^{\text{Syn}}$ than its (approximately) baseline value. Although not shown here, codimension-one continuations of the baseline equilibrium with respect to variations in $\tau_{\text{EE}}^{\text{Syn}} = \tau_{\text{IE}}^{\text{Syn}}$, but with fixed values of $\tau_{\text{EI}}^{\text{Syn}} = \tau_{\text{II}}^{\text{Syn}}$, still show an increase in the level of excitation at higher values of $\tau_{\text{EE}}^{\text{Syn}} = \tau_{\text{IE}}^{\text{Syn}}$ and a shift in the balance of excitation and inhibition to hyper-excitation. However, as it can be seen in Figure 3f, at large values of inhibitory synaptic decay time constants, including the baseline value $\tau_{\text{EI}}^{\text{Syn}} = \tau_{\text{II}}^{\text{Syn}} = 8.3$ ms, this shift to hyper-excitation does not drive the network's dynamics to an oscillatory regime. Both an oscillatory state and a non-oscillatory hyper-excited state can be observed in the network activity when $\tau_{\text{EE}}^{\text{Syn}} = \tau_{\text{IE}}^{\text{Syn}}$ is increased from small values to large values while $\tau_{\text{EI}}^{\text{Syn}} = \tau_{\text{II}}^{\text{Syn}}$ is at about $4 - 7$ ms. It should be noted, however, that the level of stable hyper-excitation observed at large values of excitatory synaptic decay time constants corresponds to conductance ratios $M_{G_{\text{EE}}^{\text{Syn}}}/M_{G_{\text{EI}}^{\text{Syn}}}$ that may not be biologically plausible.

Importantly, the curve of Hopf bifurcation points in Figure 3f implies that the baseline parameter values $\tau_{\text{EE}}^{\text{Syn}} = \tau_{\text{IE}}^{\text{Syn}} = 1.7$ ms and $\tau_{\text{EI}}^{\text{Syn}} = \tau_{\text{II}}^{\text{Syn}} = 8.3$ ms are critical values for network stability, in the sense that the network remains quite responsive at these values, while relatively small reduction in the mean decay time constant of inhibitory neurons transitions the network's dynamics to an oscillatory state.

Effect of synaptic quantal conductances

The quantal conductance of a synapse is the peak membrane conductance change in a neuron caused by receiving a single spike from a presynaptic neuron. Therefore, changes in the quantal conductance of a synapse directly modulate the strength of the synapse. This is also implied from the mean synaptic conductance equations (5) and (6). It should be noted that, as described in the Materials and methods section, the quantal conductances Q_{XY}, $X, Y \in \{E, I\}$, that are incorporated in the mean-field model we use here are in fact quantal conductances per connection, that means, they are equal to the quantal conductances per synapse times the number of synapses per connection. As a result, Q_{XY}, $X, Y \in \{E, I\}$, in our study can be altered both by changes in the number of synapses and by modulations of the synaptic peak conductances.

The mean synaptic conductances (5) and (6) change with synaptic quantal conductances in the same way as they change with synaptic decay time constants. However, standard deviation and autocorrelation time constants of membrane potential fluctuations change differently with synaptic quantal conductances. In general, similar to the effect of increases in $\tau_{EI}^{Syn} = \tau_{II}^{Syn}$, the response curves given in Figure 2b show that the gain and excitability of both inhibitory and excitatory neurons decrease with increasing $Q_{EI}^{Syn} = Q_{II}^{Syn}$. Therefore, we expect that the effects on network balance caused by varying synaptic quantal conductances to be similar to the effects caused by changing synaptic decay time constants. This can be seen in Figure 4, which shows bifurcation diagrams of key biophysical quantities with inhibitory quantal conductances $Q_{EI}^{Syn} = Q_{II}^{Syn}$ varied as the bifurcation parameter.

In a rather similar way to what we observed in Figure 3 when $\tau_{EI}^{Syn} = \tau_{II}^{Syn}$ was changed, we also see here that when the value of inhibitory quantal conductances $Q_{EI}^{Syn} = Q_{II}^{Syn}$ increases the following changes occur in the steady-state values of different quantities of the model: the mean firing rates of both excitatory and inhibitory populations decrease concurrently; the ratio of mean excitatory to mean inhibitory synaptic conductances decreases; the absolute value of mean synaptic currents decreases; the ratio of the absolute value of mean excitatory to the absolute value of mean inhibitory synaptic currents in the excitatory population decreases; the mean electrochemical driving force of the inhibitory synapses in both populations decreases; and the mean driving force of the excitatory synapses increases. As a result, the overall excitation-inhibition balance in the network shifts toward over-inhibition as the inhibitory synapses are strengthened by increasing $\tau_{EI}^{Syn} = \tau_{II}^{Syn}$.

In contrast to the high sensitivity of the network's stability to reductions in decay time constants of the inhibitory neurons, the bifurcation diagrams of Figure 4 imply that the stability of the network's equilibrium is relatively robust to decreases in the value of inhibitory quantal conductances. Comparing the curves of equilibrium values for the ratios $M_{G_{EE}^{Syn}}/M_{G_{EI}^{Syn}}$ and $|M_{I_{EE}^{Syn}}|/|M_{I_{EI}^{Syn}}|$ shown here in Figures 4b and 4c, respectively, with those shown in Figures 3b and 3c, we can see that significantly higher level of overall excitation can be maintained in the network at low values of $Q_{EI}^{Syn} = Q_{II}^{Syn}$ before a transition in the network's dynamics to an oscillatory regime occurs. However, the loss of stability of the network's equilibrium at very low values of $Q_{EI}^{Syn} = Q_{II}^{Syn}$ appears to be more critical. Stable delta-band oscillations can exists only for very short range of values of $Q_{EI}^{Syn} = Q_{II}^{Syn}$.

The neuronal response curves given in Figure 2b show that changes in excitatory and inhibitory synaptic quantal conductances have opposite impacts on the gain and excitability of the neurons. Unlike the non-intuitive dynamics that we observed in Figure 3f when decay time constants of the synapses were jointly changed, the curve of Hopf bifurcation points shown in Figure 4f, when both $Q_{EI}^{Syn} = Q_{II}^{Syn}$ and $Q_{EE}^{Syn} = Q_{IE}^{Syn}$ are free to change, shows an

Figure 4: **Effect of variations in synaptic quantal conductances on the long-term mean-field activity of the network.** All parameter values of the mean-field model, except for the synaptic quantal conductances that are explicitly specified in each graph, are set to their baseline values given in Table 1. The model is driven by external inputs of constant mean frequency $r_{\mathrm{EE}}^{\mathrm{Ext}} = r_{\mathrm{IE}}^{\mathrm{Ext}} = 1$ Hz. The same description as given in Figure 3 holds for the graphs presented here, with the only difference being that here the bifurcation parameter for codimension-one continuation is the inhibitory quantal conductances $Q_{\mathrm{EI}}^{\mathrm{Syn}} = Q_{\mathrm{II}}^{\mathrm{Syn}}$, with the baseline value of 12 nS, and the additional bifurcation parameter for codimension-two continuation is the excitatory quantal conductances $Q_{\mathrm{EE}}^{\mathrm{Syn}} = Q_{\mathrm{IE}}^{\mathrm{Syn}}$, with the baseline value of 3 nS.

intuitively expected dynamics. Increasing the value of excitatory quantal conductances results in a rise in the level of excitation in the network, which then requires stronger inhibitory synapses for maintaining the network stability and avoiding transitions to oscillatory states. As a result, the Hopf bifurcation point occurs at higher values of $Q_{\mathrm{EI}}^{\mathrm{Syn}} = Q_{\mathrm{II}}^{\mathrm{Syn}}$ when $Q_{\mathrm{EE}}^{\mathrm{Syn}} = Q_{\mathrm{IE}}^{\mathrm{Syn}}$ is increased. Moreover, with any fixed values of inhibitory quantal conductances, a phase transition occurs in the network's dynamics if the quantal conductances of excitatory synapses increase beyond a Hopf bifurcation value. However, the baseline parameter values $Q_{\mathrm{EE}}^{\mathrm{Syn}} = Q_{\mathrm{IE}}^{\mathrm{Syn}} = 3$ nS and $Q_{\mathrm{EI}}^{\mathrm{Syn}} = Q_{\mathrm{II}}^{\mathrm{Syn}} = 12$ nS appear to be significantly below the curve of Hopf bifurcation points in Figure 3f, implying that the network stability at the baseline state is sufficiently robust to changes in both inhibitory and excitatory synaptic quantal conductances.

Effect of synaptic reversal potentials

Changes in reversal potentials of synapses modulate the electrochemical driving force of the synapses and also directly affect the mean membrane potential of the neurons across the network, as implied from (8). It can also be seen through (9) and (10) that changes in synaptic reversal potentials also affect the standard deviation and autocorrelation time constants of neuronal membrane potential fluctuations. Therefore, the characteristic response curves of the neurons and the dynamics of overall excitation-inhibition balance in the network can be significantly affected by changes in the reversal potential of the synapses. To show such impacts on the global dynamics of the network, we perform bifurcation analysis with respect to variations in synaptic reversal potentials, as we have performed with respect to variations in the other synaptic parameters. That is, we first continue the baseline stable equilibrium of the network with respect to variations in the value of inhibitory reversal potential V_I^{Syn}, and then extend the results by performing a codimension-two continuation of the detected Hopf bifurcation points by additionally allowing variation of the excitatory reversal potential V_E^{Syn}. The results are shown in Figure 5.

The neuronal response curves in Figure 2 show that, within a wide range of plausible firing rate frequencies, the gain of inhibitory and excitatory neurons increases with both increasing and decreasing changes in the value of V_I^{Syn} from its baseline value, although the increase in the gain of inhibitory neurons caused by a reduction in V_I^{Syn} is less pronounced. This may suggest that the mean firing rate of the inhibitory and excitatory populations will increase with V_I^{Syn} changes in either directions. However, the curves of steady-state mean firing rates shown in Figure 5a do not confirm this intuitive prediction. Although the steady-state value of p_I increases when V_I^{Syn} decreases below its baseline value, as well as when V_I^{Syn} increases sufficiently above its baseline value, the steady-state value of p_E is strictly decreasing with increasing V_I^{Syn}. In particular, these profiles of variations in p_E and p_I have counter-intuitive relations with the steady-state values of the mean membrane potentials. Since V_E^{Syn} is fixed at the constant value of 0 mV in the codimension-one analysis presented here, the curves of excitatory driving forces shown in Figure 5d imply that, on average, both excitatory and inhibitory neuronal populations monotonically become more depolarized with increases in V_I^{Syn}, which is a pattern of variation not directly implied from the mean firing rate of the neurons.

An explanation for the network interactions that could result in the steady-state curves described above can be as follows. Although the neurons become further depolarized with increases in V_I^{Syn}, the driving forces of inhibitory synapses, $|M_{V_X} - V_I^{Syn}|$, $X \in \{E, I\}$, shown in Figure 5d keeps decreasing to very small values as V_I^{Syn} increases to large values. This results in significant reduction in inhibitory synaptic currents at large values of V_I^{Syn}, despite the rise in the firing rates of the inhibitory neurons due to their increased gain. As a result, a stable balance of excitatory and inhibitory synaptic currents can be achieved if the mean firing rate of excitatory neurons significantly reduces to near zero, as shown in Figure 5a. It should be noted, however, that in the studies presented here a constant level of excitation is always provided to all neurons of the network by the excitatory external inputs of constant frequency $r_{EE}^{Ext} = r_{IE}^{Ext} = 1$ Hz. In the opposite direction, when V_I^{Syn} decreases to low values, the mean driving force of inhibitory synapses increases to sufficiently large values, which allow for inhibitory synaptic currents that are large enough to be balanced by larger excitatory synaptic currents. Therefore, a stable balance of currents can be achieved at low values of V_I^{Syn}, with both p_E and p_I taking relatively large values due to the increased neuronal gains. This increase in the mean firing rates of the neurons with decreases in V_I^{Syn} to sufficiently low

Figure 5: **Effect of variations in synaptic reversal potentials on the long-term mean-field activity of the network.** All parameter values of the mean-field model, except for the synaptic reversal potentials that are explicitly specified in each graph, are set to their baseline values given in Table 1. The model is driven by external inputs of constant mean frequency $r_{\text{EE}}^{\text{Ext}} = r_{\text{IE}}^{\text{Ext}} = 1$ Hz. The same description as given in Figure 3 holds for the graphs presented here, with the only difference being that here the bifurcation parameter for codimension-one continuation is the inhibitory reversal potential $V_{\text{I}}^{\text{Syn}}$, with the baseline value of -80 mV, and the additional bifurcation parameter for codimension-two continuation is the excitatory reversal potential $V_{\text{E}}^{\text{Syn}}$, with the baseline value of 0 nS.

values can be seen in Figure 5a.

The steady-state value of the ratio $M_{G_{\text{EE}}^{\text{Syn}}}/M_{G_{\text{EI}}^{\text{Syn}}}$ between mean excitatory and inhibitory conductances, as shown in Figure 5b, decreases with increasing $V_{\text{I}}^{\text{Syn}}$. Consistent with the explanation provided above, this implies that the balance of excitation and inhibition in the network moves monotonically with $V_{\text{I}}^{\text{Syn}}$, to over-inhibition for large values of $V_{\text{I}}^{\text{Syn}}$, and to over-excitation for small values of $V_{\text{I}}^{\text{Syn}}$. However, unlike what we observed in Figures 3c and 4c for changes in synaptic decay time constants and quantal conductances, here the monotonic shift in the level of excitation-inhibition balance cannot be correctly implied from the ratio $|M_{I_{\text{EE}}^{\text{Syn}}}|/|M_{I_{\text{EI}}^{\text{Syn}}}|$ between synaptic currents in the excitatory population, as shown in Figure 5c.

The bifurcation diagrams of Figure 5 further reveal the presence of two Hopf bifurcation points. When $V_{\text{I}}^{\text{Syn}}$ decreases below the Hopf point denoted by green dots in Figure 5, the

network's dynamics transitions to a stable oscillatory behavior. Figure 5e shows that the frequency of these oscillations are in the delta band. However, the emergent delta oscillations here are of significantly lower magnitude than those arising from changes in inhibitory synaptic decay time constants, as shown in Figure 3. Moreover, the curve of stable limit cycles calculated here vanishes at a second Hopf point when V_I^{Syn} decreases further. This implies that reductions in the value of inhibitory synaptic reversal potentials is less critical to the stability of the network. In fact, a stable yet high level of excitation can be present in the network at low values of V_I^{Syn}.

Finally, codimension-two continuation of the two Hopf bifurcation points detected in the analysis described above implies that a qualitatively similar network dynamics is observed at a wide range of values of excitatory synaptic reversal potentials. This is shown in Figure 5f. Unlike changes with respect to V_I^{Syn}, the neuronal response curves in Figure 2c show monotonic changes in the gain of neurons with respect to V_E^{Syn}. When V_E^{Syn} decreases, the gain of neurons decreases significantly. The resulting reduction in the level of excitation then allows for further reduction in V_I^{Syn} before network transition to an oscillatory regime. Therefore, the two Hopf bifurcation points occur at lower values of V_I^{Syn} when V_E^{Syn} decreases.

Structural contributors to the balance of excitation and inhibition

The mean-field model we use in our study represents a network of randomly connected neurons with relatively sparse connectivity. The dynamics of such network can be significantly affected by the structural factors that determine the overall topology of the network. These factors include the relative number of excitatory and inhibitory neurons in the network and the density of connectivity between the neurons. By employing bifurcation analysis similar to our previous investigations on the effect of physiological parameters intrinsic to synapses and neurons, we analyze how the steady-state balance of excitation and inhibition obtained at the baseline network structure is altered by changes in each of the structural factors determining the overall level of recurrent neuronal interactions in the network.

Effect of the ratio between the number of inhibitory and excitatory neurons

Knowing the crucial role of inhibitory neurons in stabilizing the excitatory interactions across the network, the ratio between the total number of inhibitory and excitatory neurons is expected to have significant impacts on the excitation-inhibition balance. Here, with a fixed number of neurons N = 10000, we change the inhibitory proportion N_I/N of the total network population and observe how the steady-state mean-field activities in the network change accordingly. The resulting codimension-one bifurcation diagrams for different quantities of the model are shown in Figure 6.

Since N is fixed, a change in the ratio N_I/N alters both N_E and N_I. Therefore, considering (1) and (2), the mean values of both excitatory and inhibitory synaptic conductances given by (5) and (6), respectively, change with N_I/N. Standard deviation and autocorrelation time constant of membrane potential fluctuations in both excitatory and inhibitory populations are also directly affected by changes in N_I/N, as implied by (9), (10), (1), and (2). Moreover, a change in N_E and N_I directly modifies the dynamics of the covariance of firing rates, given by (14). Despite these different contributions in the equations of the model in comparison with those of the inhibitory synaptic decay time constants, the bifurcation diagrams obtained by changing N_I/N closely resemble the diagrams obtained by changing $\tau_{EI}^{Syn} = \tau_{II}^{Syn}$. This can be seen by comparing Figure 6 with Figure 3. Therefore, when N_I/N changes, the long-

Figure 6: **Effect of variations in the ratio between the number of inhibitory and excitatory neurons on the long-term mean-field activity of the network.** All parameter values of the mean-field model, except for the number of inhibitory and excitatory neurons, are set to their baseline values given in Table 1. The total number of neurons is fixed at N = 10000. The inhibitory proportion of neurons in the network, N_I/N, is varied over a reasonable range of values. The model is driven by external inputs of constant mean frequency $r_{EE}^{Ext} = r_{IE}^{Ext} = 1$ Hz. The same description as given for graphs (a)–(e) in Figure 3 holds for the graphs presented here, with the only difference being that here the bifurcation parameter for codimension-one continuation is N_I/N, with the baseline value of 0.13.

term dynamics of the network changes similarly to what we observed before as a result of changes in $\tau_{EI}^{Syn} = \tau_{II}^{Syn}$. In particular, increasing the inhibitory proportion of the total network population N_I/N from its baseline value moves the excitation-inhibition balance toward over-inhibition. Moreover, the stability of network activity appears to be critically sensitive to decreases in N_I/N, so that relatively small reduction in N_I/N causes a transition in network activity to stable oscillations in the delta frequency band.

The contribution of inhibitory neurons in regulating network activity is not only influenced by their relative population size, but also by their cellular properties and efficacy of the synapses that they make on other neurons, as we studied separately above. To see some of the joint effects of such structural and physiological factors on the stability of balanced activity in the network, we perform codimension-two continuation of the Hopf point detected in Figure 6 by additionally varying each of the inhibitory synaptic parameters as a second

| (a) | (b) | (c) |

Figure 7: **Joint effect of the synaptic parameters and the ratio between the number of inhibitory and excitatory neurons on the long-term mean-field activity of the network.** Codimension-two continuation of Hopf bifurcation points are shown in each graph. The inhibitory proportion of neurons in the network, N_I/N, is considered as one of the bifurcation parameters in all graphs. Different synaptic parameters are then chosen in each graph as the second bifurcation parameter. Arrows indicate transition from stable (dark) to unstable (light) equilibria as the parameters are varied across the curve. The points marked by dots are the same Hopf bifurcation points marked in Figures 3, 4, 5 and 6. (a) Curve of Hopf bifurcation points with decay time constant of inhibitory synapses chosen as the second bifurcation parameter. (b) Curve of Hopf bifurcation points with quantal conductance of inhibitory synapses chosen as the second bifurcation parameter. (c) Curve of Hopf bifurcation points with reversal potential of inhibitory synapses chosen as the second bifurcation parameter.

bifurcation parameter. The resulting curves of Hopf bifurcation points are shown in Figure 7.

The results discussed before showed that changes in $\tau_{EI}^{Syn} = \tau_{II}^{Syn}$ and N_I/N have very similar impacts on the network's dynamics. A decrease in $\tau_{EI}^{Syn} = \tau_{II}^{Syn}$ increases the level of excitation and causes the network to undergo a critical transition to an oscillatory dynamics at a Hopf bifurcation point. Consequently, larger numbers of inhibitory neurons relative to the number of excitatory neurons are required to maintain the stability of the network at lower values of $\tau_{EI}^{Syn} = \tau_{II}^{Syn}$. This explains the profile of the curve of Hopf points shown in Figure 7a. Disregarding the irregularly folded segment of the curve shown in Figure 7b—which can be due to the distortions observed in Figure 2b in the semi-analytically calculated response curves of excitatory neurons at low values of $Q_{EI}^{Syn} = Q_{II}^{Syn}$—this curve of Hopf bifurcation points implies a generally similar effect when $Q_{EI}^{Syn} = Q_{II}^{Syn}$ and N_I/N are jointly varied. That is, a larger ratio of N_I/N is required for the stability of balanced activity at lower values of $Q_{EI}^{Syn} = Q_{II}^{Syn}$. However, the sensitivity of the Hopf points with respect to changes in $Q_{EI}^{Syn} = Q_{II}^{Syn}$ and N_I/N appears to be particularly different. The critical (bifurcation) value of N_I/N, below which the network transitions to oscillatory behavior, changes only slightly with sufficiently large values of $Q_{EI}^{Syn} = Q_{II}^{Syn}$. However, this critical ratio appears to be highly sensitive to excessively low values of $Q_{EI}^{Syn} = Q_{II}^{Syn}$.

The curve of Hopf bifurcation points shown in Figure 7c implies that both high and low values of V_I^{Syn} allow for network stability even at very low values of N_I/N, although such stable activity at low values of V_I^{Syn} will correspond to a state of stable hyper-excitation. Moreover, it can be observed that the critical value of N_I/N is very sensitive to both increases and decreases in V_I^{Syn} from its baseline value of -80 mV. Therefore, relatively small changes in the baseline value of V_I^{Syn} can remove the oscillatory activity of the network caused by a

Figure 8: **Effect of variations in the overall network connectivity density on the long-term mean-field activity of the network.** All parameter values of the mean-field model, except for the connection probabilities, are set to their baseline values given in Table 1. The model is driven by external inputs of constant mean frequency $r_{\mathrm{EE}}^{\mathrm{Ext}} = r_{\mathrm{IE}}^{\mathrm{Ext}} = 1$ Hz. All connection probabilities are set to take the same value, which is varied as the bifurcation parameter over a biologically plausible range of values. The resulting codimension-one continuation of the network equilibrium is shown in the graphs for different network quantities. Curves of equilibria are shown by solid lines. No changes in the stability of the equilibria is detected. The points marked by dots in all graphs correspond to the baseline parameter values $\mathrm{P_{EE} = P_{IE} = P_{EI} = P_{II} = 0.05}$. We show bifurcation diagrams for (a) mean excitatory firing rate p_{E} and mean inhibitory firing rate p_{I}, (b) mean synaptic conductances $M_{G_{\mathrm{EX}}^{\mathrm{Syn}}}$, $\mathrm{X} \in \{\mathrm{E, I}\}$, of the excitatory population, and the ratio between the two conductances, (c) absolute value of the mean excitatory and inhibitory synaptic currents, $|M_{I_{\mathrm{EX}}^{\mathrm{Syn}}}|$, $\mathrm{X} \in \{\mathrm{E, I}\}$, in the excitatory population, as well as the ratio between the two currents, (d) mean excitatory and inhibitory synaptic driving forces $|M_{V_{\mathrm{E}}} - \mathrm{V}_{\mathrm{X}}^{\mathrm{Syn}}|$, $\mathrm{X} \in \{\mathrm{E, I}\}$, in the excitatory population, as well as such driving forces $|M_{V_{\mathrm{I}}} - \mathrm{V}_{\mathrm{X}}^{\mathrm{Syn}}|$, $\mathrm{X} \in \{\mathrm{E, I}\}$, in the inhibitory population.

shortage of the number of inhibitory neurons in the network.

Effect of the connectivity density

The random connectivity in the local network that we study here is relatively sparse, with baseline connection probabilities $\mathrm{P_{EE} = P_{IE} = P_{EI} = P_{II} = 0.05}$ between and within excitatory and inhibitory populations. To investigate how changes in the level of sparsity in network connectivity affects the dynamic balance of excitation and inhibition, we continue the stable baseline equilibrium of the network with respect to changes in the value of overall connectivity density $\mathrm{P_{EE} = P_{IE} = P_{EI} = P_{II}}$ as a codimension-one bifurcation parameter. The results are shown in Figure 8.

The mean firing rates of both excitatory and inhibitory populations decrease concurrently

with increases in the overall connectivity density, as shown in Figure 8a. The ratio of the mean excitatory to inhibitory conductances shown in Figure 8b indicates that the network's excitation-inhibition balance moves toward over-inhibition when the network's overall connectivity density increases. On the other hand, at low values of $P_{EE} = P_{IE} = P_{EI} = P_{II}$, which correspond to highly sparse network connectivity, the network operates at a non-oscillatory state of hyper-excitation without undergoing a phase transition to a slow oscillatory regime. This is despite the significant decrease in mean synaptic conductances and synaptic currents at low values of overall connectivity density in the network, as it can be seen in Figures 8b and 8c. In general, we observe high sensitivity of the network balance to increases in the sparsity of the network from its baseline level, wheres much less sensitivity is observed when overall network connectivity becomes denser. Moreover—unlike the other cases that we have studied—we notice that no sustained oscillatory behavior emerges in the mean field activity of the network when over-excitation occurs due to increases in the sparsity of the network connectivity.

Besides changes in the overall network connectivity, population-specific changes in connection probabilities also affect the stability and balance of activity across the network. Recurrent excitatory-to-excitatory connections are necessary for sustaining the activity of cortical networks when they are involved in performing memory or cognitive processing tasks. Inhibitory-to-excitatory connections are crucial for stabilizing and regulating the excitatory activity in the network, and excitatory-to-inhibitory connections provide inhibitory neurons with sufficient excitation they need for their operation. However, the reason for the presence of a substantial amount of recurrent inhibitory-to-inhibitory connections in local cortical networks [54, 55], and the functionality of this type of connectivity, is less intuitive to explain. These recurrent inhibitory connections provide a level of disinhibition in the network that can regulate the overall inhibitory activity and prevent over-inhibition. However, the disinhibition created through these inhibitory-to-inhibitory connections also reduces the effectiveness of the inhibitory neurons in stabilizing the activity of excitatory neurons. To disambiguate functional implications of inhibitory-to-inhibitory connectivity, we investigate how varying only the density of this type of recurrent connectivity impacts the network balance. We choose the probability of inhibitory-to-inhibitory connections, P_{II}, as a bifurcation parameter and continue the baseline equilibrium state of the network. The results are shown in Figure 9.

The steady-state mean firing rate curves shown in Figure 9a confirm a significant rise in the firing rate of inhibitory neurons when the level of disinhibition in the network decreases substantially at low values of P_{II}. The resulting intensive inhibition significantly hyperpolarizes the excitatory neurons, as seen in Figure 9d, and suppresses their firing activity. This shift to hyper-inhibition in the network balance reflects clearly on the steady-state ratios $M_{G_{EE}^{Syn}}/M_{G_{EI}^{Syn}}$ and $|M_{I_{EE}^{Syn}}|/|M_{I_{EI}^{Syn}}|$, as shown in Figures 9b and 9c. In the opposite direction, however, the excessive disinhibition that results from increasing P_{II} above its baseline value yields a relatively sharp increase in the level of excitation in the network. Excitatory neurons become exceedingly depolarized and their firing rates increases significantly. This substantial rise in the activity of excitatory neurons also depolarizes the inhibitory neurons and results in a concurrent rise in their firing rate. However, as implied from Figures 9a and 9d, the strong recurrent inhibition that is generated within the inhibitory population at large values of P_{II} significantly restricts the mean depolarization level and firing rate of inhibitory neurons. This restrains the inhibitory neurons from effectively controlling the excitatory activity in the network, so that the excitation-inhibition balance moves to excessive excitation with increases in

Figure 9: **Effect of variations in the inhibitory-to-inhibitory connection probability on the long-term mean-field activity of the network.** All parameter values of the mean-field model, except for the probability of connections between inhibitory neurons, are set to their baseline values given in Table 1. The model is driven by external inputs of constant mean frequency $r_{\mathrm{EE}}^{\mathrm{Ext}} = r_{\mathrm{IE}}^{\mathrm{Ext}} = 1$ Hz. The same description as given for graphs (a)–(e) in Figure 3 holds for the graphs presented here, with the only difference being that here the bifurcation parameter for codimension-one continuation is $\mathrm{P_{II}}$, with the baseline value of 0.05.

$\mathrm{P_{II}}$. The steady-state value of the ratio $M_{G_{\mathrm{EE}}^{\mathrm{Syn}}}/M_{G_{\mathrm{EI}}^{\mathrm{Syn}}}$ shown in Figure 9b precisely indicates such a shift in the network balance.

The bifurcation diagrams of Figure 9 further reveal that, as $\mathrm{P_{II}}$ increases from its baseline value, the network's dynamics undergoes several Hopf bifurcations, at which the stability of the network equilibrium switches and sustained oscillations emerge, or vanish. In particular, stable oscillations of delta-band frequency emerge when the density of recurrent inhibitory connectivity increases slightly above its baseline value. However, these oscillations disappear for moderately higher values of recurrent inhibitory connectivity, at which a non-oscillatory state of hyper-excitation is present across the network. The results also predict the emergence of very-high amplitude stable oscillatory activity at disproportionately large values of $\mathrm{P_{II}}$. However, it should be noted that, due to the Markovian assumption used in the derivation of the mean-field model, the accuracy of the mean-field model degrades at the high firing activity associated with these oscillations. Note also that, similar to the results presented before, the curves of limit cycles originating from each Hopf bifurcation points have been continued only

up to the point where the mean-field model remains mathematically valid. This numerical termination point in the continuation of limit cycles usually occurs at parameter values for which the quantity under the square root in (9) becomes negative.

Instantaneous correlation of excitation and inhibition

The results presented so far have been obtained based on the analysis of long-term dynamics of the mean-field model, which provides information on the overall level of excitation-inhibition balance throughout the network, and how such balance is affected by changes in each of the physiological and structural parameters of the network. In experimental studies, a balance between excitation and inhibition in local cortical networks is sometimes deduced from tight correlation between the instantaneous activities of the inhibitory and excitatory populations [6, 7, 10]. Therefore, here we investigate how two distinct states of network activity that we observed in the results presented above, namely, a stable balanced state and a (functionally imbalanced) oscillatory state, reflect on the instantaneous correlation between the mean firing rates of excitatory and inhibitory populations.

We use the results we obtained above when we analyzed the effect of synaptic decay time constants. We solve the equations of the mean-field model corresponding to two different states: the baseline balanced state with $\tau_{\text{EI}}^{\text{Syn}} = \tau_{\text{II}}^{\text{Syn}} = 8.3$ ms, and a high-amplitude slow oscillatory state with $\tau_{\text{EI}}^{\text{Syn}} = \tau_{\text{II}}^{\text{Syn}} = 6.5$ ms. Note that the emergence of stable delta-band oscillations in the later state is ensured, as an inhibitory synaptic decay time constant of 6.5 ms is smaller than the critical Hopf bifurcation value of 7.06 ms obtained in the bifurcation analysis results shown in Figure 3. Therefore, with a constant input drive of $r_{\text{EE}}^{\text{Ext}} = r_{\text{IE}}^{\text{Ext}} = 1$ Hz, the solutions of the model with $\tau_{\text{EI}}^{\text{Syn}} = \tau_{\text{II}}^{\text{Syn}} = 6.5$ ms oscillate on a limit cycle, as shown in the top graph in Figure 10b.

For each of the network states described above, we generate 50 different solution curves for p_{E} and p_{I} by driving the model with 50 different external excitatory inputs $r_{\text{EE}}^{\text{Ext}} = r_{\text{IE}}^{\text{Ext}}$. The temporal profile of these inputs is generated by adding random fluctuations of amplitude 0.4 Hz to the constant baseline inputs $r_{\text{EE}}^{\text{Ext}} = r_{\text{IE}}^{\text{Ext}} = 1$ Hz. The frequency bandwidth of these fluctuations ranges approximately from 1 to 20 Hz, as described in Figure 10. To remove the effect of initial transient responses of the network, we use only the last 2 seconds of the computed solution curves for our correlation analysis.

A sample temporal profile of p_{E} and p_{I} is shown in Figure 10 for each state, along with cross-correlations between p_{E} and p_{I} for all 50 pairs of simulated mean firing rates. In both states, the sample temporal curves show a tight instantaneous correlation between the mean excitatory and inhibitory firing rates. The cross-correlation curves confirm the presence of such tight correlations in all 50 firing rate profiles. The time lags between the mean excitatory and inhibitory firing rates in both states appear to be very small: 0.50 ± 1.08 ms at the baseline balanced state, and 0.35 ± 0.74 ms at the oscillatory state, meaning that on average the inhibitory firing activity is slightly ahead of the excitatory firing activity. It should be noted though, that these tight correlations between instantaneous excitatory and inhibitory activities exist at both of the two distinct stable and oscillatory network states, despite the essentially different dynamic behavior and conditions of overall excitation-inhibition balance that we observed at these states through our bifurcation analysis.

Figure 10: **Instantaneous correlation between mean-field excitatory and inhibitory activity at baseline (balanced) and oscillatory states.** At the top of each panel, instantaneous profiles of mean excitatory and inhibitory firing rates, p_E and p_I, computed for the mean-field model with the constant input drives $r_{EE}^{Ext} = r_{IE}^{Ext} = 1$ Hz, are shown over a simulation time interval of 5000 ms. All biophysical parameters of the model take their baseline values, except for τ_{EI}^{Syn} and τ_{II}^{Syn} which take different values in each panel. The bottom right graph of each panel shows cross-correlations between mean excitatory and inhibitory firing rates over the simulation time interval of $[3000, 5000]$ ms. In each of these graphs, 50 correlation curves are shown that are obtained from simulating the model with 50 different external input drives. These external drives are generated by adding random fluctuations of amplitude 0.4 to the baseline value of $r_{EE}^{Ext} = r_{IE}^{Ext} = 1$ Hz. The temporal patterns of the fluctuations are generated by low-pass filtering 50 stochastic signals, each of which taking uniformly distributed random values at each instances of time. The passbands of the low-pass filters take 50 different values in the range from 1 Hz to 20 Hz. In each cross-correlation graph, a sample curve corresponding to input fluctuations filtered at passband frequency of 10 Hz is highlighted. The instantaneous profiles of the mean firing rates corresponding to the highlighted cross-correlation curves are shown on the bottom left side of each panel. (a) Instantaneous correlation between excitatory and inhibitory firing rates at the baseline balanced state, with baseline values of inhibitory synaptic decay time constants $\tau_{EI}^{Syn} = \tau_{II}^{Syn} = 8.3$ ms. (b) Instantaneous correlation between excitatory and inhibitory firing rates at an oscillatory state emerged in the dynamics of the model when the values of inhibitory synaptic decay time constants are reduced to $\tau_{EI}^{Syn} = \tau_{II}^{Syn} = 6.5$ ms.

Balanced and oscillatory activity in spiking networks

We performed our extensive bifurcation analyses presented above by taking the advantage of the simplicity and computational tractability of the mean-field model, which allowed for investigating changes in the long-term dynamics of the underlying local cortical network over a wide range of variations in the key physiological and structural parameters of the network, thereby observing the impacts of these factors on the balance of excitation and inhibition in the network. However, the inevitable simplifying assumptions, such as a Markovian assumption, the and semi-analytic derivations that were used in developing the mean-field model we employed in our study can raise concerns about the reliability of our predictions made based on this model. To address such concerns, to some extent, we use the network of $10,000$ spiking AdEx neurons described in the Materials and methods section to partially verify the predictions made by the mean-field model.

Figure 11: **Instantaneous correlation between the average spiking activity of excitatory and inhibitory neurons at asynchronous irregular (baseline balanced) and oscillatory bursting states.** In each panel, a rastergram of the excitatory (red) and inhibitory (blue) spiking activity in the spiking neuronal network (21)–(22) is shown on the left. For visual clarity, only the activity of a randomly selected 10 percent subset of total neurons are illustrated. The biophysical parameters of the spiking network, and the Poisson distributed input spike trains, are set as described in the Materials and methods section, except for the inhibitory synaptic decay time constants τ_{nm}^{Syn}, $n \in \mathcal{N}$, $m \in \mathcal{N}_{\mathrm{I}}$ which take different values in each panel. The middle graph in each panel shows the average firing rate of the excitatory and inhibitory neurons over the last 2 seconds of the simulation time. The graph of cross-correlation between these instantaneous mean firing activities is shown on the right side of each panel. (a) Spontaneous activity and correlation between average excitatory and inhibitory firing rates at the baseline asynchronous irregular state with baseline values of inhibitory synaptic decay time constants, $\tau_{nm}^{\mathrm{Syn}} = 8.3$ ms, $n \in \mathcal{N}$, $m \in \mathcal{N}_{\mathrm{I}}$. (b) Spontaneous activity and correlation between average excitatory and inhibitory firing rates at an oscillatory bursting state emerged as a result of a reduction in the values of inhibitory synaptic decay time constants to $\tau_{nm}^{\mathrm{Syn}} = 6.5$ ms, $n \in \mathcal{N}$, $m \in \mathcal{N}_{\mathrm{I}}$.

Specifically, we test if the results we obtained from the mean-field model on the state transition induced by inhibitory synaptic decay time constants, as well as those obtained on instantaneous correlations, can be replicated by a sufficiently large network of spiking neurons. We simulate the spiking neuronal network (21)–(22) at the same two dynamic states described in the results shown in Figure 10, namely, the state of balanced asynchronous irregular activity with the baseline parameter values $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}} = 8.3$ ms, and the oscillatory state associated with reduced inhibitory decay time constants $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}} = 6.5$ ms. The rastergram of spiking activity at each state is shown in Figure 11 for 10 percent of inhibitory and excitatory neurons.

The rastergram shown in Figure 10a demonstrates asynchronous and irregular activity at the baseline state. The mean firing rates of the excitatory and inhibitory neurons, cal-

culated using the last 2 seconds of the spike trains in order to remove the effect of initial transient activity, are 1.13 Hz and 5.84 Hz, respectively. These rates of firing activity almost precisely match the mean firing rates $p_{\mathrm{E}} = 1.15$ Hz and $p_{\mathrm{I}} = 5.71$ Hz obtained at the baseline balanced equilibrium of the mean-field model. Consistent with the predictions made through bifurcation analysis of the mean-field model, the rastergram shown in Figure 11b demonstrates the emergence of a slow oscillatory bursting state in the activity of the spiking neurons when inhibitory decay time constants are reduced to $\tau_{\mathrm{EI}}^{\mathrm{Syn}} = \tau_{\mathrm{II}}^{\mathrm{Syn}} = 6.5$ ms, a value below the Hopf bifurcation value shown in Figure 3. Although the population bursts in the rastergram do not appear completely periodically, partly because of the stochastic nature of the Poisson-distributed input spike trains, counting the total number of bursts appearing in the rastergram of Figure 11b confirms a close agreement between the frequency of oscillatory bursting observed here with that of the mean-field oscillations shown in Figure 10b. Importantly, the spiking activity we obtained here, both at the asynchronous irregular regime and at the oscillatory bursting regime, are also closely comparable with those obtained from *in vitro* experiments and detailed *in silico* reconstruction of the rat neocortical microcircuitry, as shown in [37, Figure 12].

Similar to the results shown in Figure 10, the instantaneous profiles of mean firing rates and the cross-correlation curves shown in Figure 11 indicate a tight temporal correlation between the excitatory and inhibitory spontaneous activity, in both asynchronous irregular and oscillatory bursting regimes. Consistent with the results obtained using the mean-field model, the time lag between the mean excitatory and inhibitory spiking rates obtained here are very small: 0.78 ms at the asynchronous irregular state, and 0.76 ms at the oscillatory bursting state. At both states, the spontaneous inhibitory activity appears to be slightly ahead of the excitatory activity.

Discussion

By leveraging the computational tractability of a biologically reasonable conductance-based mean-field model, we conducted a fairly comprehensive study of how variations in some of the main synaptic and structural parameters of a local cortical network—specifically characterized by the properties of the mouse and rat neocortical microcircuitry—influence the balance between overall excitation and inhibition in the network. We performed bifurcation analyses of the baseline balanced equilibrium state of the mean-field model with respect to variations in each of the key parameters of the network, namely, the synaptic (physiological) parameters such as synaptic decay time constants, synaptic quantal conductances, and synaptic reversal potentials, as well as the structural parameters such as the ratio of the number of inhibitory neurons to the total number of neurons, the density (or sparsity) of the overall network connectivity, and the inhibitory-to-inhibitory connection probability. Additionally, we used a sufficiently large network of spiking neurons to test the reliability of the predictions made based on the mean-field model. Below, we discuss the key findings of our study.

Characterization of the balanced state: The balanced state in our analysis refers to a population-level operational state of the network that satisfies certain properties: (1) the spontaneous activity of the neurons in the network are asynchronous and irregular, (2) the network is sufficiently excitable (3) the spontaneous and stimulated activity of the network remain stable, and (4) the network responds rapidly to a reasonably wide range of external stimuli. The neurons in such balanced state are mostly expected to be depolarized near their

spiking threshold.

Continuous quantification of the level of balance: The level of overall excitation and inhibition in a network is often identified in previous studies as either balanced or imbalanced. However, the degree by which the overall balance of excitation and inhibition, as we characterized above, deviates from its perfect level is continuously quantifiable. That means, we can quantify how balanced the network is, or how far the network activity is deviated from the balanced state. We performed such continuous quantification in our analysis. We first identified a baseline (well-) balanced reference state for the mean-field activity of our model by analyzing its long-term behavior in response to different levels of external excitatory inputs. Then, we investigated how this established balanced state is altered by changes in different parameters of the network. Our results confirm that the ratio between the mean values of excitatory and inhibitory synaptic conductances in the network—in reference to its value at the baseline balanced state—is a reliable measure for contentiously quantifying the level of overall excitation-inhibition balance. Mean excitatory and inhibitory conductances can be experimentally measured, both *in vitro* and *in vivo* [56], and the ratio between them has been shown to remain constant in well-balanced states [3–5, 11, 34].

Neuronal gain and excitability modulation: Increasing the decay time constant and the quantal conductance of inhibitory synapses decreases the gain (slope of the repose curve) and excitability (horizontal shift of the response curve) of the neurons. Opposite effects are observed when the decay time constant and the quantal conductance of excitatory synapses are increased. Increases in excitatory synaptic reversal potentials result in enhancements in the gains of the neurons. Changes in the inhibitory reversal potentials have non-monotonic effects on the gain of the neurons.

Effect of synaptic parameters: Increasing the value of the three main inhibitory synaptic parameters that we studied here to values above their baseline moves the excitation-inhibition balance toward over-inhibition. On the other hand, decreasing these parameters below their baseline value moves the balance toward over-excitation. Modulations of the excitatory synaptic parameters have the opposite effects on the network balance, but with slightly less critical impacts on the network stability.

Effect of the inhibitory proportion of neurons: Changes in the proportion of inhibitory neurons relative to the total number of neurons affect the excitation-inhibition balance in essentially similar way to how changes in inhibitory synaptic decay time constants do.

Effect of the density of overall network connectivity Increasing the sparsity of overall network connectivity moves the network balance toward over-excitation. When the network is very sparsely-connected, a further increase in the sparsity of the network connectivity causes a very sharp rise in the level of overall excitation, yet the network dynamics does not transition to an oscillatory regime. Increasing the density of overall network connectivity, on the other hand, slowly shifts the network balance to over-inhibition.

Transition to slow oscillatory regimes: In most cases in our results, sufficiently large deviations of the network balance toward over-excitation transition the network activity to

an oscillatory regime. The only exception we observed was the case where we increased the overall sparsity of the network connectivity, which resulted in hyper-excited yet stable non-oscillatory states of network activity. In all cases, the emerging oscillations are slow, with their frequency being in the delta band. In particular, network oscillations emerging due to reduction in the inhibitory proportion of the total network population or in the decay time constant of inhibitory synapses, as well as those emerging due to higher density of recurrent inhibitory-to-inhibitory connectivity, are of high amplitude. High-amplitude delta oscillations are strongly correlated with loss of consciousness, and are frequently observed in states such as coma, anaesthesia, generalized epileptic seizures, and slow wave sleep [17, 57]. Therefore, it can be implied from our results that transitions to slow oscillatory regimes due to shifts in the balance of excitation and inhibition—if occurred in a sufficiently large number of local cortical networks across the neocortex—can indeed correspond to a critical state transition in the brain to a state of diminished consciousness.

Criticality of the inhibitory synaptic conductances and the inhibitory proportion of neurons: The baseline values for inhibitory synaptic decay time constants, and for the ratio between the number of inhibitory neurons and the total number of neurons, are critical values for network stability. At such values, the network is in the well-balanced state as characterized above. However, relatively small reductions in these values result in a transition of network activity to a slow oscillatory regime. This, in particular, implies that the typical value of the inhibitory proportion of neurons in the network or, equivalently, the typical ratio between the number of inhibitory and excitatory neurons, is almost optimal. This optimality is in the sense that a larger inhibitory proportion results in redundant inhibition and a less excitable network, and a slightly smaller inhibitory proportion results in a phase transition and, possibly, loss of functionality. Therefore, the number of inhibitory neurons, relative to the number of excitably neurons, appears to be nearly at the minimum number required to safely stabilize the overall excitatory activity under normal conditions. This optimal value, however, is in direct correspondence with the value of inhibitory synaptic decay time constants. A larger number of inhibitory neurons is required for network stability if inhibitory synapses decay faster, and a smaller number of inhibitory neurons can maintain the stability of the network if inhibitory synaptic events last longer.

Criticality of the density of inhibitory-to-inhibitory connectivity: The density of recurrent inhibitory-to-inhibitory connectivity, relative to the density of other types of local cortical connectivity, is a crucial factor both in controlling the level of inhibitory activity in the network and in stabilizing the excitatory activity. Our results show that the baseline value for the probability of such recurrent inhibitory connectivity is critical. A moderately lower density of inhibitory-to-inhibitory connectivity results in a rather uncontrolled rise in inhibitory activity, which moves the network balance toward hyper-inhibition and yields a non-excitable network. On the contrary, a slight increase in the density of such recurrent inhibitory connectivity results in a sharp change in the balance of excitation and inhibition toward over-excitation and, subsequently, emergence of high-amplitude delta oscillations.

Robustness of network stability to changes in quantal conductances: The stability of the network is fairly robust to changes in synaptic quantal conductances. That is, only a significantly large reduction in the value of inhibitory quantal conductances, or a significantly large increase in the value of excitatory quantal conductances, can destabilize the network's

equilibrium through a Hopf bifurcation. This is in fact an expedient property for the functionality of the network. The values of synaptic quantal conductances are subject to dynamic changes due to plasticity of the synapses, which is essential in learning and memory. The network would otherwise be totally dysfunctional if it could be destabilized by such level of synaptic plasticity. This robustness of the network stability to changes in synaptic quantal conductances further allows for synaptic plasticity to play a key role in fine-tuning of the excitation-inhibition balance [8, 10, 31, 32, 58].

Joint effect of synaptic and structural parameters: Balance of excitation and inhibition is established by integrated contributions of multiple physiological and anatomical factors. We mostly analyzed the effects of these factors separately. However, our limited results on the joint effect of synaptic parameters and the relative size of the inhibitory population particularly suggest that the high-amplitude delta oscillations that emerge in a local cortical network due to a reduced number of inhibitory neurons can be effectively suppressed by a reasonable amount of increase in the decay time constant or reversal potential of the inhibitory synapses, which restores the network's stability by transitioning its dynamics through a Hopf bifurcation.

Instantaneous correlation of excitation and inhibition: Besides our extensive investigation of the overall excitation-inhibition balance through analyzing the long-term dynamics of the mean-field model, we also studied the instantaneous correlation between the spontaneous excitatory and inhibitory activity in a network, using both the mean-field model and the spiking neuronal network model. Our results reveals a tight instantaneous correlation between mean excitatory and inhibitory firing rates, with an average time lag of less that one millisecond—in both the baseline balanced and the oscillatory states. The presence of such a tight correlation is consistent with the some experimental observations, which have confirmed tightly correlated temporal variations in the activity of excitatory and inhibitory neurons in a balanced state, with a time lag of only a few millisecond between them [6, 7, 11, 59, 60]. In our simulations, on average, the mean spontaneous activity of inhibitory neurons leads that of the excitatory neurons by a fraction of millisecond, whereas experimental studies often report inhibitory activities to lag a few millisecond behind the excitatory activity, especially for evoked activities [6, 7]. The seeming discrepancy, however, is likely because the sign of the time lag between the correlated activities depends on the physical quantities measured (e.g., firing rates, postsynaptic potentials, postsynaptic currents, etc.) as well as specific distributions and timing of external inputs to each of the inhibitory and excitatory populations. In our simulations, we provided evenly distributed external excitatory spikes at a constant average rate to both populations.

Insufficiency of tight activity correlation for inferring normal balance: The tight instantaneous correlation between spontaneous excitatory and inhibitory activities is maintained at different levels of excitation-inhibition balance in our simulations, both in a network operating near a stable equilibrium, and in a network that has transitioned to a slowly-oscillating (-bursting) regime. The presence of such a tight temporal correlation has been confirmed experimentally at different dynamic states of spontaneous, evoked, and oscillatory network activity [6, 7, 54, 59]. However, the functionality of a cortical network, and its overall level of excitation-inhibition balance, can change substantially at these different states. For example, the presence of high-amplitude delta oscillations in a network can be associated

with a state of diminished consciousness, a state functionally distinctive from the alert state. Moreover, the level of overall excitation-inhibition balance in a network, when it is operating at different conditions, can deviate from its normal dynamic state, either toward more inhibition or toward more excitation. Importantly, such deviations do not necessarily reflect on the instantaneous correlation between excitatory and inhibitory activities. Therefore, we argue that a tight correlation between the temporal recordings of excitatory and inhibitory neuronal activities in a network is not a sufficiently strong evidence for inferring a functional balance of excitation and inhibition in the network. As we discussed above, our results suggests that the ratio between mean excitatory and inhibitory synaptic conductances is a more reliable measure of the level of overall excitation-inhibition balance in a local cortical network, as it continuously quantifies the amount of deviations from a normally balanced state.

Reliability of the predictions: The results we presented in this paper were predominantly obtained using a mean-field model, which naturally suffers from inaccuracies due to simplifying assumptions. However, the model is simple and yet maintains key biological details. This allowed us the capacity to fairly comprehensively investigate the contributions of various factors in the balance of excitation and inhibition in a local cortical network. Performing such an extensive study experimentally, or using detailed biological models, would be rather impractical. To test the reliability of the predictions made based on the mean-field model, we performed a sample study using a more detailed framework of a spiking network. The close agreement between the results we obtained from this spiking network and those predicted based on the mean-field model—which were also consistent with the spiking activity observed in *in vitro* experiments and in *in silico* computations of a detailed neocortical microcircuitry [37, Figure 12]—is promising. Hence, we anticipate that the observations we made in this paper can inform future experimental and theoretical studies on understanding the homeostatic mechanisms involved in regulation of the excitation-inhibition balance, and on identifying the pathological causes of short-term and long-term disturbances in this essential operating condition of local cortical networks.

Funding acknowledgment

This work was supported by the National Eye Institute of the National Institutes of Health under Award Number R00 EY030840. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Author contribution

We use the CRediT Taxonomy to describe each author's contributions to the work, as follows: Conceptualization (FS), Data Curation (FS), Formal Analysis (FS), Funding Acquisition (HC), Investigation (FS), Methodology (FS), Project Administration (FS, HC), Resources (N/A), Software (FS), Supervision (N/A), Validation (FS), Visualization (FS), Writing–Original Draft Preparation (FS), Writing–Review & Editing (FS, HC)

References

[1] E. R. Kandel, J. H. Schwartz, T. M. Jessell, S. A. Siegelbaum, and A. J. Hudspeth. *Principles of Neural Science.* McGraw-Hill, New York, NY, 5th edition, 2013.

[2] F. Shirani, W. M. Haddad, and R. de la Llave. On the global dynamics of an electroencephalographic mean field model of the neocortex. *SIAM Journal on Applied Dynamical Systems*, 16(4):1969–2029, 2017.

[3] Bilal Haider, Alvaro Duque, Andrea R. Hasenstaub, and David A. McCormick. Neocortical network activity in vivo is generated through a dynamic balance of excitation and inhibition. *Journal of Neuroscience*, 26(17):4535–4545, 2006.

[4] Yousheng Shu, Andrea Hasenstaub, and David A. McCormick. Turning on and off recurrent balanced cortical activity. *Nature*, 423(6937):288–293, 2003.

[5] Michael Wehr and Anthony M. Zador. Balanced inhibition underlies tuning and sharpens spike timing in auditory cortex. *Nature*, 426(6965):442–446, 2003.

[6] Michael Okun and Ilan Lampl. Instantaneous correlation of excitation and inhibition during ongoing and sensory-evoked activities. *Nature Neuroscience*, 11(5):535–537, 2008.

[7] N. Dehghani, A. Peyrache, B. Telenczuk, M. Le Van Quyen, E. Halgren, S. S. Cash, N. G. Hatsopoulos, and A. Destexhe. Dynamic balance of excitation and inhibition in human and monkey neocortex. *Scientific Reports*, 6:23176, 2016.

[8] Robert C. Froemke. Plasticity of cortical excitatory-inhibitory balance. *Annual Review of Neuroscience*, 38(1):195–219, 2015.

[9] Ofer Yizhar, Lief E. Fenno, Matthias Prigge, Franziska Schneider, Thomas J. Davidson, Daniel J. O'Shea, Vikaas S. Sohal, Inbal Goshen, Joel Finkelstein, Jeanne T. Paz, Katja Stehfest, Roman Fudim, Charu Ramakrishnan, John R. Huguenard, Peter Hegemann, and Karl Deisseroth. Neocortical excitation/inhibition balance in information processing and social dysfunction. *Nature*, 477(7363):171–178, 2011.

[10] T. P. Vogels, H. Sprekeler, F. Zenke, C. Clopath, and W. Gerstner. Inhibitory plasticity balances excitation and inhibition in sensory pathways and memory networks. *Science*, 334(6062):1569–1573, 2011.

[11] Sophie Denève and Christian K. Machens. Efficient codes and balanced networks. *Nature Neuroscience*, 19(3):375–382, 2016.

[12] Michael N. Shadlen and William T. Newsome. Noise, neural codes and cortical organization. *Current Opinion in Neurobiology*, 4(4):569–579, 1994.

[13] C. van Vreeswijk and H. Sompolinsky. Chaos in neuronal networks with balanced excitatory and inhibitory activity. *Science*, 274(5293):1724–1726, 1996.

[14] Michael N. Shadlen and William T. Newsome. The variable discharge of cortical neurons: Implications for connectivity, computation, and information coding. *Journal of Neuroscience*, 18(10):3870–3896, 1998.

[15] Nicolas Brunel. Dynamics of sparsely connected networks of excitatory and inhibitory spiking neurons. *Journal of Computational Neuroscience*, 8(3):183–208, 2000.

[16] Hai-yan He and Hollis T. Cline. What is excitation/inhibition and how is it regulated? A case of the elephant and the eisemen. *Journal of Experimental Neuroscience*, 13:1179069519859371, 2019.

[17] Grant Gillary and Ernst Niebur. The edge of stability: Response times and delta oscillations in balanced networks. *PLOS Computational Biology*, 12(9):1–22, 09 2016.

[18] F. Shirani. Transient neocortical gamma oscillations induced by neuronal response modulation. *Journal of computational neuroscience*, 48(1):103–122, 2020.

[19] Lu Chen, Xiling Li, Michelle Tjia, and Shruti Thapliyal. Homeostatic plasticity and excitation-inhibition balance: The good, the bad, and the ugly. *Current Opinion in Neurobiology*, 75:102553, 2022.

[20] T. P. Vogels and L. F. Abbott. Gating multiple signals through detailed balance of excitation and inhibition in spiking networks. *Nature Neuroscience*, 12(4):483–491, 2009.

[21] Brendan K. Murphy and Kenneth D. Miller. Balanced amplification: A new mechanism of selective amplification of neural activity patterns. *Neuron*, 61(4):635–648, 2009.

[22] Sukbin Lim and Mark S Goldman. Balanced cortical microcircuitry for maintaining information in working memory. *Nature Neuroscience*, 16(9):1306–1314, 2013.

[23] Sacha B. Nelson and Vera Valakh. Excitatory/inhibitory balance and circuit homeostasis in autism spectrum disorders. *Neuron*, 87(4):684–698, 2015.

[24] J. L. R. Rubenstein and M. M. Merzenich. Model of autism: increased ratio of excitation/inhibition in key neural systems. *Genes, Brain and Behavior*, 2(5):255–267, 2003.

[25] Jorge J. Palop, Jeannie Chin, Erik D. Roberson, Jun Wang, Myo T. Thwin, Nga Bien-Ly, Jong Yoo, Kaitlyn O. Ho, Gui-Qiu Yu, Anatol Kreitzer, Steven Finkbeiner, Jeffrey L. Noebels, and Lennart Mucke. Aberrant excitatory neuronal activity and compensatory remodeling of inhibitory hippocampal circuits in mouse models of Alzheimer's disease. *Neuron*, 55(5):697–711, 2007.

[26] Jorge J. Palop and Lennart Mucke. Network abnormalities and interneuron dysfunction in alzheimer disease. *Nature Reviews Neuroscience*, 17(12):777–792, 2016.

[27] Vardhan S. Dani, Qiang Chang, Arianna Maffei, Gina G. Turrigiano, Rudolf Jaenisch, and Sacha B. Nelson. Reduced cortical activity due to a shift in the balance between excitation and inhibition in a mouse model of rett syndrome. *Proceedings of the National Academy of Sciences*, 102(35):12560–12565, 2005.

[28] Yue Kris Wu, Keith B. Hengen, Gina G. Turrigiano, and Julijana Gjorgjieva. Homeostatic mechanisms regulate distinct aspects of cortical circuit dynamics. *Proceedings of the National Academy of Sciences*, 117(39):24514–24525, 2020.

[29] Zhengyu Ma, Gina G. Turrigiano, Ralf Wessel, and Keith B. Hengen. Cortical circuit dynamics are homeostatically tuned to criticality in vivo. *Neuron*, 104(4):655–664.e4, 2019.

[30] Nicolas Le Roux, Muriel Amar, Gérard Baux, and Philippe Fossier. Homeostatic control of the excitation–inhibition balance in cortical layer 5 pyramidal neurons. *European Journal of Neuroscience*, 24(12):3507–3518, 2006.

[31] Guillaume Hennequin, Everton J. Agnes, and Tim P. Vogels. Inhibitory plasticity: Balance, control, and codependence. *Annual Review of Neuroscience*, 40(1):557–579, 2017.

[32] Henning Sprekeler. Functional consequences of inhibitory plasticity: homeostasis, the excitation-inhibition balance and beyond. *Current Opinion in Neurobiology*, 43:198–203, 2017.

[33] Gina G. Turrigiano, Kenneth R. Leslie, Niraj S. Desai, Lana C. Rutherford, and Sacha B. Nelson. Activity-dependent scaling of quantal amplitude in neocortical neurons. *Nature*, 391(35):892–896, 1998.

[34] Mingshan Xue, Bassam V. Atallah, and Massimo Scanziani. Equalizing excitation–inhibition ratios across visual cortical neurons. *Nature*, 511(7511):596–600, 2014.

[35] B. Haider and D. A. McCormick. Rapid neocortical dynamics: Cellular and network mechanisms. *Neuron*, 62(2):171–189, 2009.

[36] Alfonso Renart, Jaime de la Rocha, Peter Bartho, Liad Hollender, Néstor Parga, Alex Reyes, and Kenneth D. Harris. The asynchronous state in cortical circuits. *Science*, 327(5965):587–590, 2010.

[37] Henry Markram, Eilif Muller, Srikanth Ramaswamy, Michael W. Reimann, Marwan Abdellah, Carlos Aguado Sanchez, Anastasia Ailamaki, Lidia Alonso-Nanclares, Nicolas Antille, Selim Arsever, Guy Antoine Atenekeng Kahou, Thomas K. Berger, Ahmet Bilgili, Nenad Buncic, Athanassia Chalimourda, Giuseppe Chindemi, Jean-Denis Courcol, Fabien Delalondre, Vincent Delattre, Shaul Druckmann, Raphael Dumusc, James Dynes, Stefan Eilemann, Eyal Gal, Michael Emiel Gevaert, Jean-Pierre Ghobril, Albert Gidon, Joe W. Graham, Anirudh Gupta, Valentin Haenel, Etay Hay, Thomas Heinis, Juan B. Hernando, Michael Hines, Lida Kanari, Daniel Keller, John Kenyon, Georges Khazen, Yihwa Kim, James G. King, Zoltan Kisvarday, Pramod Kumbhar, Sébastien Lasserre, Jean-Vincent Le Bé, Bruno R.C. Magalhães, Angel Merchán-Pérez, Julie Meystre, Benjamin Roy Morrice, Jeffrey Muller, Alberto Muñoz-Céspedes, Shruti Muralidhar, Keerthan Muthurasa, Daniel Nachbaur, Taylor H. Newton, Max Nolte, Aleksandr Ovcharenko, Juan Palacios, Luis Pastor, Rodrigo Perin, Rajnish Ranjan, Imad Riachi, José-Rodrigo Rodríguez, Juan Luis Riquelme, Christian Rössert, Konstantinos Sfyrakis, Ying Shi, Julian C. Shillcock, Gilad Silberberg, Ricardo Silva, Farhan Tauheed, Martin Telefont, Maria Toledo-Rodriguez, Thomas Tränkler, Werner Van Geit, Jafet Villafranca Díaz, Richard Walker, Yun Wang, Stefano M. Zaninetta, Javier DeFelipe, Sean L. Hill, Idan Segev, and Felix Schürmann. Reconstruction and simulation of neocortical microcircuitry. *Cell*, 163(2):456–492, 2015.

[38] Sami El Boustani and Alain Destexhe. A master equation formalism for macroscopic modeling of asynchronous irregular activity states. *Neural Computation*, 21(1):46–100, 2009.

[39] Y. Zerlaut, B. Teleńczuk, C. Deleuze, T. Bal, G. Ouanounou, and A. Destexhe. Heterogeneous firing rate response of mouse layer v pyramidal neurons in the fluctuation-driven regime. *The Journal of Physiology*, 594(13):3791–3808, 2016.

[40] Y. Zerlaut, S. Chemla, F. Chavane, and A. Destexhe. Modeling mesoscopic cortical dynamics using a mean-field model of conductance-based networks of adaptive exponential integrate-and-fire neurons. *Journal of Computational Neuroscience*, 44(1):45–61, 2018.

[41] Matteo di Volo, Alberto Romagnoni, Cristiano Capone, and Alain Destexhe. Biologically realistic mean-field models of conductance-based networks of spiking neurons with adaptation. *Neural Computation*, 31(4):653–680, 2019.

[42] M. Carlu, O. Chehab, L. Dalla Porta, D. Depannemaecker, C. Héricé, M. Jedynak, E. Köksal Ersöz, P. Muratore, S. Souihel, C. Capone, Y. Zerlaut, A. Destexhe, and M. di Volo. A mean-field approach to the dynamics of networks of complex neurons, from nonlinear integrate-and-fire to Hodgkin–Huxley models. *Journal of Neurophysiology*, 123(3):1042–1051, 2020.

[43] Corinne Teeter, Ramakrishnan Iyer, Vilas Menon, Nathan Gouwens, David Feng, Jim Berg, Aaron Szafer, Nicholas Cain, Hongkui Zeng, Michael Hawrylycz, and Stefan Koch, Christof Mihalas. Generalized leaky integrate-and-fire models classify multiple neuron types. *Nature Communications*, 9(1):709, 2018.

[44] A. Dhooge, W. Govaerts, Yu. A. Kuznetsov, H. G. E. Meijer, and B. Sautois. New features of the software MatCont for bifurcation analysis of dynamical systems. *Math. Comput. Model. Dyn. Syst.*, 14(2):147–175, 2008.

[45] Alexandre Kuhn, Ad Aertsen, and Stefan Rotter. Neuronal integration of synaptic input in the fluctuation-driven regime. *Journal of Neuroscience*, 24(10):2345–2356, 2004.

[46] Romain Brette and Wulfram Gerstner. Adaptive exponential integrate-and-fire model as an effective description of neuronal activity. *Journal of Neurophysiology*, 94(5):3637–3642, 2005.

[47] Allen Institute for Brain Science. Allen cell types database. `https://celltypes.brain-map.org/`, 2016. Accessed: 2022-09-01.

[48] Blue Brain Project. Blue brain portal. `https://portal.bluebrain.epfl.ch/`, 2005. Accessed: 2022-09-01.

[49] Armen Stepanyants, Luis M. Martinez, Alex S. Ferecská, and Zoltán F. Kisvárday. The fractions of short- and long-range connections in the visual cortex. *Proceedings of the National Academy of Sciences*, 106(9):3555–3560, 2009.

[50] Luke Campagnola, Stephanie C. Seeman, Thomas Chartrand, Lisa Kim, Alex Hoggarth, Clare Gamlin, Shinya Ito, Jessica Trinh, Pasha Davoudian, Cristina Radaelli, Mean-Hwan Kim, Travis Hage, Thomas Braun, Lauren Alfiler, Julia Andrade, Phillip Bohn, Rachel Dalley, Alex Henry, Sara Kebede, Alice Mukora, David Sandman, Grace Williams, Rachael Larsen, Corinne Teeter, Tanya L. Daigle, Kyla Berry, Nadia Dotson, Rachel Enstrom, Melissa Gorham, Madie Hupp, Samuel Dingman Lee, Kiet Ngo, Philip R. Nicovich, Lydia Potekhina, Shea Ransford, Amanda Gary, Jeff Goldy, Delissa McMillen, Trangthanh Pham, Michael Tieu, La'Akea Siverts, Miranda Walker, Colin Farrell, Martin Schroedter, Cliff Slaughterbeck, Charles Cobb, Richard Ellenbogen, Ryder P. Gwinn, C. Dirk Keene, Andrew L. Ko, Jeffrey G. Ojemann, Daniel L. Silbergeld, Daniel Carey, Tamara Casper, Kirsten Crichton, Michael Clark, Nick Dee, Lauren Ellingwood, Jessica Gloe, Matthew Kroll, Josef Sulc, Herman Tung, Katherine Wadhwani,

Krissy Brouner, Tom Egdorf, Michelle Maxwell, Medea McGraw, Christina Alice Pom, Augustin Ruiz, Jasmine Bomben, David Feng, Nika Hejazinia, Shu Shi, Aaron Szafer, Wayne Wakeman, John Phillips, Amy Bernard, Luke Esposito, Florence D. D'Orazi, Susan Sunkin, Kimberly Smith, Bosiljka Tasic, Anton Arkhipov, Staci Sorensen, Ed Lein, Christof Koch, Gabe Murphy, Hongkui Zeng, and Tim Jarsky. Local connectivity and synaptic dynamics in mouse and human neocortex. *Science*, 375(6585):eabj5861, 2022.

[51] Tobias C. Potjans and Markus Diesmann. The cell-type specific cortical microcircuit: Relating structure and activity in a full-scale spiking network model. *Cerebral Cortex*, 24(3):785–806, 2012.

[52] Michael Rudolph, Joe Guillaume Pelletier, Denis Paré, and Alain Destexhe. Characterization of synaptic conductances and integrative properties during electrically induced EEG-activated states in neocortical neurons in vivo. *Journal of Neurophysiology*, 94(4):2805–2821, 2005.

[53] Itamar D. Landau, Robert Egger, Vincent J. Dercksen, Marcel Oberlaender, and Haim Sompolinsky. The impact of structural heterogeneity on excitation-inhibition balance in cortical networks. *Neuron*, 92(5):1106–1121, 2016.

[54] Jeffry S. Isaacson and Massimo Scanziani. How inhibition shapes cortical activity. *Neuron*, 72(2):231–243, 2011.

[55] Jay R. Gibson, Michael Beierlein, and Barry W. Connors. Two networks of electrically coupled inhibitory neurons in neocortex. *Nature*, 402(6757):75–79, 1999.

[56] C. Monier, J. Fournier, and Y. Frégnac. In vitro and in vivo measures of evoked excitatory and inhibitory conductance dynamics in sensory cortices. *Journal of Neuroscience Methods*, 169(2):323–365, 2008.

[57] Joel Frohlich, Daniel Toker, and Martin M Monti. Consciousness among delta waves: a paradox? *Brain*, 144(8):2257–2277, 2021.

[58] Hai-Yan He, Wanhua Shen, Masaki Hiramoto, and Hollis T. Cline. Experience-dependent bimodal plasticity of inhibitory neurons in early development. *Neuron*, 90(6):1203–1214, 2016.

[59] Bassam V. Atallah and Massimo Scanziani. Instantaneous modulation of gamma oscillation frequency by balancing excitation with inhibition. *Neuron*, 62(4):566–577, 2009.

[60] Aanchal Bhatia, Sahil Moza, and Upinder Singh Bhalla. Precise excitation-inhibition balance controls gain and timing in the hippocampus. *eLife*, 8:e43415, 2019.