_A Network Pharmacology Approach to Elucidate the Anti-inflammatory Effects of Ellagic Acid

Acid

Skyler H. Hoang[1], Hue Dao[1], Emerson My Lam[2]

[1]Department of Cell Biology and Neuroscience, Rutgers University, Piscataway, New Jersey

089854, USA

[2]Faculty of Pharmacy, Ton Duc Thang University, Ho Chi Minh City-700000, Vietnam.

Corresponding email: hung.hoang@rutgers.edu

ORCID

Skyler H. Hoang – 0000-0001-8242-412X

Hue Dao – 0009-0003-3256-3043

Emerson My Lam – 0009-0001-4015-6355

Abstract

Ellagic acid (EA) is a naturally occurring polyphenolic compound found in various fruits and vegetables like strawberries, raspberries, pomegranates, and nuts such as pecans and walnuts. With its antioxidant properties, EA has shown potential health benefits, although further research is necessary to fully comprehend its effects, mechanisms, and safe and effective application as a complementary medicine. Notably, there is accumulating evidence of EA's anti-inflammatory effects; however, the precise underlying mechanism remains unclear. To investigate the anti-inflammatory properties of EA, a network pharmacology approach was employed. The study identified 52 inflammation-related targets of EA and revealed significant signaling pathways and relevant diseases associated with inflammation through GO and KEGG analysis. Furthermore, topological analysis identified 10 important targets, including AKT1, VEGFA, TNF, MAPK3, ALB, SELP, MMP9, MMP2, PTGS2, and ICAM1. Molecular docking and molecular dynamics simulations (integrated with were conducted molecular mechanics Poisson-Boltzmann), indicating that AKT1, PTGS2, VEGFA, and MAPK3 are the most likely targets of EA. In summary, this study not only confirmed the anti-inflammatory effects of EA observed in previous research but also identified the most probable targets of EA.

47 **Abbreviations**

AKT1	AKT serine/threonine kinase 1
ALB	Albumin
DMNC	Density of Maximum Neighborhood Component
EA	Ellagic Acid
EPC	Edge Percolated Component
GO	Gene Ontology
HIF	Hypoxia-inducible Factor
ICAM	Intercellular Adhesion Molecule
IL	Interleukin
KEGG	Kyoto Encyclopedia of Genes and Genomes
MAPK	Mitogen-activated Protein Kinase
MCC	Maximal Clique Centrality
MMP	Matrix Metallopeptidase
MM-PBSA	Molecular Mechanics Poisson-Boltzmann Surface Area
MNC	Maximum Neighborhood Component
NF-κB	Nuclear factor kappa B
PPI	Protein Protein Interaction
PTGS	Prostaglandin-endoperoxide Synthase 2
SELP	Selectin P
TNF	(TNF-α) Tumor Necrosis Factor alpha
VEGFA	Vascular Endothelial Growth Factor A

Introduction

Ellagic acid (EA) is a naturally occurring polyphenolic compound with antioxidant properties found in a variety of fruits and vegetables, including strawberries, raspberries, pomegranates, and nuts such as pecans and walnuts (Baliga et al., 2019; Sharifi-Rad et al., 2022). EA has been studied for its potential health benefits, including its ability to support the immune system and protect cells from damage (Sharifi-Rad et al., 2022). Some research has suggested that EA may have anti-inflammatory and chemopreventive effects, which could potentially help to reduce the risk of chronic diseases such as heart disease and cancer (BenSaad et al., 2017; Päivärinta et al., 2006; Ridker, 2017). For instance, in a study model of culture of primary human gingival epithelial cells, EA was demonstrated to regulate the oral innate immunity via increasing the expression of chemokine ligand 5, IL-2, IL-1β and decreasing the expression of IL-6, IL-8, and TNF (Promsong et al., 2015). In terms of its potential anticancer effects, some research has suggested that ellagic acid may help to inhibit the growth of cancer cells and induce cell death in certain types of cancer, such as breast, colon, and prostate cancer (Ceci et al., 2018). In another instance, transcriptional profiling using microarray and functional analysis on Caco-2 cells exposed to EA revealed that EA could affect the expression of growth factor receptors (fibroblast growth factor receptor 2, epidermal growth factor receptor), oncogenes (K-Ras, c-Myc), and tumor suppressors (dual specificity phosphatase 6, Fos) and of genes involved in cell cycle (cyclin B1, cyclin B1 Interacting Protein 1) (González-Sarrías et al., 2009). EA is also claimed to support a variety of other health goals, including weight loss, cardiovascular health, and improved digestion (Kang et al., 2016). More research into the mechanism of this compound is needed to fully understand these potential effects and how anti-inflammatory, anti-cancer, or metabolic syndrome preventive effects of EA can be utilized. Overall, EA appears to have

potential health benefits, but more research is needed to fully understand its effects and how it

can be used safely and effectively as a complimentary medicine.

On another note, inflammation is a natural process that occurs in the body in response to

injury or infection (L. Chen et al., 2018). It is characterized by swelling, redness, warmth, and

pain, and is a crucial part of the body's immune response (Scott, 2004). When the body is injured

or infected, immune cells are activated to help fight off the infection or repair the damage (Fang

et al., 2018; Luster et al., 2005). This process involves the release of various chemicals and

signaling molecules that cause inflammation in the affected area. Inflammation is important

because it helps to protect the body from harm and promote healing. However, chronic

inflammation, which is a long-term, persistent form of inflammation, can be harmful and

contribute to the development of chronic diseases such as heart disease, diabetes, and cancer

(Furman et al., 2019). Chronic inflammation is often caused by underlying conditions such as

obesity, smoking, and high levels of stress, and can also be triggered by certain foods and

environmental factors (Bosma-den Boer et al., 2012). There is increasing evidence to suggest

that chronic inflammation is a key factor in the development of many chronic diseases. For

instance, cardiovascular diseases are often caused by inflammation in the arteries, which can lead

to the build-up of plaque and eventually cause a heart attack or stroke (Kabbany et al., 2016).

Similarly, inflammation in the pancreas may contribute to the development of diabetes, and

inflammation in the colon may increase the risk of colon cancer (Terzić et al., 2010). Studying

inflammation is important because it helps to understand the underlying causes of chronic

diseases and may lead to the development of new treatments and prevention strategies. Certain

medications, such as nonsteroidal anti-inflammatory drugs and corticosteroids, can help to

reduce inflammation and pain or may be used to treat conditions such as arthritis (Del Grossi

Moura et al., 2018). However, their misusage has been associated with adverse events such as

gastrointestinal ulcers, serious cardiovascular events, or hypertension (Vonkeman & van de Laar,

2010; Yasir et al., 2022). It should be noted that lifestyle changes such as eating a healthy diet,

exercising regularly, and managing stress may also help to reduce inflammation and lower the

risk of chronic diseases (Margină et al., 2020). EA, with accumulating evidence of anti-

inflammatory properties, can be easily found as a part of a healthy, plant-based diet (Kang et al.,

2016). This investigation is also further compelled by the fact that EA has been demonstrated to

exert glucose-lowering, antioxidant, anti-inflammatory, and anti-glycation effects in murine

diabetic models (Amor et al., 2020). Herein, a network pharmacology approach, integrated with

molecular docking and molecular dynamics simulation, is utilized to elucidate the mechanisms

underlying anti-inflammatory properties of EA.

Results and Discussion

1. Network Construction and Analysis

The retrieval of EA target proteins and target proteins for inflammation results in 157 and

886 target proteins, respectively (**Figure 1**). There were 52 overlapped proteins, representing

potential therapeutic targets of EA for its anti-inflammatory properties. A PPI network of 52

overlapped was constructed and resulting in 325 number of edges, 12.5 average node degree,

0.688 average local clustering coefficient, as well as a PPI enrichment p-value of less than 1.0e-

16, indicating that the proteins are biologically connected (**Figure 1**).

The 5 most significant GO Biological Process enriched pathways include: response to

organic substance, response to oxygen-containing compound, response to lipid, response to

chemical, and cellular response to chemical (**Figure 2A**). The 5 most significant GO Molecular

Function enriched pathways include: identical protein binding, protein binding, signaling

receptor binding, enzyme binding, and ion binding (**Figure 2A**). The 5 most significant GO

Cellular Component enriched pathways include: vesicle lumen, endomembrane system, secretory

granule, secretory granule lumen, and membrane raft (**Figure 2A**).

Interestingly, lipid regulation and inflammation have an intimate relationship.

Particularly, abnormalities in lipid profile or regulation can lead to abnormalities in

inflammatory responses, and the reverse is also true in that alterations of inflammatory signaling

(in the context of metabolic diseases such as diabetes or atherosclerosis) can result in changes of

lipid metabolism in liver, skeletal muscle, macrophage, or adipose tissue (Glass & Olefsky,

2012). In a murine basal diet model, EA supplementation decreased total cholesterol, low-

density lipoprotein, and the abundance of acetyl-CoA carboxylase while increasing high-density

lipoprotein and the abundances of phospho-hormone-sensitive lipase, carnitine palmitoyl

transferase 1B, and peroxisome proliferator-activated receptor alpha protein (Xu et al., 2021). In

a randomized double-blind clinical trial, patients with type 2 diabetes consuming 180 mg of EA

daily for 8 weeks demonstrated a decrease in inflammatory markers (i.e, TNF, and IL-6), total

cholesterol, triglycerides, as well as low-density lipoprotein and an increase in total antioxidant

capacity, glutathione peroxidase activity, as well as superoxide dismutase enzymes (Ghadimi et

al., 2021). ROS are crucial signaling molecules that play a vital role in the course of

inflammatory diseases. EA effectively reduces endothelial ROS levels and improves arterial

relaxation impairment caused by high glucose in a model of intact rat aortas and human aortic

endothelial cells activated with high glucose concentration (Rozentsvit et al., 2017). A wide body

of literature demonstrated that polyphenolic compounds can be helpful as adjuvant therapy due

to their potential anti-inflammatory effects, antioxidant activity, and inhibition of enzymes

involved in eicosanoids formation (Hussain et al., 2016). It is within reasons to speculate that EA

might counteract inflammation and tissue damage via this pathway, especially when more recent

evidence also suggests that EA, in a model of zebrafish embryonic development, might provide

protective effects against DNA oxidative damage while simultaneously increasing the embryo

survival rate and improving morphological parameters of the larvae (Mottola et al., 2020).

Interestingly, the vesicle lumen was identified by GO Cellular Compartment analysis as a

potential cellular compartment where all these processes might take place. Aforementioned

results of previous studies are highly related to cellular production and release of inflammatory

mediators, a critical component of vesicle lumen (Türk et al., 2010). Previous network

pharmacology utilizing bioinformatics also predicted vesicle lumens as potential target of EA in

the model of endometriosis (Wu et al., 2022) and beta amyloid 25-35 peptide-induced

neurotoxicity (Li et al., 2022). Interestingly, EA was also predicted by GO Cellular Function to

potentially be involved in ion binding process. Within the scope of inflammation, H+/K+

ATPase activity, known to be stimulated by inflammatory and other factors and by the bacterium

Helicobacter pylori (Z. Zhang et al., 2020), was demonstrated to be inhibited by EA hog gastric

with a 50% inhibition at 2.1×10^{-6} M (Murakami et al., 1991), postulating a compelling case to

study the intersection of EA, inflammatory mediators, vesicle lumens, and ion exchange process.

The 30 most significant KEGG enriched pathways are illustrated in **Figure 2B**. Notably,

cancer-related pathways include: pathways in cancer, prostate cancer, microRNAs in cancer,

chemical carcinogenesis, pancreatic cancer, acute myeloid leukemia, chronic myeloid leukemia,

bladder cancer, breast cancer, and small cell lung cancer (**Figure 2B**). Cancer and inflammation

have a uniquely inseparable relationship: cancer growth and progression are strongly associated

with inflammation, which, in turn, can affect cancer formation, metastasis or response to therapy

(Singh et al., 2019; Zhao et al., 2021). Recent evidence from *in vitro* and *in vivo* studies have

suggested that EA may be able to inhibit tumor cell proliferation, induce apoptosis, break DNA

binding to carcinogens, block virus infection, and disturb inflammation, angiogenesis, and drug-

resistance processes, which are all important factors in the development and progression of

cancer (H.-M. Zhang et al., 2014). Other pathways relevant to inflammation also include

Hepatitis B, FoxO, TNF, and HIF-1 signaling pathways. In conjunction, it was demonstrated that

EA could effectively inhibit Hepatitis B-infected HepG2 2.2.15 cells from secreting hepatitis B

e-antigen (Shin et al., 2005). In a model of high glucose-induced injury in rat mesangial cells,

EA was demonstrated to inhibit the activation of the PI3K/Akt signaling pathway and reduce the

expression levels of downstream transcription factor FOXO3a (W. Lin et al., 2021). In a model

of 1,2-dimethylhydrazine-induced rat colon carcinogenesis, EA downregulated iNOS, PTGS2,

TNF and IL6 via inhibition of NF-kB and demonstrated chemopreventive effects on colon

carcinogenesis (Umesalma & Sudhandiran, 2010). Lastly, in a study using *in vitro* model of lung

cancer (i.e., HOP62 and H1975 cell lines) EA decreased both mRNA expression of target genes

of HIF-1 as well as the protein level of HIF-1, whose inhibition may enhance the effectiveness of

a vast variety of innovative and established anticancer medicines (Duan et al., 2020).

To further determine critical proteins in EA's anti-inflammatory activity, topological

analysis was performed (**Figure 2C-F**) using cytoHubba MCC, DMNC, MNC, and EPC

algorithms. The MCC algorithm determined PTGS2, MMP9, ALB, and TNF as the 4 most

essential proteins of the network (**Figure 2C**). Alternatively, the DMNC algorithm determined

ICAM1, MMP2, SELP, and MMP9 as the 4 most essential proteins of the network (**Figure 2D**).

Furthermore, the MNC algorithm determined MAPK3, TNF, AKT1, and VEGFA as the 4 most

essential proteins of the network (**Figure 2E**). Lastly, the EPC algorithm determined MAPK3,

TNF, AKT1, and VEGFA as the 4 most essential proteins of the network (**Figure 2F**). In

essence, per the recommendation of Chin et al., 2014, combining the results of these 4

algorithms determined that the top 10 most important proteins of the network are PTGS2,

MMP9, MMP2, ALB, TNF, ICAM1, SELP, MAPK3, VEGFA, and AKT1. PTGS2 is an enzyme

that plays a role in bio-transforming arachidonic acid to prostaglandins, which are signaling

molecules that have various functions in the body, including the regulation of inflammation

(Simon, 1999). MMP9, in the context of inflammation, is often upregulated and released by both

immune cells (i.e, neutrophils and monocytes) as well as endothelial or fibroblast cells (Al-Sadi

et al., 2021; Yabluchanskiy et al., 2013). Notably, MMP9 was demonstrated maintain epithelial

barrier function and preserve mucosal lining in colitis associated cancer in a model of murine

colonic epithelium and CaCo2BBE intestinal cell line (Pujada et al., 2017). The upregulation and

release of MMP9 can eventually contribute to the breakdown of extracellular matrix proteins and

other proinflammatory molecules, leading to tissue damage. In a similar manner, MMP2 is a

protease that can cleave several extracellular matrix proteins, including collagens, laminins, and

elastins and is important for various physiological processes, such as tissue repair and

remodeling, as well as pathological processes such as cancer cell invasion and metastasis

(Maybee et al., 2022; Song et al., 2021). Inadequacy of ALB is associated with inflammatory

conditions (Soeters et al., 2019); a clinical study demonstrated that ALB is negatively correlated

with inflammatory indices (i.e, C-reactive protein and white blood cell count), implying a

potential role of ALB in the regulation of inflammatory and immune responses (Sheinenzon et

al., 2021). TNF mediates inflammation by activating immune cells and inducing the expression

of other cytokines, chemokines, and adhesion molecules and eventually clears infections and in

the process of tissue repair after injury, but TNF can also be the driving force of chronic

inflammation, leading to tissue damage and the development of autoimmune diseases (Chu,

2013; Kalliolias & Ivashkiv, 2016). Within the same vein, EA reversed the release of IL-6 and

IL-8 induced by TNF in Caco-2 cell (Iglesias et al., 2020). ICAM1 is a type of cell adhesion

molecule that is expressed on the surface of various types of cells, including epithelial cells,

endothelial cells, and immune cells and plays a critical role in the inflammatory response by

mediating the adhesion of leukocytes to the endothelium (Bui et al., 2020). SELP is a type of cell

adhesion molecule that is expressed on the surface of stimulated endothelial cells and activated

platelets; SELP can mediate the interaction between leukocytes and endothelial cells, allowing

leukocytes to "roll" along the surface of the endothelium, which is important for the recruitment

of leukocytes to sites of inflammation (M. Chen & Geng, 2006). MAPK3 plays a key role in

regulating various cellular processes, including cell proliferation, differentiation, and survival

and has been shown to be upregulated in immune cells during inflammation, and its activation

also is associated with the production of pro-inflammatory mediators, such as cytokines and

chemokines (Lucas et al., 2022). In inflammation, VEGFA is important for recruiting of immune

cells to the site of inflammation, forming of new blood vessels, maintaining the integrity of

blood vessels, wound healing, and the development of the placenta during pregnancy; however,

excessive, or uncontrolled production of VEGFA can contribute to the development of certain

diseases, such as cancer (Scaldaferri et al., 2009; Shaik-Dasthagirisaheb et al., 2013). Lastly,

activation of AKT1 is known to lead to the activation of NF-κB, which in turn upregulates and

encodes cytokines, chemokines, and adhesion molecules that play critical roles in inflammatory

responses (Luthra et al., 2008; Zhou et al., 2013). EA might very well directly interact with these

targets to exert its anti-inflammatory effects, as previous studies demonstrated that EA could

affect the gene as well as protein expression of these targets (Attilio et al., 2010; Kowshik et al.,

2014; C. Lin et al., 2019; Pattanayak et al., 2017; Umesalma & Sudhandiran, 2010; Yu et al.,

2007). Therefore, molecular docking and molecular dynamics simulations were utilized to

validate said speculation.

2. Molecular Docking and Molecular Dynamics Simulation Studies

245   **Table 1.** Results of molecular docking studies, sorted by binding affinities from strongest to

246   weakest.

Target	Binding Affinity (kcal/mol)	Predicted Dissociation Constant (pM)	Contact surface (Å^2)
AKT1 (PDB: 6HHG)	-9.894	55,924.5	249.81
MAPK3 (PDB: 4QTB)	-9.452	117,921.3	254.77
PTGS2 (PDB: 5F19)	-9.359	137,962.8	255.40
MMP9 (PDB: 1L6J)	-8.916	291,397.3	230.30
ALB (PDB: 6M4R)	-8.502	586,072.6	265.98
MMP2 (PDB: 7XJO)	-7.111	6,131,516.5	208.28
SELP (PDB: 1G1S)	-6.900	8,754,532.0	159.0
ICAM1 (PDB: 1IAM)	-6.871	10,054,020.0	169.00
TNF (PDB: 2AZ5)	-6.479	17,816,814.0	173.73
VEGFA (PDB: 4KZN)	-6.160	30,525,314.0	154.23

Results of molecular docking (binding affinity, predicted Ki, and contact surface) are

summarized in **Table 1**. In brief, from the molecular docking results, EA had the strongest

binding affinities for AKT1, MAPK3, PTGS2 (with -9.894, -9.452, and -9.359 kcal/mol,

respectively) and weakest binding affinities for ICAM1, TNF, and VEGFA (with -6.871, -6.479,

-6.160 kcal/mol, respectively) (**Table 1**).However, the crucial analyses of this study rely on the

ligand contact analysis (**Figure 3**) and MM/PBSA-calculated binding energies (**Figure 4A**) by

means of molecular dynamics simulations. EA was able to maintain continuous contacts with all

important target protein throughout 100 ns of molecular dynamics simulations (**Figure 3**). From

MM-PBSA binding energy calculations, EA demonstrated strongest binding affinities with

AKT1, PTGS2, VEGFA, MAPK3 with -82.37 ± 159.8, -52.06 ± 171.4, -19.24 ±79.02, and -

13.98 ± 149.6 kJ/mol, respectively (**Figure 4A**). EA demonstrated weaker binding with ALB,

MMP9, ICAM1, TNF, MMP2, and SELP with -9.597 ± 223.9, 16.84 ± 169.2, 29.11 ± 169.2,

46.92 ± 80.85, 54.92 ± 81.22, and 59.51 ± 92.92 kJ/mol, respectively (**Figure 4A**). However,

these results do not necessarily negate ALB, MMP9, ICAM1, TNF, MMP2, and SELP as

potential targets of EA, as it only highlights that AKT1, PTGS2, VEGFA, and MAPK3 are the

most likely targets of EA. However, further analyses for the complexes with the highest

MM/PBSA binding energies somewhat corroborate this result (**Figure 4B**). EA showed binding

energies towards MAP3K and VEGFA that are comparable to known binders, except AKT1 and

PTGS2. Specifically, the binding energies of 10DEBC (for AK± 171.2 andT1), resveratrol (for

MAP3K), ibuprofen (for PTGS2), and sorafenib (for VEGFA) are -265.2 ± 161.2, -24.12 ±

145.2, -215.1 ± 171.2, and -41.76 ± 81.92. However, this does not necessarily suggest thar EA is

a non-binder for AKT1 or PTGS2, as it could potentially act a mild modulator, providing

complementary or preventative effects. Further molecular dynamics simulation analyses are

provided as Figure S1-S5.

To reiterate, AKT1, PTGS2, VEGFA, MAPK3 had been demonstrated to play critical

roles in progression of cancer, inflammation, HIF-1 signaling, and TNF signaling (Faurschou &

Gniadecki, 2008; Fojtík et al., 2021; Guadagni et al., 2007; Lee et al., 2010; C.-C. Lin et al.,

2004; Ramakrishnan et al., 2014; Siddappa et al., 2015; Stegeman et al., 2016). Thus, they,

through topological analysis and molecular dynamics simulations, are most likely to be targets of

EA for its anti-inflammatory activities and are included in the illustration of putative anti-

inflammatory mechanism of EA (**Figure 5**). Particularly, TNF-alpha, regulated by NF-kB and

responsible for the progression of cancer (Guadagni et al., 2007), was demonstrated by to

stimulate PI3K-Akt in keratinocyte cell lines HaCaT and A431 cell lines, eventually resulting in

enhanced mutagenesis and tumor development (Faurschou & Gniadecki, 2008). Compellingly,

ellagic acid was demonstrated to provide protective effects against high glucose-induced injury

in in rat mesangial cells via the PI3K/Akt/FOXO3a signaling pathway (W. Lin et al., 2021),

which again further supports the findings of this study with predictive nature. Hypoxia and mild

hypoxia cellular environments were demonstrated to increase PI3K-Akt and MAPK signaling

and decrease PI3K-Akt signaling (respectively) in human pluripotent stem cells, demonstrating a

unique but intimate relationship between hypoxia and PI3K-Akt and MAPK signaling (Fojtík et

al., 2021). Within the same vein, in in vivo experiments using human esophageal epithelial cells

(EPC2-hTERT), hypoxia conditions also upregulate PTGS2, but hypoxia conditions with

cotreatment of IL-1β prevented said upregulation, demonstrating an intimate relationship

between hypoxia, PTGS2, and inflammation. HIF-1 signaling pathway is also known for being

the upstream regulator of VEGF (which is a critical biomarker for infection, cancer, and

cardiovascular diseases). Furthermore, hypoxic conditions also induced AKT1 in neck squamous

cell carcinoma and non-small-cell lung cancer cell lines (Stegeman et al., 2016). The nuance is

also further extended, as it was previously demonstrated that MAP3K can potentially regulate

ovulation by targeting PTGS2 (also a target of HIF-1 signaling pathway) in murine ovulating

follicles (Siddappa et al., 2015). In short, we believe that EA might play a critical role in

attenuating multiple "ends" or "streams" of these interconnected relationships. However, due to

its, low hydrolysis and lipophilic and hydrophobic nature, EA is not readily absorbed in

circulation but rather concentrated in the intestine (Mohammadinejad et al., 2022). We believe

the intestine would be a great target for the investigation of animal or human models for the

effects of EA dietary consumption on inflammation or diseases with inflammatory signatures.

Conclusion

In summary, a network pharmacology approach was used to understand the anti-inflammatory

effects of EA. It was revealed that there are 52 targets of EA that are related to inflammation.

Topological analysis determined 10 important targets that are AKT1, VEGFA, TNF, MAPK3,

ALB, SELP, MMP9, MMP2, PTGS2, and ICAM1. Molecular docking and molecular dynamics

simulations helped determine AKT1, PTGS2, VEGFA, MAPK3 as the most probable targets of

EA. Computational studies need further experimental validations; however, this study

corroborates with previous studies and provides potential targets and mechanisms for future

research.

Methods

1.   Retrieval of EA target proteins

314   Target proteins of EA were retrieved from SwissTargetPrediction (Daina et al., 2019) and

SymMap v2 (Wu et al., 2019) in UniProt ID format. String-db.org (Szklarczyk et al., 2021) was

used to exclude any duplicate and resulted in a total of 157 EA target proteins.

2.   Retrieval of target proteins for inflammation

DisGeNET (Piñero et al., 2019) was used to retrieve target inflammation proteins in UniProt ID

format via the following accession numbers: C0021368 ("inflammation"), C0234251

("inflammatory pain"), and C1290884 ("inflammatory disorder"). String-db.org was also used to

exclude any duplicate and resulted in a total of 886 target proteins for inflammation.

3.   Constructing PPI network of EA target proteins for anti-inflammatory properties

After two networks of 157 EA target proteins and 886 target proteins for inflammation were

constructed on string-db.org, they were then sent to Cytoscape 3.9.1 (Shannon et al., 2003) to

find mutual proteins between the two networks via the Merge-Intersection function, resulting in

52 EA target proteins that are also relevant for its anti-inflammatory properties. This

visualization of this network and its basic topological analysis were then provided through

string-db.org at a moderate confidence score of 0.55. Within string-db.org, GO enrichment

analysis was also retrieved; 5 selected enriched terms with the lowest false discovery rate for

each GO enrichment category (Biological Processes, Molecular Function, Cellular Component)

were presented. On the other hand, ShinyGO 0.76.3 (Ge et al., 2020) was used to perform and

visualize KEGG enrichment analysis.

4.   Topological analysis of the PPI

Both local-based (i.e, MCC, DMNC, and MNC) and global-based (i.e, EPC) algorithms from the

cytoHubba plug-in of Cytoscape were used to determine the most critical target proteins. The

mathematical basis for these algorithms can be found at Chin et al., 2014. A protein is deemed

significant if it ranks among the top four with the highest score in at least one category (i.e,

MCC, DMNC, MNC, and EPC).

5.   Molecular Docking

YASARA Structure (Land & Humble, 2018) was used to perform both molecular docking and

molecular dynamics simulations. The dock_run.mcr script, which utilized Vina algorithm, was

employed to perform molecular docking for EA towards important protein targets for

inflammation. For each protein target, 100 docking runs were performed, and the docked pose

with the most negative binding affinity (ΔG) in kcal/mol was selected for further validation

through molecular dynamics simulations.

6.   Molecular Dynamics Simulations

The md_run.mcr script was employed to perform molecular dynamics simulations for docked

complexes to investigate the interactions between EA and its critical target proteins. In brief, the

important parameters for the simulations are as follows: pH is 7.4; ion concentration is 0.9%

NaCl; temperature is 298 K; water density is 0.997 g/ml; pressure control mode is solvent probe;

cell shape is cube with periodic boundary; force field is AMBER14. The simulation is speed is

'normal,' in which a mixed-multiple timestep (of 2*1.25 femtoseconds) algorithm combined

with LINCS algorithm is utilized. Eventually, analyses of contacts between EA and target

proteins as well as movement RMSD of EA were retrieved through the md_analyze.mcr script.

Binding energies derived from MM/PBSA were calculated using md_analyzebindenergy.mcr

script, based on the following premise:

$$\Delta G\text{-binding} = \Delta G\text{-complex} - (\Delta G\text{-protein} + \Delta G\text{-ligand}) \text{ (in kJ/mol)}$$

In which, ΔG-complex is calculated based on the solvation and potential energies of the

complex; ΔG-protein is calculated based on the solvation and potential energies of the protein,

and ΔG-ligand is calculated based on the solvation and potential energies of the ligand. More

negative energies indicate better binding; positive energies do not indicate no binding. Top 4

complex with the highest MM/PBSA binding are further compared with known binders for

validation.

Data Availability

All predictive and enrichment data can be found in the Supplementary Excel File online. Sheet 1

corresponds to DisGeNet C0021368 ("inflammation"). Sheet 2 corresponds to C0234251

("inflammatory pain"). Sheet 3 corresponds to C1290884 ("inflammatory disorder"). Sheet 4

corresponds to SymMap target of EA. Sheet 5 corresponds to SwissTargetPrediction target of

EA. Sheet 6 corresponds to 52 targets of EA for inflammation. Sheet 7 corresponds to GO

Biological Process enrichment analysis. Sheet 8 corresponds to GO Cellular Component

enrichment analysis. Sheet 9 corresponds to GO Molecular Function enrichment analysis. Sheet

10 corresponds to KEGG enrichment analysis.

Figure Captions

Figure 1. Venn diagram showing the overlap between 157 target proteins of ellagic acid (EA)

and 886 target proteins for inflammation as well as the protein-protein interaction network of 52

overlap proteins via string-db.org.

Figure 2. Enrichment analysis and topological analysis of the protein-protein interaction.

network. (A) Gene Ontology (GO) enrichment analysis. (B) Kyoto Encyclopedia of Genes and

Genomes (KEGG) enrichment analysis. Topological analysis to determine important proteins of

the network via (C) Maximal Clique Centrality (MCC) (D) Density of Maximum Neighborhood

Component (DMNC) (E) Maximum Neighborhood Component (MNC) (F) Edge Percolated

Component (EPC)

Figure 3. Residue contacts between ellagic acid (EA) and important proteins of the network

determined through topological analysis demonstrating that EA can interact with putative target

100% of the time of the simulations. Red, green, and yellow colors denote hydrophobic

interaction, hydrogen bonding, and the simultaneous combination of hydrophobic interaction and

hydrogen bonding, respectively.

Figure 4. (A) Binding energies calculated by molecular mechanics Poisson-Boltzmann surface

area (MM-PBSA) method, represented in violin plots in comparison with EA-protein complexes

with the highest molecular docking binding energies. (B) Binding energies calculated by

molecular mechanics Poisson-Boltzmann surface area (MM-PBSA) method, represented in

violin plots in comparison with known binders.

Figure 5. Illustrations of EA affecting inflammation, HIF-1 signaling, TNF signaling, and cancer

pathways.

Figure S1. Molecular dynamics simulation analyses for AKT1 complexes. RMSD: root-mean-

square distance.

Figure S2. Molecular dynamics simulation analyses for MAP3K complexes. RMSD: root-mean-

square distance

Figure S3. Molecular dynamics simulation analyses for PTGS2 complexes. RMSD: root-mean-

square distance

Figure S4. Molecular dynamics simulation analyses for VEGFA complexes. RMSD: root-mean-

square distance.

Figure S5. Binding modes of EA (red) and known binders (blue) towards their target proteins

(yellow) at the 100th ns of the simulations.

Author contribution

SHH planned the study, performed bioinformatics analysis, and wrote the original draft. SHH,

EML, and HD revised the manuscript and performed additional bioinformatics analysis. All

authors have read the manuscript and agreed upon submission.

Disclosure Statement

The authors declare no conflict of interest and being the sole contributor of this work as well as

no funding is associated with this work.

References

Al-Sadi, R., Engers, J., Haque, M., King, S., Al-Omari, D., & Ma, T. Y. (2021). Matrix
Metalloproteinase-9 (MMP-9) induced disruption of intestinal epithelial tight junction
barrier is mediated by NF-κB activation. *PLOS ONE, 16*(4), e0249544.
https://doi.org/10.1371/journal.pone.0249544

Amor, A. J., Gómez-Guerrero, C., Ortega, E., Sala-Vila, A., & Lázaro, I. (2020). Ellagic Acid as a
Tool to Limit the Diabetes Burden: Updated Evidence. *Antioxidants (Basel, Switzerland)*,
9(12), 1226. https://doi.org/10.3390/antiox9121226

Attilio, P., Merritt, C., Sims, J., Kane, N., & O'Sullivan, J. (2010). The effect of ellagic acid on
platelet activation as measured by the quantification of P-selectin using flow cytometry.
AANA Journal, 78(6), 453–459.

Baliga, M. S., Shivashankara, A. R., Venkatesh, S., Bhat, H. P., Palatty, P. L., Bhandari, G., &
Rao, S. (2019). Phytochemicals in the Prevention of Ethanol-Induced Hepatotoxicity. In
Dietary Interventions in Liver Disease (pp. 79–89). Elsevier.
https://doi.org/10.1016/B978-0-12-814466-4.00007-0

BenSaad, L. A., Kim, K. H., Quah, C. C., Kim, W. R., & Shahimi, M. (2017). Anti-inflammatory
potential of ellagic acid, gallic acid and punicalagin A&B isolated from Punica granatum.
BMC Complementary and Alternative Medicine, *17*(1), 47.
https://doi.org/10.1186/s12906-017-1555-0

Bosma-den Boer, M. M., van Wetten, M.-L., & Pruimboom, L. (2012). Chronic inflammatory
diseases are stimulated by current lifestyle: How diet, stress levels and medication
prevent our body from recovering. *Nutrition & Metabolism*, *9*(1), 32.
https://doi.org/10.1186/1743-7075-9-32

Bui, T. M., Wiesolek, H. L., & Sumagin, R. (2020). ICAM-1: A master regulator of cellular

responses in inflammation, injury resolution, and tumorigenesis. *Journal of Leukocyte*

Biology, *108*(3), 787–799. https://doi.org/10.1002/JLB.2MR0220-549R

Ceci, C., Lacal, P. M., Tentori, L., De Martino, M. G., Miano, R., & Graziani, G. (2018).

Experimental Evidence of the Antitumor, Antimetastatic and Antiangiogenic Activity of

Ellagic Acid. *Nutrients*, *10*(11), 1756. https://doi.org/10.3390/nu10111756

Chen, L., Deng, H., Cui, H., Fang, J., Zuo, Z., Deng, J., Li, Y., Wang, X., & Zhao, L. (2018).

Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget*,

9(6), 7204–7218. https://doi.org/10.18632/oncotarget.23208

Chen, M., & Geng, J.-G. (2006). P-selectin mediates adhesion of leukocytes, platelets, and

cancer cells in inflammation, thrombosis, and cancer growth and metastasis. *Archivum*

Immunologiae et Therapiae Experimentalis, *54*(2), 75–84.

https://doi.org/10.1007/s00005-006-0010-6

Chin, C.-H., Chen, S.-H., Wu, H.-H., Ho, C.-W., Ko, M.-T., & Lin, C.-Y. (2014). cytoHubba:

Identifying hub objects and sub-networks from complex interactome. *BMC Systems*

Biology, *8 Suppl 4*(Suppl 4), S11. https://doi.org/10.1186/1752-0509-8-S4-S11

Chu, W.-M. (2013). Tumor necrosis factor. *Cancer Letters*, *328*(2), 222–225.

https://doi.org/10.1016/j.canlet.2012.10.014

Daina, A., Michielin, O., & Zoete, V. (2019). SwissTargetPrediction: Updated data and new

features for efficient prediction of protein targets of small molecules. *Nucleic Acids*

Research, *47*(W1), W357–W364. https://doi.org/10.1093/nar/gkz382

Del Grossi Moura, M., Cruz Lopes, L., Silva, M. T., Barberato-Filho, S., Motta, R. H. L., &

Bergamaschi, C. de C. (2018). Use of steroid and nonsteroidal anti-inflammatories in the

treatment of rheumatoid arthritis: Systematic review protocol. *Medicine*, *97*(41), e12658.

https://doi.org/10.1097/MD.0000000000012658

Duan, J., Li, Y., Gao, H., Yang, D., He, X., Fang, Y., & Zhou, G. (2020). Phenolic compound

ellagic acid inhibits mitochondrial respiration and tumor growth in lung cancer. *Food &*

Function, *11*(7), 6332–6339. https://doi.org/10.1039/D0FO01177K

Fang, P., Li, X., Dai, J., Cole, L., Camacho, J. A., Zhang, Y., Ji, Y., Wang, J., Yang, X.-F., &

Wang, H. (2018). Immune cell subset differentiation and tissue inflammation. *Journal of*

Hematology & Oncology, *11*(1), 97. https://doi.org/10.1186/s13045-018-0637-x

Faurschou, A., & Gniadecki, R. (2008). TNF-α stimulates Akt by a distinct aPKC-dependent

pathway in premalignant keratinocytes. *Experimental Dermatology*, *17*(12), 992–997.

https://doi.org/10.1111/j.1600-0625.2008.00740.x

Fojtík, P., Beckerová, D., Holomková, K., Šenfluk, M., & Rotrekl, V. (2021). Both Hypoxia-

Inducible Factor 1 and MAPK Signaling Pathway Attenuate PI3K/AKT via Suppression of

Reactive Oxygen Species in Human Pluripotent Stem Cells. *Frontiers in Cell and*

Developmental Biology, *8*, 607444. https://doi.org/10.3389/fcell.2020.607444

Furman, D., Campisi, J., Verdin, E., Carrera-Bastos, P., Targ, S., Franceschi, C., Ferrucci, L.,

Gilroy, D. W., Fasano, A., Miller, G. W., Miller, A. H., Mantovani, A., Weyand, C. M.,

Barzilai, N., Goronzy, J. J., Rando, T. A., Effros, R. B., Lucia, A., Kleinstreuer, N., &

Slavich, G. M. (2019). Chronic inflammation in the etiology of disease across the life

span. *Nature Medicine*, *25*(12), 1822–1832. https://doi.org/10.1038/s41591-019-0675-0

Ge, S. X., Jung, D., & Yao, R. (2020). ShinyGO: A graphical gene-set enrichment tool for

animals and plants. *Bioinformatics (Oxford, England)*, *36*(8), 2628–2629.

https://doi.org/10.1093/bioinformatics/btz931

Ghadimi, M., Foroughi, F., Hashemipour, S., Rashidi Nooshabadi, M., Ahmadi, M. H., Ahadi

Nezhad, B., & Khadem Haghighian, H. (2021). Randomized double-blind clinical trial

examining the Ellagic acid effects on glycemic status, insulin resistance, antioxidant, and

inflammatory factors in patients with type 2 diabetes. *Phytotherapy Research*, *35*(2),

1023–1032. https://doi.org/10.1002/ptr.6867

Glass, C. K., & Olefsky, J. M. (2012). Inflammation and lipid signaling in the etiology of insulin

resistance. *Cell Metabolism*, *15*(5), 635–645. https://doi.org/10.1016/j.cmet.2012.04.001

González-Sarrías, A., Espín, J.-C., Tomás-Barberán, F. A., & García-Conesa, M.-T. (2009).

Gene expression, cell cycle arrest and MAPK signalling regulation in Caco-2 cells

exposed to ellagic acid and its metabolites, urolithins. *Molecular Nutrition & Food*

Research, *53*(6), 686–698. https://doi.org/10.1002/mnfr.200800150

Guadagni, F., Ferroni, P., Palmirotta, R., Portarena, I., Formica, V., & Roselli, M. (2007).

Review. TNF/VEGF cross-talk in chronic inflammation-related cancer initiation and

progression: An early target in anticancer therapeutic strategy. *In Vivo (Athens, Greece)*,

21(2), 147–161.

Hussain, T., Tan, B., Yin, Y., Blachier, F., Tossou, M. C. B., & Rahu, N. (2016). Oxidative Stress

and Inflammation: What Polyphenols Can Do for Us? *Oxidative Medicine and Cellular*

Longevity, *2016*, 7432797. https://doi.org/10.1155/2016/7432797

Iglesias, D. E., Cremonini, E., Fraga, C. G., & Oteiza, P. I. (2020). Ellagic acid protects Caco-2

cell monolayers against inflammation-induced permeabilization. *Free Radical Biology &*

Medicine, *152*, 776–786. https://doi.org/10.1016/j.freeradbiomed.2020.01.022

Kabbany, M. T., Joshi, A. A., & Mehta, N. N. (2016, July 18). *Cardiovascular Diseases in*

Chronic Inflammatory Disorders [American College of Cardiology].

https://www.acc.org/latest-in-cardiology/articles/2016/07/15/10/04/cardiovascular-

diseases-in-chronic-inflammatory-

disorders#:~:text=Patients%20with%20chronic%20inflammatory%20diseases,aging%2C

%20hypertension%2C%20and%20smoking

Kalliolias, G. D., & Ivashkiv, L. B. (2016). TNF biology, pathogenic mechanisms and emerging

therapeutic strategies. *Nature Reviews Rheumatology*, *12*(1), 49–62.

https://doi.org/10.1038/nrrheum.2015.169

Kang, I., Buckner, T., Shay, N. F., Gu, L., & Chung, S. (2016). Improvements in Metabolic

Health with Consumption of Ellagic Acid and Subsequent Conversion into Urolithins:

Evidence and Mechanisms. *Advances in Nutrition (Bethesda, Md.)*, *7*(5), 961–972.

https://doi.org/10.3945/an.116.012575

Kowshik, J., Giri, H., Kishore, T., Kesavan, R., Vankudavath, R., Reddy, G., Dixit, M., & Nagini,

S. (2014). Ellagic Acid Inhibits VEGF/VEGFR2, PI3K/Akt and MAPK Signaling Cascades

in the Hamster Cheek Pouch Carcinogenesis Model. *Anti-Cancer Agents in Medicinal*

Chemistry, *14*(9), 1249–1260. https://doi.org/10.2174/1871520614666140723114217

Land, H., & Humble, M. S. (2018). YASARA: a tool to obtain structural guidance in biocatalytic

investigations. In *Protein Engineering* (pp. 43–67). Springer.

Lee, J. J., Natsuizaka, M., Ohashi, S., Wong, G. S., Takaoka, M., Michaylira, C. Z., Budo, D.,

Tobias, J. W., Kanai, M., Shirakawa, Y., Naomoto, Y., Klein-Szanto, A. J. P., Haase, V.

H., & Nakagawa, H. (2010). Hypoxia activates the cyclooxygenase-2-prostaglandin E

synthase axis. *Carcinogenesis*, *31*(3), 427–434. https://doi.org/10.1093/carcin/bgp326

Li, H.-L., Zhang, S.-Y., Ren, Y.-S., Zhou, J.-C., Zhou, Y.-X., Huang, W.-Z., Piao, X.-H., Yang, Z.-

Y., Wang, S.-M., & Ge, Y.-W. (2022). Identification of ellagic acid and urolithins as

natural inhibitors of Aβ25-35-induced neurotoxicity and the mechanism predication using

network pharmacology analysis and molecular docking. *Frontiers in Nutrition*, *9*, 966276.

https://doi.org/10.3389/fnut.2022.966276

Lin, C., Wei, D., Xin, D., Pan, J., & Huang, M. (2019). Ellagic acid inhibits proliferation and

migration of cardiac fibroblasts by down-regulating expression of HDAC1. *The Journal of*

Toxicological Sciences, *44*(6), 425–433. https://doi.org/10.2131/jts.44.425

Lin, C.-C., Hsiao, L.-D., Chien, C.-S., Lee, C.-W., Hsieh, J.-T., & Yang, C.-M. (2004). Tumor

necrosis factor-α-induced cyclooxygenase-2 expression in human tracheal smooth

muscle cells: Involvement of p42/p44 and p38 mitogen-activated protein kinases and

nuclear factor-κB. *Cellular Signalling*, *16*(5), 597–607.

https://doi.org/10.1016/j.cellsig.2003.10.002

Lin, W., Liu, G., Kang, X., Guo, P., Shang, Y., Du, R., Wang, X., Chen, L., Yue, R., Kong, F., & Zhu, Q. (2021). Ellagic acid inhibits high glucose-induced injury in rat mesangial cells via the PI3K/Akt/FOXO3a signaling pathway. *Experimental and Therapeutic Medicine*, *22*(3), 1017. https://doi.org/10.3892/etm.2021.10449

Lucas, R. M., Luo, L., & Stow, J. L. (2022). ERK1/2 in immune signalling. *Biochemical Society Transactions*, *50*(5), 1341–1352. https://doi.org/10.1042/BST20220271

Luster, A. D., Alon, R., & von Andrian, U. H. (2005). Immune cell migration in inflammation: Present and future therapeutic targets. *Nature Immunology*, *6*(12), 1182–1190. https://doi.org/10.1038/ni1275

Luthra, P., Sun, D., Wolfgang, M., & He, B. (2008). AKT1-Dependent Activation of NF-κB by the L Protein of Parainfluenza Virus 5. *Journal of Virology*, *82*(21), 10887–10895. https://doi.org/10.1128/JVI.00806-08

Margină, D., Ungurianu, A., Purdel, C., Tsoukalas, D., Sarandi, E., Thanasoula, M., Tekos, F., Mesnage, R., Kouretas, D., & Tsatsakis, A. (2020). Chronic Inflammation in the Context of Everyday Life: Dietary Changes as Mitigating Factors. *International Journal of Environmental Research and Public Health*, *17*(11), 4135. https://doi.org/10.3390/ijerph17114135

Maybee, D. V., Ink, N. L., & Ali, M. A. M. (2022). Novel Roles of MT1-MMP and MMP-2: Beyond the Extracellular Milieu. *International Journal of Molecular Sciences*, *23*(17), 9513. https://doi.org/10.3390/ijms23179513

Mohammadinejad, A., Mohajeri, T., Aleyaghoob, G., Heidarian, F., & Kazemi Oskuee, R. (2022). Ellagic acid as a potent anticancer drug: A comprehensive review on in vitro, in vivo, in silico, and drug delivery studies. *Biotechnology and Applied Biochemistry*, *69*(6), 2323–2356. https://doi.org/10.1002/bab.2288

Mottola, F., Scudiero, N., Iovine, C., Santonastaso, M., & Rocco, L. (2020). Protective activity of

ellagic acid in counteract oxidative stress damage in zebrafish embryonic development.

Ecotoxicology and Environmental Safety, *197*, 110642.

https://doi.org/10.1016/j.ecoenv.2020.110642

Murakami, S., Isobe, Y., Kijima, H., Nagai, H., Muramatu, M., & Otomo, S. (1991). Inhibition of

gastric H+, K(+)-ATPase and acid secretion by ellagic acid. *Planta Medica*, *57*(4), 305–

308. https://doi.org/10.1055/s-2006-960103

Päivärinta, E., Pajari, A.-M., Törrönen, R., & Mutanen, M. (2006). Ellagic Acid and Natural

Sources of Ellagitannins as Possible Chemopreventive Agents Against Intestinal

Tumorigenesis in the Min Mouse. *Nutrition and Cancer*, *54*(1), 79–83.

https://doi.org/10.1207/s15327914nc5401_9

Pattanayak, R., Basak, P., Sen, S., & Bhattacharyya, M. (2017). An insight to the binding of

ellagic acid with human serum albumin using spectroscopic and isothermal calorimetry

studies. *Biochemistry and Biophysics Reports*, *10*, 88–93.

https://doi.org/10.1016/j.bbrep.2017.03.001

Piñero, J., Ramírez-Anguita, J. M., Saüch-Pitarch, J., Ronzano, F., Centeno, E., Sanz, F., &

Furlong, L. I. (2019). The DisGeNET knowledge platform for disease genomics: 2019

update. *Nucleic Acids Research*, gkz1021. https://doi.org/10.1093/nar/gkz1021

Promsong, A., Chung, W. O., Satthakarn, S., & Nittayananta, W. (2015). Ellagic acid modulates

the expression of oral innate immune mediators: Potential role in mucosal protection.

Journal of Oral Pathology & Medicine, *44*(3), 214–221. https://doi.org/10.1111/jop.12223

Pujada, A., Walter, L., Patel, A., Bui, T. A., Zhang, Z., Zhang, Y., Denning, T. L., & Garg, P.

(2017). Matrix metalloproteinase MMP9 maintains epithelial barrier function and

preserves mucosal lining in colitis associated cancer. *Oncotarget*, *8*(55), 94650–94665.

https://doi.org/10.18632/oncotarget.21841

Ramakrishnan, S., Anand, V., & Roy, S. (2014). Vascular endothelial growth factor signaling in hypoxia and inflammation. *Journal of Neuroimmune Pharmacology: The Official Journal of the Society on NeuroImmune Pharmacology*, *9*(2), 142–160. https://doi.org/10.1007/s11481-014-9531-7

Ridker, P. M. (2017). Inflammation, cardiovascular disease and cancer: Moving toward predictive medicine. *Canadian Medical Association Journal*, *189*(10), E382–E383. https://doi.org/10.1503/cmaj.161033

Rozentsvit, A., Vinokur, K., Samuel, S., Li, Y., Gerdes, A. M., & Carrillo-Sepulveda, M. A. (2017). Ellagic Acid Reduces High Glucose-Induced Vascular Oxidative Stress Through ERK1/2/NOX4 Signaling Pathway. *Cellular Physiology and Biochemistry: International Journal of Experimental Cellular Physiology, Biochemistry, and Pharmacology*, *44*(3), 1174–1187. https://doi.org/10.1159/000485448

Scaldaferri, F., Vetrano, S., Sans, M., Arena, V., Straface, G., Stigliano, E., Repici, A., Sturm, A., Malesci, A., Panes, J., Yla–Herttuala, S., Fiocchi, C., & Danese, S. (2009). VEGF-A Links Angiogenesis and Inflammation in Inflammatory Bowel Disease Pathogenesis. *Gastroenterology*, *136*(2), 585-595.e5. https://doi.org/10.1053/j.gastro.2008.09.064

Scott, A. (2004). What is "inflammation"? Are we ready to move beyond Celsus? *British Journal of Sports Medicine*, *38*(3), 248–249. https://doi.org/10.1136/bjsm.2003.011221

Shaik-Dasthagirisaheb, Y. B., Varvara, G., Murmura, G., Saggini, A., Potalivo, G., Caraffa, A., Antinolfi, P., Tetè, S., Tripodi, D., Conti, F., Cianchetti, E., Toniato, E., Rosati, M., Conti, P., Speranza, L., Pantalone, A., Saggini, R., Theoharides, T. C., & Pandolfi, F. (2013). Vascular Endothelial Growth Factor (VEGF), Mast Cells and Inflammation. *International Journal of Immunopathology and Pharmacology*, *26*(2), 327–335. https://doi.org/10.1177/039463201302600206

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., Amin, N., Schwikowski, B., & Ideker, T. (2003). Cytoscape: A software environment for integrated

models of biomolecular interaction networks. *Genome Research*, *13*(11), 2498–2504.

https://doi.org/10.1101/gr.1239303

Sharifi-Rad, J., Quispe, C., Castillo, C. M. S., Caroca, R., Lazo-Vélez, M. A., Antonyak, H.,

Polishchuk, A., Lysiuk, R., Oliinyk, P., De Masi, L., Bontempo, P., Martorell, M., Daştan,

S. D., Rigano, D., Wink, M., & Cho, W. C. (2022). Ellagic Acid: A Review on Its Natural

Sources, Chemical Stability, and Therapeutic Potential. *Oxidative Medicine and Cellular*

Longevity, *2022*, 1–24. https://doi.org/10.1155/2022/3848084

Sheinenzon, A., Shehadeh, M., Michelis, R., Shaoul, E., & Ronen, O. (2021). Serum albumin

levels and inflammation. *International Journal of Biological Macromolecules*, *184*, 857–

862. https://doi.org/10.1016/j.ijbiomac.2021.06.140

Shin, M., Kang, E., & Lee, Y. (2005). A flavonoid from medicinal plants blocks hepatitis B virus-e

antigen secretion in HBV-infected hepatocytes. *Antiviral Research*, *67*(3), 163–168.

https://doi.org/10.1016/j.antiviral.2005.06.005

Siddappa, D., Beaulieu, É., Gévry, N., Roux, P. P., Bordignon, V., & Duggavathi, R. (2015).

Effect of the transient pharmacological inhibition of Mapk3/1 pathway on ovulation in

mice. *PloS One*, *10*(3), e0119387. https://doi.org/10.1371/journal.pone.0119387

Simon, L. S. (1999). Role and regulation of cyclooxygenase-2 during inflammation. *The*

American Journal of Medicine, *106*(5), 37S-42S. https://doi.org/10.1016/S0002-

9343(99)00115-1

Singh, N., Baby, D., Rajguru, J. P., Patil, P. B., Thakkannavar, S. S., & Pujari, V. B. (2019).

Inflammation and cancer. *Annals of African Medicine*, *18*(3), 121–126.

https://doi.org/10.4103/aam.aam_56_18

Soeters, P. B., Wolfe, R. R., & Shenkin, A. (2019). Hypoalbuminemia: Pathogenesis and Clinical

Significance. *JPEN. Journal of Parenteral and Enteral Nutrition*, *43*(2), 181–193.

https://doi.org/10.1002/jpen.1451

Song, Z., Wang, J., Su, Q., Luan, M., Chen, X., & Xu, X. (2021). The role of MMP-2 and MMP-9 in the metastasis and development of hypopharyngeal carcinoma. *Brazilian Journal of Otorhinolaryngology*, *87*(5), 521–528. https://doi.org/10.1016/j.bjorl.2019.10.009

Stegeman, H., Span, P. N., Peeters, W. J., Verheijen, M. M., Grénman, R., Meijer, T. W., Kaanders, J. H., & Bussink, J. (2016). Interaction between hypoxia, AKT and HIF-1 signaling in HNSCC and NSCLC: Implications for future treatment strategies. *Future Science OA*, *2*(1), FSO84. https://doi.org/10.4155/fso.15.84

Szklarczyk, D., Gable, A. L., Nastou, K. C., Lyon, D., Kirsch, R., Pyysalo, S., Doncheva, N. T., Legeay, M., Fang, T., Bork, P., Jensen, L. J., & von Mering, C. (2021). The STRING database in 2021: Customizable protein–protein networks, and functional characterization of user-uploaded gene/measurement sets. *Nucleic Acids Research*, *49*(D1), D605–D612. https://doi.org/10.1093/nar/gkaa1074

Terzić, J., Grivennikov, S., Karin, E., & Karin, M. (2010). Inflammation and Colon Cancer. *Gastroenterology*, *138*(6), 2101-2114.e5. https://doi.org/10.1053/j.gastro.2010.01.058

Türk, G., Sönmez, M., Ceribaşi, A. O., Yüce, A., & Ateşşahin, A. (2010). Attenuation of cyclosporine A-induced testicular and spermatozoal damages associated with oxidative stress by ellagic acid. *International Immunopharmacology*, *10*(2), 177–182. https://doi.org/10.1016/j.intimp.2009.10.013

Umesalma, S., & Sudhandiran, G. (2010). Differential Inhibitory Effects of the Polyphenol Ellagic Acid on Inflammatory Mediators NF-κB, iNOS, COX-2, TNF-α, and IL-6 in 1,2-Dimethylhydrazine-Induced Rat Colon Carcinogenesis: ELLAGIC ACID REDUCES INFLAMMATION IN RAT COLON CARCINOGENESIS. *Basic & Clinical Pharmacology & Toxicology*, *107*(2), 650–655. https://doi.org/10.1111/j.1742-7843.2010.00565.x

Vonkeman, H. E., & van de Laar, M. A. F. J. (2010). Nonsteroidal Anti-Inflammatory Drugs: Adverse Effects and Their Prevention. *Seminars in Arthritis and Rheumatism*, *39*(4), 294–312. https://doi.org/10.1016/j.semarthrit.2008.08.001

Wu, Y., Zhang, F., Yang, K., Fang, S., Bu, D., Li, H., Sun, L., Hu, H., Gao, K., Wang, W., Zhou,

X., Zhao, Y., & Chen, J. (2019). SymMap: An integrative database of traditional Chinese

medicine enhanced by symptom mapping. *Nucleic Acids Research*, *47*(D1), D1110–

D1117. https://doi.org/10.1093/nar/gky1021

Wu, Y., Zhu, Y., Xie, N., Wang, H., Wang, F., Zhou, J., & Qu, F. (2022). A network

pharmacology approach to explore active compounds and pharmacological mechanisms

of a patented Chinese herbal medicine in the treatment of endometriosis. *PloS One*,

17(2), e0263614. https://doi.org/10.1371/journal.pone.0263614

Xu, Q., Li, S., Tang, W., Yan, J., Wei, X., Zhou, M., & Diao, H. (2021). The Effect of Ellagic Acid

on Hepatic Lipid Metabolism and Antioxidant Activity in Mice. *Frontiers in Physiology*, *12*,

751501. https://doi.org/10.3389/fphys.2021.751501

Yabluchanskiy, A., Ma, Y., Iyer, R. P., Hall, M. E., & Lindsey, M. L. (2013). Matrix

metalloproteinase-9: Many shades of function in cardiovascular disease. *Physiology*

(Bethesda, Md.), *28*(6), 391–403. https://doi.org/10.1152/physiol.00029.2013

Yasir, M., Goyal, A., & Sonthalia, S. (2022). Corticosteroid Adverse Effects. In *StatPearls*.

StatPearls Publishing. http://www.ncbi.nlm.nih.gov/books/NBK531462/

Yu, Y.-M., Wang, Z.-H., Liu, C.-H., & Chen, C.-S. (2007). Ellagic acid inhibits IL-1β-induced cell

adhesion molecule expression in human umbilical vein endothelial cells. *British Journal*

of Nutrition, *97*(4), 692–698. https://doi.org/10.1017/S0007114507666409

Zhang, H.-M., Zhao, L., Li, H., Xu, H., Chen, W.-W., & Tao, L. (2014). Research progress on the

anticarcinogenic actions and mechanisms of ellagic acid. *Cancer Biology & Medicine*,

11(2), 92–100. https://doi.org/10.7497/j.issn.2095-3941.2014.02.004

Zhang, Z., Bao, Y.-Y., & Zhou, S.-H. (2020). Pump Proton and Laryngeal H+/K+ ATPases.

International Journal of General Medicine, *13*, 1509–1514.

https://doi.org/10.2147/IJGM.S284952

Zhao, H., Wu, L., Yan, G., Chen, Y., Zhou, M., Wu, Y., & Li, Y. (2021). Inflammation and tumor

progression: Signaling pathways and targeted intervention. *Signal Transduction and*

Targeted Therapy, *6*(1), 263. https://doi.org/10.1038/s41392-021-00658-5

Zhou, W., Fu, X.-Q., Zhang, L.-L., Zhang, J., Huang, X., Lu, X.-H., Shen, L., Liu, B.-N., Liu, J.,

Luo, H.-S., Yu, J.-P., & Yu, H.-G. (2013). The AKT1/NF-kappaB/Notch1/PTEN axis has

an important role in chemoresistance of gastric cancer cells. *Cell Death & Disease*,

4(10), e847. https://doi.org/10.1038/cddis.2013.375

A.

GO-term	Description	Count in network	False discovery rate
Biological Processes			
GO:0010033	Response to organic substance	46 of 3011	5.36e-27
GO:1901700	Response to oxygen-containing compound	38 of 1567	1.18e-26
GO:0033993	Response to lipid	29 of 858	3.61e-22
GO:0042221	Response to chemical	47 of 4333	5.03e-22
GO:0070887	Cellular response to chemical	41 of 2919	4.58e-21
Molecular Function			
GO:0042802	Identical protein binding	29 of 1896	5.70e-13
GO:0005515	Protein binding	44 of 7026	1.11e-09
GO:0005102	Signaling receptor binding	22 of 1581	2.71e-08
GO:0019899	Enzyme binding	24 of 2239	3.71e-07
GO:0043167	Ion binding	38 of 6188	7.10e-07
Cellular Component			
GO:0031983	Vesicle lumen	11 of 330	1.29e-06
GO:0012505	Endomembrane system	33 of 4542	1.30e-06
GO:0030141	Secretory granule	15 of 845	2.09e-06
GO:0034774	Secretory granule lumen	10 of 324	4.25e-06
GO:0045121	Membrane raft	10 of 324	4.26e-06

A.

B.

Hydrogen bonds between ligands and proteins

RMSD Ligand Movement

Radius of Gyration

Root-mean-square fluctuation

Hydrogen bonds between ligands and proteins

Radius of Gyration

Radius of Gyration

Root-mean-square fluctuation

Hydrogen bonds between ligands and proteins

- PTGS2-Ibuprofen
- PTGS2-EA
- PTGS2-Apo

RMSD Ligand Movement

Radius of Gyration

Time [ns]

Root-mean-square fluctuation

AKT1

MAP3K

PTGS2

VEGFA