**Differences in expression of tumor suppressor, innate immune, inflammasome, and**

**potassium/gap junction channel host genes significantly predict viral reservoir size**

**during treated HIV infection**

**Ashok K. Dwivedi[1], David A. Siegel[1], Cassandra Thanh[2], Rebecca Hoh[1], Kristen S. Hobbs[2],**

**Tony Pan[2], Erica A. Gibson[2], Jeffrey Martin[3], Frederick Hecht[1], Christopher Pilcher[1], Jeffrey**

**Milush[2], Michael P. Busch[4], Mars Stone[4], Meei-Li Huang[5,6], Claire N. Levy[6,7], Pavitra**

**Roychoudhury[5,6], Florian Hladik[6,7], Keith R. Jerome[5,6], Timothy J. Henrich[2], Steven G.**

**Deeks[1]\* and Sulggi A. Lee[1]\*#**

[1] Department of Medicine, Division of HIV, Infectious Diseases & Global Medicine, University of

California San Francisco, 995 Potrero Avenue, San Francisco, CA 94110, USA. [2] Department of

Medicine, Division of Experimental Medicine, University of California San Francisco, CA 94110,

USA. [3] Department of Biostatistics & Epidemiology, University of California San Francisco, CA

94158, USA. [4] Vitalant Blood Bank, San Francisco, CA 94118 USA. [5] Department of Laboratory

Medicine and Pathology, University of Washington, Seattle WA 98195, USA. [6] Vaccine and

Infectious Disease Division, Fred Hutchinson Cancer Center, Seattle WA 98109, USA. [7]

Department of Obstetrics and Gynecology, University of Washington, WA, 98105 USA.

\*Co-correspondence:

sulggi.lee@ucsf.edu, steven.deeks@ucsf.edu

#Lead contact

Title Character Count (max 200): 193

Manuscript Word Count (max 9000): 5414

**Abstract**

The major barrier to an HIV cure is the persistence of infected cells that evade host immune surveillance despite effective antiretroviral therapy (ART). Most prior host genetic HIV studies have focused on identifying DNA polymorphisms (e.g., *CCR5Δ32*, MHC class I alleles) associated with viral load among untreated "elite controllers" (~1% of HIV+ individuals who are able to control virus without ART). However, there have been few studies evaluating host genetic predictors of viral control for the majority of people living with HIV (PLWH) on ART. We performed host RNA sequencing and HIV reservoir quantification (total DNA, unspliced RNA, intact DNA) from peripheral CD4+ T cells from 191 HIV+ ART-suppressed non-controllers. Multivariate models included covariates for timing of ART initiation, nadir CD4+ count, age, sex, and ancestry. Lower HIV total DNA (an estimate of the total reservoir) was associated with upregulation of tumor suppressor genes *NBL1* (q=0.012) and *P3H3* (q=0.012). Higher HIV unspliced RNA (an estimate of residual HIV transcription) was associated with downregulation of several host genes involving inflammasome (*IL1A, CSF3, TNFAIP5, TNFAIP6, TNFAIP9*, *CXCL3, CXCL10*) and innate immune (*TLR7*) signaling, as well as novel associations with potassium (*KCNJ2*) and gap junction (*GJB2*) channels, all q<0.05. Gene set enrichment analyses identified significant associations with TLR4/microbial translocation (q=0.006), IL-1β/NRLP3 inflammasome (q=0.008), and IL-10 (q=0.037) signaling. HIV intact DNA (an estimate of the "replication-competent" reservoir) demonstrated trends with thrombin degradation (*PLGLB1*) and glucose metabolism (*AGL*) genes, but data were (HIV intact DNA detected in only 42% of participants). Our findings demonstrate that among treated PLWH, that inflammation, innate immune responses, bacterial translocation, and tumor suppression/cell proliferation host signaling play a key role in the maintenance of the HIV reservoir during ART. Further data are needed to validate these findings, including functional genomic studies, and expanded epidemiologic studies in female, non-European cohorts.

Abstract Word Count (max 300): 299

**Short Title (max 70 characters: 43):** Host gene expression and HIV reservoir size

**Keywords:** HIV reservoir, host genetics, RNA sequencing, epidemiology

**Author Summary (max word count 150-200):** 154

Although lifelong HIV antiretroviral therapy (ART) suppresses virus, the major barrier to an HIV

cure is the persistence of infected cells that evade host immune surveillance despite effective

ART, "the HIV reservoir." HIV eradication strategies have focused on eliminating residual virus to

allow for HIV remission, but HIV cure trials to date have thus far failed to show a clinically

meaningful reduction in the HIV reservoir. There is an urgent need for a better understanding of

the host-viral dynamics during ART suppression to identify potential novel therapeutic targets for

HIV cure. This is the first epidemiologic host gene expression study to demonstrate a significant

link between HIV reservoir size and several well-known immunologic pathways (e.g., IL-1$\beta$, TLR7,

TNF-$\alpha$ signaling pathways), as well as novel associations with potassium and gap junction

channels (Kir2.1, connexin 26). Further data are needed to validate these findings, including

functional genomic studies and expanded epidemiologic studies in female, non-European

cohorts.

**CONFLICTS:** The authors do not have a commercial or other association that might pose a

conflict of interest.

**FUNDING:** This work was supported in part by the National Institutes of Health: K23GM112526

(SAL), the DARE Collaboratory (U19 AI096109; SGD), the Division of Intramural Research of the

National Institutes (MC), UM1 AI126623 (KRJ), and NIH/NIAID R01A141003 (TJH). This work

was also supported by the amfAR Research Consortium on HIV Eradication a.k.a. ARCHE

(108072-50-RGRL; SGD) and a Collaboration for AIDS Vaccine Discovery (CAVD) grant from the

Bill & Melinda Gates Foundation (INV-008500), the Reservoir Assay Validation and Evaluation

Network (RAVEN) Study Group. The funders had no role in the study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

**Reprints:** Reprint requests can be directed to Dr. Sulggi Lee, the corresponding author (contact

information above).

**Acknowledgements:**

The authors wish to acknowledge the participation of all the study participants who contributed to

this work as well as the clinical research staff of the SCOPE and Options who made this research

possible. All funders had no role in study design, data collection and analysis, decision to publish,

or preparation of the manuscript. All authors provided critical feedback in finalizing the report. SAL

and DAS conceived and designed the study with critical feedback from SGD, TJH, and DAS.

SGD, JM, FH, CP, RH, and SAL coordinated the collection, management, and quality control

processes for the cohort clinical data and SGD, JM, FH, CP, MPB, MS provided biospecimens.

SAL, CT, JM, KB, TP, EAG performed participant sample processing, SAL performed the RNA

sequencing assays, and AKD performed quality control analyses and the association analyses

for the study under the guidance of SAL and DAS. SAL, CT, and KH performed the qPCR HIV

reservoir assays (total DNA, unspliced RNA) in the lab of TH.  CNL and MLH performed the

ddPCR HIV reservoir assay (intact DNA) in the labs of FH, KRJ, and HPK.  AKD, PR, DAS, TJH,

and SAL analyzed these HIV reservoir data in relation to host transcriptomic and clinical

phenotype data. AKD, SAL, and DAS wrote the report. All authors provided critical feedback in

finalizing the manuscript.

## Introduction

Despite several unique cases of possible HIV remission [1-3], there is still no HIV vaccine or cure. The major barrier to a cure is the persistence of infected cells that evade host immune surveillance despite effective antiretroviral therapy (ART). Modern antiretroviral therapy (ART) has transformed HIV disease into a treatable chronic disease for individuals who have access to, and are able to maintain, viral suppression [4]. However, ART alone does not eliminate persistent virus in most individuals [5, 6]. HIV cure trials aimed at reactivating and eliminating the HIV reservoir have thus far failed to show a clinically meaningful reduction in the HIV reservoir [7-12]. There is an urgent need to bridge drug discovery with a deeper understanding of host-viral dynamics. Although several host factors have been shown to influence the size of the "HIV reservoir", such as the timing of ART initiation after initial HIV infection [13-16], maximum pre-ART viral load [17], ethnicity [17], and sex [17-20], there are few published human genomic and transcriptomic epidemiologic studies describing potential host factors influencing HIV persistence during treated infection.

Prior host genome wide association studies (GWAS) have focused on predictors of viral control (during untreated HIV disease), identifying key mutations in the C-C chemokine receptor type 5 gene (*CCR5Δ32*) and the human Major Histocompatibility Complex (MHC) human leukocyte antigen (HLA)-B and -C regions, that influence viral setpoint [21-24]. Recently our group reported these mutations (*CCR5Δ32 and HLA -B\*57:01)* are associated with smaller HIV reservoir size [25]. However, mRNA expression of DNA variation is complex; the basal and/or the conditional expression of these genes in multicellular organisms are influenced by external controls (alternative splicing, polyadenylation, regulatory enhancers, etc.) which may differ by cell type and tissue [26-28]. The limited number of host gene expression studies during HIV infection (e.g., RNA sequencing) have compared gene expression between distinct clinical HIV groups. For example, one prior study compared gene expression among HIV "controllers" (individuals able to control virus in the absence of therapy) versus "non-controllers" [29]. Another study

compared HIV non-controllers initiating ART "early" (<6 months from HIV infection) versus "later"

(≥6 months after infection) [30]. However, no epidemiologic study has examined quantitative

measures of the HIV reservoir size in relation to differences in host gene expression.

Here, we performed a cross-sectional study of 191 ART-suppressed HIV+ non-controllers

to identify differentially expressed host genes in relation to three measures of the peripheral CD4+

T cell reservoir: HIV cell-associated "intact" DNA (an estimate of the frequency of potentially

"replication-competent" virus with intact HIV genomes) [31], as well as total DNA (approximates

intact + defective HIV DNA) and unspliced (full length transcript prior to alternative splicing) RNA.

Increased expression of two putative tumor suppressor genes, *NBL1* and *P3H3*, was associated

with smaller total HIV reservoir size (tDNA). Higher HIV usRNA was associated with

downregulation of 17 host genes, including genes involved in pathogen pattern recognition

(*TLR7*), inflammasome cytokine activation (*IL1A, CSF3, TNFAIP65, TNFAIP6, TNFAIP9*), and

chemokine production (*CXCL3, CXCL10*). Higher usRNA also demonstrated a novel association

with *KCNJ2*, a gene encoding for an inwardly rectifying potassium (Kir2.1) channel which has

been shown to enhance HIV entry and release into host cells [32], as well as with *GJB2*, which

encodes for a gap junction channel which facilitates cell-cell signaling (e.g., K+, Ca+, ATP) that

has been implicated in cell-cell HIV transfer [33, 34]. These data add to the limited literature on

host genetic predictors of the HIV reservoir and suggest that checks on cell proliferation might

limit the total HIV reservoir size while a more "active" reservoir may stimulate host innate immune

responses and inflammation during treated HIV disease. Further data are needed to validate

these findings, including functional genomic studies using CRISPR-cas9 editing and longitudinal

samples allowing causal inferences, as well as expanded studies in female, non-European

cohorts.

**Results**

**Study population**

HIV+ ART-suppressed non-controllers were sampled from the UCSF SCOPE and Options

cohorts (**Supplemental Fig 1**). The final 191 study participants were mostly male (96%) with a

median age of 47 years and included individuals treated during early (within 6 months) or more

chronic (>6 months after) HIV infection (**Table 1**). At the time of biospecimen collection,

participants were ART-suppressed for a median of 5.1 years with a median nadir CD4+ T cell

count of 352 cells/mm$^3$ and maximum pre-ART HIV RNA of 5.1 log$_{10}$copies/mL. As expected, our

U.S.-based study population was diverse (**Fig 1)**. Thus, all results are shown for the total study

population (adjusted for ancestry using principal components [35]), as well as restricted to the

largest homogenous ancestral subgroup (Europeans), in order to enhance the ability to detect

statistically significant genetic associations.

**Measures of the HIV reservoir size were correlated with each other**

Most of the HIV reservoir consists of cells harboring defective virus – i.e., cells that harbor

HIV that is unable to go on to produce virions [36, 37], and yet the "replication-competent"

reservoir is a major target of HIV eradication strategies [31, 38, 39]. Thus, there is currently no

"gold standard" for measuring the HIV reservoir [40, 41]. Here, we performed three measures of

the HIV reservoir from peripheral CD4+ T cells: total DNA (tDNA), unspliced RNA (usRNA), and

HIV intact DNA. To estimate the frequency of the "replication-competent" reservoir, we performed

a multiplexed droplet digital PCR (ddPCR) assay to quantify the frequency of cells with "intact"

HIV sequences (i.e., likely to generate transcripts leading to virion production) [31, 40, 42]. HIV

"total" (i.e., defective+intact ) DNA and "unspliced" RNA (full-length HIV RNA) were also quantified

using a separate, in-house quantitative polymerase chain reaction (qPCR) TaqMan assay [43].

HIV usRNA was statistically significantly correlated with tDNA (R=0.58, P=4.8x10$^{-19}$) and intact

DNA (R=0.24, P=1.9x10$^{-3}$). However, HIV intact DNA was undetectable in 48% of our measured

samples while total DNA was measurable in 95% of samples, which may have influenced the lack

of association between tDNA and intact DNA in our study population (**Fig 2**).

**Earlier ART initiation and nadir CD4+ T cell count were associated with HIV reservoir size**

Consistent with prior work [17, 37, 41], our study found that clinical factors previously shown to influence the size of the HIV reservoir were significantly associated with HIV reservoir measures quantified in our cohort. Earlier timing of ART initiation (<6 months from infection) was statistically significantly associated with lower levels of HIV intact DNA (R=0.21; p=6.5x10$^{-3}$), tDNA (R=0.26; p=3.3x10$^{-4}$), and usRNA (R=0.29; p=7.0x10$^{-5}$) (**Fig 3**). Nadir CD4+ T cell count was associated with larger total HIV DNA reservoir size (R=-0.28; P=6.9x10$^{-5}$), as well as higher levels of HIV usRNA (R=-0.28; p=9.6x10$^{-5}$) and HIV intact DNA (R=-0.23; p=0.002). We did not observe a statistically significant association between HIV reservoir measures and other clinical factors: duration of ART suppression, age, or pre-ART HIV viral load, and we were unable to evaluate differences by sex/gender given low frequencies of female and transgender participants in our study.

**Increased expression of tumor suppressor genes was associated with total HIV DNA reservoir size while higher HIV usRNA was associated with downregulation of host inflammatory and innate immune genes**

A total of 19,912 genes out of 60,719 were included for downstream differential gene expression analyses. In multivariate models adjusted for age, sex, nadir CD4+ T cell count, timing of ART initiation, ancestry (PCs), and residual variability (probabilistic estimation of expression residuals, PEERs), larger total HIV DNA reservoir size was statistically significantly associated with downregulation of two host tumor suppressor genes while higher HIV usRNA levels were associated decreased expression of 17 host genes involved in inflammation and innate immunity.

We observed that upregulation of tumor suppressor genes, *NBL1* and *P3H3, was* associated with smaller total HIV DNA reservoir size (**Supplemental Table 1**). For each fold-increase in gene expression of *NBL1* or *P3H3*, there was a statistically significant decrease in

HIV total DNA (*NBL1*: -1.8%, q=0.012; *P3H3*: -1.6%, q=0.012). However, we observed the

strongest associations between HIV reservoir size and host gene expression were with HIV

unspliced RNA, largely reflecting the "transcriptionally active" HIV reservoir [44, 45]. A total of 17

host genes were inversely associated with HIV usRNA, including *KCNJ2* (-9.7%, q=0.003) which

encodes for an inwardly rectifying potassium channel that has been shown to regulate HIV-1 entry

and release [32], as well as *GJB2* (-7.1%, q=0.012), which encodes for a gap junction protein that

facilitate cell-cell communication, potentially also cell-cell HIV transfer [46, 47]. In addition to these

novel associations, HIV usRNA was also associated with several host genes involved in

proinflammatory cytokine signaling and inflammasome activation (*IL1A*: -9.6%, q=0.012, *CSF3*: -

7.5%, q=0.013; *TNFAIP6*: -7.6%, q=0.016, *TNFAIP9*: -6.9%, q=0.031, *TNFAIP5*: -5.9%, q=0.043),

innate immune responses (*TLR7*: -7.1%, q=0.016), and chemokine production (*CXCL3*: -7.2%,

q=0.043; *CXCL10*: -9.2%, q=0.049) (**Table 2, Supplement Table 2**). Given the large number of

gene hits for HIV usRNA, we also performed network analyses to better visualize immunologic

pathways identified from the differential gene expression analysis (q<0.25). We applied the

ClueGo network analysis application, which clustered the large number of genes into biologically

relevant, interpretable clusters [48]. These analyses highlighted several key pathways involving

inflammasome activation [49-52] and bacterial translocation [53-55] – e.g., genes involved in

NLRP3 (NOD-, LRR- and pyrin domain-containing protein 3) inflammasome activation, IL-1β, toll-

like receptor 4, lipopolysaccharide (LPS), and IL-17 signaling (**Fig 4**). Using unbiased gene set

enrichment analyses (GSEA), all genes in the transcriptome were rank-ordered by p-value to

identify gene sets enriched for each HIV reservoir measurement. These analyses demonstrated

that HIV total DNA was associated with complement activation and humoral immune response

pathways, but these associations were only observed in the subgroup with the largest sample

size, individuals of European ancestry (**Supplement Table 3**). HIV usRNA was again strongly

associated with gene sets involving proinflammatory signaling and microbial translocation

("Response to Bacterium", q=7.5x10$^{-5}$; "Cellular Response to Lipopolysaccharide", q=0.006), IL-

1 signaling ("Interleukin-1 beta production", q=0.008; "Regulation of Interleukin-1 Production",

q=0.008), and cytokine production ("Tumor Necrosis Factor Production", q=0.006; "Tumor

Necrosis Factor Superfamily Cytokine Production", q=0.006; "Regulation of Tumor Necrosis

Factor Production", q=0.008). In addition, several gene sets related to IL-10 signaling ("regulation

of interleukin-10 production", q=0.037, "Interleukin-10 production", q=0.041) – an anti-

inflammatory pathway associated HIV immune dysregulation and persistence [56-58] – were also

significantly associated with HIV usRNA (q=0.04) (**Fig 5, Supplement Table 4**).

**HIV intact DNA was undetectable in over half of the samples but was significantly**

**associated with gene sets involving neutrophil activation in the European subgroup**

HIV intact DNA was undetectable in over half of our measured samples while total DNA

was measurable in 95% of samples (**Fig 2**). Hence, the statistical power to detect differential gene

expression (DGE) associations was much lower for this assay compared to the other reservoir

measures. By performing GSEA (a method that aggregates several genes into immunologically

relevant "gene sets" to test for an association with HIV reservoir size), we were able to enhance

the ability to detect potential associations with HIV intact DNA. In the differential gene expression

analysis, among the European ancestry subgroup, we observed a positive trend (q<0.25)

between HIV intact DNA and two genes, *PLGLB1* (+6.0%, q=0.23), which encodes for a protein

that inhibits thrombus degradation, and *AGL* (+0.9%, q=0.23), encoding for an enzyme involved

in glycogen degradation (**Supplemental Table 5**). GSEA demonstrated that gene sets involving

neutrophil activation ("Neutrophil Degranulation", q=0.046; "Neutrophil Activation Involved in

Immune Response", q=0.046; "Leukocyte Activation"; q=0.046) were significantly associated with

HIV intact DNA, while gene sets reflecting myeloid-mediated immunity ("Myeloid Leukocyte

Mediated Immunity"; q=0.058; "Myeloid Cell Activation Involved in Immune Response"; q=0.060)

demonstrated a slight trend among European ancestry individuals (**Supplement Table 6**).

**Discussion**

In the largest population-based transcriptomic HIV reservoir study to date, among HIV+ ART-suppressed non-controllers, we identified host genetic predictors (e.g., tumor suppressor genes) might act as "checks" on cell proliferation, potentially limiting the total HIV reservoir size. We also observed several associations with host genes indicating that a more "transcriptionally active" HIV reservoir [44, 45] may promote downregulation of potentially harmful host innate immune and proinflammatory responses. Given our cross-sectional study design, further functional and longitudinal epidemiologic studies are needed to determine potential causal relationships between host gene expression and HIV reservoir size. Nonetheless, these findings suggest that even during suppressive ART, ongoing host-pathogen dynamics maintain a delicate balance between a healthy host immune system and a persistent viral reservoir.

Differential gene expression analyses demonstrated that increased expression of tumor suppressor genes *NBL1* and *P3H3* was associated with a smaller total HIV DNA reservoir size. We observed statistically significant associations between two tumor suppressor genes, *NBL1* and *P3H3*, and HIV total DNA. Given the known function of these genes, these findings may suggest that increased expression of these host genes might impact the total HIV reservoir size, possibly by restricting cellular proliferation (**Supplemental Fig 2**). Alternatively, the association of these two tumor suppressor genes with total HIV reservoir size might suggest that cells harboring provirus integrated in genes promoting cell survival are selected for over time during ART suppression, as shown in HIV integration studies [59, 60]. *NBL1*, also known as neuroblastoma suppressor of tumorigenicity 1, is a transcription factor that belongs to the DAN (differential screening-selected gene aberrant in neuroblastoma) family of proteins [61, 62] and is involved in the negative regulation of cell cycle (G1/S transition) [63-66]. Interestingly, *NBL1* was differentially expressed in an RNA-seq *ex vivo* analysis of CD4+ T cells from rhesus macaques (after HIV-1 Env immunization and antibody co-administration) among groups that were treated with immune checkpoint modulators (CTLA-4, PD-1, and CTLA-4 + PD-1 Ab-treated), suggesting

that NBL1 may be a potential pathway by which the cell cycle might be disrupted to enhance HIV-

1 Env antibody responses [67]. *P3H3* encodes for Prolyl 3-Hydroxylase 3, which functions as a

collagen prolyl hydroxylase (vital for collagen biosynthesis) that affects properties of the

extracellular matrix and alters cellular behavior [68-71]. Prior studies suggest that *P3H3* plays a

role as a tumor suppressor in breast, lymphoid, and other cancers [72-74]. Additional genes

identified from the gene set enrichment analysis suggest that ongoing humoral immunity [75] and

complement activation [76] may further contribute to maintaining the total HIV reservoir size

during ART suppression (**Supplemental Table 3**).

HIV unspliced RNA was strongly associated with differential expression of several host

genes previously associated with HIV disease, including genes involving inflammasome

activation and inflammatory cytokine signaling (*IL1A, CSF3, TNFAIP5, TNFAIP6, TNFAIP9, IL10*),

chemokine signaling (*CXCL3, CXCL10*), and innate immune response pathogen pattern

recognition (*TLR7, TLR4*) (**Table 2, Supplemental Table 2, Figure 5, Supplement Table 5**).

Given the known function of these genes and pathways, these findings might reflect that a more

"transcriptionally active" HIV reservoir leads to host downregulation of potentially harmful

proinflammatory signaling pathways during chronic treated HIV disease (**Supplemental Fig 3a**).

This would be consistent with prior data demonstrating that excessive inflammation and immune

activation predicts increased morbidity and mortality in HIV+ individuals despite effective ART [44,

45, 77-80].

Interleukin (IL)-1 is a potent proinflammatory cytokine that regulates inflammation,

triggering a cascade of inflammatory mediators via the NOD-like receptor family pyrin domain

containing 3 (NRLP3) inflammasome activation pathway [81, 82]. IL-1 is an "upstream" pro-

inflammatory inducer of IL-6 [83], which is the strongest biomarker predicting serious non-AIDS

morbidity (e.g., myocardial infarction, stroke, malignancy) [84-87] and mortality [80, 86-89] among

HIV-infected ART-suppressed individuals. In our analysis, higher HIV usRNA was associated with

decreased expression of *IL6* (-7.4%, q=0.062), *IL1A* (-9.6%, q=0.012), and *CSF3* (-7.5%,

q=0.013) (**Table 2, Supplemental Table 2**); the latter encodes for granulocyte colony stimulating

factor 3, G-CSF, a member of the IL-6 superfamily of cytokines [90] that modulates cytokine

production, differentiation, and induction of Treg cells [91]. Our pathway-based analyses

demonstrated a strong association between HIV usRNA and several genes in the IL-1β/NRLP3

inflammasome pathway (**Fig 4**), demonstrating for the first time, a link between this key

immunologic pathway and the HIV reservoir.

Additional cytokines that were statistically significantly associated with HIV usRNA were

several genes in the tumor necrosis (TNF)-α family: *TNFAIP5*, *TNFAIP6* and *TNFAIP9*. These

genes encode for proteins regulating pro- and anti-inflammatory cellular signal transduction,

differentiation, and apoptosis [92-95]. *TNFIAP5* encodes for a pattern recognition receptor that is

induced in response to TNF-α, but also in response to toll-like receptor engagement and IL-1β

signaling [96, 97], signals that are modulated through the NF-κB pathway [98]. *TNFAIP6* encodes

for another TNF-α protein which functions as an anti-inflammatory protein [99, 100], is induced by

IL-1 (upon LPS-stimulation) [101, 102], and interacts with *TNFAIP5* [103, 104]. *TNFAIP9*, also

known as *STEAP4* (six transmembrane epithelial antigen of prostate 4), has been shown to

negatively regulate NF-κB, STAT-3 signaling, and IL-6 production [105, 106]. Several chemokines

were also inversely associated with HIV usRNA, again suggesting that a more "transcriptionally

active" HIV reservoir might promote downregulation of host proinflammatory responses during

long-term ART suppression. *CXCL3* and *CXCL10*, which encode for critical chemokines involved

in the recruitment of neutrophils [107] and activated Th1 lymphocytes [108] to sites of

inflammation respectively, were associated with a 7.2% and 9.2% decrease in gene expression

per two-fold increase in HIV usRNA. CXCL3 regulates monocyte migration [109, 110], neutrophils

chemoattraction [111-113], and angiogenesis [114], and is induced by proinflammatory IL-17

[115, 116]. *CXCL10* encodes for IP-10 (interferon gamma-induced protein 10) which recruits

activated Th1 lymphocytes to sites of infection [117-119] and in HIV, signals through TLR7/9-

dependent pathways [119], predicts HIV disease progression [120, 121], correlates with acute

HIV seroconversion [122], and promotes HIV latency [123, 124]. Finally, in the gene set

enrichment analysis, we also observed a statistically significant association between IL-10

signaling and HIV usRNA (**Fig 5, Supplement Table 4**). IL-10 is an immunosuppressive cytokine

that plays an essential role in limiting the host immune response to pathogens and regulating the

magnitude and duration of inflammation to prevent damage to the host [125]. IL-10 is broadly

expressed by many immune cells, but cell type-specific signals also exist; IL-10 production is

tightly regulated by changes in the chromatin structure, *IL10* gene transcription, and post-

transcriptional regulatory mechanisms [126]. IL-10 has been associated with HIV immune

dysregulation, e.g., impaired CD4+ T cell activation [58], and more recently, IL-10 has been shown

to play a critical role in the maintenance of viral persistence [56, 57]. Among ART-suppressed

PLWH, higher levels of IL-10 measured in blood and lymph nodes were significantly associated

with HIV reservoir size (HIV integrated DNA) [57]. In SIV infected macaques, plasma IL-10 and

IL-10 gene expression was associated with viral reservoir size (SIV DNA) in blood and lymph

nodes, and *in vivo* neutralization of soluble IL-10 was shown to reduce B cell follicle maintenance

[56].

There was also a statistically significant inverse association between HIV usRNA and

genes associated with the host innate immune response (e.g., *TLR7*), while gene set enrichment

also identified TLR4, associated with microbial translocation, to be significantly associated with

HIV usRNA. These findings support the idea that HIV reservoir may not be entirely "quiescent"

during ART and that ongoing residual viral transcription contributes to harmful persistent host

immune activation even during ART suppression. *TLR7* encodes for a member of toll-like receptor

family of genes which plays critical role in pathogen recognition, activation of the innate immune

response, and functions as a bridge between innate and adaptive immunity [127]. TLR7 is a

pattern recognition receptor that can sense HIV single-stranded RNA (ssRNA) [128, 129]. TLR7

agonist administration has been associated with delayed viral rebound [130] and reduced viral

reservoirs in non-human primate studies [131]. A human clinical trial of the TLR7 agonist GS-

9620 recently demonstrated a delay in viral rebound in HIV controllers after cessation of ART

(NCT05281510) [132]. Interestingly, given that *TLR7* is located on the X chromosome, host *TLR7*

transcriptional activity has been linked to acute viremia in HIV+ women (linked to type I interferon

production) [133] as well as with enhanced innate immune function (i.e., plasmacytoid dendritic

cell IFN-$\alpha$ and TNF-$\alpha$ production) [134]. Validation of our findings in female HIV+ cohorts will be

critical for determining whether the host-viral dynamics described in our predominantly male study

population are more pronounced in women, exhibiting a TLR7 signaling "dose-response" effect

due to differential X inactivation in females [133]. Finally, another host innate pattern recognition

receptor statistically significantly associated with HIV usRNA was *TLR4*, which was demonstrated

in the pathway-based analyses linking *TLR4* to several gene sets involved in LPS-mediated

signaling and IL-17 production (**Fig 4**). Our data add to prior studies linking bacterial gut

translocation, systemic inflammation, immune activation, and HIV persistence [135-140].

The most statistically significant association with HIV usRNA was a novel association with

*KCNJ2,* a gene that encodes for an inwardly rectifying potassium channel, Kir2.1. These

potassium ion channels have been shown in prior lab studies to regulate HIV-1 entry and release

[32]. In our analyses, a two-fold increase in HIV transcription (q=0.003) was associated with a

9.7% decrease in *KCNJ2* expression, as well as an 8.4% decrease in *KCNJ2-AS1* (encodes for

KCNJ2 antisense RNA 1) expression (q=0.012). Potassium channels, including inwardly rectifying

K+ channels, have been shown to regulate the life cycle of various viruses (e.g., Ebola [141], SIV

[142]). Tight regulation of potassium ion concentrations have been shown to play a critical role in

HIV-1 virus production in CD4+ T cells in cell culture models [143]. HIV Nef protein has been

shown to increase K+ concentrations in cells [144], and in turn, changes in K+ concentration have

been shown to regulate the HIV life cycle (e.g., viral entry, replication, and release) [32]. A small

molecule inhibitor against Kir 2.1 has been recently identified [145]. The observed association

between HIV usRNA and *KCNJ2*, as well with its antisense RNA, *KCNJ2-AS1*, might then suggest

a potential novel mechanism – targeting specific types of potassium channels – to reduce the HIV

reservoir size.

We also observed a novel association with *GJB2*, which encodes for gap junction beta 2

protein (also known as *CX26*, encoding for connexin 26). Gap junction proteins act as cell-cell

communication channels to transport signaling molecules (e.g., $K^+$, Ca+, ATP) [33, 34], and

HIV-1 is thought to exploit these communication channels to disseminate infection as well as

associated inflammation even in the absence of viral replication [46, 47]. As with *KCNJ2*, the

observed association with *GJB2*, might suggest potential novel targets for limiting the HIV

reservoir size.

We did not observe statistically significant associations with HIV intact DNA and host

genes in the total study population. However, HIV intact DNA was undetectable in 48% of our

measured samples, while for example, total DNA was measurable in 95% of samples (**Fig 2**).

With so many samples below the limit of detection for intact DNA, the statistical power to detect

differential gene expression is much lower for this assay than for the other HIV reservoir assays

included in our study [146, 147]. Thus, we performed additional analyses restricted to the largest

homogenous ancestral population (European ancestry subgroup) and performed pathway

analyses to aggregate individual genes into immunologically relevant "gene sets" to test for an

association with HIV reservoir size. In this way, we were able to enhance the ability to detect

trends with HIV intact DNA. Higher HIV intact DNA was marginally associated with upregulation

of *AGL* (involved in glycogen metabolism) [148-150] and *PLGLB1* (involved in thrombin clot

degradation) [151-153] in the European ancestry subgroup (**Supplement Table 5**). Glycogen

degradation involves breaking down stored glucose for immediate release and availability, and it

has also been shown to play a key role in regulating the inflammatory immune response [150].

Antibody glycosylation has also been associated with inflammation-associated disease [150,

154], as well as time-to-viral rebound after ART interruption, a clinical definition of HIV reservoir

size [148-150]. Besides its role in thrombolysis, *PLGLB1* has previously been associated with a

replication-competent expanded HIV-1 clone described in a patient with squamous cell carcinoma

(AMBI-1 integration) [155]; here it is associated with HIV intact DNA, which estimates the

replication-competent HIV reservoir. Thus, the trends with *AGL* and *PLGLB1*, if further validated,

might reflect that a larger "replication-competent" HIV reservoir contribute to vascular and

metabolic complications that have been previously reported in HIV+ ART-suppressed individuals

[44, 45, 77-80] (**Supplemental Fig 4**).

The study has several limitations that deserve mention. First, although the HIV reservoir

has been shown to be relatively stable over time [17, 156, 157], our cross-sectional design

provides a "snapshot" of the HIV reservoir after a median of 5.1 years of ART suppression and

makes interpretation of the gene associations challenging. However, based on the known

functions of the top gene hits, we conclude that some of the host genes identified in our analyses

might reflect potential drivers of the HIV reservoir size (**Supplemental Fig 2**), while other host

genes represent the impact of persistent HIV (**Supplemental Fig 3-4**). Indeed, the true *in vivo*

associations might involve more complex feedback pathways between the HIV reservoir and host

responses. Second, as is characteristic of our San Francisco-based HIV+ population, our study

included mostly males of European ancestry. We accounted for this using well-established

methods to adjust for population stratification bias [35, 158], as well as the use of PEERs, which

help account for residual variance that often hampers RNA-seq data [159]. Nonetheless, it is

important that these results be replicated in larger studies, especially those including women and

individuals from different ethnic backgrounds. Third, the majority of the HIV reservoir persists in

lymphoid tissues, not in the periphery [160]. However, recent data suggests that the tissue

compartment largely reflects (and is the likely source of) the peripheral compartment [161-163].

Thus, it will be important to determine whether the results from our study are generalizable to the

tissue HIV reservoir in future studies. Finally, we specifically chose to exclude HIV "elite"

controllers in our study, since most people living with HIV do not fall within the ~1% of the HIV+

population able to suppress virus in the absence of therapy. Instead, the focus of our study was

to determine other (uninvestigated) host gene expression associated with the HIV reservoir

(signals that might be lost amidst a study population enriched for previously reported strong

genetic effects, such as with HLA and/or *CCR5Δ32*).

Overall, our findings describe novel and immunologically relevant host genetic

associations with the HIV+ reservoir. These include potential mechanisms inhibiting cell

proliferation to limit the size of the overall HIV reservoir, as well as compensatory host

downregulation of harmful persistent innate immune activation and inflammation (e.g., toll-like

receptor, IL-1β/NRLP3 inflammasome, microbial translocation, IL-10 signaling etc.). Finally, the

strongest association with HIV transcription was with *KCNJ2*, a potential novel mechanism by

which the host restricts residual HIV propagation via inwardly rectifying potassium channels.

Additional studies are needed to validate these findings using approaches functionally and

epidemiologically like CRISPR-Cas9 editing and expanding these studies to include more diverse

patient populations, including female and non-European ancestry individuals, using longitudinal

samples.

**Materials and Methods**

**Study Participants**

HIV+ ART-suppressed non-controllers from the UCSF SCOPE and Options HIV+ cohorts

were included in the study. Inclusion criteria were laboratory-confirmed HIV-1 infection, availability

of cryopreserved peripheral blood mononuclear cells (PBMCs), and plasma HIV RNA levels below

the limit of assay quantification for at least 24 months at the time of biospecimen collection. We

excluded individuals HIV "elite controllers" to focus on genetic variants that drive HIV persistence

among non-controllers during ART suppression but also analyzed previously reported strong

genetic effects associated with HIV+ elite control [164-166]. The estimated date of detected

infection (EDDI) was calculated for each study participant to determine recency of infection in

relation to ART initiation using the Infection Dating Tool (https://tools.incidence-

estimation.org/idt/) [167]. Additional exclusion criteria were potential factors that might influence

HIV reservoir quantification, including recent hospitalization, infection requiring antibiotics,

vaccination, or exposure to immunomodulatory drugs in the six months prior to sampling timepoint.

The research was approved by the UCSF Committee on Human Research (CHR), and all

participants provided written informed consent.

**HIV Reservoir Quantification**

Cryopreserved PBMCs were enriched for CD4+ T cells (StemCell, Vancouver, Canada),

and DNA and RNA were extracted from CD4+ T cells using the AllPrep Universal Kit (Qiagen,

Hilden, Germany). Cell-associated total HIV DNA and unspliced RNA were quantified by an in-

house quantitative polymerase chain reaction (qPCR) TaqMan assay using HIV-1 long terminal

repeat (LTR)-specific primers as previously described [43]. Participant specimens were assayed

with up to 800 ng of total cellular RNA or DNA in replicate reaction wells and copy number

determined by extrapolation against a 7-point standard curve (1–10,000 copies/second)

performed in triplicate. HIV intact DNA was quantified by targeting five regions on the HIV

genome, including highly conserved regions and positions that are frequently deleted or

hypermutated [31]. Optimized restriction enzyme digestion was used to prepare the genomic DNA

for droplet formation while minimizing the amount of shearing within the viral genome. The

protocol targeted 5 regions in the HIV genome across two droplet digital PCR (ddPCR) assays.

Droplet generation and thermocycling were performed according to manufacturer instructions.

This multiplex ddPCR assay allowed the analysis of potentially replication-competent ("intact")

proviral genomes by quantifying the number of droplets positive for 3 targets per assay. Two

targets in a housekeeping gene (*RPP30*) were used to quantify all cells, and a target in the T cell

receptor D gene (*TRD*) was used to normalize the HIV copy numbers per 1x10$^6$ CD4+ T cells. A

DNA shearing index (DSI) was then calculated, and mathematically corrected for residual DNA

shearing as measured by *RPP30* targets to calculate the estimated number of intact proviral

genomes per million CD4+ T cells after correcting for shearing [42].

**Host RNA sequencing**

A separate aliquot of the extracted RNA from CD4+ T cells was then used to perform host

RNA sequencing. HTStream pre-processing pipeline (s4hts.github.io/htstream/) was used for

removing PCR duplicates, adapters, N characters, PolyA/T sequences, Phix contaminants, and

poor-quality sequences (with quality score <20 with sliding window of 10 base pairs). The quality

of raw reads was assessed using FastQC [168]. All samples had a per base quality score and

sequence quality score >30. RNA-seq reads were then mapped to the human genome (GRCh38)

[169] with a corresponding annotation file from the GENCODE project [170]. Alignment and gene

quantification were performed using the STAR alignment tool and its quantification protocols [171-

173]. Gene expression was converted to counts per million (CPM). To normalize the distribution

of expression values across the experiment, the trimmed mean of M-values (TMM) [174] was

used for sample-specific adjustment. Low-expressed genes (<1 CPM for all samples) were

removed. The mean-variance trend was estimated [175] to assign observational weights based

on predicted variance on log2-counts per million (log-CPM) using the Limma-Voom pipeline [176].

**Differential Gene Expression Analysis**

Multivariate linear models were fit for each of the three measures of the HIV reservoir size

using the Limma-Voom workflow [175, 176], a quantitative weighting method that utilizes variance

modeling to accommodate for residual technical and/or biological heterogeneity [175]. For all

analyses, in order to account for potential population stratification bias (i.e., systematic differences

in results due to ancestry rather than association of genes with disease) we used well-established

methods to account for this by (1) calculating and including the first five principal components

(PCs) as covariates in the multivariate models [35] and (2) performing sensitivity analyses among

the largest subgroup, individuals of European ancestry. Eigenvalues were calculated to generate

genetic principal components (PC) to adjust for ancestry [35]. Multivariate models also included

covariates for sex, age, timing of ART initiation, and nadir CD4+ T cell count (duration of ART

suppression and maximum pre-ART viral load did not significantly improve the fit of the models

and were not included as covariates in the final models), as well as PEERs (probabilistic

estimation of expression residuals) to control for additional systematic sources of bias [159].

Model fit was assessed using a lambda genomic coefficient close to 1 [177]. Statistical

significance was determined using a false discovery rate (FDR) q-value threshold of <0.05.

**Gene Set Enrichment Analyses and Network Analyses**

For each of the three HIV reservoir measures, we also performed gene set enrichment analyses

(GSEA) to more broadly evaluate whether specific immune pathways were linked to each HIV

reservoir measurement. Genes from the entire transcriptome were first rank-ordered by q-values

from the differential gene expression analysis for each HIV reservoir measure, and then the rank-

ordering was used to identify immunologic pathways that were enriched from our dataset, using

the Gene Ontology Biological Processes (GO-BP) database [178]. For the HIV usRNA analyses,

for which there were several statistically significant differentially expressed genes (even after

multiple-testing), we performed network analyses to better cluster and visualize the statistically

significant results. Using ClueGo, a network analysis application [48], only statistically significant

and marginally significant genes (q<0.25) were included to calculate Kappa statistics that allowed

more meaningful visualization of potential biologically relevant pathways (**Fig 4**).

**FIGURE LEGENDS**

**Figure 1.** Principal component analysis (PCA) plot of the population structure. Principal component analysis (PCA) plot of the population structure of the full study cohort (a). Secondary PCA plot of the European ancestry subpopulation only (b) defined by the dashed box in the lower left of panel (a). Genetic PCs were calculated from genetic data from our whole exome analysis [25]. Most of the population was of European ancestry (bottom left of) (a) some continued variability. Some continued variability was observed in European ancestry subgroup (b). Self-identified race/ethnicity shown in the legend. Frequencies for participants were recorded as: White/European American (62%), Black/African American (14%), Hispanic/Latino (11%), Mixed Ethnicity/Multiracial (6%), Asian (4%), Pacific Islander (2%), Native American (<1%), and Middle Eastern (<1%).

**Figure 2.** Correlations between three measures of HIV reservoir size. HIV unspliced RNA (usRNA) was significantly correlated with (a) HIV Total DNA (tDNA) and (b) HIV intact DNA; (c) tDNA and intact DNA were not correlated with one another.

**Figure 3.** Measures of the HIV reservoir from peripheral CD4+ T cells were associated with timing of ART initiation. Panels A-D correspond to HIV tDNA, HIV usRNA, RNA/DNA, and Intact DNA, respectively. Spearman correlation and corresponding p-value are shown in each case. Earlier timing of ART initiation (<6 months from infection) was statistically significantly associated with smaller HIV intact DNA, tDNA, and usRNA.

**Figure 4.** Network analysis of the top differentially expressed genes (see **Table 2** and **Supplemental Table 2**) associated with HIV unspliced RNA demonstrated that the top significant genes mapped to immunologic pathways involving bacterial translocation (e.g., TLR4 signaling, activated by bacterial lipopolysaccharide, LPS) and pro-inflammatory responses (e.g., IL-1β

signaling, NLRP3 inflammasome assembly, Th2 cell cytokine production). A Benjamini-Hochberg

false discovery rate (FDR) of q<0.05 was used to generate nodes (circles) based on kappa scores

≥0.4. The size of the nodes reflects the enrichment significance of the terms, and the different

colors represent distinct functional groups.

**Figure 5.** Network analysis of the top statistically significant gene sets associated with HIV

unspliced RNA (see **Supplement Table 4**). Gene sets related to immunologic pathways involving

bacterial translocation (e.g., response to bacterium, LPS-mediated signaling pathway), and

inflammatory signaling (e.g., IL-1β, IL-6, IL-10, TNF-$\alpha$), were significantly associated with HIV

usRNA. A Benjamini-Hochberg false discovery rate (FDR) of q<0.05 was used to generate nodes

(circles) based on kappa scores ≥0.4. The size of the nodes reflects the enrichment significance

of the terms, and the different colors represent distinct functional groups.

**References:**

1. Turk G, Seiger K, Lian X, Sun W, Parsons EM, Gao C, et al. A Possible Sterilizing Cure of HIV-1 Infection Without Stem Cell Transplantation. Ann Intern Med. 2022;175(1):95-100. Epub 2021/11/17. doi: 10.7326/L21-0297. PubMed PMID: 34781719; PubMed Central PMCID: PMCPMC9215120.

2. Hutter G, Nowak D, Mossner M, Ganepola S, Mussig A, Allers K, et al. Long-term control of HIV by CCR5 Delta32/Delta32 stem-cell transplantation. N Engl J Med. 2009;360(7):692-8. Epub 2009/02/14. doi: 10.1056/NEJMoa0802905. PubMed PMID: 19213682.

3. Gupta RK, Abdul-Jawad S, McCoy LE, Mok HP, Peppa D, Salgado M, et al. HIV-1 remission following CCR5Delta32/Delta32 haematopoietic stem-cell transplantation. Nature. 2019;568(7751):244-8. Epub 2019/03/06. doi: 10.1038/s41586-019-1027-4. PubMed PMID: 30836379; PubMed Central PMCID: PMCPMC7275870.

4. Ndung'u T, McCune JM, Deeks SG. Why and where an HIV cure is needed and how it might be achieved. Nature. 2019;576(7787):397-405. doi: 10.1038/s41586-019-1841-8. PubMed PMID: WOS:000504660500088.

5. Katlama C, Deeks SG, Autran B, Martinez-Picado J, Van Lunzen J, Rouzioux C, et al. Barriers to a cure for HIV: new ways to target and eradicate HIV-1 reservoirs. The Lancet. 2013;381(9883):2109-17.

6. Esté JA, Cihlar T. Current status and challenges of antiretroviral research and therapy. Antiviral research. 2010;85(1):25-33.

7. Gay CL, Kuruc JD, Falcinelli SD, Warren JA, Reifeis SA, Kirchherr JL, et al. Assessing the impact of AGS-004, a dendritic cell-based immunotherapy, and vorinostat on persistent HIV-1 Infection. Sci Rep. 2020;10(1):5134. Epub 2020/03/22. doi: 10.1038/s41598-020-61878-3. PubMed PMID: 32198428; PubMed Central PMCID: PMCPMC7083965.

8. Fidler S, Stohr W, Pace M, Dorrell L, Lever A, Pett S, et al. Antiretroviral therapy alone versus antiretroviral therapy with a kick and kill approach, on measures of the HIV reservoir in participants with recent HIV infection (the RIVER trial): a phase 2, randomised trial. Lancet. 2020;395(10227):888-98. Epub 2020/02/23. doi: 10.1016/S0140-6736(19)32990-3. PubMed PMID: 32085823.

9. Gutierrez C, Serrano-Villar S, Madrid-Elena N, Perez-Elias MJ, Martin ME, Barbas C, et al. Bryostatin-1 for latent virus reactivation in HIV-infected patients on antiretroviral therapy. AIDS.

2016;30(9):1385-92. Epub 2016/02/19. doi: 10.1097/QAD.0000000000001064. PubMed PMID:

26891037.

10.    Vibholm L, Schleimann MH, Hojen JF, Benfield T, Offersen R, Rasmussen K, et al. Short-Course

Toll-Like Receptor 9 Agonist Treatment Impacts Innate Immunity and Plasma Viremia in Individuals With

Human Immunodeficiency Virus Infection. Clin Infect Dis. 2017;64(12):1686-95. Epub 2017/03/23. doi:

10.1093/cid/cix201. PubMed PMID: 28329286; PubMed Central PMCID: PMCPMC5849129.

11.    Riddler SA, Para M, Benson CA, Mills A, Ramgopal M, DeJesus E, et al. Vesatolimod, a Toll-like

Receptor 7 Agonist, Induces Immune Activation in Virally Suppressed Adults Living With Human

Immunodeficiency Virus-1. Clin Infect Dis. 2021;72(11):e815-e24. Epub 2020/10/13. doi:

10.1093/cid/ciaa1534. PubMed PMID: 33043969.

12.    Elliott JH, McMahon JH, Chang CC, Lee SA, Hartogensis W, Bumpus N, et al. Short-term

administration of disulfiram for reversal of latent HIV infection: a phase 2 dose-escalation study. Lancet

HIV. 2015;2(12):e520-9. Epub 2015/11/29. doi: 10.1016/S2352-3018(15)00226-X. PubMed PMID:

26614966; PubMed Central PMCID: PMCPMC5108570.

13.    Chun TW, Justement JS, Moir S, Hallahan CW, Maenza J, Mullins JI, et al. Decay of the HIV

reservoir in patients receiving antiretroviral therapy for extended periods: implications for eradication of

virus. J Infect Dis. 2007;195(12):1762-4. Epub 20070502. doi: 10.1086/518250. PubMed PMID:

17492591.

14.    Hocqueloux L, Avettand-Fènoël V, Jacquot S, Prazuck T, Legac E, Mélard A, et al. Long-term

antiretroviral therapy initiated during primary HIV-1 infection is key to achieving both low HIV reservoirs

and normal T cell counts. Journal of Antimicrobial Chemotherapy. 2013;68(5):1169-78. doi:

10.1093/jac/dks533.

15.    Etemad B, Esmaeilzadeh E, Li JZ. Learning From the Exceptions: HIV Remission in Post-

treatment Controllers. Front Immunol. 2019;10:1749. Epub 20190724. doi: 10.3389/fimmu.2019.01749.

PubMed PMID: 31396237; PubMed Central PMCID: PMCPMC6668499.

16.    Fajnzylber J, Sharaf R, Hutchinson JN, Aga E, Bosch RJ, Hartogensis W, et al. Frequency of post

treatment control varies by antiretroviral therapy restart and viral load criteria. AIDS. 2021;35(13):2225-7.

doi: 10.1097/QAD.0000000000002978. PubMed PMID: 34127579; PubMed Central PMCID:

PMCPMC8490281.

17.    Bachmann N, von Siebenthal C, Vongrad V, Turk T, Neumann K, Beerenwinkel N, et al.

Determinants of HIV-1 reservoir size and long-term dynamics during suppressive ART. Nat Commun.

2019;10(1):3193. Epub 2019/07/22. doi: 10.1038/s41467-019-10884-9. PubMed PMID: 31324762;

PubMed Central PMCID: PMCPMC6642170.

18.    Gianella S, Rawlings SA, Nakazawa M, Chaillon A, Strain M, Layman L, et al. Sex differences in

HIV Persistence and Reservoir Size during Aging. Clin Infect Dis. 2021. Epub 20211006. doi:

10.1093/cid/ciab873. PubMed PMID: 34612493.

19.    Gandhi M, Bacchetti P, Miotti P, Quinn TC, Veronese F, Greenblatt RM. Does Patient Sex Affect

Human Immunodeficiency Virus Levels? Clinical Infectious Diseases. 2002;35(3):313-22. doi:

10.1086/341249.

20.    Deeks SG, Bacchetti P, Chomont N, Karn J, Lewin SR, Solomon A, et al. Sex-Based Differences

in Human Immunodeficiency Virus Type 1 Reservoir Activity and Residual Immune Activation. The

Journal of Infectious Diseases. 2019;219(7):1084-94. doi: 10.1093/infdis/jiy617.

21.    de Roda Husman AM, Koot M, Cornelissen M, Keet IP, Brouwer M, Broersen SM, et al.

Association between CCR5 genotype and the clinical course of HIV-1 infection. Ann Intern Med.

1997;127(10):882-90. Epub 1998/02/12. doi: 10.7326/0003-4819-127-10-199711150-00004. PubMed

PMID: 9382366.

22.    Dean M, Carrington M, Winkler C, Huttley GA, Smith MW, Allikmets R, et al. Genetic restriction of

HIV-1 infection and progression to AIDS by a deletion allele of the CKR5 structural gene. Hemophilia

Growth and Development Study, Multicenter AIDS Cohort Study, Multicenter Hemophilia Cohort Study,

San Francisco City Cohort, ALIVE Study. Science. 1996;273(5283):1856-62. Epub 1996/09/27. doi:

10.1126/science.273.5283.1856. PubMed PMID: 8791590.

23.    Rappaport J, Cho YY, Hendel H, Schwartz EJ, Schachter F, Zagury JF. 32 bp CCR-5 gene

deletion and resistance to fast progression in HIV-1 infected heterozygotes. Lancet. 1997;349(9056):922-

3. Epub 1997/03/29. doi: 10.1016/S0140-6736(05)62697-9. PubMed PMID: 9093257.

24.    Kulkarni S, Lied A, Kulkarni V, Rucevic M, Martin MP, Walker-Sperling V, et al. CCR5AS lncRNA

variation differentially regulates CCR5, influencing HIV disease outcome. Nat Immunol. 2019;20(7):824-

34. Epub 2019/06/19. doi: 10.1038/s41590-019-0406-1. PubMed PMID: 31209403; PubMed Central

PMCID: PMCPMC6584055.

25.    Siegel DA, Thanh C, Wan E, Hoh R, Hobbs K, Pan T, et al. Host Variation in Interferon, MHC

Class I, Glycosylation, and Viral Transcription Genes Predict HIV Persistence. 2022. doi:

10.1101/2021.10.31.466670.

26.    Francesconi M, Lehner B. The effects of genetic variation on gene expression dynamics during

development. Nature. 2014;505(7482):208-11. Epub 2013/11/26. doi: 10.1038/nature12772. PubMed

PMID: 24270809.

27.    Williams RB, Chan EK, Cowley MJ, Little PF. The influence of genetic variation on gene

expression. Genome Res. 2007;17(12):1707-16. Epub 2007/12/08. doi: 10.1101/gr.6981507. PubMed

PMID: 18063559.

28.    Yazar S, Alquicira-Hernandez J, Wing K, Senabouth A, Gordon MG, Andersen S, et al. Single-

cell eQTL mapping identifies cell type-specific genetic control of autoimmune disease. Science.

2022;376(6589):eabf3041. Epub 2022/04/08. doi: 10.1126/science.abf3041. PubMed PMID: 35389779.

29.    Shahbaz S, Jovel J, Elahi S. Differential transcriptional and functional properties of regulatory T

cells in HIV-infected individuals on antiretroviral therapy and long-term non-progressors. Clin Transl

Immunology. 2021;10(5):e1289. Epub 20210526. doi: 10.1002/cti2.1289. PubMed PMID: 34094548;

PubMed Central PMCID: PMCPMC8155695.

30.    Petkov S, Chiodi F. Distinct transcriptomic profiles of naive CD4+T cells distinguish HIV-1

infected patients initiating antiretroviral therapy at acute or chronic phase of infection. Genomics.

2021;113(6):3487-500. doi: 10.1016/j.ygeno.2021.08.014. PubMed PMID: WOS:000703998100004.

31.    Levy CN, Hughes SM, Roychoudhury P, Reeves DB, Amstuz C, Zhu H, et al. A highly

multiplexed droplet digital PCR assay to measure the intact HIV-1 proviral reservoir. Cell Rep Med.

2021;2(4):100243. Epub 2021/05/06. doi: 10.1016/j.xcrm.2021.100243. PubMed PMID: 33948574;

PubMed Central PMCID: PMCPMC8080125.

32.     Dubey RC, Mishra N, Gaur R. G protein-coupled and ATP-sensitive inwardly rectifying potassium

ion channels are essential for HIV entry. Scientific reports. 2019;9(1):1-9.

33.     Sáez JC, Nicholson B. Connexin and Pannexin Based Channels in the Nervous System. From

Molecules to Networks2014. p. 257-83.

34.     Goodenough DA, Paul DL. Gap Junctions. Cold Spring Harbor Perspectives in Biology.

2009;1(1):a002576-a. doi: 10.1101/cshperspect.a002576.

35.     Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components

analysis corrects for stratification in genome-wide association studies. Nature genetics. 2006;38(8):904-9.

Epub 2006/07/25. doi: 10.1038/ng1847. PubMed PMID: 16862161.

36.     Ho YC, Shan L, Hosmane NN, Wang J, Laskey SB, Rosenbloom DI, et al. Replication-competent

noninduced proviruses in the latent reservoir increase barrier to HIV-1 cure. Cell. 2013;155(3):540-51.

Epub 2013/11/19. doi: 10.1016/j.cell.2013.09.020. PubMed PMID: 24243014; PubMed Central PMCID:

PMCPMC3896327.

37.     Bruner KM, Murray AJ, Pollack RA, Soliman MG, Laskey SB, Capoferri AA, et al. Defective

proviruses rapidly accumulate during acute HIV-1 infection. Nat Med. 2016;22(9):1043-9. Epub

2016/08/09. doi: 10.1038/nm.4156. PubMed PMID: 27500724; PubMed Central PMCID:

PMCPMC5014606.

38.     Procopio FA, Fromentin R, Kulpa DA, Brehm JH, Bebin AG, Strain MC, et al. A Novel Assay to

Measure the Magnitude of the Inducible Viral Reservoir in HIV-infected Individuals. EBioMedicine.

2015;2(8):872-81. Epub 2015/10/02. doi: 10.1016/j.ebiom.2015.06.019. PubMed PMID: 26425694;

PubMed Central PMCID: PMC4563128.

39.     Eriksson S, Graf EH, Dahl V, Strain MC, Yukl SA, Lysenko ES, et al. Comparative analysis of

measures of viral reservoirs in HIV-1 eradication studies. PLoS Pathog. 2013;9(2):e1003174. doi:

10.1371/journal.ppat.1003174. PubMed PMID: 23459007; PubMed Central PMCID: PMCPMC3573107.

40.     Bruner KM, Wang Z, Simonetti FR, Bender AM, Kwon KJ, Sengupta S, et al. A quantitative

approach for measuring the reservoir of latent HIV-1 proviruses. Nature. 2019;566(7742):120-5. doi:

10.1038/s41586-019-0898-8. PubMed PMID: 30700913.

41.    Eriksson S, Graf EH, Dahl V, Strain MC, Yukl SA, Lysenko ES, et al. Comparative analysis of

measures of viral reservoirs in HIV-1 eradication studies. PLoS pathogens. 2013;9(2):e1003174. Epub

2013/03/06. doi: 10.1371/journal.ppat.1003174. PubMed PMID: 23459007; PubMed Central PMCID:

PMC3573107.

42.    Levy CN, Hughes SM, Roychoudhury P, Amstuz C, Zhu H, Huang ML, et al. HIV reservoir

quantification by five-target multiplex droplet digital PCR. STAR Protoc. 2021;2(4):100885. Epub

20211011. doi: 10.1016/j.xpro.2021.100885. PubMed PMID: 34693363; PubMed Central PMCID:

PMCPMC8517383.

43.    Malnati MS, Scarlatti G, Gatto F, Salvatori F, Cassina G, Rutigliano T, et al. A universal real-time

PCR assay for the quantification of group-M HIV-1 proviral load. Nat Protoc. 2008;3(7):1240-8. Epub

2008/07/05. doi: 10.1038/nprot.2008.108. PubMed PMID: 18600229.

44.    Astorga-Gamaza A, Buzon MJ. The active human immunodeficiency virus reservoir during

antiretroviral therapy: emerging players in viral persistence. Curr Opin HIV AIDS. 2021;16(4):193-9. doi:

10.1097/COH.0000000000000685. PubMed PMID: 33973900.

45.    Pierson T, McArthur J, Siliciano RF. Reservoirs for HIV-1: mechanisms for viral persistence in the

presence of antiviral immune responses and antiretroviral therapy. Annu Rev Immunol. 2000;18:665-708.

doi: 10.1146/annurev.immunol.18.1.665. PubMed PMID: 10837072.

46.    Malik S, Eugenin EA. Role of Connexin and Pannexin containing channels in HIV infection and

NeuroAIDS. Neuroscience Letters. 2019;695:86-90. doi: 10.1016/j.neulet.2017.09.005.

47.    Valdebenito S, Barreto A, Eugenin EA. The role of connexin and pannexin containing channels in

the innate and acquired immune response. Biochimica et Biophysica Acta (BBA) - Biomembranes.

2018;1860(1):154-65. doi: 10.1016/j.bbamem.2017.05.015.

48.    Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, et al. ClueGO: a

Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks.

Bioinformatics. 2009;25(8):1091-3. Epub 20090223. doi: 10.1093/bioinformatics/btp101. PubMed PMID:

19237447; PubMed Central PMCID: PMCPMC2666812.

49.    Mantovani A, Dinarello CA, Molgora M, Garlanda C. Interleukin-1 and Related Cytokines in the

Regulation of Inflammation and Immunity. Immunity. 2019;50(4):778-95. Epub 2019/04/18. doi:

10.1016/j.immuni.2019.03.012. PubMed PMID: 30995499; PubMed Central PMCID: PMCPMC7174020.

50.    Gabay C, Lamacchia C, Palmer G. IL-1 pathways in inflammation and human diseases. Nat Rev

Rheumatol. 2010;6(4):232-41. Epub 2010/02/24. doi: 10.1038/nrrheum.2010.4. PubMed PMID:

20177398.

51.    Dinarello CA. Immunological and inflammatory functions of the interleukin-1 family. Annu Rev

Immunol. 2009;27:519-50. Epub 2009/03/24. doi: 10.1146/annurev.immunol.021908.132612. PubMed

PMID: 19302047.

52.    Zheng D, Liwinski T, Elinav E. Inflammasome activation and regulation: toward a better

understanding of complex mechanisms. Cell Discov. 2020;6:36. Epub 2020/06/19. doi: 10.1038/s41421-

020-0167-x. PubMed PMID: 32550001; PubMed Central PMCID: PMCPMC7280307.

53.    Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, et al. Microbial

translocation is a cause of systemic immune activation in chronic HIV infection. Nat Med.

2006;12(12):1365-71. Epub 2006/11/23. doi: 10.1038/nm1511. PubMed PMID: 17115046.

54.    Vujkovic-Cvijin I, Dunham RM, Iwai S, Maher MC, Albright RG, Broadhurst MJ, et al. Dysbiosis of

the gut microbiota is associated with HIV disease progression and tryptophan catabolism. Sci Transl Med.

2013;5(193):193ra91. Epub 2013/07/12. doi: 10.1126/scitranslmed.3006438. PubMed PMID: 23843452;

PubMed Central PMCID: PMCPMC4094294.

55.    Somsouk M, Estes JD, Deleage C, Dunham RM, Albright R, Inadomi JM, et al. Gut epithelial

barrier and systemic inflammation during chronic HIV infection. AIDS. 2015;29(1):43-51. Epub

2014/11/12. doi: 10.1097/QAD.0000000000000511. PubMed PMID: 25387317; PubMed Central PMCID:

PMCPMC4444362.

56.    Harper J, Ribeiro SP, Chan CN, Aid M, Deleage C, Micci L, et al. Interleukin-10 contributes to

reservoir establishment and persistence in SIV-infected macaques treated with antiretroviral therapy. J

Clin Invest. 2022;132(8). Epub 2022/03/02. doi: 10.1172/JCI155251. PubMed PMID: 35230978; PubMed

Central PMCID: PMCPMC9012284.

57. Ribeiro SP, Aid M, Dupuy FP, Chan CN, Hultquist JF, Delage C, et al. IL-10 driven memory T cell

survival and Tfh differentiation promote HIV

2 persistence. BioRxiv. 2022. doi: doi: https://doi.org/10.1101/2021.02.26.432955.

58. Said EA, Dupuy FP, Trautmann L, Zhang Y, Shi Y, El-Far M, et al. Programmed death-1-induced

interleukin-10 production by monocytes impairs CD4+ T cell activation during HIV infection. Nature

medicine. 2010;16(4):452-9. Epub 2010/03/09. doi: 10.1038/nm.2106. PubMed PMID: 20208540.

59. Chomont N, El-Far M, Ancuta P, Trautmann L, Procopio FA, Yassine-Diab B, et al. HIV reservoir

size and persistence are driven by T cell survival and homeostatic proliferation. Nat Med. 2009;15(8):893-

900. Epub 2009/06/23. doi: 10.1038/nm.1972. PubMed PMID: 19543283; PubMed Central PMCID:

PMCPMC2859814.

60. Wagner TA, McLaughlin S, Garg K, Cheung CY, Larsen BB, Styrchak S, et al. HIV latency.

Proliferation of cells with HIV integrated into cancer genes contributes to persistent infection. Science.

2014;345(6196):570-3. Epub 20140710. doi: 10.1126/science.1256304. PubMed PMID: 25011556;

PubMed Central PMCID: PMCPMC4230336.

61. Kattamuri C, Luedeke DM, Nolan K, Rankin SA, Greis KD, Zorn AM, et al. Members of the DAN

family are BMP antagonists that form highly stable noncovalent dimers. J Mol Biol. 2012;424(5):313-27.

Epub 20121009. doi: 10.1016/j.jmb.2012.10.003. PubMed PMID: 23063586; PubMed Central PMCID:

PMCPMC3509953.

62. Hung WT, Wu FJ, Wang CJ, Luo CW. DAN (NBL1) specifically antagonizes BMP2 and BMP4

and modulates the actions of GDF9, BMP2, and BMP4 in the rat ovary. Biol Reprod. 2012;86(5):158, 1-9.

Epub 20120531. doi: 10.1095/biolreprod.111.096172. PubMed PMID: 22357543.

63. Ozaki T, Nakamura Y, Enomoto H, Hirose M, Sakiyama S. Overexpression of DAN gene product

in normal rat fibroblasts causes a retardation of the entry into the S phase. Cancer Res. 1995;55(4):895-

900. PubMed PMID: 7850806.

64. Sung JC, McCarthy S, Turner J, Li CG, Yeatman TJ. The NBL1 tumor suppressor gene is

downregulated in colon cancer by promoter methylation. Journal of the American College of Surgeons.

2004;199(3). doi: 10.1016/j.jamcollsurg.2004.05.200.

65.    Hayashi T, Sentani K, Oue N, Ohara S, Teishima J, Anami K, et al. The search for secreted

proteins in prostate cancer by the Escherichia coli ampicillin secretion trap: expression of NBL1 is highly

restricted to the prostate and is related to cancer progression. Pathobiology. 2013;80(2):60-9. Epub

20120829. doi: 10.1159/000341396. PubMed PMID: 22948749.

66.    Nakamura Y, Ozaki T, Ichimiya S, Nakagawara A, Sakiyama S. Ectopic expression of DAN

enhances the retinoic acid-induced neuronal differentiation in human neuroblastoma cell lines. Biochem

Biophys Res Commun. 1998;243(3):722-6. doi: 10.1006/bbrc.1998.8112. PubMed PMID: 9500977.

67.    Bradley T, Kuraoka M, Yeh CH, Tian M, Chen H, Cain DW, et al. Immune checkpoint modulation

enhances HIV-1 antibody induction. Nat Commun. 2020;11(1):948. Epub 2020/02/23. doi:

10.1038/s41467-020-14670-w. PubMed PMID: 32075963; PubMed Central PMCID: PMCPMC7031230.

68.    O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference sequence

(RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. Nucleic

acids research. 2016;44(D1):D733-D45.

69.    Rappu P, Salo AM, Myllyharju J, Heino J. Role of prolyl hydroxylation in the molecular

interactions of collagens. Essays in biochemistry. 2019;63(3):325-35.

70.    Shah R, Smith P, Purdie C, Quinlan P, Baker L, Aman P, et al. The prolyl 3-hydroxylases P3H2

and P3H3 are novel targets for epigenetic silencing in breast cancer. British journal of cancer.

2009;100(10):1687-96.

71.    Li Y, Chen Y, Ma Y, Nenkov M, Haase D, Petersen I. Collagen prolyl hydroxylase 3 has a tumor

suppressive activity in human lung cancer. Experimental cell research. 2018;363(1):121-8.

72.    Hatzimichael E, Dasoula A, Shah R, Syed N, Papoudou-Bai A, Coley HM, et al. The prolyl-

hydroxylase EGLN3 and not EGLN1 is inactivated by methylation in plasma cell neoplasia. Eur J

Haematol. 2010;84(1):47-51. Epub 2009/09/10. doi: 10.1111/j.1600-0609.2009.01344.x. PubMed PMID:

19737309.

73.    Hatzimichael E, Lo Nigro C, Lattanzio L, Syed N, Shah R, Dasoula A, et al. The collagen prolyl

hydroxylases are novel transcriptionally silenced genes in lymphoma. Br J Cancer. 2012;107(8):1423-32.

Epub 2012/09/08. doi: 10.1038/bjc.2012.380. PubMed PMID: 22955849; PubMed Central PMCID:

PMCPMC3494450.

74.    Shah R, Smith P, Purdie C, Quinlan P, Baker L, Aman P, et al. The prolyl 3-hydroxylases P3H2

and P3H3 are novel targets for epigenetic silencing in breast cancer. Br J Cancer. 2009;100(10):1687-96.

Epub 2009/05/14. doi: 10.1038/sj.bjc.6605042. PubMed PMID: 19436308; PubMed Central PMCID:

PMCPMC2696763.

75.    Maggi E, Almerigogna F, Del Prete G, Romagnani S. Abnormal B cell helper activity by virus-

infected human CD4+ T cells. Seminars in Immunology. 1993;5(6):449-55. doi: 10.1006/smim.1993.1051.

76.    Liszewski MK, Kolev M, Le Friec G, Leung M, Bertram PG, Fara AF, et al. Intracellular

complement activation sustains T cell homeostasis and mediates effector differentiation. Immunity.

2013;39(6):1143-57. Epub 20131205. doi: 10.1016/j.immuni.2013.10.018. PubMed PMID: 24315997;

PubMed Central PMCID: PMCPMC3865363.

77.    Kaplan RC, Sinclair E, Landay AL, Lurain N, Sharrett AR, Gange SJ, et al. T cell activation

predicts carotid artery stiffness among HIV-infected women. Atherosclerosis. 2011;217(1):207-13. Epub

20110315. doi: 10.1016/j.atherosclerosis.2011.03.011. PubMed PMID: 21492857; PubMed Central

PMCID: PMCPMC3139014.

78.    Lederman MM, Funderburg NT, Sekaly RP, Klatt NR, Hunt PW. Residual immune dysregulation

syndrome in treated HIV infection. Adv Immunol. 2013;119:51-83. doi: 10.1016/B978-0-12-407707-

2.00002-3. PubMed PMID: 23886064; PubMed Central PMCID: PMCPMC4126613.

79.    Hunt PW. HIV and Inflammation: Mechanisms and Consequences. Curr Hiv-Aids Rep.

2012;9(2):139-47. doi: 10.1007/s11904-012-0118-8. PubMed PMID: WOS:000208974200005.

80.    Hunt PW, Sinclair E, Rodriguez B, Shive C, Clagett B, Funderburg N, et al. Gut epithelial barrier

dysfunction and innate immune activation predict mortality in treated HIV infection. The Journal of

Infectious Diseases. 2014;210(8):1228-38. Epub 2014/04/24. doi: 10.1093/infdis/jiu238. PubMed PMID:

24755434; PubMed Central PMCID: PMC4192038.

81.    Agostini L, Martinon F, Burns K, McDermott MF, Hawkins PN, Tschopp J. NALP3 forms an IL-

1beta-processing inflammasome with increased activity in Muckle-Wells autoinflammatory disorder.

Immunity. 2004;20(3):319-25. PubMed PMID: 15030775.

82.     Dinarello CA. Interleukin-1 in the pathogenesis and treatment of inflammatory diseases. Blood.

2011;117(14):3720-32. Epub 2011/02/10. doi: 10.1182/blood-2010-07-273417. PubMed PMID:

21304099; PubMed Central PMCID: PMCPMC3083294.

83.     Ridker PM. From C-Reactive Protein to Interleukin-6 to Interleukin-1: Moving Upstream To

Identify Novel Targets for Atheroprotection. Circ Res. 2016;118(1):145-56. doi:

10.1161/CIRCRESAHA.115.306656. PubMed PMID: 26837745; PubMed Central PMCID:

PMCPMC4793711.

84.     Kuller LH, Tracy R, Belloso W, De Wit S, Drummond F, Lane HC, et al. Inflammatory and

coagulation biomarkers and mortality in patients with HIV infection. PLoS Med. 2008;5(10):e203. PubMed

PMID: 18942885.

85.     Sunil M, Nigalye M, Somasunderam A, Martinez ML, Yu X, Arduino RC, et al. Unchanged Levels

of Soluble CD14 and IL-6 Over Time Predict Serious Non-AIDS Events in HIV-1-Infected People. AIDS

Res Hum Retroviruses. 2016;32(12):1205-9. doi: 10.1089/AID.2016.0007. PubMed PMID: 27344921;

PubMed Central PMCID: PMCPMC5175436.

86.     Tenorio AR, Zheng Y, Bosch RJ, Krishnan S, Rodriguez B, Hunt PW, et al. Soluble markers of

inflammation and coagulation but not T-cell activation predict non-AIDS-defining morbid events during

suppressive antiretroviral treatment. The Journal of Infectious Diseases. 2014;210(8):1248-59. Epub

2014/05/06. doi: 10.1093/infdis/jiu254. PubMed PMID: 24795473; PubMed Central PMCID:

PMC4192039.

87.     Grund B, Baker JV, Deeks SG, Wolfson J, Wentworth D, Cozzi-Lepri A, et al. Relevance of

Interleukin-6 and D-Dimer for Serious Non-AIDS Morbidity and Death among HIV-Positive Adults on

Suppressive Antiretroviral Therapy. PLoS One. 2016;11(5):e0155100. doi:

10.1371/journal.pone.0155100. PubMed PMID: 27171281; PubMed Central PMCID: PMCPMC4865234.

88.     Lee SL, Byakwaga H, Boum Y, Burdo T, Williams KC, Lederman MM, et al. Immunologic

Pathways that Predict Mortality in HIV-Infected Ugandans Initiating ART. The Journal of Infectious

Diseases. 2017;In Review.

89.     So-Armah KA, Tate JP, Chang CH, Butt AA, Gerschenson M, Gibert CL, et al. Do Biomarkers of

Inflammation, Monocyte Activation, and Altered Coagulation Explain Excess Mortality Between HIV

Infected and Uninfected People? J Acquir Immune Defic Syndr. 2016;72(2):206-13. doi:

10.1097/QAI.0000000000000954. PubMed PMID: 27824677.

90.     Unver N, McAllister F. IL-6 family cytokines: Key inflammatory mediators as biomarkers and

potential therapeutic targets. Cytokine & Growth Factor Reviews. 2018;41:10-7. doi:

10.1016/j.cytogfr.2018.04.004.

91.     Rutella S, Zavala F, Danese S, Kared H, Leone G. Granulocyte colony-stimulating factor: a novel

mediator of T cell tolerance. The Journal of Immunology. 2005;175(11):7085-91.

92.     Jacques P, Elewaut D. Tumor necrosis factor α-induced proteins: Natural brakes on

inflammation. Arthritis & Rheumatism. 2012;64(12):3831-4. doi: 10.1002/art.34664.

93.     Sethi JK, Hotamisligil GS. Metabolic Messengers: tumour necrosis factor. Nat Metab.

2021;3(10):1302-12. Epub 20211014. doi: 10.1038/s42255-021-00470-z. PubMed PMID: 34650277.

94.     Heir R, Stellwagen D. TNF-Mediated Homeostatic Synaptic Plasticity: From in vitro to in vivo

Models. Front Cell Neurosci. 2020;14:565841. Epub 20200930. doi: 10.3389/fncel.2020.565841. PubMed

PMID: 33192311; PubMed Central PMCID: PMCPMC7556297.

95.     Gough P, Myles IA. Tumor Necrosis Factor Receptors: Pleiotropic Signaling Complexes and

Their Differential Effects. Front Immunol. 2020;11:585880. Epub 20201125. doi:

10.3389/fimmu.2020.585880. PubMed PMID: 33324405; PubMed Central PMCID: PMCPMC7723893.

96.     Breviario F, d'Aniello EM, Golay J, Peri G, Bottazzi B, Bairoch A, et al. Interleukin-1-inducible

genes in endothelial cells. Cloning of a new gene related to C-reactive protein and serum amyloid P

component. J Biol Chem. 1992;267(31):22190-7. PubMed PMID: 1429570.

97.     Porte R, Davoudian S, Asgari F, Parente R, Mantovani A, Garlanda C, et al. The Long Pentraxin

PTX3 as a Humoral Innate Immunity Functional Player and Biomarker of Infections and Sepsis. Frontiers

in Immunology. 2019;10. doi: 10.3389/fimmu.2019.00794.

98.     Doni A, Stravalaci M, Inforzato A, Magrini E, Mantovani A, Garlanda C, et al. The Long Pentraxin

PTX3 as a Link Between Innate Immunity, Tissue Remodeling, and Cancer. Frontiers in Immunology.

2019;10. doi: 10.3389/fimmu.2019.00712.

99.    Dyer DP, Salanga CL, Johns SC, Valdambrini E, Fuster MM, Milner CM, et al. The Anti-

inflammatory Protein TSG-6 Regulates Chemokine Function by Inhibiting Chemokine/Glycosaminoglycan

Interactions. Journal of Biological Chemistry. 2016;291(24):12627-40. doi: 10.1074/jbc.M116.720953.

100.    Dyer DP, Thomson JM, Hermant A, Jowitt TA, Handel TM, Proudfoot AEI, et al. TSG-6 Inhibits

Neutrophil Migration via Direct Interaction with the Chemokine CXCL8. The Journal of Immunology.

2014;192(5):2177-85. doi: 10.4049/jimmunol.1300194.

101.    Lee TH, Klampfer L, Shows TB, Vilcek J. Transcriptional regulation of TSG6, a tumor necrosis

factor- and interleukin-1-inducible primary response gene coding for a secreted hyaluronan-binding

protein. J Biol Chem. 1993;268(9):6154-60. PubMed PMID: 8454591.

102.    Mittal M, Tiruppathi C, Nepal S, Zhao YY, Grzych D, Soni D, et al. TNFalpha-stimulated gene-6

(TSG6) activates macrophage phenotype transition to prevent inflammatory lung injury. Proc Natl Acad

Sci U S A. 2016;113(50):E8151-E8. Epub 20161128. doi: 10.1073/pnas.1614935113. PubMed PMID:

27911817; PubMed Central PMCID: PMCPMC5167170.

103.    Nauta AJ, Bottazzi B, Mantovani A, Salvatori G, Kishore U, Schwaeble WJ, et al. Biochemical

and functional characterization of the interaction between pentraxin 3 and C1q. European Journal of

Immunology. 2003;33(2):465-73. doi: 10.1002/immu.200310022.

104.    Garlanda C, Bottazzi B, Bastone A, Mantovani A. Pentraxins at the crossroads between innate

immunity, inflammation, matrix deposition, and female fertility. Annu Rev Immunol. 2005;23:337-66. doi:

10.1146/annurev.immunol.23.021704.115756. PubMed PMID: 15771574.

105.    Takai C, Matsumoto I, Inoue A, Umeda N, Tanaka Y, Kurashima Y, et al. Specific overexpression

of tumour necrosis factor-α-induced protein (TNFAIP)9 in CD14+CD16− monocytes in patients with

rheumatoid arthritis: comparative analysis with TNFAIP3. Clinical and Experimental Immunology.

2015;180(3):458-66. doi: 10.1111/cei.12606.

106.    Inoue A, Matsumoto I, Tanaka Y, Umeda N, Tanaka Y, Mihara M, et al. Murine tumor necrosis

factor α-induced adipose-related protein (tumor necrosis factor α-induced protein 9) deficiency leads to

arthritis via interleukin-6 overproduction with enhanced NF-κB, STAT-3 signaling, and dysregulated

apoptosis of macrophages. Arthritis & Rheumatism. 2012;64(12):3877-85. doi: 10.1002/art.34666.

107.    Groves D, Jiang Y. Chemokines, a family of chemotactic cytokines. Critical Reviews in Oral

Biology & Medicine. 1995;6(2):109-18.

108.    Foley JF, Yu C-R, Solow R, Yacobucci M, Peden KWC, Farber JM. Roles for CXC Chemokine

Ligands 10 and 11 in Recruiting CD4+T Cells to HIV-1-Infected Monocyte-Derived Macrophages,

Dendritic Cells, and Lymph Nodes. The Journal of Immunology. 2005;174(8):4892-900. doi:

10.4049/jimmunol.174.8.4892.

109.    Smith DF, Galkina E, Ley K, Huo Y. GRO family chemokines are specialized for monocyte arrest

from flow. Am J Physiol Heart Circ Physiol. 2005;289(5):H1976-84. Epub 20050603. doi:

10.1152/ajpheart.00153.2005. PubMed PMID: 15937099.

110.    Ahuja SK, Murphy PM. The CXC Chemokines Growth-regulated Oncogene (GRO) α, GROβ,

GROγ, Neutrophil-activating Peptide-2, and Epithelial Cell-derived Neutrophil-activating Peptide-78 Are

Potent Agonists for the Type B, but Not the Type A, Human Interleukin-8 Receptor. Journal of Biological

Chemistry. 1996;271(34):20545-50. doi: 10.1074/jbc.271.34.20545.

111.    Ahuja SK, Murphy PM. The CXC chemokines growth-regulated oncogene (GRO) alpha,

GRObeta, GROgamma, neutrophil-activating peptide-2, and epithelial cell-derived neutrophil-activating

peptide-78 are potent agonists for the type B, but not the type A, human interleukin-8 receptor. J Biol

Chem. 1996;271(34):20545-50. doi: 10.1074/jbc.271.34.20545. PubMed PMID: 8702798.

112.    Rudack C, Maune S, Eble J, Schroeder J-M. The Primary Role in Biologic Activity of the

Neutrophil Chemokines IL-8 and GRO-αin Cultured Nasal Epithelial Cells. Journal of Interferon &

Cytokine Research. 2003;23(2):113-23. doi: 10.1089/107999003321455507.

113.    Metzemaekers M, Gouwy M, Proost P. Neutrophil chemoattractant receptors in health and

disease: double-edged swords. Cellular & Molecular Immunology. 2020;17(5):433-50. doi:

10.1038/s41423-020-0412-0.

114.    Zaja-Milatovic S, Richmond A. CXC chemokines and their receptors: a case for a significant

biological role in cutaneous wound healing. Histol Histopathol. 2008;23(11):1399-407. doi: 10.14670/HH-

23.1399. PubMed PMID: 18785122; PubMed Central PMCID: PMCPMC3140405.

115.     Al-Alwan LA, Chang Y, Baglole CJ, Risse P-A, Halayko AJ, Martin JG, et al. Autocrine-regulated

airway smooth muscle cell migration is dependent on IL-17–induced growth-related oncogenes. Journal

of Allergy and Clinical Immunology. 2012;130(4):977-85.e6. doi: 10.1016/j.jaci.2012.04.042.

116.     Chen C-L, Wang Y, Huang C-Y, Zhou Z-Q, Zhao J-J, Zhang X-F, et al. IL-17 induces antitumor

immunity by promoting beneficial neutrophil recruitment and activation in esophageal squamous cell

carcinoma. OncoImmunology. 2017;7(1). doi: 10.1080/2162402x.2017.1373234.

117.     Rotondi M, Chiovato L, Romagnani S, Serio M, Romagnani P. Role of chemokines in endocrine

autoimmune diseases. Endocr Rev. 2007;28(5):492-520. Epub 20070502. doi: 10.1210/er.2006-0044.

PubMed PMID: 17475924.

118.     Smit MJ, Verdijk P, van der Raaij-Helmer EM, Navis M, Hensbergen PJ, Leurs R, et al. CXCR3-

mediated chemotaxis of human T cells is regulated by a Gi- and phospholipase C-dependent pathway

and not via activation of MEK/p44/p42 MAPK nor Akt/PI-3 kinase. Blood. 2003;102(6):1959-65. Epub

20030515. doi: 10.1182/blood-2002-12-3945. PubMed PMID: 12750173.

119.     Simmons RP, Scully EP, Groden EE, Arnold KB, Chang JJ, Lane K, et al. HIV-1 infection induces

strong production of IP-10 through TLR7/9-dependent pathways. AIDS. 2013;27(16):2505-17. Epub

2013/10/08. doi: 10.1097/01.aids.0000432455.06476.bc. PubMed PMID: 24096630; PubMed Central

PMCID: PMCPMC4288813.

120.     Gray CM, Liovat A-S, Rey-Cuillé M-A, Lécuroux C, Jacquelin B, Girault I, et al. Acute Plasma

Biomarkers of T Cell Activation Set-Point Levels and of Disease Progression in HIV-1 Infection. PLoS

ONE. 2012;7(10). doi: 10.1371/journal.pone.0046143.

121.     Valverde-Villegas JM, de Medeiros RM, Ellwanger JH, Santos BR, Melo MG, Almeida SEM, et al.

High CXCL10/IP-10 levels are a hallmark in the clinical evolution of the HIV infection. Infect Genet Evol.

2018;57:51-8. Epub 2017/11/11. doi: 10.1016/j.meegid.2017.11.002. PubMed PMID: 29122683.

122.     Pastor L, Casellas A, Carrillo J, Alonso S, Parker E, Fuente-Soro L, et al. IP-10 Levels as an

Accurate Screening Tool to Detect Acute HIV Infection in Resource-Limited Settings. Sci Rep.

2017;7(1):8104. Epub 2017/08/16. doi: 10.1038/s41598-017-08218-0. PubMed PMID: 28808319;

PubMed Central PMCID: PMCPMC5556096.

123.    Wang Z, Yin X, Ma M, Ge H, Lang B, Sun H, et al. IP-10 Promotes Latent HIV Infection in Resting

Memory CD4(+) T Cells via LIMK-Cofilin Pathway. Front Immunol. 2021;12:656663. Epub 2021/08/28.

doi: 10.3389/fimmu.2021.656663. PubMed PMID: 34447368; PubMed Central PMCID:

PMCPMC8383741.

124.    Cameron PU, Saleh S, Sallmann G, Solomon A, Wightman F, Evans VA, et al. Establishment of

HIV-1 latency in resting CD4+ T cells depends on chemokine-induced changes in the actin cytoskeleton.

Proc Natl Acad Sci U S A. 2010;107(39):16934-9. Epub 2010/09/15. doi: 10.1073/pnas.1002894107.

PubMed PMID: 20837531; PubMed Central PMCID: PMCPMC2947912.

125.    Ouyang W, O'Garra A. IL-10 Family Cytokines IL-10 and IL-22: from Basic Science to Clinical

Translation. Immunity. 2019;50(4):871-91. Epub 2019/04/18. doi: 10.1016/j.immuni.2019.03.020. PubMed

PMID: 30995504.

126.    Saraiva M, O'Garra A. The regulation of IL-10 production by immune cells. Nat Rev Immunol.

2010;10(3):170-81. Epub 2010/02/16. doi: 10.1038/nri2711. PubMed PMID: 20154735.

127.    Werling D, Jungi TW. TOLL-like receptors linking innate and adaptive immune response.

Veterinary Immunology and Immunopathology. 2003;91(1):1-12. doi: 10.1016/s0165-2427(02)00228-3.

128.    Diebold SS, Kaisho T, Hemmi H, Akira S, Reis e Sousa C. Innate antiviral responses by means of

TLR7-mediated recognition of single-stranded RNA. Science. 2004;303(5663):1529-31. Epub 20040219.

doi: 10.1126/science.1093616. PubMed PMID: 14976261.

129.    Heil F, Hemmi H, Hochrein H, Ampenberger F, Kirschning C, Akira S, et al. Species-specific

recognition of single-stranded RNA via toll-like receptor 7 and 8. Science. 2004;303(5663):1526-9. Epub

2004/02/21. doi: 10.1126/science.1093620. PubMed PMID: 14976262.

130.    Borducchi EN, Liu J, Nkolola JP, Cadena AM, Yu W-H, Fischinger S, et al. Antibody and TLR7

agonist delay viral rebound in SHIV-infected monkeys. Nature. 2018;563(7731):360-4. doi:

10.1038/s41586-018-0600-6.

131.    Lim S-Y, Osuna CE, Hraber PT, Hesselgesser J, Gerold JM, Barnes TL, et al. TLR7 agonists

induce transient viremia and reduce the viral reservoir in SIV-infected rhesus macaques on antiretroviral

therapy. Science Translational Medicine. 2018;10(439). doi: 10.1126/scitranslmed.aao4521.

132.    SenGupta D, Brinson C, DeJesus E, Mills A, Shalit P, Guo S, et al. The TLR7 agonist

vesatolimod induced a modest delay in viral rebound in HIV controllers after cessation of antiretroviral

therapy. Sci Transl Med. 2021;13(599). Epub 2021/06/25. doi: 10.1126/scitranslmed.abg3071. PubMed

PMID: 34162752.

133.    Azar P, Mejia JE, Cenac C, Shaiykova A, Youness A, Laffont S, et al. TLR7 dosage

polymorphism shapes interferogenesis and HIV-1 acute viremia in women. JCI Insight. 2020;5(12). Epub

2020/06/20. doi: 10.1172/jci.insight.136047. PubMed PMID: 32554924; PubMed Central PMCID:

PMCPMC7406249.

134.    Abbas F, Cenac C, Youness A, Azar P, Delobel P, Guery JC. HIV-1 infection enhances innate

function and *TLR7* expression in female plasmacytoid dendritic cells. Life Sci Alliance. 2022;5(10).

Epub 2022/10/23. doi: 10.26508/lsa.202201452. PubMed PMID: 36271499; PubMed Central PMCID:

PMCPMC9441429.

135.    Klatt NR, Funderburg NT, Brenchley JM. Microbial translocation, immune activation, and HIV

disease. Trends Microbiol. 2013;21(1):6-13. Epub 20121011. doi: 10.1016/j.tim.2012.09.001. PubMed

PMID: 23062765; PubMed Central PMCID: PMCPMC3534808.

136.    Younas M, Psomas C, Reynes C, Cezar R, Kundura L, Portales P, et al. Microbial Translocation

Is Linked to a Specific Immune Activation Profile in HIV-1-Infected Adults With Suppressed Viremia. Front

Immunol. 2019;10:2185. Epub 20190913. doi: 10.3389/fimmu.2019.02185. PubMed PMID: 31572392;

PubMed Central PMCID: PMCPMC6753629.

137.    Nganou-Makamdop K, Talla A, Sharma AA, Darko S, Ransier A, Laboune F, et al. Translocated

microbiome composition determines immunological outcome in treated HIV infection. Cell.

2021;184(15):3899-914 e16. Epub 20210707. doi: 10.1016/j.cell.2021.05.023. PubMed PMID: 34237254;

PubMed Central PMCID: PMCPMC8316372.

138.    Massanella M, Fromentin R, Chomont N. Residual inflammation and viral reservoirs: alliance

against an HIV cure. Curr Opin HIV AIDS. 2016;11(2):234-41. doi: 10.1097/COH.0000000000000230.

PubMed PMID: 26575148; PubMed Central PMCID: PMCPMC4743501.

139.    Zilberman-Schapira G, Zmora N, Itav S, Bashiardes S, Elinav H, Elinav E. The gut microbiome in

human immunodeficiency virus infection. BMC Med. 2016;14(1):83. Epub 2016/06/04. doi:

10.1186/s12916-016-0625-3. PubMed PMID: 27256449; PubMed Central PMCID: PMCPMC4891875.

140.    Zevin AS, McKinnon L, Burgener A, Klatt NR. Microbial translocation and microbiome dysbiosis in

HIV-associated immune activation. Curr Opin HIV AIDS. 2016;11(2):182-90. Epub 2015/12/19. doi:

10.1097/COH.0000000000000234. PubMed PMID: 26679414; PubMed Central PMCID:

PMCPMC4752849.

141.    Penny CJ, Vassileva K, Jha A, Yuan Y, Chee X, Yates E, et al. Mining of Ebola virus entry

inhibitors identifies approved drugs as two-pore channel pore blockers. Biochim Biophys Acta Mol Cell

Res. 2019;1866(7):1151-61. Epub 20181105. doi: 10.1016/j.bbamcr.2018.10.022. PubMed PMID:

30408544; PubMed Central PMCID: PMCPMC7114365.

142.    Dobson SJ, Mankouri J, Whitehouse A. Identification of potassium and calcium channel inhibitors

as modulators of polyomavirus endosomal trafficking. Antiviral Res. 2020;179:104819. Epub 20200508.

doi: 10.1016/j.antiviral.2020.104819. PubMed PMID: 32389733; PubMed Central PMCID:

PMCPMC7205714.

143.    Choi B, Gatti PJ, Haislip AM, Fermin CD, Garry RF. Role of potassium in human

immunodeficiency virus production and cytopathic effects. Virology. 1998;247(2):189-99.

144.    Choi B, Fermin CD, Comardelle AM, Haislip AM, Voss TG, Garry RF. Alterations in intracellular

potassium concentration by HIV-1 and SIV Nef. Virol J. 2008;5:60. Epub 20080519. doi: 10.1186/1743-

422X-5-60. PubMed PMID: 18489774; PubMed Central PMCID: PMCPMC2396157.

145.    Wu M, Wang H, Yu H, Makhina E, Xu J, Dawson ES, et al. A potent and selective small molecule

Kir2.1 inhibitor.  Probe Reports from the NIH Molecular Libraries Program. Bethesda (MD)2010.

146.    Regoes RR, Rosenbloom DIS, Bacchetti P, Stone M, Deng X, Bosch RJ, et al. Assessing intra-

lab precision and inter-lab repeatability of outgrowth assays of HIV-1 latent reservoir size. PLOS

Computational Biology. 2019;15(4). doi: 10.1371/journal.pcbi.1006849.

147.    Stone M, Rosenbloom DIS, Bacchetti P, Deng X, Dimapasoc M, Keating S, et al. Assessing the

Suitability of Next-Generation Viral Outgrowth Assays to Measure Human Immunodeficiency Virus 1

Latent Reservoir Size. J Infect Dis. 2021;224(7):1209-18. doi: 10.1093/infdis/jiaa089. PubMed PMID:

32147687; PubMed Central PMCID: PMCPMC8514180.

148.    Offersen R, Yu W-H, Scully EP, Julg B, Euler Z, Sadanand S, et al. HIV Antibody Fc N-Linked

Glycosylation Is Associated with Viral Rebound. Cell Reports. 2020;33(11). doi:

10.1016/j.celrep.2020.108502.

149.    Ma T, McGregor M, Giron L, Xie G, George AF, Abdel-Mohsen M, et al. Single-cell glycomics

analysis by CyTOF-Lec reveals glycan features defining cells differentially susceptible to HIV. eLife.

2022;11. doi: 10.7554/eLife.78870.

150.    Giron LB, Palmer CS, Liu Q, Yin X, Papasavvas E, Sharaf R, et al. Non-invasive plasma glycomic

and metabolic biomarkers of post-treatment control of HIV. Nature Communications. 2021;12(1). doi:

10.1038/s41467-021-24077-w.

151.    Frank SL, Klisak I, Sparkes RS, Lusis AJ. A gene homologous to plasminogen located on human

chromosome 2q11-p11. Genomics. 1989;4(3):449-51. doi: 10.1016/0888-7543(89)90356-x. PubMed

PMID: 2714803.

152.    Morioka H, Morii T, Vogel T, Hornicek FJ, Weissbach L. Interaction of plasminogen-related

protein B with endothelial and smooth muscle cells in vitro. Exp Cell Res. 2003;287(1):166-77. doi:

10.1016/s0014-4827(03)00137-x. PubMed PMID: 12799192.

153.    Ichinose A, Espling ES, Takamatsu J, Saito H, Shinmyozu K, Maruyama I, et al. Two types of

abnormal genes for plasminogen in families with a predisposition for thrombosis. Proc Natl Acad Sci U S

A. 1991;88(1):115-9. doi: 10.1073/pnas.88.1.115. PubMed PMID: 1986355; PubMed Central PMCID:

PMCPMC50760.

154.    Ma J, Wei K, Liu J, Tang K, Zhang H, Zhu L, et al. Glycogen metabolism regulates macrophage-

mediated acute inflammatory responses. Nature Communications. 2020;11(1). doi: 10.1038/s41467-020-

15636-8.

155.    Maldarelli F, Wu X, Su L, Simonetti FR, Shao W, Hill S, et al. HIV latency. Specific HIV integration

sites are linked to clonal expansion and persistence of infected cells. Science. 2014;345(6193):179-83.

Epub 20140626. doi: 10.1126/science.1254194. PubMed PMID: 24968937; PubMed Central PMCID:

PMCPMC4262401.

156.    Peluso MJ, Bacchetti P, Ritter KD, Beg S, Lai J, Martin JN, et al. Differential decay of intact and

defective proviral DNA in HIV-1-infected individuals on suppressive antiretroviral therapy. JCI Insight.

2020;5(4). Epub 2020/02/12. doi: 10.1172/jci.insight.132997. PubMed PMID: 32045386; PubMed Central

PMCID: PMCPMC7101154.

157.    Falcinelli SD, Kilpatrick KW, Read J, Murtagh R, Allard B, Ghofrani S, et al. Longitudinal

Dynamics of Intact HIV Proviral DNA and Outgrowth Virus Frequencies in a Cohort of Individuals

Receiving Antiretroviral Therapy. J Infect Dis. 2021;224(1):92-100. Epub 2020/11/21. doi:

10.1093/infdis/jiaa718. PubMed PMID: 33216132; PubMed Central PMCID: PMCPMC8253129.

158.    Gogarten SM, Sofer T, Chen H, Yu C, Brody JA, Thornton TA, et al. Genetic association testing

using the GENESIS R/Bioconductor package. Bioinformatics. 2019;35(24):5346-8. Epub 2019/07/23. doi:

10.1093/bioinformatics/btz567. PubMed PMID: 31329242; PubMed Central PMCID: PMCPMC7904076.

159.    Stegle O, Parts L, Piipari M, Winn J, Durbin R. Using probabilistic estimation of expression

residuals (PEER) to obtain increased power and interpretability of gene expression analyses. Nature

protocols. 2012;7(3):500.

160.    Pantaleo G, Graziosi C, Butini L, Pizzo PA, Schnittman SM, Kotler DP, et al. Lymphoid organs

function as major reservoirs for human immunodeficiency virus. Proc Natl Acad Sci U S A.

1991;88(21):9838-42. Epub 1991/11/01. doi: 10.1073/pnas.88.21.9838. PubMed PMID: 1682922;

PubMed Central PMCID: PMCPMC52816.

161.    Evering TH, Mehandru S, Racz P, Tenner-Racz K, Poles MA, Figueroa A, et al. Absence of HIV-1

evolution in the gut-associated lymphoid tissue from patients on combination antiviral therapy initiated

during primary infection. PLoS Pathog. 2012;8(2):e1002506. Epub 2012/02/10. doi:

10.1371/journal.ppat.1002506. PubMed PMID: 22319447; PubMed Central PMCID: PMCPMC3271083.

162.    Josefsson L, von Stockenstrom S, Faria NR, Sinclair E, Bacchetti P, Killian M, et al. The HIV-1

reservoir in eight patients on long-term suppressive antiretroviral therapy is stable with few genetic

changes over time. Proc Natl Acad Sci U S A. 2013;110(51):E4987-96. Epub 2013/11/28. doi:

10.1073/pnas.1308313110. PubMed PMID: 24277811; PubMed Central PMCID: PMCPMC3870728.

163.    Imamichi H, Degray G, Dewar RL, Mannon P, Yao M, Chairez C, et al. Lack of

compartmentalization of HIV-1 quasispecies between the gut and peripheral blood compartments. J Infect

Dis. 2011;204(2):309-14. Epub 2011/06/16. doi: 10.1093/infdis/jir259. PubMed PMID: 21673043; PubMed

Central PMCID: PMCPMC3114472.

164.    Ramirez CM, Sinclair E, Epling L, Lee SA, Jain V, Hsue PY, et al. Immunologic profiles

distinguish aviremic HIV-infected adults. AIDS. 2016;30(10):1553-62. Epub 2016/02/09. doi:

10.1097/QAD.0000000000001049. PubMed PMID: 26854811; PubMed Central PMCID:

PMCPMC5679214.

165.    Boufassa F, Lechenadec J, Meyer L, Costagliola D, Hunt PW, Pereyra F, et al. Blunted response

to combination antiretroviral therapy in HIV elite controllers: an international HIV controller collaboration.

PLoS One. 2014;9(1):e85516. Epub 2014/01/28. doi: 10.1371/journal.pone.0085516. PubMed PMID:

24465584; PubMed Central PMCID: PMCPMC3894966.

166.    Emu B, Sinclair E, Hatano H, Ferre A, Shacklett B, Martin JN, et al. HLA class I-restricted T-cell

responses may contribute to the control of human immunodeficiency virus infection, but such responses

are not always necessary for long-term virus control. J Virol. 2008;82(11):5398-407. Epub 2008/03/21.

doi: 10.1128/JVI.02176-07. PubMed PMID: 18353945; PubMed Central PMCID: PMCPMC2395228.

167.    Grebe E, Facente SN, Bingham J, Pilcher CD, Powrie A, Gerber J, et al. Interpreting HIV

diagnostic histories into infection time estimates: analytical framework and online tool. BMC Infect Dis.

2019;19(1):894. Epub 2019/10/28. doi: 10.1186/s12879-019-4543-9. PubMed PMID: 31655566; PubMed

Central PMCID: PMCPMC6815418.

168.    Andrews S. FastQC: a quality control tool for high throughput sequence data. 2010. 2017.

169.    Schneider VA, Graves-Lindsay T, Howe K, Bouk N, Chen H-C, Kitts PA, et al. Evaluation of

GRCh38 and de novo haploid genome assemblies demonstrates the enduring quality of the reference

assembly. Genome Research. 2017;27(5):849-64. doi: 10.1101/gr.213611.116.

170.    Harrow J, Frankish A, Gonzalez JM, Tapanari E, Diekhans M, Kokocinski F, et al. GENCODE:

the reference human genome annotation for The ENCODE Project. Genome research. 2012;22(9):1760-

74.

171.    Dobin A, Gingeras TR. Mapping RNA-seq reads with STAR. Current protocols in bioinformatics.

2015;51(1):11.4. 1-.4. 9.

172.    Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a

reference genome. BMC bioinformatics. 2011;12(1):1-16.

173.    Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal

RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.

174.    Robinson MD, Oshlack A. A scaling normalization method for differential expression analysis of

RNA-seq data. Genome biology. 2010;11(3):1-9.

175.    Law CW, Chen Y, Shi W, Smyth GK. voom: Precision weights unlock linear model analysis tools

for RNA-seq read counts. Genome biology. 2014;15(2):1-17.

176.    Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression

analyses for RNA-sequencing and microarray studies. Nucleic acids research. 2015;43(7):e47-e.

177.    Devlin B, Roeder K. Genomic control for association studies. Biometrics. 1999;55(4):997-1004.

178.    Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R package for comparing biological themes

among gene clusters. Omics: a journal of integrative biology. 2012;16(5):284-7.

1151

**Table 1.** Descriptive statistics for the study population of 191 HIV-infected ART-suppressed non-controllers. Median frequencies (with interquartile ranges) are shown below unless otherwise specified.

| Descriptive Characteristic | Total (N=191) | Early-Treated[a] (N=54) | Later-Treated[a] (N=137) |
|---|---|---|---|
| Male (%)[b] | 183 (96%) | 54 (100%) | 129 (94%) |
| Age (years) | 47 (13) | 44 (12) | 47 (13) |
| Nadir CD4+ T cell count (cells/mm3) | 352 (251) | 522 (346) | 304 (190) |
| Maximum pre-ART HIV RNA (log$_{10}$copies/mL) | 5.1 (0.9) | 5.6 (0.7) | 5.0 (0.8) |
| Duration of ART suppression (years) | 5.1 (4.2) | 6.0 (4.3) | 4.7 (4.2) |
| Timing of ART initiation (years) | 2.0 (4.6) | 0.20 (0.19) | 3.5 (4.3) |
| HIV intact DNA (log$_{10}$copies/10$^6$ CD4+ T cells) | 1.3 (1.0) | 1.3 (0.5) | 1.9 (1.0) |
| HIV total DNA (log$_{10}$copies/10$^6$ CD4+ T cells) | 1.0 (1.3) | 0.4 (1.3) | 1.2 (1.3) |
| HIV unspliced RNA (log$_{10}$copies/10$^6$ CD4+ T cells) | 3.2 (0.8) | 3.0 (0.8) | 3.3 (0.7) |
| HIV RNA/DNA | 2.3 (1.0) | 2.4 (1.0) | 2.3 (0.9) |

[a] Early-treated = Individuals who initiated ART within 6 months of the date of detected HIV infection; later-treated = Individuals who initiated ART after 6 months of date of detected HIV infection.
[b] Absolute frequencies (with percent)

**Figure 1.** Principal component analysis (PCA) plot of the population structure. Principal component analysis (PCA) plot of the population structure of the full study cohort (a). Secondary PCA plot of the European ancestry subpopulation only (b) defined by the dashed box in the lower left of panel (a). Genetic PCs were calculated from genetic data from our whole exome analysis [25]. Most of the population was of European ancestry (bottom left of) (a) some continued variability. Some continued variability was observed in European ancestry subgroup (b). Self-identified race/ethnicity shown in the legend. Frequencies for participants were recorded as: White/European American (62%), Black/African American (14%), Hispanic/Latino (11%), Mixed Ethnicity/Multiracial (6%), Asian (4%), Pacific Islander (2%), Native American (<1%), and Middle Eastern (<1%).

**Figure 2.** Correlations between three measures of HIV reservoir size. HIV unspliced RNA (usRNA) was significantly correlated with (a) HIV Total DNA (tDNA) and (b) HIV intact DNA; (c) tDNA and intact DNA were not correlated with one another.

**Figure 3.** Measures of the HIV reservoir from peripheral CD4+ T cells were associated with timing of ART initiation. Panels A-D correspond to HIV tDNA, HIV usRNA, RNA/DNA, and Intact DNA, respectively. Spearman correlation and corresponding p-value are shown in each case. Earlier timing of ART initiation (<6 months from infection) was statistically significantly associated with smaller HIV intact DNA, tDNA, and usRNA.

**Table 2.** Differentially expressed host genes in relation to $\log_{10}$copies of HIV unspliced RNA (usRNA) in the full cohort (top panel) and the European ancestry subpopulation (bottom panel), at a Benjamini-Hochberg false discovery rate (FDR) of q<0.05. HIV usRNA was significantly associated with downregulation of 17 host genes, including KCNJ2, a novel association with a gene encoding for an inwardly rectifying potassium (Kir2.1) channel (which may enhance HIV entry and release into host cells [32]), gap junction (GJB2) as well as genes involved in pathogen pattern recognition (TLR7), inflammasome cytokine activation (IL1A, CSF3, TNFAIP5, TNFAIP9, TNFAIP9), and chemokine production (CXCL3, CXCL10). An additional list of host genes associated with HIV usRNA at an FDR q<0.25 is shown in **Supplemental Table 2**.

| HIV Unspliced RNA | | | | | | |
|---|---|---|---|---|---|---|
| Gene | Gene Name | p[a] | q[b] | FC[c] | % Change[d] | Description |
| **Full Cohort** | | | | | | |
| KCNJ2 | Potassium Inwardly Rectifying Channel Subfamily J Member 2, kir2.1 | 1.49E-07 | 0.003 | 0.903 | -9.7 | KCNJ2, encodes for an inwardly rectifying potassium channel (Kir2.1). Inwardly rectifying potassium ion channels can regulate HIV 1 entry and release into host cells [32]. Tight regulation of potassium ion concentrations has been shown to play a critical role in HIV-1 virus production in CD4+ T cells in cell culture models [143]. |
| IL1A | Interleukin-1 alpha | 1.55E-06 | 0.012 | 0.904 | -9.6 | IL-1 is a potent proinflammatory cytokine that regulates inflammation by triggering a cascade of inflammatory mediators via NRLP3 inflammasome activation pathway [81, 82]. IL-1 is an "upstream" pro-inflammatory inducer of interleukin-6 (IL-6) [83], which is the strongest biomarker for non-AIDS morbidity (e.g., myocardial infarction, stroke, malignancy) [84-87] and mortality [80, 86-89] among HIV-infected ART-suppressed individuals in resource-rich countries. |
| GJB2 | Gap Junction Protein Beta 2 | 2.26E-06 | 0.012 | 0.929 | -7.1 | GJB2 encodes for gap junction beta 2 protein, or connexin 26 (CX26), which acts as a communication channel between cells, facilitating transport of signaling molecules (calcium and cyclic AMP, ATP), and charged ions (K+, Ca+) [33, 34]. HIV-1 is thought to exploit these communication channels to disseminate infection and associated inflammation even in the absence of viral replication [46, 47]. |
| KCNJ2-AS1 | KCNJ2 antisense RNA 1 | 2.49E-06 | 0.012 | 0.916 | -8.4 | KCNJ2-AS1 is a long non-coding RNA (lncRNA) encodes for KCNJ2 antisense RNA 1. |
| AC034199.1 | AC034199.1 | 2.88E-06 | 0.012 | 0.946 | -5.4 | Novel transcript: no data found in literature in association with HIV or immune response. |
| CSF3 | Colony Stimulating Factor 3 (G-CSF) | 3.93E-06 | 0.013 | 0.925 | -7.5 | CSF3 encodes for granulocyte stimulating factor 3 (G-CSF), a member of the IL-6 superfamily of cytokines (78) and is also a growth factor for Neutrophils [90]. CSF3 modulates the function of CD4+ T cells by regulating cytokine production, their differentiation, and Treg induction [91]. |
| TNFAIP6 | Tumor Necrosis Factor alpha induced protein 6 | 5.99E-06 | 0.016 | 0.924 | -7.6 | TNFAIP6 encodes for TNF-stimulated gene 6 protein (TSG-6), which, like TNFAIP5 (TSG-5), is induced by tumor necrosis factor α (TNF-α) and interleukin-1 (IL-1) in response to lipopolysaccharide (LPS)-stimulation [101, 102]. |

| Gene | Name | | | | | Description |
|---|---|---|---|---|---|---|
| TLR7 | Toll Like Receptor 7 | 6.59E-06 | 0.016 | 0.929 | -7.1 | TLR7 is a member of toll-like receptor family of genes which plays critical role in pathogen recognition, activation of innate immune response and functions as a bridge between innate and adaptive immunity [127]. TLR7, is a pattern recognition receptor that can sense HIV single-stranded RNA (ssRNA) in endosomes [128, 129]. |
| MRAS | Muscle RAS oncogene homolog | 1.07E-05 | 0.024 | 0.945 | -5.5 | MRAS encodes for a protein in the Ras family of small GTPases which functions as signal transducers in cellular processes. |
| TNFAIP9 | Tumor Necrosis Factor alpha induced protein 9 | 1.53E-05 | 0.031 | 0.931 | -6.9 | TNFAIP9, encodes for TNF-α induced protein 9 (TSG-9). It is also known as STEAP4 (six transmembrane epithelial antigen of prostate 4) involved in negative regulation of NF-κB, STAT-3 signaling, and IL-6 production [105, 106]. |
| MIR3945HG | MIR3945 Host Gene | 2.09E-05 | 0.038 | 0.942 | -5.8 | MIR3945HG is an interferon stimulated lncRNA. |
| DAPK1-IT1 | DAPK1 Intronic Transcript 1 | 2.72E-05 | 0.043 | 0.950 | -5.0 | DAPK1-IT1 is a lncRNA transcribed from the death associated protein kinases 1 (DAPK1). |
| OR2B11 | Olfactory Receptor Family 2 Subfamily B Member 11 | 2.96E-05 | 0.043 | 0.939 | -6.1 | OR2B11 is a member of G-protein-coupled receptors (GPCR) responsible for the recognition and G protein-mediated transduction of odorant signals. |
| CXCL3 | C-X-C Motif Chemokine Ligand 3 | 3.03E-05 | 0.043 | 0.928 | -7.2 | CXCL3 is a member of CXC subfamily called cytokine-induced neutrophil chemoattractant (CINCs). CXCL3 is involved in adhesion and migration of monocytes [109, 110], neutrophils chemoattraction [111-113], and angiogenesis [114]. |
| TNFAIP5 /PTX3 | TNF Alpha-Induced Protein 5 (TNFAIP5), Pentraxin-related protein (PTX3), Tumor Necrosis Factor-Inducible Protein TSG-14 (TSG14). | 3.27E-05 | 0.043 | 0.941 | -5.9 | TNFAIP5 is a pattern recognition receptor (PRRs) that is induced in response to TNF- α, TLR engagement and IL-1β signaling [96, 97]. |
| RRN3P4 | RRN3 Pseudogene 4 | 3.97E-05 | 0.049 | 0.950 | -5.0 | Pseudogene |
| CXCL10 | C-X-C Motif Chemokine Ligand 10 | 4.21E-05 | 0.049 | 0.908 | -9.2 | CXCL10 encodes for IP-10 (interferon gamma-induced protein 10) (interferon gamma-induced protein 10) which recruits activated Th1 lymphocytes to sites of infection [117-119] and in HIV, signals through TLR7/9-dependent pathways [119], predicts HIV disease progression [120, 121], correlates with acute HIV seroconversion [122], and promotes of HIV latency [123, 124]. |

## European Ancestry Subpopulation

| Gene | Name | | | | | Description |
|---|---|---|---|---|---|---|
| TLR7[a] | Toll Like Receptor 7 | 1.48E-06 | 0.018 | 0.906 | -9.4 | TLR7 is a member of toll-like receptor family of genes which plays critical role in pathogen recognition, activation of innate immune response and functions as a bridge between innate and adaptive immunity [127]. TLR7, is a pattern recognition receptor that can sense HIV single-stranded RNA (ssRNA) in endosomes [128, 129]. |
| GJB2[a] | Gap Junction Protein Beta 2 | 2.70E-06 | 0.018 | 0.909 | -9.1 | GJB2 encodes for gap junction beta 2 protein, or connexin 26 (CX26), which acts as a communication channel between cells, facilitating transport of signaling molecules (calcium and cyclic AMP, ATP), and charged ions (K+, Ca+) [33, 34]. HIV-1 is thought to exploit these communication channels to disseminate infection and associated inflammation even in the absence of viral replication [46, 47]. |
| AC034199.1[a] | Ac034199.1 | 3.09E-06 | 0.018 | 0.930 | -7.0 | novel transcript |
| PPP1R17 | Protein Phosphatase 1 Regulatory Subunit 17, | 4.20E-06 | 0.018 | 0.934 | -6.6 | PPP1R17 (Protein Phosphatase 1 Regulatory Subunit 17) is a substrate for cGMP-dependent protein kinase. |

| IGSF6 | Immunoglobulin Superfamily Member 6 | **4.73E-06** | 0.018 | 0.972 | **-2.8** | Immunoglobulin superfamily member 6 (IGSF6) is also known as downregulated by activation (DORA). |
|---|---|---|---|---|---|---|
| AL133163.2 | Al133163.2 | **7.81E-06** | 0.025 | 0.945 | **-5.5** | novel transcript |

[a] p = two sided p-value.
[b] q = two-sided false discovery rate (FDR) Benjamini-Hochberg q-value.
[c] FC = fold-change in host gene expression per two-fold change in copies of HIV from multivariate model adjusted for age, sex, nadir CD4+ T cell count, timing of ART initiation, ancestry (PCs), and residual variability (probabilistic estimation of expression residuals, PEERs).
[d] % Change = percent change in host gene expression per two-fold change in copies of HIV.
[e] % also significant in full cohort analysis.

**Figure 4.** Network analysis of the top differentially expressed genes (see **Table 2** and **Supplemental Table 2**) associated with HIV unspliced RNA demonstrated that the top significant genes mapped to immunologic pathways involving bacterial translocation (e.g., TLR4 signaling, activated by bacterial lipopolysaccharide, LPS) and pro-inflammatory responses (e.g., IL-1β signaling, NLRP3 inflammasome assembly, Th2 cell cytokine production). A Benjamini-Hochberg false discovery rate (FDR) of q<0.05 was used to generate nodes (circles) based on kappa scores ≥0.4. The size of the nodes reflects the enrichment significance of the terms, and the different colors represent distinct functional groups.

**Figure 5.** Network analysis of the top statistically significant gene sets associated with HIV unspliced RNA (see **Supplement Table 4**). Gene sets related to immunologic pathways involving bacterial translocation (e.g., response to bacterium, LPS-mediated signaling pathway), and inflammatory signaling (e.g., IL-1β, IL-6, IL-10, TNF-α), were significantly associated with HIV usRNA. A Benjamini-Hochberg false discovery rate (FDR) of q<0.05 was used to generate nodes (circles) based on kappa scores ≥0.4. The size of the nodes reflects the enrichment significance of the terms, and the different colors represent distinct functional groups.