TITLE

Analysis of CDPK1 targets identifies a trafficking adaptor complex that regulates
microneme exocytosis in *Toxoplasma*

AUTHORS

Alex W Chan[1,2], Malgorzata Broncel[3], Nicole Haseley[1], Sundeep Chakladar[1], Elena Andree[1], Alice
L Herneisen[1,2], Emily Shortt[1], Moritz Treeck[3], Sebastian Lourido[1,2]*

7
8 [1] Whitehead Institute for Biomedical Research, Cambridge, MA, USA
9 [2] Biology Department, Massachusetts Institute of Technology, Cambridge, MA, USA
10 [3] Signaling in Apicomplexan Parasites Laboratory, The Francis Crick Institute, London, UK
11 * Correspondence: lourido@wi.mit.edu
12

ABSTRACT

Apicomplexan parasites use Ca^{2+}-regulated exocytosis to secrete essential virulence factors
from specialized organelles called micronemes. Ca^{2+}-dependent protein kinases (CDPKs) are
required for microneme exocytosis; however, the molecular events that regulate trafficking and
fusion of micronemes with the plasma membrane remain unresolved. Here, we combine sub-
minute resolution phosphoproteomics and bio-orthogonal labeling of kinase substrates in
Toxoplasma gondii to identify 163 proteins phosphorylated in a CDPK1-dependent manner. In
addition to known regulators of secretion, we identify uncharacterized targets with predicted
functions across signaling, gene expression, trafficking, metabolism, and ion homeostasis. One
of the CDPK1 targets is a putative HOOK activating adaptor. In other eukaryotes, HOOK
homologs form the FHF complex with FTS and FHIP to activate dynein-mediated trafficking of
endosomes along microtubules. We show the FHF complex is partially conserved in *T. gondii*,
consisting of HOOK, an FTS homolog, and two parasite-specific proteins (TGGT1_306920 and
TGGT1_316650). CDPK1 kinase activity and HOOK are required for the rapid apical trafficking
of micronemes as parasites initiate motility. Moreover, parasites lacking HOOK or FTS display
impaired microneme protein secretion, leading to a block in the invasion of host cells. Taken
together, our work provides a comprehensive catalog of CDPK1 targets and reveals how
vesicular trafficking has been tuned to support a parasitic lifestyle.
31

INTRODUCTION

Ca^{2+}-regulated exocytosis is ubiquitous among eukaryotes. This signaling paradigm regulates
an array of processes such as neurotransmitter release in neurons, hormone secretion in
endocrine cells, and histamine secretion in mast cells (Pang and Südhof, 2010). Ca^{2+}-regulated
exocytosis is also critical for apicomplexan parasites that are the causative agents of rampant,
life-threatening diseases including malaria, toxoplasmosis, and cryptosporidiosis (Havelaar et
al., 2015). Central to their pathogenesis is their ability to transition from intracellular replicative

stages to extracellular motile stages, which involves a unique form of cellular movement called
gliding, egress from the primary host cell, and invasion into a new host. These processes are
driven by the Ca^{2+}-regulated exocytosis of apicomplexan-specific membrane-bound organelles
called micronemes and rhoptries.

The sequential exocytosis of micronemes and rhoptries is required to promote extracellular
motile stages of the parasite (Blader et al., 2016; Carruthers and Sibley, 1997). Micronemes are
localized to the parasite apex and their positioning is dependent on cortical microtubules,
ultrastable filaments that polymerize down the length of the parasite (Chen et al., 2015; Leung et
al., 2017; Wang et al., 2021). Exocytosis of microneme cargo enables host cell rupture by
releasing perforin-like proteins during egress and translocation of exposed adhesins required for
gliding and attachment to new host cells (Carruthers and Sibley, 1997; Carruthers and Tomley,
2008; Kafsack et al., 2009). Multiple microneme proteins, including the associated cysteine
repeat modular proteins (CRMP) complex, are also required to trigger the exocytosis of rhoptries
upon host cell contact (Kessler et al., 2008; Sidik et al., n.d.; Singer et al., 2022; Sparvoli et al.,
2022). Rhoptry proteins include effectors that modulate host responses and transmembrane
proteins that are embedded into the host plasma membrane to enable active invasion (Bradley
and Sibley, 2007; Lamarque et al., 2011, 2014; Ong et al., 2010; Tyler and Boothroyd, 2011).
Intracellular Ca^{2+} release is necessary and sufficient to trigger the rapid trafficking and exocytosis
of micronemes (Carruthers et al., 1999a, 1999b; Endo et al., 1982; Sidik et al., 2016b). While Ca^{2+}
is also necessary for rhoptry discharge, exocytosis relies on additional cellular processes such
as microneme exocytosis (Coleman et al., 2018; Segev-Zarko et al., 2022). While the exocytosis
of micronemes and rhoptries is known to be critical for parasite motility, the mechanisms linking
Ca^{2+} signaling to their trafficking and fusion to the plasma membrane are still unclear.

Ca^{2+} signals in apicomplexans are primarily transduced by Ca^{2+}-dependent protein kinases
(CDPKs) (Billker et al., 2009; Farrell et al., 2012; Garrison et al., 2012; Kumar et al., 2017; Lourido
et al., 2012, 2010; Lourido and Moreno, 2015; Luo et al., 2005; Márquez-Nogueras et al., 2021;
McCoy et al., 2012; Sebastian et al., 2012). CDPKs are serine/threonine protein kinases that are
unique to apicomplexans and plants. CDPKs are activated by directly binding to Ca^{2+}, in contrast
to Ca^{2+}/calmodulin-dependent protein kinases (CaMKs) found in animals, which are indirectly
regulated by Ca^{2+}-bound calmodulin (CaM) (Ojo et al., 2010; Wernimont et al., 2010). CDPKs are
crucial for apicomplexan infection—yet are absent from mammals—making them attractive drug
targets; however, their mechanisms of action are still not well understood at a cellular and
molecular level. In *T. gondii*, Ca^{2+}-dependent protein kinase 1 (CDPK1) is required for the Ca^{2+}-
regulated exocytosis of micronemes, impacting all steps of parasite motility including egress,
gliding, and invasion (Lourido et al., 2010). Small-molecule competitive inhibitors against CDPK1
have been identified and have shown some activity against *T. gondii* (Doggett et al., 2014;
Johnson et al., 2012; Lourido et al., 2013b, 2010; Winzer et al., 2015). Identifying the signaling
pathways regulated by CDPK1 could reveal the pathways controlling microneme and rhoptry
exocytosis.

The Ca^{2+} that activates CDPK1 and other cellular processes is released from intracellular stores
following cyclic nucleotide–mediated activation of protein kinase G (PKG) (Bisio and Soldati-
Favre, 2019; Brown et al., 2017; Sidik et al., 2016a). This process can be artificially induced by
treating parasites with cGMP specific phosphodiesterase (PDE) inhibitors zaprinast or BIPPO
that indirectly activate PKG (**Figure 1A**)(Lourido et al., 2012; Nofal et al., 2022; Sidik et al., 2016a;
Yuasa et al., 2005). PDE inhibition activates PKG within seconds and triggers the Ca^{2+} and lipid
signaling nodes controlling parasite motility, including the exocytosis of micronemes (**Figure**
1A)(Lourido et al., 2012; Yuasa et al., 2005). The identity of the CDPK1 substrates that contribute
to the regulation of microneme exocytosis remains unknown. Determining CDPK1-dependent
phosphorylation is challenging because the signaling pathways controlling parasite motility are
rapid and integrate signals from multiple kinases.
92

Global phosphoproteomic studies found that Ca^{2+}-dependent phosphorylation included proteins
involved in signal transduction, motility, exocytosis, cytoskeleton, in addition to many proteins
with unknown functions; however, the functional relevance and organization of these proteins—
especially CDPK1—in regulating exocytosis remain unclear (Herneisen et al., 2022; Invergo et
al., 2017; McCoy et al., 2012; Nofal et al., 2022; Treeck et al., 2014, 2011). Here, we utilized time-
resolved phosphoproteomics and chemical genetics to identify 163 proteins phosphorylated by
CDPK1. Our comprehensive analysis of CDPK1 not only identified phosphoregulation of known
factors involved in parasite motility, but also revealed new regulators of exocytosis. We identified
a conserved HOOK complex that is phosphorylated by CDPK1 and is required for microneme
exocytosis by mediating the rapid trafficking of micronemes during parasite motility.
103

RESULTS

Identifying CDPK1-dependent phosphorylation with sub-minute resolution

We examined the effect of CDPK1 on the phosphoproteome. To control the expression of
CDPK1, we endogenously tagged the kinase with a C-terminal auxin-inducible degron (AID) for
rapid conditional knockdown upon treating parasites with auxin (**Figure 1A**)(Brown et al., 2018;
Shortt et al., 2022; Smith et al., 2022). CDPK1 was robustly depleted from extracellular parasites
following auxin treatment for 3.5 hrs (**Figure 1B**). We compared vehicle- and auxin-treated
parasites at four timepoints in the 5 min following zaprinast stimulation (0, 9, 30, and 300 s) to
capture the earliest changes following Ca^{2+} flux (**Figure 1C**). We used TMTpro labeling to
multiplex 16 samples enabling analysis of a complete time course comparing vehicle- and auxin-
treated parasites in biological duplicate in a single LC-MS/MS experiment (Li et al., 2020).
Samples treated with DMSO instead of zaprinast served as a control. We enriched for
phosphopeptides using sequential metal-oxide affinity chromatography (SMOAC) (Tsai et al.,
2014). In total, we generated four datasets: an enriched zaprinast phosphoproteome [1], a
zaprinast proteome (unenriched) [2], an enriched DMSO phosphoproteome [3], and a DMSO
proteome (unenriched) [4]. Of the 4,255 parasite proteins quantified in the zaprinast proteome
by LC-MS/MS, CDPK1 was the only protein with a greater than two-fold depletion in parasites

treated with auxin (**Figure 1D**). The remainder of the proteome was largely stable (median $\log_2$-
fold change = -0.002 ± 0.089 M.A.D).

We first quantified peptide abundances across time relative to the 0 sec vehicle-treated time
point to identify CDPK1-dependent phosphorylation, which yielded kinetic profiles of individual
phosphopeptides in vehicle and auxin conditions (**Figure 1C**). The zaprinast phosphoproteome
included 2,570 phosphorylated proteins, represented by 10,594 unique phosphopeptides
quantified across both biological replicates (**Figure 1E**). We calculated the area under the curve
(AUC) of individual phosphopeptide profiles for vehicle ($AUC_{vehicle}$) and auxin (AUC_{auxin}) conditions.
To identify the subset of phosphopeptides exhibiting CDPK1-dependent phosphorylation, we
calculated the difference between AUC values ($AUC_{difference}$) by comparing the distribution of
$AUC_{difference}$ values in the zaprinast phosphoproteome to a null distribution derived from the
DMSO phosphoproteome (**Figure 1—figure supplement 1**). This approach allowed us to
account for the variability among AUCs under conditions that did not exhibit dynamic changes.
We identified 74 unique CDPK1-dependent phosphopeptides across both biological replicates
(Group A), belonging to 69 proteins (**Figure 1E–F**). Additionally, we identified peptides that were
zaprinast responsive—despite being CDPK1 independent—by comparing the distribution of
$AUC_{vehicle}$ values in the zaprinast phosphoproteome to a null distribution derived from the DMSO
phosphoproteome, resulting in 809 unique phosphopeptides representing 501 proteins. We
utilized projection-based clustering to sort kinetic profiles of CDPK1-independent peptides into
three groups, in which Group B and Group C contained phosphopeptides that increased in
abundance and Group D contained phosphopeptides that decreased in abundance (Thrun and
Ultsch, 2021)(**Figure 1G**, **Figure 1—figure supplement 1**) (Thrun and Ultsch, 2021). Significant
changes at the peptide level are attributed to altering levels of phosphorylation as opposed to
protein levels. Of the 543 proteins exhibiting zaprinast-dependent phosphorylation, 484 were
quantified at the proteome level. When comparing the 300 to 0 sec time points, zaprinast-
dependent proteins displayed no significant changes (median $\log_2$-fold change = 0 ± 0.074
M.A.D). This was consistent with the rapid time scale of these experiments and the overall
stability of the 4,255 proteins quantified in the proteome (median $\log_2$-fold change = 0.01 ± 0.089
M.A.D).

Gene ontology analysis identified several classes of genes that may be relevant to the regulation
of zaprinast responses within the different groups (**Figure 1H**). CDPK1-independent
phosphorylation and dephosphorylation was prevalent amongst proteins regulating cyclic
nucleotide signaling (ACβ, GC, UGO, PDE1, PDE2, PDE7, PDE9, PDE10, and PKG), lipid
signaling (PI4K, PI4P5K, PI-PLC, DGK1, and PAP1), and Ca^{2+} signaling (CDPK2A and CDPK3).
CDPK1-independent phosphorylation was also observed on downstream proteins regulating
microneme exocytosis (APH and DOC2.1), rhoptry exocytosis (ARO, AIP, PL3, and NdP2), gliding
motility (AKMT and MyoA), and homeostasis/biogenesis (NHE3, VHA1, NST2, and ERK7). MyoA
phosphorylation was observed at S20 and S21, which was previously shown to be
phosphorylated by CDPK3 and required for gliding motility (Gaji et al., 2015). SCE1
phosphorylation was observed at S225, which was previously shown to be phosphorylated by

CDPK3 and required to relieve inhibition during Ca^{2+}-stimulated egress (McCoy et al., 2017). Our
analysis highlights temporally-resolved changes in the phosphoproteome in response to
zaprinast stimulation. We identify a subset of proteins phosphorylated by CDPK1 that likely
represent factors involved in motility-related exocytosis, which we discuss in further detail below.
167

Myristoylation modulates CDPK1 activity and alters its interacting partners

CDPK1 was found to be myristoylated on Gly2 (Broncel et al., 2020). Myristoylation results in
lipid modifications on proteins that can impact membrane targeting and protein-protein
interactions (Martin et al., 2011; Wright et al., 2014). The myristoylated proteome of *T. gondii* was
recently characterized using metabolic tagging and enrichment (Broncel et al., 2020). In this
approach, parasites were grown in the presence of a myristic acid analogue containing a terminal
alkyne group (YnMyr) that allowed enrichment of labeled proteins through click chemistry (Heal
et al., 2011). We assessed whether myristoylation of CDPK1 affects the kinase's function. We
validated myristoylation of CDPK1 using myristoylation-dependent pull downs followed by
immunoblot detection and MS analysis (**Figure 2—figure supplement 1**). To investigate the role
of myristoylation on CDPK1 function, an inducible knock-down strain (iKD) was generated by
introducing a mAID-Myc tag at its C terminus (**Figure 2—figure supplement 1**). Auxin-
dependent depletion was confirmed using immunoblot (**Figure 2—figure supplement 1**).
Depletion of CDPK1 abolished ionophore-induced egress, as expected from previous results
(**Figure 2—figure supplement 1**)(Lourido et al., 2010). Next, we complemented the iKD
parasites by introducing HA-tagged WT (cWT) or myristoylation defective (cMut, G2A) copies of
CDPK1 into the *UPRT* locus (**Figure 2A**). We verified correct integration of both complementation
constructs (**Figure 2—figure supplement 1**), and confirmed their equivalent and constitutive
expression, as well as the auxin sensitivity of the endogenous mAID-tagged CDPK1 (**Figure 2B**).
As predicted, the cMut allele was not myristoylated, based on acylation-dependent pull downs
and immunoblotting (**Figure 2C**).
189

Given that myristoylation is frequently reported to facilitate membrane association, we examined
the localization of cWT and cMut CDPK1 by immunofluorescence (**Figure 2D**). No clear
differences were detected between the punctate cytosolic patterns of cWT and cMut. We
explored possible effects of myristoylation on the subcellular fractionation of CDPK1 resolved
using differential centrifugation (**Figure 2E**). First, we evaluated the fractionation pattern of the
endogenous, myristoylated CDPK1. YFP-expressing parasites were metabolically labeled with
Myr or YnMyr and lysed in a hypotonic buffer to preserve intact membrane structures. Next,
lysates were fractionated to generate a low-speed pellet and supernatant at 16,000 x *g*. The low-
speed supernatant was fractionated further into a high-speed pellet and supernatant at 100,000
x *g*. Click reaction based pull down and immunoblotting were used to resolve myristoylation-
dependent partitioning. In contrast to the doubly acylated GAP45, which was present exclusively
in the low-speed pellet, myristoylated CDPK1 was observed in the low-speed pellet and
supernatant, and the high-speed pellet (**Figure 2E**), suggesting a potential association with
membranous structures or higher molecular weight complexes. We next used both the cWT and
cMut lines to elucidate any myristoylation-dependent changes to CDPK1 localization. While cWT

could be found predominantly in the high-speed pellet, loss of myristoylation released cMut into
the high-speed supernatant confirming an association with membranous structures or large
protein complexes (**Figure 2F**).
208
To evaluate the role of CDPK1 myristoylation in parasite fitness, we performed plaque assays
comparing the various strains (**Figure 2G**). In the presence of the endogenous copy of CDPK1,
both complemented lines developed normally. However, upon auxin-mediated depletion of
endogenous CDPK1, cMut plaque size substantially decreased. This finding demonstrates that
one or more steps of the *T. gondii* lytic cycle are negatively affected by the loss of CDPK1
myristoylation. In light of CDPK1's known function, we next explored whether CDPK1
myristoylation might impact the parasite's ability to egress from host cells. In the absence of
auxin, cWT, cMut, and the parental line (iKD) egressed within two min of ionophore stimulation
(**Figure 2—figure supplement 1**). While cWT parasites maintained similar egress kinetics
following auxin treatment, cMut parasites showed a significant delay after two min of treatment
(**Figure 2H**). This egress delay was overcome by six min, suggesting that CDPK1 myristoylation
is important for ionophore-induced egress, but not essential.
221
Finally, we wanted to examine whether myristoylation of CDPK1 affects its interactions with other
proteins, which may occur by direct binding to the kinase or by indirect association with a shared
membrane structure. We performed immunoprecipitation mass spectrometry (IP-MS) on HA-
tagged CDPK1 from cWT, cMut, and untagged TIR1 parasites after hypotonic lysis. CDPK1 was
the most significantly enriched protein across both cWT and cMut pulldowns when compared
to untagged controls (**Figure 2I**). Two additional proteins were significantly enriched along with
CDPK1: NUP134 (TGGT1_240510) and SRS36D (TGGT1_292280). Comparing the cWT and
cMut pulldowns, we found two proteins preferentially associated with cWT: NUDIX hydroxylase
(TGGT1_227450) and a 14-3-3 protein (TGGT1_269960) (**Figure 2J**). Loss of myristoylation
appears to enhance interactions with three different proteins: ROM4 (TGGT1_268590), a putative
T complex protein 1 alpha subunit (TGGT1_229990), and IMP dehydrogenase (TGGT1_233110).
Genome-wide knockout screen data suggest that NUP134 and the putative T complex protein
1 are the only putative interacting partners required for parasite fitness (Sidik et al., 2016b).
NUP134 binds and co-localizes with the nuclear pore complex, whereas the putative T complex
protein 1 may function as an ATP-dependent chaperone (Courjol et al., 2017). CDPK1 is localized
primarily to the cytoplasm of the parasite and partially in the nucleus, which could explain its
interactions with NUP134 (Lourido et al., 2010; Ojo et al., 2010). ROM4 is an integral membrane
protease required to shed secreted microneme proteins through proteolysis, and its knockdown
impaired gliding motility and invasion of host cells (Buguliskis et al., 2010). ROM4 localizes to
the plasma membrane of parasites but was enriched with cytosolic CDPK1 and was not
observed in our sub-minute resolution phosphoproteomics. Our pulldowns did not reveal a
membrane compartment association for myristoylated CDPK1 but did identify interacting
proteins enriched in cytosolic and myristoylated CDPK1. Given the annotated functions of these
interacting proteins, the factors are unlikely to participate in the motility-related functions of

CDPK1. Additional studies will be required to understand how myristoylation influences CDPK1
activity.
248
Identification of direct CDPK1 targets through thiophosphorylation

Our phosphoproteomic time course experiment identified proteins exhibiting CDPK1-dependent
phosphorylation in live parasites, which includes direct and indirect substrates of the kinase.
CDPK1 is unusual among apicomplexan and metazoan kinases in that it contains a glycine at its
gatekeeper residue, resulting in an expanded ATP-binding pocket that can accommodate bulky
ATP analogues. CDPK1 can use bulky analogues like N6-furfuryladenosine (kinetin)-5′-O-[3-
thiotriphosphate] (KTPγS) to thiophosphorylate its substrates in parasite lysates (Lourido et al.,
2013a). Thiophosphorylated substrates can subsequently be enriched with an iodoacetyl resin
(Blethrow et al., 2008). Specificity can be assessed by comparison to mutant parasites harboring
a G128M gatekeeper mutation in CDPK1 (CDPK1^M) that retains kinase activity but prevents it
from using KTPγS. Previous studies using lysates identified a small set of six putative CDPK1-
dependent substrates; however, these studies lost the target specificity conferred by the
subcellular context and lacked the accuracy and sensitivity of quantitative proteomic
approaches (Lourido et al., 2013a).
263
We implemented several modifications to the thiophosphorylation procedure that improved
identification of CDPK1 targets (**Figure 3A**)(Rothenberg et al., 2016). First, we used stable
isotope labeling of amino acids in cell culture (SILAC) to directly compare thiophosphorylation in
WT (CDPK1^G) and mutant (CDPK1^M) parasites. Second, we used parasites semi-permeabilized
with the bacterial toxin aerolysin—as opposed to the preparations of detergent-lysed parasites
used in prior methods. Aerolysin forms 3-nm pores in the plasma membrane which permit the
diffusion of small molecules but not proteins, enabling us to perform labeling reactions without
drastically disrupting the concentration or localization of proteins (Iacovache et al., 2006). Lastly,
we prevented non-specific extracellular substrate labeling due to premature lysis by treating
semi-permeabilized parasites with 1B7, a nanobody that allosterically inhibits CDPK1 but does
not enter the cytosol of semi-permeabilized parasites (**Figure 3B**)(Ingram et al., 2015).
275
Thiophosphorylation experiments were performed in biological triplicate comparing CDPK1^G and
CDPK1^M parasites. We assessed thiophosphorylation labeling by immunoblot, identifying an
array of proteins specifically labeled in CDPK1^G, but not CDPK1^M parasites (**Figure 3C**). MS
analyses quantified the abundance of peptides in CDPK1^G relative to CDPK1^M parasites,
identifying 734 unique peptides across three biological replicates (**Figure 3D**). Samples from
CDPK1^G parasites were significantly enriched in phosphorylated peptides, consistent with
CDPK1-mediated thiophosphorylation of targets (**Figure 3E**). 123 peptides across 104 proteins
were likely direct substrates of CDPK1. GO enrichment analysis did not reveal any pathways
relevant to the function of CDPK1 in exocytosis (**Figure 3F**). While our approach largely maintains
kinases in their subcellular context, aerolysin treatment disrupts native ion concentrations and
detaches the plasma membrane from the inner membrane complex (IMC) (Wichroski et al., 2002).

We therefore proceeded to consider the thiophosphorylated substrates in the context of the
time-resolved phosphoproteomics.
289
CDPK1 targets participate in pathways controlling parasite motility

Considering the thiophosphorylation and time-resolved phosphoproteomics, we arrived at a
prioritization scheme to identify 163 proteins phosphorylated in a CDPK1-dependent manner
(**Figure 3G**). Proteins were divided into five classes based on the overlap of phosphorylated sites
between both approaches. Class 1 contains five proteins for which the same phosphorylated
site was identified in both the time course and thiophosphorylation experiments. Class 2 contains
four proteins for which phosphorylated sites identified across both approaches were within 50
amino acid residues of one another. Class 3 contains two proteins that were enriched by
thiophosphorylation and were also CDPK1-dependent in the time course, but the identified sites
were more than 50 residues apart. Class 4 contains 93 proteins that were exclusively enriched
by thiophosphorylation. Lastly, Class 5 contains 59 proteins that were CDPK1-dependent
exclusively in the time course and are likely indirect targets of the kinase. Proteins within each
class were further stratified by fitness scores reported from genome-wide knockout screens,
with lower scores representing genes more essential for parasite fitness in cell culture (Sidik et
al., 2016b). Of the 163 targets of CDPK1, 72 proteins across all classes also displayed changes
in zaprinast-dependent phosphorylation at distinct sites that were independent of CDPK1. This
overlap suggests that some proteins modified by CDPK1 are also regulated by additional kinases
and phosphatases.
308
We hypothesized that regulators of exocytosis would be found among the targets of CDPK1. Of
the 163 protein targets, 38 have been previously localized and/or functionally characterized
(indicated with asterisks; **Figure 3G**). Of these 38 proteins, 13 have been implicated in regulating
motile stages of the parasite (**Figure 3H, Figure 3—figure supplement 1**). Among the
candidates associated with microneme exocytosis, centrin 2 (CEN2) and RNG2 have previously
been localized to tubulin-based structures in the apical complex of the parasite, and their
knockdown is sufficient to inhibit secretion of microneme proteins and block host cell invasion
(Katris et al., 2014; Lentini et al., 2019; Leung et al., 2019). Candidates associated with rhoptry
exocytosis include the armadillo repeats only protein (ARO), ARO-interacting protein (AIP),
palmitoyl acyltransferase DHHC7, and patatin-like phospholipase (PL3) (Beck et al., 2013;
Mueller et al., 2016, 2013; Wilson et al., 2020). ARO and DHHC7 both localize to the cytosolic
face of rhoptries, where they influence the recruitment of AIP and ACβ. Knockdown of either
DHHC7 or ARO disrupts the localization of mature rhoptries, inhibiting the secretion of rhoptry
proteins required for invasion. Another potential CDPK1 target, the apical complex lysine
methyltransferase (AKMT), rapidly relocalizes from the apical complex to the parasite body
during Ca^{2+}-regulated motility, and its knockdown impairs the gliding motility of parasites
required for invasion and egress (Heaslip et al., 2011). Other putative CDPK1 targets participate
in homeostasis and biogenesis of secretory organelles, such as the vacuolar type Na^+/H^+
exchanger (NHE3), dynamin-related protein B (DrpB), V-ATPase a1 (VHA1), and GDP-fucose
transporter (NST2). NHE3-knockout parasites exhibit sensitivity to osmotic shock and

dysregulated cytosolic Ca^{2+}, resulting in reduced secretion of microneme proteins and an
inhibition of invasion (Francia et al., 2011). DrpB was previously identified as a CDPK1 target,
and its depletion results in severe defects in the biogenesis of micronemes and rhoptries
(Breinich et al., 2009; Lourido et al., 2013a). VHA1 and NST2 participate in the maturation of
microneme and rhoptry proteins (Bandini et al., 2019; Stasic et al., 2019). Lastly, signaling
proteins regulating intracellular cGMP levels, such as the guanylate cyclase (GC) and the unique
GC organizer (UGO), were also phosphorylated in a CDPK1-dependent manner (Bisio et al.,
2019; Brown and Sibley, 2018; Yang et al., 2019). CDPK1 acts downstream of GC-mediated
production of cGMP, which may suggest regulatory feedback on Ca^{2+} release, as has been
suggested for CDPK3 (Nofal et al., 2022). These data demonstrate that identification of CDPK1
targets can uncover proteins involved in Ca^{2+}-regulated exocytosis.
340
To identify new factors involved in Ca^{2+}-regulated secretion, we prioritized Class 1 and Class 2
candidates. Seven candidates in this category lack functional annotation or have been
associated with apical structures: TGGT1_227610, TGGT1_221470, TGGT1_235160,
TGGT1_254870, KinesinB (TGGT1_273560), a small nuclease (TGGT1_310060), and
TGGT1_289100 (MIC18). TGGT1_221470 was previously identified as a CDPK1 substrate after
a pull down from parasite lysates but remained functionally uncharacterized (Lourido et al.,
2013a). KinesinB localizes to cortical microtubules, TGGT1_254870 localizes to the apical
complex, and TGGT1_289100 co-localizes with micronemes, each of which are parasite
structures relevant to exocytosis (Butler et al., 2014; Leung et al., 2017; Long et al., 2017). Of
these candidates, only TGGT1_227610 and TGGT1_289100 appear to be required for parasite
fitness (Long et al., 2017; Sidik et al., 2016b). Despite prior annotation as a microneme protein,
TGGT1_289100 lacks a signal peptide and is localized to the cytosol by spatial proteomics
(Barylyuk et al., 2020; Butler et al., 2014). Instead, TGGT1_289100 is predicted to have an N-
terminal Hook domain and extensive coiled-coil domains (Simm et al., 2015; Söding et al., 2005)
with homology to activating adaptors that bind to endosomes and activate super-processive
dynein-mediated trafficking towards the minus end of microtubules (**Figure 4A**) (Bielska et al.,
2014; Guo et al., 2016; Zhang et al., 2014). The predicted polarity of cortical microtubules and
their association with micronemes support a model where the vesicles are trafficked by dyneins
towards the apical end of the parasite (Chen et al., 2015; Leung et al., 2017; Wang et al., 2021).
We observed extensive CDPK1-dependent phosphorylation between the Hook and coiled-coil
domains of TGGT1_289100 (S167, S177, and S189–191), consistent with potential regulation of
the putative adaptor. These data motivated functional characterization of this factor—henceforth
referred to as HOOK.
364
A Hook domain protein phosphorylated by CDPK1 regulates parasite invasion

To study the role of HOOK during the acute stages of the parasite, we generated a conditional
knockdown by fusing an HA-mAID tag to its N terminus (AID-HOOK; **Figure 4—Figure
supplement 1**). HOOK was depleted from parasites following 24–40 hrs of auxin treatment, as
determined by immunoblotting or immunofluorescence microscopy (**Figure 4B–C**). In contrast
to previous observations that localized HOOK to micronemes, we observed only partial co-

localization with micronemes with a majority of HOOK localized to the cytosol (**Figure 4C**).
Plaque formation was impaired when AID-HOOK parasites were grown in the presence of auxin,
consistent with the strong effect on parasite fitness reported from genome-wide knockout
screens (**Figure 4D**)(Sidik et al., 2016b). These results indicate that HOOK is required during the
acute stages of *T. gondii*. Parasite replication was not affected following 24 hrs of auxin
treatment (**Figure 4E**). Microneme and rhoptry biogenesis were also unaffected following 24 hrs
of auxin treatment immunofluorescence analysis (**Figure 4C**, **Figure 4—figure supplement 1**).
These data suggest HOOK functions during stages associated with parasite motility, such as
parasite invasion and egress. Parasites depleted of HOOK displayed reduced invasion efficiency,
consistent with that of CDPK1-depleted parasites (**Figure 4F**). We noted that even in the absence
of auxin treatment AID-HOOK parasites required longer incubations to reach comparable levels
of invasion to wildtype, suggesting that manipulation of the N terminus partially affects HOOK
function.
384
HOOK depletion only partially recapitulates the effects of CDPK1 loss. Parasites depleted of
CDPK1 were unable to egress—as documented previously (Lourido et al., 2012, 2010)—whereas
depletion of HOOK had no effect on parasite egress (**Figure 4G**). Finally, we also examined
whether the stability of CDPK1 affected HOOK expression by tagging the C terminus of HOOK
with a 3xHA tag in CDPK1 cKD parasites (HOOK-3xHA). HOOK localization and abundance was
unaffected by depleting parasites of CDPK1 for 24 hrs as determined by immunofluorescence
(**Figure 4H**). Taken together, our results indicate that HOOK is required for invasion of host cells,
but dispensable for egress.
393

HOOK forms a complex that regulates parasite invasion

Opisthokont HOOK proteins function in complexes to activate dynein-mediated trafficking of
vesicular cargo along microtubules (Bielska et al., 2014; Christensen et al., 2021; Gillingham et
al., 2014; Guo et al., 2016; Xu et al., 2008; Yao et al., 2014). In *D. melanogaster* and mammals,
HOOK proteins have been shown to form dimers and bind FTS and FHIP via a C-terminal region
that interacts with vesicular cargo (Christensen et al., 2021; Krämer and Phistry, 1996; Lee et al.,
2018; Xu et al., 2008). To identify analogous binding partners of HOOK in *T. gondii*, we performed
IP-MS on lysates from HOOK-3xHA parasites and an untagged control (**Figure 5A**). Three
proteins showed comparable enrichment to HOOK-3xHA: TGGT1_264050, TGGT1_316650, and
TGGT1_306920. Like FTS, TGGT1_264050 bears homology to E2 ubiquitin-conjugating
enzymes, but lacks the catalytic cysteine required for enzymatic activity at position 162. Because
of these shared features and their homology, we designated TGGT1_264050 as FTS.
406

To confirm that FTS forms a complex with HOOK, we investigated its localization and performed
reciprocal IPs. We fused a 3xHA tag to the C terminus of FTS and observed a similar localization
to HOOK by immunofluorescence: primarily cytosolic with partial overlap with micronemes
(**Figure 5B**). To identify binding partners of FTS, we performed IP-MS on FTS-3xHA parasite
lysates and compared protein enrichment to the HOOK-3xHA IP (**Figure 5C**). Four proteins were

significantly enriched in the reciprocal IP experiments—HOOK, FTS, TGGT1_316650, and
TGGT1_306920—confirming the four-member complex.
414
To determine whether FTS functions in the same pathway as HOOK, we generated a conditional
knockdown by fusing an AID-HA tag to the C terminus of FTS (FTS-AID; **Figure 5—Figure**
supplement 1). FTS was readily depleted from parasites following 40 hrs of auxin treatment
(**Figure 5D**). FTS depletion completely blocked plaque formation (**Figure 5E**). Similar to HOOK,
depletion of FTS had no observable effect on parasite replication or microneme biogenesis
(**Figure 5—figure supplement 1**, **Figure 5F**). During the motile stages of the parasite, FTS
depletion resulted in a block in invasion efficiency but had no effect on egress, phenocopying
the effects of HOOK depletion (**Figure 5G**, **Figure 5H**). FTS appeared to tolerate tagging better
than HOOK, since the FTS-AID parasites behaved normally in the absence of auxin. These data
suggest that HOOK and FTS form a functional complex required for parasite invasion.
425
We pursued enzyme-catalyzed proximity labeling to complement IP-MS studies and capture
transient protein interactions (Branon et al., 2018). In this approach, cells express a protein of
interest tagged with the promiscuous biotin ligase TurboID. Addition of a biotin substrate to live
cells results in biotinylation within a few nanometers of the TurboID-tagged protein, in a matter
of minutes. Biotinylated proteins can be enriched from cellular lysates using streptavidin-affinity
purification and identified using MS. Proximity labeling was recently used to characterize the
cargo diversity of distinct human FHF complexes in human cells (Christensen et al., 2021). To
identify additional proteins that interact with the *T. gondii* HOOK complex, we performed
proximity labeling after fusing TurboID to the C terminus of FTS, which was more amenable to
tagging compared to HOOK (**Figure 5I**, **Figure 5—figure supplement 1**). As a control, we used
parasites expressing a cytosolic mNeonGreen-TurboID that would broadly label cytosolic
proteins and identify non-specific interactions (**Figure 5—figure supplement 1**).
438
Of the 14 proteins significantly enriched in FTS-TurboID parasites, members of the HOOK
complex (HOOK, FTS, TGGT1_306920, and TGGT1_316650) were the top four most enriched
proteins. In humans, proximity labeling studies suggest FTS and FHIP bind at the C terminus of
the HOOK dimer to mediate FHIP binding to RAB5 endosomes (Christensen et al., 2021). Our
FTS proximity labeling results suggest that TGGT1_306920 and TGGT1_316650 may also bind
in this manner. Among the other significantly-enriched proteins, PC and ACC1 encode
carboxylases known to be covalently modified by biotin and likely represent non-specific
enrichment. DHFR enrichment is likely due to its use as a selectable marker when generating the
FTS-TurboID strain. TGGT1_294610 and TGGT1_280770 are a putative histone lysine
methyltransferase and regulator of chromosome condensation (RCC1)-repeat containing
protein, respectively, and have no characterized functions. Of the remaining enriched proteins
that lack annotations, TGGT1_221180 was previously localized to micronemes by spatial
proteomics and contains a transmembrane domain, suggesting a molecular link between the
HOOK complex and micronemes (Barylyuk et al., 2020).
453

The HOOK complex promotes invasion by regulating microneme exocytosis

To determine whether the HOOK complex is required for microneme exocytosis, we directly measured secretion of the microneme protein MIC2. Microneme exocytosis exposes integral membrane proteins like MIC2 that function as adhesins during gliding motility. These adhesins subsequently undergo proteolytic cleavage and are released into the supernatant, which can be analyzed by immunoblot (Carruthers et al., 2000). Quantification of the proportion of secreted MIC2 was performed by generating a standard curve from unstimulated lysates. Strikingly, parasites depleted of either member of the HOOK complex were severely impaired in their ability to secrete microneme proteins (**Figure 5J**). Inhibition of the complex led to a similar degree of secretion impairment as CDPK1 depletion. Together, these data suggest that HOOK and FTS form a functional complex that is required for microneme exocytosis.

Microneme trafficking depends on CDPK1 activity and HOOK

We hypothesized that if the HOOK complex is required for sustained microneme exocytosis, then depletion of HOOK would inhibit trafficking of micronemes during parasite motility. Dynamic relocalization of micronemes can be visualized using live microscopy of parasites expressing the integral microneme protein CLAMP fused to mNeonGreen (CLAMP-mNG) (**Figure 6A**)(Sidik et al., 2016b). To trigger microneme exocytosis, parasites are treated with zaprinast to stimulate a rise in intracellular Ca^{2+} and activate CDPK1. Dynamic relocalization of micronemes is defined as the CLAMP-mNG signal that concentrates at the apical end of the parasite over time. To quantify microneme relocalization, mNG signal is measured, over time, along the apical-basal axis of individual parasites within a vacuole. We scored microneme relocalization by calculating the difference in maximum apical intensity comparing the instant of zaprinast addition to the time point just prior to parasite egress of the uninhibited zaprinast-stimulated controls (**Figure 6A**).

We first assessed whether CDPK1 kinase activity is required for microneme relocalization using the specific inhibitor 3-MB-PP1 (Lourido et al., 2010). We incubated parasites with 3-MB-PP1 for 30 min, imaged for 1 min to establish a baseline, and then stimulated with zaprinast. Parasites pre-treated with DMSO relocalized micronemes as expected; however, microneme relocalization was blocked when CDPK1 was inhibited by 3-MB-PP1 (**Figure 6B-D**). These results indicate microneme relocalization depends on CDPK1 kinase activity.

We next assessed whether HOOK is also required for microneme relocalization. We fused an mNG reporter to the C terminus of CLAMP in AID-HOOK parasites and TIR1 parasites as a control (**Figure 6—Figure supplement 1**). We treated parasites with auxin for 24 hrs prior to stimulation with zaprinast. Microneme relocalization was unaffected in TIR1/CLAMP-mNG parasites treated with auxin (**Figure 6E–F**). By contrast, microneme relocalization did not occur in intracellular AID-HOOK parasites regardless of auxin treatment (**Figure 6G–H**). We attributed the lack of relocalization in vehicle-treated parasites to the hypomorphism resulting from N-terminal tagging, which is consistent with the delayed invasion and reduced plaque size documented for this strain. To determine whether vehicle-treated AID-HOOK parasites relocalized micronemes over longer periods, we examined CLAMP-mNG localization in

extracellular parasites after egress. While vehicle-treated AID-HOOK parasites were able to
relocalize micronemes, parasites depleted of HOOK exhibited a striking microneme localization
dispersed throughout the cytosol (**Figure 6I–J**). As expected, TIR1/CLAMP-mNG parasites
maintained the apical relocalization of micronemes regardless of auxin treatment in extracellular
parasites. We used an alternative metric for microneme relocalization to account for the
disorganized localization of micronemes in parasites depleted of HOOK. We quantified the
percent of the total CLAMP-mNG signal localized to the apical end versus the remaining body
of the parasite (**Figure 6K**). The apical region cutoffs were experimentally derived from vehicle-
treated TIR1/CLAMP-mNG parasites and defined as the apical 12.5% of the parasite. While a
majority of CLAMP-mNG signal was localized to the apical region of TIR1/CLAMP-mNG
parasites, most of the CLAMP-mNG signal observed in parasites depleted of HOOK was found
in the parasite body. Micronemes rapidly adopt an aberrant localization in the absence of HOOK
during parasite motility. These results indicate that CDPK1 kinase activity and HOOK are
required for microneme trafficking during parasite motility.
510
DISCUSSION

In this study, we sought to identify new regulators of Ca^{2+}-mediated exocytosis in *T. gondii* by
studying the targets of a key regulator, the kinase CDPK1. We identified 163 proteins
phosphorylated in a CDPK1-dependent manner using sub-minute resolution
phosphoproteomics and thiophosphorylation for direct substrate capture. We determined that
myristoylation of CDPK1 contributes to the kinase's function during the lytic cycle. 13 of the
identified CDPK1 targets have previously described functions in parasite motility, revealing
possible points of regulation within relevant signaling pathways. Furthermore, we characterized
a new regulator of microneme exocytosis called HOOK, which forms a stable complex with FTS
and two other proteins. Homologs of HOOK and FTS participate in dynein-mediated vesicular
trafficking in other organisms. In *T. gondii*, knockdown of HOOK or FTS blocked invasion of host
cells and altered rapid microneme trafficking during Ca^{2+}-regulated motility. Overall, we show
how studying parasite signaling pathways can illuminate the cellular adaptations that support
parasitism.

Over the past decade, several efforts have sought to characterize phosphorylation within
apicomplexans. Improvements in phosphopeptide enrichment and mass spectrometry have
enabled global characterization of the phosphoproteomes from *T. gondii* tachyzoites and
P. falciparum schizonts, trophozoites, and ring-stages(Lasonder et al., 2012; Pease et al., 2013;
Solyakov et al., 2011; Treeck et al., 2011). Coupling such approaches with genetic and
pharmacological tools has enabled the deconvolution of kinase-specific effects from the vast
global phosphorylation program(Blomqvist et al., 2020; Fang et al., 2017; Invergo et al., 2017;
Nofal et al., 2022; Pease et al., 2018; Treeck et al., 2014), revealing the activation of MyoA-
mediated gliding by *Tg*CDPK3, the roles of *Pb*CDPK4 and *Pb*SRPK1 during male
gametogenesis, and the erythrocytic-stage effects of *Pf*CDPK1, *Pf*PK7, and *Pf*CDPK5(Blomqvist
et al., 2020; Fang et al., 2017; Gaji et al., 2015; Invergo et al., 2017; Pease et al., 2018; Tang et
al., 2014). Attributing individual phosphorylation events to specific kinases can be complicated

by cumulative changes or adaptation of the signaling networks in knockouts. Chemical-genetic
approaches can more precisely inhibit the activity of a given kinase at the time of the assay,
while controlling for off-target effects, and have been used to characterize *Pf*PKG-dependent
phosphorylation (Alam et al., 2015; Brochet et al., 2014). Kinetically-resolved
phosphoproteomics has enabled a more nuanced understanding of signaling cascades during
parasite motility(Herneisen et al., 2022; Invergo et al., 2017; Nofal et al., 2022). While such
technical advances have led to more in-depth phosphoproteome mapping, identifying the direct
targets of a given kinase remains a key challenge to structuring the observed changes into
signaling pathways.
547
To resolve direct and indirect effects on the phosphoproteome for *T. gondii* CDPK1, we
combined two complementary approaches: temporally-resolved phosphoproteomics and bio-
orthogonal labeling of direct substrates. We implemented several methodological advances.
Rapid conditional knockdown using the auxin-inducible degron system resulted in rapid and
specific depletion of the kinase of interest for the phosphoproteomic studies(Brown et al., 2018).
Additionally, quantification and coverage by mass spectrometry were improved through TMTpro
multiplexing and ion mobility spectrometry (FAIMS)(Bekker-Jensen et al., 2020; Van Vranken et
al., n.d.). We also improved on the low coverage observed in previous thiophosphorylation
experiments with parasite lysates(Fang et al., 2017; Lourido et al., 2013a) by maintaining a more
native signaling environment using aerolysin semi-permeabilization, SILAC-based peptide
quantification, and shortened labeling times(Rothenberg et al., 2016). These approaches
characterize proteome-wide phosphorylation kinetics in live parasites and reveal direct kinase-
substrate relationships for CDPK1.
561
Myristoylation of CDPK1 contributes to its function. Contrary to previous reports describing the
kinase as cytosolic or nuclear(Ojo et al., 2010; Pomel et al., 2008), we demonstrate that
myristoylated CDPK1 is at least partially associated with structures that can be fractionated from
the cytoplasm by differential centrifugation. Loss of myristoylation led to CDPK1 release from
the insoluble fraction. The ortholog of CDPK1 in *P. berghei*, *Pb*CDPK4, also displays
myristoylation-specific functions during male gametogenesis: myristoylated *Pb*CDPK4 is critical
for the first genome replication, whereas the non-myristoylated *Pb*CDPK4 is important for the
completion of gametogenesis(Fang et al., 2017). Myristoylation could impact CDPK1's ability to
access certain targets efficiently. However, such interactions appear to be too weak or transient
to be captured by immunoprecipitation. Mutating the myristoylation site of CDPK1 only modestly
affected ionophore-induced egress. However, minor effects in these key transitions may be
magnified over repeated lytic cycles, which may explain the more substantial impact of losing
CDPK1 myristoylation on plaque formation. Alternatively, other kinases may compensate for
decreased CDPK1 activity, particularly under hyperactivated conditions like stimulated egress.
A plausible candidate for compensation is CDPK2A, which also appears to be myristoylated,
despite lacking an N-terminal MG motif, and was recently shown to display epistatic interactions
with CDPK1(Broncel et al., 2020; Shortt et al., 2022).
579

CDPK1 plays a critical role in the transition from the replicative intracellular stages to motile
extracellular stages (Lourido et al., 2010; Shortt et al., 2022). During this transition parasites must
execute rapid cellular changes that involve the exocytosis of micronemes and rhoptries,
reorganization of the cytoskeleton, gliding motility, and maintenance of ion homeostasis(Blader
et al., 2015). CDPK1 has also been suggested to control the actomyosin system and extrusion
of the conoid(Tosetti et al., 2019). Our detailed target analysis suggests CDPK1 regulates—and
possibly helps coordinate—multiple pathways. We revealed over a hundred CDPK1 targets with
diverse predicted functions, including many relevant to phenotypes dependent on CDPK1. In
addition to the effects on microneme trafficking examined in this study, our results point to a
direct link between CDPK1 and rhoptry exocytosis. Ca^{2+} has been implicated in rhoptry
discharge, as the rhoptry-localized Ca^{2+}-binding FER2 is required for rhoptry
exocytosis(Coleman et al., 2018). We observed a preponderance of CDPK1-dependent
phosphorylation on proteins regulating rhoptry exocytosis, including DHHC7, ARO, and AIP.
Phosphorylation on ARO was observed near the N-terminal acylation sites required for rhoptry
targeting and may regulate the bundling and positioning of mature rhoptries during motile
stages(Mueller et al., 2016). However, formally demonstrating the relationship between
Ca^{2+}/CDPK1 and rhoptry discharge is complicated by the dependency of the latter on the
secretion of certain microneme proteins (Ben Chaabene et al., 2021; Carruthers and Sibley,
1997). *T. gondii* tachyzoites have several rhoptries, yet only two are docked for exocytosis at a
given time (Aquilini et al., 2021; Mageswaran et al., 2021; Segev-Zarko et al., 2022). Regulating
the activity of ARO during motile stages may influence the ability to mobilize and re-dock
rhoptries in preparation for invasion. Considering how different CDPK1 substrates function, we
expect the various phenotypes associated with CDPK1 will depend on distinct sets of
substrates.
604
Although CDPK1 is known to mediate microneme exocytosis (Lourido et al., 2010), the precise
molecular events controlling this process have remained elusive. In this study, we determined
that CDPK1 activity is required for the Ca^{2+}-stimulated trafficking of micronemes to the apical
end. This phenotype depends on HOOK, one of the CDPK1 targets uncovered by both global
phosphoproteomics and thiophosphorylation studies. Despite prior annotation as a microneme
protein(Butler et al., 2014), HOOK likely functions as an activating adaptor in *T. gondii* based on
homology to other Hook proteins and the functional characterization we have performed. In
particular, immunoprecipitation of HOOK identified three interacting proteins: FTS,
TGGT_316650, and TGGT1_306920. In *H. sapiens*, *A. nidulans*, and *D. melanogaster*, Hook
proteins complex with FTS and FHIP (the FHF complex), to link cargo to the dynein machinery
for trafficking along microtubules(Bielska et al., 2014; Gillingham et al., 2014; Guo et al., 2016;
Xu et al., 2008; Yao et al., 2014). FHF complexes have been shown to recognize early endosomes
via FHIP binding to RAB5(Bielska et al., 2014; Guo et al., 2016; Zhang et al., 2014). Homology to
FHIP was lacking in TGGT1_316650 and TGGT1_306920. However, genome-wide knockout
screens indicate TGGT1_306920 is critical for parasite fitness, leading us to speculate that it may
mediate cargo binding in a manner structurally distinct from known adaptors. Additional work
will be needed to elucidate the function of the HOOK complex and to understand how it mediates
trafficking of apicomplexan-specific organelles.

Our results place the HOOK complex downstream of CDPK1, regulating the stimulated
relocalization of micronemes to the apical end of parasites. However, the HOOK complex only
partially explains CDPK1's role during egress. Whereas CDPK1 depletion blocks egress entirely,
parasites depleted of HOOK or FTS exit host cells, despite subsequent defects in microneme
secretion and invasion. Microneme discharge is required for egress, supplying adhesins for
gliding motility and the perforin PLP1 to rupture the parasitophorous vacuole membrane(Kafsack
et al., 2009). It therefore appears that the initial round of microneme discharge during egress
depends on CDPK1, and only subsequent rounds require the HOOK complex. Indeed, a fraction
of micronemes are already found docked at the apical complex prior to the transition from the
replicative to the motile stages, and may constitute the first round of microneme
exocytosis(Mageswaran et al., 2021; Sun et al., 2022). Dynein-mediated trafficking might
become important immediately following egress when continuous microneme discharge is
necessary, as is the case during gliding motility and invasion. Consistent with this hypothesis,
depletion of the dynein light chain 8a (DLC8a) yielded similar phenotypes to HOOK and FTS
knockdown: intact egress but dysfunctional microneme protein secretion, gliding motility, and
invasion(Lentini et al., 2019). Loss of DLC8a also led to defects in rhoptry positioning, which
were not observed in HOOK knockdowns(Lentini et al., 2019), so other activating adaptors may
be involved in rhoptry trafficking. Our results suggest that the HOOK complex has been
specifically adapted for microneme trafficking in *T. gondii*.

Several aspects of microneme trafficking remain to be determined. The precise nature of the
dynein complex is poorly understood. Eukaryotes typically express a single cytoplasmic dynein
heavy chain. In *T. gondii*, TGGT1_294550 is the top candidate for the cytoplasmic dynein heavy
chain (DHC) as it contains the necessary domains, is conserved among apicomplexans, and is
required for parasite fitness (Lentini et al., 2019; Sidik et al., 2016b). Other putative DHCs in the
T. gondii genome are likely axonemal dyneins required for flagellar function in sexual-stage
parasites. It also remains unknown how the HOOK complex binds to micronemes. In *H. sapiens*
and *D. melanogaster*, RAB5 on vesicles interacts with FHIP in the HOOK complex(Bielska et al.,
2014; Gillingham et al., 2014; Guo et al., 2016; Xu et al., 2008; Yao et al., 2014). We speculate
that TGGT1_306920 may serve the role of FHIP within the HOOK complex but its binding partner
on micronemes remains unknown. RAB5A and RAB5C have been implicated in the biogenesis
of micronemes, but their roles during exocytosis have not been explored(Kremer et al., 2013).
Understanding how micronemes are recognized may elucidate how cargo specificity is achieved
and regulated.

Ca^{2+} signaling has been tuned to support apicomplexan-specific cellular processes but
identifying and integrating effectors within the pathway has remained a major bottleneck.
Signaling effectors conserved amongst eukaryotes are often integrated into apicomplexan
cellular pathways in novel or unusual ways. Here, we identified candidate downstream effectors
of the Ca^{2+}-regulated kinase CDPK1 by monitoring proteome-wide protein phosphorylation. In
addition to creating a catalog of candidates for future characterization, we identified,
characterized, and integrated the HOOK complex downstream of CDPK1 within the Ca^{2+}

signaling network controlling microneme exocytosis. Lastly, our results are the first to implicate

activating adaptors as critical factors for the pathogenesis of an apicomplexan organism by

promoting exocytic trafficking.

MATERIALS & METHODS

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background (*T. gondii*)	TIR1	PMID: 28465425		RH/TIR1/ΔKU80/ΔHXGPRT
Strain, strain background (*T. gondii*)	CDPK1-AID	doi: https://doi.org/10.1101/2022.07.19.500742		RH/TIR1/ΔKU80/ΔHXGPRT/CDPK1-V5-mNeonGreen-mAID-Ty
Strain, strain background (*T. gondii*)	iKD CDPK1	This paper	TGGT1_301440	RH/TIR1/ΔKU80/ΔHXGPRT/CDPK1-mAID-Myc/HXGPRT
Strain, strain background (*T. gondii*)	cWT CDPK1	This paper	TGGT1_301440	RH/TIR1/ΔKU80/ΔHXGPRT/CDPK1-mAID-Myc/HXGPRT/CDPK1-HA-T2A-GFP
Strain, strain background (*T. gondii*)	cMut CDPK1	This paper	TGGT1_301440	RH/TIR1/ΔKU80/ΔHXGPRT/CDPK1-mAID-Myc/HXGPRT/CDPK1(G2A)-HA-T2A-mCherry
Strain, strain background (*T. gondii*)	Δku80Δhxgprt YFP			RH/ΔKU80/ΔHXGPRT/YFP
Strain, strain background (*T. gondii*)	CDPK1G	PMID: 19218426		RH/ΔKU80/ΔHXGPRT
Strain, strain background (*T. gondii*)	CDPK1M	PMID: 23149386	TGGT1_301440	RH/ΔKU80/ΔHXGPRT/CDPK1(G128M)-Myc

Strain, strain background (*T. gondii*)	AID-HOOK	This paper	TGGT1_289100	RH/TIR1/ΔKU80/ΔHXGPRT/HA-mAID-HOOK
Strain, strain background (*T. gondii*)	CDPK1-AID/HOOK-3xHA	This paper	TGGT1_289100	RH/TIR1/ΔKU80/ΔHXGPRT/CDPK1-V5-mNeonGreen-mAID-Ty/HOOK-V5-3HA/HXGPRT
Strain, strain background (*T. gondii*)	FTS-AID	This paper	TGGT1_264050	RH/TIR1/ΔKU80/ΔHXGPRT/FTS-V5-mAID-HA/HXGPRT
Strain, strain background (*T. gondii*)	FTS-3xHA	This paper	TGGT1_264050	RH/TIR1/ΔKU80/ΔHXGPRT/FTS-V5-3HA/HXGPRT
Strain, strain background (*T. gondii*)	FTS-TurboID	This paper	TGGT1_264050	RH/TIR1/ΔKU80/ΔHXGPRT/FTS-TurboID-Ty/DHFR
Strain, strain background (*T. gondii*)	cytosolic mNG-TurboID	This paper		RH/TIR1/ΔKU80/ΔHXGPRT/pMIC2-mNeonGreen-TurboID-Ty
Strain, strain background (*T. gondii*)	CLAMP-mNG	PMID: 27594426	TGGT1_265790	RH/ΔKU80/ΔHXGPRT/CLAMP-mNeonGreen
Strain, strain background (*T. gondii*)	TIR1/CLAMP-mNG	This paper	TGGT1_265790	RH/TIR1/ΔKU80/ΔHXGPRT/CLAMP-mNeonGreen
Strain, strain background (*T. gondii*)	TIR1/CLAMP-mNG/AID-HOOK	This paper	TGGT1_265790	RH/TIR1/ΔKU80/ΔHXGPRT/HA-mAID-HOOK/CLAMP-mNeonGreen
Cell line (*Homo sapiens*)	Human Foreskin Fibroblasts (HFFs)	ATCC	SCRC-1041	
Antibody	Mouse polyclonal anti-CDPK1	other		Provided by Matthew Child and Matt Bogyo. WB

				(1:3000). Only used in Figure 2.
Antibody	Mouse monoclonal clone 4A6 anti-Myc	Millipore	Cat# 05-724, RRID:AB_11211891	WB (1:1000)
Antibody	Rat monoclonal clone 3F10 anti-HA	Roche	Cat# 11867423001, RRID:AB_390919	WB (1:1000); IFA (1:1000). Only used in Figure 2.
Antibody	Mouse monoclonal clone TP3 anti-*Toxoplasma*	Abcam	Cat# ab8313, RRID:AB_306466	WB (1:1000)
Antibody	Rabbit polyclonal anti-GAP45	other		Provided by Peter Bradley Lab. WB (1:1000). Only used in Figure 2.
Antibody	Mouse monoclonal anti-GFP	Roche	Cat# 11814460001, RRID:AB_390913	WB (1:1000)
Antibody	Rabbit polyclonal anti-mCherry	Abcam	Cat# ab167453, RRID:AB_2571870	WB (1:1000)
Antibody	Rabbit monoclonal anti-SAG1	other		Provided by John Boothroyd Lab; WB (1:10,000). Only used in Figure 2.
Antibody	Rabbit monoclonal clone 51-8 anti-thiophosphate ester	Abcam	Cat# ab92570, RRID:AB_10562142	WB (1:5000)

Antibody	Mouse monoclonal clone 12G10 anti-tubulin	Developmental Studies Hybridoma Bank at the University of Iowa	RRID: AB_1157911	Provided by Marc-Jan Gubbels Lab. WB (1:2000)
Antibody	Guinea pig monoclonal anti-CDPK1	Covance	Custom antibody	WB (1/50,000)
Antibody	Rabbit monoclonal (C29F4) anti-HA	Cell Signaling Technology	Cat# 3724, RRID:AB_1549585	WB (1:1000); IFA (1:1600)
Antibody	Mouse monoclonal clone 6D10 anti-MIC2	PMID: 10799515		Provided by L. David Sibley Lab. WB (1:5000); IFA (1:2000)
Antibody	Mouse anti-ROP1	other		Provied by Peter Bradley Lab. IFA (1:2000)
Antibody	Mouse polyclonal anti-SAG1	PMID: 3183382		Provided by L. David Sibley Lab. IFA (1:500). Used for invasion assays.
Antibody	Rabbit polyclonal anti-GAP45	Lampire Biological Laboratory		Provided by R. Drew Etheridge Lab. IFA (1:1000). Used for invasion assays.
Antibody	Mouse monoclonal (16B12) anti-HA	BioLegend	Cat# 901533, RRID: AB_2565005	WB (1:1000)
Antibody	Rabbit polyclonal clone WU1614 anti-ALD	PMID: 16923803		Provided by L. David Sibley Lab. WB (1:10,000)
Antibody	Rabbit polyclonal anti-GAP45	other		Provided by Dominique Soldati-Favre Lab. WB (1:5000); IFA (1:5000)

Antibody	Peroxidase-AffiniPure polyclonal Goat Anti-Rabbit IgG (H+L)	Jackson ImmunoResearch Laboratories	Cat# 111-035-003, RRID:AB_2313567	WB (1/5000)
Antibody	Alexa Fluor 594 anti-rabbit	Life Technologies		IFA (1:1000)
Antibody	Alexa Fluor 488 anti-mouse	Life Technologies		IFA (1:1000)
Antibody	Alexa Fluor 488 anti-rabbit	Life Technologies		IFA (1:1000)
Antibody	Alexa Fluor 594 anti-mouse	Life Technologies		IFA (1:1000)
Antibody	IRDye 800CW Donkey anti-Guinea Pig IgG Secondary Antibody	LICOR	LICOR:926-32411	WB (1:10,000)
Antibody	IRDye 680RD Donkey anti-Guinea Pig IgG Secondary Antibody	LICOR	LICOR:926-68077	WB (1:10,000)
Antibody	IRDye 800CW Goat anti-Mouse IgG1-Specific Secondary Antibody	LICOR	LICOR:926-32350	WB (1:10,000)

Antibody	IRDye 680LT Goat anti-Mouse IgG Secondary Antibody	LICOR	LICOR:926-68020	WB (1:10,000)
Antibody	IRDye 800CW Goat anti-Rabbit IgG Secondary Antibody	LICOR	LICOR:926-32211	WB (1:10,000)
Antibody	IRDye 680LT Goat anti-Rabbit IgG Secondary Antibody	LICOR	LICOR:926-68021	WB (1:10,000)
Antibody	Alpaca clone 1B7 anti-CDPK1	PMID: 26305940	n/a	
Peptide, recombinant protein	IRDye 680RD Streptavidin	LICOR	LICOR:926-68079	WB (1:3,000)
Peptide, recombinant protein	Aerolysin	PMID: 26584919	n/a	
Peptide, recombinant protein	Sequencing Grade Modified Trypsin	Promega	Promega:V5113	
Chemical compound, drug	3-Indoleacetic acid (auxin)	Sigma Aldrich	Sigma Aldrich:I2886-5G	
Chemical compound, drug	Compound 1	PMID: 12455981	n/a	
Chemical compound, drug	Hoechst 33258	Santa Cruz	Santa Cruz:sc-394039	Egress assay (1:4000)
Chemical compound, drug	Hoechst 33342	Invitrogen	Invitrogen: H3570	IFA (1:20,000)

Chemical compound, drug	DAPI (4',6-Diamidino-2-Phenylindole, Dihydrochloride)	Invitrogen	Invitrogen: D1306	
Chemical compound, drug	Prolong Diamond	Thermo Fisher	Thermo Fisher:P36965	
Chemical compound, drug	zaprinast	Calbiochem	Calbiochem: 684500	
Chemical compound, drug	A23187	Calbiochem	Calbiochem: 100105	
Chemical compound, drug	Myristic acid	Tokyo Chemical Industry	Cat# M0476	
Chemical compound, drug	Alkyne-myristic acid (YnMyr)	Iris Biotech	Cat# RL-2055	
Chemical compound, drug	Trypsin cleavable capture reagent	PMID: 25807930; PMID: 32618271		This reagent was first reported as RTB in PMID: 25807930.
Chemical compound, drug	L-Proline for SILAC	Thermo Fisher Scientific	Thermo Fisher Scientific:88211	
Chemical compound, drug	Fetal Bovine Serum, dialyzed, US origin, One Shot™ format	Thermo Fisher Scientific	Thermo Fisher Scientific:A3382001	
Chemical compound, drug	DMEM for SILAC	Thermo Fisher Scientific	Thermo Fisher	

			Scientific:8 8364	
Chemical compound, drug	L-Arginine-HCl for SILAC	Thermo Fisher Scientific	Thermo Fisher Scientific:8 9989	
Chemical compound, drug	L-Lysine-2HCl for SILAC	Thermo Fisher Scientific	Thermo Fisher Scientific:8 9987	
Chemical compound, drug	L-Arginine-HCl, 13C6, 15N4 for SILAC	Thermo Fisher Scientific	Thermo Fisher Scientific:8 9990	
Chemical compound, drug	L-Lysine-2HCl, 13C6, 15N2 for SILAC	Thermo Fisher Scientific	Thermo Fisher Scientific:8 8209	
Chemical compound, drug	p-nitrobenzyl mesylate (PNBM)	Epitomics	Epitomics # 3700-1	
Chemical compound, drug	OXONE®, monopersulf ate compound	Sigma Aldrich	Sigma Aldrich: 228036-5G	
Chemical compound, drug	N6-furfuryladen osine (kinetin)-5′-O-[3-thiotriphosp hate] sodium salt (KTPγS)	Axxora	Axxora:BL G-F008-05	
Chemical compound, drug	Adenosine 5′-triphosphate disodium	Sigma Aldrich	Sigma Aldrich:A64 19-1G	

	salt hydrate (ATP)			
Chemical compound, drug	Guanosine 5′-Triphosphate, Disodium Salt (GTP)	Calbiochem	Millipore:371701	
Chemical compound, drug	β-Casein from bovine milk	Sigma Aldrich	Sigma Aldrich:C6905	
Chemical compound, drug	Biotin	Sigma Aldrich	Sigma Aldrich: B4501-1G	
Chemical compound, drug	PP1 Analog III, 3-MB-PP1	Calbiochem	Sigma Aldrich:529582	
Commercial assay or kit	S-trap micro	Protifi	Protific:C02-micro-80	
Commercial assay or kit	Pierce Quantitative Fluorometric Peptide Assay	Thermo Fisher Scientific	Thermo Fisher Scientific:23290	
Commercial assay or kit	TMTpro™ 16plex Label Reagent Set	Thermo Fisher Scientific	Thermo Fisher Scientific:A44522	
Commercial assay or kit	EasyPep™ MS Sample Prep Kits - Maxi	Thermo Fisher Scientific	Thermo Fisher Scientific:A45734	
Commercial assay or kit	High-Select TiO2 Phosphopeptide Enrichment Kit	Thermo Fisher Scientific	Thermo Fisher Scientific:A32993	

Commercial assay or kit	High-Select™ Fe-NTA Phosphopeptide Enrichment Kit	Thermo Fisher Scientific	Thermo Fisher Scientific:A32992	
Commercial assay or kit	Pierce High pH Reversed-Phase Peptide Fractionation Kit	Thermo Fisher Scientific	Thermo Fisher Scientific:84868	
Commercial assay or kit	Dynabeads® MyOne™ Streptavidin C1	Invitrogen	Cat# 65001	
Commercial assay or kit	Bio-Rad DC assay	Bio-Rad	Bio-Rad:500011 16	
Commercial assay or kit	Sep-Pak C18 Plus Short Cartridge, 360 mg Sorbent per Cartridge, 55-105 µm	Waters	Waters:WAT020515	
Commercial assay or kit	SulfoLink™ Coupling Resin	Thermo Fisher Scientific	Thermo Fisher Scientific:20401	
Commercial assay or kit	Radiance Plus	Azure Biosystems	VWR:10147-298	

	Chemiluminescent Substrate			
Commercial assay or kit	Pierce™ Anti-HA Magnetic Beads	Thermo Scientific	Thermo Scientific:88836	
Commercial assay or kit	Pierce™ Streptavidin Magnetic Beads	Thermo Scientific	Thermo Scientific:88817	
Recombinant DNA reagent	All plasmids used in this study are listed in Supplementary file 10			
Sequence-based reagent	All primers and oligonucleotides used in this study are listed in Supplementary file 10			
Software, algorithm	Proteome Discoverer 4.2	Thermo Fisher		
Software, algorithm	MaxQuant (version 1.5.0.25 and 1.5.2.8)	PMID: 19029910	RRID:SCR_014485	Free software for searching of mass spectrometry acquisition files
Software, algorithm	Perseus (version 1.5.0.9)	PMID: 27348712	RRID:SCR_015753	Free software for processing of MaxQuant output files

Software, algorithm	Scaffold DIA	Proteome Software		
Software, algorithm	R version 4.0	R Foundation for Statistical Computing		
Software, algorithm	Prism 8	GraphPad		
Software, algorithm	Fiji	PMID: 22743772		
Software, algorithm	ilastik	PMID: 31570887	ilastik.org	
Software, algorithm	HHPRED	PMID: 29258817		
Software, algorithm	SnapGene	other	https://www.snapgene.com/	
Software, algorithm	ToxoDB	PMID: 18003657	ToxoDB.org	
other	DMEM, high glucose (media)	Life Technologies	Life Technologies:11965118	
other	HALT protease inhibitor	Thermo Fisher	Thermo Fisher:87786	Materials and Methods: lysis buffer
other	HALT protease and phosphatase inhibitor	Thermo Fisher	Thermo Fisher:PI78440	Materials and Methods: lysis buffer
other	Benzonase	Sigma Aldrich	Sigma Aldrich:E1014	Materials and Methods: lysis buffer

Cell culture

T. gondii parasites were grown in human foreskin fibroblasts (HFFs, ATCC SRC-1041) maintained in DMEM (GIBCO 11965118) supplemented with 3% inactivated fetal calf serum and

677   10 µg/mL gentamicin (Thermo Fisher Scientific), referred to as media. When noted, DMEM was
678   supplemented with 10% inactivated fetal bovine serum (IFS) and 10 µg/mL gentamicin, referred
679   to as 10% IFS media. Parasites and HFFs were grown at 37°C/5% CO_2 unless indicated
680   otherwise.
681
Parasite transfection and strain construction

Genetic background of parasite strains
T. gondii RH strains were used as genetic backgrounds for this study. All strains contain the
Δku80Δhxgprt mutations to facilitate homologous recombination (Huynh and Carruthers, 2009).
TIR1 expresses the TIR1-FLAG ubiquitin ligase and the CAT enzyme conferring chloramphenicol
resistance (Brown et al., 2017).
688
Transfection
Parasites were pelleted at 1000 x *g* for 5 to 10 min and resuspended with Cytomix (10 mM KPO_4,
120 mM KCl, 0.15 mM $CaCl_2$, 5 mM $MgCl_2$, 25 mM HEPES, 2 mM EDTA, 2 mM ATP, and 5 mM
glutathione) and combined with DNA to a final volume of 400 µL. Parasites were electroporated
using an ECM 830 Square Wave electroporator (BTX) in 4 mm cuvettes with the following setting:
1.7 kV, 2 pulses, 176 µs pulse length, and 100 msec interval.
695
Endogenous tagging of CDPK1 (CDPK1-AID)
CDPK1-AID was generated in the study (Shortt et al., 2022). Briefly, *V5-mNG-mAID-Ty* was PCR
amplified from pBM050 (V5-TEV-mNG-mAID-Ty; GenBank: OM640006) to attach homology
arms to *TGGT1_301440* (*CDPK1*). DNA was co-transfected with a Cas9 expression plasmid
targeting *CDPK1* in TIR1 parasites. mNG-expressing parasites were isolated using FACS and
subcloned by limiting dilution. Tagging was confirmed by PCR, flow cytometry, and
immunoblotting.
703
Endogenous tagging of CDPK1 (iKD) and complementation (cWT and cMut)
Generating the inducible CDPK1 knock-down strain (iKD CDPK1). The pTUB1_YFP_mAID_3HA
vector was amplified by inverse PCR using primer pair P1/P2 to substitute the 3xHA tag
sequence for a Myc tag encoding sequence(Brown et al., 2017). A Cas9 expression plasmid
targeting the 3'UTR of *CDPK1* was generated by inverse PCR on pSag1_Cas9-U6_sgUPRT
using primers P3/P4(Shen et al., 2014). The sequence encoding *mAID-Myc-HXGPRT* was PCR
amplified using primer pair P5/P6 and co-transfected into TIR1 parasites with the Cas9 expression
plasmid. Recombinant parasites were selected 24 hrs post transfection by addition of
mycophenolic acid (MPA; 25 µg/mL) and xanthine (XAN; 50 µg/mL) to culture medium. Lines
were cloned, and successful 5' and 3' integration of the *mAID-Myc-HXGPRT* cassette was
confirmed using primer pairs P30/P31 and P32/P33. Absence of WT was confirmed using primers
P34/P35.
716
Generating the cWT and cMut CDPK1 complementation strain. To generate the
complementation construct, pUPRT_CDPK1_ HA_T2A_GFP (GenBank: Pending), the *CDPK1*

5'UTR was amplified from genomic DNA using primer pair P7/P8. Recodonized *CDPK1* cDNA-
HA sequence was synthesized (GeneArt strings, Life Technologies) and amplified with
appropriate overhangs using primers P9/P10. Sequence encoding T2A-GFP was amplified from
an in-house unpublished plasmid using primer pair P11/P12. The three resulting fragments were
Gibson cloned into the PacI-linearized pUPRT_HA vector(Reese et al., 2011). To generate the
complementation construct, pUPRT_CDPK1(G2A)_HA_T2A_mCherry (GenBank: Pending), the
CDPK1 5'UTR was amplified from genomic DNA using primer pair P13/P8. Recodonized *CDPK1-*
HA was amplified from pUPRT_CDPK1_ HA_T2A_GFP with appropriate overhangs using
primers P14/P15. Primers P13/P15 were used to introduce a G2A point mutation within *CDPK1*.
Sequence encoding *T2A-mCherry* was amplified from an in-house unpublished plasmid using
primer pair P11/P16. The three resulting PCR amplicons were Gibson cloned into the PacI-
linearized pUPRT_HA vector. Complementation plasmids were linearized with AclI and
individually co-transfected with the Cas9 expression plasmid targeting the *UPRT* locus.
Transgenic parasites were subjected to 5'-fluo-2'-deoxyuridine (FUDR) selection (5 µM) 24 hrs
post transfection. Resistant parasites were cloned, and successful 5' and 3' integration was
confirmed using primer pairs P36/P37 and P38/P39, respectively. Disruption of the endogenous
UPRT locus was confirmed using primer pair P40/P41.
736
Endogenous tagging of genes
Genes were endogenously tagged using the previously described high-throughput tagging (HiT)
strategy (Smith et al., 2022). Cutting units specific to each gene were purchased as gene
fragments (IDT gBlocks; P22 and P23) and integrated with the following empty HiT vector
backbones via Gibson assembly: pALH086 (V5-mAID-HA; GenBank: ON312869), pALH047 (V5-
3HA; GenBank: ON312868), and pALH173 (TurboID-Ty; GenBank: Pending).
743
Between 30 and 50 µg of each vector was linearized with BsaI and co-transfected with 50 µg of
the pSS014 Cas9 expression plasmid (GenBank: OM640002). Vectors targeting *TGGT1_289100*
(*HOOK*) were transfected into TIR1 or CDPK1-AID. Vectors targeting *TGGT1_264050* (*FTS*) were
transfected into TIR1. Parasites underwent drug selection for approximately 1 week in 10% IFS
media with 3 µM pyrimethamine or 25 µg/mL mycophenolic acid and 50 µg/mL xanthine,
followed by subcloning into 96-well plates. Single clones were screened for tag expression by
PCR, immunofluorescence, or immunoblot. PCR validation was performed using the primers
P44/P45 (FTS-AID) and P46/P47 (FTS-TurboID).
752
Endogenous tagging of HOOK N terminus
AID-HOOK N-terminal tagging was generated by PCR amplifying the gene fragment encoding
the *HA-mAID* with homology arms to *HOOK* (IDT gBlock; P17) using the primers P18/19. A
sgRNA targeting *HOOK* was assembled into the pSS013 Cas9 expression plasmid (GenBank:
OM640003) using the primers P20/P21. 10 µg of the PCR product was transfected with 50 µg of
the Cas9 expression plasmid. Parasites were subcloned into 96-well plates. Single clones were
screened using PCR primers P42/P43 and validated by immunofluorescence and immunoblot.
760
TIR1/pMIC2-mNeonGreen-TurboID-Ty

pMIC2-mNG-TurboID-Ty was amplified with primers P24/P25 from pALH184 (pMIC2-mNG-
TurboID-Ty-3'DHFR; GenBank: Pending) with homology arms to the 5′ and 3′ ends of a defined,
neutral genomic locus (Markus et al., 2019). Amplified DNA was co-transfected with the Cas9
expression plasmid targeting the neutral locus (pBM041; GenBank: MN019116). mNeonGreen-
expressing parasites were isolated by FACS and subcloned in 96-well plates. Expression was
confirmed by fluorescence.
768
Endogenous tagging of CLAMP
mNeonGreen DNA was amplified using PCR from pGL015 (V5-mNG-mAID-Ty; GenBank:
OM640005) with homology arms targeting the C terminus of *TGGT1_265790* (*CLAMP*) using the
primers P26/P27. A sgRNA targeting *CLAMP* was assembled into the pSS013 Cas9 expression
plasmid (GenBank: OM640003) using the primers P28/P29. Between 5 and 10 µg of
mNeonGreen DNA was co-transfected with 50 µg of the Cas9 expression plasmid targeting
CLAMP into TIR1 and AID-HOOK. Parasites were grown in 10% IFS media until lysing the HFF
monolayer, followed by FACS-mediated isolation of mNeonGreen expressing parasites and
subcloning into 96-well plates. Single clones were screened for mNeonGreen expression by PCR
using primers P48/P49 and live microscopy.
779
Sub-minute phosphoproteomics
Parasite harvest and treatment
T. gondii tachyzoites from the RH strain CDPK1-AID were used to infect nine 15 cm dishes. 3.75
x 10^7 parasites were used to infect each dish in 20 mL of media. Approximately 24 hrs later, the
media of eight dishes was replaced with 15 mL of media containing 1 µM of compound 1 to
block egress and synchronize parasites by inhibiting PKG (Donald et al., 2006; Hopp et al., 2012;
Taylor et al., 2010). The media of the ninth dish was replaced with media containing DMSO. On
day 2, parasites were harvested when the monolayer of HFFs in the DMSO dish were
approximately 80% lysed. The eight dishes treated with compound 1 were washed once with 10
mL of warm 1X PBS, once with 30 mL of warm 1X PBS, and incubated with 10 mL of warm
FluoroBrite (Fluorobrite DMEM A1896701, 4 mM glutamine, 10 µg/mL gentamicin) for 10 min at
37°C. Infected HFFs were scraped and passed through a 27G needle to mechanically liberate
parasites and passed through a 5 µm filter. Parasites were pelleted at 1000 x *g* at 4°C for 10 min,
resuspended in 10 mL of FluoroBrite and a 1:250 dilution was used to count parasites
(approximately 2.25 x 10^9 total parasites). Parasites were pelleted at 1000 x *g* at 4°C for 7 min
and resuspended in 2 mL of FluoroBrite. 1 mL of parasites were diluted into a total volume of 40
mL of FluoroBrite containing either 500 µM of auxin or vehicle of 1X PBS and incubated at 37°C
for 3.5 hrs. 50 µL of auxin- or vehicle-treated parasites and untagged TIR1 parasites were
analyzed by flow cytometry (Miltenyi MACSQuant VYB) to detect mNeonGreen fluorescence.
After confirming depletion of CDPK1, parasites were pelleted at 1000 x *g* for 10 min at room
temperature and resuspended in 210 µL of 500 µM auxin or vehicle of PBS diluted in FluoroBrite.
To obtain a time course, the 0 sec time point was first collected by mixing 16 µL of parasites
with 4 µL of 5X DMSO (0.5% DMSO in FluoroBrite) and immediately lysed with 20 µL of 2X Lysis
Buffer (10% SDS, 100 mM TEAB pH 7.5, 2 mM $MgCl_2$, and 2X HALT protease and phosphatase

inhibitors). To obtain the 9, 30, and 300 sec time points, 80 µL of parasites in 1.5 mL tubes were
incubated in a ThermoMixer (Eppendorf) set to 37°C. The parasites were stimulated by adding
20 µL of warmed 5X zaprinast (500 µM zaprinast in FluoroBrite) or a vehicle of 5X DMSO. At each
time point, 20 µL of stimulated parasites were transferred directly into 20 µL of 2X Lysis Buffer
to quench the reaction. Complete time courses were collected sequentially in the following order:
auxin-treated stimulated with zaprinast, vehicle-treated stimulated with zaprinast, auxin-treated
stimulated with DMSO, and vehicle-treated stimulated with DMSO. Lysates were treated with
benzonase at a final concentration of 5 units/µL to remove DNA and were immediately subjected
to protein cleanup and digestion.
813
Protein cleanup and digestion
Proteins were prepared for mass spectrometry using a modified version of the S-trap protocol
(Protifi). Proteins in lysates were reduced with 5 mM TCEP for 10 min at 55°C and alkylated with
15 mM MMTS for 10 min at room temperature. The lysates were acidified to a final concentration
of 1.2% v/v phosphoric acid. A 6X volume of S-trap binding buffer (90% methanol, 100 mM
TEAB, pH 7.55) was mixed to each sample to precipitate proteins. The solution was loaded onto
S-trap micro columns (Protifi) and spun at 4,000 x *g* for 1 min until all the solution had passed
through the column. The columns were washed four times with 150 µL of S-trap binding buffer
and centrifuged at 4,000 x *g* for 1 min between each wash. Proteins were digested on-column
with 0.75 µg of trypsin (Promega) in 50 mM TEAB pH 8.5 overnight at 37°C in a humidified
incubator. Digested peptides were eluted in three steps at 4,000 x *g* for 1 min: 40 µL of 50 mM
TEAB, 40 µL of 0.2% formic acid, and 35 µL of 50% acetonitrile/0.2% formic acid. The peptide
concentrations of eluted peptides were quantified using the Pierce Fluorometric Peptide Assay
(Thermo Fisher Scientific) according to manufacturer's instructions. The remaining samples were
frozen in liquid nitrogen and lyophilized.
829
TMTpro labeling
Lyophilized peptides were resuspended in 100 mM TEAB pH 8.5 to peptide concentrations of
1.6 µg/µL. TMTpro reagents (Thermo Fisher Scientific; A44522 LOT# VI306829) were
resuspended in acetonitrile to 25 µg/µL. 80 µg of peptides in 50 µL were combined with 250 µg
of TMTpro reagent in 10 µL to achieve approximately a 3:1 label:peptide weight/weight ratio
(Zecha et al., 2019). The TMTpro labels were assigned to minimize reporter ion interference and
inter batch variability in the following scheme: Replicate 1 auxin: 0, 9, 30, 300 sec (126, 128C
130C, 132C); Replicate 1 vehicle: 0, 9, 30, 300 sec (127N, 129N, 131N, 133N); Replicate 2 auxin:
0, 9, 30, 300 sec (127C, 129C, 131C, 133C); Replicate 2 vehicle: 0, 9, 30, 300 sec (128N, 130N,
132N, 134) (Brenes et al., 2019). The zaprinast and DMSO samples were incubated for 1 hr at
room temperature shaking at 400 rpm. Unreacted TMTpro reagent was quenched with
hydroxylamine at a final concentration of 0.2%. The samples were pooled, acidified to 3% with
formic acid, and were processed using the EasyPep Maxi Sample Prep column (Thermo Fisher
Scientific) according to the manufacturer's instructions. 5% of the eluate volume was reserved
as the unenriched proteome sample. The remaining eluted peptides were frozen in liquid nitrogen
and lyophilized.
846

Phosphopeptide enrichment

Phosphopeptides were enriched using the Sequential enrichment from Metal Oxide Affinity
Chromatography (SMOAC) protocol according to manufacturer instructions (Tsai et al., 2014).
First, the High-Select TiO$_2$ Phosphopeptide Enrichment Kit (Thermo Fisher Scientific) was used
to enrich phosphopeptides from lyophilized TMTpro-labeled samples. The flow-through and
contents of the first wash were pooled, frozen in liquid nitrogen, and lyophilized, along with the
eluate. Second, the High-Select Fe-NTA Phosphopeptide Enrichment Kit (Thermo Fisher
Scientific) was used to enrich phosphopeptides from the pooled flow-through and first wash
from the previous enrichment. The eluted phosphopeptides were frozen in liquid nitrogen and
lyophilized.

Fractionation

The enriched and unenriched samples were fractionated with the Pierce High pH Reversed-
Phase Peptide Fractionation Kit (Thermo Fisher Scientific) according to manufacturer
instructions for TMT-labeled peptides. The acetonitrile wash was omitted for enriched samples
to prevent loss of phosphopeptides. The eluted peptides from the High-Select TiO$_2$
Phosphopeptide Enrichment and High-Select Fe-NTA Phosphopeptide Enrichment were pooled
prior to fractionation. 100 µg of unenriched samples were fractionated. 8 fractions were collected
for each TMTpro set: zaprinast phosphoproteome (enriched) [1], zaprinast proteome
(unenriched) [2], DMSO phosphoproteome (enriched) [3], and DMSO proteome (unenriched) [4].
The fractions were frozen in liquid nitrogen and lyophilized.

MS data acquisition

Lyophilized peptides were resuspended in approximately 15 µL (enriched) or 50 µL (unenriched)
of 0.1% formic acid and were analyzed on an Exploris 480 Orbitrap mass spectrometer equipped
with a FAIMS Pro source (Bekker-Jensen et al., 2020) connected to an EASY-nLC 1200
chromatography system using 0.1% formic acid as Buffer A and 80% acetonitrile/0.1% formic
acid as Buffer B. Peptides were loaded onto a heated analytical column (ES900, Thermo,
PepMap RSLC C18 3 µm, 100 Å, 75 µm x 15 cm, 40°C) via trap column (164946, Thermo,
Acclaim PepMap C18 3 µm, 100 Å, 75 µm x 20 mm nanoViper). Peptides were separated at 300
nL/min. Enriched samples were separated on a gradient of 5–20% B for 110 min, 20–28% B for
10 min, 28–95% B for 10 min, 95% B for 10 min, 95–2% B for 2 min, 2% B for 2 min, 2–98% B
for 2 min, 98% B for 2 min, 98–2% B for 2 min, and 2% B for 2 min. Unenriched samples were
separated on a gradient of 5–25% B for 110 min, 25–40% B for 10 min, 40–95% B for 10 min,
95% B for 10 min, 95–2% B for 2 min, 2% B for 2 min, 2–98% B for 2 min, 98% B for 2 min, 98–
2% B for 2 min, and 2% B for 2 min. The orbitrap and FAIMS were operated in positive ion mode
with a positive ion voltage of 1800V; with an ion transfer tube temperature of 270°C; using a
standard FAIMS resolution and compensation voltage of -50 and -65V, an inner electrode
temperature of 100°C and outer electrode temperature 80°C with 4.5 mL/min carrier gas. DDA
analysis was performed with a cycle time of 1.5 sec. Full scan spectra were acquired in profile
mode at a resolution of 60,000, with a scan range of 400–1400 m/z, 300% AGC target, maximum
injection time of 50 msec, intensity threshold of 5 x 10^4, 2–5 charge state, dynamic exclusion of
30 sec, mass tolerance of 10 ppm, purity threshold of 70%, and purity window of 0.7. MS2

spectra were generated with a HCD collision energy of 32 at a resolution of 45,000 using
TurboTMT settings with a first mass at 110 m/z, an isolation window of 0.7 m/z, 200% AGC
target, and maximum injection time of 120 msec.
893
Phosphoproteomic time course analysis
Raw files were analyzed in Proteome Discoverer 2.4 (Thermo Fisher Scientific) to generate peak
lists and protein and peptide IDs using Sequest HT (Thermo Fisher Scientific) and the ToxoDB
release 49 GT1 protein database. The maximum missed cleavage sites for trypsin was limited to
2. Precursor and fragment mass tolerances were 10 ppm and 0.02 Da, respectively. The
following modifications were included in the search: dynamic oxidation (+15.995 Da; M), dynamic
phosphorylation (+79.966 Da; S,T,Y), dynamic acetylation (+42.011 Da; N-terminus), static
TMTpro (+304.207 Da; any N-terminus), static TMTpro (+304.207 Da; K), and static methylthio
(+45.988 Da; C). TMTpro 16plex isotope correction values were accounted for (Thermo Fisher
Scientific; A44522 LOT# VI306829). Peptides identified in each sample were filtered by
Percolator to achieve a maximum FDR of 0.01 (Käll et al., 2008a, 2008b, 2007). Site localization
scores were generated using ptmRS, with phosphoRS and use of diagnostic ions set to true
(Taus et al., 2011). Reporter ion quantification used an integration tolerance of 20 ppm on the
most confident centroid. For reporter ion quantification, unique peptides were quantified using
a co-isolation threshold of 50, and average reporter signal-to-noise ratio of 10. Abundances were
normalized on the total protein amount. Protein level and peptide level ratios were generated for
each time point relative to 0 sec vehicle-treated parasites stimulated with DMSO.
911
Exported peptide and protein abundance files from Proteome Discoverer 2.4 were loaded into R
(version 4.1.1). To determine CDPK1-dependent phosphorylation, $\log_2$ ratios for peptide
abundances in DMSO and zaprinast-treated samples derived from Proteome Discoverer were
used. Only phosphorylated peptides quantified across all time points were used for analysis.
Area under the curve (AUC) values were calculated for individual peptides undergoing vehicle
(AUC_{veh}) and auxin (AUC_{auxin}) treatment using trapezoidal integration. AUC differences (AUC_{diff})
were calculated by taking the difference between AUC_{veh} and AUC_{auxin} values. The distribution of
zaprinast AUC_{diff} values were tested against a null distribution derived from the DMSO AUC_{diff}
values to calculate *z-scores* and *p-values* using a two-tailed *t*-test. Replicates 1 and 2 were
analyzed independently and phosphopeptides with *p*-values < 0.05 across both replicates were
determined to be CDPK1-dependent (Group A). Peptides exhibiting phosphorylation
independent of CDPK1 (Group B, C, and D) were determined similarly, but compared the
distribution of zaprinast $AUC_{vehicle}$ values to a null distribution of DMSO $AUC_{vehicle}$ values and
excluded phosphopeptides already determined to be CDPK1-dependent. CDPK1-independent
phosphopeptides were clustered into Groups B, C, and D using projection-based clustering
(Thrun and Ultsch, 2021).
928
Gene ontology enrichment

Gene ontology terms were obtained for all genes present in the enriched zaprinast
phosphoproteome from ToxoDB.org (Computed GO function and GO function IDs). Gene

ontology IDs within each group of zaprinast-dependent genes (Groups A-D) were tested for
enrichment against the entire enriched zaprinast phosphoproteome. Enrichment p values were
generated using a hypergeometric test. Enrichment ratios were calculated by dividing the gene
ratio (overlap/signatures) by the relative frequency of gene sets (gene sets/background). Only
GO IDs with significant enrichment and an overlap of 2 were plotted. Thiophosphorylation
enrichment was performed similarly.

Metabolic tagging, click reaction, pull down, and Western blotting

Metabolic tagging and cell lysis
Upon infection of HFF monolayers the medium was removed and replaced by fresh culture media
supplemented with 25 µM YnMyr (Iris Biotech) or Myr (Tokyo Chemical Industry). The parasites
were then incubated for 16 hrs, washed with PBS (2x) and lysed on ice using a lysis buffer (PBS,
0.1% SDS, 1% Triton X-100, EDTA-free complete protease inhibitor (Roche Diagnostics)).
Lysates were kept on ice for 20 min and centrifuged at $17,000 \times g$ for 20 min to remove insoluble
material. Supernatants were collected and protein concentration was determined using a BCA
protein assay kit (Pierce).

Click reaction and pull down
Lysates were thawed on ice. Proteins (100–300 µg) were taken and diluted to 1 mg/mL using the
lysis buffer. A click mixture was prepared by adding reagents in the following order and by
vortexing between the addition of each reagent: a capture reagent (stock solution 10 mM in
water, final concentration 0.1 mM), $CuSO_4$ (stock solution 50 mM in water, final concentration 1
mM), TCEP (stock solution 50 mM in water, final concentration 1 mM), TBTA (stock solution 10
mM in DMSO, final concentration 0.1 mM) (Heal et al., 2011). Capture reagent used herein was
the Trypsin cleavable reagents (Broncel et al., 2020). Following the addition of the click mixture
the samples were vortexed (room temperature, 1 hr), and the reaction was stopped by addition
of EDTA (final concentration 10 mM). Subsequently, proteins were precipitated
(chloroform/methanol, 0.25:1, relative to the sample volume), the precipitates isolated by
centrifugation ($17,000 \times g$, 10 min), washed with methanol (1 x 400 µL) and air dried (10 min). The
pellets were then resuspended (final concentration 1 mg/mL, PBS, 0.4 % SDS) and the
precipitation step was repeated to remove excess of the capture reagent. Next, samples were
added to 15 µL of pre-washed (0.2 % SDS in PBS (3 x 500 µL)) Dynabeads® MyOneTM
Streptavidin C1 (Invitrogen) and gently vortexed for 90 min. The supernatant was removed, and
the beads were washed with 0.2 % SDS in PBS (3 x 500 µL).

SDS-PAGE and Western blotting
Beads were supplemented with 2% SDS in PBS (20 µL) and 4x SLB (Invitrogen), boiled (95°C,
10 min), centrifuged (1,000 x g, 2 min) and loaded on 10% or 4–20% SDS-PAGE gel (Bio-Rad).
Following electrophoresis (60 min, 160V), gels were briefly washed with water and proteins were
transferred (25 V, 1.3 A, 7 min) onto nitrocellulose membranes (Bio-Rad) using Bio-Rad Trans
Blot Turbo Transfer system. After brief wash with PBS-T (PBS, 0.1% Tween-20) membranes
were blocked (5% milk, TBS-T, 1h) and incubated with primary antibodies (5% milk, TBS-T,

overnight, 4°C) at the following dilutions: rat anti-HA (1:1000; Roche Diagnostics), mouse anti-
Myc (1:1000; Millipore), rabbit anti-Gra29 (1:1000; Moritz Treeck Lab), rabbit anti-SFP1 (1:1000;
Moritz Treeck Lab), mouse anti-T. gondii [TP3] (1:1000; Abcam), mouse anti-CDPK1 (1:3000;
John Boothroyd Lab), rabbit anti-SAG1 (1:10,000; John Boothroyd Lab), rabbit anti-GAP45
(1:1000; Peter Bradley Lab), mouse anti-GFP (1:1000, Roche Diagnostics) and rabbit anti-
mCherry (1:1000, Abcam). Following washing (TBS-T, 3x) membranes were incubated with IR
dye-conjugated secondary antibodies (1:10,000, 5% milk, TBS-T, 1 hr) and after a final washing
step imaged on a LiCOR Odyssey imaging system (LI-COR Biosciences).
982

MS detection of myristoylated CDPK1

984 Mass spectrometry proteomics methods and data for myristoylated CDPK1 are available from
985 the ProteomeXchange Consortium via the PRIDE partner repository (ID PXD019677) and the
986 associated publication (Broncel et al., 2020).
987

Depletion of mAID tagged CDPK1 (iKD)

989 Parasites were treated with 500 µM auxin or equivalent volume of vehicle (ethanol) for at least 2
990 hrs prior to Western blot analysis.
991

Subcellular fractionation

RH Δ*ku80*Δ*hxgprt* YFP expressing parasites were metabolically tagged with 25 µM Myr or YnMyr
for 16 hrs. Following a PBS wash, the parasites were syringe lysed in Endo buffer (44.7 mM
K_2SO_4, 10 mM $MgSO_4$, 106 mM sucrose, 5 mM glucose, 20 mM Tris–H_2SO_4, 3.5 mg/mL BSA,
pH 8.2) and collected by centrifugation (512 x *g*, 10 min). The parasites were then lysed in 300
µL of cold hypotonic buffer (10 mM HEPES, pH 7.5) supplemented with protease inhibitors
(Roche), passed through 25G needle (5x) and left on ice for 40 min. Next, lysates were pelleted
by centrifugation (16,000 x *g*, 20 min, 4 °C) and the resulting supernatant was subjected to an
additional high speed (100,000 x *g*, 1 h, 4 °C) centrifugation step. To avoid the loss of the high-
speed pellet, only half of the supernatant was removed at this point. Each fraction was then
taken up in 0.4% (final) SDS HEPES, clicked to a capture reagent and pulled down as described
above. Myristoylation-dependent partitioning was revealed by SDS-PAGE and Western blotting.

Myristoylation-dependent fractionation for CDPK1 complemented WT and Mut lines: parasites
were seeded 24 hrs prior experiment. Following a PBS wash, the parasites were syringe lysed in
Endo buffer (44.7 mM K_2SO_4, 10 mM $MgSO_4$, 106 mM sucrose, 5 mM glucose, 20 mM Tris–
H_2SO_4, 3.5 mg/mL BSA, pH 8.2) and collected by centrifugation (512 x *g*, 10 min). The parasites
were then lysed in 300 µL of cold hypotonic buffer (10 mM HEPES, pH 7.5) supplemented with
protease inhibitors (Roche), passed through 25G needle (5x) and left on ice for 40 min. Next,
lysates were pelleted by centrifugation (100,000 x *g*, 1 hr, 4 °C), the supernatant was removed,
and cytosolic proteins precipitated with methanol/chloroform. Proteins from the pellet and
supernatant fractions were dissolved in 2% (final) SDS PBS and myristoylation-dependent
partitioning was revealed by SDS-PAGE and Western blotting.

Immunoprecipitation of cWT and cMUT CDPK1

Parasite harvest

cWT, cMUT, and TIR1 (untagged) parasites were infected onto confluent HFFs in 15 cm dishes. At 1-day post-infection (dpi), 50 µM auxin or vehicle was added to deplete endogenous mAID-tagged CDPK1 and 1 µM compound 1 was added to block egress until parasites were ready to harvest. At 2 dpi, infected HFFs were washed twice with PBS to wash out drugs. Parasites were mechanically released in Endo buffer with a 27G needle, passed through a 5 µm filter, and spun at 1000 x *g* for 10 min. Parasite pellets were resuspended in a cold hypotonic buffer with 1X HALT protease inhibitors to parasite concentrations of 1.1 x 10^9 tg/mL, passed through a 27G needle five times, and incubated on ice for 1 hr to complete hypotonic lysis. The samples were spun at 1000 x *g* for 5 min to pellet unlysed parasites and the supernatant was saved. NaCl was added to a final concentration of 150 mM and this was used as the immunoprecipitation input.

Immunoprecipitation

25 µL of anti-HA magnetic beads (Thermo) were used per condition. Beads were washed twice with a wash buffer (10 mM HEPES, 150 mM NaCl, pH 7.5). To begin pulldown, parasite lysate was used to resuspend washed beads and incubated for 1 hr rotating at room temperature. Beads were washed four times with a wash buffer. Proteins were eluted by resuspending beads in 20 µL of 1X S-trap sample buffer (5% SDS, 50 mM TEAB, pH 7.5) and incubated at 70°C for 10 min. The eluate was collected for MS sample processing and analysis. Results are representative of two independent experiments.

Protein cleanup and digestion

Proteins were prepared for mass spectrometry as described above in "Sub-minute phosphoproteomics - Protein cleanup and digestion". Eluted peptides were frozen in liquid nitrogen, lyophilized, and stored at -80°C until MS analysis.

MS data acquisition

Lyophilized peptides were resuspended in 20 µL of 0.1% formic acid and were analyzed on an Exploris 480 Orbitrap mass spectrometer equipped with a FAIMS Pro source (Bekker-Jensen et al., 2020) connected to an EASY-nLC chromatography system using 0.1% formic acid as Buffer A and 80% acetonitrile/0.1% formic acid as Buffer B. Peptides were separated at 300 nL/min on a gradient of 1–6% B for 1 min, 6–21% B for 41 min and 30 sec, 21–36% B for 20 min and 45 sec, 36–50% B for 10 min and 15 sec, 100% B for 14 min and 30 sec, 100–2% B for 3 min, 2% B for 3 min, 2–98% B for 3 min, and 98% B for 3 min. The orbitrap and FAIMS were operated in positive ion mode with a positive ion voltage of 1800V; with an ion transfer tube temperature of 270°C; using a standard FAIMS resolution and an inner and outer electrode temperature of 100°C with 4.5 mL/min carrier gas. Samples were analyzed twice in DIA mode with a compensation mode of -50 and -65V. Full scan spectra were acquired in profile mode at a resolution of 120,000, with a scan range of 400–1000 m/z, 300% AGC target, and auto mode for maximum injection time. MS2 spectra for the DIA scan were generated with a isolation window of 20 m/z with a 0 m/z window overlap, 30 scan events, a HCD collision energy of 30 at a

1056 resolution of 30,000, first mass at 200 m/z, precursor mass range of 400–1000 m/z, and a
1057 standard AGC target and automatically determined maximum injection time.
1058
Immunoprecipitation data analysis
DIA-MS samples were analyzed using Scaffold DIA (2.0.0). DIA-MS data files were converted to
mzML format using ProteoWizard (3.0.19254) (Chambers et al., 2012). Analytic samples were
aligned based on retention times and individually searched against
dku80_FAIMS_DIA_90min_autoIT.blib with a peptide mass tolerance of 10 ppm and a fragment
mass tolerance of 0.02 Da. The ToxoDB release 46 GT1 protein database was used for protein
identification. The following modifications were included in the search: dynamic oxidation
(+15.995 Da; M), dynamic phosphorylation (+79.966 Da; S,T,Y), and static methylthio (+45.988
Da; C). The digestion enzyme was trypsin with a maximum of 2 missed cleavage sites allowed.
Only peptides with charges in the range of 2 to 3 and length in the range 6 to 30 were considered.
Peptides identified in each sample were filtered by Percolator to achieve a maximum FDR of
0.01 (Käll et al., 2008a, 2008b, 2007). Individual search results were combined, and peptide
identifications were assigned posterior error probabilities and filtered to an FDR threshold of 0.01
by Percolator. Peptide quantification was performed by Encyclopedia (0.9.2). For each peptide,
the 5 highest quality fragment ions were selected for quantification. Proteins that contained
similar peptides and could not be differentiated based on MS/MS analysis were grouped to
satisfy the principles of parsimony. Protein groups were thresholded to achieve a protein FDR
less than 1%. Significance values were derived from *t*-tests across two replicates and adjusted
with Benjamini-Hochberg correction with an FDR of 0.05. Exported protein abundance files from
Scaffold DIA were loaded into R (version 4.1.1).
1079
Thiophosphorylation of CDPK1 substrates
Parasite harvest and treatment
T. gondii tachyzoites from the RH strain (CDPK1[G] and CDPK1[M]) were passaged twice across 4
days in SILAC media in T12.5 flasks. CDPK1[G] parasites were grown in "heavy" SILAC media
(DMEM 88364, 10% dialyzed FBS, 0.1 mg/mL $^{13}C^{15}N$ L-arginine, 0.1 mg/mL $^{13}C^{15}N$ L-lysine) and
CDPK1[M] parasites were grown in "light" SILAC media (DMEM 88364, 10% dialyzed FBS, 0.1
mg/mL L-Arginine, 0.1 mg/mL L-Lysine). On day 4, confluent HFFs in 15 cm dishes were infected
with CDPK1[G] and CDPK1[M] parasites in SILAC media. On day 6, extracellular parasites were
harvested by filtering through a 5 µm filter and pelleted at 1000 x *g* for 7 min at 4°C. Parasites
were washed once in 1X intracellular buffer (ICB) (137 mM KCl, 5 mM NaCl, 20 mM HEPES, 10
mM $MgCl_2$, pH 7.5 KOH) and then resuspended in 400 µL of 1X ICB. Parasites were semi-
permeabilized after the addition of 400 µL of 6 µg/mL aerolysin diluted in 1X ICB and incubated
at 37°C for 10 min. After semi-permeabilization, 400 µL of 4X Ca^{2+} solution (16 mM CaEGTA, 100
ng/mL 1B7 nanobody in 1X ICB) followed by 400 µL of 4X ATP solution (4 mM GTP, 0.4 mM
ATP, 0.2 mM KTPγS, 1X HALT protease and phosphatase inhibitor in 1X ICB). The kinase
reaction was initiated by incubating parasites at 30°C for 5 min. Parasites were pelleted at 1000
x *g* at 4°C for 10 min. Parasite pellets were resuspended in 250 µL of 1X lysis buffer (10% TritonX-
100, 1X HALT protease and phosphatase inhibitor, 10 mM K_2EGTA in 1X ICB).

1098

Protein quantification and thiophosphorylation immunoblotting

1099

Proteins in lysates were quantified using DC assay (BioRad) utilizing BSA as a protein standard
and a diluent (150 mM NaCl, 20 mM Tris pH 7.6) to prepare standard curves and dilution series.
Thiophosphorylation was verified using immunoblot by first incubating 3 µL of sample with p-
nitrobenzyl mesylate diluted in 1X ICB at a final concentration at 2 mM for 2 hrs at room
temperature. 5X Laemmli sample buffer was added (see "Immunoblotting" for recipe) and
samples were boiled for 10 min. Samples were resolved on a home-made polyacrylamide gel
(5% stacking, 15% resolving, 15-well, 0.75 mm) and transferred overnight at 4°C. Nitrocellulose
membranes were blocked with 5% milk in TBS-T for 1 hr. Primary antibody incubations were
performed with an anti-thiophosphate ester antibody (rabbit 51-8; 1:5000) and anti-tubulin
(mouse 12G10; 1:2000) for 1 hr at room temperature. Secondary incubations were performed
with LI-COR antibodies (rabbit 680, mouse 800; 1:10,000) for 1 hr at room temperature. Imaging
was performed using a LI-COR Odyssey.

1112

Protein cleanup and digestion

1113

Proteins were precipitated using methanol chloroform extraction. For 250 µL of lysate, 800 µL
methanol and 200 µL of chloroform was added, followed by vortexing. After adding 600 µL of
water, the sample was vortexed and centrifuged at max speed for 5 min at 4°C. After removing
the supernatant without disrupting the precipitate, 375 µL of methanol was added and vortexed.
Samples were centrifuged at max speed for 15 min at 4°C. The protein pellet was allowed to air
dry after removing the supernatant. Dried protein pellets were resuspended in 200 µL of 8M urea.
1 mg of protein (determined from protein quantification; see above) from CDPK1^G and CDPK1^M
parasites were pooled for a total of 2 mg of protein. To digest protein, 5X volume of 1X Trypsin
Digest Buffer (100 mM ammonium acetate pH 8.9, 1mM CaCl$_2$, 2 mM TCEP) was added to the
sample followed by 40 µg of sequencing grade trypsin (Promega). Proteins were digested
overnight rotating at room temperature. To prepare samples for desalting, glacial acetic acid was
added to 10% (v/v) and debris was briefly spun down. A C18 Sep-Pak Plus cartridge (Waters)
was prepared using a syringe pump and 10 mL syringe with three 10 mL washes at a flow rate
of 2 mL/min: 0.1% acetic acid, 90% acetonitrile/0.1% acetic acid, and 0.1% acetic acid. The
sample was loaded in a 5 mL syringe at a flow rate of 0.5 mL/min. The syringe and cartridge
were washed with 5 mL of 0.1% acetic acid at a flow rate of 0.5 mL/min. A final wash was
performed using 10 mL of 0.1% acetic acid at a flow rate of 2 mL/min. Peptides were eluted with
4.5mL of 40% acetonitrile/0.1% acetic acid at a flow rate of 0.5 mL/min. Samples were spun in
a speed vac for 3 hrs until samples could be pooled into a single 2 mL tube. Samples were frozen
in liquid nitrogen and lyophilized overnight. Lyophilized peptides were stored at -80°C.

1134

Thiophosphate enrichment

1135

To enrich for thiophosphorylated peptides, 400 µL of SulfoLink Coupling Resin slurry was used
(Thermo). Incubations were performed in the dark using aluminum foil due to the light sensitivity
of the resin. The resin was washed twice with 1 mL of binding buffer (25 mM HEPES, 50%
acetonitrile, pH 7.0 NaOH + HCl) rotating for 5 min followed by a 1000 rpm spin for 10 sec. The
supernatant was removed without disturbing the pelleted resin. The beads were blocked with

1mL of blocking buffer (binding buffer with 25 μg/mL β-casein and 2 mM TCEP) and rotated for
5 min at room temperature. After pelleting the resin and removing the supernatant, the resin was
resuspended with lyophilized peptides dissolved in 500 μL of the blocking buffer. Samples were
incubated overnight at room temperature. Samples were spun at 1000 rpm for 10 sec and the
supernatant was spun again, lyophilized, and stored at -80°C prior to MS analysis. The beads
received the following series of 1 mL washes for 5 min rotating and spun at 1000 rpm for 10 sec:
twice with binding buffer with 2 mM TCEP, once with quenching buffer (25 mM HEPES, 50%
acetonitrile, 5 mM DTT, pH 8.5 NaOH), once with binding buffer with 2 mM TCEP, once with 5%
formic acid (no rotation), once with binding buffer with 2 mM TCEP, and three times with 0.1%
acetic acid. The resin was no longer light sensitive after incubation with the quenching buffer. To
elute captured peptides, resin was resuspended in 500 μL OXONE (2 mg/mL potassium
monopersulfate) and rotated for 5 min at room temperature. The supernatant containing eluted
peptides was collected after spinning at 1000 rpm for 10 sec. OXONE was removed from the
sample using C18 spin columns (Pierce) according to manufacturer instructions. A total of 4
washes with the equilibrium/wash buffer was performed. Peptides were spun in a speed vac
until dry and stored at -80°C until MS analysis.
1157
MS data acquisition
The samples were resuspended in 10–20 μL of 0.1% formic acid for MS analysis and were
analyzed on a Q-Exactive HF-X Orbitrap mass spectrometer connected to an EASY-nLC 1200
chromatography system using 0.1% formic acid as Buffer A and 80% acetonitrile/0.1% formic
acid as Buffer B. Peptides were loaded onto a analytical column (column: PF360-75-15-N-5,
New Objective, 360 μm OD, 75 μm ID, 15 μm Tip PicoFrit Emitter; resin: 04A-4506, Phenomenex
Aeris Peptide, C18 1.7 μm) via trapping column (column: 360 um OD, 100 um ID; resin: AA12S11,
YMC Gel ODS-A, C18 10 μm). Peptides were separated at 300 nL/min on a gradient of 2% B for
5 min, 2–25% B for 100 min, 25–40% B for 20 min, 40–100% B for 1 min, and 100% B for 12
min. The orbitrap was operated in positive ion mode with a positive ion voltage of 2700V with an
ion transfer tube temperature of 300°C. Full scan spectra were acquired in profile mode at a
resolution of 60,000, with a scan range of 375 to 1600 m/z, 1×10^6 AGC target, maximum
injection time of 50 msec, intensity threshold of 4×10^5, dynamic exclusion of 13 sec, and 20
data dependent scans (DDA Top 20). MS2 spectra were generated with a HCD collision energy
of 27 at a resolution of 15,000, first mass at 100 m/z, isolation window of 1.5 m/z, an AGC target
of 1×10^5 with a maximum injection time of 20 msec, and scan range of 200 to 2000 m/z.
1174
Thiophosphorylation analysis
Raw files were analyzed in Proteome Discoverer 2.2 (Thermo Fisher Scientific) to generate peak
lists and protein and peptides identifications using Sequest HT (Thermo Fisher Scientific) and
the ToxoDB release 34 GT1 protein database. The maximum missed cleavage sites for trypsin
was limited to 2. The following modifications were included in the search: dynamic oxidation
(+15.995 Da; M), dynamic phosphorylation (+79.966 Da; S,T,Y), dynamic $^{13}C_6^{15}N_4$ (+10.008 Da;
R), dynamic $^{13}C_6^{15}N_2$ (+8.014 Da; K), and dynamic acetylation (+42.011 Da; N-terminus). Site
localization scores were generated using ptmRS, with phosphoRS and use of diagnostic ions
set to true. SILAC 2plex (Arg10, Lys8) method was used for relative quantification of protein and

unique peptide abundances. For peptide level analysis, ratios of unique peptides comparing
CDPK1^G and CDPK1^M were generated and low abundance resampling (5%) was used to impute
missing values. For protein level analysis, ratios of proteins were determined from the summed
abundance of unique peptides comparing CDPK1^G and CDPK1^M strains.
1188
Peptide and protein level data were exported from Proteome Discoverer for analysis in R. For
enriched peptides, abundance ratios were calculated within each replicate for high confidence
peptides (CDPK1^G/CDPK1^M) and normalized to the median abundance ratio of the whole
proteome peptides derived from the flow through samples. Normalized abundance ratios in the
flow through proteome samples were calculated by dividing abundance ratios by the median
abundance ratios within each replicate. For enriched peptides, an average $\log_2$ normalized
abundance ratio was calculated from three replicates. Significantly enriched peptides were
calculated using a one-tailed *t*-test using the normalized abundance ratios for three replicates.
A non-linear significance threshold was calculated using the function $(y = |4/x|)$, where
significantly enriched peptides had product value of $-\log_{10}(p)$*(mean $\log_2$ abundance ratio) greater
than 4.
1200
Immunoblotting

Samples were combined with 5X Laemmli sample buffer (10% SDS, 50% glycerol, 300mM Tris
HCl pH 6.8, 0.05% bromophenol blue, 5% beta-mercaptoethanol) and were incubated at 95°C
for 10 min. The samples were run on precast 4–15% or 7.5% SDS gels (Bio-Rad) and were
transferred overnight onto a nitrocellulose membrane in transfer buffer (25mM TrisHCl, 192 mM
glycine, 0.1% SDS, 20% methanol) at 4°C. Blocking was performed with 5% milk in PBS for 1
hr rocking at room temperature. Antibody incubations were performed with 5% milk in TBS-T for
1 hr rocking at room temperature. Three 5 min TBS-T washes were performed before and after
secondary antibody incubations rocking at room temperature. After a final PBS wash, imaging
was performed using a LI-COR Odyssey.
1211
1212 For immunoblot detection of the HA tag in AID-HOOK and FTS-AID after auxin-mediated
1213 depletion, secondary antibody incubations were performed with anti-rabbit HRP antibodies
1214 (Jackson ImmunoResearch) and detected with chemiluminescence (Azure) for increased
1215 sensitivity. Imaging was performed using a Bio-Rad Gel Doc XR.
1216
Immunofluorescence analysis

HFFs were seeded onto coverslips and grown until confluence. Confluent HFFs were infected
with parasites. Approximately 2 hrs later, media was exchanged with media containing either 50
µM auxin or vehicle solution of PBS. At 24 hrs post-infection, the media was aspirated, and the
coverslips were washed with PBS three times before fixation with 4% formaldehyde in PBS for
20 min. Following three washes in PBS, the fixed cells were permeabilized with 0.25% Triton X-
100 for 15 min. After three washes in PBS, the coverslips were incubated in a blocking solution
(5% IFS/ 5% NGS in PBS) for 10 min at room temperature. Coverslips were incubated for 1–2
hrs with primary antibody diluted in blocking solution. Anti-GAP45 or anti-MIC2 was used as a

parasite counterstain. After three washes in PBS, the coverslips were incubated in blocking
solution for 5 min, followed by secondary antibody diluted in blocking solution containing
Hoechst 33342 for 1 hr. The coverslips were washed three times in PBS, once in water, and
finally mounted with Prolong Diamond overnight at room temperature. Microscope images were
acquired with the Nikon Ti Eclipse and NIS Elements software package.
1231
Immunofluorescence analysis - Figure 2
Parasite-infected HFF monolayers grown on glass coverslips were fixed with 3% formaldehyde
for 15 min prior to washing with PBS. Fixed cells were then permeabilized (PBS, 0.1% Triton X-
100, 10 min), blocked (3% BSA in PBS, 1 hr), and labeled with anti-HA (1:1000, 1hr; Roche). HA-
tagged CDPK1 in the cWT and cMut lines was visualized with secondary goat antibodies (1:2000,
1 hr; Life Technologies) conjugated to Alexa Fluor 594 and 488, respectively. Cytosolic GFP
(cWT) and mCherry (cMut) were used as parasite counterstains. Nuclei were visualized with the
DNA stain (DAPI; Sigma) added at 5 µg/mL with the secondary antibody. Stained coverslips were
mounted on glass slides with Slowfade (Life Technologies) and imaged on a Nikon Eclipse Ti-U
inverted fluorescent microscope. Images were analyzed using Nikon NIS Elements imaging
software.
1243
Plaque Assays

600 and 1200 parasites were used to infect 6-well plates of HFFs in 10% IFS media. At 1 dpi,
media was exchanged with 10% IFS media containing either 50 µM auxin or vehicle solution of
PBS. Parasites were allowed to grow undisturbed for 8 days total. Plates were washed with PBS
and fixed for 10 min with 100% ethanol at room temperature. After removing ethanol, plates
were allowed to dry prior to staining with crystal violet solution for 30 min to 1 hr. Plates were
washed 3 times with PBS, once with water, and allowed to dry before scanning.
1251
1252 *Plaque assays - Figure 2*
1253 Parasites were harvested by syringe lysis, counted, and 200 parasites were seeded on confluent
1254 HFF monolayers grown in 24-well plates (Falcon). Wells were treated with 500 µM auxin or
1255 vehicle (ethanol) and plaques were allowed to form for 5 days. Plaque formation was assessed
1256 by inspecting the methanol fixed and 0.1% crystal violet stained HFF monolayers.
1257
Invasion assays

Confluent HFFs seeded in T12.5 flasks were infected with parasites. Approximately 2 hrs later,
the media was exchanged for 10% IFS media containing either 50 µM auxin or vehicle solution
of PBS. Parasites were harvested at 2 dpi and the media was exchanged for 1% IFS in invasion
media (DMEM Sigma D2902, 20 mM HEPES, pH 7.4) with auxin or vehicle PBS added. Confluent
HFFs seeded in clear-bottomed 96-well plates were infected with 2×10^5 extracellular parasites.
After centrifuging the plate at 290 x *g* for 5 min to synchronize invasion, the infected 96-well plate
was incubated by floating on a water bath at 37°C/5% CO_2 for 90 min. Wells were washed once
with PBS before fixing infected HFFs with 4% formaldehyde in PBS for 20 min at room
temperature. Wells were washed three times with a wash buffer (1% NGS in PBS) and then

incubated in a blocking buffer (5% IFS and 1% NGS in PBS) overnight at 4°C. To stain
extracellular parasites, wells were incubated with anti-SAG1 antibody for 30 min at room
temperature. After washing wells three times with a wash buffer, fixed HFFs were permeabilized
in 0.25% TritonX-100 in blocking buffer for 8 min room temperature. After washing the wells
three times with a wash buffer, the wells were incubated in a blocking buffer for 10 min at room
temperature, followed by anti-GAP45 antibody for 30 min at room temperature to label all
parasites. After three washes in a wash buffer, wells were incubated with a secondary antibody
solution containing Hoechst for 30 min. After three washes in PBS, wells were imaged using a
Biotek Cytation 3.
1277
Egress assays

Confluent HFFs seeded in glass bottomed 35 mm dishes (Ibidi or Mattek) were infected with
approximately 2 x 10^5 parasites. Approximately 2 hrs later, the media was exchanged for 10%
IFS media containing either 50 µM auxin or vehicle solution of PBS. At 24 hrs post-infection, the
media was replaced with 1% IFS in Ringer's (155 mM NaCl, 2 mM $CaCl_2$, 3 mM KCl, 1 mM
$MgCl_2$, 3 mM NaH_2PO_4, 10 mM HEPES, 10 mM glucose, pH 7.4). Zaprinast (final concentration
500 µM) was added into the dishes after 15 sec of imaging and imaging of infected HFFs
continued for a total of 10 min. Images were acquired using a Nikon Ti Eclipse with an enclosure
maintained at 37°C. The number of intracellular vacuoles was quantified at 0 min and 10 min.
Results are the sum of four fields of view per condition and are representative of three
independent experiments.
1289
Egress assays - Figure 2
Parasites were added to HFF monolayer and grown for 24 hrs in a 96 well plate. The wells were
then treated with 500 µM auxin or an equivalent volume of vehicle (ethanol) for 2 hrs and then
washed with PBS (2x). The media was exchanged for 100 µl Ringer's solution (155 mM NaCl, 3
mM KCl, 2 mM $CaCl_2$, 1 mM $MgCl_2$, 3 mM NaH_2PO_4, 10 mM HEPES, 10 mM glucose) and the
plate was placed on a heating block to maintain the temperature at 37°C. To artificially induce
egress, 50 µL of Ringer's solution containing 24 µM ionophore (8 µM final, Thermo) was added
to each well. At specified time points the cells were fixed by adding 33 µL 16% formaldehyde
(3% final) for 15 min. Cells were washed in PBS (3x) and stained with rabbit anti-TgCAP 1:2000
(Hunt et al., 2019) followed by goat anti-rabbit Alexa Fluor 488 (1:2000) and DAPI (5 µg/mL).
Automated image acquisition of 25 fields per well was performed on a Cellomics Array Scan VTI
HCS reader (Thermo Scientific) using a 20x objective. Image analysis was performed using the
Compartmental Analysis BioApplication on HCS Studio (Thermo Scientific). Egress levels were
determined in triplicate for three independent assays. Vacuole counts were normalized to $t = 0$
to determine how many intact vacuoles had remained after egress. The results were statistically
tested using one-way ANOVA with Tukey's multiple comparison test in GraphPad Prism 7. The
data are presented as mean ± s.d.
1307

Replication assays

Confluent HFFs seeded on coverslips were infected with parasites. Parasites were centrifuged at 290 x g for 5 min to synchronize invasion of host cells. Approximately 2 hrs later, the media was exchanged for 10% IFS media containing either 50 µM auxin or vehicle solution of PBS. At 24 hrs post-infection, the media was aspirated and fixed as described in "Immunofluorescence Analysis". Coverslips were incubated in primary and secondary antibodies for 30 min. Anti-GAP45 antibody was used to visualize parasites. Microscope images were acquired with the Nikon Ti Eclipse and NIS Elements software package. Tiled images were acquired in a four-by-four manner using a 40x objective. Parasites per vacuole were quantified for at least 100 vacuoles for three biological replicates.

Immunoprecipitation of HOOK and FTS complex

Parasite harvest
HOOK-3xHA, CDPK1-AID (parental), FTS-3xHA and TIR1 (parental) parasites were infected onto confluent HFFs in 15 cm dishes. After the parasites lysed the HFF monolayer (approximately 40 hrs post-infection), extracellular parasites were passed through 5 µm filters and washed twice with DMEM by pelleting at 1000 x g for 10 min. Parasite pellets were resuspended in 1X Dynein Lysis Buffer (DLB) (30 mM HEPES, 50 mM KOAc1, 2 mM MgOAc, 10% glycerol, 1 mM EGTA pH 8.0, 1 mM DTT, 0.5 mM ATP, 125 units/mL benzonase, 1X HALT protease and phosphatase inhibitor, and 1% NP-40 IGEPAL CA 630) to achieve a concentration of 6.67×10^8 parasites/mL (Redwine et al., 2017). Parasites were lysed on ice for 10 min and vortexed periodically to facilitate lysis. Lysates were centrifuged at 1000 x g for 5 min at room temperature to pellet unlysed parasites and the lysate supernatant was collected and used as the immunoprecipitation input.

Immunoprecipitation
25 µL of anti-HA magnetic beads (Thermo Scientific 88836) was used for 200 µL of parasite lysate. Beads were aliquoted and washed three times with a wash buffer (30 mM HEPES, 50 mM KOAc1, 2 mM MgOAc, 10% glycerol, 1 mM EGTA pH 8.0, 1 mM DTT, 0.5 mM ATP, and 0.01% NP-40 IGEPAL CA 630) using a magnetic rack. To begin the pulldown, 200 µL of parasite lysate was used to resuspend washed beads. Tubes were rotated at 4°C for 1 hr (FTS-AID and TIR1) and 3 hrs (HOOK-3xHA and CDPK1-AID). Beads were resuspended in 250 µL of wash buffer, transferred to a new tube, and received two additional washes with the same volume. Proteins were eluted by resuspending beads in 22 µL of 1X S-trap sample buffer (5% SDS, 50 mM TEAB, pH 7.5) and incubated at 70°C for 10 min. The eluate was collected for MS sample processing and analysis. Results are representative of three independent experiments.

Protein cleanup and digestion
Proteins were prepared for mass spectrometry as described above in "Sub-minute phosphoproteomics - Protein cleanup and digestion". Eluted peptides were frozen in liquid nitrogen, lyophilized, and stored at -80°C until MS analysis.

MS data acquisition

Lyophilized peptides were resuspended in 25 µL of 0.1% formic acid and were analyzed on an Exploris 480 Orbitrap mass spectrometer equipped with a FAIMS Pro source (Bekker-Jensen et al., 2020) connected to an EASY-nLC chromatography system using 0.1% formic acid as Buffer A and 80% acetonitrile/0.1% formic acid as Buffer B. Peptides were separated at 300 nL/min on a gradient of 1–6% B for 1 min, 6–21% B for 41 min and 30 sec, 21–36% B for 20 min and 45 sec, 36–50% B for 10 min and 15 sec, 100% B for 14 min and 30 sec, 100–2% B for 3 min, 2% B for 3 min, 2–98% B for 3 min, and 98% B for 3 min. The orbitrap and FAIMS were operated in positive ion mode with a positive ion voltage of 1800V; with an ion transfer tube temperature of 270°C; using a standard FAIMS resolution and compensation voltage of -50 and -65V, an inner and outer electrode temperature of 100°C with 4.5 mL/min carrier gas. Full scan spectra were acquired in profile mode at a resolution of 60,000, with a scan range of 350-1400 m/z, 300% AGC target, 25 msec maximum injection time, intensity threshold of 5×10^3, 2–6 charge state, dynamic exclusion of 20 sec, 15 data dependent scans (DDA Top 15), and mass tolerance of 10 ppm. MS2 spectra were generated an HCD collision energy of 30 at a resolution of 15,000, first mass at 110 m/z, with an isolation window of 1.3 m/z, and an automatically determined AGC target and maximum injection time in standard and auto mode.

Immunoprecipitation MS analysis

Raw files were analyzed in Proteome Discoverer 2.4 (Thermo Fisher Scientific) to generate peak lists and protein and peptides identifications using Sequest HT (Thermo Fisher Scientific) and the ToxoDB release 49 GT1 protein database. The maximum missed cleavage sites for trypsin was limited to 2. The following modifications were included in the search: dynamic oxidation (+15.995 Da; M), dynamic phosphorylation (+79.966 Da; S,T,Y), dynamic acetylation (+42.011 Da; N-terminus), and static methylthio (+45.988 Da; C). Label free quantification of proteins was performed using summed abundances from unique peptides. Abundances were normalized on the total peptide amount. Pairwise ratios were calculated for protein abundances comparing strains and conditions. Significance values were derived from *t*-tests across three replicates and adjusted with Benjamini-Hochberg correction. Exported protein abundance files from Proteome Discoverer 2.4 were loaded into R (version 4.1.1).

Proximity labeling FTS-TurboID

Parasite harvest and treatment

FTS-TurboID and mNG-TurboID (cytosolic control) parasites were infected onto confluent HFFs in 15 cm dishes. After the parasites lysed the HFF monolayer (approximately 40 hrs post-infection), extracellular parasites were passed through 5 µm filters and washed twice with DMEM by pelleting at 1000 x *g* for 10 min. At least 1×10^8 Parasites were resuspended in 200 µL of DMEM. 200 µL of biotin (final concentration 500 µM) or vehicle DMSO in DMEM was mixed with parasites. Tubes were incubated in a water bath at 37°C for 5 min. Parasites were pelleted at 12,000 x *g* for 1 min at 4°C for a total of three 1 mL PBS washes. Parasites were resuspended in a RIPA NP-40 lysis buffer (10 mM Tris-HCl pH 7.5, 140 mM NaCl, 1% NP-40 IGEPAL CA 630, 0.1% sodium deoxycholate, 0.1% SDS, 1x HALT protease inhibitor, and 125 units/mL

1392 benzonase) at a parasite concentration of 5 x 10^8 parasites/mL. Parasites were lysed at room
1393 temperature for 15 min and stored at -20°C. Results are representative of three independent
1394 experiments.
1395
Biotinylated protein enrichment
Streptavidin magnetic beads were washed three times with a RIPA NP-40 lysis buffer. Thawed
lysates were spun at 16,000 x *g* for 5 min at 4°C and the supernatant was used as the pulldown
input. 20 µL of streptavidin magnetic beads was used for 200 µL of lysates (1 x 10^8 parasites).
Beads were incubated with lysate rotating for 1 hr. Beads received a series of 1 mL washes:
twice with the RIPA NP-40 lysis buffer, once with 1 M KCl, once with 0.1M Na_2CO_3 (pH 11), once
with 2 M urea in 10 mM Tris pH 8.0), and twice with RIPA NP-40 lysis buffer. Proteins were eluted
by incubating beads with 2.5 mM biotin in S-trap sample buffer (5% SDS, 50 mM TEAB, pH 7.5)
for 10 min at 95°C.
1405
1406 *Protein cleanup and digestion*
1407 Proteins were prepared for mass spectrometry as described above in "Sub-minute
1408 phosphoproteomics - Protein cleanup and digestion". Eluted peptides were frozen in liquid
1409 nitrogen, lyophilized, and stored at -80°C until MS analysis.
1410
MS data acquisition
Lyophilized peptides were resuspended in 25 µL of 0.1% formic acid and were analyzed on an
Exploris 480 Orbitrap mass spectrometer equipped with a FAIMS Pro source (Bekker-Jensen et
al., 2020) connected to an EASY-nLC chromatography system using 0.1% formic acid as Buffer
A and 80% acetonitrile/0.1% formic acid as Buffer B. Peptides were separated at 300 nL/min on
a gradient of 2% B for 1 min, 2–25% B for 41 min, 25–40% B for 6 min, 40–100% B for 12 min,
100–2% B for 3 min, 2% B for 3 min, 2–98% B for 3 min, and 98% B for 3 min. The orbitrap and
FAIMS were operated in positive ion mode with a positive ion voltage of 1800V; with an ion
transfer tube temperature of 270°C; using a standard FAIMS resolution and compensation
voltage of -50 and -65V, an inner and outer electrode temperature of 100°C with 4.5 mL/min
carrier gas. Full scan spectra were acquired in profile mode at a resolution of 60,000, with a scan
range of 350–1400 m/z, 300% AGC target, 25 msec maximum injection time, intensity threshold
of 5 x 10^3, 2–6 charge state, dynamic exclusion of 20 sec, 15 data dependent scans (DDA Top
15), and mass tolerance of 10 ppm. MS2 spectra were generated with a HCD collision energy of
30 at a resolution of 15,000, first mass at 110 m/z, with an isolation window of 1.3 m/z, and a
normalized AGC target of 200% with an automatically determined maximum injection time.
1427
Proximity labeling MS analysis
Raw files were analyzed in Proteome Discoverer 2.4 (Thermo Fisher Scientific) to generate peak
lists and protein and peptides identifications using Sequest HT (Thermo Fisher Scientific) and
the ToxoDB release 49 GT1 protein database. The maximum missed cleavage sites for trypsin
was limited to 2. The following modifications were included in the search: dynamic oxidation
(+15.995 Da; M), dynamic phosphorylation (+79.966 Da; S,T,Y), dynamic biotinylation (+226.078
Da; K, N-terminus), dynamic acetylation (+42.011 Da; N-terminus), and static methylthio

(+45.988 Da; C). Label free quantification of proteins was performed using summed abundances
from unique peptides. Abundances were normalized on the total peptide amount. Pairwise ratios
were calculated for protein abundances comparing strains and conditions. Significance values
were derived from t-tests across three replicates and adjusted with Benjamini-Hochberg
correction. Exported protein abundance files from Proteome Discoverer 2.4 were loaded into R
(version 4.1.1).
1441

MIC2 secretion assays

Parasite harvest and treatment
Extracellular parasites were harvested in chilled DMEM and washed twice in DMEM after
centrifugation at 1000 x g for 10 min at 4°C. Parasites were resuspended at a concentration of
6 x 10^8 parasites/mL in cold DMEM. 3 x 10^7 parasites were aliquoted into round-bottom 96-well
plates. An additional aliquot of parasites was lysed in 5X Laemmli sample buffer (see
"Immunoblotting" for recipe) to obtain the total parasite lysate used to determine total MIC2
levels. Secretion was stimulated in plates with IFS and ethanol in DMEM (final concentration 3%
IFS and 1% ethanol) or a vehicle solution of DMEM. Plates were incubated by floating on a water
bath at 37°C/5% CO_2 for 90 min. Plates were spun at 1000 x g for 5 min at 4°C to separate
parasites from secreted proteins. Supernatants were transferred to a new well and spun again.
The final supernatant was mixed with a 5X Laemmli sample buffer, boiled at 95°C for 10 min,
and stored at -20°C along with total lysates until immunoblot analysis.
1455

Immunoblot analysis and quantification
To quantify MIC2 protein levels, a standard curve derived from the total parasite lysate was
generated alongside supernatants containing secreted microneme proteins. The standard curve
was derived from 3-fold serial dilutions of the total parasite lysate (undiluted, 1:3, 1:9, and 1:27)
in a 1X Laemmli sample buffer. The serial dilutions and supernatants of auxin- and vehicle-
treated parasites of a single strain were loaded onto the same precast 4–15% gel (Bio-Rad).
Subsequent immunoblotting steps were performed as described in "Immunoblotting". Anti-MIC2
was used to detect total and secreted MIC2 proteins. Secreted MIC2 proteins have a lower
molecular weight due to parasite-mediated proteolytic cleavage. Anti-CDPK1 was used as a
loading control for total parasite lysate and to reveal any parasite lysis or carry over in the
supernatant. Immunoblots confirming the depletion of tagged proteins were also collected.
Imaging was performed on a LI-COR Odyssey at high resolution for quantification.
1468

Immunoblot quantification was performed in Fiji on unadjusted inverted images. For a single
immunoblot, lane profiles of uniform dimensions were generated for MIC2 signal. Background-
subtracted signal intensity was measured as an area using the line and wand tool. A standard
curve was derived from the quantified MIC2 signal of the dilution series. Standard curves across
all conditions and replicates had R^2 values greater than 0.90. Secreted MIC2 signals were within
the linear range of the standard curve and were used to calculate the percent of total MIC2
secreted. Results are representative of three independent experiments for each parasite strain.
1476

Microneme relocalization

Parasites expressing the CLAMP-mNG reporter were grown in HFFs in glass-bottomed 35mm dishes (Ibidi and Mattek) for approximately 20 hrs. For 3-MB-PP1 treatment, media was exchanged 30 min prior to live microscopy for 3% IFS in Ringer's buffer (155 mM NaCl, 2mM $CaCl_2$, 3mM KCl, 1mM $MgCl_2$, 3mM NaH_2PO_4, 10mM HEPES, 10 mM glucose, pH 7.4) containing either 3 µM 3-MB-PP1 or vehicle solution of DMSO. At approximately 1 min after beginning live microscopy, parasites were stimulated with zaprinast (500 µM final concentration) or a vehicle solution of DMSO in corresponding Ringer's buffer. For TIR1/CLAMP-mNG and AID-HOOK/CLAMP-mNG parasite, media was exchanged 2 hrs after infection for 10% IFS media containing either 50 µM auxin or vehicle solution of PBS. Media was exchanged for 3% IFS in Ringer's buffer just prior to live microscopy. At approximately 30 sec after beginning live microscopy, parasites were stimulated with zaprinast (500 µM final concentration) or a vehicle solution of DMSO in corresponding Ringer's buffer. Images were recorded every 5–7 sec until egress or approximately 5 min using a Nikon Ti Eclipse with an enclosure maintained at 37°C.

Microneme relocalization was quantified using built-in commands from ImageJ (v. 1.53e). To determine the distribution of fluorescent micronemes at a specific time frame, a line was drawn from the parasite's apical end to its basal end. The "plot profile" command was applied to this line to calculate fluorescence intensity at regularly spaced intervals (0.13 microns) across the parasite on unadjusted images. This process was repeated for each time frame of interest and for each parasite in the analyzed vacuole. Regression analysis was performed on relocalization data from each time frame, resulting in logarithmic regression plots that display how fluorescence intensity correlates with distance from the parasite's apical end. Microneme relocalization is shown as SuperPlots to simultaneously visualize the median relocalization of an entire vacuole and the relocalization of individual parasites within each vacuole (Lord et al., 2020).

Microneme localization in extracellular parasites

Microneme localization in extracellular parasites was quantified using a custom-built, image analysis pipeline. ImageJ's built-in commands were used to measure the length of the parasite's major axis. Based on the patterns of vehicle-treated TIR1/CLAMP-mNG data, 1/8th of the parasite closest to the apical end was deemed the apical region. The remaining region was deemed the body. Using the Interactive Learning and Segmentation Toolkit (ilastik v. 1.3.3) (Berg et al., 2019), the apical region and body of parasites were isolated from each other and from the surrounding media. Ilastik's pixel and object quantification tools were then used to determine the total intensity of fluorescence in each of these regions. Comparing these values to each other yielded the percentage of fluorescent micronemes present in each of these sections in relation to the total fluorescence present in the parasite. This process was repeated for several extracellular parasites in each of the conditions tested.

ACKNOWLEDGEMENTS

We thank all the members of the Lourido laboratory for helpful discussions and support, especially Tyler Smith for aiding with the tagging vector design and Christopher Giuliano for insights during data analysis. We thank Forest M. White for generous support in developing the mass spectrometry methodologies used in this study and for helpful discussions. For data in Figure 2, we thank Matthew Child and Matt Bogyo for the CDPK1 antibody, Peter Bradley for the GAP45 antibody, John Boothroyd for the SAG1 antibody. We thank Marc-Jan Gubbels for the alpha-tubulin antibody, L. David Sibley for the MIC2, SAG1, and ALD antibody, Peter Bradley for the ROP1 antibody, Drew Etheridge for the GAP45 antibody, and Dominique Soldati-Favre for the GAP45 antibody. This work relied on VEuPathDB.org and we thank all contributors to this resource. This research was supported by funds from National Institutes of Health grants to SL (R01AI144369) and MB and MT (R01AI123457), a National Science Foundation Graduate Research Fellowship to AWC and ALH (174530). MT received funding from the Francis Crick Institute which receives its core funding from Cancer Research UK (CC2132), the UK Medical Research Council (CC2132), and the Wellcome Trust (CC2132). The Science Technology Proteomics Platform at the Francis Crick Institute received funding from Cancer Research UK (CC0199), the UK Medical Research Council (CC0199), and the Wellcome Trust (CC0199). For the purpose of Open Access, the authors have applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

COMPETING INTERESTS

The authors declare no competing interests.

1541
Figure 1. Identifying CDPK1-dependent phosphorylation with sub-minute resolution. (A)
Stimulating parasites with zaprinast triggers Ca^{2+}-mediated activation of CDPK1, resulting in the
secretion of microneme proteins (red) required for motile stages of the parasite. Conditional
knockdown (cKD) of CDPK1 endogenously tagged with mNeonGreen-mAID-Ty (green) after

auxin treatment. **(B)** Flow cytometry of mNeonGreen (mNG) fluorescence in extracellular CDPK1
cKD or parental TIR1 parasites treated with vehicle or auxin for 3.5 hrs. **(C)** Schematic of
phosphoproteomic time course. Parasites were harvested prior to CDPK1 depletion with auxin
for 3.5 hrs, followed by stimulation with zaprinast or vehicle (DMSO). Samples were collected at
0, 9, 30, and 300 sec. The experiment was performed in biological replicates. Samples were
labeled with TMTpro, pooled for analysis, and phosphopeptides were enriched using SMOAC
prior to LC-MS/MS. Four sets of samples were generated: enriched phosphoproteomes for
zaprinast [1] and DMSO [3], and proteomes for zaprinast [2] and DMSO [4]. Mock reporter ion
intensities enabling relative quantification for a given peptide are shown to illustrate fold-change
of unique phosphopeptide abundances during zaprinast stimulation. CDPK1-dependent
phosphorylation is determined by calculating the area under the curve (AUC) difference between
vehicle and auxin treatment conditions. **(D)** Protein abundances in the zaprinast proteome set
[2] at 300 sec comparing vehicle- and auxin-treated CDPK1 cKD parasites. **(E)** UpSet plot for
the number of phosphopeptides identified in the enriched zaprinast phosphoproteome [1] across
individual replicates. Phosphopeptides exhibiting CDPK1-dependent phosphorylation with $p <$
0.05 are indicated. **(F)** Scatter plot of $AUC_{difference}$ values of enriched zaprinast phosphopeptides
[1] across biological replicates. Significance was determined by comparing the distribution of
$AUC_{difference}$ values from zaprinast phosphopeptides [1] to a null distribution of DMSO
phosphopeptides [3] for individual replicates. **(G)** CDPK1-dependent and zaprinast-dependent
phosphopeptide abundances over time. Ratios of zaprinast-treated samples relative to the
vehicle-treated (no auxin) $t = 0$ samples. Median ratios of a group (solid lines). Individual
phosphopeptides (opaque lines). CDPK1-dependent phosphopeptides (Group A) determined as
described in F. Zaprinast-dependent phosphopeptides (Groups B, C, and D) were determined
by comparing the distribution of $AUC_{vehicle}$ values from zaprinast phosphopeptides [1] to a null
distribution of DMSO phosphopeptides [3]. Groups were determined by projection-based
clustering. **(H)** GO terms enriched among phosphopeptides undergoing a significant change
after zaprinast stimulation. Significance was determined using a hypergeometric test.

1573
Figure 2. Myristoylation modulates CDPK1 activity and alters its interacting partners. (A)
Complementation strategy used to evaluate the functional importance of CDPK1 myristoylation.
See Figure 2—figure supplement 1 for the construction of the iKD line. **(B)** Immunoblot
demonstrating the auxin-dependent depletion of endogenous CDPK1 in the iKD, cWT, and cMut
parasites (Myc) as well as equivalent expression of the complements (HA). *T. gondii* (toxo)
antibody was used as a loading control. **(C)** Biochemical validation of complemented lines by
YnMyr-dependent pull down. Enrichment of WT and Mut complements (HA). The inducible
endogenous CDPK1 (Myc) and *T. gondii* (toxo) antibody was used as enrichment and loading
controls, respectively. **(D)** Localization of the complemented versions of CDPK1 and
corresponding cytosolic reporters within cWT (GFP) and cMut (mCherry) by
immunofluorescence. **(E)** Myristoylation-dependent subcellular partitioning of CDPK1.
Localization of YnMyr-enriched CDPK1 was evaluated using differential centrifugation. The
partitioning into pellet [P] and supernatant [S] fractions was detected by immunoblot (CDPK1)
and compared to doubly acylated GAP45. GFP and SAG1 were used as S and P controls,
respectively. As only half of the supernatant fraction was removed from the high-speed pellet
(100,000 x *g*), the GFP signal is present in the latter. **(F)** Partitioning of complemented WT and
mutant CDPK1 after high speed centrifugation (HA). *T. gondii* (toxo) antibody was used as a P

control whereas GFP and mCherry were used as S controls for cWT and cMut, respectively. **(G)** Plaque assays demonstrating that myristoylation of CDPK1 is important for the lytic cycle of *T. gondii*. **(H)** Lack of CDPK1 myristoylation delays ionophore-induced egress from host cells. Each data point is an average of $n = 3$ biological replicates, error bars represent standard deviation. Significance calculated using 1-way ANOVA with Tukey's multiple comparison test. See Figure 2—figure supplement 1 for vehicle controls. **(I)** Immunoprecipitation-MS (IP-MS) of CDPK1-HA in cWT, cMut, and untagged TIR1 parasites across $n = 2$ biological replicates. Significantly enriched proteins (red) for proteins with more than three unique peptides, $p_{cWT} < 0.05$, and $p_{cMut} < 0.05$, and $\log_2$ fold-change > 1 across both pull-downs; t-tests and Benjamini-Hochberg corrected. **(J)** IP-MS fold-enrichment comparing cWT and cMut pull-downs. Significantly enriched proteins (red) for proteins with more than 3 unique peptides, $p < 0.05$, and $\log_2$ fold-change > 1 or < -1; t-test and Benjamini-Hochberg corrected.

1603
Figure 3. Identifying the direct substrates of CDPK1. (A) Schematic describing a strategy to
identify direct substrates of CDPK1. WT (CDPK1[G]) and mutant (G128M; CDPK1[M]) parasites were
grown in SILAC media for multiplexed quantitation. Extracellular parasites were semi-
permeabilized with aerolysin, enabling diffusion of small molecules but not proteins. CDPK1
substrate labeling was initiated by treating semi-permeabilized parasites with Ca[2+], KTPγS, ATP,
and 1B7. While CDPK1 in both WT and mutant parasites can utilize ATP to phosphorylate

substrates, only WT parasites can use KTPγS to thiophosphorylate substrates.
Thiophosphorylated peptides were specifically enriched and the remaining flow-through was
saved for whole proteome analysis. Enriched and whole proteome samples were analyzed by
LC-MS/MS. **(B)** 1B7 nanobody treatment inhibits non-specific extracellular kinase activity of
CDPK1. Thiophosphorylated substrates were detected in lysates using an anti-thiophosphate
ester antibody immunoblot. Extracellular CDPK1 activity (lane 1) was blocked by 1B7 (lane 2).
Aerolysin treatment resulted in intracellular labeling (lane 5) that was unaffected by 1B7 (lane 6).
(C) Thiophosphorylation performed in aerolysin-treated parasites comparing WT (CDPK1^G) and
mutant (CDPK1^M) strains. Detection was performed as in B. Tubulin was used as a loading
control. **(D)** Heatmap quantification of peptides using LC-MS/MS. Fold-change of peptide
abundance shown as a ratio of WT (CDPK1^G) to mutant (CDPK1^M) abundances. Experiment was
performed in n = 3 biological replicates. **(E)** Abundances of unique peptides after
thiophosphorylation in CDPK1^G and CDPK1^M parasites across n = 3 biological replicates.
Significantly enriched phosphorylated peptides are colored in red (-$\log_{10}(p)$*fold-change > 4),
one-tailed t-test. **(F)** GO terms enriched among significant phosphopeptides from E. Significance
was determined using a hypergeometric test. **(G)** Putative targets of CDPK1 determined by sub-
minute phosphoproteomics and thiophosphorylation of direct substrates. For a given CDPK1
target gene, the presence of a unique peptide phosphorylated in a CDPK1-dependent manner
(column 1) is indicated if identified in the time course (green) and/or thiophosphorylation
(magenta). The presence of additional unique phosphorylated peptides exhibiting zaprinast-
dependent effects (column 2) is indicated if the peptide was phosphorylated (orange) or
dephosphorylated (blue). Numbered boxes indicate multiple unique peptides. Fitness scores
(column 3) obtained from genome-wide KO screen data (blues). Lower scores indicate gene is
required for lytic stages of the parasite. Gene names (left), TGGT1 gene IDs (right). Gene names
with asterisks (*) are associated with published data. **(H)** Signaling diagram describing parasite
motility. Proteins exhibiting CDPK1-dependent phosphorylation by either sub-minute
phosphoproteomics or thiophosphorylation are indicated (green). Proteins exhibiting CDPK1-
independent phosphorylation (red) or dephosphorylation (blue) are indicated.
1638

1639
Figure 4. HOOK is required for host cell invasion, but dispensable for egress. (A) Schematic
of *T. gondii* and *H. sapiens* HOOK protein domains. HOOK domain (blue), coiled-coil domain
(yellow), sites phosphorylated by CDPK1 (red). **(B)** Immunoblot of HOOK conditional knockdown
parasites (AID-HOOK) after auxin treatment for 40 hrs compared to untagged TIR1 parasites.
CDPK1 was used as a loading control. **(C)** AID-HOOK is visualized in fixed intracellular parasites
by immunofluorescence after auxin treatment for 24 hrs. Hoechst and MIC2 are used as
counterstains. **(D)** Plaque assays of host cells infected with TIR1 or AID-HOOK parasites for 8
days in auxin. Host cells are stained with crystal violet. **(E)** Replication assays of host cells
infected with TIR1 or AID-HOOK parasites in auxin for 24 hrs. Parasites per vacuole were

quantified from immunofluorescence on fixed intracellular parasites. $p > 0.9$. Two-way ANOVA.
(F) Invasion assays of untagged TIR1, CDPK1-AID, and AID-HOOK parasites treated auxin for
40 hrs. Medians are plotted for $n = 3$ biological replicates (different shades of gray); n.s., $p >$
0.05, Welch's t-test. **(G)** Parasite egress stimulated with zaprinast following treatment with auxin
for 24 hrs. Egress was monitored by live microscopy. Percent egress plotted for $n = 3$ biological
replicates, n.s., $p > 0.05$, Welch's t-test. **(H)** HOOK tagged with a C-terminal 3xHA in CDPK1
cKD parasites (CDPK1-AID) visualized in fixed intracellular parasites by immunofluorescence as
in D.
1657

1658
Figure 5. The HOOK complex is required for microneme exocytosis. (A) IP-MS of HOOK-
3xHA or untagged parasites. Protein abundances determined by LC-MS/MS are shown for *n* =
3 biological replicates. Significantly enriched proteins (red) based on more than 3 unique
peptides and *p* < 0.05; ANOVA and Benjamini-Hochberg corrected. **(B)** FTS-3xHA visualized in
fixed intracellular parasites by immunofluorescence after treatment with auxin for 24 hrs.
Hoechst and MIC2 are used as counterstains. **(C)** Reciprocal IP-MS of HOOK-3xHA and FTS-
3xHA. FTS is tagged with a C-terminal 3xHA epitope at the endogenous locus (FTS-3xHA). IP
enrichment is shown as the fold-change of protein abundances in tagged versus untagged

strains determined by LC-MS/MS across $n = 3$ biological replicates. Significantly enriched proteins (red), for proteins with more than 3 unique peptides, $p_{HOOK} < 0.05$, and $p_{FTS} < 0.05$; ANOVA and Benjamini-Hochberg corrected. **(D)** Immunoblot of FTS cKD parasites. FTS is tagged with an C-terminal mAID-HA at its endogenous locus (FTS-AID) and treated with auxin for 40 hrs. ALD is used as a loading control. **(E)** Plaque assays of host cells infected with TIR1 or FTS-AID parasites for 8 days in auxin. Host cells are stained with crystal violet. **(F)** Micronemes are visualized in fixed intracellular FTS-AID and TIR1 parasites by immunofluorescence after treatment auxin for 24 hrs. Hoechst and GAP45 are used as counterstains. **(G)** Invasion assays of untagged TIR1, AID-HOOK, and FTS-AID parasites treated auxin for 40 hrs. Medians are plotted for $n = 3$ biological replicates (different shades of gray), n.s., $p > 0.05$, Welch's t-test. **(H)** Parasite egress stimulated zaprinast following auxin treatment for 24 hrs. Egress was monitored by live microscopy. Percent egress plotted for $n = 3$ biological replicates, n.s., $p > 0.05$, Welch's t-test. **(I)** Proximity labeling MS of FTS using TurboID (FTS-TurboID) compared to a cytosolic TurboID control (cytosolic mNeonGreen-TurboID). Protein abundances determined by LC-MS/MS are shown for $n = 3$ biological replicates. Significantly enriched proteins in FTS-TurboID are colored in red (red and blue), unique peptides > 3, ratio > 1, $p < 0.05$, ANOVA and Benjamini-Hochberg corrected. **(J)** Microneme protein secretion assays of parasites treated with auxin for 40 hrs. Extracellular parasites are stimulated with 1% ethanol (EtOH) and 3% IFS for 1.5 hrs. Percent MIC2 secreted is plotted for $n = 3$ biological replicates, n.s., $p > 0.05$, Welch's t-test.

Figure 6. CDPK1 activity and HOOK are required for microneme trafficking during parasite motility stages. (A) Schematic to analyze microneme trafficking during parasite motile stages. Intracellular parasites expressing microneme protein CLAMP endogenously tagged with mNeonGreen (CLAMP-mNG). Parasites are treated either with 3-MB-PP1 (inhibit CDPK1) or auxin (for conditional knockdown). Live microscopy was performed to detect CLAMP-mNG signal over time. Zaprinast was added at 1 min or 30 sec to stimulate microneme relocalization

to the apical end of the parasite. Fluorescence intensities across the apical-basal axis of each individual parasite within a vacuole was measured across time. Microneme relocalization was quantified by calculating the difference of maximum CLAMP intensity between time points preceding drug addition and egress. **(B)** Maximum intensity projections at single time points of CLAMP-mNG parasites treated with 3 μM 3-MB-PP1 or vehicle and zaprinast. **(C)** Relative fluorescence intensity of CLAMP-mNG signal across the apical-basal axis of parasites in B. Zaprinast (red) or vehicle (blue). Splines mean intensity for all parasites in each vacuole are shown with SD shaded. **(D)** Microneme relocalization. SuperPlots showing vacuole median peak differences are displayed as triangles. Individual parasites are displayed as circles. Replicates are differentially shaded, n.s., $p > 0.05$, unpaired t-test. **(E)** Maximum intensity projections at single time points of TIR1/CLAMP-mNG parasites treated with auxin and stimulated with zaprinast. **(F)** Microneme relocalization was quantified for TIR1/CLAMP-mNG parasites as in D. **(G)** Maximum intensity projections at single time points of AID-HOOK/CLAMP-mNG parasites treated auxin and stimulated with zaprinast. **(H)** Microneme relocalization was quantified for AID-HOOK/CLAMP-mNG parasites as in D. **(I)** Maximum intensity projections of extracellular TIR1/CLAMP-mNG and AID-HOOK/CLAMP-mNG parasites. **(J)** Percent of extracellular parasites in I with WT CLAMP-mNG localization, n.s., $p > 0.05$, Welch's t-test. **(K)** Percent total CLAMP-mNG signal intensity in the apical versus body of extracellular parasites, n.s., $p > 0.05$, Welch's t-test.

1714
Figure 1—figure supplement 1. Zaprinast-dependent phosphoproteome. (A) Distribution of
$AUC_{difference}$ values for individual peptides in the enriched DMSO phosphoproteome and enriched
zaprinast phosphoproteome across two biological replicates. **(B)** Heatmap of zaprinast-
dependent phosphopeptide abundance ratios across time relative to the vehicle t_0 interval during
auxin or vehicle treatment. Peptides are organized by CDPK1-dependent phosphopeptides
(Group A) and CDPK1-independent phosphopeptides (Group B, C, D).
1721

1722

Figure 2—figure supplement 1. CDPK1 myristoylation, inducible knockdown, and complementation. (A) YnMyr-dependent pull down confirming myristoylation of CDPK1. GRA2 antibody was used as a loading control. **(B)** MS2 fragmentation spectra indicating myristoylation of Gly2 of CDPK1 after YnMyr-dependent pull down. **(C)** PCR analysis confirming correct integration of the mAID cassette at the C terminus of endogenous *CDPK1* in the TIR1 line. **(D)** Immunoblot validation of auxin-dependent depletion of CDPK1 in the iKD line using the anti-Myc antibody and the anti-toxo antibody as a loading control. The band at 75 kDa represents anti-Myc-related background. **(E)** Conditional depletion of CDPK1 abolishes ionophore-induced egress from host cells. Intracellular parasites were treated with auxin or vehicle (EtOH) for 2 hrs and egress was initiated by addition of 8 µM A23187. The number of intact vacuoles was monitored over the course of 6 min. Each data point is an average of two biological replicates, each in technical triplicate, error bars represent standard deviation. **(F)** PCR analysis confirming correct integration of the complementation constructs encoding the WT (cWT) and myristoylation mutant (cMut) copies of *CDPK1* at the *UPRT* locus of the iKD line. Primers are indicated by arrows. Base pairs (bp). **(G)** In the absence of auxin, both cWT and cMut parasites egress from host cells within 2 min post-ionophore treatment.

1739
Figure 3—figure supplement 1. Factors controlling parasite motility. (A) Expanded list of factors involved in parasite motile stages from Fig. 3H.
1742

1743
1744 **Figure 4—figure supplement 1. Extended analysis of HOOK knockdown. (A)** PCR analysis
1745 confirming correct integration of the HA-mAID cassette at the N terminus of endogenous *HOOK*
1746 (*TGGT1_289100*) in the TIR1 line. **(B)** Rhoptries (ROP1) are visualized in fixed intracellular
1747 parasites by immunofluorescence after treatment with auxin for 24 hrs. Hoechst and GAP45 are
1748 used as counterstains.
1749

Figure 5—figure supplement 1. Extended analysis of FTS knockdown, proximity labeling, and microneme protein secretion. (A) PCR analysis confirming correct integration of the mAID-HA cassette at the C terminus of endogenous *FTS* (*TGGT1_264050*) in the TIR1 line. **(B)** Uncropped immunoblot shown in Figure 5D confirming C-terminal tagging of FTS. The band at ~70 kDa represents anti-HA/anti-ALD-related background present in all conditions. **©** Replication assays of host cells infected with TIR1 or FTS-AID parasites in auxin for 24 hrs. Parasites per vacuole were quantified from immunofluorescence on fixed intracellular parasites. $p > 0.9$. Two-way ANOVA. **(D)** Live microscopy of HFFs infected with parasites expressing cytosolic mNeonGreen-TurboID as the cytosolic control for proximity labeling. **(E)** PCR analysis confirming presence of TurboID-Ty cassette in the TIR1 line. **(F)** Immunoblot detection of biotinylated proteins in FTS-TurboID and cytosolic mNG-TurboID parasites treated with 500 µM of biotin or a vehicle of DMSO. Biotinylated proteins detected with a labeled streptavidin. anti-CDPK1 antibody was used as a loading control. **(G)** Serial dilution of total parasite lysate for cKD

1764   strains for TIR1, CDPK1, HOOK, and FTS used in microneme protein secretion assays in Figure
1765   5J to generate standard curves.
1766

1767

Figure 6—figure supplement 1. Extended analysis of FTS knockdown, proximity labeling, and microneme protein secretion. (A) PCR analysis confirming correct integration of the mNeonGreen reporter at the C terminus of endogenous *CLAMP* (*TGGT1_265790*) in the TIR1 line. **(B)** PCR analysis confirming correct integration of the mNeonGreen reporter at the C terminus of endogenous *CLAMP* in the AID-HOOK line.

1773

SUPPLEMENTARY FILES

Video 1. Representative image series of parasites expressing endogenously tagged CLAMP-mNG following pre-treatment with DMSO and stimulation with DMSO.

Video 2. Representative image series of parasites expressing endogenously tagged CLAMP-mNG following pre-treatment with DMSO and stimulation with 500 µM zaprinast.

Video 3. Representative image series of parasites expressing endogenously tagged CLAMP-mNG following pre-treatment with 3 µM 3-MB-PP1 and stimulation with 500 µM zaprinast.

Video 4. Representative image series of parasites expressing endogenously tagged TIR1/CLAMP-mNG following pre-treatment with vehicle and stimulation with DMSO.

Video 5. Representative image series of parasites expressing endogenously tagged TIR1/CLAMP-mNG following pretreatment with vehicle and stimulation with 500 µM zaprinast.

Video 6. Representative image series of parasites expressing endogenously tagged TIR1/CLAMP-mNG following pre-treatment with auxin and stimulation with 500 µM zaprinast.

Video 7. Representative image series of parasites expressing endogenously tagged AID-HOOK/CLAMP-mNG following pre-treatment with vehicle and stimulation with DMSO.

Video 8. Representative image series of parasites expressing endogenously tagged AID-HOOK/CLAMP-mNG following pretreatment with vehicle and stimulation with 500 µM zaprinast.

Video 9. Representative image series of parasites expressing endogenously tagged AID-HOOK/CLAMP-mNG following pre-treatment with auxin and stimulation with 500 µM zaprinast.

Supplementary file 1. Sub-minute phosphoproteomics time course peptide and abundance assignments from Proteome Discoverer 2.4 and analysis for the zaprinast enriched samples.

      Supplementary file 1.1. RAW peptide and abundance assignments.

      Supplementary file 1.2. Replicate 1 analysis with $AUC_{difference}$ and p values.

      Supplementary file 1.3. Replicate 2 analysis with $AUC_{difference}$ and p values.

      Supplementary file 1.4. Replicate 1 and 2 analysis with $AUC_{vehicle}$ values for CDPK1-independent phosphorylation.

      Supplementary file 1.5. Group assignments for zaprinast-dependent phosphopeptides.

Supplementary file 2. Sub-minute phosphoproteomics time course protein and abundance assignments from Proteome Discoverer 2.4 for the zaprinast proteome.

Supplementary file 3. Sub-minute phosphoproteomics time course peptide and abundance assignments from Proteome Discoverer 2.4 and analysis for the DMSO enriched samples.

      Supplementary file 3.1. RAW peptide and abundance assignments.

      Supplementary file 3.2. Replicate 1 analysis with $AUC_{difference}$ values.

      Supplementary file 3.3. Replicate 2 analysis with $AUC_{difference}$ values.

      Supplementary file 3.4. Replicate 1 and 2 analysis with $AUC_{vehicle}$ values for CDPK1-independent phosphorylation.

Supplementary file 4. Sub-minute phosphoproteomics time course protein and abundance assignments from Proteome Discoverer 2.4 for the DMSO proteome.

Supplementary file 5. CDPK1 immunoprecipitation protein and abundance assignments from Scaffold DIA.

Supplementary file 6. Thiophosphorylation of CDPK1 substrates. Peptide and abundance assignments from Proteome Discoverer 2.4 and analysis.

      Supplementary file 6.1. RAW peptide and abundance assignments.

Supplementary file 6.2. Analyzed peptides with enrichment significance values.

Supplementary file 6.3. Analyzed flowthrough peptides used for normalization.

Supplementary file 7. Thiophosphorylation of CDPK1 substrates. Protein and abundance

assignments from Proteome Discoverer 2.4 and analysis.

Supplementary file 8. HOOK and FTS immunoprecipitation protein and abundance

assignments from Proteome Discoverer 2.4.

Supplementary file 9. FTS proximity labeling protein and abundance assignments from

Proteome Discoverer 2.4.

Supplementary file 10. Sequences and accessions of oligonucleotides and plasmids used in

this study.

1828

SOURCE DATA

Figure 2—source data 1. This file contains source data that was used to generate the blot in

Figure 2. Myc, LICOR. Figure 2B.

Figure 2—source data 2. This file contains source data that was used to generate the blot in

Figure 2. HA, LICOR. Figure 2B.

Figure 2—source data 3. This file contains source data that was used to generate the blot in

Figure 2. toxo, LICOR. Figure 2B.

Figure 2—source data 4. This file contains source data that was used to generate the blot in

Figure 2. HA, LICOR. Figure 2C.

Figure 2—source data 5. This file contains source data that was used to generate the blot in

Figure 2. Myc, LICOR. Figure 2C.

Figure 2—source data 6. This file contains source data that was used to generate the blot in

Figure 2. toxo, LICOR. Figure 2C.

Figure 2—source data 7. This file contains source data that was used to generate the blot in

Figure 2. CDPK1, GFP, LICOR. Figure 2E.

Figure 2—source data 8. This file contains source data that was used to generate the blot in

Figure 2. GAP45, SAG1, LICOR. Figure 2E.

Figure 2—source data 9. This file contains source data that was used to generate the blot in

Figure 2. HA, LICOR. Figure 2F.

Figure 2—source data 10. This file contains source data that was used to generate the blot in

Figure 2. toxo, LICOR. Figure 2F.

Figure 2—source data 11. This file contains source data that was used to generate the blot in

Figure 2. GFP, LICOR. Figure 2F.

Figure 2—source data 12. This file contains source data that was used to generate the blot in

Figure 2. mCherry, LICOR. Figure 2F.

Figure 2—source data 13. This file contains the annotated source data that were used to

generate blots in Figure 2.

Figure 2—source data 14. This file contains source data that was used to make the graph

presented in Figure 2. GraphPad Prism data for Figure 2H.

Figure 3—source data 1. This file contains source data that was used to generate the blot in

Figure 3. Thiophosphate ester, LICOR 700 channel.

Figure 3—source data 2. This file contains source data that was used to generate the blot in

Figure 3. Thiophosphate ester, LICOR 700 channel.

Figure 3—source data 3. This file contains source data that was used to generate the blot in

Figure 3. TUB1, LICOR 800 channel.

Figure 3—source data 4. This file contains the annotated source data that were used to
generate blots in Figure 3.
Figure 4—source data 1. This file contains source data that was used to generate the blot in
Figure 4. HA-AID-HOOK, chemiluminescence.
Figure 4—source data 2. This file contains source data that was used to generate the blot in
Figure 4. HA-AID-HOOK, colorimetric, ladder.
Figure 4—source data 3. This file contains source data that was used to generate the blot in
Figure 4. CDPK1, LICOR 700 channel.
Figure 4—source data 4. This file contains the annotated source data that were used to
generate blots in Figure 4.
Figure 4—source data 5. This file contains source data that was used to make the graph
presented in Figure 4. GraphPad Prism data for Figure 4E.
Figure 4—source data 6. This file contains source data that was used to make the graph
presented in Figure 4. GraphPad Prism data for Figure 4F.
Figure 4—source data 7. This file contains source data that was used to make the graph
presented in Figure 4 and Figure 5. GraphPad Prism data for Figure 4G and 5H.
Figure 5—source data 1. This file contains source data that was used to generate the blot in
Figure 5. FTS-AID-HA, chemiluminescence.
Figure 5—source data 2. This file contains source data that was used to generate the blot in
Figure 5. FTS-AID-HA, colorimetric, ladder.
Figure 5—source data 3. This file contains source data that was used to generate the blot in
Figure 5. ALD, LICOR 700 channel.
Figure 5—source data 4. This file contains source data that was used to generate the blot in
Figure 5 and Figure 5—figure supplement 1. MIC2, LICOR 800 channel. (TIR1).
Figure 5—source data 5. This file contains source data that was used to generate the blot in
Figure 5 and Figure 5—figure supplement 1. MIC2, LICOR 800 channel. (CDPK1-AID).
Figure 5—source data 6. This file contains source data that was used to generate the blot in
Figure 5 and Figure 5—figure supplement 1. MIC2, LICOR 800 channel. (AID-HOOK).
Figure 5—source data 7. This file contains source data that was used to generate the blot in
Figure 5 and Figure 5—figure supplement 1. MIC2, LICOR 800 channel. (FTS-AID).
Figure 5—source data 8. This file contains the annotated source data that were used to
generate blots in Figure 5.
Figure 5—source data 9. This file contains source data that was used to make the graph
presented in Figure 5. GraphPad Prism data for Figure 5G.
Figure 5—source data 10. This file contains source data that was used to make the graphs
presented in Figure 5. GraphPad Prism data for Figure 5J.
Figure 6—source data 1. This file contains source data that was used to make the graph
presented in Figure 6. GraphPad Prism data for Figure 5C.
Figure 6—source data 2. This file contains source data that was used to make the graph
presented in Figure 6. GraphPad Prism data for Figure 5D.
Figure 6—source data 3. This file contains source data that was used to make the graph
presented in Figure 6. GraphPad Prism data for Figure 5F.
Figure 6—source data 4. This file contains source data that was used to make the graph
presented in Figure 6. GraphPad Prism data for Figure 5H.
Figure 6—source data 5. This file contains source data that was used to make the graph
presented in Figure 6. GraphPad Prism data for Figure 5J.
Figure 6—source data 6. This file contains source data that was used to make the graph
presented in Figure 6. GraphPad Prism data for Figure 5K.

Figure 2—figure supplement 1—source data 1. This file contains source data that was used to generate the blot in Figure 2—figure supplement 1. CDPK1, LICOR.

Figure 2—figure supplement 1—source data 2. This file contains source data that was used to generate the blot in Figure 2—figure supplement 1. GRA2, LICOR.

Figure 2—figure supplement 1—source data 3. This file contains source data that was used to generate the blot in Figure 2—figure supplement 1. Myc, LICOR.

Figure 2—figure supplement 1—source data 4. This file contains source data that was used to generate the blot in Figure 2—figure supplement 1. toxo, LICOR.

Figure 2—figure supplement 1—source data 5. This file contains source data that was used to generate the DNA gel in Figure 2—figure supplement 1. iKD integration.

Figure 2—figure supplement 1—source data 6. This file contains source data that was used to generate the DNA gel in Figure 2—figure supplement 1. cWT, cMut integration.

Figure 2—figure supplement 1—source data 7. This file contains the annotated source data that were used to generate blots in Figure 2—figure supplement 1.

Figure 2—figure supplement 1—source data 8. This file contains source data that was used to make the graph presented in Figure 2. GraphPad Prism data for Figure 2—figure supplement 1E.

Figure 2—figure supplement 1—source data 9. This file contains source data that was used to make the graph presented in Figure 2. GraphPad Prism data for Figure 2—figure supplement 1G.

Figure 4—figure supplement 1—source data 1. This file contains source data that was used to generate the DNA gel in Figure 4—figure supplement 1. AID-HOOK integration.

Figure 4—figure supplement 1—source data 2. This file contains the annotated source data that were used to generate blots in Figure 4—figure supplement 1.

Figure 5—figure supplement 1—source data 1. This file contains source data that was used to generate the DNA gel in Figure 5—figure supplement 1. FTS-AID integration.

Figure 5—figure supplement 1—source data 2. This file contains source data that was used to generate the DNA gel in Figure 5—figure supplement 1. FTS-TurboID tag.

Figure 5—figure supplement 1—source data 3. This file contains source data that was used to generate the blot in Figure 5—figure supplement 1. Biotin, LICOR 700 channel.

Figure 5—figure supplement 1—source data 4. This file contains source data that was used to generate the blot in Figure 5—figure supplement 1. CDPK1, LICOR 800 channel.

Figure 5—figure supplement 1—source data 5. This file contains source data that was used to generate the blot in Figure 5—figure supplement 1. CDPK1, LICOR 700 channel. (TIR1 standard).

Figure 5—figure supplement 1—source data 6. This file contains source data that was used to generate the blot in Figure 5—figure supplement 1. GAP45, LICOR 700 channel. (CDPK1-AID standard).

Figure 5—figure supplement 1—source data 7. This file contains source data that was used to generate the blot in Figure 5—figure supplement 1. CDPK1, LICOR 700 channel. (AID-HOOK standard).

Figure 5—figure supplement 1—source data 8. This file contains source data that was used to generate the blot in Figure 5—figure supplement 1. CDPK1, LICOR 700 channel. (FTS-AID standard).

Figure 5—figure supplement 1—source data 9. This file contains the annotated source data that were used to generate blots in Figure 5—figure supplement 1. CDPK1, LICOR 700 channel. (FTS-AID standard).

1959 **Figure 5—figure supplement 1—source data 10.** This file contains source data that was used
1960 to make the graph presented in Figure 5—figure supplement 1. GraphPad Prism data for
1961 Figure 5—figure supplement 1C.
1962 **Figure 6—figure supplement 1—source data 1.** This file contains source data that was used
1963 to generate the DNA gel in Figure 6—figure supplement 1. CLAMP-mNeonGreen integration.
1964 (TIR1/CLAMP-mNG).
1965 **Figure 6—figure supplement 1—source data 2.** This file contains source data that was used
1966 to generate the DNA gel in Figure 6—figure supplement 1. CLAMP-mNeonGreen integration.
1967 (AID-HOOK/CLAMP-mNG).
1968 **Figure 6—figure supplement 1—source data 3.** This file contains the annotated source data
1969 that were used to generate blots in Figure 6—figure supplement 1.
1970
1971 ## MATERIALS AVAILABILITY STATEMENT

1972 Proteomics data will be deposited in ProteomeXchange Consortium via the PRIDE partner
1973 repository. Accession numbers will be listed in the appropriate Methods section. The sequences
1974 of cloning vectors generated for this study have been or will be deposited in GenBank, as listed
1975 in **Supplementary File 10**. Custom analysis scripts in the R computing language are available
1976 upon request. Strains generated for this study are available upon request.

1977
REFERENCES

Alam MM, Solyakov L, Bottrill AR, Flueck C, Siddiqui FA, Singh S, Mistry S, Viskaduraki M, Lee
K, Hopp CS, Chitnis CE, Doerig C, Moon RW, Green JL, Holder AA, Baker DA, Tobin AB.
2015. Phosphoproteomics reveals malaria parasite Protein Kinase G as a signalling hub
regulating egress and invasion. *Nat Commun* **6**:7285.
Aquilini E, Cova MM, Mageswaran SK, Dos Santos Pacheco N, Sparvoli D, Penarete-Vargas
DM, Najm R, Graindorge A, Suarez C, Maynadier M, Berry-Sterkers L, Urbach S, Fahy PR,
Guérin AN, Striepen B, Dubremetz J-F, Chang Y-W, Turkewitz AP, Lebrun M. 2021. An
Alveolata secretory machinery adapted to parasite host cell invasion. *Nat Microbiol* **6**:425–
434.
Bandini G, Leon DR, Hoppe CM, Zhang Y, Agop-Nersesian C, Shears MJ, Mahal LK, Routier
FH, Costello CE, Samuelson J. 2019. O-Fucosylation of thrombospondin-like repeats is
required for processing of microneme protein 2 and for efficient host cell invasion by
Toxoplasma gondii tachyzoites. *J Biol Chem* **294**:1967–1983.
Barylyuk K, Koreny L, Ke H, Butterworth S, Crook OM, Lassadi I, Gupta V, Tromer E, Mourier
T, Stevens TJ, Breckels LM, Pain A, Lilley KS, Waller RF. 2020. A Comprehensive
Subcellular Atlas of the Toxoplasma Proteome via hyperLOPIT Provides Spatial Context
for Protein Functions. *Cell Host Microbe* **28**:752–766.e9.
Beck JR, Fung C, Straub KW, Coppens I, Vashisht AA, Wohlschlegel JA, Bradley PJ. 2013. A
Toxoplasma palmitoyl acyl transferase and the palmitoylated armadillo repeat protein
TgARO govern apical rhoptry tethering and reveal a critical role for the rhoptries in host
cell invasion but not egress. *PLoS Pathog* **9**:e1003162.
Bekker-Jensen DB, Martínez-Val A, Steigerwald S, Rüther P, Fort KL, Arrey TN, Harder A,
Makarov A, Olsen JV. 2020. A Compact Quadrupole-Orbitrap Mass Spectrometer with
FAIMS Interface Improves Proteome Coverage in Short LC Gradients. *Mol Cell Proteomics*
19:716–729.

Ben Chaabene R, Lentini G, Soldati-Favre D. 2021. Biogenesis and discharge of the rhoptries:
Key organelles for entry and hijack of host cells by the Apicomplexa. *Mol Microbiol*
115:453–465.
Berg S, Kutra D, Kroeger T, Straehle CN, Kausler BX, Haubold C, Schiegg M, Ales J, Beier T,
Rudy M, Eren K, Cervantes JI, Xu B, Beuttenmueller F, Wolny A, Zhang C, Koethe U,
Hamprecht FA, Kreshuk A. 2019. ilastik: interactive machine learning for (bio)image
analysis. *Nat Methods* **16**:1226–1232.
Bielska E, Schuster M, Roger Y, Berepiki A, Soanes DM, Talbot NJ, Steinberg G. 2014. Hook is
an adapter that coordinates kinesin-3 and dynein cargo attachment on early endosomes. *J*
Cell Biol **204**:989–1007.
Billker O, Lourido S, Sibley LD. 2009. Calcium-dependent signaling and kinases in
apicomplexan parasites. *Cell Host Microbe* **5**:612–622.
Bisio H, Lunghi M, Brochet M, Soldati-Favre D. 2019. Phosphatidic acid governs natural egress
in Toxoplasma gondii via a guanylate cyclase receptor platform. *Nat Microbiol* **4**:420–428.
Bisio H, Soldati-Favre D. 2019. Signaling Cascades Governing Entry into and Exit from Host
Cells by Toxoplasma gondii. *Annu Rev Microbiol* **73**:579–599.
Blader I, Coleman B, Chen C, Gubbels M, Hill C. 2016. The lytic cycle of Toxoplasma gondii:
15 years later, 463-485.
Blader IJ, Coleman BI, Chen C-T, Gubbels M-J. 2015. Lytic Cycle of Toxoplasma gondii: 15
Years Later. *Annu Rev Microbiol* **69**:463–485.
Blethrow JD, Glavy JS, Morgan DO, Shokat KM. 2008. Covalent capture of kinase-specific
phosphopeptides reveals Cdk1-cyclin B substrates. *Proc Natl Acad Sci U S A* **105**:1442–
1447.
Blomqvist K, Helmel M, Wang C, Absalon S, Labunska T, Rudlaff RM, Adapa S, Jiang R, Steen
H, Dvorin JD. 2020. Influence of Plasmodium falciparum Calcium-Dependent Protein
Kinase 5 (PfCDPK5) on the Late Schizont Stage Phosphoproteome. *mSphere*.
doi:10.1128/msphere.00921-19
Bradley PJ, Sibley LD. 2007. Rhoptries: an arsenal of secreted virulence factors. *Curr Opin*
Microbiol **10**:582–587.
Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, Feldman JL, Perrimon N,
Ting AY. 2018. Efficient proximity labeling in living cells and organisms with TurboID. *Nat*
Biotechnol **36**:880–887.
Breinich MS, Ferguson DJP, Foth BJ, van Dooren GG, Lebrun M, Quon DV, Striepen B,
Bradley PJ, Frischknecht F, Carruthers VB, Meissner M. 2009. A dynamin is required for
the biogenesis of secretory organelles in Toxoplasma gondii. *Curr Biol* **19**:277–286.
Brenes A, Hukelmann J, Bensaddek D, Lamond AI. 2019. Multibatch TMT Reveals False
Positives, Batch Effects and Missing Values. *Mol Cell Proteomics* **18**:1967–1980.
Brochet M, Collins MO, Smith TK, Thompson E, Sebastian S, Volkmann K, Schwach F,
Chappell L, Gomes AR, Berriman M, Rayner JC, Baker DA, Choudhary J, Billker O. 2014.
Phosphoinositide Metabolism Links cGMP-Dependent Protein Kinase G to Essential Ca2+
Signals at Key Decision Points in the Life Cycle of Malaria Parasites. *PLoS Biol*
12:e1001806.
Broncel M, Dominicus C, Vigetti L, Nofal SD, Bartlett EJ, Touquet B, Hunt A, Wallbank BA,
Federico S, Matthews S, Young JC, Tate EW, Tardieux I, Treeck M. 2020. Profiling of
myristoylation in Toxoplasma gondii reveals an N-myristoylated protein important for host
cell penetration. *Elife* **9**. doi:10.7554/eLife.57861
Brown KM, Long S, Sibley LD. 2018. Conditional Knockdown of Proteins Using Auxin-inducible
Degron (AID) Fusions in Toxoplasma gondii. *Bio Protoc* **8**. doi:10.21769/BioProtoc.2728
Brown KM, Long S, Sibley LD. 2017. Plasma Membrane Association by N-Acylation Governs

PKG Function in Toxoplasma gondii. *MBio* **8**. doi:10.1128/mBio.00375-17

Brown KM, Sibley LD. 2018. Essential cGMP Signaling in Toxoplasma Is Initiated by a Hybrid P-Type ATPase-Guanylate Cyclase. *Cell Host Microbe* **24**:804–816.e6.

Buguliskis JS, Brossier F, Shuman J, Sibley LD. 2010. Rhomboid 4 (ROM4) affects the processing of surface adhesins and facilitates host cell invasion by Toxoplasma gondii. *PLoS Pathog* **6**:e1000858.

Butler CL, Lucas O, Wuchty S, Xue B, Uversky VN, White M. 2014. Identifying novel cell cycle proteins in Apicomplexa parasites through co-expression decision analysis. *PLoS One* **9**:e97625.

Carruthers VB, Giddings OK, Sibley LD. 1999a. Secretion of micronemal proteins is associated with toxoplasma invasion of host cells. *Cell Microbiol* **1**:225–235.

Carruthers VB, Moreno SN, Sibley LD. 1999b. Ethanol and acetaldehyde elevate intracellular [Ca2+] and stimulate microneme discharge in Toxoplasma gondii. *Biochem J* **342 (Pt 2)**:379–386.

Carruthers VB, Sherman GD, Sibley LD. 2000. The Toxoplasma adhesive protein MIC2 is proteolytically processed at multiple sites by two parasite-derived proteases. *J Biol Chem* **275**:14346–14353.

Carruthers VB, Sibley LD. 1997. Sequential protein secretion from three distinct organelles of Toxoplasma gondii accompanies invasion of human fibroblasts. *Eur J Cell Biol* **73**:114–123.

Carruthers VB, Tomley FM. 2008. Microneme proteins in apicomplexans. *Subcell Biochem* **47**:33–45.

Chambers MC, Maclean B, Burke R, Amodei D, Ruderman DL, Neumann S, Gatto L, Fischer B, Pratt B, Egertson J, Hoff K, Kessner D, Tasman N, Shulman N, Frewen B, Baker TA, Brusniak M-Y, Paulse C, Creasy D, Flashner L, Kani K, Moulding C, Seymour SL, Nuwaysir LM, Lefebvre B, Kuhlmann F, Roark J, Rainer P, Detlev S, Hemenway T, Huhmer A, Langridge J, Connolly B, Chadick T, Holly K, Eckels J, Deutsch EW, Moritz RL, Katz JE, Agus DB, MacCoss M, Tabb DL, Mallick P. 2012. A cross-platform toolkit for mass spectrometry and proteomics. *Nat Biotechnol* **30**:918–920.

Chen C-T, Kelly M, Leon J de, Nwagbara B, Ebbert P, Ferguson DJP, Lowery LA, Morrissette N, Gubbels M-J. 2015. Compartmentalized Toxoplasma EB1 bundles spindle microtubules to secure accurate chromosome segregation. *Mol Biol Cell* **26**:4562–4576.

Christensen JR, Kendrick AA, Truong JB, Aguilar-Maldonado A, Adani V, Dzieciatkowska M, Reck-Peterson SL. 2021. Cytoplasmic dynein-1 cargo diversity is mediated by the combinatorial assembly of FTS-Hook-FHIP complexes. *Elife* **10**. doi:10.7554/eLife.74538

Coleman BI, Saha S, Sato S, Engelberg K, Ferguson DJP, Coppens I, Lodoen MB, Gubbels M-J. 2018. A Member of the Ferlin Calcium Sensor Family Is Essential for Toxoplasma gondii Rhoptry Secretion. *MBio* **9**. doi:10.1128/mBio.01510-18

Courjol F, Mouveaux T, Lesage K, Saliou J-M, Werkmeister E, Bonabaud M, Rohmer M, Slomianny C, Lafont F, Gissot M. 2017. Characterization of a nuclear pore protein sheds light on the roles and composition of the Toxoplasma gondii nuclear pore complex. *Cell Mol Life Sci* **74**:2107–2125.

Doggett JS, Ojo KK, Fan E, Maly DJ, Van Voorhis WC. 2014. Bumped kinase inhibitor 1294 treats established Toxoplasma gondii infection. *Antimicrob Agents Chemother* **58**:3547–3549.

Donald RGK, Zhong T, Wiersma H, Nare B, Yao D, Lee A, Allocco J, Liberator PA. 2006. Anticoccidial kinase inhibitors: identification of protein kinase targets secondary to cGMP-dependent protein kinase. *Mol Biochem Parasitol* **149**:86–98.

Endo T, Sethi KK, Piekarski G. 1982. Toxoplasma gondii: Calcium Ionophore A23187-mediated
exit of trophozoites from infected murine macrophages. *Exp Parasitol* **53**:179–188.
Fang H, Klages N, Baechler B, Hillner E, Yu L, Pardo M, Choudhary J, Brochet M. 2017.
Multiple short windows of calcium-dependent protein kinase 4 activity coordinate distinct
cell cycle events during Plasmodium gametogenesis. *Elife* **6**. doi:10.7554/eLife.26524
Farrell A, Thirugnanam S, Lorestani A, Dvorin JD, Eidell KP, Ferguson DJP, Anderson-White
BR, Duraisingh MT, Marth GT, Gubbels M-J. 2012. A DOC2 protein identified by
mutational profiling is essential for apicomplexan parasite exocytosis. *Science* **335**:218–
221.
Francia ME, Wicher S, Pace DA, Sullivan J, Moreno SNJ, Arrizabalaga G. 2011. A Toxoplasma
gondii protein with homology to intracellular type Na^+/H^+ exchangers is important for
osmoregulation and invasion. *Exp Cell Res* **317**:1382–1396.
Gaji RY, Johnson DE, Treeck M, Wang M, Hudmon A, Arrizabalaga G. 2015. Phosphorylation
of a Myosin Motor by TgCDPK3 Facilitates Rapid Initiation of Motility during Toxoplasma
gondii egress. *PLoS Pathog* **11**:e1005268.
Garrison E, Treeck M, Ehret E, Butz H, Garbuz T, Oswald BP, Settles M, Boothroyd J,
Arrizabalaga G. 2012. A forward genetic screen reveals that calcium-dependent protein
kinase 3 regulates egress in Toxoplasma. *PLoS Pathog* **8**:e1003049.
Gillingham AK, Sinka R, Torres IL, Lilley KS, Munro S. 2014. Toward a comprehensive map of
the effectors of rab GTPases. *Dev Cell* **31**:358–373.
Guo X, Farías GG, Mattera R, Bonifacino JS. 2016. Rab5 and its effector FHF contribute to
neuronal polarity through dynein-dependent retrieval of somatodendritic proteins from the
axon. *Proc Natl Acad Sci U S A* **113**:E5318–27.
Havelaar AH, Kirk MD, Torgerson PR, Gibb HJ, Hald T, Lake RJ, Praet N, Bellinger DC, de Silva
NR, Gargouri N, Speybroeck N, Cawthorne A, Mathers C, Stein C, Angulo FJ,
Devleesschauwer B, World Health Organization Foodborne Disease Burden Epidemiology
Reference Group. 2015. World Health Organization Global Estimates and Regional
Comparisons of the Burden of Foodborne Disease in 2010. *PLoS Med* **12**:e1001923.
Heal WP, Wright MH, Thinon E, Tate EW. 2011. Multifunctional protein labeling via enzymatic
N-terminal tagging and elaboration by click chemistry. *Nat Protoc* **7**:105–117.
Heaslip AT, Nishi M, Stein B, Hu K. 2011. The motility of a human parasite, Toxoplasma gondii,
is regulated by a novel lysine methyltransferase. *PLoS Pathog* **7**:e1002201.
Herneisen AL, Li Z-H, Chan AW, Moreno SNJ, Lourido S. 2022. Temporal and thermal profiling
of the Toxoplasma proteome implicates parasite Protein Phosphatase 1 in the regulation
of Ca2+-responsive pathways. *Elife* **11**. doi:10.7554/eLife.80336
Hopp CS, Bowyer PW, Baker DA. 2012. The role of cGMP signalling in regulating life cycle
progression of Plasmodium. *Microbes Infect* **14**:831–837.
Hunt A, Russell MRG, Wagener J, Kent R, Carmeille R, Peddie CJ, Collinson L, Heaslip A,
Ward GE, Treeck M. 2019. Differential requirements for cyclase-associated protein (CAP)
in actin-dependent processes of Toxoplasma gondii. *Elife* **8**. doi:10.7554/eLife.50598
Huynh M-H, Carruthers VB. 2009. Tagging of endogenous genes in a Toxoplasma gondii strain
lacking Ku80. *Eukaryot Cell* **8**:530–539.
Iacovache I, Paumard P, Scheib H, Lesieur C, Sakai N, Matile S, Parker MW, van der Goot FG.
2006. A rivet model for channel formation by aerolysin-like pore-forming toxins. *EMBO J*
25:457–466.
Ingram JR, Knockenhauer KE, Markus BM, Mandelbaum J, Ramek A, Shan Y, Shaw DE,
Schwartz TU, Ploegh HL, Lourido S. 2015. Allosteric activation of apicomplexan calcium-
dependent protein kinases. *Proc Natl Acad Sci U S A* **112**:E4975–84.

Invergo BM, Brochet M, Yu L, Choudhary J, Beltrao P, Billker O. 2017. Sub-minute
Phosphoregulation of Cell Cycle Systems during Plasmodium Gamete Formation. *Cell Rep*
21:2017–2029.
Johnson SM, Murphy RC, Geiger JA, DeRocher AE, Zhang Z, Ojo KK, Larson ET, Perera BGK,
Dale EJ, He P, Reid MC, Fox AMW, Mueller NR, Merritt EA, Fan E, Parsons M, Van Voorhis
WC, Maly DJ. 2012. Development of Toxoplasma gondii calcium-dependent protein kinase
1 (TgCDPK1) inhibitors with potent anti-toxoplasma activity. *J Med Chem* **55**:2416–2426.
Kafsack BFC, Pena JDO, Coppens I, Ravindran S, Boothroyd JC, Carruthers VB. 2009. Rapid
membrane disruption by a perforin-like protein facilitates parasite exit from host cells.
Science **323**:530–533.
Käll L, Canterbury JD, Weston J, Noble WS, MacCoss MJ. 2007. Semi-supervised learning for
peptide identification from shotgun proteomics datasets. *Nat Methods* **4**:923–925.
Käll L, Storey JD, MacCoss MJ, Noble WS. 2008a. Assigning significance to peptides identified
by tandem mass spectrometry using decoy databases. *J Proteome Res* **7**:29–34.
Käll L, Storey JD, Noble WS. 2008b. Non-parametric estimation of posterior error probabilities
associated with peptides identified by tandem mass spectrometry. *Bioinformatics* **24**:i42–
8.
Katris NJ, van Dooren GG, McMillan PJ, Hanssen E, Tilley L, Waller RF. 2014. The apical
complex provides a regulated gateway for secretion of invasion factors in Toxoplasma.
PLoS Pathog **10**:e1004074.
Kessler H, Herm-Götz A, Hegge S, Rauch M, Soldati-Favre D, Frischknecht F, Meissner M.
2008. Microneme protein 8--a new essential invasion factor in Toxoplasma gondii. *J Cell*
Sci **121**:947–956.
Krämer H, Phistry M. 1996. Mutations in the Drosophila hook gene inhibit endocytosis of the
boss transmembrane ligand into multivesicular bodies. *J Cell Biol* **133**:1205–1215.
Kremer K, Kamin D, Rittweger E, Wilkes J, Flammer H, Mahler S, Heng J, Tonkin CJ, Langsley
G, Hell SW, Carruthers VB, Ferguson DJP, Meissner M. 2013. An overexpression screen of
Toxoplasma gondii Rab-GTPases reveals distinct transport routes to the micronemes.
PLoS Pathog **9**:e1003213.
Kumar S, Kumar M, Ekka R, Dvorin JD, Paul AS, Madugundu AK, Gilberger T, Gowda H,
Duraisingh MT, Keshava Prasad TS, Sharma P. 2017. PfCDPK1 mediated signaling in
erythrocytic stages of Plasmodium falciparum. *Nat Commun* **8**:63.
Lamarque M, Besteiro S, Papoin J, Roques M, Vulliez-Le Normand B, Morlon-Guyot J,
Dubremetz J-F, Fauquenoy S, Tomavo S, Faber BW, Kocken CH, Thomas AW, Boulanger
MJ, Bentley GA, Lebrun M. 2011. The RON2-AMA1 interaction is a critical step in moving
junction-dependent invasion by apicomplexan parasites. *PLoS Pathog* **7**:e1001276.
Lamarque MH, Roques M, Kong-Hap M, Tonkin ML, Rugarabamu G, Marq J-B, Penarete-
Vargas DM, Boulanger MJ, Soldati-Favre D, Lebrun M. 2014. Plasticity and redundancy
among AMA-RON pairs ensure host cell entry of Toxoplasma parasites. *Nat Commun*
5:4098.
Lasonder E, Green JL, Camarda G, Talabani H, Holder AA, Langsley G, Alano P. 2012. The
Plasmodium falciparum schizont phosphoproteome reveals extensive phosphatidylinositol
and cAMP-protein kinase A signaling. *J Proteome Res* **11**:5323–5337.
Lee I-G, Olenick MA, Boczkowska M, Franzini-Armstrong C, Holzbaur ELF, Dominguez R.
2018. A conserved interaction of the dynein light intermediate chain with dynein-dynactin
effectors necessary for processivity. *Nat Commun* **9**:986.
Lentini G, Dubois DJ, Maco B, Soldati-Favre D, Frénal K. 2019. The roles of Centrin 2 and
Dynein Light Chain 8a in apical secretory organelles discharge of Toxoplasma gondii.

Traffic **20**:583–600.

Leung JM, He Y, Zhang F, Hwang Y-C, Nagayasu E, Liu J, Murray JM, Hu K. 2017. Stability
and function of a putative microtubule-organizing center in the human parasite
Toxoplasma gondii. *Mol Biol Cell* **28**:1361–1378.

Leung JM, Liu J, Wetzel LA, Hu K. 2019. Centrin2 from the human parasite Toxoplasma gondii
is required for its invasion and intracellular replication. *J Cell Sci* **132**.
doi:10.1242/jcs.228791

Li J, Van Vranken JG, Pontano Vaites L, Schweppe DK, Huttlin EL, Etienne C, Nandhikonda P,
Viner R, Robitaille AM, Thompson AH, Kuhn K, Pike I, Bomgarden RD, Rogers JC, Gygi
SP, Paulo JA. 2020. TMTpro reagents: a set of isobaric labeling mass tags enables
simultaneous proteome-wide measurements across 16 samples. *Nat Methods* **17**:399–
404.

Long S, Anthony B, Drewry LL, David Sibley L. 2017. A conserved ankyrin repeat-containing
protein regulates conoid stability, motility and cell invasion in Toxoplasma gondii. *Nature*
Communications. doi:10.1038/s41467-017-02341-2

Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK. 2020. SuperPlots: Communicating
reproducibility and variability in cell biology. *J Cell Biol* **219**. doi:10.1083/jcb.202001064

Lourido S, Jeschke GR, Turk BE, David Sibley L. 2013a. Exploiting the Unique ATP-Binding
Pocket of *Toxoplasma* Calcium-Dependent Protein Kinase 1 To Identify Its Substrates.
ACS Chemical Biology. doi:10.1021/cb400115y

Lourido S, Moreno SNJ. 2015. The calcium signaling toolkit of the Apicomplexan parasites
Toxoplasma gondii and Plasmodium spp. *Cell Calcium* **57**:186–193.

Lourido S, Shuman J, Zhang C, Shokat KM, Hui R, Sibley LD. 2010. Calcium-dependent
protein kinase 1 is an essential regulator of exocytosis in Toxoplasma. *Nature* **465**:359–
362.

Lourido S, Tang K, Sibley LD. 2012. Distinct signalling pathways control Toxoplasma egress
and host-cell invasion. *EMBO J* **31**:4524–4534.

Lourido S, Zhang C, Lopez MS, Tang K, Barks J, Wang Q, Wildman SA, Shokat KM, Sibley LD.
2013b. Optimizing small molecule inhibitors of calcium-dependent protein kinase 1 to
prevent infection by Toxoplasma gondii. *J Med Chem* **56**:3068–3077.

Luo S, Ruiz FA, Moreno SNJ. 2005. The acidocalcisome Ca2+-ATPase (TgA1) of Toxoplasma
gondii is required for polyphosphate storage, intracellular calcium homeostasis and
virulence. *Mol Microbiol* **55**:1034–1045.

Mageswaran SK, Guérin A, Theveny LM, Chen WD, Martinez M, Lebrun M, Striepen B, Chang
Y-W. 2021. In situ ultrastructures of two evolutionarily distant apicomplexan rhoptry
secretion systems. *Nat Commun* **12**:4983.

Markus BM, Bell GW, Lorenzi HA, Lourido S. 2019. Optimizing Systems for Cas9 Expression in
Toxoplasma gondii. *mSphere* **4**. doi:10.1128/mSphere.00386-19

Márquez-Nogueras KM, Hortua Triana MA, Chasen NM, Kuo IY, Moreno SN. 2021. Calcium
signaling through a transient receptor channel is important for Toxoplasma gondii growth.
Elife **10**. doi:10.7554/eLife.63417

Martin DDO, Beauchamp E, Berthiaume LG. 2011. Post-translational myristoylation: Fat
matters in cellular life and death. *Biochimie* **93**:18–31.

McCoy JM, Stewart RJ, Uboldi AD, Li D, Schröder J, Scott NE, Papenfuss AT, Lehane AM,
Foster LJ, Tonkin CJ. 2017. A forward genetic screen identifies a negative regulator of
rapid Ca2+-dependent cell egress (MS1) in the intracellular parasite Toxoplasma gondii. *J*
Biol Chem **292**:7662–7674.

McCoy JM, Whitehead L, van Dooren GG, Tonkin CJ. 2012. TgCDPK3 regulates calcium-

dependent egress of Toxoplasma gondii from host cells. *PLoS Pathog* **8**:e1003066.

Mueller C, Klages N, Jacot D, Santos JM, Cabrera A, Gilberger TW, Dubremetz J-F, Soldati-
Favre D. 2013. The Toxoplasma protein ARO mediates the apical positioning of rhoptry
organelles, a prerequisite for host cell invasion. *Cell Host Microbe* **13**:289–301.

Mueller C, Samoo A, Hammoudi P-M, Klages N, Kallio JP, Kursula I, Soldati-Favre D. 2016.
Structural and functional dissection of Toxoplasma gondii armadillo repeats only protein. *J*
Cell Sci **129**:1031–1045.

Nofal SD, Dominicus C, Broncel M, Katris NJ, Flynn HR, Arrizabalaga G, Botté CY, Invergo BM,
Treeck M. 2022. A positive feedback loop mediates crosstalk between calcium, cyclic
nucleotide and lipid signalling in calcium-induced Toxoplasma gondii egress. *PLoS Pathog*
18:e1010901.

Ojo KK, Larson ET, Keyloun KR, Castaneda LJ, Derocher AE, Inampudi KK, Kim JE, Arakaki
TL, Murphy RC, Zhang L, Napuli AJ, Maly DJ, Verlinde CLMJ, Buckner FS, Parsons M, Hol
WGJ, Merritt EA, Van Voorhis WC. 2010. Toxoplasma gondii calcium-dependent protein
kinase 1 is a target for selective kinase inhibitors. *Nat Struct Mol Biol* **17**:602–607.

Ong Y-C, Reese ML, Boothroyd JC. 2010. Toxoplasma Rhoptry Protein 16 (ROP16) Subverts
Host Function by Direct Tyrosine Phosphorylation of STAT6*. *J Biol Chem* **285**:28731–
28740.

Pang ZP, Südhof TC. 2010. Cell biology of Ca2+-triggered exocytosis. *Curr Opin Cell Biol*
22:496–505.

Pease BN, Huttlin EL, Jedrychowski MP, Dorin-Semblat D, Sebastiani D, Segarra DT, Roberts
BF, Chakrabarti R, Doerig C, Gygi SP, Chakrabarti D. 2018. Characterization of
Plasmodium falciparum Atypical Kinase PfPK7- Dependent Phosphoproteome. *J*
Proteome Res **17**:2112–2123.

Pease BN, Huttlin EL, Jedrychowski MP, Talevich E, Harmon J, Dillman T, Kannan N, Doerig C,
Chakrabarti R, Gygi SP, Chakrabarti D. 2013. Global Analysis of Protein Expression and
Phosphorylation of Three Stages of *Plasmodium falciparum* Intraerythrocytic Development.
Journal of Proteome Research. doi:10.1021/pr400394g

Pomel S, Luk FCY, Beckers CJM. 2008. Host cell egress and invasion induce marked
relocations of glycolytic enzymes in Toxoplasma gondii tachyzoites. *PLoS Pathog*
4:e1000188.

Redwine WB, DeSantis ME, Hollyer I, Htet ZM, Tran PT, Swanson SK, Florens L, Washburn
MP, Reck-Peterson SL. 2017. The human cytoplasmic dynein interactome reveals novel
activators of motility. *Elife* **6**. doi:10.7554/eLife.28257

Reese ML, Zeiner GM, Saeij JPJ, Boothroyd JC, Boyle JP. 2011. Polymorphic family of injected
pseudokinases is paramount in Toxoplasma virulence. *Proc Natl Acad Sci U S A*
108:9625–9630.

Rothenberg DA, Gordon EA, White FM, Lourido S. 2016. Identification of Direct Kinase
Substrates Using Analogue-Sensitive Alleles. *Methods Mol Biol* **1355**:71–84.

Sebastian S, Brochet M, Collins MO, Schwach F, Jones ML, Goulding D, Rayner JC,
Choudhary JS, Billker O. 2012. A Plasmodium calcium-dependent protein kinase controls
zygote development and transmission by translationally activating repressed mRNAs. *Cell*
Host Microbe **12**:9–19.

Segev-Zarko L-A, Dahlberg PD, Sun SY, Pelt DM, Kim CY, Egan ES, Sethian JA, Chiu W,
Boothroyd JC. 2022. Cryo-electron tomography with mixed-scale dense neural networks
reveals key steps in deployment of Toxoplasma invasion machinery. *PNAS Nexus*
1:gac183.

Shen B, Brown KM, Lee TD, David Sibley L. 2014. Efficient Gene Disruption in Diverse Strains
of Toxoplasma gondii Using CRISPR/CAS9. *mBio*. doi:10.1128/mbio.01114-14

Shortt E, Hackett CG, Stadler RV, Ward GE, Lourido S. 2022. CDPK2A and CDPK1 form a
signaling module upstream of Toxoplasma motility. *bioRxiv*.
doi:10.1101/2022.07.19.500742
Sidik SM, Hortua Triana MA, Paul AS, El Bakkouri M, Hackett CG, Tran F, Westwood NJ, Hui
R, Zuercher WJ, Duraisingh MT, Moreno SNJ, Lourido S. 2016a. Using a Genetically
Encoded Sensor to Identify Inhibitors of Toxoplasma gondii Ca2+ Signaling. *J Biol Chem*
291:9566–9580.
Sidik SM, Huet D, Ganesan SM, Huynh M-H, Wang T, Nasamu AS, Thiru P, Saeij JPJ,
Carruthers VB, Niles JC, Lourido S. 2016b. A Genome-wide CRISPR Screen in
Toxoplasma Identifies Essential Apicomplexan Genes. *Cell* **166**:1423–1435.e12.
Sidik SM, Valleau D, Acevedo-Sánchez Y, Godoy LC, Pasaje CFA, Huynh M-H, Carruthers VB,
Niles JC, Lourido S. n.d. A conserved complex of microneme proteins mediates rhoptry
discharge in*Toxoplasma*. doi:10.1101/2022.11.28.518173
Simm D, Hatje K, Kollmar M. 2015. Waggawagga: comparative visualization of coiled-coil
predictions and detection of stable single α-helices (SAH domains). *Bioinformatics* **31**:767–
769.
Singer M, Simon K, Forné I, Meissner M. 2022. A central protein complex essential for Invasion
in Toxoplasma gondii. *bioRxiv*. doi:10.1101/2022.02.24.481622
Smith TA, Lopez-Perez GS, Herneisen AL, Shortt E, Lourido S. 2022. Screening the
Toxoplasma kinome with high-throughput tagging identifies a regulator of invasion and
egress. *Nat Microbiol* **7**:868–881.
Söding J, Biegert A, Lupas AN. 2005. The HHpred interactive server for protein homology
detection and structure prediction. *Nucleic Acids Res* **33**:W244–8.
Solyakov L, Halbert J, Alam MM, Semblat J-P, Dorin-Semblat D, Reininger L, Bottrill AR, Mistry
S, Abdi A, Fennell C, Holland Z, Demarta C, Bouza Y, Sicard A, Nivez M-P, Eschenlauer S,
Lama T, Thomas DC, Sharma P, Agarwal S, Kern S, Pradel G, Graciotti M, Tobin AB,
Doerig C. 2011. Global kinomic and phospho-proteomic analyses of the human malaria
parasite Plasmodium falciparum. *Nature Communications*. doi:10.1038/ncomms1558
Sparvoli D, Delabre J, Penarete-Vargas DM, Kumar Mageswaran S, Tsypin LM, Heckendorn J,
Theveny L, Maynadier M, Mendonça Cova M, Berry-Sterkers L, Guérin A, Dubremetz J-F,
Urbach S, Striepen B, Turkewitz AP, Chang Y-W, Lebrun M. 2022. An apical membrane
complex for triggering rhoptry exocytosis and invasion in Toxoplasma. *EMBO J*
41:e111158.
Stasic AJ, Chasen NM, Dykes EJ, Vella SA, Asady B, Starai VJ, Moreno SNJ. 2019. The
Toxoplasma Vacuolar H+-ATPase Regulates Intracellular pH and Impacts the Maturation
of Essential Secretory Proteins. *Cell Rep* **27**:2132–2146.e7.
Sun SY, Segev-Zarko L-A, Chen M, Pintilie GD, Schmid MF, Ludtke SJ, Boothroyd JC, Chiu W.
2022. Cryo-ET of *Toxoplasma* parasites gives subnanometer insight into tubulin-based
structures. *Proc Natl Acad Sci U S A* **119**. doi:10.1073/pnas.2111661119
Tang Q, Andenmatten N, Hortua Triana MA, Deng B, Meissner M, Moreno SNJ, Ballif BA, Ward
GE. 2014. Calcium-dependent phosphorylation alters class XIVa myosin function in the
protozoan parasite Toxoplasma gondii. *MBoC* **25**:2579–2591.
Taus T, Köcher T, Pichler P, Paschke C, Schmidt A, Henrich C, Mechtler K. 2011. Universal
and confident phosphorylation site localization using phosphoRS. *J Proteome Res*
10:5354–5362.
Taylor HM, McRobert L, Grainger M, Sicard A, Dluzewski AR, Hopp CS, Holder AA, Baker DA.
2010. The malaria parasite cyclic GMP-dependent protein kinase plays a central role in
blood-stage schizogony. *Eukaryot Cell* **9**:37–45.
Thrun MC, Ultsch A. 2021. Using Projection-Based Clustering to Find Distance- and Density-

Based Clusters in High-Dimensional Data. *J Classification* **38**:280–312.

Tosetti N, Dos Santos Pacheco N, Soldati-Favre D, Jacot D. 2019. Three F-actin assembly centers regulate organelle inheritance, cell-cell communication and motility in Toxoplasma gondii. *Elife* **8**. doi:10.7554/eLife.42669

Treeck M, Sanders JL, Elias JE, Boothroyd JC. 2011. The phosphoproteomes of Plasmodium falciparum and Toxoplasma gondii reveal unusual adaptations within and beyond the parasites' boundaries. *Cell Host Microbe* **10**:410–419.

Treeck M, Sanders JL, Gaji RY, LaFavers KA, Child MA, Arrizabalaga G, Elias JE, Boothroyd JC. 2014. The calcium-dependent protein kinase 3 of toxoplasma influences basal calcium levels and functions beyond egress as revealed by quantitative phosphoproteome analysis. *PLoS Pathog* **10**:e1004197.

Tsai C-F, Hsu C-C, Hung J-N, Wang Y-T, Choong W-K, Zeng M-Y, Lin P-Y, Hong R-W, Sung T-Y, Chen Y-J. 2014. Sequential phosphoproteomic enrichment through complementary metal-directed immobilized metal ion affinity chromatography. *Anal Chem* **86**:685–693.

Tyler JS, Boothroyd JC. 2011. The C-terminus of Toxoplasma RON2 provides the crucial link between AMA1 and the host-associated invasion complex. *PLoS Pathog* **7**:e1001282.

Van Vranken, Pontano Vaites, Schweppe. n.d. TMTpro reagents: a set of isobaric labeling mass tags enables simultaneous proteome-wide measurements across 16 samples. *Nature*.

Wang X, Fu Y, Beatty WL, Ma M, Brown A, Sibley LD, Zhang R. 2021. Cryo-EM structure of cortical microtubules from human parasite Toxoplasma gondii identifies their microtubule inner proteins. *Nat Commun* **12**:3065.

Wernimont AK, Artz JD, Finerty P Jr, Lin Y-H, Amani M, Allali-Hassani A, Senisterra G, Vedadi M, Tempel W, Mackenzie F, Chau I, Lourido S, Sibley LD, Hui R. 2010. Structures of apicomplexan calcium-dependent protein kinases reveal mechanism of activation by calcium. *Nat Struct Mol Biol* **17**:596–601.

Wichroski MJ, Melton JA, Donahue CG, Tweten RK, Ward GE. 2002. Clostridium septicum alpha-toxin is active against the parasitic protozoan Toxoplasma gondii and targets members of the SAG family of glycosylphosphatidylinositol-anchored surface proteins. *Infect Immun* **70**:4353–4361.

Wilson SK, Heckendorn J, Martorelli Di Genova B, Koch LL, Rooney PJ, Morrissette N, Lebrun M, Knoll LJ. 2020. A Toxoplasma gondii patatin-like phospholipase contributes to host cell invasion. *PLoS Pathog* **16**:e1008650.

Winzer P, Müller J, Aguado-Martínez A, Rahman M, Balmer V, Manser V, Ortega-Mora LM, Ojo KK, Fan E, Maly DJ, Van Voorhis WC, Hemphill A. 2015. In Vitro and In Vivo Effects of the Bumped Kinase Inhibitor 1294 in the Related Cyst-Forming Apicomplexans Toxoplasma gondii and Neospora caninum. *Antimicrob Agents Chemother* **59**:6361–6374.

Wright MH, Clough B, Rackham MD, Rangachari K, Brannigan JA, Grainger M, Moss DK, Bottrill AR, Heal WP, Broncel M, Serwa RA, Brady D, Mann DJ, Leatherbarrow RJ, Tewari R, Wilkinson AJ, Holder AA, Tate EW. 2014. Validation of N-myristoyltransferase as an antimalarial drug target using an integrated chemical biology approach. *Nature Chemistry*. doi:10.1038/nchem.1830

Xu L, Sowa ME, Chen J, Li X, Gygi SP, Harper JW. 2008. An FTS/Hook/p107FHIP Complex Interacts with and Promotes Endosomal Clustering by the Homotypic Vacuolar Protein Sorting Complex. *MBoC* **19**:5059–5071.

Yang L, Uboldi AD, Seizova S, Wilde M-L, Coffey MJ, Katris NJ, Yamaryo-Botté Y, Kocan M, Bathgate RAD, Stewart RJ, McConville MJ, Thompson PE, Botté CY, Tonkin CJ. 2019. An apically located hybrid guanylate cyclase–ATPase is critical for the initiation of Ca2+ signaling and motility in Toxoplasma gondii. *J Biol Chem* **294**:8959–8972.

Yao X, Wang X, Xiang X. 2014. FHIP and FTS proteins are critical for dynein-mediated
transport of early endosomes in Aspergillus. *Mol Biol Cell* **25**:2181–2189.
Yuasa K, Mi-Ichi F, Kobayashi T, Yamanouchi M, Kotera J, Kita K, Omori K. 2005. PfPDE1, a
novel cGMP-specific phosphodiesterase from the human malaria parasite Plasmodium
falciparum. *Biochem J* **392**:221–229.
Zecha J, Satpathy S, Kanashova T, Avanessian SC, Kane MH, Clauser KR, Mertins P, Carr SA,
Kuster B. 2019. TMT Labeling for the Masses: A Robust and Cost-efficient, In-solution
Labeling Approach. *Mol Cell Proteomics* **18**:1468–1478.
Zhang J, Qiu R, Arst HN, Peñalva MA, Xiang X. 2014. HookA is a novel dynein–early endosome
linker critical for cargo movement in vivo. *J Cell Biol* **204**:1009–1026.

2402