Predicting the functional effect of compound heterozygous genotypes from large scale
variant effect maps

Michael Xie[1,2], Amy Sirr[1], Russell S. Lo[1], Gareth A. Cromie[1], Michelle Tang[1], Julee Ashmead[1],
Katherine Owens[1,3], J. Nathan Kutz[3], Richard N. McLaughlin Jr.[1] and Aimée M. Dudley[1,2]*

[1] Pacific Northwest Research Institute, Seattle, Washington, USA

[2] Molecular Engineering Graduate Program, University of Washington, Seattle, Washington, USA

[3] Department of Applied Mathematics, University of Washington, Seattle, Washington, USA

Keywords: Yeast, Serine Deficiency, Neu-Laxova Syndrome, PSAT1, Inborn Errors of Metabolism

*Corresponding Author:

Aimée M. Dudley, Ph.D.

Pacific Northwest Research Institute

720 Broadway

Seattle, WA 98122

aimee.dudley@gmail.com

Abstract

Loss of function mutations in *PHGDH*, *PSAT1*, and *PSPH* cause a set of rare, autosomal recessive diseases known as Neu-Laxova syndrome (NLS) or serine-deficiency disorders. The diseases present with a broad range of phenotypes, including lethality, severe neurological manifestations, seizures, and intellectual disability. However, because L-serine supplementation, especially if started early in pregnancy, can ameliorate and in some cases even prevent symptoms, knowledge of pathogenic variants is highly actionable. Recently, our laboratory established a yeast-based assay for human *PSAT1* function. We have now applied it at scale to assay the functional impact of 1,914 SNV-accessible amino acid substitutions. Our results agree well with clinical interpretations and protein structure-function relationships, supporting the use of our data as functional evidence under the ACMG interpretation guidelines. In addition to assaying the functional impact of individual variants in yeast haploid cells, we can assay pairwise combinations of *PSAT1* alleles that recapitulate human genotypes, including compound heterozygotes, in yeast diploids. Results from this diploid assay successfully distinguish patient genotypes from those of healthy carriers and agree well with disease severity. Finally, we present a linear model that can accurately predict biallelic function in diploids using information from the individual allele measurements in haploids. Taken together, our work provides an example of how large-scale functional assays in model systems can be powerfully applied to the study of a rare disease.

Background:

High throughput sequencing technologies have accelerated the discovery of human genetic variation. Data from the Genome Aggregation Database (gnomAD) [1] suggest that the average person harbors approximately 200 rare (allele frequency <0.1%) protein coding variants [2]. However, leveraging genomic information for the diagnosis and treatment of medically actionable diseases is currently limited by the relatively small number of alleles for which clinical interpretation is readily available. Most of the ~550K missense variants in ClinVar [3], even those in well-studied disease genes, are variants of uncertain significance (VUS). The difficulty of variant interpretation is compounded further in rare diseases, and patients often face long diagnostic odysseys or remain undiagnosed.

Inborn errors of metabolism (IEM), genetic alterations that disrupt metabolic homeostasis, are a relatively large class of rare diseases. Over 700 IEMs [4] have been described and new disorders continue to be discovered [5]. Although they are rare individually, collectively these diseases are common, with estimates suggesting that IEMs may affect 1 in 800-2600 live births [6,7]. Many are severe and manifest early in life, but some are also medically actionable and timely diagnosis can prevent the onset of irreversible damage. Disruptions of metabolic pathways can lead to toxic levels of substrate accumulation or deficiency of an essential product. Therapeutic strategies often focus on amending these imbalances. For example, dietary restriction is prescribed for the treatment of urea cycle disorders, phenylketonuria, and galactosemia, while dietary supplementation is prescribed for homocystinuria and pyridoxine-dependent epilepsy [8–10].

Serine deficiency disorders are an emerging group of IEMs with promising case studies describing successful treatment with dietary supplementation (**Figure 1**). Impairment of any of the three L-serine biosynthesis pathway enzymes, phosphoglycerate dehydrogenase (PGDH; encoded by *PHGDH*), phosphoserine aminotransferase (PSAT; encoded by *PSAT1*), and phosphoserine phosphatase (PSP; encoded by *PSPH*), result in systemic serine deficiency [11]. Serine metabolism is central to numerous biological processes, including the synthesis of proteins, nucleotides, and phospholipids, as well as the formation of the neuromodulators D-serine and glycine. As a result, serine biosynthesis defects negatively impact nervous system development and function, with clinical manifestations that include microcephaly, seizures, intellectual disability, and neuropathies. Case studies have demonstrated that oral serine supplementation can reduce and, in some cases, prevent the onset of these severe symptoms that typically manifest very early in life [12–16] Prompt diagnosis is crucial, as the impact of therapeutic intervention, including prenatal dietary supplementation, is greatest before patients become symptomatic and irreversible neurological damage occurs.

Figure 1: Key developments in the study of serine deficiency disorders, a set rare but treatable inborn errors of metabolism.

Unfortunately, like many rare diseases, diagnosing serine deficiency disorders is difficult in the absence of a family history of disease. One challenge is that the disorders display a broad phenotypic spectrum. At one extreme, Neu-Laxova syndrome (NLS) is a severe developmentally lethal congenital disease, while serine deficiency ranges from growth defects and severe neurological manifestations in infants to milder forms with intellectual disability, developmental delay, or progressive neuropathies that can manifest in children or adults [11,17]. Although it has been hypothesized that residual enzyme activity may explain the phenotypic variability [18–20] within serine deficiency disorders, functional studies regarding this subject are sparse and the natural history of the disorders are largely unknown. A second challenge is the difficulty of biochemical diagnosis, which requires measurement of amino acid levels in patient serum or cerebral spinal fluid compared to closely age-matched controls [17,21].

Identifying the presence of pathogenic sequence variants through whole exome/genome sequencing represents an attractive alternative for diagnosing serine deficiency disorders. However, this approach relies on having clinical interpretations for rare variants, which are generally not available. Instead, within the current disease literature, clinical sequencing has primarily been used as a tool by experts to provide molecular confirmation once a patient has become symptomatic. High-throughput approaches to variant interpretation that can be applied to rare variants are needed to increase the success rate of sequencing-based diagnostics. While computational prediction methods can be applied at scale, their relatively high error rates limit their utility (examples in [22]). Functional assays that can quantitatively

measure variant effects on protein function [23] are an alternative approach that can be used
as part of the criteria for variant interpretation established by the American College of Medical
Genetics (ACMG) [24]. Improvements in DNA synthesis technology have enabled time-and-cost
effective methods for building large variant libraries. When combined with high-throughput
phenotyping methods, functional assays can comprehensively assess variants that have been
identified in the human population as well as those that may arise in the future [25,26].

Recently, our laboratory established a yeast-based assay for human *PSAT1* function [27]. Here,
we apply it at scale to quantify the functional impact of 1,914 amino acid substitutions that are
accessible via a single nucleotide variant (SNV) in the human *PSAT1* coding sequence. Our
results agree well with clinically interpreted alleles and with protein structure-function
relationships, supporting the use of our data as functional evidence under the ACMG
interpretation guidelines. In addition to assaying the functional impact of single variants in
yeast haploids, we demonstrate that we can apply our assay to yeast diploids with pairwise
combinations of *PSAT1* alleles that recapitulate human genotypes. The results of the diploid
functional assay distinguish patient genotypes from those of healthy carriers and agree with the
stratification of patient genotypes by disease severity. Finally, we developed a mathematical
model that can accurately predict biallelic function in diploids from pairwise combinations of
individual allele activity measured experimentally in haploids. Taken together, our work
provides an example of how large-scale functional assays in model systems can be powerfully
applied in the study of rare disease and to inform future diagnostic efforts.

Methods

Strain library construction

All *Saccharomyces cerevisiae* strains used in this study (**Supplementary Table S1** and **S2**) were
derived from the isogenic lab strains FY4 (MAT**a**) and FY5 (MATα) [28]. Unless otherwise noted,
strains were grown in rich medium (YPD, 1% yeast extract, 2% peptone, and 2% glucose) or
minimal medium (SD, without amino acids, 2% glucose) using standard media conditions and
methods for yeast genetic manipulation [29].

Methodology concerning the design and construction of the yeast codon-optimized version
(*yPSAT1*) of human *PSAT1* isoform 1, the wild type (*yPSAT1*) strain, the deletion (*ser1Δ0*) strain,
and individual variant strains were previously described in Sirr *et al* [27]. Here, we describe the
design and construction of a variant library ~10 fold larger and modifications of the growth
assay for high throughput as follows:

The variant library was designed to capture the amino acid substitutions resulting from all SNV-
accessible missense mutations across 369 codons, excluding the start and stop, in the human
PSAT1 isoform 1 cDNA sequence (CCDS6660.1; Consensus Coding Sequence database [30]. The

9 possible single nucleotide variants at each codon resulted in 4-7 unique amino acid
substitutions. *yPSAT1* variants encoding the complete set of these unique amino acid
substitutions (n = 2,182) were synthesized (Twist Biosciences) in *yPSAT1* as an oligonucleotide
library in which each well of a 96 well plate contained all amino acid substitutions (n=4-7) at a
given amino acid.

Individual wells of these plates were amplified to approximately 500 ng of DNA by 15-cycles of
high-fidelity PCR (Phusion High-Fidelity DNA Polymerase; Thermo Scientific) and transformed
into a MAT**a** haploid deletion strain (*ser1Δ0*) using standard methods. Single colonies from the
yeast transformations were isolated such that 6,335 individual transformants, each encoding a
single amino acid substitution, were arrayed into 96-well plates containing rich medium.
Because individual transformants are isolated and maintained as separate stocks (one strain per
well in a 96-well plate), each strain is an independently constructed biological isolate of the
variant it contains. For downstream phenotype normalization, each library plate also contained
replicates of the same control strains: 2 deletion (*ser1Δ0)* and 4 wild type (*yPSAT1*) strains.

All of the original *yPSAT1* variant strains in the large-scale library are of a single mating type
(MAT**a**) (**Supplementary Table S1**). To generate diploid strains harboring combinations of
yPSAT1 variants were first introduced into a strain of the opposite mating type and the resulting
yPSAT1 MATα strains were mated in a pairwise manner to target *yPSAT1* MAT**a** strains using
standard methods [31]. The resulting diploid strains (**Supplementary Table S2**) harboring
combinations of *yPSAT1* alleles (missense variant, wild type, or null) were then arrayed in an
alternating checkerboard pattern that minimizes the influence of nutrient competition from
neighboring colonies during phenotype measurement.

Variant library sequence confirmation

For each *yPSAT1* transformant, we know which codon is mutated (target codon), but not which
specific variant is present. To determine this, we used a custom MinION (Oxford Nanopore
Technologies) sequencing pipeline, as described in Lo *et al* [32]. Briefly, individual
transformants were pooled in groups of 12, so that no target codon was represented more than
once in a single pool. The pool was then sequenced and at each target codon the most frequent
variant codon was identified (candidate variant) as well as the second most frequent variant. An
error model was then developed for each variant.

Candidate variants then underwent quality control. Variants passed if they were supported by
at least 15 reads, if the observed frequency of the variant was at least 3.3x the estimated error
frequency for that variant and if no missense or nonsense secondary mutations were present.
In addition, candidates were rejected when the second most frequent variant was both
enriched (>=10-fold) relative to its error frequency and was observed at >30% of the frequency
of the candidate variant.

Growth assays

Before phenotyping, the set of variant strains generated in this study were extended to include the haploid strains carrying missense variants constructed and sequence confirmed by Sirr *et al* [27]. These additional strains were re-arrayed to match the phenotyping layout used in this study, with at least 2 isolates of each variant. Additional control plates included a plate with wild type at every position, and plates consisting entirely of D100A or A99V (with control positions) *yPSAT1* strains, which are ClinVar pathogenic alleles encoding missense variants.

A Biomek i7 robot outfitted with a V&P 96-pin head was utilized to pin strain plate libraries between different culture media. Strains were initially grown to saturation in rich medium and then pinned onto solid medium utilizing glycerol as the central carbon source (YPG,1% yeast extract, 2% peptone, 2% glycerol) to remove any yeast cells lacking mitochondria (petite). Strains were then pinned back into rich medium and grown to saturation. Each plate was then pinned in replicate (n=3-6) onto solid minimal medium, which lacks serine, and grown for 3 days at 30°C. A mounted Canon PowerShot SX10 IS compact digital camera was used to take images (ISO200, f4.5, 1/40s exposure) every 24 hours for three days under consistent lighting, camera to subject distance, and zoom. Each plate was labeled with a custom code39 barcode that was included in the frame of view. Images were acquired as jpg files.

Image-based growth quantification

Each plate image was processed using PyPl8 (https://github.com/lacyk3/PyPl8) to extract features from each strain patch in each barcoded agar plate as follows: First, the barcode within each image was detected and decoded to rename each file using the corresponding plate name, replicate number, condition, and timepoint. Next, each image was cropped into 96 square tiles, segmented, and each replica pinned patch was identified using Otsu's method or circle detection. Finally, the sum of the gray scale pixel intensities within each strain patch (pixelsum), was extracted and used as the metric for growth estimation.

Growth data fitting and normalization

For haploid strains, raw phenotypic values were normalized, quality control filters were applied to each isolate, and a final relative growth estimate for each variant was determined, as described in Lo *et al* [32]. Briefly, normalization steps were carried out to account for the effects of plate-to-plate variation, relative growth of neighboring patches, and plate edge effects. Pin effect normalization did not reduce noise and was omitted. Isolates with (nonsynonymous) secondary mutations were removed from the dataset as were all isolates of variants that showed a high degree of variation in isolate-to-isolate growth values. This left a final filtered dataset of 5,164 independent isolates. Finally, a linear model was used to estimate

the relative growth of each genotype, on a scale with growth of null controls set to 0 and
growth of wild type set to 1.

A similar approach was applied to phenotypic values extracted from the diploid growth assay,
although neighbor effects were assumed negligible because of the checkerboard pinning
arrangement. For normalization, a linear model was used to simultaneously estimate the
effects of genotype, plate edge positioning and plate-to-plate variation on growth. Genotype
effects were rescaled to set homozygous null to 0 and homozygous wild type to 1.

Predicting diploid growth

We developed a linear model to predict the growth of diploid strains based on which pair of
yPSAT1 variants is present, using individual growth estimates of each allele in haploids. In this
model, for each diploid, the (haploid) growth estimates of the lower and higher growth alleles
were labelled as minimum and maximum (j, k), respectively. In cases of homozygous
combinations, the minimum and maximum were equal. Strains carrying a single copy of the
wild type allele (*yPSAT1*) or null (*ser1Δ0*) had their respective haploid estimates set equal to 1
and 0. To predict diploid growth (d) from the more impaired (x_j) and less impaired (x_k) alleles,
we performed an ordinary least squares regression to fit an additive pairwise model (d = a + bx_j
+ cx_k). We also compared this model to two simpler regression models. In the first model,
diploid growth was predicted from the mean of the haploid estimates ($x_j = x_k$). In the second,
diploid growth was predicted from the higher growing haploid allele only, i.e. complete
dominance ($x_j = 0$). Leave one out cross validation was used to assess model performance in
the best model.

Next, we evaluated the performance of experimental versus predicted growth as a binary
classifier for identifying genotypes matching those of patients diagnosed with *PSAT1*-related
serine deficiency disorders vs carrier parents. A logistic regression model indicated a threshold
(p=0.5) decision value of 72% and 73% for experimental and predicted diploid growth
respectively.

Protein structure and conservation analysis

Structural features were derived from the crystal structure of human phosphoserine
aminotransferase isoform 1 complexed with pyridoxal phosphate cofactor (PDB: 3e77; subunit
L17-L370, homodimer biological assembly). Secondary structure features were extracted from
this crystal structure using DSSP software [33,34]. AlphaFold [35,36] was used to visualize the
predicted structure of the missing 16 N-terminal residues. Molecular illustrations were created
by using the PyMol Molecular Graphics System Version 2.3.2 (Schrödinger). Tools from the
publicly available PyMol script and plugin repository were used to determine interfacial
residues and charge distribution. Dimer interface residues were defined as residues at which

their subunit-only based surface areas are larger than their complex-based surface areas (cutoff
value = 0.5). Macromolecular electrostatics were estimated using the Adaptive Poisson-
Boltzmann Solver plugin. Precomputed evolutionary conservation scores and grades, derived
from multiple sequence alignments of amino acid homologs with known structure, for subunit
A of PDB 3e77 were extracted from the ConSurf Database [38,39].

Results

Surveying the functional impact of large-scale missense variation in *PSAT1*

Our laboratory had previously established a yeast-based complementation assay that leveraged
the ability of the human *PSAT1* coding sequence to functionally replace its yeast ortholog, *SER1*
[27]. In this assay, growth on minimal medium lacking serine provides a quantitative readout of
PSAT1 activity, allowing the functional impact of protein coding variants to be assessed. Variant
impact is expressed on a relatively intuitive scale of activity between the level associated with
no activity (that of a complete gene deletion) and that conferred by the wild type human
protein coding sequence (*yPSAT1*). Variants causing a reduction in PSAT activity could have
their effect via decreases in enzymatic activity, protein stability, or a combination of both.

In this study, we applied the assay at scale to measure the effect of thousands of amino acid
substitutions as follows. First, with the exception of the translational start and stop codons, we
identified all unique amino acid substitutions (n = 2,182) that were accessible via a SNV across
the full length of human *PSAT1* isoform 1 cDNA (1,113 bp). Variant codons encoding each
amino acid substitution were then introduced into the yeast codon-optimized version of *PSAT1*
(*yPSAT1*). The resulting variant library was transformed into a haploid *SER1* deletion strain
(*ser1Δ0*) and integrated in single copy at the *SER1* locus of the yeast genome, under the control
of the endogenous *SER1* transcriptional promoter and terminator. Next, transformants were
individually arrayed in 96-well plate format, and the identity of the variant codon present in
each transformant was determined by Oxford Nanopore MinION sequencing (**Methods**). Strains
that harbored secondary mutations in the protein coding sequence were removed from further
consideration. The arrayed strain library was then grown, in triplicate, on solid medium lacking
serine and imaged after three days at 30°C. Data from these images were used to measure the
growth of each strain, relative to wild type (*yPSAT1*) and null (*ser1Δ0*), using a custom
automated image analysis pipeline and normalization software (**Methods**).

Here, we report the functional impact of 1,914 amino acid substitutions in *PSAT1*
(**Supplementary Table S3**), corresponding to ~88% of all unique SNV-accessible amino acid
substitutions (n=2,182) in the human *PSAT1* cDNA sequence. A subset of these substitutions
(n=196) was also assayed in our previous study [27] which sought to quantify the functional
impact of the full set of *PSAT1* missense variants described in ClinVar, gnomAD, or the clinical
literature at that time. Reanalysis in the current study allowed us to assess these amino acid

substitutions using the same data processing pipeline as the 1,718 new substitutions, thereby
placing them on a common scale to facilitate direct comparison. Despite the use of an improved
analysis pipeline, there was excellent agreement (R^2=0.85) between the normalized growth
estimates of the 196 substitutions in the two studies.

Among the full set of 1,914 amino acid substitutions, the distribution of growth values is
bimodal, with a large peak centered approximately on the value of the wild type *yPSAT1* strain
(normalized growth=1) and a smaller peak centered on the value of the null (deletion) strain
(normalized growth=0) (**Figure 2**). These results are consistent with a large group of protein
coding variants showing little or no functional impairment relative to wild type, and a smaller
group behaving as complete loss of function alleles, comparable to the null control. The
remaining protein coding variants showed varying levels of less severe functional impairment
(**Figure 2**).

Figure 2. Distribution of variant effect for 1,914 PSAT1 unique SNV-accessible missense amino acid substitutions.
Frequency of experimentally measured yeast growth values scaled relative to wild type yPSAT1 (normalized
growth=1) and null (normalized growth=0).

Mapping functional effects to protein structure and conservation

To provide a better understanding of the impact of missense substitutions on protein function,
we examined the results of our assay in the context of mechanistic, structural, and evolutionary
features of the human phosphoserine aminotransferase isoform 1 (PSAT) protein. PSAT belongs
to the large family of fold type I pyridoxal phosphate (PLP; active form of vitamin b6) cofactor-
dependent enzymes, and more specifically, the class IV group of aminotransferases [40,41] In
PLP-dependent enzymes the cofactor serves two ubiquitous functions in catalysis: to form a
reactive Schiff base covalent linkage with the substrate amino group and to act as an electron
sink to stabilize intermediate states [42–44]. Reaction specificity is determined by the
conformation of the active site as well as how residues interact with PLP and the substrates. For

aminotransferases, these enzymes typically form functional homomultimers with interfacial
PLP-containing active sites, where transamination reactions proceed via substrate binding,
amino group transfer, and product release [41].

PSAT catalyzes the reversible conversion of glutamate to α-ketoglutarate and 3-
phosphohydroxypyruvate (3-PHP) to phosphoserine [45]. The crystal structure PSAT complexed
with PLP (PDB: 3e77; residues L17-L370) is characterized by a s-shaped homodimer assembly in
which two separate PLP-containing active sites reside along the interface. An overall positive
charge distribution is also localized near these active sites (**Figure 3.B**). Each subunit consists of
two domains with mixed α/β fold; a large N-terminal catalytic domain (L17-G263), which forms
the dimer interface and binds PLP, and a small C-terminal domain (A264-L370) (**Figure 3.A**).
Although the 16 N-terminal amino acids (M1-K16) are not included in the crystal structure,
AlphaFold [35,36] predicts these residues to be placed between the two domains, proximal to
the active site (**Supplemental Figure S1**).

The overall structural organization and active site architecture of PSAT are conserved with
other members of the class IV family of aminotransferases [41], and phosphoserine
aminotransferases from eukaryotic and prokaryotic organisms [46]. Evolutionary conservation
scores for PSAT (pdb 3e77; subunit A) from the ConSurf DataBase [38,39], which are derived
from sequence alignments of amino acid homologs with known structure, illustrate that more
conserved (negative ConSurf score) regions localize near the active sites (**Figure 4.B**).
Furthermore, structural analysis of related phosphoserine aminotransferases [47–50] with
additional crystallographic experiments in substrate (or analog) bound states (not currently
available for human PSAT), have identified substrate binding residues that are highly conserved.

Figure 3. Overall subunit organization and homodimer assembly of human PSAT. **A)** Ribbon diagram representation
of the secondary structure of a subunit of huPSAT (PDB 3e77; L17-L370) complexed with PLP (green) in stick

Some of our knowledge of human PSAT protein structure-function is currently based on studies
of non-human phosphoserine aminotransferases. Our dataset of 1,914 PSAT variants
characterizes the functional impact of a distribution of 4-7 amino acid substitutions at each
position across the entire length of the human PSAT protein (370 aa) (**Figure 4.A**). These
substitutions display a range of activity from complete loss of function (scores ≤ 5%) to not
impaired relative to wild type (scores ≥ 95%). Impaired variants were non-randomly distributed
across the length of the protein. We identified several positions where the majority of SNV-
accessible variants exhibited some degree of functional impairment. In particular and
consistent with expectation, substitutions in conserved active site residues (**Figure 4.C**) were
often associated with more severe loss of activity (median scores ≤ 35%). However, not all
substitutions at highly conserved residues exhibited loss of function in our assay, and not all
substitutions at less conserved residues had activities close to wild type.

Figure 4. Missense variant effect map across the length of human PSAT (PDB: 3e77; L17-L370) compared to structural features and conservation. **A)** The bottom most plot represents the rank ordered distribution of variant growth at each amino acid position. Each stacked bar depicts the effect size of missense variants, grouped into normalized mean haploid growth values bins of 30% (righthand colorbar), and ordered from most impaired (top) to least impaired (bottom) growth bins. Variants that are not impaired relative to wild type (≥ 95%) are colored as white. Variants with growth values that were impaired relative to wild type (≤ 95%) are colored in shades of pink, with darker shades indicating more impaired growth. Grey fields correspond to missing data. The height of each color in each stacked bar represents the number of variants of a given growth bin, over the total number of possible variants at that position. Overlayed above are the secondary structure of PSAT, ConSurf evolutionary conservation scores [38,39], subunit domains and functional residues. Residues implicated in substrate binding (yellow), formation of the dimer interface (black), or cofactor binding (red) are depicted by circles. Alpha-helices (α; cylinder), 3₁₀-helices (η; cylinder), beta-sheets (β; arrows) are colored in gray and labeled, with turns shown as upward half-coils. **B-C)** Subunit surface colored by **B)** the ConSurf evolutionary conservation [38,39] grade and **C)** median yeast growth estimate per position. Two PLP cofactors are shown in each surface representation, to indicate the other interfacial active site bound to the opposite subunit (not depicted).

Functional impact of missense variation at the active site

To compare our results to known structural features of human PSAT, we examined the impact of amino acid substitutions at active site residues known bind the PLP cofactor, and a subset of conserved residues that have been found to bind substrates in other phosphoserine aminotransferases. First, we considered 12 residues that directly interacted with functional groups of PLP: the aldehyde group, the pyridoxal ring, and the phosphate group (**Figure 5.A** and **Figure 5.B**). The role of the aldehyde group in forming the initial Schiff base linkage with the catalytic lysine, and the role of the pyridoxal ring in stabilizing reaction intermediates though delocalization, are well established for PLP-dependent enzymes [41,43,44]. Each of the six SNV-accessible missense substitutions at the catalytic lysine (K200) resulted in null activity (scores ≤ 5%) (**Figure 5.C**). For ring coordination residues involved in pi-stacking (W107), or hydrogen bonding to the phenol (T156, S179) or nitrogen (D176, S178) groups of PLP, the majority of tested substitutions (15/23) displayed null-like activity (**Figure 5.C**). Notably, substitutions between the biochemically similar amino acids of threonine and serine at these positions resulted in less-impaired growth scores (T156S = 40%, S178T = 52%, S179T = 81%). Only two positions, S178 and S179, displayed median variant growth above the null level but most substitutions were still strongly impaired (S178 median score = 16%, S179 median score = 40%).

The role of the phosphate group of PLP is less well characterized for PLP-dependent enzymes. It may act as anchor point for the protein [51], or even directly interact with nearby substrates in [44]. In PSAT, six interfacial residues, G78, G79, C80, Q199, N241*, and T242*(asterisk indicates residues from opposite subunit), bind to the phosphate group (**Figure 5.B**). All substitutions at G78, G79, and Q199 displayed functional impairment (≤ 95%), with Q199E (score = 28%) and G79A (score = 78%) being the only two variants with activity above null (>5%) (**Figure 5.C**). Broader distributions of variant effects are observed at N241* and T242*, with two variants N241H (score = 103%), T242S (score = 108%) displaying no functional impairment relative to wild type (**Figure 5.C**). Interestingly, most substitutions at the C80 residue display wild type like activity (C80Y|S|G|W) with the only exception being C80R (growth score = 80%) (**Figure 5.C**). This residue (C80) is typically found as a serine, threonine (not SNV-accessible in human), or arginine in other organisms (ConSurf DataBase [38,39] profile for 3e77 subunit A). It has been hypothesized that C80 contributes to the higher phosphoserine substrate affinity (Km = 5 µM) observed in PSAT relative to other phosphoserine aminotransferases [46], and possibly the positive charge introduction represented by C80R negatively impacts this binding.

However, like most aminotransferases, the specific mechanisms of substrate binding are less understood for PSAT. As an alternative, we considered 5 substrate binding residues that were identified in *Escherichia coli* [47], *Bacillus alcalophilus* [48], and *Arabidopsis thaliana* [49] phosphoserine aminotransferases that had crystal structures of phosphoserine or α-methyl-L-glutamate (analog) bound states (not available for human PSAT). Amino acid substitutions at all

but one of these substrate binding residues, corresponding to H335, R336, R342, H44* and
R45* (asterisk indicates residues from opposite subunit) in human PSAT, show strong loss of
function (**Figure 5.D and Figure 5.E**). The exception is R336, which displayed activity similar to
wild type for all tested substitutions. Thus, R336 may be a conserved residue that does not
directly participate in substrate stabilization in PSAT but does in other orthologs.

Figure 5. Functional impact of amino acid substitutions at active site residues. Stick representations for residues in
one subunit are colored in cyan, residues from the opposite subunit in the homodimer assembly are in yellow, and
the PLP cofactor in green in all panels where the crystal structure of human PSAT complexed with PLP (PDB: 3e77)
is visualized. **A-B)** Organization of PLP cofactor binding residues. Hydrogen bonding is depicted by dashed black
lines and nearby water molecules are represented by white spheres. **A)** Residues that bind or coordinate PLP's
heterocyclic ring, with the catalytic lysine shown covalently bound to PLP. **B)** Residues that bind to the phosphate
group of PLP. **C)** The distribution of variant growth scores for cofactor binding residues. Each circle is colored
according to the amino acid that it is substituted as shown (Mut AA). Asterisks here denote amino acids from the

other subunit. The median growth score for each position is shown as a horizontal grey bar. **D)** Relative position of
highly conserved substrate binding residues to PLP, in the unbound (no substrate) crystal structure of human PSAT.
E) The distribution of functional effects for these conserved substrate binding residues, with the same coloring
scheme and annotation as in **C**.

Mapping functional effects to clinical interpretations

Although as a rare disease there are a limited number of pathogenic alleles, a number of those
alleles reside at these functionally important residues. We next compared our results to clinical
classifications for all corresponding missense variants in ClinVar. We expected that
substitutions derived from known pathogenic variants would produce strong reductions in PSAT
activity in our assay, while those derived from known benign variants would have activities
comparable to wild type or display only weak reductions in activity. Of the 90 amino acid
substitutions derived from variants currently in ClinVar, three (S179L, A99V, and D100A) are
derived from variants annotated as pathogenic/likely pathogenic, two (I123V and P295R) from
variants annotated as benign/likely-benign, and three (V149M, A234S, and R306C) from
variants with conflicting interpretations [3] (November 2022). Since our previous study [27],
while the small number of missense alleles with definitive clinical significance calls remained
largely unchanged, the number of missense VUS increased nearly 12-fold (from 7 to 82).
Consistent with expectation and our previous results [27], the pathogenic/likely pathogenic
substitutions all resulted in substantial functional impairment in our assay, with normalized
growth values of 82% or below. Similarly, the benign/likely-benign substitutions demonstrated
little impairment, with normalized growth values above 91% (**Figure 6**).

To facilitate the use of our functional data as supporting evidence for human sequence variant
annotation [24], we used the activities of the substitutions with associated clinical significance
calls in a simple thresholding approach, similar that that used in our previous study [27]. We
defined a functionally impaired range for scores ≤82% growth in our assay, which is less than or
equal to the scores of substitutions derived from known pathogenic variants (**Figure 6**). We also
defined a functionally unimpaired range for scores ≥91%, which is greater than or equal to the
scores of substitutions derived from known benign variants (**Figure 6**). Because gain of function
mutations have not been associated with serine deficiency disorders, we included values
greater than wild type in the unimpaired range. Finally, because there are no clinically
annotated variants associated with substitutions having scores between 82% and 91%, this
range of the assay is of unknown clinical significance, and we have labeled this range
indeterminate. Within these ranges of our assay, 534 (28%) of all amino acid substitutions are
functionally impaired, 1,100 (57%) are functionally unimpaired, and 280 (15%) are
indeterminant (**Figure 6**). Of the 80 ClinVar missense variants currently annotated as VUS that
were tested in our assay, 22 result in substitutions that fall in our impaired range, 41 in our
unimpaired range and 17 in the intermediate range. Thus, our dataset provides functional

information for a large number of variants that lack clinical interpretation, including 77% of the missense VUS currently listed in ClinVar.

Figure 6: Mutational scan of 1,914 PSAT1 unique SNV-accessible missense amino acid substitutions. Rank ordered (lowest to highest) normalized variant growth. Black dots represent the mean normalized growth estimate for each substitution, with light grey bars indicating standard errors. All current ClinVar annotations associated with tested substitutions are labelled above their growth estimate as benign/likely-benign (green), pathogenic/likely-pathogenic (magenta), conflicting interpretations (yellow), or as an asterisk below their growth estimate if they are of uncertain significance (dark grey). The boxed panel depicts the stratification of normalized growth estimates, based on thresholds derived from ClinVar clinical annotation, into functionally unimpaired (green), indeterminate (gray), or functionally impaired (magenta) ranges.

Experimental models of homozygous and compound heterozygous genotypes

Loss of function mutations in *PSAT1* cause disease that ranges from severe (Neu-Laxova syndrome 2, NLS2) to milder forms (PSAT deficiency, PSATD) of the disorder. These are collectively referred to as NLS2/PSATD, or individually when clinical severity is discussed specifically. Because NLS2/PSATD is an autosomal recessive disease, clinical manifestation depends on the enzymatic function conferred by the combination of *PSAT1* alleles in the patient's genome. The ability to generate stable diploid yeast cells by mating haploids allows us to model diploid human *PSAT1* genotypes in yeast and quantitively assess the function of allele pairs. As in our previous study [27], we sought to use our growth assay to functionally assess pairwise combinations of protein-coding variants across all reported unique patient (and carrier parent) genotypes currently available (**Supplemental Table S4**). Since our previous study [27], six additional reports [20,52–56] have added descriptions of 20 new NLS2/PSATD patients and

twelve unique genotypes to the disease literature (**Supplemental Table S4**). To experimentally
measure the functional impact of these allele combinations in our yeast assay, we constructed
diploid strains harboring *yPSAT1* allele combinations encoding the same pair of PSAT1 amino
acid sequences as the human genotypes. We then assayed the growth of these strains relative
to homozygous wild type (*yPSAT1*) and null (*ser1Δ0*) on solid minimal medium lacking serine.
Previously constructued *yPSAT1* diploid strains encoding homozygous A99V or S43R, and the
compound heterzygote A99V / S179L [27], were also included for comparision. Although the
growth of some strains modelling patient genotypes were close to those recpitulating some
carrier genotypes, all modelled patient genotypes (**Supplemental Table S5**) displayed reduced
growth relative to their respective carrier parents and the homozygous wildtype (**Figure 7.B**).
Consistent with our previous study [27], we also observed good agreement between the degree
of *PSAT1* functional impairment in our diploid assay and disease severity (**Figure 7.B**). Diploids
corresponding to genotypes from NLS2 patients had normalized growth from 0% to 77%, and
diploids corresponding to genotypes from PSATD patients had a higher normalized growth
range of 70% to 88%.

Figure 7: Modeling PSAT activity in patient genotypes. Both bar plots show the mean normalized growth estimate
and standard errors. **A)** Haploid growth estimates for novel (purple) [20,52,54,56] and established [18,57] (pink)
missense alleles in patient amino acid substitutions. **B)** Diploid yeast models of patient trios. Carrier parents are
shown in light and dark grey, and the offspring in magenta. Recurrent carrier genotypes are repeated for ease of
reference relative to the offspring. The clinical severity (NLS2 or PSATD) associated with patient genotypes is
indicated. Asterisks denote offspring genotypes that have been reported in multiple unrelated families.

Our assay results are also consistent with clinical stratification within the group of NLS2 and
PSATD patients. For PSATD, the A15P / A15P homozygote is the least impaired in our assay
(77% growth) and is also the mildest form of PSATD reported to date. A recent report [56]
described two unrelated teenagers that, other than congenital or childhood ichthyosis, were

developmentally normal and displayed neuropathy onset at ages 16 and 17. For NLS2, the two
genotypes with the lowest growth in our assay (G79W / G79W and C245R / C245R) correspond
to patients displaying severe NLS2 [20]. Three of the remaining NLS2 genotypes modelled here
(Y70N / R342W, A99V / S179L, A99V / R61W) display higher growth in our assay and
correspond to patients described as having moderate NLS expression [18,20,52]. Among these,
Y70N / R342W was the most impaired in our assay (18% growth) and the patient had a
postnatal survival of 8 weeks [52], while the individual harboring R61W / A99V (77% growth,
least impaired) had the longest NLS2-associated postnatal survival described to date (4 months)
[20]. Patients with genotypes corresponding to the A99V homozygote (60% growth), displayed
variable postnatal survival ranging from 1 day to 9 weeks [18–20]. Together, these results
highlight the potential value of our model organism assay for variant interpretation in the
context of a diploid *PSAT1* genotype.

Predicting biallelic effects from individual allele measurements

Experimentally assessing the level of PSAT activity associated with all allele-pairs would require
the construction of 1.83M diploid yeast strains. As a more feasible alternative, we evaluated
whether a relationship exists between individual haploid estimates and their resulting diploid
growth in combination. For homozygotes, a linear relationship was seen between haploid and
diploid growth estimates (**Figure 7.A** and **Figure 7.B** and **Supplemental Figure S2**; R^2=0.93). To
extend this analysis to heterozygotes, we fit a model to our experimental diploid dataset that
includes patient and carrier genotypes (n = 23), predicting diploid growth as a linear
combination of the allele with the higher and the allele with the lower haploid growth values
(**Figure 8.A**). This model assumed a uniform degree of dominance of the higher growing allele
over the lower growing allele and explained 97% of variance in the growth of the diploids. To
compare how this model performed relative to alternatives with simpler assumptions, we
generated models that assumed either complete dominance of the higher-growing allele or an
equal impact of both alleles (**Figure 8.B** and **Figure 8.C**). Our full model performed significantly
better than either the alternative models (Anova, Df=1; p=<5.4x10^{-6} and p<2.3x10^{-6},
respectively). Leave one out cross validation of the pairwise model displayed an excellent ability
to predict diploid growth from haploid measurements (RMSE = 0.0698 and MAE=0.0566).

Figure 8: Experimental and predicted biallelic growth values as clinical classifiers. Observed versus fitted values
shown for three linear regression models (pairwise additive **(A)**, simple dominant **(B)**, and simple average **(C)**)
predicting diploid growth (allele pairs) as a function of haploid growth (single alleles). Each circle represents a
unique biallelic combination. Patient, carrier, and control strains are labeled as indicated. A diagonal line of perfect
correspondence (1:1 ratio) and the coefficient of determination (R^2) for each model are included for ease of
comparison. Decision boundaries for observed (72%) versus predicted (73%) diploid growth as a binary classifier
for identifying patient genotypes modeled in our diploid assay are shown as horizontal and vertical dotted grey
lines in **(A)**, respectively. Confidence intervals (95%) for each model are shaded in purple. **D)** Heatmap of the
predicted diploid growth value for all possible pairwise combinations of haploid allele measurements. **E)** Frequency
of predicted diploid growth values (in 5% bins) for all ~1.8M combinations of alleles. The decision boundary of 73%
is shown as a vertical dotted line.

We next compared the results of applying logistic regression to experimental and predicted
diploid growth values to produce binary classifiers distinguishing (low growing) genotypes from
patients diagnosed with NLS2/PSATD (n=9) versus (high growing) carrier parent genotypes in
our trio models (n=10). The threshold growth values at the decision boundary (p=0.5) were
similar for both approaches: 72% for experimental and 73% for predicted diploid growth.
Although the decision boundary using predicted values performed better and misclassified only
one patient genotype (A15P / A15P) versus a total of 4 for the model using experimental values
(**Figure 8.A**). It is likely the haploid estimates used to make predictions on diploid growth were
less noisy as they were generated from a larger number of measurements compared to our set
of experimental diploids. These results suggest that diploid growth values predicted from
haploid measurements perform well in clinical classification of human *PSAT1* genotypes.

On this basis, we computed the predicted diploid growth from all unique pairwise combinations
(~1.8M) of haploid estimates from 1,914 *yPSAT1* missense alleles, wild type (*yPSAT1*), and null
(*ser1Δ0*) using the pairwise-additive model (**Figure 8.A**). We then classified whether the
resulting values fell above or below the disease classification boundary of 73% growth. A large
majority (89%) of genotypes fell above the decision boundary threshold, suggesting these
genotypes would not lead to NLS2/PSATD (**Figure 8.D** and **Figure 8.E**). However, over 200K
yPSAT1 biallelic combinations displayed predicted diploid growth below the classification
threshold, consistent with these genotypes resulting in disease (**Figure 8.E**).

Discussion

Here we presented a comprehensive functional analysis of SNVs in human PSAT. While this data
shows strong agreement with the structure-function features of the enzyme, clinical
interpretation relies both on knowledge of functional impact of a variant and the relationship to
clinical data. Specifically, we leveraged the small number of variants with existing clinical
interpretation to provide evidence for the likely clinical impact of other variants in similar
ranges of the assay. As additional clinical information become available through identification
of new patients and additional research into the natural history of the disease, the ability to
make clinical inferences using our dataset will improve. In the three years since our previous

study [27], additional patients were described in the disease literature. Not only was there good agreement between the results of our assay and disease severity, but this additional clinical information also allowed us to extend the range of our assay associated with clinical outcomes.

Given that the therapeutic potential for this medically actionable disorder is highest before patients become symptomatic, clinical sequencing can be a strong diagnostic tool if variant interpretation is also informative. Sequencing based diagnostics are becoming more widely applied in newborns, prenatally [58], and as a part of programs like Undiagnosed Disease Program [59,60] or Deciphering Developmental Disorders [61]. The most recent AMCG guidelines [62] recommend exome or genome sequencing as a first or second tier option for patients displaying early neonatal (<1 year) congenital abnormalities, or for patients exhibiting developmental delay or intellectual disability before the age of 18. Furthermore, large scale genome sequencing efforts like the All of US research program and UK biobank [63] are aggregating a depth of population level genomic and longitudinal health data. Given these developments, studies such as ours that provide information both about individual alleles and the functional impact of allele pairs will play an increasingly important role in realizing the promise of precision medicine.

References

1. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. Nature. Nature Research; 2020;581:434–43.

2. Gudmundsson S, Singer-Berk M, Watts NA, Phu W, Goodrich JK, Solomonson M, et al. Variant interpretation using population databases: Lessons from gnomAD. Hum Mutat [Internet]. 2021; Available from: https://onlinelibrary.wiley.com/doi/10.1002/humu.24309

3. Landrum MJ, Lee JM, Benson M, Brown GR, Chao C, Chitipiralla S, et al. ClinVar: Improving access to variant interpretations and supporting evidence. Nucleic Acids Res. Oxford University Press; 2018;46:D1062–7.

4. Illsinger S, Das AM. Impact of selected inborn errors of metabolism on prenatal and neonatal development. IUBMB Life. 2010;62:403–13.

5. Ferreira CR, Rahman S, Keller M, Zschocke J, Abdenur J, Ali H, et al. An international classification of inherited metabolic disorders (ICIMD). J Inherit Metab Dis. John Wiley and Sons Inc; 2021;44:164–77.

6. Sanderson S, Green A, Preece MA, Burton H. The incidence of inherited metabolic disorders in the West Midlands, UK. Arch Dis Child. 2006;91:896–9.

7. Dionisi-Vici C, Rizzo C, Burlina AB, Caruso U, Sabetta G, Uziel G, et al. Inborn errors of metabolism in the Italian pediatric population: a national retrospective survey. J Pediatr. 2002;140:321–7.

8. Gambello MJ, Li H. Current strategies for the treatment of inborn errors of metabolism.
Journal of Genetics and Genomics. Institute of Genetics and Developmental Biology; 2018. p.
61–70.

9. Morris AAM, Kožich V, Santra S, Andria G, Ben-Omran TIM, Chakrapani AB, et al. Guidelines
for the diagnosis and management of cystathionine beta-synthase deficiency. J Inherit Metab
Dis. Springer Netherlands; 2017. p. 49–74.

10. Stockler S, Plecko B, Gospe SM, Coulter-Mackie M, Connolly M, van Karnebeek C, et al.
Pyridoxine dependent epilepsy and antiquitin deficiency. Mol Genet Metab. 2011;104:48–60.

11. El-Hattab AW. Serine biosynthesis and transport defects. Mol Genet Metab. Academic Press
Inc.; 2016. p. 153–9.

12. de Koning TJ, Klomp LWJ, Oppen ACCV, Beemer PFA, Dorland L, Berg IETV den, et al.
Prenatal and early postnatal treatment in 3-phosphoglycerate-dehydrogenase deficiency.
Lancet. Elsevier B.V.; 2004;364:2221–2.

13. Jaeken J, Detheux M, van Maldergem L, Foulon M, Carchon H, van Schaftingen E, et al. 3-
Phosphoglycerate dehydrogenase deficiency: an inborn error of serine biosynthesis. Archives
ofDisease in Childhood. 1996.

14. Jaeken J, Detheux M, Fryns JP, Collet JF, Alliet P, van Schaftingen E. Phosphoserine
phosphatase deficiency in a patient with Williams syndrome. J Med Genet. BMJ Publishing
Group; 1997;34:594–6.

15. de Koning TJ, Duran M, van Maldergem L, Pineda M, Dorland L, Gooskens R, et al.
Congenital microcephaly and seizures due to 3-phosphoglycerate dehydrogenase de¢ciency:
Outcome of treatment with amino acids. J. Inherit. Metab. Dis. 2002.

16. Hart CE, Race V, Achouri Y, Wiame E, Sharrard M, Olpin SE, et al. Phosphoserine
aminotransferase deficiency: A novel disorder of the serine biosynthesis pathway. Am J Hum
Genet. University of Chicago Press; 2007;80:931–7.

17. de Koning TJ. Amino acid synthesis deficiencies. J Inherit Metab Dis. Springer Netherlands;
2017. p. 609–20.

18. Acuna-Hidalgo R, Schanze D, Kariminejad A, Nordgren A, Kariminejad MH, Conner P, et al.
Neu-laxova syndrome is a heterogeneous metabolic disorder caused by defects in enzymes of
the l-serine biosynthesis pathway. Am J Hum Genet. Cell Press; 2014;95:285–93.

19. El-Hattab AW, Shaheen R, Hertecant J, Galadari HI, Albaqawi BS, Nabil A, et al. On the
phenotypic spectrum of serine biosynthesis defects. J Inherit Metab Dis. Springer Netherlands;
2016;39:373–81.

20. Abdelfattah F, Kariminejad A, Kahlert AK, Morrison PJ, Gumus E, Mathews KD, et al.
Expanding the genotypic and phenotypic spectrum of severe serine biosynthesis disorders. Hum
Mutat. John Wiley and Sons Inc.; 2020;41:1615–28.

21. Moat S, Carling R, Nix A, Henderson M, Briddon A, Prunty H, et al. Multicentre age-related
reference intervals for cerebrospinal fluid serine concentrations: Implications for the diagnosis
and follow-up of serine biosynthesis disorders. Mol Genet Metab. 2010;101:149–52.

22. Sun S, Yang F, Tan G, Costanzo M, Oughtred R, Hirschman J, et al. An extended set of yeast-
based functional assays accurately identifies human disease mutations. Genome Res.
2016;26:670–80.

23. Starita LM, Ahituv N, Dunham MJ, Kitzman JO, Roth FP, Seelig G, et al. Variant
Interpretation: Functional Assays to the Rescue. Am J Hum Genet. Cell Press; 2017. p. 315–25.

24. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, et al. Standards and guidelines for
the interpretation of sequence variants: A joint consensus recommendation of the American
College of Medical Genetics and Genomics and the Association for Molecular Pathology.
Genetics in Medicine. Nature Publishing Group; 2015;17:405–24.

25. Gasperini M, Starita L, Shendure J. The power of multiplexed functional analysis of genetic
variants. Nat Protoc. Nature Publishing Group; 2016. p. 1782–7.

26. Weile J, Roth FP. Multiplexed assays of variant effects contribute to a growing genotype–
phenotype atlas. Hum Genet. Springer Verlag; 2018. p. 665–78.

27. Sirr A, Lo RS, Cromie GA, Scott AC, Ashmead J, Heyesus M, et al. A yeast-based
complementation assay elucidates the functional impact of 200 missense variants in human
PSAT1. J Inherit Metab Dis. John Wiley and Sons Inc.; 2020;43:758–69.

28. Winston F, Dollard C, Ricupero-Hovasse SL. Construction of a set of convenient
saccharomyces cerevisiae strains that are isogenic to S288C. Yeast. Chichester, UK: John Wiley
& Sons, Ltd; 1995;11:53–5.

29. Rose M, Winston F, and Hieter P. Methods in Yeast Genetics: A Laboratory Course Manual.
Cold Spring Harbor Laboratory Press; 1990.

30. Farrell CM, O'Leary NA, Harte RA, Loveland JE, Wilming LG, Wallin C, et al. Current status
and new features of the Consensus Coding Sequence database. Nucleic Acids Res.
2014;42:D865–72.

31. Illuxley C, Green ED, Dunbam I. Rapid assessment of S. cerevisiae mating type by PCR.
Trends in Genetics. 1990;6:236.

32. Lo RS, Cromie GA, Tang M, Teng K, Owens K, Sirr A, et al. The functional impact of 1,570
SNP-accessible missense variants in human OTC. Available from:
https://doi.org/10.1101/2022.10.26.513893

33. Touw WG, Baakman C, Black J, te Beek TAH, Krieger E, Joosten RP, et al. A series of PDB-
related databanks for everyday needs. Nucleic Acids Res. 2015;43:D364-8.

34. Kabsch W, Sander C. Dictionary of protein secondary structure: pattern recognition of
hydrogen-bonded and geometrical features. Biopolymers. 1983;22:2577–637.

35. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate
protein structure prediction with AlphaFold. Nature. 2021;596:583–9.

36. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold
Protein Structure Database: massively expanding the structural coverage of protein-sequence
space with high-accuracy models. Nucleic Acids Res. 2022;50:D439–44.

37. Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, et al. Biopython: freely available
Python tools for computational molecular biology and bioinformatics. Bioinformatics.
2009;25:1422–3.

38. Goldenberg O, Erez E, Nimrod G, Ben-Tal N. The ConSurf-DB: pre-calculated evolutionary
conservation profiles of protein structures. Nucleic Acids Res. 2009;37:D323-7.

39. ben Chorin A, Masrati G, Kessel A, Narunsky A, Sprinzak J, Lahav S, et al. ConSurf-DB: An
accessible repository for the evolutionary conservation patterns of the majority of PDB
proteins. Protein Sci. 2020;29:258–67.

40. Murtas G, Marcone GL, Sacchi S, Pollegioni L. L-serine synthesis via the phosphorylated
pathway in humans. Cellular and Molecular Life Sciences. Springer Science and Business Media
Deutschland GmbH; 2020. p. 5131–48.

41. Koper K, Han SW, Pastor DC, Yoshikuni Y, Maeda HA. Evolutionary origin and functional
diversification of aminotransferases. Journal of Biological Chemistry. American Society for
Biochemistry and Molecular Biology Inc.; 2022.

42. John RA. Pyridoxal phosphate-dependent enzymes. Biochim Biophys Acta. 1995;1248:81–
96.

43. Eliot AC, Kirsch JF. Pyridoxal phosphate enzymes: Mechanistic, structural, and evolutionary
considerations. Annu Rev Biochem. 2004. p. 383–415.

44. Liang J, Han Q, Tan Y, Ding H, Li J. Current advances on structure-function relationships of
pyridoxal 5'-phosphate-dependent enzymes. Front Mol Biosci. Frontiers Media S.A.; 2019.

45. Youn BAEK J, Youn JUN D, Taub D, Ho KIM Y. Characterization of human phosphoserine
aminotransferase involved in the phosphorylated pathway of L-serine biosynthesis. Biochem. J.
2003.

46. Singh RK, Kumar D, Gourinath S. Phosphoserine Aminotransferase has Conserved Active Site
from Microbes to Higher Eukaryotes with Minor Deviations. Protein Pept Lett. Bentham Science
Publishers Ltd.; 2021;28:996–1008.

47. Hester G, Stark W, Moser M, È rg Kallen Zora Markovic Â-Housley J, Jansonius JN. Crystal
Structure of Phosphoserine Aminotransferase from Escherichia coli at 2.3 A Ê Resolution:
Comparison of the Unligated Enzyme and a Complex with a a a-Methyl-L-Glutamate. J. Mol.
Biol. 1999.

48. Battula P, Dubnovitsky AP, Papageorgiou AC. Structural basis of l-phosphoserine binding to
Bacillus alcalophilus phosphoserine aminotransferase. Acta Crystallogr D Biol Crystallogr.
2013;69:804–11.

49. Sekula B, Ruszkowski M, Dauter Z. Structural analysis of phosphoserine aminotransferase
(isoform 1) from arabidopsis Thaliana– the enzyme involved in the phosphorylated pathway of
serine biosynthesis. Front Plant Sci. Frontiers Media S.A.; 2018;9.

50. Singh RK, Tomar P, Dharavath S, Kumar S, Gourinath S. N-terminal residues are crucial for
quaternary structure and active site conformation for the phosphoserine aminotransferase
from enteric human parasite E. histolytica. Int J Biol Macromol. Elsevier B.V.; 2019;132:1012–
23.

51. Denesyuk AI, Denessiouk KA, Korpela T, Johnson MS. Functional attributes of the phosphate
group binding cup of pyridoxal phosphate-dependent enzymes. J Mol Biol. Academic Press;
2002;316:155–72.

52. Ni C, Cheng RH, Zhang J, Liang JY, Wei RQ, Li M, et al. Novel and recurrent PHGDH and
PSAT1 mutations in Chinese patients with Neu-Laxova syndrome. European Journal of
Dermatology. John Libbey Eurotext; 2019;29:641–6.

53. Shapira Zaltsberg G, McMillan HJ, Miller E. Phosphoserine aminotransferase deficiency:
imaging findings in a child with congenital microcephaly. Journal of Maternal-Fetal and
Neonatal Medicine. Taylor and Francis Ltd; 2020;33:1033–5.

54. Debs S, Ferreira CR, Groden C, Kim HJ, King KA, King MC, et al. Adult diagnosis of congenital
serine biosynthesis defect: A treatable cause of progressive neuropathy. Am J Med Genet A.
John Wiley and Sons Inc; 2021;185:2102–7.

55. Serrano Olave A, Padín López A, Martín Cruz M, Monís Rodríguez S, Narbona Arias I, Jiménez
López JS. Prenatal Diagnosis of Neu–Laxova Syndrome. Diagnostics. MDPI AG; 2022;12:1535.

56. Shen Y, Peng Y, Huang P, Zheng Y, Li S, Jiang K, et al. Juvenile-onset PSAT1-related
neuropathy: A milder phenotype of serine deficiency disorder. Front Genet. Frontiers Media SA;
2022;13.

57. Brassier A, Valayannopoulos V, Bahi-Buisson N, Wiame E, Hubert L, Boddaert N, et al. Two
new cases of serine deficiency disorders treated with l-serine. European Journal of Paediatric
Neurology. W.B. Saunders Ltd; 2016;20:53–60.

58. Fu F, Li R, Yu Q, Wang D, Deng Q, Li L, et al. Application of exome sequencing for prenatal
diagnosis of fetal structural anomalies: clinical experience and lessons learned from a cohort of
1618 fetuses. Genome Med. BioMed Central Ltd; 2022;14.

59. Gahl WA, Wise AL, Ashley EA. The Undiagnosed Diseases Network of the National Institutes
of Health: A National Extension. JAMA. 2015;314:1797–8.

60. Macnamara EF, D'Souza P, Tifft CJ. The undiagnosed diseases program: Approach to
diagnosis. Ferreira C, editor. Transl Sci Rare Dis [Internet]. 2019;4:179–88. Available from:
https://www.medra.org/servlet/aliasResolver?alias=iospress&doi=10.3233/TRD-190045

61. Firth H v, Wright CF, DDD Study. The Deciphering Developmental Disorders (DDD) study.
Dev Med Child Neurol. 2011;53:702–3.

62. Manickam K, McClain MR, Demmer LA, Biswas S, Kearney HM, Malinowski J, et al. Exome
and genome sequencing for pediatric patients with congenital anomalies or intellectual
disability: an evidence-based clinical guideline of the American College of Medical Genetics and
Genomics (ACMG). Genetics in Medicine. Springer Nature; 2021;23:2029–37.

63. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, et al. The UK Biobank resource
with deep phenotyping and genomic data. Nature. 2018;562:203–9.

Declarations

780 No human or animal subjects were used in this work.

Competing interests

782 A.M.D. is a scientific advisor to and has a financial interest in Fenologica Biosciences, Inc., a
783 company that develops instrumentation and analytical tools for the analysis of microbial
784 colonies.

Funding

786 This work was funded in part by NIH/ NIGMS award R01 GM134274 to A.M.D.

Acknowledgements

788 We thank Douglas Fowler, Maitreya Dunham, Fowzan Alkuraya, and Ayman El Hattab for
789 helpful discussions.