Homeodomain-interacting protein kinase maintains neuronal homeostasis during normal *Caenorhabditis elegans* aging and systemically regulates longevity from serotonergic and GABAergic neurons

Authors: Maria I. Lazaro-Pena[1], Adam B. Cornwell[1], Carlos A. Diaz-Balzac[2], Ritika Das[1,3,4], Zachary C. Ward[1], Nicholas Macoretta[3], Juilee Thakar[1,5,6], Andrew V. Samuelson[1*]

Affiliations:
[1] Department of Biomedical Genetics, University of Rochester Medical Center, Rochester, New York, USA
[2] Division of Endocrinology, Diabetes and Metabolism, Department of Medicine, University of Rochester Medical Center, Rochester, New York, USA
[3] Department of Biology, University of Rochester, Rochester, New York, USA
[4] Department of Cell Biology, Skirball Institute of Biomolecular Medicine, NYU School of Medicine, New York, New York, USA.
[5] Department of Biostatistics and Computational Biology, University of Rochester Medical Center, 601 Elmwood Avenue, Rochester, NY 14642, USA
[6] Department of Microbiology and Immunology, University of Rochester Medical Center, 601 Elmwood Avenue, Rochester, NY 14642, USA

* Corresponding author. Andrew_Samuelson@URMC.Rochester.edu

Competing Interest Statement: The authors have no competing interests.

Abstract

Aging and the age-associated decline of the proteome is determined in part through neuronal control of evolutionarily conserved transcriptional effectors, which safeguard homeostasis under fluctuating metabolic and stress conditions by regulating an expansive proteostatic network. We have discovered the *Caenorhabditis elegans* homeodomain-interacting protein kinase (HPK-1) acts as a key transcriptional effector to preserve neuronal integrity, function, and proteostasis during aging. Loss of *hpk-1* results in drastic dysregulation in expression of neuronal genes, including genes associated with neuronal aging. During normal aging *hpk-1* expression increases throughout the nervous system more broadly than any other kinase. Within the aging nervous system, *hpk-1* induction overlaps with key longevity transcription factors*,* which suggests *hpk-1* expression mitigates natural age-associated physiological decline. Consistently, pan-neuronal overexpression of *hpk-1* extends longevity, preserves proteostasis both within and outside of the nervous system, and improves stress resistance. Neuronal HPK-1 improves proteostasis through kinase activity. HPK-1 functions cell non-autonomously within serotonergic and GABAergic neurons to improve proteostasis in distal tissues by specifically regulating distinct components of the proteostatic network. Increased serotonergic HPK-1 enhances the heat shock response and survival to acute stress. In contrast, GABAergic HPK-1 induces basal autophagy and extends longevity, which requires *mxl-2* (MLX), *hlh-30* (TFEB), and *daf-16* (FOXO). Our work establishes *hpk-1* as a key neuronal transcriptional regulator critical for preservation of neuronal function during aging. Further, these data provide novel insight as to how the nervous system partitions acute and chronic adaptive response pathways to delay aging by maintaining organismal homeostasis.

Keywords: Neuronal cell-nonautonomous control, gene expression, proteostasis, longevity, aging, stress response, autophagy, homeodomain interacting protein kinase

Introduction

The gradual decline of function within the proteome (proteostasis) is a characteristic of aging, which precipitates the onset and progression of a growing number of age-associated diseases (Balch et al., 2008; Gidalevitz et al., 2006; Gidalevitz et al., 2010; Kikis et al., 2010; Morimoto, 2011, 2019; Sala et al., 2017; Taylor and Dillin, 2011): Alzheimer's disease (AD), Parkinson's disease (PD), Huntington's disease, and Amyotrophic Lateral Sclerosis (ALS) are neurodegenerative diseases driven by genetic alterations that typically predispose a mutant protein isoform to aggregate and have toxic gain of function properties on the neuronal proteome. Outside of the nervous system age-associated proteostatic decline also leads to disease, for example: the onset of diabetes mellitus (Jaisson and Gillery, 2014); thus the decline in proteostasis during normal aging is not limited to the nervous system. A growing number of studies suggest that aging is not simply the result of the stochastic accumulation of damage, but is determined through genetics and coordinated mechanisms across tissues and cell types (Labbadia and Morimoto, 2015; Lazaro-Pena et al., 2022; Lopez-Otin et al., 2013). Proteostatic decline corresponds with the age-associated breakdown of a large proteostasis network (PN), which integrates stress-responsive control of protein folding, degradation, and translation in response to myriad cell intrinsic and non-autonomous signals. And yet, while major components of the proteostatic network have been identified, it is still poorly understood how multi-cellular organisms maintain systemic proteostasis within and across tissues to delay aging.

The relatively simple metazoan animal *Caenorhabditis elegans* (*C. elegans*) is a premier model system to learn how proteostasis is coordinated across cell- and tissue-

types in response to myriad signals. Discoveries of cell non-autonomous signaling first

made in *C. elegans* have evolutionarily conserved components, and revealed that the

nervous system acts as a key cell non-autonomous regulator of organismal proteostasis

and longevity (reviewed in (Miller et al., 2020)). For example, a pair of thermosensory

neurons systemically regulate the heat shock response (HSR) in a serotonin-dependent

manner (Prahlad et al., 2008; Tatum et al., 2015). In contrast, a second regulatory

component from the GABAergic and cholinergic systems normally limits muscle cell

proteostasis (Garcia et al., 2007; Silva et al., 2013).

The first discovery of a putative link between the transcriptional co-factor

homeodomain-interacting protein kinase (HPK-1) and longevity was from our genome-

wide RNAi screen that identified 103 genes that were essential for decreased insulin-

like signaling (ILS) to extend longevity (Samuelson et al., 2007). In *C. elegans, hpk-1*

preserves proteostasis, stress response and organismal longevity (Berber et al., 2016;

Das et al., 2017; Samuelson et al., 2007). Additionally, we have previously shown that

HPK-1 extends longevity through distinct genetic pathways defined by HSF-1, and the

target of rapamycin complex 1 (TORC1) (Das et al., 2017). Activation by either

metabolic or genotoxic stressors has been observed from yeast to mammals,

suggesting that this family of transcriptional cofactors arose early in evolution to couple

metabolic and stress signaling (Hofmann et al., 2003; Kim et al., 2006; Lee et al., 2008;

Shojima et al., 2012).

In general, mammalian HIPK family members regulate the activity of transcription

factors (TFs), chromatin modifiers, signaling molecules and scaffolding proteins in

response to cellular stress (Calzado et al., 2007; de la Vega et al., 2012; Song and Lee,

2003a, b). For example, nutrient stress, such as glucose deprivation can activate Hipk1

and Hipk2 (Ecsedy et al., 2003; Garufi et al., 2013; Song and Lee, 2003b). Conversely,

hyperglycemia triggers HIPK2 degradation via the proteasome (Baldari et al., 2017).

We sought to identify the tissue- and cell-types in which the HPK-1 transcriptional

circuits act to extend longevity, increase resistance to acute proteotoxic stress, and

which PN components are regulated across tissues. We have discovered that HPK-1

functions as a key regulator of the proteostatic response, originating in the nervous

system of *C. elegans*. We find that *hpk-1* is the most broadly upregulated kinase during

normal *C. elegans* aging, predominantly within the nervous system, and acts to

preserves neuronal function and proteostasis within the nervous system. Further, HPK-

1 kinase activity within the nervous system initiates cell non-autonomous signals that

trigger protective peripheral responses to maintain organismal homeostasis. Responses

can qualitatively differ depending on the neuronal type from which they arise:

serotonergic HPK-1 activity increases thermotolerance and protects the proteome from

acute stress by enhancing the HSR, without increasing lifespan or altering basal

autophagy. In contrast, GABAergic HPK-1 activity induces autophagy to extend

longevity, without altering thermotolerance. HPK-1 overexpression is sufficient to induce

autophagy gene expression, and autophagosome formation requires *hlh-30* (TFEB),

mxl-2 (MLX), and *daf-16* (FOXO) transcription factors. We posit that serotonergic HPK-1

signaling responds to acute proteotoxic stress, whereas GABAergic HPK-1 responds to

chronic metabolic changes mediated through target of rapamycin complex 1 (TORC1)

(Das et al., 2017). Yet, each of these distinct adaptive responses is sufficient to activate

the PN and improve proteostasis outside of the nervous system. Our findings reveal

HPK-1 as a key neuronal regulator in coordinating adaptive metabolic and stress

response pathways across tissues to delay aging throughout the animal.

Results

HPK-1 acts from neuronal tissue to promote longevity and health span.

To precisely determine where HPK-1 functions in the regulation of longevity, we

overexpressed *hpk-1* within neuronal (*rab-3p*), muscle (*myo-3p*), intestine (*ges-1p*) or

hypodermal (*dpy-7p*) tissues to identify where activity was sufficient to increase lifespan.

Increased neuronal expression of *hpk-1* was sufficient to increase lifespan by 17%,

which is comparable to constitutive overexpression of *hpk-1* throughout the soma

(*Figure 1A*) (Das et al., 2017). In contrast, transgenic animals overexpressing *hpk-1* in

either muscular, intestinal, or hypodermal cells had no significant change in lifespan

(*Figure 1B-D*). These results were somewhat unexpected, as we previously found that

loss of *hpk-1* solely within either neuronal, intestinal, or hypodermal cells is sufficient to

shorten lifespan (Das et al., 2017). *hpk-1* is broadly expressed during embryogenesis

and larval development, but HPK-1 protein is only expressed within the nervous system

in adult animals (Berber et al., 2013; Berber et al., 2016; Das et al., 2017; Raich et al.,

2003). Furthermore, *hpk-1* and orthologs are known to have roles in development,

differentiation and cell fate-specification (Berber et al., 2013; Blaquiere and Verheyen,

2017; Calzado et al., 2007; Hattangadi et al., 2010; Rinaldo et al., 2007; Rinaldo et al.,

2008; Steinmetz et al., 2018). Therefore, we conclude that HPK-1 functions primarily

within the nervous system to regulate aging.

We performed a locomotion assay to test whether neuronal *hpk-1* could rescue

an age-associated decline in movement, a readout of *C. elegans* healthspan (Berber et

al., 2016; Das et al., 2017; Herndon et al., 2002; Huang et al., 2004; Samuelson et al.,

2007). As expected, *hpk-1(pk1393)* null mutants displayed a reduced frequency of body

bends at day 2 of adulthood (*Figure 1E*). Increased expression of *hpk-1* in neurons was

sufficient to partially mitigate the locomotion defect of *hpk-1* null mutants in response to

active stimuli. Next, we tested whether restoring *hpk-1* exclusively within the nervous

system would prevent the premature age-associated decline in movement of *hpk-1* null

mutant animals. Indeed, restoring *hpk-1* neuronal expression rescued the premature

age-associated decline in movement of *hpk-1* null mutant animals (*Video 1*). Thus,

HPK-1 activity within the nervous system is sufficient to improve healthspan. Lastly,

neuronal *hpk-1* rescue was sufficient to restore a wild-type lifespan in *hpk-1* null mutant

animals (*Figure 1F*). It should be noted this same transgene increases wild-type lifespan

(*Figure 1A*), which suggests HPK-1 retains essential functions for adult healthspan and

lifespan outside of the nervous system; we posit these functions are related to

development as HPK-1 is no longer expressed outside of the nervous system in adult

animals (Das et al., 2017).

We next asked whether neuronal *hpk-1* expression would be sufficient to rescue

the ability of animals to survive acute stress. We and others previously found that in the

absence of *hpk-1*, animals are vulnerable to thermal stress (Berber et al., 2016; Das et

al., 2017). Restoring *hpk-1* expression solely within the nervous system was sufficient to

rescue the reduced thermotolerance of *hpk-1* null mutant animals (*Figure 1G*).

Collectively, these findings are consistent with the notion that *hpk-1* primarily acts within

the nervous system to regulate aging and stress resistance.

hpk-1 expression increases broadly during normal aging.

We sought to determine whether *hpk-1* expression changes during normal aging.

hpk-1 mRNA expression increases in wild-type animals during normal aging between

day 2 and 10 or adulthood (*Figure 2A*). Adult *C. elegans* are composed of 959 somatic

cells, of which 302 are neuronal; all somatic cells are all post-mitotic and arise from an

invariant lineage. A recent comprehensive single-cell analysis of age-associated

changes in gene expression throughout the *C. elegans* identified 211 unique "cell-

clusters" within the aging soma: groups of cells within a lineage or tissue defined by

gene expression profiles (Roux et al., 2022). The "*C. elegans* Aging Atlas" analysis

identified 124 neuronal cell clusters and obtained age-associated changes in gene

expression for 122, with high concordance of overall gene expression between neuron

clusters to a previous project, which mapped the complete gene expression profiles of

the *C. elegans* nervous system in late larval animals (The CeNGEN project (Taylor et

al., 2021)). The *C. elegans* Aging Atlas provides the means to investigate age-

associated changes in gene expression at an unprecedented level of resolution.

Using the *C. elegans* Aging Atlas dataset, in which differentially expressed genes

were identified within each cell cluster in old versus young animals, we find that *hpk-1*

expression increases during aging within 48 cell clusters: broadly within the nervous

system and in eleven non-neuronal cell clusters (*Figure 2B--figure supplement 1*, see

Materials and methods). With aging, neuronal expression of *hpk-1* increased within

interneurons, motor neurons, sensory neurons, and all neuronal sub-types (cholinergic,

dopaminergic, GABAergic, serotonergic, and glutamatergic). We found no cell-clusters

in which *hpk-1* is downregulated during normal aging. It is well known that many

longevity-associated TFs increase lifespan when overexpressed and shorten lifespan

when lost (reviewed in (Denzel et al., 2019)), which infers these TFs might be

upregulated, or become chronically activated, during normal aging. A major finding of

the *C. elegans* Aging Atlas study was that many longevity-associated TFs are

upregulated during normal aging in more than 200 cell-type clusters (Roux et al., 2022).

Increased expression of longevity-associated TFs is thought to reflect an age-

associated adaptive response to dysregulation of homeostatic mechanisms and

accumulating damage (Roux et al., 2022). We posit *hpk-1* neuronal expression

increases during normal aging in wild-type animals to mitigate accumulating damage

and cellular dysfunction.

During normal aging, *hpk-1* and longevity-associated TFs are upregulated in

shared subsets of neurons.

We next assessed whether any TFs showed patterns of spatial upregulation

similar to the age-associated increase in expression of *hpk-1*, and then segregated

those cell types into neuronal and non-neuronal subgroups. Seventy-one TFs were

differentially upregulated with *hpk-1* during aging within three or more neuron clusters,

thirty within non-neuronal clusters, with an overlap of twenty-one TFs between neurons

and non-neurons (*Figure 2B--figure supplement 2* and *Supplementary File 5*). Overall,

twenty-six of these TFs have previously been implicated in longevity. There are ten TFs

broadly upregulated with aging in the nervous system that overlap with *hpk-1*

(significant differential expression in 9 or more neuronal clusters), six of which have

previously been linked to longevity: *daf-16* (FOXO, overlaps in 19/37 neuronal clusters)*,*

hlh-30 (TFEB, 18/37)*, dpy-22* (mediator complex subunit 12L, 15/37), *gei-3* (10/37), *skn-*

1 (NRF2, 10/37)*,* and *hif-1* (hypoxia inducible factor 1, 10/37) (*Figure 2C*) (An and

Blackwell, 2003; Chen et al., 2009; Fawcett et al., 2015; Kenyon et al., 1993; Lapierre et al., 2013; Lin et al., 1997; Ogg et al., 1997; Roux et al., 2022; Suriyalaksh et al., 2022; Tacutu et al., 2012; Zhang et al., 2009). Conversely, 12 TFs exhibited an inverse correspondence, and were largely downregulated with aging in the same neurons where *hpk-1* is normally upregulated (significant differential expression in 9 or more neuronal clusters) (*Figure 2D*); only two of which have previously been implicated in longevity: *xbp-1* (X-box binding protein 1) a key regulator of the endoplasmic reticulum unfolded protein response (ER-UPR), and *mdl-1* (MAX dimerization protein) (Johnson et al., 2014; Nakamura et al., 2016; Riesen et al., 2014). Of note, *mdl-1* is a Myc-family member with opposing roles in longevity from *mxl-2* (Mlx) (Johnson et al., 2014), and *mxl-2* is required for *hpk-1* to extend longevity. (Das et al., 2017).

We tested whether we could verify changes in age-associated gene expression for several candidate longevity-associated transcription factors identified in the Aging Atlas study. Age-synchronized wild-type animals at either day 1 or 11 of adulthood were harvested and mRNA levels of *hpk-1, xbp-1* and *gei-3* were assessed. As expected, both *hpk-1* and *gei-1* were significantly induced in older animals, and conversely *xbp-1* expression was decreased (*Figure 2E*)**.** We conclude that we can recapitulate results obtained from the Aging Atlas study and confirm that longevity-associated TFs are significantly differentially expressed during normal aging. Next we determined whether *hpk-1* expression is significantly induced within the *C. elegans* nervous system during normal aging; we assessed age-associated changes neuronal *hpk-1* mRNA levels *in vivo* (*hpk-1p::mCherry*). As expected, we find that *hpk-1* is significantly upregulated within isolated *C. elegans* neurons during normal aging (*Figure 2F).*

In non-neuronal clusters we found a much smaller subset of TFs with

corresponding positive age-associated fold-changes. Only six of these TFs have been

implicated in longevity: *fkh-9* (FOXG), *sea-2*, *zfh-2* (ZFHX3) (Walter et al., 2011), *xbp-1,*

mdl-1, and *hlh-6* (achaete-scute family bHLH transcription factor 3) (*Figure 2--figure*

supplement 1).

Within neuronal cell-clusters, *hpk-1* and the aforementioned longevity-associated

TFs are broadly expressed throughout the nervous system by late larval development,

with the exception of the nematode specific gene *gei-3*, for which overlap is restricted to

a subset of neurons (*Figure 2--figure supplement 3*)*.* TFs with age-associated increases

in expression in clusters overlapping with *hpk-1* are associated with positive regulation

of adult lifespan, heat and oxidative stress response, bacterial innate immunity,

chromatin remodeling, development, mitophagy, axon regeneration, neddylation, and

ubiquitination (*Figure 2C*). TFs with decreased expression in aging in *hpk-1*-upregulated

neuronal cell clusters have been implicated in the ER unfolded protein response and the

innate immune response (*Figure 2D*). Interestingly, when assessed at the level of

individual time points rather than "old" and "young" groups, we find that the age-

associated up-regulation of *hpk-1* and expression changes of longevity-associated TFs

does not necessarily occur in a tightly-linked manner suggestive of a single upstream

regulatory process (see Appendix 1). Rather, we posit *hpk-1* is induced within the

nervous system along with key longevity transcription factors during normal aging in

response to accumulating stressors and collapsing homeostatic mechanisms to mitigate

the physiological effects of aging.

hpk-1 is the most broadly upregulated kinase during normal aging

We were surprised to find that *hpk-1* mRNA expression increased during aging,

as kinase regulation at the level of gene expression seemed atypical. Therefore, using

the *C. elegans* Aging Atlas dataset we assessed whether any of the 438 *C. elegans*

kinases (Zaru et al., 2017) are differentially expressed during aging: 115 kinases are

upregulated and conversely 120 are downregulated in at least one cell cluster, while

254 kinases do not change expression (*Figure 2--figure supplement 4, Supplementary*

File 6). *hpk-1* is upregulated in more cell-clusters than any other kinase, while *hpk-1* is

upregulated in 48, the next most broadly upregulated kinase is *gcy-28* (natriuretic

peptide receptor 1, 38 cell-clusters) (Li and van der Kooy, 2018; Shinkai et al., 2011;

Tsunozaki et al., 2008). Only 14 kinases are downregulated in 10 or more cell-clusters.

Conversely, 20 kinases are upregulated in ten or more cell-clusters (*Figure 2--figure*

supplement 4). In addition to *hpk-1* and *gcy-28*, five kinases previously linked to

longevity are upregulated: *nipi-3* (a Tribbles pseudokinase that regulates SKN-1 (Wu et

al., 2021))*, aak-1* (AMP-activated kinase catalytic subunit alpha-2) (Lee et al., 2009),

scd-2 (ALK receptor tyrosine kinase) (Shen et al., 2007), *rsks-1* (ribosomal protein S6

kinase) (Hansen et al., 2007; Pan et al., 2007), and *smg-1* (SMG1) (Masse et al., 2008).

We previously demonstrated that HPK-1 is an essential component of TORC1-

mediated longevity and autophagy (Das et al., 2017). Interestingly we find that only two

of 11 TOR-associated genes: *rsks-1* and *aak-1,* are upregulated during aging in three or

more cell-clusters (Blackwell et al., 2019; Hindupur et al., 2015; Pan et al., 2007). We

find *rsks-1* is exclusively upregulated outside of the nervous system and has poor

overlap with *hpk-1* (2/13). *aak-1* is primarily but not exclusively upregulated outside of

the nervous system during aging, with some overlap with *hpk-1*. In future studies it will

be interesting to dissect whether HPK-1 intersects with these components of TORC1

signaling within and/or outside of the nervous system.

HPK-1 preserves integrity of the *C. elegans* nervous system.

Based on the neuronal HPK-1 capacity to promote longevity, we sought to

determine whether HPK-1 preserves the anatomy and physiology of the nervous system

during aging. To assess age-associated changes in neuroanatomy of ventral and dorsal

cord motor neurons (VD and DD), we used the *unc-25p::GFP* cell-specific reporter

(*Figure 3A*) (Cinar et al., 2005; Kraemer et al., 2003). Aged populations of *hpk-1* mutant

animals displayed more than a 3-fold difference in VD axonal breaks and more than a 2-

fold difference in DD axonal breaks compared to wild type animals (*Figure 3A-B, D*).

These age-associated breaks could be rescued by neuronal expression of *hpk-1* (*Figure*

3C). Importantly, young wild-type and *hpk-1* mutant animals did not show neuronal

breaks (*Figure 3B*). Thus, loss of *hpk-1* does not disrupt axon formation during

development, rather HPK-1 preserves axonal integrity in aging motor neurons.

In *C. elegans,* locomotion is controlled by acetylcholine-releasing excitatory

motor neurons and GABA (γ-aminobutyric acid)-releasing-inhibitory motor neurons

(Jorgensen and Nonet, 1995; Richmond and Jorgensen, 1999). Thus, locomotion is a

useful readout to assess neurotransmission. To examine the role of *hpk-1* in synaptic

transmission, we performed a paralysis assay after acetylcholinesterase inhibition with

aldicarb, where mutations that impair cholinergic signaling cause a reduction in rate of

paralysis (Mahoney et al., 2006; Oh and Kim, 2017). Therefore, mutants with lower

synaptic transmission undergo reduced paralysis (i.e., animals are resistant), while

mutants with higher synaptic transmission exhibit greater sensitivity. Interestingly, *hpk-1*

mutants showed increased aldicarb resistance by L4/YA (*Figure 3E*), suggesting that

loss of *hpk-1* reduces cholinergic synaptic transmission and that HPK-1 normally

preserves neuronal function.

C. elegans serotonin (5-HT) signaling in conjunction with dopamine (DA)

modulates locomotion; animals treated with exogenous 5-HT exhibit paralysis over time

(Hardaker et al., 2001; Omura et al., 2012). In a paralysis assay with exogenous 5-HT,

we observed that late L4/YA *hpk-1* mutant animals paralyzed sooner: at 2 minutes of 5-

HT treatment, 63% of mutants were immobilized compared to 25% of wild type animals

immobilized (*Figure 3F*). Based on the aforementioned findings, we conclude that HPK-

1 prevents an age-associated decline in axonal and synaptic integrity.

Loss of *hpk-1* results in neuronal dysregulation of neuronal gene expression

We sought to gain deeper insight into how loss of *hpk-1* negatively impacted the

C. elegans nervous system. To identify early transcriptional dysregulation events that

occur due to the absence of *hpk-1*, we conducted bulk RNA-sequencing of day 2 adult

wild-type and *hpk-1(pk1393)* null mutant animals, an age at which axonal breaks have

yet to occur (*Figure 3B, D*), but after neuronal function has begun to decline (*Figure 3E,*

F). Differential expression analysis identified 2201 total genes with significant

expression changes with loss of *hpk-1*, 1853 upregulated and 348 downregulated

(adjusted p-value < 0.05 and $\log_2$ foldchange magnitude ≥ 1) (*Figure 4A* and

Supplementary File 10). Functional enrichment with over-representation analysis was

performed for the up- and down-regulated genes separately (GOSeq, adjusted p-value

< 0.01). Among down-regulated genes, we find 25 genes implicated in the innate

immune response, as well as TGF-beta and WNT signaling (five ubiquitin ligase

complex component genes), glycosphingolipid metabolism, and surfactant metabolism.

(*Figure 4B*). By contrast, more than 50 GO terms and pathways were enriched for the

upregulated genes. The most significantly enriched single term reveals a substantial

number of cuticle and collagen associated genes as dysregulated (*Figure 4C*).

Upregulation of some collagens and ECM-remodeling processes have been associated

with longevity in *C. elegans* (Ewald et al., 2015; Rahimi et al., 2022). Alternatively,

aberrant upregulation could lead to stiffening of the cuticle, a hallmark of aged *C.*

elegans and some progeric mutants (Ewald et al., 2015; Rahimi et al., 2022).

A striking number of other functions significantly enriched in the *hpk-1*

upregulated genes from across GO, KEGG, and Reactome are related to neuronal

function and signaling, including: neuropeptide and neurotransmitter pathways,

mechanosensation, chemotaxis, and axon guidance, among others (*Figure 4C*). To

explore further, we looked at the *hpk-1* null upregulated genes in the context of a

curated compendium of genes with functions that are important for the nervous system

(Hobert, 2013), as well as genes identified as uniquely expressed in neurons or

significantly enriched in neurons from tissue-specific bulk RNA-Seq (Kaletsky et al.,

2018).

Among 1496 genes known, or expected through homology, to function in the

nervous system, 264 are upregulated with loss of *hpk-1*, representing a much larger

overlap than expected by chance (hypergeometric test p-value < 0.0001)

(*Supplementary File 11*). Genes with higher fold-change of expression in *hpk-1* mutants

were associated with ciliated sensory neurons (motor proteins, sensory cilia transport),

calcium binding, and neuropeptides. We find that 55 of these neuron-associated genes

are normally upregulated with age in wild-type animals in at least one cell cluster in the

C. elegans Aging Atlas, 87 are downregulated with age, with the remaining having

mixed or no significant age-associated changes. This suggests that while loss of *hpk-1*

results in expression of some genes associated with aging, the bulk of differentially

expressed genes at day 2 of adulthood are consistent with dysregulation of neuronal

gene expression. Refined genetic perturbation within the nervous system coupled with

single cell RNA-sequencing over time would be required to fully reveal the impact of

HPK-1 function on age-associated changes in gene expression.

We next investigated how many of these genes are predominantly- or

exclusively- expressed in neurons, based on a set of 880 neuron-enriched and neuron-

specific genes (Kaletsky et al., 2018). We found that 15% of all *hpk-1* upregulated

genes are enriched or specific to neurons in wild-type animals (283 genes,

hypergeometric test p-value < 0.0001) whereas only five *hpk-1* down-regulated genes

normally exhibit such neuron-restricted expression. To determine what functions are

over-represented among the upregulated neuronally expressed genes, we ran Gene

Ontology enrichment analysis. Representing these results as a network, and clustering

terms which have overlapping gene associations, we identified some major themes:

neurotransmitter and ion transport, synaptic components and signaling, axon and

neuron projection components, feeding and locomotion behavior, mechanosensation

and touch-responsive behavior are all upregulated in the absence of *hpk-1* (*Figure 4D*).

Among all genes upregulated in the *hpk-1* null mutants, the largest group correspond to

genes related to neuropeptide encoding, processing and receptors. Neuropeptides are

short sequences of amino acids important for neuromodulation of animal behavior,

broadly expressed in sensory, motor and interneurons. Of the 120 neuropeptide-

encoding genes in the *C. elegans* genome, 40 (33%) are upregulated in the *hpk-1* null

mutants. Nine of these are insulin-related (*ins*), which suggest increased *hpk-1*

expression during normal aging or *hpk-1* overexpression may positively affect longevity

through decreased *ins* neuropeptide expression. We surmise that many of the neuronal

genes that are induced in the absence of *hpk-1* might be compensatory responses to

maintain neuronal function. We conclude loss of *hpk-1* upregulates expression of a

number of neuronal genes including those involved in: neuropeptide signaling,

neurotransmission, release of synaptic vesicles, calcium homeostasis, and function of

sensory neurons (Appendix 2).

Neuronal HPK-1 regulates proteostasis cell-intrinsically and cell non-

autonomously via neurotransmission.

We hypothesized that HPK-1 maintains neuronal function by fortifying the neural

proteome. We measured polyglutamine toxic aggregate formation in neuronal cells

(*rgef-1p::Q40::YFP*) of animals overexpressing *hpk-1* pan-neuronally (*rab-3p*). These

animals showed a decreased number of fluorescent foci within the nerve ring (*Figure*

5Ai - iii, B). To assess whether this was associated with improved neuronal function, we

assessed alterations in locomotion capacity (body bends) after placement in liquid.

Increased neuronal expression of *hpk-1* significantly increased the average number of

body bends (17.7 ± 1 vs 12.5 ± 1.2 bends/30sec), suggesting that neuronal expression

of *hpk-1* improves proteostasis in the nervous system and mitigates locomotion defects

associated with the decline in proteostasis (*Figure 5C*).

We next tested whether HPK-1 was necessary to preserve proteostasis within

neurons. We crossed the *rgef-1::Q40::YFP* reporter with neuronal enhanced RNAi

background (*sid-1(pk3321);unc-119p::sid-1*) and assessed whether neuronal

inactivation of *hpk-1* hastened age-associated proteostatic decline. Animals with

neuronal *hpk-1* inactivation had a significant increase in foci number within neurons and

a decreased number of body bends in liquid (19.7 $\pm$ 0.8 vs 25.5 $\pm$ 1.1 bends/30sec)

(*Figure 5Aiv, D, E*). Thus, HPK-1 functions cell intrinsically to maintain neuronal

proteostasis.

We next assessed whether neuronal HPK-1 activates a cell non-autonomous

signal to regulate proteostasis in distal tissues. We used a proteostasis reporter

expressed exclusively in muscle tissue (*unc-54p::Q35::YFP*); animals display an age-

associated accumulation of fluorescent foci, which are proteotoxic aggregates, and

become paralyzed as muscle proteome function collapses (Lazaro-Pena et al., 2021;

Morley et al., 2002). Animals with increased neuronal *hpk-1* expression had significantly

decreased fluorescent foci in muscle tissue (*Figure 5Fi-ii*). For instance, at day 3 of

adulthood, animals overexpressing neuronal HPK-1 had an average of 21 $\pm$ 0.9 polyQ

aggregates, while control animals had an average of 26 $\pm$ 1.1 polyQ aggregates (*Figure*

5G). In *C. elegans*, the age-related progressive proteotoxic stress of muscle polyQ

expression is pathological and results in paralysis. Of note, increased *hpk-1* expression

in neurons was sufficient to reduce paralysis; at day 13 of adulthood only 61% $\pm$ 0.10 of

neuronal *hpk-1* overexpressing animals were paralyzed compared to 81% $\pm$ 0.08 of

paralyzed control animals (*Figure 5H*). Thus, increased neuronal HPK-1 activity is

sufficient to improve proteostasis cell non-autonomously.

We confirmed that the kinase activity is essential for HPK-1 to improve

proteostasis. We find the kinase domain of HPK-1 is highly conserved from yeast to

mammals, both at the amino acid level and predicted tertiary structure, based on

multiple sequence alignments, AlphaFold structure prediction, and visualization with

PyMol (*Figure 5--figure supplement 1*) (DeLano, 2002; Jumper et al., 2021; Varadi et

al., 2022). We mutated two highly conserved residues within the kinase domain, K176A

and D272N, which are known to be essential for the kinase activity of HPK-1 orthologs

(He et al., 2010), and tested whether these kinase-dead mutant versions could improve

proteostasis when neuronally overexpressed. In contrast to wild-type HPK-1 (*Figure 5G,*

H), neither mutant had any effect on either polyglutamine foci accumulation (*Figure 5--*

figure supplement 1C, D) or paralysis (*Figure 5--figure supplement 1E, F*) in muscle

cells (*unc-54p::Q35::YFP*); thus, increased *hpk-1* expression improves proteostasis

through kinase activity.

We tested whether neuronal HPK-1 is required for maintaining proteostasis in

distal muscle cells. Neuronal inactivation of *hpk-1* (*sid-1(pk3321); unc-119p::sid-1; unc-*

54p::Q35::YFP; hpk-1(RNAi)) was sufficient to hasten the collapse of proteostasis,

significantly increasing the number of polyQ aggregates (45 ± 1 vs 32 ± 1.1 at day 4)

and paralysis rate ($84\% \pm 0.04$ vs $14\% \pm 0.04$ at day 9) (*Figure 5Fiii, I, J*). Thus, HPK-1

function within the nervous system impacts proteostasis in distal muscle tissue, which

implies that HPK-1 coordinates organismal health through cell non-autonomous

mechanisms.

To identify the signaling mechanism through which HPK-1 regulates distal

proteostasis, we tested whether either neuropeptide or neurotransmitter transmission

are essential for cell non-autonomous regulation of proteostasis (using *unc-31* and *unc-*

13 mutant animals*, respectively Figure 5K*). Neuronal expression of *hpk-1* in *unc-31*

mutants improved proteostasis in the absence of neuropeptide transmission and dense

core vesicles, as increased neuronal *hpk-1* expression still delayed the age-associated

accumulation of aggregates within muscle cells. In contrast, in the absence of

neurotransmitter function, neuronal expression of *hpk-1* in *unc-13* mutants failed to

improve proteostasis in muscle tissue (*Figure 5L--figure supplement 2*). We conclude

that HPK-1 mediated longevity-promoting cell non-autonomous signaling occurs through

neurotransmission.

Neuronal HPK-1 induces molecular chaperones and autophagy in distal tissues.

We sought to identify the components of the PN that are cell non-autonomously

regulated via HPK-1 activity in the nervous system. We tested whether increasing levels

of neuronal *hpk-1* alters the expression of molecular chaperones (*hsp-16.2p::GFP*).

Increased neuronal expression of *hpk-1* in wild type animals did not alter the basal

expression of molecular chaperones (*Figure 6A*). In contrast, after heat shock,

increased neuronal *hpk-1* expression resulted in hyper-induction of *hsp-16.2p::GFP*

expression, particularly in the pharynx and head muscles, but surprisingly not within

intestinal cells (*Figure 6B-E*). We conclude that while *hpk-1* is required for the broad

induction of molecular chaperones (Das et al., 2017), neuronal HPK-1 primes

inducibility of the HSR within specific tissues, rather than in a systemic manner

throughout the organism.

We next tested whether increased neuronal HPK-1 activity is sufficient to induce

autophagy. Using two LGG-1/Atg8 fluorescent reporter strains to visualize

autophagosome levels in neurons (*rgef-1p::GFP::LGG-1),* hypodermal seam cells and

intestine (*lgg-1p::GFP::LGG-1)* (Kumsta et al., 2017), we found that increased neuronal

hpk-1 expression significantly induces autophagy in neurons and hypodermal seam cell

(*Figure 6F-I*), but not in the intestine (*Figure 6J, K*). Sustained autophagy and

transcriptional activation of autophagy genes are important components for somatic

maintenance and organismal longevity (Lapierre et al., 2015). Consistently *hpk-1* is

required for the induction of autophagy gene expression induced by TORC1 inactivation

(Das et al., 2017). We tested whether neuronal overexpression of *hpk-1* is sufficient to

induce autophagy gene expression, and found that increased neuronal overexpression

induces expression of *atg-18* (WD repeat domain, phosphoinositide interacting 2) and

bec-1 (Beclin 1) (*Figure 6L*). Thus, increased HPK-1 activity within the nervous system

is sufficient to induce basal levels of autophagy both cell-intrinsically and non-

autonomously, which we posit occurs through increased autophagy gene expression to

extend longevity.

GABAergic and Serotonergic HPK-1 differentially regulate longevity and acute

stress signaling.

hpk-1 is broadly expressed within the *C. elegans* adult nervous system (Das et

al., 2017), but the neuronal cell-types responsible for extending longevity are unknown.

We overexpressed *hpk-1* in different subtypes of neurons and assessed changes in

muscle proteostasis (*unc-54p::Q35::YFP*)*.* Increased *hpk-1* expression in glutamatergic

or dopaminergic neurons was not sufficient to decrease the formation of fluorescent foci

or reduce paralysis (*Figure 7--figure supplement 1*). While it is possible that *hpk-1*

overexpression in these neuronal subtypes failed to reach a threshold effect, these

results suggest that HPK-1 does not act within these neuronal subtypes to cell non-

autonomously regulate proteostasis. In contrast, increased expression of *hpk-1* in either

serotonergic (*tph-1p::hpk-1*) or GABAergic (*unc-47p::hpk-1*) neurons was sufficient to

improve proteostasis in muscle tissue (*Figure 7A-D*). Accordingly, we found that *hpk-1*

is expressed in serotonergic and GABAergic neurons, based on colocalization between

hpk-1 and neuronal cell-type specific reporters (*Figure 7E, F--figure supplement 2*),

consistent with the notion that HPK-1 initiates pro-longevity signals from these neurons.

We sought to determine whether signals initiated from serotonergic and

GABAergic neurons induced similar or distinct components of the PN in distal tissues,

and assessed the overall consequence on organismal health. Since pan-neuronal

expression of *hpk-1* was sufficient to induce autophagy, improve thermotolerance, and

increase lifespan, we chose to dissect whether these three phenotypes were regulated

within distinct neuronal cell-types. Increased HPK-1 serotonergic signaling was

sufficient to improve thermotolerance, without altering lifespan or autophagy (*Figure 8A-*

D). In contrast, increased *hpk-1* expression in GABAergic neurons did not alter

thermotolerance, but induced autophagy and slightly but significantly increased lifespan

(*Figure 8E-H*).

To identify the mechanism through which increased neuronal HPK-1 activity

induces autophagy in distal tissues, we examined serotonergic neurons. As expected,

loss of *tph-1,* which encodes the *C. elegans* tryptophan hydroxylase 1 essential for

biosynthesis of serotonin (Sze et al., 2000), failed to block the induction of autophagy by

increased neuronal *hpk-1* expression (*Figure 8I*, *tph-1(mg280);rab-3p::hpk-1*)*. unc-30*

encodes the ortholog of human paired like homeodomain 2 (PITX2) and is essential for

the proper differentiation of type-D inhibitory GABAergic motor neurons (Jin et al., 1994;

McIntire et al., 1993). Consistent with our previous result demonstrating that only

GABAergic expression of *hpk-1* promotes autophagy, loss of *unc-30* was sufficient to

completely abrogate the induction of autophagy in seam cells after increased pan-

neuronal expression of *hpk-1* (*Figure 8I*, *unc-30(ok613);rab-3p::hpk-1*). We conclude

that HPK-1 activity in serotonergic neurons promotes systemic thermotolerance,

whereas HPK-1 in GABAergic neurons promotes autophagy. We surmise that the two

neuronal subtypes are likely to contribute synergistically to the extension of longevity.

HPK-1 regulates autophagy gene expression and across diverse species HIPK

family members directly regulate transcription factors (Sardina et al., 2023), therefore

we sought to identify transcription factors essential for HPK-1 overexpression within

GABAergic neurons to induce autophagy. *pha-4* (FOXA), *mxl-2* (MLX), *hlh-30* (TFEB),

daf-16 (FOXO) and *hsf-1* are all required in specific contexts for the induction of

autophagy, and longevity (Hansen et al., 2008; Kumsta et al., 2017; Lapierre et al.,

2013; Lapierre et al., 2015; Nakamura et al., 2016; O'Rourke and Ruvkun, 2013).

Furthermore, we previously found that *hsf-1, mxl-2, or pha-4* are all necessary for the

increased lifespan conferred by *hpk-1* overexpression throughout the soma (Das et al.,

2017). Using feeding based RNAi, we found that *hpk-1* overexpression in GABAergic

neurons still significantly induced autophagosomes in seam cells after loss of *hsf-1* or

pha-4 (*Figure 8J*); thus HPK-1 induction of autophagy is independent of *hsf-1* and *pha-4*

and suggests the *pha-4* requirement for HPK-1 mediated longevity is independent of

autophagy gene regulation (at least in hypodermal seam cells). In contrast, after

inactivation of *mxl-2*, *hlh-30* or *daf-16,* GABAergic overexpression of *hpk-1* failed to

significantly increase autophagosomes (*Figure 8J*); thus HPK-1 mediated induction of

autophagy requires *mxl-2*, *hlh-30*, and *daf-16*. We were puzzled to find that the average

number of autophagosome in control animals (i.e., lacking *hpk-1* overexpression) varied

after gene inactivation (*Figure 8--figure supplement 1A*). To rigorously test whether *hpk-*

1 overexpression required *hlh-30* for the induction of autophagosomes, we crossed the

hlh-30 null mutation into animals overexpressing *hpk-1* within GABAergic neurons (*unc-*

47p::hpk-1;hlh-30(tm1978);lgg-1p::GFP::LGG-1; we find that *hlh-30* is required for the

induction of autophagosomes in seam cells when *hpk-1* is overexpressed in GABAergic

neurons (*Figure 8--figure supplement 1B).*

We sought to fortify our previous findings that: 1) TORC1 negatively regulates

hpk-1 under basal conditions; 2) *hpk-1* is essential for decreased TORC1 signaling to

increase longevity and induce autophagy; and 3) HPK-1-mediated longevity requires

mxl-2, the ortholog of mammalian Mlx transcription factor (Das et al., 2017). From these

data, we predicted that TOR-mediated longevity might require *mxl-2* via *hpk-1.* Indeed,

we find that inactivation of either *daf-15* (Raptor) or *let-363* (TOR), increased lifespan

and required *mxl-2*, as *mxl-2(tm1516)* null mutant animals failed to increase lifespan in

the absence of TORC1 (*Figure 8--figure supplement 2*). MXL-2 functions as a

heterodimer with MML-1 (Pickett et al., 2007), which have similar roles in *C. elegans*

longevity (Johnson et al., 2014). Interestingly, one of the human homologs of MML-1,

Carbohydrate Response Element Binding Protein

(ChREBP/MondoB/MLXIPL/WBSCR14), maps within a deleted region in Williams-

Beuren Syndrome patients, which among other symptoms causes metabolic

dysfunction, silent diabetes, and premature aging (Pober, 2010). This further supports

our model and is consistent with other studies linking the Myc-family of transcription

factors to longevity, autophagy, and TORC1 signaling (Johnson et al., 2014; Nakamura

et al., 2016; Samuelson et al., 2007). Collectively, we conclude that HPK-1 regulates

serotonergic signals to improve survival in acute thermal stress, likely through HSF-1,

and initiates GABAergic signals to extend longevity through increased autophagy, which

is linked to decreased TORC1 signaling and regulation of the Myc TFs.

Discussion

It has been known that a core group of TFs integrates various types of metabolic

and stress signals; moreover, this group of TFs can extend longevity when

overexpressed, and conversely hasten aging when lost (reviewed in (Denzel et al.,

2019)), but whether the activity or expression of these longevity-associated

transcriptional regulators are activated by signals of increasing damage or dysfunction

during normal aging was unknown. Recently, single-cell RNA sequencing has revealed

that these longevity-associated TFs increase in expression during aging in wild-type

animals across multiple somatic tissues (Roux et al., 2022). This comprehensive

empirical dataset provides foundational support for a key theme in aging research:

during aging in wild-type animals, increasing damage and dysfunction is mitigated by

the activation of stress response programs. Our findings that: 1) *hpk-1* expression is

upregulated during normal aging in more neurons and non-neuronal cells than any other

kinase, and 2) that *hpk-1* is induced in cell-clusters that overlap with longevity-

associated TFs, including *daf-16, hlh-30, skn-1* and *hif-1*, implies that HPK-1 is a key

transcriptional regulator that mitigates aging. We posit this response is activated in the

nervous system during normal aging in wild-type animals to mitigate increasing stress

and the breakdown of homeostatic mechanisms within the nervous system.

Consistently, the presence of human neurodegenerative disorders, a severe form of

neuronal proteotoxic stress, correlates with activation of human *hpk-1* expression: for

example*,* human HIPK2 is activated in frontal cortex samples of patients with AD, in

motor neurons of ALS patients, and in mouse models of ALS, (Lee et al., 2016). Thus,

our results suggest that overexpression of *hpk-1* in the *C. elegans* nervous system may

extend longevity by preemptively delaying the accumulation of age-associated damage

and dysfunction to the proteome (*Figure 9A*). However, *hpk-1* homologs in more

complex metazoans can also promote cell death, a key homeostatic component of

tissue regeneration, indicating that chronic activation of HIPKs is not always beneficial

(Calzado et al., 2007; Lee et al., 2016; Wang et al., 2020; Zhang et al., 2020). Whether

activated HIPKs will fortify proteostasis and extend longevity, or promote cell death is

likely to be dependent on many factors, including the specific homolog, levels of

expression, physiological context, genetic background, and cell-type.

We find that *hpk-1* prevents age-associated decline of axonal and synaptic

integrity, which together preserve neuronal function during the aging process and thus

also promote longer healthspan. In support of this concept, adult *hipk2 -/-* mice display

normal neurogenesis but manifest delayed-onset severe psychomotor and behavioral

deficiencies, as well as degeneration of dopaminergic neurons, reminiscent of

Parkinson's Disease (Zhang et al., 2007). Loss of *hpk-1* results in dysregulation of

neuronal gene expression by early adulthood; some changes are consistent with

neuronal aging but in other cases we observe an inverse correlation. For example, one

third of all neuropeptides are induced in the absence of *hpk-1*; yet increased

neuropeptide signaling has been associated with extended *C. elegans* longevity and is

considered protective in AD (Chen et al., 2018; Hodge et al., 2022; Petrella et al., 2019).

One possibility is in the absence of *hpk-1*, attenuated inducibility of the proteostatic

network accelerates age-associated cellular dysfunction (*Figure 9A*), and upregulation

of neuropeptide signaling compensates in the nervous system for the loss of

proteostatic response. This concept is consistent with our finding that increasing

neuronal HPK-1 activity does not require neuropeptides to improve proteostasis in distal

tissues. An alternative possibility for the connection between protective effects of HPK-1

and neuropeptides in neurons is that HPK-1 may reduce insulin signaling by

suppressing the processing of *ins* peptides, thereby decreasing ILS to activate DAF-16

(FOXO) transcriptional quality control programs. For example, loss of *egl-3*, a

neuropeptide processing gene, extends longevity by inactivating processing of insulins

(Hamilton et al., 2005), and *egl-3* is upregulated in *hpk-1* loss of function.

Impairment of the proteostatic network after loss of *hpk-1* results in broad

upregulation of gene expression, which may be provide additional levels of

compensatory mechanisms to mitigate neuronal dysfunction. For example, loss of *hpk-1*

induces expression of intraflagellar transport (IFT) components in sensory cilia.

Mutations within sensory neurons can alter longevity (Alcedo and Kenyon, 2004; Apfeld

and Kenyon, 1999), and decreased IFT during *C. elegans* aging impairs sensory

perception and metabolism (Zhang et al., 2021). Deficiencies in primary cilia impairs

neurogenesis in AD transgenic mice, resulting in accumulation of $A\beta_{42}$ and tau protein

and disease progression (Armato et al., 2013; Chakravarthy et al., 2012; Ma et al.,

2022; Morelli et al., 2017). However, our bulk RNA-seq analysis cannot decipher

changes in gene expression at the level of individual neurons after loss of *hpk-1. hpk-1* and orthologs are known to have roles in development, differentiation and cell fate-specification (Berber et al., 2013; Blaquiere and Verheyen, 2017; Calzado et al., 2007; Hattangadi et al., 2010; Rinaldo et al., 2007; Rinaldo et al., 2008; Steinmetz et al., 2018), therefore we cannot preclude the possibility that HPK-1 may function to broadly limit neuronal gene expression during the course of normal development and differentiation, thereby specifying neuronal transcriptional programs. Mis-differentiation of the nervous system in the absence of *hpk-1* could result in both intracellular transcriptional noise and cell non-autonomous noise across cell-types and tissues.

We show that HPK-1 expression in serotonergic neurons activates a cell non-autonomous signal to prime induction of molecular chaperones in muscle cells, whereas HPK-1 expression in GABAergic neurons signals induces autophagy in hypodermal tissue (*Figure 9B*). Our findings place HPK-1 within a selective cohort of transcriptional regulators within the *C. elegans* nervous system that are activated in response to metabolic, environmental, and intrinsic cues; thereby initiating distinct cell non-autonomous signals to regulate components of the PN in peripheral tissues (Douglas et al., 2015; Durieux et al., 2011; Frakes et al., 2020; Imanikia et al., 2019; Prahlad et al., 2008; Prahlad and Morimoto, 2011; Silva et al., 2013; Tatum et al., 2015; Taylor and Dillin, 2013). Loss of serotonin production in *tph-1* mutant animals has a minimal impact on lifespan at basal temperature (Murakami and Murakami, 2007). However, simultaneous loss of both serotonin and dopamine biosynthesis shortens lifespan (Murakami and Murakami, 2007). Serotonin biosynthesis is also required for several conditions that extend longevity: animals treated with an atypical antidepressant

serotonin antagonist (mianserin) (Petrascheck et al., 2007), changes in food perception

(Entchev et al., 2015), and neuronal cell non-autonomous induction of the mitochondrial

unfolded protein response (Berendzen et al., 2016; Zhang et al., 2018). In contrast, loss

of either of the serotonin receptors *ser-1* or *ser-4* have opposing effects on lifespan,

suggesting antagonistic activity between serotonin receptors (Murakami and Murakami,

2007).

Surprisingly, we find that increased HPK-1 activity in serotonergic neurons does

not increase lifespan, but is sufficient to improve muscle proteostasis and amplify the

expression of molecular chaperones that maintain proteostasis within the cytosol and

nucleus. Interestingly, neuronal HSF-1 regulation of the HSR through thermosensory

circuits is also separable from lifespan (Douglas et al., 2015). Serotonin signaling and

the serotonin receptor *ser-1* are essential for the induction of the HSR (Prahlad et al.,

2008; Tatum et al., 2015). ADF serotoninergic neurons receive upstream stress signals

from the thermosensory circuit, consisting of the amphid sensory neurons (AFD) and

the AIY interneurons, which regulate HSF-1 activity in the germline in a serotonin-

dependent manner (Tatum et al., 2015). However, increased serotonergic HPK-1

activity is not equivalent to altered of thermosensory signals: *hpk-1* improves

proteostasis within body wall muscles but not intestinal cells through neurotransmitters

and independent from dense core vesicles. In contrast, loss of the guanylyl cyclase

(*gcy-8*) in AFD thermosensory neurons or the LIM-homeobox transcription factor (*ttx-3*)

in AIY interneurons improves intestinal proteostasis through dense core vesicle

neurosecretion (i.e., *unc-31*). *hpk-1* is expressed in AFD, but only at very low levels in

AIY neurons (data not shown and (Taylor et al., 2021)), and *hpk-1* is essential for the

induction of the HSR within intestinal cells (Das et al., 2017). Whether *hpk-1* activity

within thermosensory neurons functions within the canonical thermosensory-serotonin

circuit to regulate the intestinal HSR or proteostasis is not mutually exclusive from our

findings and will be interesting to pursue in future studies. Our results suggest

serotonergic HPK-1 is an important regulator of the HSR, potentially functioning in

parallel to the thermosensory-serotonin circuit to restore proteostasis in response to

acute thermal stress.

We posit that increased serotonergic HPK-1 activity further activates HSF-1 in

response to heat (*Figure 9B*). This would be consistent with known

hyperphosphorylation of HSF-1 and amplification of the HSR in response to increasingly

severe heat stress (reviewed in (Lazaro-Pena et al., 2022)). Interestingly, HPK-1 limits

SUMOylation of HSF-1, SUMO limits the heat shock response in both *C. elegans* and

mammals, and severe heat shock results in HSF1 deSUMOylation (Das et al., 2017;

Hietakangas et al., 2003; Hietakangas et al., 2006). To our surprise, while increased

neuronal *hsf-1* expression increases longevity (Douglas et al., 2015; Morley and

Morimoto, 2004), we could find no study that has identified specific neuronal cell-types

in which increased HSF-1 activity is sufficient to extend longevity or enhance the HSR.

Whether or not HPK-1 regulates HSF-1 within the nervous system and/or in peripheral

muscle cells through cell non-autonomous signals will be interesting to determine in

future studies. HPK-1 co-localizes with HSF-1 in neurons, HPK-1 protein is stabilized

after heat shock, *hpk-1* is essential for induction of the HSR, and *hsf-1* and *hpk-1* are

mutually interdependent to extend longevity (Das et al., 2017) (*Figure 9B*).

Increased HPK-1 activity in GABAergic neurons induces autophagy in peripheral

cells, improves proteostasis and longevity. Whether HPK-1 functions in GABAergic

neurons through alteration of GABA or another neurotransmitter is unknown, but proper

differentiation of GABAergic neurons is essential for neuronal HPK-1 activation of

autophagy in distal tissue. We posit HPK-1 cell non-autonomous signaling occurs via

antagonism of GABA signaling. Loss of GABA increases *C. elegans* lifespan (Chun et

al., 2015; Yuan et al., 2019). However, not all neurotransmitters regulate lifespan, as

only mutants for the GABA biosynthetic enzyme, *unc-25*, increase lifespan. This

GABAergic-mediated control in lifespan seems to be mediated by the *gbb-1*

metabotropic GABA$_B$ receptor (Chun et al., 2015). Interestingly, GABA signaling inhibits

autophagy and causes oxidative stress in mice, which is mitigated via Tor1 inhibition

(Lakhani et al., 2014). In fact, we previously found that neuronal expression of HPK-1 is

inhibited by TORC1 and *hpk-1* is required for TORC1 inactivation to extend longevity

and the induction of autophagy gene expression (Das et al., 2017). Both TORC1

inactivation and *hpk-1* overexpression requires the Myc-family member *mxl-2* to extend

longevity (Das et al., 2017; Johnson et al., 2014; Nakamura et al., 2016) (*Figure 8--*

figure supplement 2). Consistently, TORC1 also acts within neurons to regulate aging

(Zhang et al., 2019); our results suggest that GABAergic neurons might be the specific

neuronal cell-type through which alterations in TORC1 activity regulate longevity.

Collectively, our results position HPK-1 activity in GABAergic neurons as the likely

integration point for cell non-autonomous regulation of autophagy via TORC1 (*Figure*

9B).

A growing number of longevity-associated transcription factors have been linked

to autophagy, including *pha-4* (FOXA), *mxl-2* (MLX), *hlh-30* (TFEB), *daf-16* (FOXO) and

hsf-1 (Hansen et al., 2008; Kumsta et al., 2017; Lapierre et al., 2013; Lapierre et al.,

2015; Nakamura et al., 2016; O'Rourke and Ruvkun, 2013). We find that *hlh-30, daf-16,*

and *hlh-30* are all required for the induction of autophagosomes when *hpk-1* is

overexpressed in GABAergic neurons. We posit these transcription factors respond

within the distal hypodermal seam cells in response to a neuronal signal initiated by

HPK-1, as neurons are largely refractory to RNAi. However, this does not preclude cell

intrinsic neuronal interactions; future refined spatial genetic analysis will be required to

definitively decipher the precise tissues of action. DAF-16 and HLH-30 have previously

been shown to function in combination to promote stress resistance and longevity (Lin

et al., 2018). Analogously, the MML-1::MXL-2 complex and HLH-30 activate each other

and are downstream effectors of longevity after TORC1 inhibition (*mml-1* encodes the

MondoA/ChREBP ortholog, an obligate interactor with MXL-2) (Nakamura et al., 2016).

Based on our discovery that all three of these longevity-associated transcription factors

are essential for GABAergic HPK-1 activity to induce autophagy, it is tempting to

speculate HPK-1 acts as a key mediator of pro-longevity signaling after TORC1

inhibition through coordinated activation of MML-1::MXL-2, HLH-30, and DAF-16

(*Figure 9B*).

This work reveals a new level of specificity in the capacity of the nervous system

to initiate adaptive responses by triggering distinct components of the proteostatic

network across cell types. Results elucidating how neuronal cell-types generate

heterotypic signals to coordinate the maintenance of homeostasis are just beginning to

emerge. For example, a previous study discovered that activating non-autonomous

signaling of the endoplasmic reticulum unfolded protein response (ER-UPR) via the

XBP-1 transcription factor can be initiated from serotonergic neurons and dopaminergic

neurons; these signals are unique and converge to activate distinct branches of the ER-

UPR within the same distal tissues (intestinal cells) (Higuchi-Sanabria et al., 2020). In

contrast, our work demonstrates differential, yet specific, activation of cytosolic

components of the PN in separate peripheral tissues through neurotransmitters. Our

work suggests the nervous system partitions neuronal cell-types to coordinate the

activation of complementary proteostatic mechanisms across tissues, despite utilizing

the same transcriptional regulator within each neuronal cell-type. One possibility is that

HPK-1 regulates specific adaptive responses based on the unique epigenetic states

between differentiated neuronal subtypes, which would be consistent with recent

findings: during development and differentiation, transcriptional rewiring of cells set

chaperoning capacities and alter usage of mechanisms that maintain protein quality

control (Nisaa and Ben-Zvi, 2021; Shemesh et al., 2021). Our work reveals HPK-1 as a

novel transcriptional regulator within the nervous system of metazoans, which integrates

diverse stimuli to coordinate specific adaptive responses that promote healthy aging.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or Reference**	**Identifiers**	**Additional Information**
genetic reagent (*C. elegans*)	N2	N2 (CGCM)		*Wild type*
genetic reagent (*C. elegans*)	AVS392	EK273		*hpk-1(pk1393) X*
genetic reagent (*C. elegans*)	AVS543	This paper	*artEx41*	*artEx41 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)]*
genetic reagent (*C. elegans*)	AVS609	This paper	*artEx43*	*artEx43 [myo-3p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1*
genetic reagent (*C. elegans*)	AVS614	This paper	*artEx48*	*artEx48 [dpy-7p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1*
genetic reagent (*C. elegans*)	AVS627	This paper	*artEx52*	*artEx52 [ges-1p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1*
genetic reagent (*C. elegans*)	AVS420	This paper	*artEx41*	*hpk-1(pk1393) X; artEx41 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)]*
genetic reagent (*C. elegans*)	AVS602	CZ13799	*juIs76*	*juIs76 [unc-25p::GFP + lin-15(+)] II*
genetic reagent (*C. elegans*)	AVS608	This paper	*juIs76*	*hpk-1(pk1393); juIs76 [unc-25p::GFP + lin-15(+)] II*
genetic reagent (*C. elegans*)	AVS752	This paper	*artEx41*	*artEx41 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)]; hpk-1(pk1393); juIs76 [unc-25p::GFP + lin-15(+)] II*
genetic reagent (*C. elegans*)	AVS832	This paper	*rmIs132; artEx58*	*rmIs132 [unc-54p::Q35::YFP] I; artEx58 [rab-3p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1*
genetic reagent (*C. elegans*)	AVS833	This paper	*rmIs132; artEx59*	*rmIs132 [unc-54p::Q35::YFP] I; artEx59 [rab-3p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 2*
genetic reagent (*C. elegans*)	AVS834	This paper	*rmIs132; artEx60*	*rmIs132 [unc-54p::Q35::YFP] I; artEx60 [rab-3p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 3*
genetic reagent (*C. elegans*)	AVS835	This paper	*rmIs132; artEx61*	*rmIs132 [unc-54p::Q35::YFP] I; artEx61 [rab-3p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 4*
genetic reagent (*C. elegans*)	AVS744	This paper	*rmIs132; artEx75*	*artEx75 (rab-3p::hpk-1(K176A) + pCFJ90 (myo-2p::mCherry); rmIs132 [Punc-54::Q35::YFP] I line 1*
genetic reagent (*C. elegans*)	AVS748	This paper	*rmIs132; artEx79*	*artEx79 (rab-3p::hpk-1(D272N) + pCFJ90 (myo-2p::mCherry); rmIs132 [Punc-54::Q35::YFP] I line 1*
genetic reagent (*C. elegans*)	AVS562	This paper	*rmIs132; artEx41*	*rmIs132 [unc-54p::Q35::YFP] I; artEx41 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)] line 5*
genetic reagent (*C. elegans*)	AVS563	This paper	*rmIs110; artEx41*	*rmIs110 [F25B3.3p::Q40::YFP]; artEx41 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)] line 1*
genetic reagent (*C. elegans*)	AVS837	This paper	*rmIs110; artEx92*	*rmIs110 [F25B3.3p::Q40::YFP]; artEx92 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)] line 2*
genetic reagent (*C. elegans*)	AVS838	This paper	*rmIs110; artEx93*	*rmIs110 [F25B3.3p::Q40::YFP]; artEx93 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)] line 3*
genetic reagent (*C. elegans*)	AVS839	This paper	*rmIs110; artEx94*	*rmIs110 [F25B3.3p::Q40::YFP]; artEx94 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)] line 4*
genetic reagent (*C. elegans*)	AVS214	HC196		*sid-1(qt9) V*
genetic reagent (*C. elegans*)	AVS265	TU3401	*uIs69*	*sid-1(pk3321)V; uIs69 [unc-119p::sid-1; myo-2p::mCherry]V*
genetic reagent (*C. elegans*)	AVS540	This paper	*rmIs132; uIs69*	*rmIs132 [unc-54p::Q35::YFP] I; sid-1(pk3321)V; uIs69[unc-119p::sid-1; myo-2p::mCherry]V*
genetic reagent (*C. elegans*)	AVS541	This paper	*rmIs110; uIS69*	*rmIs110 [F25B3.3p::Q40::YFP]; sid-1(pk3321)V; uIs69[unc-119p::sid-1; myo-2p::mCherry]V*
genetic reagent (*C. elegans*)	AVS713	This paper	*rmIs132; artEx41*	*rmIs132 [unc-54p::Q35::YFP] I; artEx41(rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)]; unc-31(e169)V*
genetic reagent (*C. elegans*)	AVS810	This paper	*rmIs132; artEx41*	*rmIs132 [unc-54p::Q35::YFP] I; artEx41 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)]; unc-13(e51)I*

genetic reagent (*C. elegans*)	AVS84	TJ375	*gpIs1*	*gpIs1 [hsp-16.2p::GFP] IV*
genetic reagent (*C. elegans*)	AVS397	This paper	*gpIs1; artEx35*	*gpIs1 [hsp-16.2p::GFP]; artEx35 [sur-5p::hpk-1::CFP +pCFJ90 (myo-2p::mCherry)]*
genetic reagent (*C. elegans*)	AVS399	This paper	*gpIs1; artEx33*	*gpIs1 [hsp-16.2p::GFP]; artEx33 [rab-3p::hpk-1::CFP +pCFJ90 (myo-2p::mCherry)]*
genetic reagent (*C. elegans*)	AVS715	This paper	*artEx41; sqIs24*	*artEx41 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)]; sqIs24 [rgef-1p::GFP::lgg-1 + unc-122p::RFP]*
genetic reagent (*C. elegans*)	AVS716	This paper	*artEx41; sqIs13*	*artEx41 [rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry)]; sqIs13 [lgg-1p::GFP::lgg-1 + odr-1p::RFP]*
genetic reagent (*C. elegans*)	AVS682	This paper	*rmIs132; artEx62*	*artEx62 [tph-1p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1; rmIs132 [unc-54p::Q35::YFP] I*
genetic reagent (*C. elegans*)	AVS683	This paper	*rmIs132; artEx63*	*artEx63 [tph-1p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 2; rmIs132 [unc-54p::Q35::YFP] I*
genetic reagent (*C. elegans*)	AVS685	This paper	*rmIs132; artEx65*	*artEx65 [unc-47p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1; rmIs132 [unc-54p::Q35::YFP] I*
genetic reagent (*C. elegans*)	AVS686	This paper	*rmIs132; artEx66*	*artEx66 [unc-47p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 2; rmIs132 [unc-54p::Q35::YFP] I*
genetic reagent (*C. elegans*)	AVS691	This paper	*rmIs132; artEx71*	*artEx71 [eat-4p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1; rmIs132 [unc-54p::Q35::YFP] I*
genetic reagent (*C. elegans*)	AVS692	This paper	*rmIs132; artEx72*	*artEx72 [eat-4p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 2; rmIs132 [unc-54p::Q35::YFP] I*
genetic reagent (*C. elegans*)	AVS693	This paper	*rmIs132; artEx73*	*artEx73 [cat-2p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1; rmIs132 [unc-54p::Q35::YFP] I*
genetic reagent (*C. elegans*)	AVS694	This paper	*rmIs132; artEx74*	*artEx74 [cat-2p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 2; rmIs132 [unc-54p::Q35::YFP] I*
genetic reagent (*C. elegans*)	AVS809	This paper	*sqIs13; artEx62*	*artEx62 [tph-1p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1; sqIs13 [lgg-1p::GFP::lgg-1 + odr-1p::RFP]*
genetic reagent (*C. elegans*)	AVS794	This paper	*sqIs13; artEx65*	*artEx65 [unc-47p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1; sqIs13 [lgg-1p::GFP::lgg-1 + odr-1p::RFP]*
genetic reagent (*C. elegans*)	AVS709	This paper	*artEx62*	*artEx62 [tph-1p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1*
genetic reagent (*C. elegans*)	AVS710	This paper	*artEx65*	*artEx65 [unc-47p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1*
genetic reagent (*C. elegans*)	AVS888	This paper	*aetEx99*	*artEx99 [unc-47p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 2*
genetic reagent (*C. elegans*)	AVS816	This paper	*sqIs13; artEx41*	*artEx41(rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry); sqIs13 [lgg-1p::GFP::lgg-1 + unc-122p::RFP]; unc-30(ok613)*
genetic reagent (*C. elegans*)	AVS872	This paper	*sqIs13; artEx41*	*artEx41(rab-3p::hpk-1::CFP + pCFJ90 (myo-2p::mCherry); sqIs13 [lgg-1p::GFP::lgg-1 + unc-122p::RFP]; tph-1(mg280)*
genetic reagent (*C. elegans*)	AVS836	This paper	*artEx91*	*artEx91 [hpk-1p::GFP + tph-1p::mCherry] line 1*
genetic reagent (*C. elegans*)	AVS757	This paper	*artEx87*	*artEx87 [hpk-1p::mCherry + unc-47p::GFP] line 1*
genetic reagent (*C. elegans*)	AVS001	CB1370		*daf-2(e1370)*
genetic reagent (*C. elegans*)	AVS022	GR1309		*daf-2(e1370); daf-16(mgDf47)*
genetic reagent (*C. elegans*)	AVS495	CL6264	*uIs60*	*uIs60 [unc-119p::YFP + unc119p::sid-1]; eri-1(mg366)*
genetic reagent (*C. elegans*)	AVS817	This paper	*uIs60; artIs1*	*artIs1 [sur-5p::HPK-1::CFP + pCFJ90 (myo-2p::m-cherry)]; uIs60 [unc-119p::YFP + unc119p::sid-1]; eri-1(mg366)*
genetic reagent (*C. elegans*)	AVS488	This paper		*mxl-2(tm1516)*
genetic reagent (*C. elegans*)	AVS1000	This paper	*sqIs13; artEx65*	*artEx65 [unc-47p::hpk-1 + pCFJ90 (myo-2p::mCherry)] line 1; sqIs13 [lgg-1p::GFP::lgg-1 + odr-1p::RFP]; hlh-30(tm1978)*
Recombinant DNA reagent	pAVS1	This paper	plasmid	*rab-3p, hpk-1* cDNA, pPD95.75 backbone
Recombinant DNA reagent	pAVS2	This paper	plasmid	*dyp-7p, hpk-1* cDNA, pPD95.75 backbone

Recombinant DNA reagent	pAVS3	This paper	plasmid	*myo-3p, hpk-1* cDNA, pPD95.75 backbone
Recombinant DNA reagent	pAVS4	This paper	plasmid	*cat-2p, hpk-1* cDNA, pPD95.75 backbone
Recombinant DNA reagent	pAVS5	This paper	plasmid	*eat-4p, hpk-1* cDNA, pPD95.75 backbone
Recombinant DNA reagent	pAVS6	This paper	plasmid	*tph-1p, hpk-1* cDNA, pPD95.75 backbone
Recombinant DNA reagent	pAVS7	This paper	plasmid	*unc-47p, hpk-1* cDNA, pPD95.75 backbone
Recombinant DNA reagent	pAVS8	This paper	plasmid	*rab-3p, hpk-1(K176A)*, derived from pAVS1
Recombinant DNA reagent	pAVS9	This paper	plasmid	*rab-3p, hpk-1(D272N)*, derived from pAVS1
Recombinant DNA reagent	pPD95.75	Addgene	Plasmid #1494	
Sequence-based reagent	*atg-18F*	This paper	PCR primer	5'-ACTTGAGAAAACGGAAGGTGTT
Sequence-based reagent	*atg-18R*	This paper	PCR primer	5'-TGATAGCATCGAACCATCCA
Sequence-based reagent	*cdc-42F*	This paper	PCR primer	5'-AGCCATTCTGGCCGCTCTCG
Sequence-based reagent	*cdc-42R*	This paper	PCR primer	5'-GCAACCGCTTCTCGTTTGGC
Sequence-based reagent	*bec-1F*	This paper	PCR primer	5'-TTTTGTTGAAAGAGCTCAAGGA
Sequence-based reagent	*bec-1R*	This paper	PCR primer	5'-CAACCAGTGAATCAGCATGAA
Sequence-based reagent	*hpk-1F*	This paper	PCR primer	5'-AGTATGCACAGCTCCATCAC
Sequence-based reagent	*hpk-1R*	This paper	PCR primer	5'-CCATTATTGGGACCGGAACA
Sequence-based reagent	*xbp-1F*	This paper	PCR primer	5'-TGCCTTTGAATCAGCAGTGG
Sequence-based reagent	*xbp-1R*	This paper	PCR primer	5'-ACCGTCTGCTCCTTCCTCAATG
Sequence-based reagent	*gei-3F*	This paper	PCR primer	5'-AAGTCCGAGTCGCTGAACAC
Sequence-based reagent	*gei-3R*	This paper	PCR primer	5'- ATGCCTGAATGCTGACGCTC
Chemical compound, drug	isopropylthiogalactoside (IPTG)	GoldBio	I2481c-100	
Chemical compound, drug	TRIzol reagent	Life Technologies	Catalog: 15596026	
Chemical compound, drug	FUdR	Fisher/Alfa Aesar	CAS 50-91-9	
Chemical compound, drug	Aldicarb	Fluka analytical	# 33386	
Chemical compound, drug	serotonin (5-HT)	Sigma	H9623-100mg	
Commercial assay kit	QuikChange II XL Site-Directed Mutagenesis Kit	Agilent	Catalog #200521	
Commercial assay kit	RNeasy Plus Mini Kit	Qiagen	Cat. No.:74034	
Commercial assay kit	cDNA synthesis kit	BioRad	#1708890	
Commercial assay kit	PerfeCTa SYBR green FastMix	Quantabio	# 101414-276	
Commercial assay kit	TruSeq Stranded mRNA	Illumina		
Software, algorithm	Prism	GraphPad	version 7	

Software, algorithm	AxioVision		v4.8.2.0	
Software, algorithm	FastQC	Andrews, 2010		
Software, algorithm	Trimmomatic	Bolger et al., 2014		
Software, algorithm	STAR 2.4.2a	Dobin et al., 2013		
Software, algorithm	featureCounts	Liao et al., 2014	version 1.4.6-p5	
Software, algorithm	RSEM	Li and Dewey, 2011		
Software, algorithm	R statistical software environment	Team, 2013	version 4.0.2	
Software, algorithm	DESeq2	Love et al., 2014		
Software, algorithm	GOSeq	Young et al., 2010		
Software, algorithm	MuDataSeurat	Hao et al., 2021		

C. elegans strains and details

759 All strains were maintained at 20°C on standard NGM plates with OP50. For all

760 experiments, animals were grown in 20x concentrated HT115 bacteria seeded on 6cm

761 RNAi plates. Details on the strains, mutant alleles, and transgenic animals used in this

762 study are listed in the Key Resources Table.

Generation of transgenic strains

To assemble tissue-specific constructs for increased *hpk-1* expression, the *hpk-1* cDNA

was amplified and cloned under control of the following promoters: hypodermal *dpy-7p*

(Gilleard et al., 1997), body wall muscle *myo-3p* (Okkema et al., 1993), pan-neuronal

rab-3p (Nonet et al., 1997), dopaminergic neurons *cat-2p* (Lints and Emmons, 1999),

glutamatergic neurons *eat-4p* (Lee et al., 1999), serotonergic neurons *tph-1p* (Sze et al.,

2000), and g-aminobutyric acid (GABA)ergic neurons *unc-47p* (Gendrel et al., 2016). In

brief, promoter sequences were subcloned from existing plasmids into the *Prab-3p::hpk-*

1 plasmid. To create kinase domain point mutations *K176A* and *D272N*, the *Prab-*

3p::hpk-1 plasmid was used and mutations were performed with a QuikChange II XL

Site-Directed Mutagenesis Kit (Agilent).

All the assembled plasmids were validated by sequencing prior to microinjection. These

constructs were injected at 5 ng/ml together with *myo-2p::mCherry* at 5 ng/ml as co-

injection marker and *pBlueScript* at 90ng/ml as DNA carrier.

RNAi feeding

The *hpk-1* RNAi clone was originated from early copies of *E. coli* glycerol stocks of the

comprehensive RNAi libraries generated in the Ahringer and Vidal laboratories. The

control empty vector (L4440) and *hpk-1* RNAi colonies were grown overnight in Luria

broth with 50 µg ml–1 ampicillin and then seeded onto 6cm RNAi agar plates containing

5 mM isopropylthiogalactoside (IPTG) to induce dsRNA expression overnight at room

temperature. RNAi clones used in this study were verified by DNA sequencing and

subsequent BLAST analysis to confirm their identity.

RT-qPCR analysis

For measurement of gene induction, animals were synchronized and grown at 20°C,

then isolated at the specified age. For time course of *hpk-1* mRNA upregulation,

animals were harvested at specified time points (D2, D4, D6, D8 and D10). For *hpk-1*

mRNA regulation at D11 of adulthood, animals were synchronized, grown at 20°C and

harvested at D1 and D11 of adulthood. For autophagy genes upregulation, animals

were synchronized, grown at 20°C and harvested at D1 of adulthood. After harvesting

animals, RNA extraction was followed by using TRIzol reagent (Life Technologies)

followed by RNeasy Plus Mini Kit (Qiagen). RNA concentration was measured using a

Nanodrop, and RNA preparations were reverse transcribed into cDNA using the BioRad

cDNA synthesis kit (#1708890) as per manufacturer's protocol. Quantitative real time

PCR was performed using PerfeCTa SYBR green FastMix (Quantabio) with three

technical replicates for each condition. Primer sets with at least one primer spanning the

exon were used to amplify the gene of interest. *cdc-42* mRNA levels were used for

normalization. Fold change in mRNA levels was determined using ΔΔ Ct method (Livak

and Schmittgen, 2001). Primer sequences can be found in the Key Resources Table.

Lifespan analysis

Traditional lifespan assays were performed essentially as described in: (Das et al.,

2017; Lee et al., 2003). Briefly, animals were synchronized by egg prep bleaching

followed by hatching in M9 solution at 20°C overnight. The L1 animals were seeded

onto 6cm plates with HT115 bacteria and allowed to develop at 20°C. At L4 stage,

FUdR was added to a final concentration of 400 μM. Viability was scored every day or

every other day as indicated in each figure. Prism 7 was used to for statistical analyses

and the *P* values were calculated using log-rank (Mantel-Cox) method.

For lifespan with RNAi-mediated knockdown of TOR (*Figure 8--figure*

supplement 2), animals for the *let-363* (TOR RNAi) condition were first raised on empty

vector RNAi during development, and transferred to plates with *let-363* RNAi bacteria at

L4 to avoid developmental arrest at L3. In addition to having IPTG present in the media

of the RNAi plates, an additional 200uL of 0.2 M IPTG was added directly to the

bacterial lawn of all plates and allowed to dry immediately before adding animals. The

assays were otherwise performed as just described.

For lifespan assays in *Figure 1F*, assay plates and animal populations were

prepared as just described, and then lifespan was determined from analysis of time-

series image data collected for a given animal at approximately hour intervals

throughout adult life on modified Epson V800 flatbed scanners with our instance of the

C. elegans "lifespan machine" (Oswal et al., 2021; Stroustrup et al., 2013).

In all cases: scoring of viability was blinded with respect to genotype and values

obtained at previous timepoints, and animals that died due to rupturing, desiccation on

the side of the plate or well, or clear body morphology defects were censored.

Thermotolerance assay.

Survival assays at high temperature were conducted as previously described in

(Johnson et al., 2014). In brief, synchronized L1 animals were allowed to develop at

20°C and FUdR was added at the L4 stage. At day 1 adulthood, animals were moved to

35°C for a period of 6 and 8 hours. Animals were allowed to recover for 2 hours at 20°C,

and viability was scored. In all cases, scoring of viability was blinded with respect to

genotype and values obtained at previous timepoints; animals that died due to

rupturing, desiccation on the side of the plate or well, or had clear body morphology

defects were censored. Statistical testing between pairs of conditions for differences in

the number of foci was performed using student's *t* test analysis.

Induction of *hsp-16.2p*::*GFP* after heat shock.

The *hsp-16.2p*::*GFP* animals were heat shocked on day 1 of adulthood for 1 hour at

35°C and imaged after 4 hours of recovery. Images were acquired using a Zeiss Axio

Imager M2m microscope with AxioVision v4.8.2.0 software. The GFP a.u. values from

the head and intestine of the animals were acquired using the area selection feature

from AxioVision software. Two independent trials with a minimum of 20 animals per

experiment were performed.

Measurement of autophagosome formation using the *GFP::LGG-1/Atg8* reporter.

GFP::LGG-1/Atg8 foci formation was visualized as described in (Kumsta et al., 2017).

Briefly, L4 stage and day 1 adult animals were raised on HT115 bacteria at 20°C and

imaged using a Zeiss Axio Imager M2m microscope with AxioVision v4.8.2.0 software at

63X magnification. Two independent trials where at least 20 seam cells from 15-20

different animals were scored for *GFP::LGG-1* puncta accumulation. In all cases scoring

of puncta was blinded with respect to genotype.

Measurement of neurodegeneration in motor neurons.

Animals carrying the *unc-25p::GFP* reporter strain to visualize VD and DD motor

neurons were synchronized and allowed to developed at 20°C and FUdR was added at

L4 stage. At L4, D2 and D9 of adulthood, animals were imaged in a Zeiss Axio Imager

M2m microscope and the number of animals showing axonal breaks in the VD and DD

motor neurons were scored. Axonal breaks were defined as an area of discontinued

fluorescence in either the ventral or dorsal nerve cords.

Polyglutamine aggregation in neurons and locomotion analyses.

The visualization and quantification of the progressive decline in proteostasis in the

nervous system (specifically the nerve ring) was performed as described in (Brignull et

al., 2006; Lazaro-Pena et al., 2021). In brief, synchronized *rgef-1p*::*polyQ40*::*YFP* L4

animals were treated with FUdR. Then, z-stack images from the nerve ring were

acquired using a Zeiss Axio Imager M2m microscope with AxioVision v4.8.2.0 software

at 63X magnification. The fluorescent foci form compressed *z*-stack images from 20

animals per technical replicate were scored blind every other day from four to eight days

of adulthood.

To assess locomotion capacity, at day two of adulthood, 40 animals were

transferred to a drop of 10ul of M9 solution and the number of body bends performed in

a period of 30 seconds from each animal was scored (as described in (Lazaro-Pena et

al., 2021)). Statistical testing between pairs of conditions for differences in the number

of foci was performed using student's *t* test analysis.

Polyglutamine aggregation in muscle and paralysis analyses.

The visualization and quantification of the progressive decline in proteostasis in muscle

tissue was performed as described in (Lazaro-Pena et al., 2021; Morley et al., 2002).

Briefly, synchronized *unc-54p*::*Q35*::*YFP* L4 animals were treated with FUdR. Then,

fluorescent foci from 20 animals per technical replicate were scored blind daily from

days one to three of adulthood. To assess paralysis, at days 5, 7, 9, and 11 of

adulthood, prodded animals that responded with head movement (and were therefore

still alive) but no backward or repulsive motion were scored as paralyzed (as described

in (Lazaro-Pena et al., 2021)). Statistical testing between pairs of conditions for

differences in the number of foci was performed using student's *t* test analysis. Two to

five independent transgenics lines were tested.

Aldicarb assay.

The aldicarb assay was performed as described in (Mahoney et al., 2006; Oh and Kim,

2017). Plates containing Aldicarb 1mM were prepared the day before the assay and

stored at 4°C. Animals were synchronized by picking at L4 stage and placing them in a

plate with HT115 bacteria. After 24 hours, one aldicarb plate per strain were brought out

of 4°C and a small drop of OP50 was placed in the center of the plate a let dry for 30

minutes. Approximately 25 animals were placed on each plate, and the number of

paralyzed animals were scored every hour up to 5 hours. The assay was performed at

room temperature.

Exogenous 5-HT immobilization assay.

Animals were synchronized by picking at L4 stage and placing them in a plate with

HT115 bacteria. After 24 hours, serotonin (5-HT) was dissolved in M9 buffer to a 20mM

concentration (Hart, 2006). Twenty worms were placed on a glass plate well with 400ul

of 20mM serotonin for 10 minutes. The same number of control animals were placed in

400ul of M9 buffer. The locomotion of worms (mobilize or immobilized) was annotated

at the following time points: 2, 4, 6, 8, and 10 minutes. The assay was performed at

room temperature.

Sample preparation and sequencing for RNA-Seq.

10 cm RNAi plates were seeded with 1 mL of 10x concentrated HT115 EV bacteria from

an overnight culture, and allowed to dry for 1-2 days. N2 or *hpk-1(pk1393)*

Approximately 1,000 synchronized L1 animals were added to each plate (~3000

animals per condition across multiple plates). At the L4 stage, 600 μL of 4 mg/mL FUdR

was added. Animals were kept at 20 °C for the duration of the experiment.

Day 2 adult animals were collected in M9 buffer, washed 2x in M9, and a final

wash with DEPC-treated RNase-free water. Approximately two times the volume of the

animal pellet of TRIzol® reagent was added to each animal preparation, followed by

brief mixing and freezing overnight at -80 °C. Tubes were then allowed to partially thaw,

and were vortexed for five minutes to assist with disrupting the cuticle. Samples were

transferred to new tubes, and 200 μL of chloroform per 1 mL of TRIzol was added,

followed by 20s vortexing; the tubes settled at RT for 10 minutes. Supernatants were

transferred to new tubes, and an equal amount of 100% ethanol was added and mixed

before proceeding to column purification with the Qiagen RNAeasy Mini kit according

the manufacturer's instructions. Samples were eluted with 30 – 50 µL of DEPC-treated

water, and checked for initial concentration and quality with a Nanodrop ND-1000

spectrophotometer. Biological replicate samples were prepared from independently

synchronized populations of animals.

Isolated RNA was provided to the University of Rochester Genomics Research

Center for library preparation and sequencing. Prior to library preparation, RNA integrity

of all samples was confirmed on an Agilent 2100 Bioanalyzer. Libraries for sequencing

were prepared with the Illumina TruSeq Stranded mRNA kit, according to the

manufacturer's instructions. Quality of the resulting libraries was checked again with a

Bioanalyzer prior to sequencing to ensure integrity. Sequencing was performed on an

Illumina HiSeq2500 v4, yielding an average of approximately 31 million single-ended

100 bp reads per sample. Quality of the output was summarized with FastQC (Andrews,

2010) and reads were trimmed and filtered with Trimmomatic to remove adapter

sequence and any low-quality content occasionally observed toward the ends of reads

(Bolger et al., 2014). After filtering out low-quality reads, an average of 30 million reads

per sample remained, and were used for the rest of the analysis.

Analysis of *hpk-1* null and wild-type control RNA-Seq dataset.

RNA-Seq reads were aligned to the *C. elegans* genome (assembly WBcel235) with

STAR 2.4.2a (Dobin et al., 2013) using gene annotation from Ensembl (version 82)

(Cunningham et al., 2022). An average of 90.2% of reads per sample were uniquely and

unambiguously mapped to the genome. These alignments were used as input to

featureCounts (version 1.4.6-p5) for gene-level counts (Liao et al., 2014). Ambiguous or

multi-mapping reads, comprising an average of approximately 9% of the reads per

sample, were not included in the gene-level count results. Transcript-level quantification

was also performed with RSEM (Li and Dewey, 2011) to obtain TPM (Transcripts Per

Million) expression estimates.

Further analysis was performed primarily in the R statistical software

environment (version 4.0.2) (Team, 2013), utilizing custom scripts and incorporating

packages from Bioconductor. Both count and TPM expression matrices were filtered to

remove genes with low or no expression, keeping genes with at least 10 read counts in

at least one sample. Genes were further filtered to keep only those annotated as protein

coding, pseudogene, ncRNA, lincRNA, antisense, or snRNA, which could have been

included in the sequenced libraries after poly-A selection; 96.3% of genes included in

the final set are protein-coding. Gene identifiers have been updated as necessary from

newer versions of WormBase (WS284) for integration with recent resources and

datasets.

Differential expression analysis was performed with DESeq2, and the optional

fold-change shrinkage procedure was applied to moderate fold-changes for low-

expression genes (Love et al., 2014). Genes significantly differentially expressed

between *hpk-1(pk1393)* and N2 animals were taken as those with FDR-adjusted p-

value less than 0.05 and absolute value of $\log_2$ foldchange of at least 1 (on a linear

scale, down- or up-regulated by at least two-fold). Functional and over-representation

analysis for differentially expressed genes was performed with GOSeq (Young et al.,

2010)using gene sets and pathways from the Gene Ontology (Harris et al., 2004),

KEGG (Kanehisa and Goto, 2000), and Reactome (Jassal et al., 2020).

Integration with public tissue-specific and single-cell RNA-Seq datasets.

C. elegans Aging Atlas and CenGen single-cell datasets

We obtained differential expression results for old (days 8, 11, 15) vs young (days 1, 3,

5) animals across 200 cell clusters from the Aging Atlas single-cell RNA-Seq project

(Roux et al., 2022). Significantly differentially expressed genes within each cluster were

taken as those with adjusted p-value < 0.05, and $\log_2$ fold-change magnitude of at least

0.5 (approximately 1.4 fold up- or down-regulated on a linear scale). Additional neuron

identity information was obtained by mapping Aging Atlas neuron clusters to clusters

identified in late-larval stage animals in the CeNGEN single-cell neuronal transcriptome

project (Taylor et al., 2021), assisted by correlations between clusters of the two

datasets already provided with the Aging Atlas. Chemical signaling subtypes of neurons

were determined by expression of marker genes commonly used in the field for

fluorescent labeling of neurons: *tph-1* for serotonergic, *unc-47* for GABAergic, *eat-4* for

glutamatergic, *cat-2* for dopaminergic, *cat-2* for cholinergic. Where a high correlation

had been established between a cluster in the Aging Atlas and CeNGEN, a signaling

type was assigned if the cluster expressed the associated marker gene at TPM greater

than 75 based on the provided pre-normalized expression matrix available from the

CeNGEN website. Ten Aging Atlas neuron clusters did not have an obvious

corresponding cluster in the CeNGEN dataset; for these clusters neuron signaling class

was based on marker genes having average cluster counts > 0.5 across all cells in a

cluster and, where possible, a consensus of the six Aging Atlas timepoints.

Normalized single-cell aging time-course gene expression was obtained by

processing raw counts in the H5AD file from the Aging Atlas resource website

(http://c.elegans.aging.atlas.research.calicolabs.com/data). Briefly, the H5AD data from

SCANPy was read in and converted into a Seurat object with MuDataSeurat

(https://github.com/PMBio/MuDataSeurat) (Hao et al., 2021). SCTransform, as

implemented in Seurat, was used to normalize raw counts for library size differences,

yielding log-scale normalized expression values for each cell that were used for further

analysis.

Genes uniquely expressed in neurons, or significantly enriched compared to

other tissues were obtained from tissue-specific bulk RNA-Seq reported by (Kaletsky et

al., 2018) due to the additional dynamic range provided over current single-cell

sequencing technology.

C. elegans kinome, transcription factors, neuronal genome, and longevity-associated

genes

A collection of *C. elegans* kinases was assembled based on Zaru *et al.* 2017 (Zaru et

al., 2017) and cross-referenced against current gene annotation in WormBase version

WS284 (Harris et al., 2020) to keep only protein-coding genes and remove

pseudogenes or "dead" genes. This produced a list of 438 kinases in the *C. elegans*

kinome (*Figure 2--figure supplement 4, Supplementary File 5*). *C. elegans* transcription

factors were identified from a previously published compendium (Nye et al., 1989).

Genes with known or predicted functions important to neurons, and their associated

roles, were assembled from tables curated in (Hobert, 2013). Genes with lifespan

phenotypes primarily from perturbation studies in *C. elegans* utilizing RNAi or mutant

strains were assembled from annotation in WormBase (WS282) as well as the GenAge

database (de Magalhaes and Toussaint, 2004).

Statistical analysis for *in vivo* assays.

Unless otherwise specified, GraphPad PRISM version 7 software was used for

statistical analyses. Data were considered statistically significant when P value was

lower than 0.05. In figures, asterisks denote statistical significance as (*, $P < 0.05$, **, P

< 0.001, ****, $P < 0.0001$) as compared to appropriate controls. N number listed in figure

legend indicates the number of one representative experiment among all biological

trials.

Appendices.

Appendix 1. While single-cell RNA-Seq is a powerful approach already, well-

documented challenges remain with respect to consistency in cell captures and

differentiating biological noise from technical artifacts such as "drop-outs" from low

numbers of average reads per cell (Hwang et al., 2018, 2021; Ke et al., 2022; Nayak

and Hasija, 2021)(refs). We attempted to identify gene-gene correlations within cell

clusters across individual ages based on the *C. elegans* Aging Atlas dataset and found

that inconsistency in the number of cells per cluster across timepoints limits our ability to

identify many robust and significant associations- such variability is likely technical in

nature, as it is not globally consistent across cell clusters for a given timepoint. In most

cases, when aggregated into "young" and "old" sample groups, as was done for

differential expression analysis in the original Aging Atlas analysis (Roux et al., 2022)

the groups represent comparable numbers of total cells for most clusters. The trends we

have observed at the level of individual time-points indicate that the age-associated up-

regulation of *hpk-1* does not always occur at the same time point across cell clusters

and is not necessarily tightly linked to the upregulation of longevity-associated

transcription factors in the same cluster (*Appendix 1 Figure 1*). Thus, the age-

associated upregulation of pro-longevity factors in many cell clusters may not be due to

a highly coordinated regulatory process, but rather a distributed response to

accumulating local stress or damage. More work will be necessary to resolve if such

late-life expression changes indeed contribute to maintenance of wild-type lifespan and

healthspan, and the mechanism of the changes thereof (e.g., increased transcription,

down-regulation of miRNA targeting, epigenetic changes) and the factors that mediate

these changes.

1034
Appendix 1--Figure 1. *hpk-1* and longevity-associated transcription factors are
up-regulated in overlapping cell-clusters, but timing of the age-associated
changes varies between cell clusters.
Per-age expression summary for cell clusters with four or more differentially expressed
longevity-associated transcription factors, of the 48 cell clusters in which *hpk-1* is
upregulated in old vs young animals. Each sub-plot shows log-scale normalized scRNA-
Seq expression (mean and SEM) for the indicated cluster. Only genes which were
significantly differentially expressed in the cluster are shown (old vs young animals, up-
or down-regulated). Primary data for this analysis was generated in the *C. elegans*
Aging Atlas (Roux et al., 2022).

1045

Appendix 2. Loss of *hpk-1* upregulates expression of a number of neuronal genes

including those involved in: neuropeptide signaling, neurotransmission, release of

synaptic vesicles, calcium homeostasis, and function of sensory neurons. In addition to

upregulation of *ins* neuropeptides, 12 FMRFamide-related (flp) peptides, 19

neuropeptide-like proteins (*nlp*), nine neuropeptide maturation genes (e.g., *egl-3*

(proprotein convertase subtilisin/kexin type 2) and *egl-21*(carboxypeptidase E)), and

nine putative GPCR neuropeptide receptors were also significantly upregulated in *hpk-1*

null mutants. Neurotransmitters are signaling molecules that trigger a response by

binding to ionotropic or metabotropic neurotransmitter receptors in recipient neurons or

tissues; we find that mainly the latter are upregulated in *hpk-1* null mutant animals,

including: acetylcholine (*gar-1, gar-2* and *gar-3*), GABA (*gbb-2*) and glutamate (*mgl-3*

and uncharacterized *C30A5.10*). In total nine metabotropic and two ionotropic glutamate

receptors, 14 vesicular glutamate transport family genes, nine acetylcholine receptor

ligand-gated ion channels, and nine GABA receptor ligand-gated ion channels were

upregulated. Additionally, *ace-2*, *ace-3* and *cha-1*, involved in acetylcholine degradation

and synthesis (respectively), were upregulated. *comt-4*, a catechol-O-methyltransferase

expected to be involved in the degradation of dopamine, was significantly down-

regulated.

1064   Synaptic transmission, the formation and release of synaptic vesicles, carry

neurotransmitters and rely on proteins necessary for docking, priming and fusing to the

axon terminal plasma membrane, which releases cargo into the synaptic cleft.

Interestingly, loss of *hpk-1* causes the upregulation of synaptic proteins such as *unc-13*,

unc-10 and *rab-3*, which are essential for vesicle priming, recruitment and sequestration

of vesicles, respectively. The process of vesicle release is highly dependent on a Ca^{2+}

signal detected by synaptotagmins proteins, which promotes vesicle fusion to the

plasma membrane (Deak, 2014). In *hpk-1* null mutant animals, we observed

upregulation of synaptotagmin *snt-1*, *snt-3* and *snt-4* Ca^{2+} sensor proteins for vesicle

release.

1074   Neuronal function is regulated by changes in calcium levels: a decrease in

calcium uptake reduces neuronal function by a reducing neurotransmitter release

(Scriabine and Kazda, 1989). In contrast, increasing calcium influx triggers activation of

proteases and phospholipases, resulting in cell damage and cell death (Cheung et al.,

1986; Siesjo, 1989). Voltage-gated Ca^{2+} channels (VGCCs) are composed of 24

transmembrane alpha-1 subunits and associate with auxiliary beta-1 and alpha-2-delta

subunits. *hpk-1* null mutant animals have increased expression of two of five alpha-1

subunits (*egl-19* and *unc-2*), and both alpha-2-delta subunits (*unc-36* and *tag-180*).

Previous research describes the role of *egl-19* voltage-gated Ca^{2+} channel in the

induction of the heat stress response (Silva et al., 2013): moderate upregulation of

acetylcholine signaling improves muscle proteostasis by the activation of HSF-1,

triggered by the *egl-19*-dependent Ca^{2+} influx.

Neuronally expressed cilia are essential for cell motility, sensory perception and

signaling. The assembly and maintenance of sensory cilia depends on IFT: bidirectional

movement of cargo to and from the base and tip of the axoneme. The *C. elegans* IFT

machinery is composed of: anterograde IFT motors, heterotrimeric kinesin-II and

homodimeric OSM-3, retrograde dynein motors; and the IFT-A, -B and BBSome particle

complexes (Hao et al., 2009; Scholey et al., 2004). Loss of *hpk-1* induces the

expression of a high number of IFT machinery components including: *klp-11* (kinesin II

motor), *che-3* (dynein motor), three and seven members of the IFT-A and -B complexes

(respectively), and five members of BBSome complex. In mammals, there is a link

between primary cilia signaling and age-related brain disease, including

neurodegenerative disorders (Ma et al., 2022).

Data availability.

All data supporting the findings of this study are available within the article, its Supplementary Information files, or the peer-review file. RNA-Seq data for *hpk-1(pk1393)* and N2 is available from GEO, accession GSE220744. Data analysis scripts have been deposited at: https://github.com/samuelsonlab-urmc/hpk1_manuscript_2023.

Acknowledgments.

We would like to thank members of the Samuelson laboratory, past and present, for their thoughtful insight and assistance related to this project, especially Drs. Rachel Kitt and Sara Farrell. We would like to thank members of the Western New York Worm Group for their input and discussions. We would like to thank Dr. Doug Portman (URMC) and his laboratory for strains and advice. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). We would like to personally thank Drs. Dirk Bohmann (URMC), Gary Ruvkun (Massachusetts General Hospital/Harvard Medical School), Arjumand Ghazi (University of Pittsburgh), Seung-Jae Lee (Korea Advanced Institute of Science and Technology) and Mary Wines-Samuelson for critical reading of this manuscript. We would like to thank Drs. Cynthia Kenyon and David Kelley (Calico Life Sciences) for providing us the full scRNA-seq results for the old vs young differential expression analysis results for each cell cluster in the *C. elegans* Aging Atlas. We would like to thank Dr. John Ashton for assistance and the University of Rochester Genomics Research Center for conducting RNA-sequencing. Research reported in this publication was supported by the National Institute On Aging of the National Institutes of Health under Award Number RF1AG062593. CADB was supported by F32HD105323. The

1121   content is solely the responsibility of the authors and does not necessarily represent the

1122   official views of the National Institutes of Health.

1123

 Figure, Figure Supplement, Video, and File Legends
1125

Figure 1. Neuronal HPK-1 extends longevity and promotes healthspan.
Lifespan curves of animals overexpressing *hpk-1* (red line) in the nervous system **(A)**,
muscle **(B)**, intestine **(C)** and hypodermis **(D)** compared with control non-transgenic
siblings (blue line). (n≥175, log-rank test, *P* < 0.0001 for A, n≥78 for B, n≥82 for C and
n≥105 for D). Lifespan graphs are representative of two biological replicates. **(E)**
Frequency of body bends of *wild type, rab-3p::hpk-1* and *hpk-1(pk1393); rab-3p::hpk-1*
day 2 adult animals (n≥48). **(F)** Lifespan curves of wild-type (blue), *hpk-1(pk1393)* (red),
and *hpk-1(pk1393); rab-3p::hpk-1* animals (pink). **(G)** Survival of day 1 adult animals
subjected to heat shock treatment. Graph represents one of the two individual trials
(n≥66). *T* test analysis with **P < 0.01, ***P < 0.001 and ***P < 0.0001. See
Supplementary Files 1, 2 and 3 for details and additional trials.
1137

Figure 2. During normal aging *hpk-1* is the most broadly upregulated kinase and
overlaps with key longevity-associated transcription factors within the nervous
system.
(A) Relative expression of *hpk-1* mRNA during normal aging, as measured by RT-
qPCR**.** *T*-test analysis with *P < 0.05, **P < 0.01, ***P < 0.001 and ****P < 0.0001. **(B)**
Venn diagram depicting transcription factors (TFs) with significant age-associated
differential expression with *hpk-1* within three or more of the 36 neuronal and/or 11 non-
neuronal cell clusters (yellow and blue, respectively), along with TFs that have
previously been implicated in *C. elegans* longevity (red). **(C, D)** Heat map of neuronal
cell clusters in which *hpk-1* expression increases during normal aging along with most
broadly co-expressed TFs (≥ 9 cell-clusters). Positive (red) and negative (blue) fold
change with aging of a given TF are shown in (C) and (D), respectively, grouped by
average across the indicated clusters. Neuronal types, subtypes, cell-cluster or
individual neuronal pair, as well as whether a TF has previously been implicated in
aging is indicated. Primary data for this analysis was generated in the *C. elegans* Aging
Atlas (Roux et al., 2022) and significant differences in expression with aging were
determined by filtering differential expression results from a comparison of old vs young
animals, based on adjusted p-values <0.05 and a $\log_2$ foldchange magnitude of at least
0.5 (between days 8,11,15 and days 1,3,5). **(E)** Relative expression of *hpk-1, xbp-1,*
and gei-3 mRNA of day 11 adult animals, compared to day 1 and as measured by RT-
qPCR. *T* test analysis with **P < 0.01 and ***P < 0.001. **(F)** Representative images of
hpk-1 expression within the nervous system of day 1 and 11 adult animals (*hpk-*
1p::mCherry; unc-47p::GFP). See *Figure 7F* and Figure 7- *figure supplement 1A-C* for
representative images of whole animals. (n≥5). **(G, H)** Functions and pathways
associated with two or more of the TFs listed in (C) and (D), respectively. **(I)** *hpk-1* is
differentially expressed with age in more cell clusters than any other kinase, and mostly
in neurons. Heatmap shows $\log_2$ foldchange in old vs young animals for kinases with
differential expression in ≥10 cell clusters. Primary data for this analysis was generated
in the *C. elegans* Aging Atlas (Roux et al., 2022) and filtered for significant age-
associated expression changes as described in (C, D)*.* See *Supplementary Files 4* for
additional RT-qPCR details, and *5* and *6* for dataset.
1169

Figure 3. Neuronal HPK-1 maintains neuronal integrity during aging.
(A) Fluorescent micrographs of control and *hpk-1(pk1393)* 9 day old animals expressing GFP in GABAergic DD and VD motor neurons. Asterisks (*) indicate axonal breaks in the motor neurons. **(B)** Percentage of wild type and *hpk-1* mutant animals with neurodegeneration in VD motor neurons (n$\geq$23). **(C)** Percentage of *hpk-1* rescue animals with neurodegeneration in VD motor neurons (n$\geq$10). **(D)** Percentage of wild type and *hpk-1* mutant animals with neurodegeneration in DD motor neurons (n$\geq$20). Percentage of immobilized animals after exposure to aldicarb (n$\geq$67) **(D)** and to 20mM 5-HT (n$\geq$38) **(E)**. *T* test analysis with **P < 0.01 and ***P < 0.001. See *Supplementary Files 7, 8 and 9* for details and additional trials.

Figure 4. Loss of *hpk-1* results in broad dysregulation of neuronal gene expression.
(A) Differential expression analysis for RNA-Seq of day 2 adult *hpk-1* null mutant animals (n=3) compared to N2 wild-type animals (n=3) identified 2,201 genes with significantly altered expression with loss of *hpk-1* (adjusted p-value < 0.05 and |log$_2$ foldchange| $\geq$ 1), 84% of which were upregulated in the mutants. The volcano plot illustrates the criteria applied for selecting genes with significant and substantial expression changes, shown as blue dots: vertical dashed lines are the foldchange threshold and the horizontal dashed line is the p-value threshold. **(B, C)** Enrichment for functions and pathways associated with the 348 downregulated genes **(B)** or the 1853 upregulated genes **(C)** in *hpk-1* animals yields innate immune response as the most significantly over-represented term for downregulated genes, but reveals a large number of significant associations to pathways and processed for the upregulated genes- many of which are associated with neurons and neuronal signaling. Each bar represents a functional term or pathway from the Gene Ontology (GO), KEGG, or Reactome databases, as indicated. Bar colors show the –log$_{10}$ transformed enrichment p-value. The number in each bar is the size of the overlap between the set of differentially expressed genes and the genes in the term or pathway. The x-axis indicates the fraction of the total genes in the term or pathway that were differentially expressed. All results shown have significant enrichment after adjustment for multiple testing. **(D)** Functional enrichment for 283 genes upregulated with *hpk-1* loss that are specific to, or enriched in, *C. elegans* neurons shows broad dysregulation of functions associated with neurons including neurotransmitter transport and release, cell-cell junctions, axons, neuropeptides, and sensory processes, among others. 880 genes uniquely expressed or significantly enriched in neurons were derived from bulk tissue-specific RNA-Seq profiling results in (Kaletsky et al., 2018) and 32% of these are significantly upregulated in *hpk-1* null animals. See *Supplementary Files 10 and 11* for dataset and additional analysis.

Figure 5. Neuronal HPK-1 prevents the decline in proteostasis through neurotransmitter release.
(A) Nomarski (*i*) and fluorescent (*ii-iv*) micrographs of animals expressing polyglutamine fluorescent reporter within the nervous system. (**B** and **C**) Quantification of foci in the nerve ring (B) and body bend frequency (C) (n$\geq$21 for B and n$\geq$36 for C). (**D** and **E**) Quantification of foci in the nerve ring (D) and frequency of body bends (E) in neuronal

enhanced RNAi animals (n≥19 for D and n≥39 for E). **(F)** Fluorescent micrographs of animals expressing the polyglutamine fluorescent reporter in muscle. (**G** and **H**) Quantification of foci in muscle (G) and paralysis rate (H). Graphs are representative of 5 individual transgenic lines (n≥18 for G and n≥23 for H). (**I** and **J**) Quantification of foci in muscle (I) and paralysis rate (J) in neuronal enhanced RNAi animals (n≥22 for I and n≥73 for J). **(K)** Neuropeptides and neurotransmitters are essential for neuronal signaling; vesicle release depends on *unc-31* and *unc-13*, respectively. (**L**) Quantification of relative *polyQ* fluorescent foci in muscle cells after *rab-3p::hpk-1* overexpression in control, *unc-31(e169)* and *unc-13(e51)* animals; see Figure 5- *figure supplement 2* for absolute values and statistical analysis. *T* test analysis with **P < 0.01 and ***P < 0.001. See *Supplementary Files 12, 2, and 13* for details and additional trials.

Figure 6. Neuronal HPK-1 induces the proteostatic network in distal tissues.
(**A**) Fluorescent micrographs of animals expressing *hsp-16.2p::GFP* in basal conditions. (**B** and **C**) Fluorescent micrographs (B) and fluorescent density quantification (C) of control and *rab-3p::hpk-1* day 1 adult animals expressing *hsp-16.2p::GFP* after heat shock (35°C) for 1 hour at (n≥15). (**D** and **E**) Fluorescent micrographs and densitometry quantification of the intestinal fluorescence of *hsp-16.2p::GFP* after heat shock (n≥24). (**F** and **G**) Fluorescent micrographs and quantification of fluorescent puncta in the nerve ring of control and *rab-3p:hpk-1* L4 animals expressing *rgef-1p::GFP::LGG-1/Atg8* (n≥21). **(H** and **I)** Fluorescent micrographs (H) and quantification (I) of puncta in seam cells of control and *rab-3p::hpk-1* animals expressing *lgg-1p::GFP::LGG-1/Atg8* (n≥43). (**J** and **K**) Fluorescent micrographs (J) and quantification (K) of puncta in proximal intestinal cells of control and *rab-3p::hpk-1* animals expressing the *lgg-1p:GFP::LGG-1/Atg8* autophagosome reporter (n≥17). **(L)** Expression of autophagy genes *atg-18* and *bec-1* via RT-qPCR in *rab-3p::HPK-1* animals compared to non-transgenic controls. *T*-test analysis with *P < 0.05, **P < 0.01, ***P < 0.001 and ****P < 0.0001. See *Supplementary Files 4, 14 and 15* for details and additional trials.

Figure 7. Expression of HPK-1 in serotonergic and GABAergic prevents the decline in proteostasis in muscle tissue.
(**A** and **B**) Quantification of foci (A) and paralysis rate (B) of animals expressing *hpk-1* in serotonergic neurons (n≥17 for A and n≥28 for B). (**C** and **D**) Quantification of foci (C) and paralysis rate (D) of animals expressing *hpk-1* in GABAergic neurons (n≥19 for C and n≥27 for D). **(E)** Representative fluorescent micrographs of *hpk-1* expression in serotonergic neurons. **(F)** Representative fluorescent micrographs of *hpk-1* expression in GABAergic neurons. *T* test analysis performed to each time point with *P < 0.05 and **P < 0.01. See *Supplemental Files 12 and 13* for details and additional trials.

Figure 8. Serotonergic and GABAergic HPK-1 signaling activates distinct adaptive responses to regulate stress resistance and longevity.
(A) Survival of serotonergic HPK-1 expressing day 1 adult animals after heat shock (n≥32). **(B)** Lifespan of serotonergic HPK-1 expressing animals (n≥76). (**C** and **D**) Representative fluorescent micrographs (C) and quantification (D) of autophagosomes in hypodermal cells of serotonergic HPK-1 day 1 adult animals (n≥80). **(E)** Survival of

GABAergic HPK-1 expressing day 1 adult animals after heat shock (n≥47). **(F)** Lifespan
of GABAergic HPK-1 expressing animals (n≥119). **(G** and **H)** Representative fluorescent
micrographs (G) and quantification of autophagosomes in hypodermal cells of
GABAergic HPK-1 day 1 adult animals (n≥64). **(I)** Quantification of autophagosomes in
hypodermal cells of mutant animals of serotonin and GABAergic signaling (*tph-1* and
unc-30, respectively), expressing pan-neuronal *hpk-1* (day 1 adults, n≥ 67). **(J)** Relative
change in number of puncta in seam cells of *unc-47p::hpk-1* animals expressing *lgg-*
1p::GFP::LGG-1/Atg8, compared to non-transgenic controls after indicated gene
inactivation (n≥28) for all conditions); see Figure 8- *figure supplement 1* for absolute
values and statistical analysis. *T* test analysis with *$*P < 0.05$, **$**P < 0.01$ and ****$****P <$
0.0001. See *Supplementary Files 1, 3, and 15* for details and additional trials.
1273
Figure 9. Differential regulation of the proteostatic network through HPK-1 activity
in serotonergic and GABAergic neurons.
(A) During normal aging *hpk-1* expression increases broadly throughout the nervous
system in response to accumulating damage and dysfunction. Overexpression of *hpk-1*
within the nervous system fortifies proteostasis by priming and preemptively activating
mechanisms that delay the progression of aging. In the absence of *hpk-1* the
proteostatic network is compromised, resulting in neuronal dysfunction and increased
expression compensatory mechanisms to maintain homeostasis (see text for details).
(B) HPK-1 activity in serotonergic and GABAergic neurons initiates distinct adaptive
responses, either of which improve proteostasis in a cell non-autonomous manner.
Serotonergic HPK-1 protects the proteome from acute heat stress, while GABAergic
HPK-1 fortifies the proteome by regulating autophagy activity in response to metabolic
stress, such as changes in TORC1 activity. The MXL-2, HLH-30 (TFEB), and DAF-16
(FOXO) transcription factors are all required for hypodermal induction of autophagy in
response to increased HPK-1 activity in GABAergic neurons. Gray indicates predicted
interactions based on our prior genetic analysis (Das et al., 2017). MML-1 and MXL-2
heterodimerize and encode the orthologs of the Myc-family of transcription factors
Mondo A/ChREBP and Mlx, respectively.
1292
1293 **Video 1. HPK-1 functions within the nervous system to regulates healthspan.**
1294 Video of wild-type, *hpk-1(pk1393)* null animals, and *hpk-1* null animals with neuronal
1295 expression of *hpk-1* (*rab-3p::HPK-1*). Videos were generated from consecutive images
1296 taken approximately every 15 minutes over the course of the animals lifespan using the
1297 Lifespan Machine method (Stroustrup et al., 2013).
1298
Figure 2--Figure Supplement 1. During normal aging *hpk-1* is differentially
expressed along with transcription factors in cell clusters corresponding to
tissues outside of the nervous system.
Heat map of non-neuronal cell clusters in which *hpk-1* expression increases during
normal aging along with TFs differentially expressed in 3 or more neuron cell-clusters.
Fold-changes of a given TF from old versus young animals are shown, upregulation
with aging in red and downregulation with aging in blue. Neuronal types,
neurotransmitter subtypes are indicated in the top, names of the cell cluster or individual
neuronal pair are given at the bottom, and whether a TF has previously been implicated

in aging is indicated by the black and gold on the left side (genes with gold bars have
previously published aging-associated phenotypes). Primary data for this analysis was
generated in the *C. elegans* Aging Atlas (Roux et al., 2022) and differential expression
with aging was based on adjusted p-values <0.05 and a $\log_2$ FC magnitude of at least
0.5 (between days 8,11,15 and days 1,3,5). Note *hpk-1* is included for reference and is
a transcriptional co-factor, but is not a *bona fide* transcription factor. See *Supplementary*
Files 5 and 6 for dataset.
1315
Figure 2--Figure Supplement 2. *Hpk-1* and all transcription factors differentially
expressed in overlapping cell clusters within the *C. elegans* nervous system
during aging.
Heat map of neuronal cell clusters in which *hpk-1* expression increases during normal
aging along with TFs differentially expressed in 3 or more neuron cell-clusters. Fold-
changes of a given TF from old versus young animals are shown, upregulation with
aging in red and downregulation with aging in blue. Neuronal types, neurotransmitter
subtypes are indicated in the top, names of the cell cluster or individual neuronal pair
are given at the bottom, and whether a TF has previously been implicated in aging is
indicated by the black and gold on the left side (genes with gold bars have previously
published aging-associated phenotypes). Primary data for this analysis was generated
in the *C. elegans* Aging Atlas (Roux et al., 2022) and differential expression with aging
was based on adjusted p-values <0.05 and a $\log_2$ FC magnitude of at least 0.5 (between
days 8,11,15 and days 1,3,5). Note *hpk-1* is included for reference and is a
transcriptional co-factor, but is not a *bona fide* transcription factor. See *Supplementary*
Files 5 and 6 for dataset.
1332
Figure 2--Figure Supplement 3. *hpk-1* is broadly co-expressed with longevity-
associated transcription factors in late larval stage animals.
Vertical bars show the number of neurons expressing TFs or *hpk-1* in the intersection
indicated by the filled dots. The number of neurons each individual gene is found
expressed is given by the horizontal bars, "set size". Primary data for this analysis was
generated in the CenGEN project, expression was determined based on their "medium"
stringency threshold of average TPM per neuron type ≥ 0.1 (Taylor et al., 2021).
1340
Figure 2--Figure Supplement 4. Only a limited subset of kinases change
expression during normal aging.
Heat map of 184 kinases that change expression during normal aging in any cell cluster
in the aging atlas dataset, of the 438 total kinases identified in *C. elegans*. Increased
(red) and decreased (blue) expression with aging of a given kinase is shown. Major
tissue types, kinase classes, as well as whether a kinase has previously been
implicated in aging is indicated in the top and side bars, respectively. Primary data for
this analysis was generated in the *C. elegans* Aging Atlas (Roux et al., 2022) and
differential expression with aging was based on adjusted p-values <0.05 and a $\log_2$ FC
magnitude of at least 0.5 (between days 8,11,15 and days 1,3,5). See *Supplementary*
Files 5 and 6 for dataset.
1352

Figure 5--Figure Supplement 1. HPK-1 kinase activity is necessary to improve proteostasis.

(A) Multiple sequence alignment of the HPK-1 kinase domain across evolutionarily distant species. Alignment was conducted using Clustal Omega (Sievers et al., 2011), with sequence IDs: HIPK2 (NP_073577.3), HPK-1 (F20B6.8a), HIPK2 (*M. musculus*, NP_001281073.1), HIPK (*D. melanogaster*, CG17090), HPK-1 (*C. briggsae*, CBP26904), and Yak1 (SGDID:S000003677). **(B)** Overlap between the predicted tertiary structures of the kinase domain of *C. elegans* HPK-1 (green), *S. cerevisiae* Yak1 (pink), and *H. sapiens* HIPK2 (blue). The K176A and D272N mutations are indicated by a red triangle and purple star, respectively, falling within the catalytic domain (He et al., 2010). Predicted structures for these genes were downloaded from AlphaFold (Jumper et al., 2021; Varadi et al., 2022): HPK-1(Q8MQ70 HIPK_CAEEL), Yak1 (A0A7C8V8P7_ORBOL), (H7BXX9_HUMAN). Highly conserved regions observed using multiple sequence alignment where then aligned in 3D using PyMol alignment tool (DeLano, 2002). **(C-F)** Quantification of foci in muscle (C, D) and paralysis rate (E, F) of transgenic animals with neuronal overexpression of *hpk-1(K176A)* (C, E) and *hpk-1(D272N)* (D, F), respectively. (n≥20 for C, n≥20 for D, n≥48 for E and n≥30 for F). *T* test analysis was performed between transgenic and non-transgenic animals at every time point. See *Supplementary Files 12 and 13* for details.

Figure 5--Figure Supplement 2. HPK-1 cell-nonautonomous regulation of proteostasis depends on neurotransmitter signaling.

(**A**) Quantification of *polyQ* foci in muscle of wild type and neuropeptide secretion mutants, *unc-31(e169)* (n≥21), **(B)** Quantification of foci in muscle of wild type and neurotransmitters secretion mutants, *unc-13(e51)* (n≥21). *T* test analysis with **$P < 0.01$ and ***$P < 0.001$. See *Supplementary File 12* for details.

Figure 7--Figure Supplement 1. *Hpk-1* expression in glutamatergic and dopaminergic neurons is not sufficient to prevent the decline in proteostasis.

(A and **B)** Quantification of number of foci (A) and paralysis rate (B) of animals expressing *hpk-1* in the glutamatergic neurons (*eat-4p::hpk-1*; green line) and control non-transgenic siblings (blue line) (n≥18 for A and n≥29 for B). **(C** and **D)** Quantification of number of foci (C) and paralysis rate (D) of animals expressing *hpk-1* in the dopaminergic neurons (*cat-2p::hpk-1;* orange line) and control non-transgenic siblings (blue line) (n≥16 for C and n≥34 for D). See *Supplementary Files 12 and 13* for details and additional trials.

Figure 7--Figure Supplement 2. *Hpk-1* is expressed in GABAergic and serotonergic neurons. Representative fluorescent micrographs of day 1 adult animals co-expressing *hpk-1p::mCherry* and *unc-47p::GFP* **(A-C)**; and *hpk-1p::GFP* and *tph-1p::mCherry* **(D-F)**.

Figure 8--Figure Supplement 1. HLH-30 (TFEB), MXL-2 (Mlx), and DAF-16 (FoxO) are required for HPK-1 mediated induction of autophagy.

(A) Quantification of puncta in seam cells of *unc-47p::hpk-1* and non-transgenic control animals expressing *lgg-1p::GFP::LGG-1/Atg8* after either RNAi treatment with empty

vector (ev), *hsf-1, mxl-2, pha-4, hlh-30,* or *daf-16* (n$\geq$28). *T* test analysis with ****$P <$ 0.0001. **(B)** Quantification of puncta in seam cells of *unc-47p::hpk-1* and non-transgenic control animals expressing *lgg-1p::GFP::LGG-1/Atg8* in otherwise wild-type or *hlh-30(tm1978)* mutant animals at L4 (left) or day 1 of adulthood (right) (n$\geq$57). *T* test analysis with ****$P < 0.0001$. See *Supplementary File 15* for details and additional trials.

Figure 8--Figure Supplement 2. ***mxl-2* is required for decreased TORC1 signaling to increase longevity.**
Lifespan of wild-type (N2, darker shades) or *mxl-2(tm1516)* null mutant animals (lighter shades) treated with either empty vector (black/gray), *let-363(RNAi)* (blues), or *daf-15(RNAi)* (reds). Lifespan graph is representative of one trial with two replicates. Curves were plotted using WormLife (https://github.com/samuelsonlab-urmc/wormlife) (Cornwell et al., 2018). See *Supplementary File 1* for details and additional trial.

Supplementary File 1. *Lazaro-Pena SF01 Lifespan assays.xlsx.* Trials, data, strains, genotypes, number of animals, and statistics for lifespan data presented in Figures 1a-d, f, 8b,f, and 8-S2.

Supplementary File 2. *Lazaro-Pena SF02 Body bends assays.xlsx.* Experimental data, strains, genotypes, number of animals and statistical analysis for data presented in Figures 1e, and 5c,e.

Supplementary File 3. *Lazaro-Pena SF03 Thermotolerance assays.xlsx.* Trials, data, analysis, strains, genotypes, number of animals, and statistics for thermotolerance data presented in Figures 1g, and 8a,e.

Supplementary File 4. *Lazaro-Pena SF04 RT-qPCR assays.xlsx.* Trials, data, analysis, strains, genotypes, RNAi conditions, and statistics for RT-qPCR expression analysis presented in Figures 2a,e, and 6l.

Supplementary File 5. *Lazaro-Pena SF05 TFs DE with aging in hpk-1 DE clusters.xlsx.* Transcription factors differentially expressed in old vs young animals in the same cell clusters where *hpk-1* has significant expression changes with age in the Aging Atlas dataset. This includes a summary of which clusters exhibit upregulation and which exhibit downregulation for a TF, and if the gene has been previously shown to have aging or lifespan associated phenotypes in RNAi or loss-of-function mutant experiments. Primary data was from the *C. elegans* Aging Atlas, and association to aging phenotypes was determined based on curated annotations from WormBase and GenAge. File is associated with Figures 2, 2-S1, 2-S2, and 2-S4.

Supplementary File 6. *Lazaro-Pena SF06 C elegans kinome aging DE.xlsx.* Kinases differentially expressed in old vs young animals out of any of the 200 clusters with DE analysis results from the Aging Atlas dataset. Out of 438 kinases identified in *C. elegans*, 186 have significant age-associated expression changes in at least one Aging Atlas cell cluster. For these kinases, we summarized the number of age-associated up- and down-regulated changes, if they have been previously identified as having aging or

lifespan phenotypes in loss-of-function experiments, and the fold-changes in *hpk-1* null vs WT animals for significant changes in our bulk RNA-Seq experiment. Primary data was from the *C. elegans* Aging Atlas, and association to aging phenotypes was determined based on curated annotations from WormBase and GenAge. File is associated with Figures 2, 2-S1, 2-S2, and 2-S4.

Supplementary File 7. *Lazaro-Pena SF07 Neurodegeneration assay.xlsx.* Experimental data, strains, genotypes, number of animals/cells, age and statistical analysis for data presented in Figures 3b-d.

Supplementary File 8. *Lazaro-Pena SF08 Aldicarb assay.xlsx.* Experimental data, strains, genotypes, number of animals, hours on aldicarb and statistical analysis for data presented in Figure 3e.

Supplementary File 9. *Lazaro-Pena SF09 Serotonin assay.xlsx.* Experimental data, strains, genotypes, number of animals, minutes on 5-HT and statistical analysis for data presented in Figure 3f.

Supplementary File 10. *Lazaro-Pena SF10 hpk-1(0) and N2 D2 RNA-seq.xlsx.* Differential expression analysis results for RNA-Seq of *hpk-1(pk1393)* null mutant animals compared to wildtype, presented in this manuscript. *Sheet 1:* differential expression for all 16,828 genes considered to be expressed in any sample in the experiment. *Sheet 2:* the subset of genes with significant expression changes between groups (adjusted p-value < 0.05, log2 fold-change magnitude ≥ 1). Additional columns in both sheets include a summary of the raw and normalized expression for each group, and the human homologs from OrthoList, where applicable. File is associated with Figure 4.

Supplementary File 11. *Lazaro-Pena SF11 hpk-1 DE genes with neuron-associated function annotations.xlsx.* Of the genes significantly differentially expressed in *hpk-1(pk1393)* null mutant animals vs wildtype, we found 280 overlap a set of genes with important functions in neurons (Hobert, O. (2018). *The neuronal genome of Caenorhabditis elegans*). For these genes, table includes a summary of their function, the differential expression fold-change and raw expression level in the *hpk-1(pk1393)* vs wildtype RNA-Seq comparison, and the clusters from the Aging Atlas dataset with significant age-associated expression changes (where applicable). File is associated with Figure 4.

Supplementary File 12. *Lazaro-Pena SF12 Polyglutamine assays.xlsx.* Trials, data, analysis, strains, genotypes, number of animals, age and statistical analysis of polyglutamine foci formation data presented in Figures 5b,d,g,i,l, 7a,c, 5-S1c,d, -S6a,b, and 7 -S1a,c.

Supplementary File 13. *Lazaro-Pena SF13 Paralysis assays.xlsx.* Trials, data, analysis, strains, genotypes, number of animals, age and statistical analysis of data for

1490 paralysis onset in animals expressing *polyQ::YFP* presented in Figures 5h,j, 7b,d, 5-
1491 S1e,f, and 7-S1b,d.
1492
1493 **Supplementary File 14.** *Lazaro-Pena SF14 Induction of molecular chaperone.xlsx.*
1494 Trials, fluorescent quantification, analysis, strains, genotypes, number of animals, heat
1495 shock conditions, and statistical analysis of data presented in Figure 6c,e.
1496
1497 **Supplementary File 15.** *Lazaro-Pena SF15 Induction of autophagosomes.xlsx.*
1498 Trials, autophagosome quantification, analysis, strains, genotypes, number of cells, age
1499 and statistical analysis of data presented in Figures 6g,i,k, 8d,h,I,j, and 8-S1.
1500

References

Alcedo, J., and Kenyon, C. (2004). Regulation of C. elegans longevity by specific gustatory and olfactory neurons. Neuron *41*, 45-55.

An, J.H., and Blackwell, T.K. (2003). SKN-1 links C. elegans mesendodermal specification to a conserved oxidative stress response. Genes Dev *17*, 1882-1893.

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data. [Online].

Apfeld, J., and Kenyon, C. (1999). Regulation of lifespan by sensory perception in Caenorhabditis elegans. Nature *402*, 804-809.

Armato, U., Chakravarthy, B., Pacchiana, R., and Whitfield, J.F. (2013). Alzheimer's disease: an update of the roles of receptors, astrocytes and primary cilia (review). International journal of molecular medicine *31*, 3-10.

Balch, W.E., Morimoto, R.I., Dillin, A., and Kelly, J.W. (2008). Adapting proteostasis for disease intervention. Science *319*, 916-919.

Baldari, S., Garufi, A., Granato, M., Cuomo, L., Pistritto, G., Cirone, M., and D'Orazi, G. (2017). Hyperglycemia triggers HIPK2 protein degradation. Oncotarget *8*, 1190-1203.

Berber, S., Llamosas, E., Thaivalappil, P., Boag, P.R., Crossley, M., and Nicholas, H.R. (2013). Homeodomain interacting protein kinase (HPK-1) is required in the soma for robust germline proliferation in C. elegans. Developmental dynamics : an official publication of the American Association of Anatomists *242*, 1250-1261.

Berber, S., Wood, M., Llamosas, E., Thaivalappil, P., Lee, K., Liao, B.M., Chew, Y.L., Rhodes, A., Yucel, D., Crossley, M.*, et al.* (2016). Homeodomain-Interacting Protein Kinase (HPK-1) regulates stress responses and ageing in C. elegans. Scientific reports *6*, 19582.

Berendzen, K.M., Durieux, J., Shao, L.W., Tian, Y., Kim, H.E., Wolff, S., Liu, Y., and Dillin, A. (2016). Neuroendocrine Coordination of Mitochondrial Stress Signaling and Proteostasis. Cell *166*, 1553-1563 e1510.

Blackwell, T.K., Sewell, A.K., Wu, Z., and Han, M. (2019). TOR Signaling in Caenorhabditis elegans Development, Metabolism, and Aging. Genetics *213*, 329-360.

Blaquiere, J.A., and Verheyen, E.M. (2017). Homeodomain-Interacting Protein Kinases: Diverse and Complex Roles in Development and Disease. Curr Top Dev Biol *123*, 73-103.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics *30*, 2114-2120.

Brignull, H.R., Moore, F.E., Tang, S.J., and Morimoto, R.I. (2006). Polyglutamine proteins at the pathogenic threshold display neuron-specific aggregation in a pan-neuronal Caenorhabditis elegans model. J Neurosci *26*, 7597-7606.

Calzado, M.A., Renner, F., Roscic, A., and Schmitz, M.L. (2007). HIPK2: a versatile switchboard regulating the transcription machinery and cell death. Cell Cycle *6*, 139-143.

Chakravarthy, B., Gaudet, C., Menard, M., Brown, L., Atkinson, T., Laferla, F.M., Ito, S., Armato, U., Dal Pra, I., and Whitfield, J. (2012). Reduction of the immunostainable length of the hippocampal dentate granule cells' primary cilia in 3xAD-transgenic mice producing human Abeta(1-42) and tau. Biochem Biophys Res Commun *427*, 218-222.

Chen, D., Thomas, E.L., and Kapahi, P. (2009). HIF-1 modulates dietary restriction-mediated lifespan extension via IRE-1 in Caenorhabditis elegans. PLoS Genet *5*, e1000486.

Chen, X.Y., Du, Y.F., and Chen, L. (2018). Neuropeptides Exert Neuroprotective Effects in Alzheimer's Disease. Front Mol Neurosci *11*, 493.

Cheung, J.Y., Leaf, A., and Bonventre, J.V. (1986). Mitochondrial function and intracellular calcium in anoxic cardiac myocytes. The American journal of physiology *250*, C18-25.

Chun, L., Gong, J., Yuan, F., Zhang, B., Liu, H., Zheng, T., Yu, T., Xu, X.Z., and Liu, J. (2015). Metabotropic GABA signalling modulates longevity in C. elegans. Nature communications *6*, 8828.

Cinar, H., Keles, S., and Jin, Y. (2005). Expression profiling of GABAergic motor neurons in Caenorhabditis elegans. Curr Biol *15*, 340-346.

Cornwell, A.B., Llop, J.R., Salzman, P., Thakar, J., and Samuelson, A.V. (2018). The Replica Set Method: A High-throughput Approach to Quantitatively Measure Caenorhabditis elegans Lifespan. Journal of visualized experiments : JoVE.

Cunningham, F., Allen, J.E., Allen, J., Alvarez-Jarreta, J., Amode, M.R., Armean, I.M., Austine-Orimoloye, O., Azov, A.G., Barnes, I., Bennett, R.*, et al.* (2022). Ensembl 2022. Nucleic Acids Res *50*, D988-D995.

Das, R., Melo, J.A., Thondamal, M., Morton, E.A., Cornwell, A.B., Crick, B., Kim, J.H., Swartz, E.W., Lamitina, T., Douglas, P.M.*, et al.* (2017). The homeodomain-interacting protein kinase HPK-1 preserves protein homeostasis and longevity through master regulatory control of the HSF-1 chaperone network and TORC1-restricted autophagy in Caenorhabditis elegans. PLoS Genet *13*, e1007038.

de la Vega, L., Grishina, I., Moreno, R., Kruger, M., Braun, T., and Schmitz, M.L. (2012). A redox-regulated SUMO/acetylation switch of HIPK2 controls the survival threshold to oxidative stress. Mol Cell *46*, 472-483.

de Magalhaes, J.P., and Toussaint, O. (2004). GenAge: a genomic and proteomic network map of human ageing. FEBS Lett *571*, 243-247.

Deak, F. (2014). Neuronal vesicular trafficking and release in age-related cognitive impairment. J Gerontol A Biol Sci Med Sci *69*, 1325-1330.

DeLano, W.L. (2002). Pymol: An open-source molecular graphics tool. CCP4 Newsl Protein Crystallogr *40*, 82-92.

Denzel, M.S., Lapierre, L.R., and Mack, H.I.D. (2019). Emerging topics in C. elegans aging research: Transcriptional regulation, stress response and epigenetics. Mech Ageing Dev *177*, 4-21.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

Douglas, P.M., Baird, N.A., Simic, M.S., Uhlein, S., McCormick, M.A., Wolff, S.C., Kennedy, B.K., and Dillin, A. (2015). Heterotypic Signals from Neural HSF-1 Separate Thermotolerance from Longevity. Cell Rep *12*, 1196-1204.

Durieux, J., Wolff, S., and Dillin, A. (2011). The cell-non-autonomous nature of electron transport chain-mediated longevity. Cell *144*, 79-91.

Ecsedy, J.A., Michaelson, J.S., and Leder, P. (2003). Homeodomain-interacting protein kinase 1 modulates Daxx localization, phosphorylation, and transcriptional activity. Mol Cell Biol 23, 950-960.

Entchev, E.V., Patel, D.S., Zhan, M., Steele, A.J., Lu, H., and Ch'ng, Q. (2015). A gene-expression-based neural code for food abundance that modulates lifespan. Elife 4, e06259.

Ewald, C.Y., Landis, J.N., Porter Abate, J., Murphy, C.T., and Blackwell, T.K. (2015). Dauer-independent insulin/IGF-1-signalling implicates collagen remodelling in longevity. Nature 519, 97-101.

Fawcett, E.M., Hoyt, J.M., Johnson, J.K., and Miller, D.L. (2015). Hypoxia disrupts proteostasis in Caenorhabditis elegans. Aging Cell 14, 92-101.

Frakes, A.E., Metcalf, M.G., Tronnes, S.U., Bar-Ziv, R., Durieux, J., Gildea, H.K., Kandahari, N., Monshietehadi, S., and Dillin, A. (2020). Four glial cells regulate ER stress resistance and longevity via neuropeptide signaling in C. elegans. Science 367, 436-440.

Garcia, S.M., Casanueva, M.O., Silva, M.C., Amaral, M.D., and Morimoto, R.I. (2007). Neuronal signaling modulates protein homeostasis in Caenorhabditis elegans post-synaptic muscle cells. Genes Dev 21, 3006-3016.

Garufi, A., Ricci, A., Trisciuoglio, D., Iorio, E., Carpinelli, G., Pistritto, G., Cirone, M., and D'Orazi, G. (2013). Glucose restriction induces cell death in parental but not in homeodomain-interacting protein kinase 2-depleted RKO colon cancer cells: molecular mechanisms and implications for tumor therapy. Cell death & disease 4, e639.

Gendrel, M., Atlas, E.G., and Hobert, O. (2016). A cellular and regulatory map of the GABAergic nervous system of C. elegans. Elife 5.

Gidalevitz, T., Ben-Zvi, A., Ho, K.H., Brignull, H.R., and Morimoto, R.I. (2006). Progressive disruption of cellular protein folding in models of polyglutamine diseases. Science 311, 1471-1474.

Gidalevitz, T., Kikis, E.A., and Morimoto, R.I. (2010). A cellular perspective on conformational disease: the role of genetic background and proteostasis networks. Curr Opin Struct Biol 20, 23-32.

Gilleard, J.S., Barry, J.D., and Johnstone, I.L. (1997). cis regulatory requirements for hypodermal cell-specific expression of the Caenorhabditis elegans cuticle collagen gene dpy-7. Mol Cell Biol 17, 2301-2311.

Hamilton, B., Dong, Y., Shindo, M., Liu, W., Odell, I., Ruvkun, G., and Lee, S.S. (2005). A systematic RNAi screen for longevity genes in C. elegans. Genes Dev 19, 1544-1555.

Hammarlund, M., Nix, P., Hauth, L., Jorgensen, E.M., and Bastiani, M. (2009). Axon regeneration requires a conserved MAP kinase pathway. Science 323, 802-806.

Hansen, M., Chandra, A., Mitic, L.L., Onken, B., Driscoll, M., and Kenyon, C. (2008). A role for autophagy in the extension of lifespan by dietary restriction in C. elegans. PLoS Genet 4, e24.

Hansen, M., Taubert, S., Crawford, D., Libina, N., Lee, S.J., and Kenyon, C. (2007). Lifespan extension by conditions that inhibit translation in Caenorhabditis elegans. Aging Cell 6, 95-110.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee, M.J., Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell 184, 3573-3587 e3529.

Hardaker, L.A., Singer, E., Kerr, R., Zhou, G., and Schafer, W.R. (2001). Serotonin modulates locomotory behavior and coordinates egg-laying and movement in Caenorhabditis elegans. J Neurobiol 49, 303-313.

Harris, M.A., Clark, J., Ireland, A., Lomax, J., Ashburner, M., Foulger, R., Eilbeck, K., Lewis, S., Marshall, B., Mungall, C., et al. (2004). The Gene Ontology (GO) database and informatics resource. Nucleic Acids Res 32, D258-261.

Harris, T.W., Arnaboldi, V., Cain, S., Chan, J., Chen, W.J., Cho, J., Davis, P., Gao, S., Grove, C.A., Kishore, R., et al. (2020). WormBase: a modern Model Organism Information Resource. Nucleic Acids Res 48, D762-D767.

Hart, A.C. (2006). Behavior. In The C elegans Research Community, WormBook.

Hattangadi, S.M., Burke, K.A., and Lodish, H.F. (2010). Homeodomain-interacting protein kinase 2 plays an important role in normal terminal erythroid differentiation. Blood 115, 4853-4861.

He, Q., Shi, J., Sun, H., An, J., Huang, Y., and Sheikh, M.S. (2010). Characterization of Human Homeodomain-interacting Protein Kinase 4 (HIPK4) as a Unique Member of the HIPK Family. Molecular and cellular pharmacology 2, 61-68.

Herndon, L.A., Schmeissner, P.J., Dudaronek, J.M., Brown, P.A., Listner, K.M., Sakano, Y., Paupard, M.C., Hall, D.H., and Driscoll, M. (2002). Stochastic and genetic factors influence tissue-specific decline in ageing C. elegans. Nature 419, 808-814.

Hietakangas, V., Ahlskog, J.K., Jakobsson, A.M., Hellesuo, M., Sahlberg, N.M., Holmberg, C.I., Mikhailov, A., Palvimo, J.J., Pirkkala, L., and Sistonen, L. (2003). Phosphorylation of serine 303 is a prerequisite for the stress-inducible SUMO modification of heat shock factor 1. Mol Cell Biol 23, 2953-2968.

Hietakangas, V., Anckar, J., Blomster, H.A., Fujimoto, M., Palvimo, J.J., Nakai, A., and Sistonen, L. (2006). PDSM, a motif for phosphorylation-dependent SUMO modification. Proc Natl Acad Sci U S A 103, 45-50.

Higuchi-Sanabria, R., Durieux, J., Kelet, N., Homentcovschi, S., de Los Rios Rogers, M., Monshietehadi, S., Garcia, G., Dallarda, S., Daniele, J.R., Ramachandran, V., et al. (2020). Divergent Nodes of Non-autonomous UPR(ER) Signaling through Serotonergic and Dopaminergic Neurons. Cell Rep 33, 108489.

Hindupur, S.K., Gonzalez, A., and Hall, M.N. (2015). The opposing actions of target of rapamycin and AMP-activated protein kinase in cell growth control. Cold Spring Harbor perspectives in biology 7, a019141.

Hobert, O. (2013). The neuronal genome of Caenorhabditis elegans. WormBook, 1-106.

Hodge, F., Bajuszova, V., and van Oosten-Hawle, P. (2022). The Intestine as a Lifespan- and Proteostasis-Promoting Signaling Tissue. Front Aging 3, 897741.

Hofmann, T.G., Stollberg, N., Schmitz, M.L., and Will, H. (2003). HIPK2 regulates transforming growth factor-beta-induced c-Jun NH(2)-terminal kinase activation and apoptosis in human hepatoma cells. Cancer Res 63, 8271-8277.

Huang, C., Xiong, C., and Kornfeld, K. (2004). Measurements of age-related changes of physiological processes that predict lifespan of Caenorhabditis elegans. Proc Natl Acad Sci U S A 101, 8084-8089. Epub 2004 May 8012.

Hwang, B., Lee, J.H., and Bang, D. (2018). Single-cell RNA sequencing technologies and bioinformatics pipelines. Experimental & molecular medicine 50, 1-14.

Hwang, B., Lee, J.H., and Bang, D. (2021). Author Correction: Single-cell RNA sequencing technologies and bioinformatics pipelines. Experimental & molecular medicine 53, 1005.

Imanikia, S., Ozbey, N.P., Krueger, C., Casanueva, M.O., and Taylor, R.C. (2019). Neuronal XBP-1 Activates Intestinal Lysosomes to Improve Proteostasis in C. elegans. Curr Biol 29, 2322-2338 e2327.

Jaisson, S., and Gillery, P. (2014). Impaired proteostasis: role in the pathogenesis of diabetes mellitus. Diabetologia *57*, 1517-1527.

Jassal, B., Matthews, L., Viteri, G., Gong, C., Lorente, P., Fabregat, A., Sidiropoulos, K., Cook, J., Gillespie, M., Haw, R.*, et al.* (2020). The reactome pathway knowledgebase. Nucleic Acids Res *48*, D498-D503.

Jin, Y., Hoskins, R., and Horvitz, H.R. (1994). Control of type-D GABAergic neuron differentiation by C. elegans UNC-30 homeodomain protein. Nature *372*, 780-783.

Johnson, D.W., Llop, J.R., Farrell, S.F., Yuan, J., Stolzenburg, L.R., and Samuelson, A.V. (2014). The Caenorhabditis elegans Myc-Mondo/Mad complexes integrate diverse longevity signals. PLoS Genet *10*, e1004278.

Jorgensen, E.M., and Nonet, M.L. (1995). Neuromuscular junctions in the nematode C. elegans. Paper presented at: seminars in DEVELOPMENTAL BIOLOGY (Elsevier).

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A.*, et al.* (2021). Highly accurate protein structure prediction with AlphaFold. Nature *596*, 583-589.

Kaletsky, R., Yao, V., Williams, A., Runnels, A.M., Tadych, A., Zhou, S., Troyanskaya, O.G., and Murphy, C.T. (2018). Transcriptome analysis of adult Caenorhabditis elegans cells reveals tissue-specific gene and isoform expression. PLoS Genet *14*, e1007559.

Kanehisa, M., and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids Res *28*, 27-30.

Ke, M., Elshenawy, B., Sheldon, H., Arora, A., and Buffa, F.M. (2022). Single cell RNA-sequencing: A powerful yet still challenging technology to study cellular heterogeneity. Bioessays *44*, e2200084.

Kenyon, C., Chang, J., Gensch, E., Rudner, A., and Tabtiang, R. (1993). A C. elegans mutant that lives twice as long as wild type. Nature *366*, 461-464.

Kikis, E.A., Gidalevitz, T., and Morimoto, R.I. (2010). Protein homeostasis in models of aging and age-related conformational disease. Advances in experimental medicine and biology *694*, 138-159.

Kim, E.A., Noh, Y.T., Ryu, M.J., Kim, H.T., Lee, S.E., Kim, C.H., Lee, C., Kim, Y.H., and Choi, C.Y. (2006). Phosphorylation and transactivation of Pax6 by homeodomain-interacting protein kinase 2. J Biol Chem *281*, 7489-7497.

Kraemer, B.C., Zhang, B., Leverenz, J.B., Thomas, J.H., Trojanowski, J.Q., and Schellenberg, G.D. (2003). Neurodegeneration and defective neurotransmission in a Caenorhabditis elegans model of tauopathy. Proc Natl Acad Sci U S A *100*, 9980-9985.

Kumsta, C., Chang, J.T., Schmalz, J., and Hansen, M. (2017). Hormetic heat stress and HSF-1 induce autophagy to improve survival and proteostasis in C. elegans. Nature communications *8*, 14337.

Labbadia, J., and Morimoto, R.I. (2015). The biology of proteostasis in aging and disease. Annu Rev Biochem *84*, 435-464.

Lakhani, R., Vogel, K.R., Till, A., Liu, J., Burnett, S.F., Gibson, K.M., and Subramani, S. (2014). Defects in GABA metabolism affect selective autophagy pathways and are alleviated by mTOR inhibition. EMBO Mol Med *6*, 551-566.

Lapierre, L.R., De Magalhaes Filho, C.D., McQuary, P.R., Chu, C.C., Visvikis, O., Chang, J.T., Gelino, S., Ong, B., Davis, A.E., Irazoqui, J.E.*, et al.* (2013). The TFEB orthologue HLH-30 regulates autophagy and modulates longevity in Caenorhabditis elegans. Nature communications *4*, 2267.

Lapierre, L.R., Kumsta, C., Sandri, M., Ballabio, A., and Hansen, M. (2015). Transcriptional and epigenetic regulation of autophagy in aging. Autophagy *11*, 867-880.

Lazaro-Pena, M.I., Cornwell, A.B., and Samuelson, A.V. (2021). Quantifying Tissue-Specific Proteostatic Decline in Caenorhabditis elegans. Journal of visualized experiments : JoVE.

Lazaro-Pena, M.I., Ward, Z.C., Yang, S., Strohm, A., Merrill, A.K., Soto, C.A., and Samuelson, A.V. (2022). HSF-1: Guardian of the Proteome Through Integration of Longevity Signals to the Proteostatic Network. Frontiers in Aging *3*.

Lee, P., Cho, B.R., Joo, H.S., and Hahn, J.S. (2008). Yeast Yak1 kinase, a bridge between PKA and stress-responsive transcription factors, Hsf1 and Msn2/Msn4. Mol Microbiol *70*, 882-895.

Lee, R.Y., Sawin, E.R., Chalfie, M., Horvitz, H.R., and Avery, L. (1999). EAT-4, a homolog of a mammalian sodium-dependent inorganic phosphate cotransporter, is necessary for glutamatergic neurotransmission in caenorhabditis elegans. J Neurosci *19*, 159-167.

Lee, S., Shang, Y., Redmond, S.A., Urisman, A., Tang, A.A., Li, K.H., Burlingame, A.L., Pak, R.A., Jovicic, A., Gitler, A.D.*, et al.* (2016). Activation of HIPK2 Promotes ER Stress-Mediated Neurodegeneration in Amyotrophic Lateral Sclerosis. Neuron *91*, 41-55.

Lee, S.J., Murphy, C.T., and Kenyon, C. (2009). Glucose shortens the life span of C. elegans by downregulating DAF-16/FOXO activity and aquaporin gene expression. Cell metabolism *10*, 379-391.

Lee, S.S., Kennedy, S., Tolonen, A.C., and Ruvkun, G. (2003). DAF-16 target genes that control C. elegans life-span and metabolism. Science *300*, 644-647.

Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics *12*, 323.

Li, N., and van der Kooy, D. (2018). Mutations in the guanylate cyclase gcy-28 neuronally dissociate naive attraction and memory retrieval. The European journal of neuroscience *48*, 3367-3378.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930.

Lin, K., Dorman, J.B., Rodan, A., and Kenyon, C. (1997). daf-16: An HNF-3/forkhead family member that can function to double the life-span of Caenorhabditis elegans. Science *278*, 1319-1322.

Lin, X.X., Sen, I., Janssens, G.E., Zhou, X., Fonslow, B.R., Edgar, D., Stroustrup, N., Swoboda, P., Yates, J.R., 3rd, Ruvkun, G.*, et al.* (2018). DAF-16/FOXO and HLH-30/TFEB function as combinatorial transcription factors to promote stress resistance and longevity. Nature communications *9*, 4400.

Lints, R., and Emmons, S.W. (1999). Patterning of dopaminergic neurotransmitter identity among Caenorhabditis elegans ray sensory neurons by a TGFbeta family signaling pathway and a Hox gene. Development *126*, 5819-5831.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402-408.

Lopez-Otin, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G. (2013). The hallmarks of aging. Cell *153*, 1194-1217.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Ma, R., Kutchy, N.A., Chen, L., Meigs, D.D., and Hu, G. (2022). Primary cilia and ciliary signaling pathways in aging and age-related
brain disorders. Neurobiology of disease *163*, 105607.
Mahoney, T.R., Luo, S., and Nonet, M.L. (2006). Analysis of synaptic transmission in Caenorhabditis elegans using an aldicarb-
sensitivity assay. Nature protocols *1*, 1772-1777.
Masse, I., Molin, L., Mouchiroud, L., Vanhems, P., Palladino, F., Billaud, M., and Solari, F. (2008). A novel role for the SMG-1 kinase
in lifespan and oxidative stress resistance in Caenorhabditis elegans. PLoS One *3*, e3354.
McIntire, S.L., Jorgensen, E., and Horvitz, H.R. (1993). Genes required for GABA function in Caenorhabditis elegans. Nature *364*,
334-337.
Miller, H.A., Dean, E.S., Pletcher, S.D., and Leiser, S.F. (2020). Cell non-autonomous regulation of health and longevity. Elife *9*.
Morelli, A., Sarchielli, E., Guarnieri, G., Coppi, E., Pantano, D., Comeglio, P., Nardiello, P., Pugliese, A.M., Ballerini, L., Matucci, R.,
et al. (2017). Young Human Cholinergic Neurons Respond to Physiological Regulators and Improve Cognitive Symptoms in an
Animal Model of Alzheimer's Disease. Front Cell Neurosci *11*, 339.
Morimoto, R.I. (2011). The heat shock response: systems biology of proteotoxic stress in aging and disease. Cold Spring Harb
Symp Quant Biol *76*, 91-99.
Morimoto, R.I. (2019). Cell-Nonautonomous Regulation of Proteostasis in Aging and Disease. Cold Spring Harbor perspectives in
biology.
Morley, J.F., Brignull, H.R., Weyers, J.J., and Morimoto, R.I. (2002). The threshold for polyglutamine-expansion protein aggregation
and cellular toxicity is dynamic and influenced by aging in Caenorhabditis elegans. Proc Natl Acad Sci U S A *99*, 10417-10422.
Morley, J.F., and Morimoto, R.I. (2004). Regulation of longevity in Caenorhabditis elegans by heat shock factor and molecular
chaperones. Mol Biol Cell *15*, 657-664. Epub 2003 Dec 2010.
Munkacsy, E., Khan, M.H., Lane, R.K., Borror, M.B., Park, J.H., Bokov, A.F., Fisher, A.L., Link, C.D., and Rea, S.L. (2016). DLK-1,
SEK-3 and PMK-3 Are Required for the Life Extension Induced by Mitochondrial Bioenergetic Disruption in C. elegans. PLoS
Genet *12*, e1006133.
Murakami, H., and Murakami, S. (2007). Serotonin receptors antagonistically modulate Caenorhabditis elegans longevity. Aging Cell
6, 483-488.
Nakamura, S., Karalay, O., Jager, P.S., Horikawa, M., Klein, C., Nakamura, K., Latza, C., Templer, S.E., Dieterich, C., and Antebi,
A. (2016). Mondo complexes regulate TFEB via TOR inhibition to promote longevity in response to gonadal signals. Nature
communications *7*, 10944.
Nayak, R., and Hasija, Y. (2021). A hitchhiker's guide to single-cell transcriptomics and data analysis pipelines. Genomics *113*, 606-
619.
Nisaa, K., and Ben-Zvi, A. (2021). Chaperone networks are shaped by cellular differentiation and identity. Trends Cell Biol.
Nonet, M.L., Staunton, J.E., Kilgard, M.P., Fergestad, T., Hartwieg, E., Horvitz, H.R., Jorgensen, E.M., and Meyer, B.J. (1997).
Caenorhabditis elegans rab-3 mutant synapses exhibit impaired function and are partially depleted of vesicles. J Neurosci *17*,
8061-8073.
Nye, K.J., Shi, Y.G., Andrews, J.M., and Wise, R. (1989). Pharmacokinetics and tissue penetration of cefepime. The Journal of
antimicrobial chemotherapy *24*, 23-28.
O'Rourke, E.J., and Ruvkun, G. (2013). MXL-3 and HLH-30 transcriptionally link lipolysis and autophagy to nutrient availability. Nat
Cell Biol *15*, 668-676.
Ogg, S., Paradis, S., Gottlieb, S., Patterson, G.I., Lee, L., Tissenbaum, H.A., and Ruvkun, G. (1997). The Fork head transcription
factor DAF-16 transduces insulin-like metabolic and longevity signals in C. elegans. Nature *389*, 994-999.
Oh, K.H., and Kim, H. (2017). Aldicarb-induced Paralysis Assay to Determine Defects in Synaptic Transmission in Caenorhabditis
elegans. Bio Protoc *7*.
Okkema, P.G., Harrison, S.W., Plunger, V., Aryana, A., and Fire, A. (1993). Sequence requirements for myosin gene expression
and regulation in Caenorhabditis elegans. Genetics *135*, 385-404.
Omura, D.T., Clark, D.A., Samuel, A.D., and Horvitz, H.R. (2012). Dopamine signaling is essential for precise rates of locomotion by
C. elegans. PLoS One *7*, e38649.
Oswal, N., Martin, O.M.F., Stroustrup, S., Matusiak Bruckner, M.A., and Stroustrup, N. (2021). A Hierarchical Process Model Links
Behavioral Aging and Lifespan in C. elegans. bioRxiv, 2021.2003.2031.437415.
Pan, K.Z., Palter, J.E., Rogers, A.N., Olsen, A., Chen, D., Lithgow, G.J., and Kapahi, P. (2007). Inhibition of mRNA translation
extends lifespan in Caenorhabditis elegans. Aging Cell *6*, 111-119.
Paradis, S., and Ruvkun, G. (1998). Caenorhabditis elegans Akt/PKB transduces insulin receptor-like signals from AGE-1 PI3
kinase to the DAF-16 transcription factor. Genes Dev *12*, 2488-2498.
Petrascheck, M., Ye, X., and Buck, L.B. (2007). An antidepressant that extends lifespan in adult Caenorhabditis elegans. Nature
450, 553-556.
Petrella, C., Di Certo, M.G., Barbato, C., Gabanella, F., Ralli, M., Greco, A., Possenti, R., and Severini, C. (2019). Neuropeptides in
Alzheimer's Disease: An Update. Current Alzheimer research *16*, 544-558.
Pickett, C.L., Breen, K.T., and Ayer, D.E. (2007). A C. elegans Myc-like network cooperates with semaphorin and Wnt signaling
pathways to control cell migration. Dev Biol *310*, 226-239.
Pober, B.R. (2010). Williams-Beuren syndrome. N Engl J Med *362*, 239-252.
Prahlad, V., Cornelius, T., and Morimoto, R.I. (2008). Regulation of the cellular heat shock response in Caenorhabditis elegans by
thermosensory neurons. Science *320*, 811-814.
Prahlad, V., and Morimoto, R.I. (2011). Neuronal circuitry regulates the response of Caenorhabditis elegans to misfolded proteins.
Proc Natl Acad Sci U S A *108*, 14204-14209.
Rahimi, M., Sohrabi, S., and Murphy, C.T. (2022). Novel elasticity measurements reveal C. elegans cuticle stiffens with age and in a
long-lived mutant. Biophys J *121*, 515-524.
Raich, W.B., Moorman, C., Lacefield, C.O., Lehrer, J., Bartsch, D., Plasterk, R.H., Kandel, E.R., and Hobert, O. (2003).
Characterization of Caenorhabditis elegans homologs of the Down syndrome candidate gene DYRK1A. Genetics *163*, 571-
580.
Richmond, J.E., and Jorgensen, E.M. (1999). One GABA and two acetylcholine receptors function at the C. elegans neuromuscular
junction. Nature neuroscience *2*, 791-797.

Riesen, M., Feyst, I., Rattanavirotkul, N., Ezcurra, M., Tullet, J.M., Papatheodorou, I., Ziehm, M., Au, C., Gilliat, A.F., Hellberg, J.*, et*
al. (2014). MDL-1, a growth- and tumor-suppressor, slows aging and prevents germline hyperplasia and hypertrophy in C.
elegans. Aging (Albany NY) *6*, 98-117.
Rinaldo, C., Prodosmo, A., Siepi, F., and Soddu, S. (2007). HIPK2: a multitalented partner for transcription factors in DNA damage
response and development. Biochemistry and cell biology = Biochimie et biologie cellulaire *85*, 411-418.
Rinaldo, C., Siepi, F., Prodosmo, A., and Soddu, S. (2008). HIPKs: Jack of all trades in basic nuclear activities. Biochim Biophys
Acta *1783*, 2124-2129.
Roux, A.E., Yuan, H., Podshivalova, K., Hendrickson, D., Kerr, R., Kenyon, C., and Kelley, D.R. (2022). The complete cell atlas of
an aging multicellular organism. bioRxiv, 2022.2006.2015.496201.
Sala, A.J., Bott, L.C., and Morimoto, R.I. (2017). Shaping proteostasis at the cellular, tissue, and organismal level. J Cell Biol *216*,
1231-1241.
Samuelson, A.V., Carr, C.E., and Ruvkun, G. (2007). Gene activities that mediate increased life span of C. elegans insulin-like
signaling mutants. Genes Dev *21*, 2976-2994.
Sardina, F., Conte, A., Paladino, S., Pierantoni, G.M., and Rinaldo, C. (2023). HIPK2 in the physiology of nervous system and its
implications in neurological disorders. Biochim Biophys Acta Mol Cell Res *1870*, 119465.
Scriabine, A., and Kazda, S. (1989). Pharmacological basis for use of calcium antagonists in hypertension. Magnesium *8*, 253-265.
Shemesh, N., Jubran, J., Dror, S., Simonovsky, E., Basha, O., Argov, C., Hekselman, I., Abu-Qarn, M., Vinogradov, E., Mauer, O.*,*
et al. (2021). The landscape of molecular chaperones across human tissues reveals a layered architecture of core and variable
chaperones. Nature communications *12*, 2180.
Shen, L.L., Wang, Y., and Wang, D.Y. (2007). Involvement of genes required for synaptic function in aging control in C. elegans.
Neurosci Bull *23*, 21-29.
Shinkai, Y., Yamamoto, Y., Fujiwara, M., Tabata, T., Murayama, T., Hirotsu, T., Ikeda, D.D., Tsunozaki, M., Iino, Y., Bargmann, C.I.*,*
et al. (2011). Behavioral choice between conflicting alternatives is regulated by a receptor guanylyl cyclase, GCY-28, and a
receptor tyrosine kinase, SCD-2, in AIA interneurons of Caenorhabditis elegans. J Neurosci *31*, 3007-3015.
Shojima, N., Hara, K., Fujita, H., Horikoshi, M., Takahashi, N., Takamoto, I., Ohsugi, M., Aburatani, H., Noda, M., Kubota, N.*, et al.*
(2012). Depletion of homeodomain-interacting protein kinase 3 impairs insulin secretion and glucose tolerance in mice.
Diabetologia *55*, 3318-3330.
Siesjo, B.K. (1989). Calcium and cell death. Magnesium *8*, 223-237.
Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Soding, J.*, et al.* (2011).
Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol *7*, 539.
Silva, M.C., Amaral, M.D., and Morimoto, R.I. (2013). Neuronal reprograming of protein homeostasis by calcium-dependent
regulation of the heat shock response. PLoS Genet *9*, e1003711.
Song, J.J., and Lee, Y.J. (2003a). Effect of glucose concentration on activation of the ASK1-SEK1-JNK1 signal transduction
pathway. J Cell Biochem *89*, 653-662.
Song, J.J., and Lee, Y.J. (2003b). Role of the ASK1-SEK1-JNK1-HIPK1 signal in Daxx trafficking and ASK1 oligomerization. J Biol
Chem *278*, 47245-47252.
Steinmetz, E.L., Dewald, D.N., and Walldorf, U. (2018). Homeodomain-interacting protein kinase phosphorylates the Drosophila
Paired box protein 6 (Pax6) homologues Twin of eyeless and Eyeless. Insect molecular biology *27*, 198-211.
Stroustrup, N., Ulmschneider, B.E., Nash, Z.M., Lopez-Moyado, I.F., Apfeld, J., and Fontana, W. (2013). The Caenorhabditis
elegans Lifespan Machine. Nature methods *10*, 665-670.
Suriyalaksh, M., Raimondi, C., Mains, A., Segonds-Pichon, A., Mukhtar, S., Murdoch, S., Aldunate, R., Krueger, F., Guimera, R.,
Andrews, S.*, et al.* (2022). Gene regulatory network inference in long-lived C. elegans reveals modular properties that are
predictive of novel aging genes. iScience *25*, 103663.
Sze, J.Y., Victor, M., Loer, C., Shi, Y., and Ruvkun, G. (2000). Food and metabolic signalling defects in a Caenorhabditis elegans
serotonin-synthesis mutant. Nature *403*, 560-564.
Tacutu, R., Shore, D.E., Budovsky, A., de Magalhaes, J.P., Ruvkun, G., Fraifeld, V.E., and Curran, S.P. (2012). Prediction of C.
elegans longevity genes by human and worm longevity networks. PLoS One *7*, e48282.
Tatum, M.C., Ooi, F.K., Chikka, M.R., Chauve, L., Martinez-Velazquez, L.A., Steinbusch, H.W., Morimoto, R.I., and Prahlad, V.
(2015). Neuronal serotonin release triggers the heat shock response in C. elegans in the absence of temperature increase.
Curr Biol *25*, 163-174.
Taylor, R.C., and Dillin, A. (2011). Aging as an event of proteostasis collapse. Cold Spring Harbor perspectives in biology *3*.
Taylor, R.C., and Dillin, A. (2013). XBP-1 is a cell-nonautonomous regulator of stress resistance and longevity. Cell *153*, 1435-1447.
Taylor, S.R., Santpere, G., Weinreb, A., Barrett, A., Reilly, M.B., Xu, C., Varol, E., Oikonomou, P., Glenwinkel, L., McWhirter, R.*, et*
al. (2021). Molecular topography of an entire nervous system. Cell *184*, 4329-4347 e4323.
Team, R.C. (2013). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing.
Tsunozaki, M., Chalasani, S.H., and Bargmann, C.I. (2008). A behavioral switch: cGMP and PKC signaling in olfactory neurons
reverses odor preference in C. elegans. Neuron *59*, 959-971.
Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan, D., Stroe, O., Wood, G., Laydon, A.*, et al.*
(2022). AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with
high-accuracy models. Nucleic Acids Res *50*, D439-D444.
Walter, L., Baruah, A., Chang, H.W., Pace, H.M., and Lee, S.S. (2011). The homeobox protein CEH-23 mediates prolonged
longevity in response to impaired mitochondrial electron transport chain in C. elegans. PLoS Biol *9*, e1001084.
Wang, S.J.H., Sinclair, D.A.R., Kim, H.Y., Kinsey, S.D., Yoo, B., Shih, C.R.Y., Wong, K.K.L., Krieger, C., Harden, N., and Verheyen,
E.M. (2020). Homeodomain-interacting protein kinase (Hipk) plays roles in nervous system and muscle structure and function.
PLoS One *15*, e0221006.
Wu, C., Karakuzu, O., and Garsin, D.A. (2021). Tribbles pseudokinase NIPI-3 regulates intestinal immunity in Caenorhabditis
elegans by controlling SKN-1/Nrf activity. Cell Rep *36*, 109529.
Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene ontology analysis for RNA-seq: accounting for selection
bias. Genome biology *11*, R14.

Yuan, F., Zhou, J., Xu, L., Jia, W., Chun, L., Xu, X.Z.S., and Liu, J. (2019). GABA receptors differentially regulate life span and
health span in C. elegans through distinct downstream mechanisms. Am J Physiol Cell Physiol *317*, C953-C963.
Zaru, R., Magrane, M., O'Donovan, C., and UniProt, C. (2017). From the research laboratory to the database: the Caenorhabditis
elegans kinome in UniProtKB. Biochem J *474*, 493-515.
Zhang, J., Pho, V., Bonasera, S.J., Holtzman, J., Tang, A.T., Hellmuth, J., Tang, S., Janak, P.H., Tecott, L.H., and Huang, E.J.
(2007). Essential function of HIPK2 in TGFbeta-dependent survival of midbrain dopamine neurons. Nature neuroscience *10*,
77-86.
Zhang, J., Shang, Y., Kamiya, S., Kotowski, S.J., Nakamura, K., and Huang, E.J. (2020). Loss of HIPK2 Protects Neurons from
Mitochondrial Toxins by Regulating Parkin Protein Turnover. J Neurosci *40*, 557-568.
Zhang, Q., Wu, X., Chen, P., Liu, L., Xin, N., Tian, Y., and Dillin, A. (2018). The Mitochondrial Unfolded Protein Response Is
Mediated Cell-Non-autonomously by Retromer-Dependent Wnt Signaling. Cell *174*, 870-883 e817.
Zhang, Y., Lanjuin, A., Chowdhury, S.R., Mistry, M., Silva-Garcia, C.G., Weir, H.J., Lee, C.L., Escoubas, C.C., Tabakovic, E., and
Mair, W.B. (2019). Neuronal TORC1 modulates longevity via AMPK and cell nonautonomous regulation of mitochondrial
dynamics in C. elegans. Elife *8*.
Zhang, Y., Shao, Z., Zhai, Z., Shen, C., and Powell-Coffman, J.A. (2009). The HIF-1 hypoxia-inducible factor modulates lifespan in
C. elegans. PLoS One *4*, e6348.
Zhang, Y., Zhang, X., Dai, Y., Song, M., Zhou, Y., Zhou, J., Yan, X., and Shen, Y. (2021). The decrease of intraflagellar transport
impairs sensory perception and metabolism in ageing. Nature communications *12*, 1789.
1865

A Neuronal overexpression

Survival %

rab-3p::hpk-1
WT

0   35

B Muscle overexpression

myo-3p::hpk-1
WT

0   35

C Intestinal overexpression

ges-1p::hpk-1
WT

0   35

D Hypodermis overexpression

dpy-7p::hpk-1
WT

0   35

Age (Days)

E

Body bends/30sec

****   ****

WT   *hpk-1(pk1393)*   *hpk-1(pk1393); rab-3p::hpk-1*

F

Survival %

hpk-1(pk1393); rab-3p::hpk-1
WT
hpk-1(pk1393)

0   25

Age (Days)

G

Survival %

***   **

WT   *hpk-1(pk1393)*   *hpk-1(pk1393); rab-3p::hpk-1*

C **hpk-1 (K176A) kinase mutation**

foci / animal
unc-54p::Q35::YFP

- control
- rab-3p::hpk-1 (K176A)

Days adulthood

D **hpk-1 (D272N) kinase mutation**

foci / animal
unc-54p::Q35::YFP

- control
- rab-3p::hpk-1 (D272N)

Days adulthood

E **hpk-1 (K176A) kinase mutation**

% paralyzed animal
(unc-54p::Q35::YFP)

- control
- rab-3p::hpk-1 (K176A)

Days adulthood

F **hpk-1 (D272N) kinase mutation**

% paralyzed animal
(unc-54p::Q35::YFP)

- control
- rab-3p::hpk-1 (D272N)

Days adulthood

A *rgef-1p::Q40::YFP*

i nerve ring

ii control

iii rab-3p::hpk-1

iv hpk-1(neRNAi)

F *unc-54p::Q35::YFP*

i control

ii rab-3p::hpk-1

iii hpk-1(neRNAi)

B #Foci / animal (*rgef-1p::Q40::YFP*) vs Days of adulthood — control, rab-3p::hpk-1

C Body bends (30sec) — control, rab-3p::hpk-1 **

D #Foci / animal (*rgef-1p::Q40::YFP*) vs Days of adulthood — empty vector, hpk-1(neRNAi) **

E Body bends (30sec) — empty vector, hpk-1(neRNAi) ****

G #Foci / animal (*unc-54p::Q35::YFP*) vs Days of adulthood — control, rab-3p::hpk-1 * * **

H Paralyzed / total animals (*unc-54p::Q35::YFP*) vs Days of adulthood — control, rab-3p::hpk-1

I #Foci / animal (*unc-54p::Q35::YFP*) vs Days of adulthood — empty vector, hpk-1(neRNAi) ****

J Paralyzed / total animals (*unc-54p::Q35::YFP*) vs Days of adulthood — empty vector, hpk-1(neRNAi) ****

K DCV — *unc-31* — Neuropeptide release; SCV — *unc-13* — Neurotransmitter release; Synapse

L Relative polyQ accumulation (aggregates in *hpk-1(ne)* to non -Tg): control 0.754, unc-31(e169) 0.84, unc-13(e51) 1.02

A

puncta / seam cell (GFP::LGG-1/Atg8)

control (ev) ・ GABAergic::hpk-1 (ev) ・ control (hsf-1) ・ GABAergic::hpk-1 (hsf-1) ・ control (mxl-2) ・ GABAergic::hpk-1 (mxl-2) ・ control (pha-4) ・ GABAergic::hpk-1 (pha-4) ・ control (hlh-30) ・ GABAergic::hpk-1 (hlh-30) ・ control (daf-16) ・ GABAergic::hpk-1 (daf-16)

**** **** ns **** ns ns

B

L4 animals ・ D1 animals

puncta / seam cell (GFP::LGG-1/Atg8)

control ・ GABAergic::hpk-1 ・ hlh-30(tm1978) ・ GABAergic::hpk-1; hlh-30(tm1978)

**** ns **** ns

A.

Young neurons → aging → Old neurons

expression of *hpk-1* & longevity-associated TF

Wild type

Age-associated increase maintains homeostasis

hpk-1 overexpression

Overexpression activates homeostatic mechanisms to preemptively mitigate damage

hpk-1 null mutant

Neuronal dysfunction, compromised proteostatic network

B.

Serotonergic neurons   GABAergic neurons

Heat →   HPK-1   HPK-1   ⊣ TORC1

Serotonin   GABA

Acute Stress signaling   Longevity signaling

Muscle tissue   HSF-1   MML-1::MXL-2/HLH-30/DAF-16   Hypodermal tissue

Molecular chaperones   Autophagy

Protein folding   Protein turnover

Proteostasis & Longevity