Population demographic history and evolutionary rescue: influence of a bottleneck event

Running Title (max 40 characters):
Demographic history and evolutionary rescue

Authors
Laure Olazcuaga[1,*], Beatrice Lincke[1], Sarah Delacey[1], Lily F. Durkee[1,2], Brett A. Melbourne[3] and Ruth A. Hufbauer [1,2]

[1] Department of Agricultural Biology, Colorado State University, Fort Collins, Colorado 80523, USA
[2] Graduate Degree Program in Ecology, Colorado State University, Fort Collins, Colorado 80523, USA
[3] Department of Ecology & Evolutionary Biology, University of Colorado, Boulder, Colorado 80309, USA

* corresponding author: Laure Olazcuaga – olaz.laure@gmail.com

Acknowledgments
L.O., B.L. and R.A.H. acknowledges support from the department Agricultural Biology at Colorado State University for the Ag Biology Undergraduate Research Fellowship. L.O., R.A.H. and B.A.M. acknowledge support from the US National Science Foundation (DEB-1930650 to R.A.H.; DEB-1930222 to B.A.M.).

Abstract

Rapid environmental change presents a significant challenge to the persistence of natural populations. Rapid adaptation that restores positive growth rate, enabling populations in decline following severe environmental change to avoid extinction, is called evolutionary rescue. Numerous studies have shown evolutionary rescue can indeed prevent extinction. Here, we extend those results by considering the demographic history of populations. To evaluate how demographic history influences evolutionary rescue, we created 84 populations of red flour beetle, *Tribolium castaneum*, with three classes of demographic history: diverse populations that did not experience a bottleneck, and populations that experienced either an intermediate or a strong bottleneck. We subjected these populations to a new and challenging environment for six discrete generations and tracked extinction and population size. Populations that did not experience a bottleneck in their demographic history avoided extinction entirely, while more than 20% of populations that experienced an intermediate or strong bottleneck went extinct. Similarly, among the extant populations at the end of the experiment, adaptation increased fitness in the novel environment the most for populations that had not experienced a bottleneck in their history. Taken together, these results highlight the importance of considering the demographic history of populations to make useful and effective conservation decisions and management strategies for populations experiencing environmental change that pushes them towards extinction.

Keywords (max 6):
adaptation, genetic variation, inbreeding depression, heterozygosity, extinction, *Tribolium castaneum*

Introduction

In the Anthropocene, populations experience frequent and intense changes to their environments (Bellard *et al.*, 2012). If environmental change is severe enough, population growth may slow to the point that population size starts to decline. If dispersal capabilities are limited or suitable environments are no longer available (Schloss *et al.*, 2012), populations must adapt to challenging environments or risk extinction (Smith *et al.*, 1989; Holt, 1990). Adaptation that is rapid enough to reverse population decline due to a challenging environment and prevent an otherwise inevitable extinction is called evolutionary rescue (Gomulkiewicz & Holt, 1995; Carlson *et al.*, 2014). This process is characterized by a U-shaped curve in population size through time: initially, population size decreases due to maladaptation to the environment, then population size increases as adaptive alleles spread through the population (Gomulkiewicz & Holt, 1995; Orr & Unckless, 2014).

Theory and experiments with model organisms show that evolutionary rescue can avert extinction. Theory clearly shows that the probability of evolutionary rescue increases with genetic variation(Gomulkiewicz & Holt, 1995; Barrett & Schluter, 2008) and suggests that it decreases with mutational load and inbreeding depression (Barrett & Kohn, 1991; Ellstrand & Elam, 1993). Despite confirmation of these effects in experiments with model organisms (Bijlsma *et al.*, 2010; Agashe *et al.*, 2011; Lachapelle & Bell, 2012; Ramsayer *et al.*, 2013; Hufbauer *et al.*, 2015), evidence of evolutionary rescue in nature remains scarce, suggesting that the conditions that allow evolutionary rescue to occur may be limited (Carlson *et al.*, 2014). One condition that is particularly important to examine experimentally is the demographic history of populations. In the wild, populations are frequently subjected to various stressful events that can lead to a temporary reduction in size, such as hunting or

habitat destruction. These reductions in population size, called bottleneck events

(Frankham *et al.*, 1999) can affect genetic variation, mutational load, and inbreeding

levels in the populations that experience them (hereafter bottlenecked populations).

Such bottleneck events, whether rare extreme events or repeated less-severe

restrictions in population size, can reduce the future adaptive potential of populations

(Nei *et al.*, 1975; Bouzat, 2010) and thus the probability of evolutionary rescue.

Bottlenecks influence the ability of populations to adapt to a future challenging

environment in two main ways (O'Brien & Evermann, 1988; Frankham *et al.*, 1999;

Reusch & Wood, 2007). First, standing genetic variation is lost by genetic drift during

bottleneck events due to an increase of stochastic events associated with small

population size (Hedrick, 2005). This reduction of genetic diversity decreases the

evolutionary potential of populations since potential future adaptive alleles are lost

(Futuyma, 1986). In general, adequate genome-wide genetic variation is important

for the conservation of natural populations (Kardos *et al.*, 2021). In addition, when

populations are small, selection is weak relative to genetic drift, thus reductions in

genetic diversity can include increases in frequency and fixation of deleterious alleles

(Lynch *et al.*, 1995), leading to populations with lower average fitness than

populations that have not experienced a bottleneck. However, given the stochastic

nature of which alleles go to fixation *versus* are lost, by chance, some beneficial

alleles may be fixed as well, potentially leading some bottlenecked populations to

have higher fitness (Bouzat, 2010). Thus, in addition to lower average fitness in

bottlenecked populations, greater variation in fitness is expected across different

bottlenecked populations compared to non-bottleneck populations.

The second main way that bottlenecks can influence the ability of populations

to adapt is through inbreeding. After bottleneck events, inbreeding increases due to

increased homozygosity of alleles common by descent (Wright, 1977; Ralls *et al.*,

1979; Frankham *et al.*, 1999; Kirkpatrick & Jarne, 2000; Fox *et al.*, 2008;

Charlesworth & Willis, 2009; Hedrick & Fredrickson, 2010). When homozygosity

increases at loci with deleterious alleles, inbreeding leads to inbreeding depression,

which can reduce fitness and increase the risk of extinction (Charlesworth &

Charlesworth, 1987; Frankham, 2005a, 2005b; O'Grady *et al.*, 2006). Inbreeding in

populations of intermediate size may also lead to purging of deleterious alleles

(Crnokrak & Barrett, 2002; Glémin, 2007; Facon *et al.*, 2011). Thus, like for genetic

drift, the inbreeding effects of bottlenecks can increase the variation in fitness among

populations.

Despite the strong evidence for a positive effect of genetic variation (Hedrick,

2005) and a negative effect of inbreeding depression (Frankham *et al.*, 1999) on the

rapid adaptation that is central to evolutionary rescue, it remains unclear what the

consequences of past bottleneck events during the demographic history are for the

future ability of a population to adapt to a new challenging environment. Given the

role of bottlenecks in reducing genetic variation and increasing genetic load

(Frankham *et al.*, 1999; Reusch & Wood, 2007), it is important to examine the effects

of bottlenecks on evolutionary rescue experimentally. In a recent study, Klerks *et al.*,

(2019) evaluated how population demographic history (bottlenecked or not)

influenced the response of experimental populations to selection for increased heat

tolerance. They found that bottlenecked populations had a 50% slower response to

selection than normal populations. This suggests an important role for bottlenecks in

the context of evolutionary rescue. Monitoring population dynamics was outside the

scope of Klerks *et al.*, (2019), and so it is not clear if the U-shaped curve

characteristic of evolutionary rescue occurred. Studying the population dynamics is,

however, necessary to understand the minimum population size that populations will

reach without going extinct or how long it will take for populations of a given size to

recover. In addition, the effects of bottleneck events in eukaryote populations are

most commonly studied with two types of populations: bottlenecked or non-

bottlenecked. The effect of bottleneck intensity has already been demonstrated in

bacterial experimental evolution, where the adaptative pathways were different

depending on whether the bottleneck was weak or intense (Vogwill *et al.*, 2016;

Garoff *et al.*, 2020; Mahrt *et al.*, 2021). Furthermore, bottlenecks can be more or less

intense in nature, which could influence subsequent population dynamics and

response to selection.

Here we study how bottleneck events of two intensities influence the process

of evolutionary rescue of populations in a severely challenging environment. We

created populations of *Tribolium castaneum* that had experienced no bottleneck in

their demographic history, an intermediate bottleneck, or a strong bottleneck, and

monitored their dynamics in a challenging environment. We evaluate how the

probability of extinction depends upon population demographic history as well as

their estimated heterozygosity throughout the experiment, and whether surviving

populations exhibit the U-shaped population trajectory characteristic of evolutionary

rescue and how that depends upon population demographic history. Finally, we

quantified adaptation to the new environment at the end of the experiment by

measuring the growth rate of all rescued populations and evaluating the evolution of

the rate of development, a life history trait that could be involved in adaptation

(Berner *et al.*, 2004).

Methods

Model system

We used *T. castaneum*, the red flour beetle, to test the effects of demographic
history on evolutionary rescue. *T. castaneum* populations were kept in 4 cm by 4 cm
by 6 cm acrylic containers with 20 g of a standard medium (95% wheat flour and 5%
brewer's yeast), which served as both habitat and a food source. We will henceforth
refer to a container and medium together as a patch. Multiple patches were used to
maintain each distinct laboratory population of *T. castaneum*, and single patches
were used in growth assays. The life cycle of *T. castaneum* was constrained to
mimic that of a seasonally breeding organism with non-overlapping generations and
a discrete dispersal phase (Melbourne & Hastings, 2008), a life history found
commonly among plants and animals. During laboratory maintenance, adult beetles
were placed in groups of 50 individuals per patch with fresh standard medium for 24
hours to reproduce and start the next generation. At the end of the reproduction
phase, adults were removed from the patches and the eggs were left to mature into
adults for an additional 34 days for a total of 35 days per generation. During
laboratory maintenance and the experiments, beetles were kept in incubators at
31°C with 54 ± 14% relative humidity. Several incubators were used, and patches
were randomized among and within incubators once each week to prevent
systematic effects of incubation conditions.

Creating populations with different demographic histories

A large diverse population was initiated from the multiple crosses of different stock
strains ("GA1", "Lab-S", "Z2", "Z4" and "Estill" from USDA stock maintenance). These
strains were originally all from samples from the US southwest, to minimize potential

genetic or cytoplasmic incompatibilities between strains. All possible pairwise

crosses indicated no reduction in fitness between any given pair (data not shown).

This diverse population was maintained and well mixed for approximately ten

generations at a population size of 10,000 individuals, distributed in four temporal

blocks with gene flow across the different temporal blocks. From this large and

diverse population, we created 60 independent replicate populations experiencing

bottlenecks in population size of two different intensities over two generations.

During the first generation of the bottleneck event, all 60 populations were reduced

to a size of two individuals. During the second generation of the bottleneck event, 24

populations were allowed to triple in size to six individuals to initiate the next

generation (hereafter called "intermediate bottleneck") and 36 populations were not

allowed to grow, and experienced a second bottleneck of two individuals, to initiate

the next generation (hereafter called "strong bottleneck"). Then these 60 populations

were maintained for six generations in the laboratory to increase population size

prior to starting the experiment (Figure 1). In contrast to the intermediate and strong

bottlenecked populations, the diverse population was maintained as a single large

population during those six generations to maintain high genetic diversity and was

used to create independent replicate populations at the beginning of the experiment

(hereafter called "diverse populations"). During the six generations, the population

size of each bottlenecked population was recorded so that the individual population

demographic history within each bottleneck category was known. This allowed us to

calculate the heterozygosity remaining relative to the source population at the start of

the experiment following Allendorf (1986). Heterozygosity is a good measure of

genetic variation, and how it decreases during a bottleneck event is well understood

(Allendorf, 1986). The expected heterozygosity at generation t (He_t), is a function of

expected heterozygosity and population size in generation t-1:

$$He_t = He_{t-1} - \left(\frac{1}{2N_{t-1}}\right) \qquad\qquad (1)$$

where He_{t-1} is the expected heterozygosity at the previous generation t-1 and N_{t-1}

is the population size at the previous generation t-1. To calculate the proportion of

expected heterozygosity remaining, expected heterozygosity of the source

population at the time of the bottleneck is defined as 1, that is He_0= 1. By tracking

population size each generation, we can calculate the proportion of expected

heterozygosity remaining each subsequent generation. This also allows us to

compare the strength of the bottlenecks imposed on our experimental populations

with other bottleneck situations. For example, the intermediate bottleneck is

comparable to a population remaining at a size of 50 individuals for about 37

generations, and the strong bottleneck is comparable to a population remaining at a

size of 50 individuals for about 57 generations.

Evolutionary rescue experiment

To test the effect of demographic history on evolutionary rescue, we compared the

probability of rescue of bottlenecked *versus* diverse populations. In total, we

compared the evolution of 36 diverse populations, 24 intermediate bottleneck

populations, and 36 strong bottleneck populations distributed over three temporal

blocks. To start the experiment, 100 individuals from each population were housed in

two patches connected via 2mm holes drilled in the sides. We challenged these

experimental populations over six discrete generations with a stressful medium that

consisted of 97% corn flour, 2.85% wheat flour, and 0.15% brewer's yeast, to

represent a resource-limited environment similar to Hufbauer *et al.* (2015). Each

generation, the adults in each population were censused completely, then the

individuals were put back equally in the two connected patches with fresh

challenging media for the 5-week life cycle as described above. For each population,

the proportion of expected heterozygosity was computed each generation (Eq. 1)

until the end of the experiment or the extinction of the population. A population was

considered extinct when the population size was 0 or 1 at census. At the end of the

experiment, the growth rate of each population was measured as the number of

offspring produced per 30 adults (replicated as much as possible depending on

population size). To evaluate potential differences in development time among the

populations that survived to the end of the experiment (hereafter, extant

populations), the composition of different developmental stages was enumerated

after 4 and 5 weeks of development by counting the number of larvae, pupae, and

adults in each patch.

Due to a protozoan parasite outbreak in the lab colonies, 12 populations

showing symptoms of contamination were excluded from the experiment (9, 1 and 2

populations for diverse-source, intermediate bottleneck, and strong bottleneck

populations respectively). With these exclusions, 23 diverse-source populations, 23

intermediate bottleneck populations and 34 strong bottleneck populations were

evaluated in the statistical analysis (Figure 1).

Statistical analysis

All statistical models were fitted in R version 3.4.2 (R Core Team, 2014). For all

models, we estimated the significance of interactions and main effects by comparing

a full model (i.e., with model terms of the same order) to a reduced model without the

interaction or main effect of interest using a Likelihood Ratio Test (LRT).

To understand how the probability of extinction in challenging environments was

influenced by demographic history, we analyzed the probability of extinction (binary

outcome) as a binomial distribution. However, none of the diverse populations went

extinct during the experiment. Thus, due to the separation problem caused by

estimated extinction probabilities exactly equal to zero, the full model including this

treatment could not be fitted (Albert & Anderson, 1984). To solve this issue, we used

a method of penalized likelihood (Firth, 1993; Heinze & Schemper, 2002). Hence, we

analyzed the probability of extinction using penalized logistic regression with Firth's

correction using the package *logistf* (Heinze *et al.*, 2022). Explanatory variables in

the model were demographic history (a categorical variable with three levels:

"diverse", "intermediate bottleneck " and "strong bottleneck") and temporal block (a

categorical variable with three levels), both as fixed effects. In addition, to test

whether time to extinction was influenced by demographic history, for the

populations that went extinct we analyzed the time to extinction (number of

generations) using a generalized linear model with a Poisson distribution and log

link. Explanatory variables in the model were the categorical demographic history

and the temporal block, again as fixed effects. We alternatively performed these

analyses using the expected heterozygosity (He_t) at the beginning of the experiment

as a continuous measure of demographic history with comparable results (we

present only the results using the model with the categorical variable).

A U-shaped curve in population size characterizes evolutionary rescue. To

evaluate how the shape of rescue curves differ by demographic history, we focused

on populations extant at the end of the experiment and analyzed size through time

using a Generalized Additive Mixed Model (GAMM; Wood, 2017) with a negative

binomial distribution to account for overdispersion using the package *mgcv* (Wood,

2022). Explanatory variables in the model were the categorical demographic history,

generation (a continuous variable from 2 to 6) modeled as a 1-dimensional smooth

effect, and the interaction between the two (thus allowing the smooth to vary by

demographic history), along with temporal block (three levels) as fixed effects and

population identity as a random effect. To model the shape of the rescue curve, the

data started at generation 2, the peak in abundance before the decline.

In addition, to quantify the variability of responses and visualize whether

outcomes of populations that experienced a bottleneck event are more variable than

of populations that did not, we computed the coefficients of variation in population

size across the three demographic histories across all generations for the extant

populations. We used the package *DescTools* to compute coefficients of variation

and their 95% confidence intervals (Signorell *et al.*, 2022). Considering the

coefficient of variation instead of the variance allows us to compare variation while

controlling for mean population size.

To test the influence of demographic history on adaptation, we evaluated

growth rate under standard density and the distribution of developmental stages of

extant populations. The growth rate (i.e., number of offspring divided by the number

of parents) is commonly analyzed with a lognormal transformation. Due to the

presence of 0 in our dataset, this direct transformation was not possible. To choose

the best distribution to analyze our data, we compared the distribution of residuals

from the growth rate analysis via an alternative lognormal transformation (log

transformation of growth rate+1), a square root transformation, and without

transformation. This preliminary analysis showed that analyzing the growth rate

without transformation was the best option. Hence growth rate was analyzed using a

linear model with a normal distribution. The development rate was evaluated by

comparing the presence of the different life stages (i.e. larva, pupa, or adult)

weighted by the number in each box, using a cumulative linkage model for ordinal

data to consider the order of appearance of each stage (larva, then pupa, then

adult), using the *ordinal* package (Christensen, 2015). For the growth rate model and

for the development rate model, explanatory variables in the model were the

categorical demographic history and temporal blocks as fixed effects, with population

identity included as a random effect to account for replication within populations. In

addition, the observation time (4 weeks or 5 weeks) was also included as a fixed

effect in the developmental rate model, as well as the replicate identity as a random

effect. We alternatively performed these analyses using the expected heterozygosity

at the end of the experiment as a continuous measure of demographic history with

comparable results (we present only the results using the model with the categorical

variable for the development stage analysis due to the absence of effect, see

results).

Results

Extinction and heterozygosity

There was a significant effect of demographic history on the probability of extinction

(LRT: χ_2^2 = 9.39, P < 0.001). Populations that had a history of a bottleneck

experienced more extinctions in the challenging environment than diverse

populations (22% and 24% of intermediate bottleneck and strong bottleneck

populations went extinct, Fig 2). The absence of bottleneck events during the

previous demographic history prevented extinction entirely (Fig. 2, left panel; 0%

extinction among diverse populations; estimate of the probability of extinction =

0.004, 95% CI [0.001, 0.05]). Considering only the demographic histories that

experienced extinctions, the probability of extinction was not significantly impacted

by bottleneck intensity during demography history (estimate of the probability of

extinction for intermediate bottlenecked populations: 0.06, [0.01; 0.27] and for strong

bottleneck populations: 0.06, [0.01; 0.28]). Among the populations that went extinct,

the number of generations to extinction were similar regardless of demographic

history, with largely overlapping confidence intervals (LRT: χ_1^2 = 0.21, P = 0.650;

estimate of the time of extinction for intermediate bottlenecked populations: 4.8

generations, 95% CI [3.1; 7.0] and for strong bottleneck populations: 5.0

generations, [3.6; 6.7] using the model including the demography history effect). For

populations that went extinct, there was not necessarily a gradual decrease in loss of

expected heterozygosity that led up to extinction (Fig. S1). Instead, heterozygosity

mostly declined markedly one generation before extinction (red-barred circles in Fig.

S1), reflecting the sharp population declines.

Population size dynamics

Following introduction into the challenging novel environment, population sizes

initially increased, due to the maternal effects from the high-quality environment the

populations were in just prior to the experiment being initiated, and then dropped

dramatically (Fig. 2). In extant populations, analysis of population size over time

reveals a U-shaped demographic curve indicative of evolutionary rescue, graphed in

Figure 3 with the predictions from the best-fit GAMM model where the shapes are

the same regardless the demographic history reflecting the most parsimonious

model (Figure 3A) and with the predictions from the full GAMM model allowing

different shapes depending on the demographic history, best representing the

experimental design (Figure 3B). These dynamics of rescue were similar regardless

of demographic history (LRT on the interaction between demographic history and

generation allowing different smooths: $\chi_6^2 = 3.0$, $P = 0.42$, meaning a lack of

evidence that the smooth are different for each demographic history; this effect is

present in model represented in Fig. 3A and is absent in the model represented in

Fig. 3B). Although the overall U-shape was apparent for all treatments, the model

estimate of population size was larger for the diverse population treatment than for

the intermediate-bottleneck and strong-bottleneck treatments (demographic history

effect: $\chi_2^2 = 15.43$, $P < 0.001$; Fig. 3A confidence interval for the diverse populations

never overlaps intermediate and strong bottleneck populations). The predicted mean

population size of the intermediate-bottleneck populations was slightly higher than

for strong-bottleneck populations, but confidence intervals overlapped for all

generations (Fig. 3A). By the last generation, the mean population size of the extant

populations was greater than the initial population size only for the diverse

populations (confidence interval not including 100 for the diverse demography

history; Figs. 2, 3A). Indeed, by the last generation, the diverse populations had

reached a mean population size of 127.9 (95% CI [107.3, 152.4]) individuals, while

the intermediate bottleneck and strong bottleneck populations had reached means of

86.8 [71.3, 105.7] individuals and 80.4 [67.3, 96.0] individuals respectively, according

to the predictions from the best-fit GAMM model (lines in Fig. 3A). These predictions

remain similar when using the full model that best represents the experimental

design (population size at the last generation for diverse populations: 143.5 [116.3,

177.2], for intermediate bottleneck: 74.6 [58.0, 96.0] and for strong bottleneck: 76.8

[61.6, 95.6]; Fig 3B). The predicted mean of the extant diverse populations was

always either above or overlapping the initial size (minimum of 96.0 individuals at 4.7

generations but with a confidence interval including 100 individuals). This was not

the case for the bottlenecked populations where the population size fell to a mean of

65.2 individuals and 60.3 individuals for the intermediate bottleneck and strong

bottleneck respectively (confidence interval not including 100 for generations 4 and 5

in Fig. 3A). These predictions remain similar when using the full model that best

represents the experimental design (minimum of the population size for diverse

populations: 99.8 [84.7, 117.6], for intermediate bottleneck: 62.0 [50.7, 75.8] and for

strong bottleneck: 58.4 [69.5, 49.0]; Fig 3B).

Coefficients of variation for population size across the three demographic

histories for almost all generations had overlapping confidence intervals (Fig 4). For

most generations, diverse populations had the lowest coefficient of variation, and at

generation 6, population sizes of the extant diverse populations were less variable

than the extant intermediate bottleneck and strong bottleneck populations ($CV_{diverse}$ =

0.29 [0.22, 0.41]; $CV_{intermediate}$ = 0.59 [0.42, 1.07], CV_{strong} = 0.51 [0.38, 0.77]; Fig. 4).

These results remained unchanged even when considering all populations, including

the populations that went extinct as having a population size of zero (results not

shown).

Growth rate and development time

Most of the extant populations for the three demographic histories had, on average

at the end of the experiment, a growth rate higher than 1, which means that more

descendants were produced than there were parents (Fig. 5). At the end of the

experiment, the growth rate was significantly different depending on the

demographic history (LRT for difference among histories: χ^2_2 = 15.64, P < 0.001).

The growth rate was higher for the diverse populations (estimate: 2.57, 95% CI

[2.20, 2.94]), than for bottlenecked populations (estimate: 1.70 [1.26, 2.13] for

intermediate and 1.87 [1.50, 2.23] for strong bottleneck populations). Overall, the

growth rate increased significantly with the heterozygosity of the populations (slope

estimate: 1.76 [0.85, 2.68]; LRT: $\chi_1^2 = 12.62$, $P < 0.001$).

The demographic history of rescued populations, had little to no influence on

developmental timing. The proportions of larvae, pupae, and adults were similar

among the three demographic histories (Fig. S2) and there was insufficient evidence

for a difference (LRT: $\chi_2^2 = 0.13$, $P = 0.94$). These proportions varied significantly

only with respect to timestep, i.e., between 4 weeks and 5 weeks (LRT: $\chi_1^2 = 2035.1$,

$P < 0.001$).

Discussion

In this study, we evaluated how the demographic history of populations influences

the probability of extinction and adaptation to a challenging environment. We found

that populations that experienced a bottleneck event in their demographic history

were more likely to go extinct, and, if they survived, were less well adapted to the

challenging environment than populations that had not experienced bottleneck

events.

Extinction associated with loss of heterozygosity for bottlenecked populations

While all diverse populations persisted throughout the six generations of the

experiment, multiple bottlenecked populations went extinct. These extinctions often

occurred in the generation following a strong decrease of the heterozygosity of the

population. However, our experimental design did not allow us to dissociate the

effect of losing heterozygosity from the effect of drop in population size.

Nonetheless, extinction only occurred in populations that underwent a bottleneck

event, which implies that decreases in heterozygosity (long- or short-term) can

strongly affect extinction.

Except for one population, none of the populations that experienced an initial

drastic drop in population size, decreasing their estimated heterozygosity below

0.45, were able to avoid extinction. This may be due to an extinction vortex, a

process in which small populations remain small due to a reinforcing negative

feedback loop that makes them vulnerable to extinction due to demographic

stochasticity (Gilpin & Soulé, 1986). The diverse populations did not decline to as

small a size as the bottlenecked populations, and perhaps this enabled them to

avoid falling into an extinction vortex. In the future, it would be interesting to know if

diverse populations that decline more sharply, due to stochasticity or experimental

manipulation, can escape more easily via rapid adaptation than populations that

have experienced bottleneck events.

Dynamics of rescue for extant populations

All treatments exhibited the U-shaped poplation size trajectory characteristic of

evolutionary rescue, when considering extant populations. The group of diverse

populations showed a higher level of adaptation, quantified by population growth

rate, than the bottleneck populations. These results confirm that rapid adaptation is

compromised following a bottleneck event (Bouzat, 2010). Our findings are

consistent with those of Klerks *et al.*, (2019) who showed that bottleneck populations

had a 50% slower response to selection for an increased heat tolerance than normal

populations. Further investigation should consider whether this difference is due to a

reduction of genetic variation in bottlenecked populations or an increase of

inbreeding depression. This information would facilitate efficient management of

populations where genetically variable and unrelated individuals are often not

available. For instance, it would increase the probability of success of a translocation

effort by targeting the individuals to be introduced (Ralls *et al.*, 2020). If the effects of

bottlenecks are primarily driven by a lack of genetic variation, individuals evolving in

different environments, even if they have a recent common ancestor, should be

considered for introductions to natural populations, to maximize the probability of

having a large pool of different alleles. However, if the effects of bottlenecks are

primarily driven by inbreeding depression, we suggest that unrelated individuals

should be introduced, even if they are all adapted to the same environment with all

the adaptive alleles to their environment potentially fixed and thus lacking genetic

variation in those traits.

Similar patterns of evolutionary rescue for intermediate and strong bottleneck

populations.

The evolutionary responses in terms of extinction, rescue, adaptive dynamics, and

traits involved were similar for populations that experienced intermediate bottlenecks

and populations that experienced strong bottlenecks. Three alternative hypotheses

can explain the general congruence between these treatments. First, the difference

in expected heterozygosity between intermediate and strong bottleneck events at the

start of the experiment was smaller than that between bottlenecked and diverse

populations (see Figure S1; averages at the start of the experiment: $He_{diverse} = 0.99$,

$He_{intermediate} = 0.68$, $He_{strong} = 0.54$). Both types of bottleneck reduced heterozygosity

substantially, but still some heterozygosity remained. It may be that in this range of

expected remaining heterozygosity, evolutionary responses to the environment are

similar. Second, populations that experienced a strong bottleneck event may have

purged deleterious alleles (Kirkpatrick & Jarne, 2000), reducing the negative fitness

effects of the bottleneck and making phenotypic outcomes more similar to the

intermediate bottleneck. The process of purging has been demonstrated in natural

populations. For instance, a recent analysis of the genomes of wild populations of

Alpine ibex that had undergone severe bottlenecks revealed a purge of highly

deleterious alleles (Grossen *et al.*, 2020). In our study, the level of adaptation to the

challenging environment may be comparable in intermediate and strong bottlenecks,

despite differences in heterozygosity, due to potential purging of deleterious alleles

in strong bottlenecks. Third, even if the phenotypic patterns seem similar, the

adaptive pathways used by these two types of populations may be different. For

instance, Vogwill *et al.*, (2016) found that antibiotic resistance evolved through highly

beneficial mutations for bacterial populations that experienced a strong bottleneck

event, whereas it evolved through a greater diversity of genetic mechanisms in

populations that experienced an intermediate bottleneck event. Differences in the

adaptive pathways may have consequences on the future evolutionary potential of

these populations.

Variance of evolutionary trajectories may depend on the demographic history

of populations.

Evolutionary responses were more variable in strong and intermediate bottlenecked

populations than in diverse populations but the confidence intervals overlapped

throughout the experiment, except for the last generation. Theoretical expectations

predict higher variance for bottlenecked populations due to sampling of the genetic

composition of each replicate population during a bottleneck event (Bouzat, 2010).

The lack of support for differing coefficients of variation at the beginning of the

experiments might be due to the fact that all populations were introduced into a new

environment and experience it in a similar way. In addition, contrary to theoretical

predictions and experimental results (Vogwill *et al.*, 2016; but see Windels *et al.*,

2021), we did not observe significantly higher coefficients of variation for strong

bottlenecks than for intermediate bottlenecks (see above for hypothetical

explanations).

Life history traits involved in the adaptation process

To better understand the different levels of adaptation between treatments, we

examined the proportions of individuals in different life stages at two time points at

the end of the experiment, to search for potential differences in development time.

We found that these proportions were on average very similar among demographic

histories. Life history traits other than development time may thus be responsible for

adaptation to the challenging environment. For instance, Agashe *et al.*, (2011)

demonstrated an increase of fecundity and egg survival, as well as a higher rate of

egg cannibalism, associated with adaptation to a new corn environment for

populations of *Tribolium castaneum*.

Conclusion and perspectives

Our findings have implications for conservation in our changing world. Managers

often have been concerned with maintaining adaptation to local environments of the

populations they manage (Dalrymple *et al.*, 2021). In the Anthropocene, where

environments are changing extremely rapidly, a focus on supporting sufficient

genetic variation for ongoing adaptation to shifting conditions is important. The

reduced ability of bottlenecked populations in this study to adapt to altered

conditions, however, suggests that more active management may be needed for

populations that have experienced bottlenecks. While genetic rescue (i.e., process of

alleviating inbreeding depression and restoring genetic diversity of inbred

populations through gene flow, Ralls *et al.*, 2020) involves additional avenues for

evolutionary rescue that were not addressed in this study, our work further supports

the substantial literature on this topic where immigrants are introduced to small

populations largely to reduce inbreeding depression but also to facilitate adaptation.

Data Archiving Statement

Data for this study are available at: to be completed after manuscript is accepted for

publication. The data and R scripts for our analyses are available at:

https://github.com/olazlaure/AnalysisEvolRescueExp

Author contributions

Conceptualization and funding acquisition: L.O. and R.A.H.;

Data acquisition: B.L., L.O. and S.D.;

Statistical analyses and writing: L.O. with inputs of B.L., S.D., L.F.D, B.A.M. and

R.A.H.

References

Agashe, D., Falk, J.J. & Bolnick, D.I. (2011) Effects of founding genetic variation on adaptation to a novel resource. *Evolution; International Journal of Organic Evolution*, **65**, 2481–2491.

Albert, A. & Anderson, J.A. (1984) On the Existence of Maximum Likelihood Estimates in Logistic Regression Models. *Biometrika*, **71**, 1–10.

Allendorf, F.W. (1986) Genetic drift and the loss of alleles versus heterozygosity. *Zoo Biology*, **5**, 181–190.

Barrett, R.D.H. & Schluter, D. (2008) Adaptation from standing genetic variation. *Trends in ecology & evolution*, **23**, 38–44.

Barrett, S.C.H. & Kohn, J.R. (1991) Genetic and evolutionary consequences of small population size in plants: implications for conservation. *Genetics and conservation of rare plants..*, 3–30.

Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. (2012) Impacts of climate change on the future of biodiversity. *Ecology letters*, **15**, 365–377.

Berner, D., Körner, C. & Blanckenhorn, W.U. (2004) Grasshopper populations across 2000 m of altitude: is there life history adaptation? *Ecography*, **27**, 733–740.

Bijlsma, R., Westerhof, M.D.D., Roekx, L.P. & Pen, I. (2010) Dynamics of genetic rescue in inbred *Drosophila melanogaster* populations. *Conservation Genetics*, **11**, 449–462.

Bouzat, J.L. (2010) Conservation genetics of population bottlenecks: the role of chance, selection, and history. *Conservation Genetics*, **11**, 463–478.

Carlson, S.M., Cunningham, C.J. & Westley, P.A.H. (2014) Evolutionary rescue in a changing world. *Trends in Ecology & Evolution*, **29**, 521–530.

Charlesworth, D. & Charlesworth, B. (1987) Inbreeding Depression and Its Evolutionary Consequences. *Annual Review of Ecology and Systematics*, **18**, 237–268.

Charlesworth, D. & Willis, J.H. (2009) The genetics of inbreeding depression. *Nature Reviews Genetics*, **10**, 783–796.

Christensen, R.H.B. (2015) ordinal: Regression Models for Ordinal Data.

Crnokrak, P. & Barrett, S.C.H. (2002) Perspective: Purging the Genetic Load: A Review of the Experimental Evidence. *Evolution*, **56**, 2347–2358.

Dalrymple, S.E., Winder, R. & Campbell, E.M. (2021) Exploring the potential for plant translocations to adapt to a warming world. *Journal of Ecology*, **109**, 2264–2270.

Ellstrand, N.C. & Elam, D.R. (1993) Population Genetic Consequences of Small Population Size: Implications for Plant Conservation. *Annual Review of Ecology and Systematics*, **24**, 217–242.

Facon, B., Hufbauer, R.A., Tayeh, A., Loiseau, A., Lombaert, E., Vitalis, R., *et al.* (2011) Inbreeding depression is purged in the invasive insect *Harmonia axyridis*. *Current Biology*, **21**, 424–427.

Firth, D. (1993) Bias reduction of maximum likelihood estimates. *Biometrika*, **80**, 27–38.

Fox, C.W., Scheibly, K.L. & Reed, D.H. (2008) Experimental Evolution of the Genetic Load and Its Implications for the Genetic Basis of Inbreeding Depression. *Evolution*, **62**, 2236–2249.

Frankham, R. (2005a) Stress and adaptation in conservation genetics. *Journal of Evolutionary Biology*, **18**, 750–755.

Frankham, R. (2005b) Genetics and extinction. *Biological Conservation*, **126**, 131–
140.
Frankham, R., Lees, K., Montgomery, M.E., England, P.R., Lowe, E.H. & Briscoe,
D.A. (1999) Do population size bottlenecks reduce evolutionary potential? *Animal*
Conservation, **2**, 255–260.
Futuyma, D.J. (1986) *Evolutionary Biology*. Sinauer Associates.
Garoff, L., Pietsch, F., Huseby, D.L., Lilja, T., Brandis, G. & Hughes, D. (2020)
Population Bottlenecks Strongly Influence the Evolutionary Trajectory to
Fluoroquinolone Resistance in Escherichia coli. *Molecular Biology and Evolution*,
37, 1637–1646.
Gilpin, M.E. & Soulé, M.E. (1986) Minimum viable populations : Processes of species
extinction. *Conservation biology: The science of scarcity and diversity*, 19–34.
Glémin, S. (2007) Mating Systems and the Efficacy of Selection at the Molecular
Level. *Genetics*, **177**, 905–916.
Gomulkiewicz, R. & Holt, R.D. (1995) When does evolution by natural selection
prevent extinction? *Evolution*, **49**, 201–207.
Grossen, C., Guillaume, F., Keller, L.F. & Croll, D. (2020) Purging of highly
deleterious mutations through severe bottlenecks in Alpine ibex. *Nature*
Communications, **11**, 1001.
Hedrick, P.W. (2005) A Standardized Genetic Differentiation Measure. *Evolution*, **59**,
1633–1638.
Hedrick, P.W. & Fredrickson, R. (2010) Genetic rescue guidelines with examples
from Mexican wolves and Florida panthers. *Conservation Genetics*, **11**, 615–626.
Heinze, G., Ploner, M., Dunkler, D., Southworth, H. & Jiricka, L. (2022) logistf: Firth's
Bias-Reduced Logistic Regression.
Heinze, G. & Schemper, M. (2002) A solution to the problem of separation in logistic
regression. *Statistics in Medicine*, **21**, 2409–2419.
Holt, R.D. (1990) The microevolutionary consequences of climate change. *Trends in*
Ecology & Evolution, **5**, 311–315.
Hufbauer, R.A., Szűcs, M., Kasyon, E., Youngberg, C., Koontz, M.J., Richards, C., *et*
al. (2015) Three types of rescue can avert extinction in a changing environment.
Proceedings of the National Academy of Sciences, **112**, 10557–10562.
Kardos, M., Armstrong, E.E., Fitzpatrick, S.W., Hauser, S., Hedrick, P.W., Miller,
J.M., *et al.* (2021) The crucial role of genome-wide genetic variation in
conservation. *Proceedings of the National Academy of Sciences*, **118**,
e2104642118.
Kirkpatrick, M. & Jarne, P. (2000) The Effects of a Bottleneck on Inbreeding
Depression and the Genetic Load. *The American Naturalist*, **155**, 154–167.
Klerks, P.L., Athrey, G.N. & Leberg, P.L. (2019) Response to Selection for Increased
Heat Tolerance in a Small Fish Species, With the Response Decreased by a
Population Bottleneck. *Frontiers in Ecology and Evolution*, **7**.
Lachapelle, J. & Bell, G. (2012) Evolutionary Rescue of Sexual and Asexual
Populations in a Deteriorating Environment. *Evolution*, **66**, 3508–3518.
Lynch, M., Conery, J. & Burger, R. (1995) Mutation accumulation and the extinction
of small populations. *American Naturalist*, **146**, 489–518.
Mahrt, N., Tietze, A., Künzel, S., Franzenburg, S., Barbosa, C., Jansen, G., *et al.*
(2021) Bottleneck size and selection level reproducibly impact evolution of
antibiotic resistance. *Nature Ecology & Evolution*, **5**, 1233–1242.
Melbourne, B.A. & Hastings, A. (2008) Extinction risk depends strongly on factors
contributing to stochasticity. *Nature*, **454**, 100–103.

Nei, M., Maruyama, T. & Chakraborty, R. (1975) The bottleneck effect and genetic
variability in populations. *Evolution*, **29**, 1–10.
O'Brien, S.J. & Evermann, J.F. (1988) Interactive influence of infectious disease and
genetic diversity in natural populations. *Trends in Ecology & Evolution*, **3**, 254–
259.
O'Grady, J.J., Brook, B.W., Reed, D.H., Ballou, J.D., Tonkyn, D.W. & Frankham, R.
(2006) Realistic levels of inbreeding depression strongly affect extinction risk in
wild populations. *Biological Conservation*, **133**, 42–51.
Orr, H.A. & Unckless, R.L. (2014) The Population Genetics of Evolutionary Rescue.
PLOS Genetics, **10**, e1004551.
R Core Team. (2014) R: A language environment for statistical computing.
Ralls, K., Brugger, K. & Ballou, J. (1979) Inbreeding and Juvenile Mortality in Small
Populations of Ungulates. *Science*, **206**, 1101–1103.
Ralls, K., Sunnucks, P., Lacy, R.C. & Frankham, R. (2020) Genetic rescue: A critique
of the evidence supports maximizing genetic diversity rather than minimizing the
introduction of putatively harmful genetic variation. *Biological Conservation*, **251**,
108784.
Ramsayer, J., Kaltz, O. & Hochberg, M.E. (2013) Evolutionary rescue in populations
of Pseudomonas fluorescens across an antibiotic gradient. *Evolutionary
Applications*, **6**, 608–616.
Reusch, T.B.H. & Wood, T.E. (2007) Molecular ecology of global change. *Molecular
Ecology*, **16**, 3973–3992.
Schloss, C.A., Nuñez, T.A. & Lawler, J.J. (2012) Dispersal will limit ability of
mammals to track climate change in the Western Hemisphere. *Proceedings of
the National Academy of Sciences*, **109**, 8606–8611.
Signorell, A., Aho, K., Alfons, A., Anderegg, N., Aragon, T., Arachchige, C., *et al.*
(2022) DescTools: Tools for Descriptive Statistics.
Smith, J.M., Chaloner, W.G. & Hallam, A. (1989) The causes of extinction.
*Philosophical Transactions of the Royal Society of London. B, Biological
Sciences*, **325**, 241–252.
Vogwill, T., Phillips, R.L., Gifford, D.R. & MacLean, R.C. (2016) Divergent evolution
peaks under intermediate population bottlenecks during bacterial experimental
evolution. *Proceedings of the Royal Society B: Biological Sciences*, **283**,
20160749.
Windels, E.M., Fox, R., Yerramsetty, K., Krouse, K., Wenseleers, T., Swinnen, J., *et
al.* (2021) Population Bottlenecks Strongly Affect the Evolutionary Dynamics of
Antibiotic Persistence. *Molecular Biology and Evolution*, **38**, 3345–3357.
Wood, S. (2022) mgcv: Mixed GAM Computation Vehicle with Automatic
Smoothness Estimation.
Wood, S.N. (2017) *Generalized additive models: an introduction with R*. Chapman &
Hall/CRC texts in statistical science. Second edition. CRC Press/Taylor & Francis
Group, Boca Raton.
Wright, S. (1977) *Evolution and the Genetics of Populations, Volume 3: Experimental
Results and Evolutionary Deductions*. University of Chicago Press, Chicago, IL.
685
686
687

Figures

Figure 1. Experimental design

695 **Figure 2. Population size of diverse, intermediate bottleneck, and strong**

696 **bottleneck populations through time.** *Red lines indicate populations that went*

697 *extinct.*

Figure 3. Population size of extant diverse populations (blue), intermediate bottleneck populations (red) and strong bottleneck populations (yellow) through time predicted (A) by the best-fit GAMM model or (B) by the full GAMM model including the interaction between time and demographic history. Each population is represented by a small dot. The lines represent the model estimate of population size for the three demographic history levels predicted by the GAMM model, from which generation 1 and the extinct populations data were excluded, and considering an average over the three temporal blocks for this representation. The best-fit GAMM model does not include an interaction between the demographic history and the time, which means that the shape of the predictions cannot differ by demographic history. The full GAMM model does include an interaction between demographic history and time, which means that the shape of the curve can differ by demographic history.The shaded and colored area around each line represents the 95% confidence interval. Points in different treatments are staggered slightly and jittered on the generation axis to reduce overlap.

Figure 4. Coefficient of variation of population size through time. *The coefficient of variation for each demographic history is represented by a point, with the 95% confidence interval represented by an error bar. Extinct populations were excluded but results were similar when they are considered. Points in different treatments are staggered slightly on the generation axis to reduce overlap.*

721
722
Figure 5. Growth rate assays for each extant population after the rescue
experiment depending on its final heterozygosity. *The mean growth rate across*
assays of each population is represented by a point, with the standard error
represented by an error bar. The points without error bars are from populations
where there was only one replicate assay to estimate the mean. The solid gray line
is the estimated mean relationship from the linear model. The shaded area around
the line represents the 95 percent confidence interval.