Divergent ecological responses to typhoon disturbance revealed via landscape-scale acoustic monitoring

Samuel R.P-J. Ross[1,2*], Nicholas R. Friedman[3,4], Kenneth L. Dudley[3], Takuma Yoshida[5], Masashi Yoshimura[5], Evan P. Economo[6], David W. Armitage[1], Ian Donohue[2]

[1]Integrative Community Ecology Unit, Okinawa Institute of Science and Technology Graduate University, Tancha, Onna-son, Okinawa, Japan.

[2]Zoology, School of Natural Sciences, Trinity College Dublin, Dublin, Ireland.

[3]Environmental Informatics Section, Okinawa Institute of Science and Technology Graduate University, Tancha, Onna-son, Okinawa, Japan.

[4]Centre for Taxonomy and Morphology, Leibniz Institute for the Analysis of Biodiversity Change, Hamburg, Germany.

[5]Environmental Science Section, Okinawa Institute of Science and Technology Graduate University, Tancha, Onna-son, Okinawa, Japan.

[6]Biodiversity & Biocomplexity Unit, Okinawa Institute of Science and Technology Graduate University, Tancha, Onna-son, Okinawa, Japan.

*Correspondence: Samuel RP-J Ross, Integrative Community Ecology Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan. E: s.ross.res@outlook.com

CRediT Author Contribution Statement | SRP-JR: Conceptualisation, Methodology, Software, Validation, Formal analysis, Investigation, Writing – Original Draft, Visualisation, Funding Acquisition; **NRF**: Conceptualisation, Software, Resources, Data Curation, Writing – Reviewing & Editing, Project Administration; **KLD**: Software, Validation, Resources, Data Curation, Visualisation, Project Administration; **TY**: Resources, Data Curation, Project Administration; **MY**: Resources, Data Curation, Project Administration; **EPE**: Resources, Data Curation, Writing – Reviewing & Editing, Project Administration, Funding Acquisition; **DWA**: Methodology, Software, Formal Analysis, Visualisation, Writing – Reviewing & Editing; **ID**: Conceptualisation, Methodology, Writing – Original Draft.

Running title: Typhoon-induced spatial ecological divergence

Key words: Soundscape, passive acoustic monitoring, sensor array, extreme weather event, spatial insurance, ecological stability, acoustic indices, Okinawa

Abstract

Climate change is increasing the frequency, intensity, and duration of extreme weather events across

the globe. Understanding the capacity for ecological communities to withstand and recover from such

events is critical. Typhoons are extreme weather events that are expected to broadly homogenise

ecosystems through structural damage to vegetation and longer-term effects of salinization. Given

their unpredictable nature, monitoring ecological responses to typhoons is challenging, particularly

for mobile animals such as birds. Here, we report spatially variable ecological responses to typhoons

across terrestrial landscapes. Using a high temporal resolution passive acoustic monitoring network

across 24 sites on the subtropical island of Okinawa, Japan, we found that typhoons elicit divergent

ecological responses among Okinawa's diverse terrestrial habitats, as indicated by increased spatial

variability of biological sound production (biophony) and individual species detections. This suggests

that soniferous communities have a diversity of pathways through which they respond to typhoons.

That is, spatial insurance effects among local ecological communities provide resilience to typhoons at

the landscape scale. Even though site-level typhoon impacts on soundscapes and bird detections

were not particularly strong, the data resolution afforded to us by monitoring at scale (high temporal

resolution, broad spatial extent) nevertheless enabled detection of spatial heterogeneity in typhoon

responses. Further, species-level responses mirrored those of acoustic indices, underscoring the

utility of such indices for revealing insight into fundamental questions concerning disturbance and

stability. Our findings demonstrate the significant potential of landscape-scale acoustic sensor

networks to capture the understudied ecological impacts of unpredictable extreme weather events.

Introduction

Climate change is increasing both the frequency and destructive potential of extreme

weather events such as typhoons (Bhatia et al., 2019; Emanuel, 2005; Kossin et al., 2020; Li &

Chakraborty, 2020). Typhoons bring exceptional winds and rainfall and can cause structural damage

to terrestrial habitats through wind-damage (Everham & Brokaw, 1996), flooding (Gardner et al.,

1991), and salinization by sea spray, particularly on islands (Elliott & Nino, 1960; Kerr, 2000). These

destructive forces can impact the mortality, composition, and dynamics of trees (Lin et al., 2020;

Morimoto et al., 2021), birds (Cely, 1991; Chevalier et al., 2019; Seki, 2005), bacteria (Ares et al.,

2020), invertebrates (Azuma et al., 1997; Willig & Camilo, 1991), and other animals (Donihue et al.,

2018; Pavelka et al., 2007; Testard et al., 2021).

Land cover may affect exposure to—and hence the ecological consequences of—extreme

weather events (Laurance, 1998). Several studies suggest that primary forest with a mix of vegetation

types and life forms is most likely to resist and recover from typhoon disturbance (Abbas et al., 2020;

Zampieri et al., 2020). Human activity and associated land use change has, therefore, considerable

potential to modify the severity and reach of typhoons (Raymond et al., 2020). For example,

deforestation through urbanisation or agricultural intensification leads to drastically simplified canopy

structure, thereby increasing typhoon exposure, as well as subjecting ecological communities to the

stressors and pollutants associated with these anthropogenic land uses (Daskalova et al., 2020;

Senzaki et al., 2020; Uchida & Ushimaru, 2014). In contrast, land abandonment might be expected to

dampen the effects of typhoons over time as intensively managed agricultural areas undergo natural

succession to a more biologically and structurally diverse system (Sirami et al., 2008; Uchida &

Ushimaru, 2014). Yet, despite the potential for land cover to moderate the impact of extreme events,

the practical difficulties associated with ecological monitoring at scale have, to date, limited

understanding of the extent of any differential impacts of typhoons across habitat types and of the

consequences of such differences for landscape-scale biodiversity or spatial processes such as

metacommunity dynamics and stability (Loreau et al., 2003; Wang et al., 2021).

Ecological stability is a central framework for considering disturbance impacts across spatial,

temporal, and organisational scales, from populations to ecosystems (Hillebrand et al., 2018; Kéfi et

al., 2019). Stability is a concept with multiple dimensions (Donohue et al., 2013; Hillebrand et al.,

2018; Pimm, 1984), including components such as resistance to and recovery from disturbance (Baert

et al., 2016; Yang et al., 2019), and the variability of ecological variables both in time and space

(Tilman et al., 2006; Wang et al., 2017). Both disturbance events and ecological responses to such

events vary across spatiotemporal scales (Clark et al., 2021; Ross et al., 2021b; Zelnik et al., 2018).

This necessitates high-resolution and long-term monitoring of ecosystems to holistically capture the

ecological impacts of infrequent extreme events such as typhoons. However, monitoring biodiversity

over large spatial and temporal scales poses considerable logistical and financial challenges.

Accordingly, most empirical studies of ecological stability are experimental (Kéfi et al., 2019), while

observational studies of disturbance typically employ space-for-time substitutions (Butsic et al.,

2017), or consider only single-time snapshots before and after disturbance (*e.g.*, Burivalova et al.,

2014). In such cases, it is extremely challenging to isolate the relevant pathways through which

disturbance events impact ecosystems in a holistic multidimensional way.

Recent advances in automation hold promise for understanding disturbance responses

through large-scale continuous monitoring of biodiversity (Keitt & Abelson, 2021; Ross et al., 2023).

Following developments in data acquisition, storage, and processing, passive acoustic monitoring of

wildlife and soundscapes is growing in popularity (Burivalova et al., 2019; Gibb et al., 2019). As sensor

networks are established to collect acoustic data autonomously (Keitt & Abelson, 2021; Sethi et al.,

2020), a diverse range of ecological studies become tractable by leveraging high-resolution acoustic

time series (*e.g.*, Deichmann et al., 2018; Lomolino et al., 2015; Rossi et al., 2017; Sueur et al., 2019; Ross et al., 2023). Studies of disturbance impacts on *soundscapes*—that is, all sound produced in an ecosystem (Pijanowski et al., 2011a, 2011b), including *biophony* (biotic sound), *geophony* (natural abiotic sound, such as rain), and *anthropophony* (human-related sound)—have recently emerged, though most commonly still make before-and-after or space-for-time comparisons (*e.g.*, Deichmann et al., 2017; Gasc et al., 2018). However, the high-resolution time series afforded by passive acoustic monitoring allows opportunistic measurement of soundscape responses to infrequent disturbance events, such as typhoons (*e.g.*, Gottesman et al., 2021), as well as documenting longer-term trends under climate change (Sueur et al., 2019). Acoustic monitoring thus provides an opportunity to overcome many of the challenges associated with studying extreme weather events, by allowing pre- and post-typhoon comparisons (Altwegg et al., 2017; Rajan et al., 2022), and capturing ecological responses to typhoons across scales in space and time (Lin et al., 2020) using a multidimensional stability framework (Donohue et al., 2013). Of the few studies that have used acoustic monitoring to capture storms or extreme events, most focused on marine soundscapes (Boyd et al., 2021; Locascio & Mann, 2005; Simmons et al., 2021), though Gottesman *et al*. (2021) recently showed that terrestrial soundscapes were less resistant than those of coral reefs to hurricane disturbance. Embedded within terrestrial soundscapes, bird vocalisations provide the opportunity to assess the impact of typhoons on critical indicator taxa (Gasc et al., 2017), while acoustic indices provide rapid information on a combination of biodiversity and other meaningful aspects of soundscape change (Bradfer-Lawrence et al., 2020; Harris et al., 2016; Rajan et al., 2022). There are, however, few studies that simultaneously assess both individual species vocalisations and acoustic indices explicitly (Ferreira et al., 2018; Ross et al., 2018).

Here, we exploit a dataset that is, to our knowledge, the highest-resolution dataset recording biological responses to an extreme weather event to date, capturing daily bird vocalisations and half-hourly acoustic indices in response to two large typhoons across 24 field sites on the island of Okinawa, Japan. We measure multiple dimensions of ecological stability for both soundscapes and individual bird species in response to a super-typhoon in September 2018, which was followed five days later by an extratropical cyclone. Our study spans Okinawa's full range of terrestrial habitats, allowing us to examine how land use can shape ecological responses to extreme events. Given that organisms are differently vulnerable to mortality and mechanical damage resulting from typhoons (Abbas et al., 2020; Zampieri et al., 2020), we expect land use to influence typhoon responses (Raymond et al., 2020).

Specifically, we test the hypotheses that typhoons (1) temporarily reduce soundscape richness and (2) bird vocalisation rates, and (3) homogenise soundscapes across sites. We also predict

that (4) natural forest habitats should have soundscapes that are more resistant to typhoons owing to

their closed canopy structure (Abbas et al., 2020; Nimmo et al., 2016). We expect to find differences

in bird species responses to typhoons, perhaps as a function of their traits (Wiley & Wunderle, 1993).

Closed canopy specialists, frugivores, granivores, and nectarivores should be most vulnerable to food

resource losses following typhoons (Chevalier et al., 2019; Wiley & Wunderle, 1993; Zhang et al.,

2016), while insectivores may benefit from increased access to prey in canopy gaps (Cely, 1991; Seki,

2005), and large-bodied or predatory birds may be especially vulnerable to typhoon-induced habitat

alteration, owing to their dependence on prey availability and habitat area (*e.g.*, Ross et al., 2019) and

their slow reproductive rates (Cely, 1991; Cohen et al., 2021; Wiley & Wunderle, 1993; see Table S1a

for summaries of our hypotheses for the focal species in this study). These hypotheses draw on the

idea that forest loss is a key catalyst of biodiversity change (Daskalova et al., 2020; Gibson et al.,

2011). This is especially pertinent given the high richness and rates of endemicity and specialism

among Okinawa's forest taxa (Inoue et al., 2019; Itô et al., 2000). Okinawa island is subject to rapid

and ongoing land use change, particularly through deforestation for urbanisation and agricultural

intensification (Ross et al., 2018; Takeuchi et al., 1981). Such land use change necessitates an explicit

focus on habitat degradation as a driver of the ecological outcomes of intensifying natural

disturbance regimes under climate change, including an increase in the frequency and destructive

potential of typhoons and extreme storms around Okinawa (A. Iwasaki, unpublished data).

Methods

Study sites and typhoon impact

This study uses data from the OKEON (Okinawa Environmental Observation Network)

Churamori Project (OKEON 美ら森プロジェクト; https://okeon.unit.oist.jp/) in Okinawa, Japan. We

use data from OKEON's 24 field sites across the island of Okinawa, representing Okinawa's full range

of land cover types (Figure 1; elevational range: 0.65 – 374.5 m). Elsewhere, we describe the

geographic variation among the sites (Ross *et al*., 2018), which were assessed using reflectance

estimates from Landsat 8 images to estimate proportional land cover for various land cover classes

within a 1,000 m circular buffer surrounding each site. This buffer size was chosen as it represents a

trade-off between local habitat structure and regional landscape features, which together shape the

diversity of highly mobile taxa such as birds (Clough *et al.*, 2009). Three pairs of sites had partly

overlapping buffer areas at this scale: Kurashiki and Tounan (6.5% overlap), OIST forest and OIST

campus (21% overlap), and Senbaru and Uehara (60% overlap). The centroid of all sites was

sufficiently distant such that no individual animals could be simultaneously detected by more than a

single recording unit, so we deemed our measurements from these sites to be suitably independent

for this study. We classified land cover into the following categories: dense closed-canopy forest;

grassland and scrubland (that is, low-medium growth coastal and disturbed vegetation, and managed

grasses); agricultural land (primarily for sugarcane); urban areas characterised by materials such as

asphalt and concrete with limited vegetation; sand and dirt with limited vegetation; freshwater

bodies; and miscellaneous land cover not described in the above categories. To deal with the

challenge of multicollinearity among land cover classes, we used an unsupervised learning approach

to identify clusters of sites with similar land cover. We used k-means clustering (optimal k = 2 clusters)

to identify sites that clearly differentiated in Principal Component Analysis (PCA) space, where the

first PCA axis captured Okinawa's primary land use gradient and explained 81.2% of the variance

among our sites (Supplementary Figure S1). The PCA loadings show that the two clusters identified

represent a distinction between sites that are primarily forested and those that are either agricultural

or urban (Figures 1b and S1), hereafter together referred to as 'developed' sites.

Acoustic data has been collected at each OKEON site since February 2017, but here we focus

on a 66-day period in 2018 surrounding the landfall of two large typhoons, Trami and Kong-Rey. Trami

passed closest to Okinawa on 29 September 2018 and was followed closely by Kong-Rey on 4 October

2018 (Japan Meteorological Agency [JMA] 2018; Figure 1a). We isolated recordings from the 30-day

periods before (*pre-disturbance* period: 30 August – 28 September 2018) and after (*post-disturbance*

period: 06 Oct – 04 Nov 2018) the typhoons made landfall, comprising a total of 771,840 minutes of

data (Figure 1). Okinawa and the Ryukyu archipelago are increasingly exposed to more frequent and

intense typhoons (A. Iwasaki, unpublished data), with annual typhoon seasons bringing disturbance

events of varying magnitude (Elliott & Nino, 1960). Typhoon Trami was the largest typhoon (tropical

cyclone) to hit Okinawa since OKEON acoustic recording began, with windspeeds reaching 183 km h^{-1}

on 29 September 2018 (JMA 2018). Trami was followed shortly after by Kong-Rey, which was less

severe, striking Okinawa as an extratropical cyclone (JMA 2018). The chosen acoustic recordings

therefore include a well-characterised pre-disturbance state (Figure S2; Ross *et al.*, 2018, 2021a),

followed by an extreme weather event and post-disturbance period during which soundscapes could

potentially recover to their pre-disturbance state (Figures 1c and 1d).

Figure 1.

typhoons

and extra

Okinawa,

2018 for

triangles are sites grouped in the forested site cluster (*n* = 10), and grey circles those in the developed cluster (*n*

= 14) based on unsupervised *k*-means clustering of land cover variables (see *Methods*). (c-d) Illustrative example

time series of the study period at the developed Chatan field site (see b), showing the dates of the typhoon

arrival (marked with coloured arrows) and the 30-day periods preceding and following typhoon impact

(delineated with dashed white bounding boxes). Dates along the X-axes span the full study period (30 Aug-04

Nov), and times along the Y-axes span 00:00-23:30 in half-hour intervals). Each grid cell then represents the

value of a detrended and normalised acoustic index for the 10-minute recording corresponding to each time-by-

date combination. To illustrate the potential for acoustic indices to reveal typhoon impacts, we show (c) the

median of the amplitude envelope (*M*; see Ross et al., 2021a), where higher values (lighter colours) represent

louder soundscapes across all frequency bands (Depraetere et al., 2012), and (d) the normalised difference

soundscape index (NDSI), where higher values (lighter colours) represent a dominance of biophony in the

soundscape, while lower values (darker colours) comprise mostly anthropophony (Kasten et al., 2012). Note the

signal of the typhoons on the soundscape, clear in (c) as an increase in total soundscape volume as the

typhoons pass Okinawa, and in (d) as a decline in the relative contribution of biophony (lighter colours) to the

soundscape following the typhoons. Geophony such as the heavy wind and rain caused by typhoons typically

produces broadband sound, so does not produce a clear signal in (d), unlike in (c).

Acoustic monitoring and data processing

Song Meter SM4 recorders (Wildlife Acoustics Inc., Concord, MA, USA) were installed at

approximately breast height (~1.3m) at each field site and were programmed to record at default gain

settings (+16 dB) via two omnidirectional microphones on a schedule of 10-minutes recording, 20-

minutes standby, with recording starting on every hour and half hour. Data were saved to an SD card

in stereo .WAV format at a sampling rate of 48-kHz. All audio data collected as part of the OKEON

Churamori Project were archived with the Okinawa Institute of Science and Technology's high-

performance computing centre.

For each 10-minute audio file, we computed three commonly used acoustic indices in R

(version 4.2.1; R Core Team 2022) using the *soundecology* package (version 1.3.3; Villanueva-Rivera &

Pijanowski 2018). We calculated the normalised difference soundscape index (NDSI) and its two

component indices, biophony ($NDSI_{Bio}$) and anthropophony ($NDSI_{Anthro}$), by first generating a

spectrogram via fast Fourier transformation (Hanning window size = 256) and splitting it into 1-kHz

frequency bands. Biophony and anthropophony are then calculated as the sum of the amplitude of all

1-kHz bands in, respectively, the 2-11 kHz and 1-2 kHz frequency ranges (Kasten et al., 2012). NDSI is

calculated as the ratio between these two components, such that higher values indicate a larger

proportion of biophony in the soundscape relative to anthropophony; NDSI scales -1 to +1, where -1

indicates complete dominance of anthropophony (low-frequency sound) whereas +1 indicates total

biophony (Kasten et al., 2012). This approach is preferable in our case over the original suggestion to

compare anthropophony with the highest amplitude frequency band from the biophony range

(Kasten et al., 2012), since it provides less weight to anthropophony and a greater focus on biophony

(S. Gage, pers. Comm.), which is important when considering biotic responses to typhoons. Choice of

acoustic indices was determined by previous work in this system showing that our focal indices

generally covary with biodiversity across the range of sonic conditions experienced in Okinawa (Ross

et al., 2021a). To facilitate comparisons among indices across studies, we normalised acoustic index

values before analysis, producing relative proportions by dividing $NDSI_{Bio}$ and $NDSI_{Anthro}$ each by their

site-specific maximum (Bradfer-Lawrence et al., 2020), and normalising NDSI as (NDSI + 1)/2, since it

ranges -1 and +1 and so cannot be scaled by its maximum to normalise values (Fairbrass et al., 2017).

We also used machine learning methods (see Ross et al., 2018) to identify and count

detections of three key focal bird species from our recordings. We used Kaleidoscope Pro (version

5.3.0; Wildlife Acoustics Inc., Concord, MA, USA) to train software recognisers for the large-billed

crow (*Corvus macrorhynchos*, ハシブトガラス in Japanese), the Japanese bush warbler (*Horornis*

diphone, ウグイス), and the Ryukyu scops-owl (*Otus elegans*, リュウキュウコノハズク). We chose

these species because of their reliable detectability in recordings across sites; they are generally

abundant, frequently vocalising, and are therefore unlikely to produce false negative detections

(though note that due to data volume, we did not explicitly quantify false negative rates). Our focal

species represent a range of life histories, habitat affinities, and vocal repertoires (Hamao, 2013;

Inoue et al., 2019; Itô et al., 2000; McWhirter et al., 1996; Ross et al., 2018), including a small-ranged

forest habitat specialist (*O. elegans*), and are therefore expected to vary in their sensitivity to

typhoons and land cover (see Table S1a for hypothesised typhoon effects on bird species detections).

Species detection algorithms often transfer poorly across sites as a result of site-specific differences in

background sonic conditions (Ross et al., 2018, 2021a), but we developed reliable detectors (≤15%

false positives on visual inspection) at 21 sites for *C. macrorhynchos*, 17 sites for *H. diphone*, and 7 of

the 10 forest sites for *O. elegans* (Table S2). Kaleidoscope Pro uses a supervised clustering approach

based on Hidden Markov Models to separate sound types. Local experts cross-checked automated

clustering of sound sources and reclassified sound clusters where necessary to refine species

recognisers. Owing to the volume of data used in this study, we did not calculate exact false positive

rates for species detections. Instead, we used Kaleidoscope Pro's 'distance-from-cluster-centroid'

measure to estimate identity confidence; larger distance values represent detections that are less

likely to be the target species. Filtering by distance-from-centroid then allows rapid removal of low

certainty detections. We chose a conservative distance filter of 0.5, though our results were

qualitatively similar under less conservative filters (Figure S3). For all analyses, half-hourly bird

detections were aggregated to daily detections to lessen the impact of individual singing events

during the day (see Table S3).

Analysis of acoustic indices

Before measuring the stability of acoustic indices through time, we detrended the normalised

acoustic index time series using a centred moving average with a three-day window size in the *R*

package *forecast* (version 8.14; Hyndman & Khandakar, 2008). We chose a three-day moving average

because increasing the temporal window size of the moving average function to five or seven days

produced qualitatively similar results at the expense of time series length and dampened soundscape

dynamics (Figure S4). We then measured four components of stability at each site for normalised and

detrended acoustic time series: temporal stability, resistance, recovery time, and spatial variability

(Table S3; Donohue et al., 2013). Temporal stability was calculated as 1 minus the coefficient of

variation (that is, the standard deviation divided by the mean) of the 30-day pre-typhoon period and

the 30-day post-typhoon period, separately. Resistance was the maximum absolute change between

the mean pre-typhoon baseline state and the maximum point of deviation from that state within 48

hours of the second typhoon passing (Hillebrand et al., 2018). Recovery time was 1 minus the time (in

hours) between the point of maximum deviation from baseline (from which resistance was measured)

and the point at which values returned to the pre-typhoon baseline (mean ± 95% confidence interval)

and stayed within them for 24 hours (White et al., 2020), though results were generally robust to

alternative window sizes (Figure S5). Temporal sampling regimes can affect the likelihood that

meaningful ecological patterns are returned from acoustic datasets (Bradfer-Lawrence et al., 2019;

Metcalf et al., 2021). To further validate our use of the entire 24-hr cycle in capturing ecologically

meaningful results, we showed that mean daily acoustic index values were consistently positively

related to those isolated from the dawn chorus, regardless of field site (Figure S6).

We calculated spatial variability per time point as the coefficient of variation (standard

deviation/mean) of acoustic index values for all 24 field sites (Table S3). Doing this at each time point

produced time series of spatial variability; spatial variability can change through time as the

soundscapes at different field sites become more similar (lower spatial variability) or less similar

(higher spatial variability), depending on a range of abiotic and biotic factors. Higher values of spatial

variability among sites following a disturbance may represent a greater diversity of potential

responses through asynchronous biomass fluxes within or among species, providing spatial insurance

through patch dynamics (Leibold et al., 2004; Loreau et al., 2003; Wang et al., 2021). To test for

potential land cover effects on spatial variability, we also calculated spatial variability among only

those sites characterised as either forested or developed (Figure S1); that is, we separately measured

the coefficient of variation of acoustic index values at the 10 forest sites and the 14 developed sites.

To aid comparison, stability components were normalised by their maximum (0-1) and defined such

that larger values represent greater stability (see Table S1b for hypothesised typhoon effects, and

Table S3 for detailed explanation of stability components and their interpretation).

All analyses were done in *R* (version 4.3.0, R Core Team 2023), using the packages *brms* and

segmented (Bürkner, 2017; Muggeo, 2008). We tested for interactive effects of typhoons and land

use on mean acoustic index states and temporal stability of indices, and for land use effects on

acoustic index resistance and recovery time. For these analyses, we fitted generalised linear mixed

effects models, with field site included as a random effect, using Stan (Stan development team 2020),

implemented via the *brm* function in *brms* (Bürkner, 2017). For all four response variables, the

modelled fixed effects included land use category (forest or developed) and typhoon state (pre- or

post-typhoon). Given their nature, resistance and recovery time were not modelled as a function of

typhoon impact. Default Hamiltonian Monte Carlo was used for the MCMC algorithm and all priors

were uninformative. As our response variables fell on the [0,1] scale, we used the Beta model family

with logit link. Model comparisons were made with leave-one-out cross validation (LOOIC)

implemented in *brms* calculated via Pareto-smoothed importance sampling (Vehtari et al., 2017). We

chose models with lowest LOOIC as best performing models (excepting cases where ΔLOOIC < 4.0,

where model selection favoured the model with fewer parameters), since lower LOOIC indicates

higher predictive accuracy. Four independent MCMC chains were run, each with a warmup phase of

5,000 iterations and sampling phase of 45,000 iterations. We inspected trace plots and density plots

visually for chain mixture and verified convergence using the Gelman-Rubin $\hat{R}$ < 1.01 and effective

sampling size statistics (Gelman & Hill, 2006). We also tested for spatial autocorrelation of model

residuals using the Moran's *I* test statistic for each fitted model (Gittleman & Kot, 1990). Moran's *I*

results were always non-significant (that is, we did not detect significant spatial autocorrelation in any

models), so we report results of the nonspatial models. Results of these models are presented as 95%

highest density intervals (credible intervals) of all chains' posterior parameter draws after the burn-in

period.

For models of spatial variability responses, we fitted break-point models of spatial variability

using the R package *mcp* (Lindeløv 2020). Break-point models fit segmented relationships between

predictor and response variables to determine whether the form of the relationship changes as a

function of the predictor variable. In our case, we modelled spatial variability as a function of time,

with increasing numbers of break points from zero to three breaks, allowing intercepts, but not

slopes, to vary between time segments. Break points were based on weakly informative prior

distributions centring around the passing of each typhoon. That is, for a single break point, prior

distributions centred on the mid-point between typhoons (around 1 Oct 2018), for two break points,

priors were centred on each typhoon separately (29 Sep and 6 Oct 2018, respectively), and for three

break points, we added a third point following the passing of the second typhoon (around 12 Oct

2018). For each model, we ran eight independent MCMC chains with a warmup phase of 10,000

iterations. We detrended acoustic index spatial variability time series with a periodicity of one week

using loess decomposition in the R package *stats* (R Core Team, 2023). We then ran all models on the

trend time series (that is, removing the noise and seasonal components), since this provided a

suitable trade-off between data resolution and model convergence. We compared models with

different numbers of break points using LOOIC, again selecting models with lowest LOOIC values as

best performing models unless a model with fewer breakpoints had ΔLOOIC < 4.0. Visual inspection of

the number, location, and agreement of breakpoints within best performing models then reveals

whether typhoons induced shifts in spatial variability; best performing models with strong agreement

of breakpoints near the typhoons suggest a higher likelihood that observed changes in model

intercept are driven by typhoons, and nonoverlapping 95% credible intervals between intercepts

provides quantitative evidence for a change in intercept.

Analysis of automated species detections

Given the lower temporal resolution of daily summed time series of bird species detections

(*i.e.*, one value per day rather than 48), we did not estimate resistance or recovery time for bird

species detections; such coarse grain time series would not produce meaningful resistance or

recovery values in a time series of this length (particularly when defining a recovery window size for

bird detections to be classed as recovered). Rather, we focused our analyses on the temporal stability

of bird detections for each species across the 30-day pre- and post-typhoon periods and the spatial

variability of detections per day across all sites, and across sites falling into each land use category

(forested versus developed). Note that the forest specialist *Otus elegans* was not detected in any

developed sites (Table S2), so for this species there is no data subset to compare between land cover

types. As automated species detections produced count data, we did not normalise raw values of bird

species detections.

As for acoustic indices, we tested for interactions between land use and typhoon effects on

the mean number of daily detections (mean state) and the temporal stability of daily detections. We

compared species effects by fitting a three-way interaction between species identity, land use, and

typhoon period (two levels: before versus after the typhoons). We specified *brms* models as

described previously, but with lognormal error distributions, which outperformed other error

structures based on LOOIC. To aid convergence, we additionally set weakly informative priors of

$N(0,2)$ for all predictor variables in both models, but otherwise opted for uninformative priors. We

measured spatial variability as the coefficient of variation of daily species detections across all 24 sites

(Table S3), and again separately for the 10 forest sites and the 14 developed sites. For spatial

variability, we fitted break-point models of spatial variability with intercept-only breaks, and used

LOOIC to compare between models with zero to three breaks. Weakly informative priors followed

break-point models of acoustic indices (above), and visual inspection of the number, location, and

agreement of breakpoints within best performing models revealed whether typhoons induced shifts

in spatial variability. Spatial variability values of species detections were already daily means, so we

did not further detrend time series.

Results

Acoustic index results

We found that NDSI was significantly lower at many sites after the typhoons (Figure 2a). This

overall pattern seemed not to be driven by an underlying change in the biophony component of NDSI

(Figure 2b), but rather by an increase in anthropophony following the typhoons (Figure 2c). There was

no land use or typhoon effect on the temporal stability, resistance, or recovery of NDSI, NDSI$_{Bio}$, or

NDSI$_{Anthro}$. However, in some cases we found differences in acoustic index values following the

typhoons (Table S4).

Figure 2. **Comparison of acoustic index mean state values before and after the typhoons.** Posterior distributions

represent 90,000 post-convergence MCMC draws of the change from pre- to post-typhoon periods, where

values below zero (grey) indicate a post-typhoon decline, and values above zero (blue) a post-typhoon increase

in mean state value. Non-zero-spanning credible intervals are marked with *, while circles indicate zero-

spanning credible intervals (no change based on the posterior distribution). Draws are shown per site, ordered

from most forested (top) to most developed (bottom) based on principle component axis 1 of the land use

dimensionality reduction (PCA; see Figure S1). Panels represent changes in mean state values for three acoustic

indices: the normalised difference soundscape index [NDSI] (a), biophony [NDSI$_{Bio}$] (b), and anthropophony

[NDSI$_{Anthro}$] (c).

When modelling the effects of typhoons on spatial variability of acoustic indices through

time, models with three break points, including at least one coinciding with the typhoons, were

chosen for all combinations of acoustic indices and land use (see Table S5). Following the typhoons,

the spatial variability of NDSI increased (Figure S7). Break point models produced breaks that closely

accompanied the typhoons, and this post-typhoon spatial divergence in NDSI was underlain by an

increase in biophony, but not anthropophony (Figures 3 and S8). Moreover, spatial variability in

biophony was higher in developed than forested sites before the typhoons (compare "pre-typhoon"

credible intervals in Table S5), and spatial variability among both groups of sites increased after the

typhoons (Table S5; Figures 3b and S8). Conversely, the spatial variability of anthropophony declined

following the typhoons (Figures 3c and S8), but was not different between forest and developed sites

(Table S5; Figure 3d).

Figure 3. Spatial variability of biophony [NDSI_Bio] and Anthropophony [NDSI_Anthro] **through time**. Time series of

$NDSI_{Bio}$ (a,b) and $NDSI_{Anthro}$ (c,d) spatial variability across all sites (a,c), and across forest (green) and developed

(purple) sites separately (b,d). Dashed lines delineate the pre- and post-typhoon periods.

 Automated species detection results

Species identity interacted with the typhoons, producing species-specific typhoon responses

(Table S4; Figure S9). Detections of *C. macrorhynchos* and *O. elegans* were similar preceding and

following the typhoons (Figures 4a and 4c), whereas *H. diphone* was detected less often after the

typhoons (Figures 4b and S10). We also found that, following the typhoons, species detections were

more stable (less variable) through time, regardless of the species considered (Figure 5; Table S4). We

found no effect of land use on the mean number of daily species detections or the temporal stability

of daily detections (Table S4; see Figure S9 for a map of mean daily detections before and after the

typhoons).

Figure 4. Comparison of mean daily species detections before and after the typhoons. Posterior distributions

representing 90,000 post-convergence MCMC draws of the change from pre- to post-typhoon periods, where

values below zero (grey) indicate a post-typhoon decline, and values above zero (blue) a post-typhoon increase

in mean daily species vocalisation detections. Non-zero-spanning credible intervals are marked with *, while

circles indicate zero-spanning credible intervals (no change based on the posterior distribution). Draws are

shown per site, ordered from most forested (top) to most developed (bottom) based on principle component

axis 1 of the land use dimensionality reduction (PCA; see Figure S1). Panels represent changes in mean daily

species detections for our three focal species: *Corvus macrorhynchos* (a), *Horornis diphone* (b), and *Otus elegans*

(c). Inferred posterior draws (automatically computed through the site random effect term) extrapolated to

field sites where species were not present (Table S2) are shown as faded distributions.

Figure 5. Comparison of temporal stability of species detections before and after the typhoons. Posterior

distributions represent 90,000 post-convergence MCMC draws of the change from pre- to post-typhoon

periods, where values below zero (grey) indicate a post-typhoon decline, and values above zero (blue) a post-

typhoon increase in the temporal stability of automated species vocalisation detections (all species). Non-zero-

spanning credible intervals are marked with *, while circles indicate zero-spanning credible intervals (no change

based on the posterior distribution). Draws are shown per site, ordered from most forested (top) to most

developed (bottom) based on principle component axis 1 of the land use dimensionality reduction (Figure S1).

See Figure S11 for posterior draws of individual species.

When modelling the effects of typhoons on spatial variability of bird detections through time,

we found species-specific typhoon effects, with detections of *Corvus macrorhynchos* and *Otus elegans*

not showing clear typhoon-related changes in spatial variability (Table S5; Figures 6a and S12).

However, across forested sites, detections of *C. macrorhynchos* showed an increase in spatial

variability coincident with the typhoons (as indicated by the selected break-point model, Figure S12b).

The spatial variability of *Otus elegans* detections was higher than of the other bird species (Table S5;

Figure 6a). In contrast to the other species, detections of *Horornis diphone* became more variable

around the onset of the typhoons across all sites (Figure 6a) and across forested sites (Figure 6b), as

supported by break-point models (Figure S12d and S12e). However, across developed sites, there was

no clear association between the typhoons and any possible break points in *H. diphone* spatial

variability (Table S5; Figures 6b and S12f).

Figure 6. **Spatial variability of automated daily species detections through time**. Time series of spatial variability

(coefficient of variation) of daily species detections across all sites (a) for each of *Corvus macrorhynchos* (grey),

Horornis diphone (blue), and *Otus elegans* (yellow), and (b) measured separately across the 10 forest (green)

and 14 developed (purple) sites for each species. Dashed lines delineate the pre- and post-typhoon periods.

Discussion

This study leverages high-resolution acoustic monitoring data from an island-wide sensor array to

record ecological responses to extreme weather events in the form of two large typhoons. We found

no land use effects on most dimensions of stability measured. However, we found that spatial

variability of biophony was greater among developed sites compared with forested ones before the

typhoons, and that the spatial variability of both biophony ($NDSI_{Bio}$) and the normalised difference

soundscape index (NDSI) increased significantly following the typhoons. Contrary to the expected

typhoon-induced homogenisation of soundscapes, our results indicate that the typhoons elicited

divergent ecological responses among Okinawa's soniferous animal communities. In support of this,

we found that detections of two of our three focal bird species, *Corvus macrorhynchos* and *Horornis*

diphone, were more spatially variable after the typhoons, in contrast to the expected post-typhoon

homogenisation through mortality or distributional shifts. This post-typhoon increase in spatial

variability of species detections was present among forested but not developed sites, suggesting

perhaps that land use and habitat change can hinder the reactive capacity of ecological communities

and their associated soundscapes. That our species-level results matched those of acoustic indices

further validates our use of acoustic indices to assess typhoon impacts in Okinawa.

We found spatial divergence of NDSI and biophony, but not anthropophony or species

detections, in developed urban and agricultural sites. This disconnect between results from the

biophony index versus automated species detections in developed sites could reflect differences in

local biodiversity in developed sites, or perhaps that differences in ambient sonic conditions affected

our ability to accurately detect species responses (Ross et al., 2021a). We did not directly measure

local biodiversity in this study, instead estimating the activity of some key focal bird species using

automated species detections. However, previous work in this system provided evidence for a loss of

rare and endemic birds under land use development, producing communities that are a nested subset

of forest bird communities in Okinawa's developed south (Ross et al., 2018). The species surveyed

here did not exhibit responses that diverged in space in developed sites following the typhoons, as

might have been expected based on biophony results in developed sites. This suggests that the spatial

divergence in the biotic component of soundscapes recorded here may be better explained by other

species (of birds or other taxa) not targeted in this study. Future work expanding on these analyses to

provide a more holistic view of the Okinawan biota should therefore prove fruitful for identifying

individual species contributions to typhoon responses. If our biophony results are indeed a product of

biotic responses to typhoons as would be expected from theory (Kasten et al., 2012), and as

supported by our finding that spatial variability of both *C. macrorhynchos* and *H. diphone* increased

among forested sites, then a post-typhoon increase in spatial variability may reflect changes to

species' patchiness. For example, Willig and Camilo (1991) described an increase in spatial patchiness

of the snail *Caracolus caracol* following Hurricane Hugo in Puerto Rico, caused by a thinning of

populations due to post-hurricane mortality.

Soundscape composition after the typhoons saw an increase in anthropophony as anticipated

(Table S1b), but not a decline in biophony as might be expected were animal populations impacted

negatively by typhoon disturbance (*e.g*., Cely, 1991; Pavelka et al., 2007). In contrast, the observed

post-typhoon increase in spatial variability in NDSI was driven by biophony rather than anthrophony.

This suggests that, while biophony may not have been affected substantially by typhoon disturbance

at the individual site level, variation in biotic responses at larger scales across field sites nonetheless

manifested as changes in the spatial variability of biophony after the typhoons. This is likely driven, at

least in part, by the observed post-typhoon spatial divergences in species vocalisations. That we did

not detect particularly strong site-level typhoon impacts, but rather saw spatial divergence in

ecological responses to typhoons across multiple sites, underscores the necessity and benefits of

monitoring at scale. Multi-site acoustic sensor arrays such as ours thus provide opportunity to

monitor both local and regional biodiversity change, in turn providing critical new insight for

conservation management (Roe et al., 2021; Sethi et al., 2020a; Van Parijs et al., 2015). The observed

post-typhoon increase in anthropophony on the other hand, likely reflects increased human activity

at developed sites as efforts to survey and repair typhoon damage got underway. Conversely, the

expected post-typhoon increase in anthropophony observed at a few forest sites may reflect changes

in sound propagation driven by vegetative structural damage and thinning of previously dense

habitats, as is often documented following large storms (Abbas et al., 2020; Elliott & Nino, 1960). We

did not measure habitat structure directly, and so the causes of increases to anthropophony following

typhoons Trami and Kong-Rey cannot be demonstrated empirically. We did, however, observe

significant damage and alterations to habitat structure at our forested field sites (T. Yoshida & M.

Yoshimura, pers. obs.). Automated bird species detections were, conversely, more stable through

time after the typhoons, suggesting disturbance may affect the consistency of species vocalisations in

Okinawa (see also Fraterrigo & Rusak, 2008), as might be expected if rare or infrequent vocalisers are

lost from soundscapes (Bradfer-Lawrence et al., 2020; Table S1b), or perhaps that typhoon-induced

changes to habitat structure allow vocalisations to travel further without attenuation, and hence be

detected more reliably by our sensors.

The focal bird species considered here differed in their responses to typhoons. Automated

vocalisation detections of the Japanese bush warbler (*Horornis diphone*) declined after the typhoons,

while, as expected, those of the large-billed crow (*Corvus macrorhynchos*) did not. Contrary to

expectation (Table S1a), however, detections of the Ryukyu scops owl (*Otus elegans*) were also

unaffected. Given that acoustic surveys cannot differentiate between cases where a species is not

producing sound and those where that species is not present (Toth et al., 2022), we cannot say with

certainty that *H. diphone* populations declined following the typhoons. Regardless, our detected post-

typhoon declines in *H. diphone* vocalisations—either through behavioural changes, distributional

shifts, or local mortality—were consistent across >80% of the field sites in which this species was

detected. We expected that insectivores, such as *H. diphone*, may not be adversely affected by the

typhoons (excepting effects of direct mortality), but rather that structural changes to habitat may

increase prey availability in canopy gaps (Cely, 1991; Seki, 2005). However, for *H. diphone* we

observed the opposite pattern. Azuma *et al*. (1997) documented considerable effects of undergrowth

removal on invertebrate communities in Okinawa. Given that *H. diphone* relies on undergrowth and

bushes for foraging (Haneda & Okabe, 1970) and typhoon disturbance has the potential to alter the

structure of this habitat (Abbas et al., 2020; Elliott & Nino, 1960), the observed decline in *H. diphone*

detections following the typhoon may therefore be due to bottom-up changes to the invertebrate

communities on which *H. diphone* feeds following typhoon damage to the undergrowth (Azuma et al.,

1997). Moreover, *H. diphone* is a small-bodied species and may thus be especially vulnerable to

weather-related mortality (Brown & Brown, 1998). In contrast, and contrary to expectation (Table

S1a), the forest specialist *O. elegans* was not detected less frequently after the typhoons, suggesting

that its habitat or foraging were unaffected by the typhoons, or perhaps that cavity nesting reduced

typhoon impact by reducing exposure to the extreme weather (Inoue et al., 2019). Overall, species

detections were more stable through time after the typhoons, though acoustic index temporal

stability was not affected by typhoons. Given that soundscapes can be more temporally stable when

human activity is restricted (Ross, 2022), the higher post-typhoon temporal stability of species

detections observed here may relate to altered human space use across Okinawa following the

typhoons.

Species-specific responses to disturbance may more generally reflect differences in life

history and other functional response traits (Suding et al., 2008), which can be useful predictors of

community dynamics, disassembly, and stability in birds (*e.g.*, Ausprey et al., 2022; Burivalova et al.,

2015; Hordley et al., 2021; Zhang et al., 2016). Similarly, different typhoon responses of vocalisation

detections among field sites may reflect differences in underlying vegetative changes as determined

by plant functional response traits. For example, Craven *et al*. (2016) found that functionally diverse

Canadian forests were dominated by trees with response traits that promoted resilience to recurrent

anthropogenic disturbance through rapid regrowth, rather than resistance to projected climate

change through drought or flood tolerance. The response traits of plants may then, in turn, determine

the structural habitat change experienced by birds and other vocalising animals (*e.g.*, Abbas et al.,

2020), as well as directly influencing sound propagation (Morton, 1975).

Though we and others have demonstrated the capacity for passive acoustic monitoring

methods to capture unpredictable extreme weather events (Gottesman et al., 2021; Simmons et al.,

2021), such methods are often limited in their ability to accurately reflect biodiversity patterns. A

recent meta-analysis reported a generally positive link between acoustic indices and biodiversity

(Alcocer et al., 2022), but one with diminishing effect sizes over time, as studies increasingly forego

appropriate validation and as their designs incorporate yet wider varieties of non-target sounds,

which can hinder the interpretability of those acoustic indices aiming to reflect biodiversity (Ross et

al., 2021a). The joint use of acoustic indices and automated species detections provides two separate

lines of evidence for typhoon-induced soundscape change; such species and soundscape methods are

still rarely used in combination despite their clear potential to provide complementary information on

ecological dynamics (*e.g.*, Ferreira et al., 2018; Ross et al., 2018). Indeed, our complementary results

linking post-typhoon spatial divergence of biophony to that of species detections underscore the

efficacy of combing both acoustic indices and individual species recognition models. That said,

building reliable vocalisation recognition algorithms remains a challenge, particularly when aiming for

transferability to different habitats or seasons, which provide a range of non-target sounds beyond

those on which algorithms may have been trained. Increasing application of deep learning to such

problems will likely help provide a solution (*e.g.*, Sethi et al., 2020b) as will continued efforts to build

labelled sound libraries from which automated species detection algorithms can be trained

(Deichmann et al., 2018).

Soundscape dynamics are frequently characterised by strong seasonal cycles (*e.g.*, Vokurková

et al., 2018). This presents a challenge when attempting to disentangle disturbance responses from

seasonal soundscape change. For example, our focal species differ in their seasonality and phenology,

meaning that natural phenological differences may be in part responsible for the differences in

species' typhoon responses we observed here. That said, *Horornis diphone* vocalisation detections

declined sharply and concomitantly with the typhoons (Figure S10). This suggests the observed post-

typhoon decline was most likely a result of the typhoons themselves, rather than a seasonal decline,

which we would expect to be much more gradual in nature. Our moving average detrend aimed to

remove as much seasonal signal as possible, though more sophisticated approaches to

deseasonalisation exist such as, for example, wavelet decomposition. Such approaches typically

require longer time series than ours, however, with clear periodic components comprising less than a

quarter of the total time series length (Cazelles et al., 2008), and so wavelet analysis was not

appropriate for our study. As the OKEON Churamori Project continues to collect soundscape

recordings from across Okinawa, advanced time series models and decomposition techniques will be

needed to process yet longer time series and periodic signals.

Our *k*-means clustering approach to distinguish field sites by their dominant land use

identified an optimal split of two clusters, separating primarily forested sites from those dominated

by developed urban or agricultural land use. However, these developed land uses can act on

ecological dynamics and stability in different ways. For example, Olivier *et al.* (2020) used citizen

science data from across France to show that agricultural intensification directly affected population,

and, in turn, community stability of birds, whereas urbanisation acted only indirectly on community

stability through changes to diversity and population asynchrony. Our study design, which was based

on unsupervised (*k*-means) site clustering by dominant land use, consequently did not allow us to

directly compare urban and agricultural field sites, despite their potential for contrasting effects on

ecological stability. Moreover, we classified sites based on land cover from a 2015 Landsat 8 image,

while habitat structure and land use are dynamic features. As such, we could not directly capture

legacy effects of historic land use on ecological communities in this study, including both

intensification—the dominant trend in Okinawa (Takeuchi et al., 1981)—and abandonment, which

can each act as driving forces structuring biodiversity (Daskalova et al., 2020; Daskalova & Kamp

2023).

Conclusion

Our study tested the capacity for land use and climate change in the form of extreme weather events

to jointly shape ecological stability. Using passive acoustic monitoring data from a landscape-scale

sensor network across Okinawa Island, we found that land use rarely modified ecological responses to

typhoons. However, soundscapes diverged across the landscape following the typhoons, contrary to

the expected typhoon-induced soundscape homogenisation. Biophony in developed sites was initially

more variable across sites, but this post-typhoon spatial divergence occurred among both forested

and developed urban and agricultural field sites. However, species detections diverged in space

among forested but not developed sites following the typhoons, suggesting that forest sites exhibited

a wider variety of pathways through which bird assemblages could respond to typhoon disturbance.

That is, land use intensification may produce ecological communities that are more homogeneous in

how they respond to disturbance (Vogel et al., 2019), while forest sites harbour communities with

greater potential for collective resilience to future disturbance through patch dynamics and rescue

effects among different local forest communities (Leibold et al., 2004). Such spatial insurance effects

have the potential to contribute to landscape-scale stability and spatial portfolio effects by affecting

population and community asynchrony (Loreau et al., 2003; Wang et al., 2021).

This study draws on prior knowledge of Okinawan biodiversity (Inoue et al., 2019; Itô et al.,

2000; McWhirter et al., 1996), the performance of passive acoustic methods in this system (Ross et

al., 2018, 2021a), and the characteristics of typhoons and land use intensification across Okinawa

Island (Elliott & Nino, 1960; Takeuchi et al., 1981), to explore fundamental ideas concerning

disturbance and stability (Ross et al., 2023). Such baseline data provides a critical backdrop against

which our results stand, allowing us to infer species and soundscape responses to the joint threats of

climate change and land use intensification from acoustic recordings of typhoons (Altwegg et al.,

2017). As longer and higher-resolution acoustic data are amassed through multi-site acoustic sensor

arrays (*e.g.*, Roe et al., 2021; Sethi et al., 2020a; Van Parijs et al., 2015), the utility of passive acoustic

monitoring to document ecological responses to extreme weather events across the globe will

become ever clearer, particularly in light of the increasing frequency and destructive potential of

extreme events in the Anthropocene.

653
Acknowledgements

Many individuals in the OKEON Churamori Project contributed to data collection, site maintenance, and

community outreach; we especially thank Masako Ogasawara, Mayuko Suwabe, Shinji Iriyama, Toshihiro

Kinjo, Izumi Maehira, Yuko Matsudo, Seiichiro Nakagawa, Shoko Suzuki, Takumi Uchima, Madoka Oguro,

Dan Warren, and Chisa Oshiro. We are grateful for the help and support provided by the Scientific

Computing and Data Analysis section of the Research Support Division at OIST. Our sincere thanks go to

the landowners, museums, local governments, and schools that host the OKEON Churamori Project field

sampling sites, and to the people of Okinawa. We also thank Marina Lašić and Rob McHenry for their

contributions to building automated species recognisers, Yvonne Buckley, Anne Magurran, and Jamie Kass

for helpful discussion, and Vito Muggeo for advice on break-point modelling in R. This work was supported

by subsidy funding to OIST, and an Irish Research Council Postgraduate Scholarship [GOIPG/2018/3023]

and Canon Foundation in Europe 2021 Research Fellowship awarded to S.R.P-J.R.

666
667 ## Conflict of Interest

668 The authors declare no conflict of interest.

669
670 ## Data availability statement

671 The data supporting the findings of this study and all R code are available via the Zenodo digital repository

672 as Ross, S. R. P.-J., Friedman, N. R., Dudley, K. L., Yoshimura, M., Yoshida, T., Economo, E. P., Armitage, D.

673 W., & Donohue, I. (2023). Data from: Divergent ecological responses to typhoon disturbance revealed via

674 landscape-scale acoustic monitoring (v0.3-review2). *Zenodo* https://doi.org/10.5281/zenodo.8133339

675
ORCID

Samuel R.P-J. Ross: https://orcid.org/0000-0001-9402-9119

Nicholas R. Friedman: https://orcid.org/0000-0002-0533-6801

Kenneth L. Dudley: https://orcid.org/0000-0003-3594-0724

Takuma Yoshida: https://orcid.org/0000-0002-2172-9488

Masashi Yoshimura: https://orcid.org/0000-0003-1468-5541

Evan P. Economo: https://orcid.org/0000-0001-7402-0432

David W. Armitage: https://orcid.org/0000-0002-5677-0501

Ian Donohue: https://orcid.org/0000-0002-4698-6448

685
References

Abbas, S., Nichol, J. E., Fischer, G. A., Wong, M. S., & Irteza, S. M. (2020). Impact assessment of a super-typhoon

on Hong Kong's secondary vegetation and recommendations for restoration of resilience in the forest

succession. *Agricultural and Forest Meteorology*, *280*, 107784.

https://doi.org/10.1016/j.agrformet.2019.107784

Alcocer, I., Lima, H., Sugai, L. S. M., & Llusia, D. (2022). Acoustic indices as proxies for biodiversity: A meta-

analysis. *Biological Reviews*. https://doi.org/10.1111/brv.12890

Altwegg, R., Visser, V., Bailey, L. D., & Erni, B. (2017). Learning from single extreme events. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *372*(1723), 20160141.

https://doi.org/10.1098/rstb.2016.0141

Ares, Á., Brisbin, M. M., Sato, K. N., Martín, J. P., Iinuma, Y., & Mitarai, S. (2020). Extreme storms cause rapid but

short-lived shifts in nearshore subtropical bacterial communities. *Environmental Microbiology*, *22*(11),

4571–4588. https://doi.org/10.1111/1462-2920.15178

Ausprey, I. J., Newell, F. L., & Robinson, S. K. (2022). Functional response traits and altered ecological niches

drive the disassembly of cloud forest bird communities in tropical montane countrysides. *Journal of*

Animal Ecology, *91*(11), 2314–2328. https://doi.org/10.1111/1365-2656.13816

Azuma, S., Sasaki, K., & Itô, Y. (1997). Effects of undergrowth removal on the species diversity of insects in

natural forests of Okinawa Hontô. *Pacific Conservation Biology*, *3*(2), 156–160.

https://doi.org/10.1071/pc970156

Baert, J. M., De Laender, F., Sabbe, K., & Janssen, C. R. (2016). Biodiversity increases functional and

compositional resistance, but decreases resilience in phytoplankton communities. *Ecology*, *97*(12),

3433–3440. https://doi.org/10.1002/ecy.1601

Bhatia, K. T., Vecchi, G. A., Knutson, T. R., Murakami, H., Kossin, J., Dixon, K. W., & Whitlock, C. E. (2019). Recent

increases in tropical cyclone intensification rates. *Nature Communications*, *10*(1), Article 1.

https://doi.org/10.1038/s41467-019-08471-z

Boyd, A. D., Gowans, S., Mann, D. A., & Simard, P. (2021). Tropical Storm Debby: Soundscape and fish sound

production in Tampa Bay and the Gulf of Mexico. *PLOS ONE*, *16*(7), e0254614–e0254614.

https://doi.org/10.1371/JOURNAL.PONE.0254614

Bradfer-Lawrence, T., Bunnefeld, N., Gardner, N., Willis, S. G., & Dent, D. H. (2020). Rapid assessment of avian

species richness and abundance using acoustic indices. *Ecological Indicators*, *115*(April), 106400–

106400. https://doi.org/10.1016/j.ecolind.2020.106400

Bradfer-Lawrence, T., Gardner, N., Bunnefeld, L., Bunnefeld, N., Willis, S. G., & Dent, D. H. (2019). Guidelines for

the use of acoustic indices in environmental research. *Methods in Ecology and Evolution*, *10*(10), 1796-

1807. https://doi.org/10.1111/2041-210X.13254

Brown, C. R., & Brown, M. B. (1998). Intense natural selection on body size and wing and tail asymmetry in cliff

swallows during severe weather. *Evolution, 52(5)*, 1461-1475. https://doi.org/10.1111/j.1558-

5646.1998.tb02027.x

Burivalova, Z., Game, E. T., & Butler, R. A. (2019). The sound of a tropical forest. *Science*, *363*(6422), 28–29.

https://doi.org/10.1126/science.aav1902

Burivalova, Z., Lee, T. M., Giam, X., Sekercioglu, Ç. H., Wilcove, D. S., & Koh, L. P. (2015). Avian responses to

selective logging shaped by species traits and logging practices. *Proceedings of the Royal Society B:*

Biological Sciences, *282*(1808). https://doi.org/10.1098/rspb.2015.0164

Burivalova, Z., Şekercioğlu, Ç. H., & Koh, L. P. (2014). Thresholds of logging intensity to maintain tropical forest

biodiversity. *Current Biology*, *24*(16), 1893–1898. https://doi.org/10.1016/j.cub.2014.06.065

Bürkner, P.-C. (2017). Brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical*

Software, *80*, 1–28.

Butsic, V., Lewis, D. J., Radeloff, V. C., Baumann, M., & Kuemmerle, T. (2017). Quasi-experimental methods

enable stronger inferences from observational data in ecology. *Basic and Applied Ecology*, *19*, 1–10.

https://doi.org/10.1016/j.baae.2017.01.005

Cazelles, B., Chavez, M., Berteaux, D., Ménard, F., Vik, J. O., Jenouvrier, S., & Stenseth, N. C. (2008). Wavelet

analysis of ecological time series. *Oecologia*, *156*(2), 287–304. https://doi.org/10.1007/s00442-008-

0993-2

Cely, J. E. (1991). Wildlife Effects of Hurricane Hugo. *Journal of Coastal Research*, 319–326.

Chevalier, M., Lindström, Å., Pärt, T., & Knape, J. (2019). Changes in forest bird abundance, community structure

and composition following a hurricane in Sweden. *Ecography*, *42*(11), 1862–1873.

https://doi.org/10.1111/ecog.04578

Clark, A. T., Arnoldi, J.-F., Zelnik, Y. R., Barabas, G., Hodapp, D., Karakoç, C., … Harpole, S. (2021). General

statistical scaling laws for stability in ecological systems. *Ecology Letters 24*(7), 1474–1486.

https://doi.org/10.1111/ele.13760

Clough, Y., Putra, D. D., Pitopang, R., & Tscharntke, T. (2009). Local and landscape factors determine functional

bird diversity in Indonesian cacao agroforestry. *Biological Conservation, 142*(5), 1032-1041.

https://doi.org/10.1016/j.biocon.2008.12.027

Cohen, J. M., Fink, D., & Zuckerberg, B. (2021). Extreme winter weather disrupts bird occurrence and abundance

patterns at geographic scales. *Ecography*, 1–13. https://doi.org/10.1111/ecog.05495

Craven, D., Filotas, E., Angers, V. A., & Messier, C. (2016). Evaluating resilience of tree communities in

fragmented landscapes: Linking functional response diversity with landscape connectivity. *Diversity and*

Distributions, *22*(5), 505–518. https://doi.org/10.1111/ddi.12423

Daskalova, G.N., & Kamp, J. (2023). Abandoning land transforms biodiversity. *Science, 380(6645)*, 581-583.

https://doi.org/10.1126/science.adf1099

Daskalova, G. N., Myers-Smith, I. H., Bjorkman, A. D., Blowes, S. A., Supp, S. R., Magurran, A. E., & Dornelas, M.

(2020). Landscape-scale forest loss as a catalyst of population and biodiversity change. *Science*,

368(June), 1341–1347. https://doi.org/10.1126/science.aba1289

Deichmann, J. L., Acevedo-Charry, O., Barclay, L., Burivalova, Z., Campos-Cerqueira, M., D'Horta, F., … Aide, T. M.

(2018). It's time to listen: There is much to be learned from the sounds of tropical ecosystems.

Biotropica, *50*(5), 713–718. https://doi.org/10.1111/btp.12593

Deichmann, J. L., Hernández-Serna, A., Delgado C., J. A., Campos-Cerqueira, M., & Aide, T. M. (2017).

Soundscape analysis and acoustic monitoring document impacts of natural gas exploration on

biodiversity in a tropical forest. *Ecological Indicators*, *74*, 39–48.

https://doi.org/10.1016/j.ecolind.2016.11.002

Depraetere, M., Pavoine, S., Jiguet, F., Gasc, A., Duvail, S., & Sueur, J. (2012). Monitoring animal diversity using

acoustic indices: Implementation in a temperate woodland. *Ecological Indicators*, *13*(1), 46–54.

https://doi.org/10.1016/j.ecolind.2011.05.006

Donihue, C. M., Herrel, A., Fabre, A. C., Kamath, A., Geneva, A. J., Schoener, T. W., Kolbe, J. J., & Losos, J. B.

(2018). Hurricane-induced selection on the morphology of an island lizard. *Nature*, *560*(7716), 88–91.

https://doi.org/10.1038/s41586-018-0352-3

Donohue, I., Petchey, O. L., Montoya, J. M., Jackson, A. L., Mcnally, L., Viana, M., … Emmerson, M. C. (2013). On

the dimensionality of ecological stability. *Ecology Letters*, *16*(4), 421–429.

https://doi.org/10.1111/ele.12086

Elliott, J. C., & Nino, Y. (1960). Okinawa's Dry Typhoons. *American Midland Naturalist*, *63*(1), 211–211.

https://doi.org/10.2307/2422941

Emanuel, K. (2005). Increasing destructiveness of tropical cyclones over the past 30 years. *Nature*, *436*(7051),

Article 7051. https://doi.org/10.1038/nature03906

Everham, E. M., & Brokaw, N. V. L. (1996). Forest damage and recovery from catastrophic wind. *The Botanical

Review 1996 62:2*, *62*(2), 113–185. https://doi.org/10.1007/BF02857920

Fairbrass, A. J., Rennett, P., Williams, C., Titheridge, H., & Jones, K. E. (2017). Biases of acoustic indices

measuring biodiversity in urban areas. *Ecological Indicators*, *83*(February), 169–177.

https://doi.org/10.1016/j.ecolind.2017.07.064

Ferreira, L. M., Oliveira, E. G., Lopes, L. C., Brito, M. R., Baumgarten, J., Rodrigues, F. H., & Sousa-Lima, R. S.

(2018). What do insects, anurans, birds, and mammals have to say about soundscape indices in a

tropical savanna. *Journal of Ecoacoustics*, *2*, PVH6YZ-PVH6YZ. https://doi.org/10.22261/JEA.PVH6YZ

Fraterrigo, J. M., & Rusak, J. A. (2008). Disturbance-driven changes in the variability of ecological patterns and

processes. *Ecology Letters*, *11*(7), 756–770. https://doi.org/10.1111/j.1461-0248.2008.01191.x

Gardner, L. R., Michener, W. K., Blood, E. R., Williams, T. M., Lipscomb, D. J., & Jefferson, W. H. (1991).

Ecological Impact of Hurricane Hugo—Salinization of a Coastal Forest on JSTOR. *Journal of Coastal

Research*, *8*, 301–317.

Gasc, A., Francomano, D., Dunning, J. B., & Pijanowski, B. C. (2017). Future directions for soundscape ecology:

The importance of ornithological contributions. *The Auk*, *134*(1), 215–228.

https://doi.org/10.1642/AUK-16-124.1

Gasc, A., Gottesman, B. L., Francomano, D., Jung, J., Durham, M., Mateljak, J., & Pijanowski, B. C. (2018).

Soundscapes reveal disturbance impacts: Biophonic response to wildfire in the Sonoran Desert Sky

Islands. *Landscape Ecology*, *33*, 1399–1415. https://doi.org/10.1007/s10980-018-0675-3

Gelman, A., & Hill, J. (2006). *Data analysis using regression and multilevel/hierarchical models*. Cambridge

university press.

Gibb, R., Browning, E., Glover-Kapfer, P., & Jones, K. E. (2019). Emerging opportunities and challenges for

passive acoustics in ecological assessment and monitoring. *Methods in Ecology and Evolution*.

https://doi.org/10.1111/2041-210X.13101

Gibson, L., Lee, T. M., Koh, L. P., Brook, B. W., Gardner, T. a., Barlow, J., … Sodhi, N. S. (2011). Primary forests are

irreplaceable for sustaining tropical biodiversity. *Nature*, *478*(7369), 378–381.

https://doi.org/10.1038/nature10425

Gittleman, J. L., & Kot, M. (1990). Adaptation: Statistics and a null model for estimating phylogenetic effects.

Systematic Zoology, *39*(3), 227–241.

Gottesman, B. L., Olson, J. C., Yang, S., Acevedo-Charry, O., Francomano, D., Martinez, F. A., … Pijanowski, B. C.

(2021). What does resilience sound like? Coral reef and dry forest acoustic communities respond

differently to Hurricane Maria. *Ecological Indicators*, *126*, 107635.

https://doi.org/10.1016/j.ecolind.2021.107635

Hamao, S. (2013). Acoustic structure of songs in island populations of the Japanese bush warbler, Cettia

diphone, in relation to sexual selection. *Journal of Ethology*, *31*(1), 9–15.

https://doi.org/10.1007/s10164-012-0341-1

Haneda, K., & Okabe, T. (1970). The life history of Cettia diphone 1. Breeding ecology. *Journal of the Yamashina*

Institute for Ornithology, *6*(1–2), 131–140.

Harris, S. A., Shears, N. T., & Radford, C. A. (2016). Ecoacoustic indices as proxies for biodiversity on temperate

reefs. *Methods in Ecology and Evolution*, *7*(6), 713–724. https://doi.org/10.1111/2041-210X.12527

Hillebrand, H., Langenheder, S., Lebret, K., Lindström, E., Östman, Ö., & Striebel, M. (2018). Decomposing

multiple dimensions of stability in global change experiments. *Ecology Letters*, *21*(1), 21–30.

https://doi.org/10.1111/ele.12867

Hordley, L. A., Gillings, S., Petchey, O. L., Tobias, J. A., & Oliver, T. H. (2021). Diversity of response and effect

traits provides complementary information about avian community dynamics linked to ecological

function. *Functional Ecology*, *35*(9), 1938–1950. https://doi.org/10.1111/1365-2435.13865

Hyndman, R. J., & Khandakar, Y. (2008). Automatic Time Series Forecasting: The forecast Package for R. *Journal*

of Statistical Software, *27*(1), 1–22. https://doi.org/10.18637/JSS.V027.I03

Inoue, T., Matsumoto, M., Yoshida, T., & Washitani, I. (2019). Spatial patterns of the Ryukyu Scops Owl's Otus

elegans breeding success and forest landscape factors on Amami-Ōshima island. *Japanese Journal of*

Ornithology, *68*(1), 19–28. https://doi.org/10.3838/JJO.68.19

Itô, Y., Miyagi, K., & Ota, H. (2000). Imminent extinction crisis among the endemic species of the forests of

Yanbaru, Okinawa, Japan. *Oryx*, *34*(4), 305–316. https://doi.org/10.1046/J.1365-3008.2000.00136.X

Japan Meteorological Agency (2018). *RSMC Tokyo-Typhoon Center*. International Number ID 1824 (Typhoon

Trami) and 1825 (Typhoon Kong-Rey). https://www.jma.go.jp/jma/jma-eng/jma-center/rsmc-hp-pub-

eg/bstve_2018_m.html [Accessed: 27 July 2020].

Kasten, E. P., Gage, S. H., Fox, J., & Joo, W. (2012). The remote environmental assessment laboratory's acoustic

library: An archive for studying soundscape ecology. *Ecological Informatics*, *12*, 50–67.

https://doi.org/10.1016/j.ecoinf.2012.08.001

Kéfi, S., Domínguez-García, V., Donohue, I., Fontaine, C., Thébault, E., & Dakos, V. (2019). Advancing our

understanding of ecological stability. *Ecology Letters*, ele.13340-ele.13340.

https://doi.org/10.1111/ele.13340

Keitt, T. H., & Abelson, E. S. (2021). Ecology in the Age of Automation. *Science*, *373*(6557), 858–859.

Kerr, A. M. (2000). Defoliation of an island (Guam, Mariana Archipelago, Western Pacific Ocean) following a

saltspray-laden "dry" typhoon. *Journal of Tropical Ecology*, *16*(6), 895–901.

https://doi.org/10.1017/S0266467400001796

Kossin, J. P., Knapp, K. R., Olander, T. L., & Velden, C. S. (2020). Global increase in major tropical cyclone

exceedance probability over the past 40 years. *Proceedings of the National Academy of Sciences*, *In*

review. https://doi.org/10.1073/pnas.1920849117

Laurance, W. F. (1998). A crisis in the making: Responses of Amazonian forests to land use and climate change.

Trends in Ecology & Evolution, *13*(10), 411–415. https://doi.org/10.1016/S0169-5347(98)01433-5

Leibold, M. A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J. M., Hoopes, M. F., … Gonzalez, A. (2004).

The metacommunity concept: A framework for multi-scale community ecology. *Ecology Letters*, *7*(7),

601–613. https://doi.org/10.1111/j.1461-0248.2004.00608.x

Li, L., & Chakraborty, P. (2020). Slower decay of landfalling hurricanes in a warming world. *Nature*, *587*(7833),

230–234. https://doi.org/10.1038/s41586-020-2867-7

Lin, T. C., Hogan, J. A., & Chang, C. T. (2020). Tropical Cyclone Ecology: A Scale-Link Perspective. *Trends in*

Ecology and Evolution, *xx*(xx), 0–10. https://doi.org/10.1016/j.tree.2020.02.012

Lindeløv, J. (2020). mcp: An R Package for Regression With Multiple Change Points. *OSF Preprints*

https://doi.org/10.31219/osf.io/fzqxv

Locascio, J. V., & Mann, D. A. (2005). Effects of Hurricane Charley on fish chorusing. *Biology Letters*, *1*(3), 362–

365. https://doi.org/10.1098/rsbl.2005.0309

Lomolino, M. V., Pijanowski, B. C., & Gasc, A. (2015). The silence of biogeography. *Journal of Biogeography*,

42(7), 1187–1196. https://doi.org/10.1111/jbi.12525

Loreau, M., Mouquet, N., Gonzalez, A., & Mooney, H. A. (2003). *Biodiversity as spatial insurance in*

heterogeneous landscapes. www.pnas.orgcgidoi10.1073pnas.2235465100

McWhirter, D. W., Ikenaga, H., Iozawa, H., Shoyama, M., & Takehara, K. (1996). A check-list of the birds of

Okinawa Prefecture with notes on recent status including hypothetical records. *Bulletin of Okinawa*

Prefectural Museum, *22*, 33–152.

Metcalf, O. C., Barlow, J., Devenish, C., Marsden, S., Berenguer, E., & Lees, A. C. (2021). Acoustic indices perform

better when applied at ecologically meaningful time and frequency scales. *Methods in Ecology and*

Evolution, 12(3), 421-431. https://doi.org/10.1111/2041-210X.13521

Mori, A. S., Furukawa, T., & Sasaki, T. (2013). Response diversity determines the resilience of ecosystems to

environmental change. *Biological reviews, 88*, 349-364.

Morimoto, J., Aiba, M., Furukawa, F., Mishima, Y., Yoshimura, N., Nayak, S., … Nakamura, F. (2021). Risk

assessment of forest disturbance by typhoons with heavy precipitation in northern Japan. *Forest*

Ecology and Management, *479*, 118521. https://doi.org/10.1016/j.foreco.2020.118521

Morton, E. S. (1975). Ecological Sources of Selection on Avian Sounds. *The American Naturalist*, *109*(965), 17–

34. https://doi.org/10.1086/282971

Nimmo, D. G., Haslem, A., Radford, J. Q., Hall, M., & Bennett, A. F. (2016). Riparian tree cover enhances the

resistance and stability of woodland bird communities during an extreme climatic event. *Journal of*

Applied Ecology, *53*(2), 449–458. https://doi.org/10.1111/1365-2664.12535

Olivier, T., Thébault, E., Elias, M., Fontaine, B., & Fontaine, C. (2020). Urbanization and agricultural

intensification destabilize animal communities differently than diversity loss. *Nature Communications*,

2020, 1–9. https://doi.org/10.1038/s41467-020-16240-6

Pavelka, M. S. M., McGoogan, K. C., & Steffens, T. S. (2007). Population Size and Characteristics of *Alouatta pigra*

Before and After a Major Hurricane. *International Journal of Primatology*, *28*(4), 919–929.

https://doi.org/10.1007/s10764-007-9136-6

Pijanowski, B. C., Farina, A., Gage, S. H., Dumyahn, S. L., & Krause, B. L. (2011a). What is soundscape ecology? An

introduction and overview of an emerging new science. *Landscape Ecology*, *26*(9), 1213–1232.

https://doi.org/10.1007/s10980-011-9600-8

Pijanowski, B. C., Villanueva-Rivera, L. J., Dumyahn, S. L., Farina, A., Krause, B. L., Napoletano, B. M., Gage, S. H.,

& Pieretti, N. (2011b). Soundscape Ecology: The Science of Sound in the Landscape. *BioScience*, *61*(3),

203–216. https://doi.org/10.1525/bio.2011.61.3.6

Pimm, S. L. (1984). The complexity and stability of ecosystems. *Nature*, *307*(5949), 321–326.

https://doi.org/10.1038/307321a0

R Core Team (2023). R: A language and environment for statistical computing. R Foundation for Statistical

Computing, Vienna, Austria. https://www.R-project.org/ [Version 4.3.0].

Rajan, S. C., Dominic, L., M, V., K, A., NP, S., & R, J. (2022). Surrogacy of post natural disaster acoustic indices for

biodiversity assessment. *Environmental Challenges*, *6*, 100420.

https://doi.org/10.1016/j.envc.2021.100420

Raymond, C., Horton, R. M., Zscheischler, J., Martius, O., AghaKouchak, A., Balch, J., Bowen, S. G., Camargo, S. J.,

Hess, J., Kornhuber, K., Oppenheimer, M., Ruane, A. C., Wahl, T., & White, K. (2020). Understanding

and managing connected extreme events. *Nature Climate Change*, *10*(7), 611–621.

https://doi.org/10.1038/s41558-020-0790-4

Roe, P., Eichinski, P., Fuller, R. A., McDonald, P. G., Schwarzkopf, L., Towsey, M., … Watson, D. M. (2021). The

Australian acoustic observatory. *Methods in Ecology and Evolution*, *12*, 1802-1808.

Ross, S. R. P.-J. (2022). A Suburban Soundscape Reveals Altered Acoustic Dynamics during the COVID-19

Lockdown. *Journal of Ecoacoustics, 6(1)*, 1–18. https://doi.org/10.35995/jea6010001

Ross, S. R. P.-J., Friedman, N. R., Dudley, K. L., Yoshimura, M., Yoshida, T., & Economo, E. P. (2018). Listening to

ecosystems: Data-rich acoustic monitoring through landscape-scale sensor networks. *Ecological*

Research, *33*(1), 135–147. https://doi.org/10.1007/s11284-017-1509-5

Ross, S. R. P.-J., Friedman, N. R., Janicki, J., & Economo, E. P. (2019). A test of trophic and functional island

biogeography theory with the avifauna of a continental archipelago. *Journal of Animal Ecology*, *88*(9),

1392–1405. https://doi.org/10.1111/1365-2656.13029

Ross, S. R. P.-J., Friedman, N. R., Yoshimura, M., Yoshida, T., Donohue, I., & Economo, E. P. (2021a). Utility of

acoustic indices for ecological monitoring in complex sonic environments. *Ecological Indicators*, *121*,

107114–107114. https://doi.org/10.1016/j.ecolind.2020.107114

Ross, S. R. P.-J., O'Connell, D. P., Deichmann, J. L., Desjonquères, C., Gasc, A., Phillips, J. N., … Burivalova, Z.

(2023a). Passive Acoustic Monitoring provides a fresh perspective on fundamental ecological

questions. *Functional Ecology 37*(4), 959-975. https://doi.org/10.1111/1365-2435.14275

Ross, S. R. P.-J., Suzuki, Y., Kondoh, M., Suzuki, K., Villa Martín, P., & Dornelas, M. (2021b). Illuminating the

intrinsic and extrinsic drivers of ecological stability across scales. *Ecological Research*, *36*(3), 364–378.

https://doi.org/10.1111/1440-1703.12214

Rossi, T., Connell, S. D., & Nagelkerken, I. (2017). The sounds of silence: Regime shifts impoverish marine

soundscapes. *Landscape Ecology*, *32*, 239–248. https://doi.org/10.1007/s10980-016-0439-x

Seki, S.-I. (2005). The effects of a typhoon (9918 Bart, 1999) on the bird community in a warm temperate forest,

Southern Japan. *Ornithological Science*, *4*(2), 117–128. https://doi.org/10.2326/osj.4.117

Senzaki, M., Barber, J. R., Phillips, J. N., Carter, N. H., Cooper, C. B., Ditmer, M. A., … Mennitt, D. J. (2020).

Sensory pollutants alter bird phenology and fitness across a continent. *Nature*, *587*, 605–609.

https://doi.org/10.1038/s41586-020-2903-7

Sethi, S. S., Ewers, R. M., Jones, N. S., Signorelli, A., Picinali, L., & Orme, C. D. L. (2020a). SAFE Acoustics: An

open-source, real-time eco-acoustic monitoring network in the tropical rainforests of Borneo. *Methods*

in Ecology and Evolution, *11*, 1182–1185. https://doi.org/10.1111/2041-210X.13438

Sethi, S. S., Jones, N. S., Fulcher, B. D., Picinali, L., Clink, D. J., Klinck, H., … Ewers, R. M. (2020b). Characterizing

soundscapes across diverse ecosystems using a universal acoustic feature set. *Proceedings of the*

National Academy of Sciences, *117*, 17049-17055.

Simmons, K. R., Eggleston, D. B., & Bohnenstiehl, D. W. R. (2021). Hurricane impacts on a coral reef soundscape.

PLoS ONE, *16*(2 February 2021), 1–27. https://doi.org/10.1371/journal.pone.0244599

Sirami, C., Brotons, L., Burfield, I., Fonderflick, J., & Martin, J.-L. (2008). Is land abandonment having an impact

on biodiversity? A meta-analytical approach to bird distribution changes in the north-western

Mediterranean. *Biological Conservation*, *141*(2), 450–459.

https://doi.org/10.1016/j.biocon.2007.10.015

Stan Development Team (2020). RStan: the R interface to Stan. R package version 2.21.2. https://mc-stan.org/

Suding, K. N., Lavorel, S., Chapin, F. S., Cornelissen, J. H. C., Díaz, S., Garnier, E., … Navas, M. L. (2008). Scaling

environmental change through the community-level: A trait-based response-and-effect framework for

plants. *Global Change Biology*, *14*(5), 1125–1140. https://doi.org/10.1111/j.1365-2486.2008.01557.x

Sueur, J., Krause, B., & Farina, A. (2019). Climate Change Is Breaking Earth's Beat. *Trends in Ecology and*

Evolution, *34*(11), 971–973. https://doi.org/10.1016/j.tree.2019.07.014

Takeuchi, K., Yoshioka, S.-I., & Fumoto, R. (1981). Land transformation on okinawa island, southwest Japan.

Geographical Reports of Tokyo Metropolitan University, *16*, 113–129.

Testard, C., Larson, S. M., Watowich, M. M., Kaplinsky, C. H., Bernau, A., Faulder, M., … Brent, L. J. N. (2021).

Rhesus macaques build new social connections after a natural disaster. *Current Biology*, *31*(11), 2299-

2309.e7. https://doi.org/10.1016/j.cub.2021.03.029

Tilman, D., Reich, P. B., & Knops, J. M. H. (2006). Biodiversity and ecosystem stability in a decade-long grassland

experiment. *Nature*, *441*(7093), 629–632. https://doi.org/10.1038/nature04742

Toth, C. A., Pauli, B. P., McClure, C. J. W., Francis, C. D., Newman, P., Barber, J. R., & Fristrup, K. (2022). A

stochastic simulation model for assessing the masking effects of road noise for wildlife, outdoor

recreation, and bioacoustic monitoring. *Oecologia*, *199*(1), 217–228. https://doi.org/10.1007/s00442-

022-05171-2

Uchida, K., & Ushimaru, A. (2014). Biodiversity declines due to abandonment and intensification of agricultural

lands: Patterns and mechanisms. *Ecological Monographs*, *84*(4), 637–658. https://doi.org/10.1890/13-

2170.1

Van Parijs, S. M., Baumgartner, M., Cholewiak, D., Davis, G., Gedamke, J., Gerlach, D., … Thompson, M. (2015).

NEPAN: A US Northeast passive acoustic sensing network for monitoring, reducing threats and the

conservation of marine animals. *Marine Technology Society Journal, 49*, 70-86.

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-

validation and WAIC. *Statistics and Computing, 27*, 1413–1432.

Villanueva-Rivera, L. J., & Pijanowski, B. C. (2018). soundecology: Soundscape Ecology. R package version 1.3.3.

https://CRAN.R-project.org/package=soundecology

Vogel, A., Manning, P., Cadotte, M. W., Cowles, J., Isbell, F., Jousset, A. L. C., … Wagg, C. (2019). Lost in trait

space: Species-poor communities are inflexible in properties that drive ecosystem functioning. In

Advances in Ecological Research (1st ed., Vol. 61, p. 131). Elsevier Ltd.

https://doi.org/10.1016/bs.aecr.2019.06.002

Vokurková, J., Motombi, F. N., Ferenc, M., Hořák, D., & Sedláček, O. (2018). Seasonality of vocal activity of a bird

community in an Afrotropical lowland rain forest. *Journal of Tropical Ecology*, *34*, 53–64.

https://doi.org/10.1017/S0266467418000056

Wang, S., Loreau, M., Arnoldi, J.-F., Fang, J., Rahman, K. Abd., Tao, S., & de Mazancourt, C. (2017). An

invariability-area relationship sheds new light on the spatial scaling of ecological stability. *Nature

Communications*, *8*(May), 15211–15211. https://doi.org/10.1038/ncomms15211

Wang, S., Loreau, M., de Mazancourt, C., Isbell, F., Beierkuhnlein, C., Connolly, J., … Craven, D. (2021). Biotic

homogenization destabilizes ecosystem functioning by decreasing spatial asynchrony. *Ecology*, *102*(6),

1–10. https://doi.org/10.1002/ecy.3332

White, L., O'Connor, N. E., Yang, Q., Emmerson, M. C., & Donohue, I. (2020). Individual species provide

multifaceted contributions to the stability of ecosystems. *Nature Ecology and Evolution*, *1*.

https://doi.org/10.1038/s41559-020-01315-w

Wiley, J. W., & Wunderle, J. M. (1993). The effects of hurricanes on birds, with special reference to Caribbean

islands. *Bird Conservation International*, *3*(4), 319–349. https://doi.org/10.1017/S0959270900002598

Willig, M. R., & Camilo, G. R. (1991). The Effect of Hurricane Hugo on Six Invertebrate Species in the Luquillo

Experimental Forest of Puerto Rico. *Biotropica*, *23*(4), 455–461. https://doi.org/10.2307/2388266

Yang, Q., Fowler, M. S., Jackson, A. L., & Donohue, I. (2019). The predictability of ecological stability in a noisy

world. *Nature Ecology and Evolution*, *3*(February), 31–33. https://doi.org/10.1038/s41559-018-0794-x

Zampieri, N. E., Pau, S., & Okamoto, D. K. (2020). The impact of Hurricane Michael on longleaf pine habitats in

Florida. *Scientific Reports*, *10*(1), 1–11. https://doi.org/10.1038/s41598-020-65436-9

Zelnik, Y. R., Arnoldi, J.-F., & Loreau, M. (2018). The Impact of Spatial and Temporal Dimensions of Disturbances

on Ecosystem Stability. *Frontiers in Ecology and Evolution*, *6*.

https://www.frontiersin.org/article/10.3389/fevo.2018.00224

Zhang, Q., Hong, Y., Zou, F., Zhang, M., Lee, T. M., Song, X., & Rao, J. (2016). Avian responses to an extreme ice

storm are determined by a combination of functional traits, behavioural adaptations and habitat

modifications. *Scientific Reports*, *6*(1), Article 1. https://doi.org/10.1038/srep22344